



HAL
open science

MULTIGRADED REGULARITY OF COMPLETE INTERSECTIONS

Marc Chardin, Navid Nemati

► **To cite this version:**

Marc Chardin, Navid Nemati. MULTIGRADED REGULARITY OF COMPLETE INTERSECTIONS. 2021. hal-03369633

HAL Id: hal-03369633

<https://hal.science/hal-03369633>

Preprint submitted on 7 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTIGRADED REGULARITY OF COMPLETE INTERSECTIONS

MARC CHARDIN AND NAVID NEMATI

ABSTRACT. V is a complete intersection scheme in a multiprojective space if it can be defined by an ideal I with as many generators as $\text{codim}(V)$. We investigate the multigraded regularity of complete intersections scheme in $\mathbb{P}^n \times \mathbb{P}^m$. We explicitly compute many values of the Hilbert functions of 0-dimensional complete intersections. We show that these values only depend upon n, m , and the bidegrees of the generators of I . As a result, we provide a sharp upper bound for the multigraded regularity of 0-dimensional complete intersections.

1. INTRODUCTION

The theory of syzygies offers a microscope for looking at systems of equations. Castelnuovo-Mumford regularity is an important invariant in commutative algebra and algebraic geometry which is strongly related to syzygies. D. Eisenbud and S. Goto in [2] showed that Castelnuovo-Mumford regularity can be obtained from the minimal free resolution. The minimal free resolution was first introduced by Hilbert to study Hilbert function. It is one of the finest invariant that we can associate with a finitely generated graded module M over a polynomial ring. Castelnuovo-Mumford regularity measures the maximum degree of the syzygies and provides a quantitative version of Serre vanishing theorem for the associated sheaf. In particular, it bounds the largest degree of the minimal generators and the smallest twist for which the sheaf is generated by its global sections. It has been used as a measure for the complexity of computational problems in algebraic geometry and commutative algebra. The two most frequent definitions of \mathbb{Z} -graded Castelnuovo-Mumford regularity are the one in terms of graded Betti numbers and the one using local cohomology.

An extension of Castelnuovo-Mumford regularity for the multigraded setting in a special case was first introduced by Hoffman and Wang in [6]. Later by Maclagan and Smith in [8], and Botbol and Chardin in [1] in a more general setting. The main motivation for studying regularity over multigraded polynomial rings was from toric geometry. Maclagan and Smith [8] developed a multigraded theory of regularity of sheaves on a simplicial toric variety \mathcal{X} with an algebraic variant defined in terms of the vanishing of graded pieces of $H_B^i(M)$, the i -th local cohomology module of M . Here B is the irrelevant ideal of the homogeneous coordinate ring of \mathcal{X} .

The conceptual difficulty of computing multigraded regularity lies in the simple fact that bounded subsets of \mathbb{Z}^r need not have single maximal or minimal elements. This makes it very hard to capture the vanishing or non-vanishing of the

2010 *Mathematics Subject Classification.* 13D02, 13D40, 14B15, 13D45.

Key words and phrases. Multigraded regularity, Hilbert function, multiprojective space, complete intersection.

multigraded pieces of local cohomology. One can define a complete intersection in projective space as a subscheme defined by as many forms as its codimension. In this case, it corresponds to complete intersection homogeneous ideals; therefore many homological invariants are determined by the degrees of the forms. One natural question that may arise is to ask the same question for the complete intersection schemes in a product of projective spaces.

Proposition 6.7 in [8] implies that the multigraded graded regularity a 0-dimensional schemes is the same as the stabilization region of the Hilbert functions. Understanding the Hilbert functions and the minimal free resolutions of the coordinate rings of points in multiprojective space is included among the list of open problems in commutative algebra found in the survey article of Peeva-Stillman[9].

Based on the above motivations, our main concern in this article is to study multigraded regularity and bigraded Hilbert function complete intersection points in $\mathbb{P}^n \times \mathbb{P}^m$. In Section 2, we start by setting our notation and define multigraded regularity. We show that if the ring is bigraded, then one can consider an alternative way to define the multigraded regularity (see Theorem 2.5).

In section 3, we study the multigraded regularity of general complete intersection schemes in $\mathbb{P}^n \times \mathbb{P}^m$. A scheme $V \subset \mathbb{P}^n \times \mathbb{P}^m$ is complete intersection if it can be defined by as many forms as its codimension. Our motivation is to show which homological invariants are determined by the degrees of these forms.

In Section 4, we focus on the case of complete intersection points in $\mathbb{P}^n \times \mathbb{P}^m$. The main result of this section is Theorem 4.3. For a complete intersection scheme of point $V \subseteq \mathbb{P}^n \times \mathbb{P}^m$ defined by f_1, \dots, f_{n+m} , many values of its bigraded Hilbert function are independent from the choices of f_i 's and they only depend upon their bidegrees. Moreover, in some regions, there is a duality among these values.

Example A. (Example 4.4) Let $S = k[x_0, x_1, x_2, y_0, y_1, y_2]$, and $I = (f_1, \dots, f_4)$ is generated by 4 bigraded forms of bidegree $(2, 2)$ such that $\text{proj}(S/I) = V$ is complete intersection scheme of points. For any $\mu \notin (2, 6) + (-\mathbb{N}, \mathbb{N}) \cup (6, 2) + (\mathbb{N}, -\mathbb{N})$, the Hilbert function of V at μ is independent from the choices of f_i 's and can be computed explicitly via Theorem 4.3.

In Section 5, we study generic complete intersection points in $\mathbb{P}^n \times \mathbb{P}^m$. The main Theorem of this section is the following

Theorem A. (Theorem 5.5) Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring over a field k of characteristics zero. If I is generated by $n + m$ generic forms of bidegree (d, e) , then the scheme V defined by I is a set of reduced points and

$$|\{\mu \in \mathbb{N}^2 \mid \text{HF}_V(\mu) \neq \text{deg}(V)\}| < \infty,$$

which means the natural projections are one-to-one.

2. MULTIGRADED REGULARITY IN BIGRADED SETTING

Notation 2.1. From now on we use the following notation in the rest of the paper. Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring where $\text{deg}(x_i) = (1, 0)$ and $\text{deg}(y_i) = (0, 1)$. Let $B_1 = (x_0, \dots, x_n)$, and $B_2 = (y_0, \dots, y_m)$ be the irrelevant ideal of \mathbb{P}^n and \mathbb{P}^m . Define $B := B_1 \cdot B_2$ the irrelevant ideals of $\mathbb{P}^n \times \mathbb{P}^m$, and $\mathfrak{m} = B_1 + B_2$ the maximal ideal of S as a standard graded ring.

We state the definition of multigraded regularity in the case of bigraded rings; for further reading, we refer readers to [1, 8].

Definition 2.2. Let M be a graded S -module. The support of the module M is

$$\text{Supp}(M) := \{\gamma \mid M_\gamma \neq 0\}.$$

Definition 2.3. (Multigraded regularity) Let $C \subseteq \mathfrak{m}$ be a finitely generated bigraded ideal of S . Define for a $\gamma \in \mathbb{Z}^2$, M is (C, γ) -**regular** if

$$\gamma + \mathbb{N}^2 \cap \cup_i \text{Supp}(H_C^i(M)) + \mathcal{F}_{i-1} = \emptyset,$$

where $\mathcal{F}_i := \{(i-1, 0), (i-2, 1), \dots, (0, i-1)\}$ for $i > 0$ and $\mathcal{F}_0 := \{0\}$, $\mathcal{F}_{-1} = -\mathcal{F}_1$, and $\mathcal{F}_i = 0$ for $i < -1$. C -regularity of M is

$$\text{reg}_C(M) := \{\gamma \in \mathbb{Z}^r \mid M \text{ is } (C, \gamma)\text{-regular}\}.$$

In particular, if $C = B$, B -regularity of M , $\text{reg}_B(M)$, is called the multigraded regularity of M .

Our approach for computing multigraded regularity relies on Mayer-Vietoris sequence to relate the various cohomology modules.

Remark 2.4. Let $S = k[x_1, \dots, x_m, y_1, \dots, y_n]$ be a bigraded polynomial ring. Define B_1, B_2, B and \mathfrak{m} as in Notation 2.1. Then we have the following complex between local cohomology modules:

$$\dots \rightarrow H_B^{i-1}(M) \rightarrow H_{\mathfrak{m}}^i(M) \rightarrow H_{B_1}^i(M) \oplus H_{B_2}^i(M) \rightarrow H_B^i(M) \rightarrow \dots$$

The following Theorem provide an alternative way to define multigraded regularity in the bigraded setting by knowing \mathfrak{m} , B_1 , and B_2 -regularity.

Theorem 2.5. Let $S = k[x_1, \dots, x_m, y_1, \dots, y_n]$ be a bigraded polynomial ring and M is a graded S -module. Adopt Notations 2.1, then

$$\text{reg}_B(M) = \text{reg}_{\mathfrak{m}}(M) \cap \text{reg}_{B_1}(M) \cap \text{reg}_{B_2}(M).$$

Proof. By analyzing the spectral sequences correspond to Čech-Koszul double complex $C_B^\bullet K_\bullet(\underline{\mathbf{x}}, \underline{\mathbf{y}}; M)$ we get

$$\text{Supp}(\text{Tor}_i^S(M, k)) \subseteq \cup_{j=0}^i \text{Supp}(H_B^j(M)) + \mathcal{E}_{i+j}$$

where \mathcal{E}_i denote the set of twists of the summands in the i -th step of the minimal free resolution of B ; see [1, Section 4.2] for more details regarding the relation between B -regularity and the support of Tor modules. By [1, Corollary 3.12], for every i we have

$$\begin{aligned} \text{Supp}(H_{\mathfrak{m}}^i(M)) &\subseteq \cup \text{Supp}(\text{Tor}_{m+n-i}^S(M, k)) + \text{Supp}(H_{\mathfrak{m}}^{m+n}(S)) \\ &\subseteq \cup \text{Supp}(\text{Tor}_{m+n-i}^S(M, k) - (m, n) - \mathbb{N}^2) \\ &\subseteq \cup \cup_{j=0}^i \text{Supp}(H_B^j(M)) + \mathcal{E}_{m+n-(i-j)} - (m, n) - \mathbb{N}^2 \\ &\subseteq \cup_{j=0}^i \text{Supp}(H_B^j(M)) + \mathcal{E}_{i-i} - \mathbb{N}^2. \end{aligned}$$

Therefore

$$\text{Supp}(H_{\mathfrak{m}}^i(M)) + \mathcal{F}_{i-1} \subseteq \cup_{j=0}^i \text{Supp}(H_B^j(M)) + \mathcal{F}_{j-1} - \mathbb{N}^2,$$

which means

$$(2.1) \quad \text{reg}_B(M) \subseteq \text{reg}_{\mathfrak{m}}(M).$$

Suppose $\mu \in \text{reg}_B(M)$. By the definition, $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \cap \text{Supp} H_B^i(M) = \emptyset$ for every i . By 2.1, $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \cap \text{Supp} H_{\mathfrak{m}}^i(M) = \emptyset$. By using Mayer-Vietoris

exact sequence one gets $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \cap (\text{Supp } H_{B_1}^i(M) \cap \text{Supp } H_{B_2}^i(M)) = \emptyset$, that implies

$$\text{reg}_B(M) \subseteq \text{reg}_m(M) \cap \text{reg}_{B_1}(M) \cap \text{reg}_{B_2}(M).$$

Suppose $\mu \in \text{reg}_m(M) \cap \text{reg}_{B_1}(M) \cap \text{reg}_{B_2}(M)$. By the definition, for every i , $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \cap (\text{Supp } H_{B_1}^i(M) \cap \text{Supp } H_{B_2}^i(M)) = \emptyset$, and $(\mu - \mathcal{F}_i + \mathbb{N}^2) \cap \text{Supp } H_m^{i+1}(M) = \emptyset$. Since $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \subset (\mu - \mathcal{F}_i + \mathbb{N}^2)$, $(\mu - \mathcal{F}_{i-1} + \mathbb{N}^2) \cap \text{Supp } H_m^{i+1}(M) = \emptyset$. The assertion follows from Mayer-Vietoris exact sequence and the definition of $\text{reg}_B(M)$. \square

3. COMPLETE INTERSECTIONS IN $\mathbb{P}^n \times \mathbb{P}^m$

We start this section by stating the definition of complete intersection scheme in a product of projective that we are using in this article.

Definition 3.1. *A subscheme $V \subseteq \mathbb{P}^n \times \mathbb{P}^m$, is a complete intersection if $V = \text{proj}(S/I)$ where I is generated by $\text{codim}(V)$ bihomogeneous elements.*

Let V be a complete intersection scheme in a projective space, then there exist ideal I defining V with as many generators as the codimension of V . In this case, ideal I is complete intersection, therefore many homological invariants are entirely determined by the degrees of the generators. Let $V = \text{proj}(S/I) \subset \mathbb{P}^n \times \mathbb{P}^m$ be a complete intersection scheme, where I is generated by $\text{codim}(V)$ forms. In this case, I is not complete intersection unless $n, m = 1$ in which case I is not the defining ideal of V (it is not B -saturated). Although the similar argument would not work in this case, but our goal is to show that many homological invariants are only depend upon the degrees of the generators of I .

Proposition 3.2. [4, Example 8.4.2] *Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring where $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Let $I = (f_1, \dots, f_{n+m})$ generated by $n + m$ forms of degree (d_i, e_i) and V be the scheme defined by I . If I is a complete intersection, then*

$$\deg(V) = \sum d_{i_1} \cdots d_{i_n} \cdot e_{j_1} \cdots e_{j_m},$$

where the sum is over all permutations $(d_{i_1}, \dots, d_{i_n}, e_{j_1}, \dots, e_{j_m})$ of $(1, \dots, n + m)$ with $i_1 < i_2 < \dots < i_n$ and $j_1 < j_2 < \dots < j_m$. In particular if $d_i = d$ and $e_i = e$ for all i , then

$$\deg(V) = \binom{n+m}{n} d^n e^m.$$

Proposition 3.3. *Let $I = (f_1, \dots, f_r)$ be a bigraded ideal defining a complete intersection of codimension r and let $\deg(f_i) = \mathbf{d}_i$. For $\mu \in \cap_i \text{reg}_B(H_i(\mathbf{K}(\mathbf{f}, S)))$,*

$$\text{HF}_{S/I}(\mu) = P(\mu),$$

where P is a polynomial that only depends upon $\mathbf{d}_1, \dots, \mathbf{d}_r$.

Proof. By the Serre Grothendieck formula [1, Proposition 4.27]

$$(3.1) \quad \text{HP}_{S/I}(\mu) = \text{HF}_{S/I}(\mu) + \sum_i (-1)^i \dim H_B^i(S/I)_\mu.$$

Hence, if $\mu \in \text{reg}_B(S/I)$ then $HP_{S/I}(\mu) = \text{HF}_{S/I}(\mu)$. Set

$$(3.2) \quad \chi(\mu) := \sum_i (-1)^i \dim(H_i(\mathbf{K}(\mathbf{f}, S))_\mu) = \sum_i (-1)^i \dim(K_i)_\mu.$$

Note that the second equality shows that χ is a function that only depends upon the degrees. Since I is complete intersection, $H_i(\mathbf{K}(\mathbf{f}, S)) = H_B^0(H_i(\mathbf{K}(\mathbf{f}, S)))$. Hence, $\text{Supp}(H_i(\mathbf{K}(\mathbf{f}, S))) \cap \text{reg}_B(H_i(\mathbf{K}(\mathbf{f}, S))) = \emptyset$. Therefore, for $\mu \in \cap_{i>0} \text{reg}_B(H_i(\mathbf{K}(\mathbf{f}, S)))$, $\chi(\mu) = \text{HF}_{S/I}(\mu)$. Hence, for $\mu \in \cap_i \text{reg}_B(H_i(\mathbf{K}(\mathbf{f}, S)))$,

$$HP_{S/I}(\mu) = \text{HF}_{S/I}(\mu) = \chi(\mu).$$

Setting $P := HP_{S/I}$ the first equality shows that $\text{HF}_{S/I}(\mu) = P(\mu)$ and the second equality shows this function only depends upon the degrees. \square

Proposition 3.4. *With the Notation 2.1, for any bigraded free S -module M ,*

$$H_B^{n+m+1}(M) \cong H_m^{n+m+2}(M).$$

Furthermore, If $n = m$ then

$$H_B^{n+1}(M) \cong H_{B_1}^{n+1}(M) \oplus H_{B_2}^{n+1}(M),$$

else if $n < m$ then,

$$H_B^{n+1}(M) \cong H_{B_1}^{n+1}(M) \quad \text{and} \quad H_B^{m+1}(M) \cong H_{B_2}^{m+1}(M).$$

Proof. Note that $H_{B_1}^a(M) = H_{B_2}^b(M) = H_m^c(M) = 0$ if $a \neq n+1$, $b \neq m+1$ and $c \neq n+m+2$. The Mayer-Vietoris exact sequence

$$\cdots \rightarrow H_m^i(M) \rightarrow H_{B_1}^i(M) \oplus H_{B_2}^i(M) \rightarrow H_B^i(M) \rightarrow H_m^{i+1}(M) \rightarrow \cdots$$

gives the results. \square

Theorem 3.5. *Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring where $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Define $B = (x_0, \dots, x_n) \cdot (y_0, \dots, y_m)$ the irrelevant ideal of $\mathbb{P}^n \times \mathbb{P}^m$. Assume $I = (f_1, \dots, f_r)$ with $\deg(f_i) = (d_i, e_i)$ and V be the scheme defined by I . If $\text{codim}(V) = r$ then*

- $\text{Supp}(H_i(\mathbf{K}(\mathbf{f}, S))) = \text{Supp}(V_{n+1+i}) \cup \text{Supp}(W_{m+1+i})$ for $i > 0$
- $\text{Supp}(H_B^i(S/I)) = \text{Supp}(V_{n+1-i}) \cup \text{Supp}(W_{m+1-i})$ for $0 \leq i \leq \dim V$,
- $\text{Supp}(H_B^{\dim V+1}(S/I)) = \{(\mu, \nu) \mid H^{\dim V}(V, \mathcal{O}_V(\mu, \nu)) \neq 0\}$
 $\subseteq \text{Supp}(V_{r-m}) \cup \text{Supp}(W_{r-n}) \cup \{\sigma - \text{Supp}(S/I)\}$,

where V_i (resp. W_i) is a subquotient of $H_{B_1}^{n+1}(K_i(\mathbf{f}, S))$ (resp. $H_{B_2}^{m+1}(K_i(\mathbf{f}, S))$), and $\sigma := \sum_i (d_i, e_i) - (n+1, m+1)$.

Proof. Consider the double complex $\mathbf{C}_B^\bullet(\mathbf{K}_\bullet(\mathbf{f}, S))$ and suppose $n \leq m$. If we start taking homologies vertically, by 3.4 in the third page we have:

$$\begin{array}{cccccc} 0 & 0 & \cdots & 0 & 0 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ V_r & V_{r-1} & \cdots & V_1 & V_0 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ W_r & W_{r-1} & \cdots & W_1 & W_0 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ M & \cdots & 0 & 0 & 0, & \end{array}$$

where

$$\begin{aligned}
M &\cong H_r(H_B^{n+m+1}(\mathbf{K}(\mathbf{f}, S))) \\
&\cong H_r(H_m^{n+m+2}(\mathbf{K}(\mathbf{f}, S))) \\
&\cong H_r((\mathbf{K}(\mathbf{f}, S)(\sigma))^*) \\
&\cong (S/I)^*(\sigma).
\end{aligned}$$

If we start taking homology horizontally, in the second page we have

$$\begin{array}{cccccc}
\cdots & 0 & H_\gamma & \cdots & H_1 & H_B^0(S/I) \\
\cdots & 0 & 0 & 0 & 0 & H_B^1(S/I) \\
\cdots & 0 & 0 & 0 & 0 & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & H_B^{\dim(V)+1}(S/I) \\
\vdots & \vdots & \vdots & \vdots & \vdots & 0,
\end{array}$$

where $\gamma := \max\{0, r - n - 1, r - m - 1\}$. The result is obtained by comparing the two abutments. \square

Definition 3.6. Let $\mathbf{d} = (d, e) \in \mathbb{N}^2$ and $r \in \mathbb{N}$, define

$$\begin{aligned}
v_i &:= i \cdot \mathbf{d} - (n + 1, 0), \\
w_i &:= i \cdot \mathbf{d} - (0, m + 1), \\
\sigma &:= r \cdot \mathbf{d} - (n + 1, m + 1).
\end{aligned}$$

Lemma 3.7. Let $I = (f_1, \dots, f_r) \subseteq S$ and $\deg(f_i) = \mathbf{d} = (d, e)$. Suppose V is the scheme defined by I , if $\text{codim}(V) = r$ then

$$\begin{aligned}
\text{Supp}(H_i(\mathbf{K}(\mathbf{f}, S))) &\subseteq v_{n+1+i} + (-\mathbb{N}, \mathbb{N}) \cup w_{m+1+i} + (\mathbb{N}, -\mathbb{N}) \quad \text{for } i > 0, \\
\text{Supp}(H_B^i(S/I)) &\subseteq v_{n+1-i} + (-\mathbb{N}, \mathbb{N}) \cup w_{m+1-i} + (\mathbb{N}, -\mathbb{N}) \quad \text{for } 0 \leq i \leq \dim V.
\end{aligned}$$

In addition, the inclusions are sharp.

Proof. With the proof of Theorem 3.5, in the second page of the spectral sequence we have for $p < n + m + 1$

$${}^v E_{p,q}^\infty = {}^v E_{p,q}^2 = \begin{cases} H_q(H_{B_1}^{n+1} \mathbf{K}(\mathbf{f}, S)) & \text{if } p = n + 1, \\ H_q(H_{B_2}^{n+1} \mathbf{K}(\mathbf{f}, S)) & \text{if } p = m + 1, \\ 0 & \text{else.} \end{cases}$$

Combining with the Theorem 3.5 shows the inclusions. The sharpness follows from the fact that for any q , v_q (resp. w_q) is in the support of $H_{B_1}^{n+1} K_q(\mathbf{f}, S)$ (resp. $H_{B_2}^{m+1} K_q(\mathbf{f}, S)$) and it is not in the support of $H_{B_1}^{n+1} K_{q-1}(\mathbf{f}, S)$ and $H_{B_1}^{n+1} K_{q+1}(\mathbf{f}, S)$ (resp. $H_{B_2}^{m+1} K_{q-1}(\mathbf{f}, S)$ and $H_{B_2}^{m+1} K_{q+1}(\mathbf{f}, S)$). \square

Corollary 3.8. Let $I = (f_1, \dots, f_r) \subseteq S$ and $\deg(f_i) = \mathbf{d} = (d, e)$ and V be the scheme defined by I . If $\text{codim}(V) = r$, then

- (1) if $\mu \notin v_i + (-\mathbb{N}, \mathbb{N}) \cup w_j + (\mathbb{N}, -\mathbb{N})$ for $i = r - m, r - m - 1$ and $j = r - n, r - n - 1$ then

$$H_B^{\dim(V)+1}(S/I)_\mu = (S/I)_{\sigma - \mu}^*.$$

Proof. Note that $\text{Supp}(H_{B_1}^{n+1}(K_i(\mathbf{f}, S))) = v_i + (-\mathbb{N}, \mathbb{N})$ and $\text{Supp} H_{B_2}^{m+1}(K_i(\mathbf{f}, S)) = w_i + (\mathbb{N}, +\mathbb{N})$ for all i . Consider the double complex $\mathbf{C}_B^\bullet(\mathbf{K}_\bullet(\mathbf{f}, S))$ and suppose $n \leq m$ as in the proof of Theorem 3.5. Since $\text{Supp}(V_i) \cap \text{Supp}(W_j) = \emptyset$, there will be no nonzero map in the spectral sequence among them. In this case, in the $n+2$ -th page we have an induced map $\psi_1 : M \rightarrow W_{r-n-1}$, and in the $m+2$ -th page we have an induced map $\psi_2 : \ker \psi_1 \rightarrow V_{r-m-1}$. If $\mu \notin v_{r-m} + (-\mathbb{N}, \mathbb{N}) \cup w_{r-n} + (\mathbb{N}, -\mathbb{N})$, $H_B^r(S/I)_\mu \cong (\ker \psi_2)_\mu$. In addition, if $\mu \notin v_{r-m-1} + (-\mathbb{N}, \mathbb{N}) \cup w_{r-n-1} + (\mathbb{N}, -\mathbb{N})$

$$H_B^{\dim(V)+1}(S/I)_\mu \cong (\ker \psi_2)_\mu = M_\mu = (S/I)_{\sigma-\mu}^*.$$

□

4. COMPLETE INTERSECTION POINTS IN $\mathbb{P}^n \times \mathbb{P}^m$

In this section we apply the results in the previous section to the case of complete intersection points in $\mathbb{P}^n \times \mathbb{P}^m$. In the rest of this section, $I = (f_1, \dots, f_{n+m})$ and V be the complete intersection scheme of points defined by I with $\mathbf{d} := \deg(f_i) = (d, e)$.

Definition 4.1. Set $\Gamma_i := \text{Supp}(H_{B_1}^{n+1}(K_{n+i}(\mathbf{f}, S))) \cup \text{Supp}(H_{B_2}^{m+1}(K_{m+i}(\mathbf{f}, S)))$. Notice that $\Gamma_i = \emptyset$ if and only if $i \geq \max\{n, m\}$. If $i \leq m$ then

$$\text{Supp}(H_{B_1}^{n+1}(K_{n+i}(\mathbf{f}, S))) = v_{n+i} + (-\mathbb{N}, \mathbb{N}),$$

and similarly for $\text{Supp}(H_{B_2}^{m+1}(K_{m+i}(\mathbf{f}, S)))$ if $i \leq n$.

Definition 4.2. For a function $F : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ define

$$\begin{aligned} F^*(a, b) &:= F(-a, -b), \\ F'(a, b) &:= F(-a, b), \text{ and} \\ F''(a, b) &:= F(a, -b). \end{aligned}$$

Theorem 4.3. Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring where $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Assume V be a complete intersection scheme of points defined by $I = (f_1, \dots, f_{n+m})$ with $\deg(f_i) = (d, e)$.

(1) If $\mu \notin \cup_{i \geq 2} \Gamma_i$ then

$$\text{HF}_{S/I}(\mu) = \chi(\mu)$$

$$\text{where } \chi((a, b)) = \sum_i (-1)^i \binom{n+m}{i} \binom{n+a-id}{n} \binom{m+b-ie}{m}.$$

(2) If $\mu \notin \Gamma_1$ then

$$\text{HF}_{S/I}(\mu) = \text{HF}_V(\mu)$$

(3) If $\mu \notin \Gamma_0$ then

$$\text{HF}_V(\mu) + \text{HF}_V(\sigma - \mu) = \deg(V) = \binom{n+m}{n} d^n e^m.$$

In particular, if $\mu \in (nd-n, (n+m)e-m) + \mathbb{N}^2 \cup ((n+m)d-n, me-m) + \mathbb{N}^2$ then $\text{HF}_V(\mu) = \deg(V)$ and $\mu \in \text{reg}_B(V)$.

(4) If $\mu \in \Gamma_0 \setminus \{\Gamma_{-1} \cup \Gamma_1\}$ then

$$\text{HF}_V(\mu) + \text{HF}_V(\sigma - \mu) = \binom{n+m}{n} (d^n e^m - \epsilon(\mu)),$$

where $\epsilon(\mu) = \text{HF}'_S(\mu - v_n) + \text{HF}''_S(\mu - w_n)$ as in the Definition 4.2.

(5) If $\mu \in \cup_{i \geq 1} \Gamma_i \setminus \{\Gamma_0\}$ then

$$\mathrm{HF}_V(\mu) = \binom{n+m}{n} d^n e^m - \chi(\sigma - \mu).$$

In particular, if $\mu \in \Gamma_1 \setminus \{\Gamma_0 \cup \Gamma_2\}$ then

$$\dim(I_V/I)_\mu = \binom{n+m}{n+1} \mathrm{HF}'_S(\mu - v_{n+1}) + \binom{n+m}{m+1} \mathrm{HF}''_S(\mu - w_{m+1}).$$

Proof. Denote the saturation of I with respect to B by J . In this case $\mathrm{HF}_V(\mu) = \dim(S/J)_\mu$ for all $\mu \in \mathbb{Z}^2$. In the proof we use these two simple fact that $\mu \in \Gamma_{i-1} \cup \Gamma_{i+1}$ yields $\mu \in \Gamma_i$ and if $\mu \in \Gamma_i$ then $\sigma - \mu \in \Gamma_{-i}$. By Proposition 3.2 the Hilbert polynomial of V is $D := \binom{n+m}{n} d^n e^m$ which in this case is equal to the $\deg(V)$. By Serre Grothendieck formula,

$$(4.1) \quad \mathrm{HF}_{S/J}(\mu) + \dim(H_B^1(S/J)_\mu) = D$$

(1) If $\mu \notin \cup_{i \geq 2} \Gamma_i$, by Lemma 3.7 $(H_i(K(\mathbf{f}, S)))_\mu = 0$ for all $i \geq 1$. Therefore by Equation 3.2, $(S/I)_\mu = \chi(\mu)$.

(2) If $\mu \notin \Gamma_1$, by Lemma 3.7, $(H_B^0(S/I))_\mu = 0$ therefore $(S/I)_\mu = (S/J)_\mu$.

(3) First note that $\mu, \sigma - \mu \notin \Gamma_0$. We claim that $\mu \notin \Gamma_{-1}$ or $\sigma - \mu \notin \Gamma_{-1}$. Suppose not, then $\mu \in \Gamma_{-1} \cap \Gamma_1$ which yields $\mu \in \Gamma_0$ which is a contradiction. So assume $\mu \notin \Gamma_{-1}$. By Corollary 3.8,

$$\dim(H_B^1(S/I)_\mu) = \mathrm{HF}_{S/I}(\sigma - \mu).$$

Since $\sigma - \mu \notin \Gamma_1$, by part (2) and Equation 3.1

$$\binom{n+m}{n} d^n e^m - \mathrm{HF}_V(\mu) \mathrm{HF}_V(\sigma - \mu).$$

The same argument works if we assume $\sigma - \mu \notin \Gamma_{-1}$.

(4) By part (1) $\mathrm{HF}_{S/I}(\mu) = \mathrm{HF}_V(\mu)$. By considering the double complex as in the proof of Theorem 3.5 and Lemma 3.7,

$$\begin{aligned} (H_n(H_{B_1}^{n+1} K(\mathbf{f}, S)))_\mu &= (H_{B_1}^{n+1}(K_n(\mathbf{f}, S)))_\mu, \text{ and} \\ (H_m(H_{B_2}^{m+1} K(\mathbf{f}, S)))_\mu &= (H_{B_2}^{m+1}(K_m(\mathbf{f}, S)))_\mu. \end{aligned}$$

By the abutment of the spectral sequence


$$(H_B^1(S/J))_\mu = \mathrm{HF}_{S/I}(\sigma - \mu) + \dim(H_{B_1}^{n+1}(K_n(\mathbf{f}, S)))_\mu + (H_{B_2}^{m+1}(K_m(\mathbf{f}, S)))_\mu.$$

Since $\sigma - \mu \notin \Gamma_1$, by parts (1) and (2), $\mathrm{HF}_{S/I}(\mu) = \mathrm{HF}_V(\mu)$ and by the definition, $\dim(H_{B_1}^{n+1}(K_n(\mathbf{f}, S)))_\mu + (H_{B_2}^{m+1}(K_m(\mathbf{f}, S)))_\mu = \binom{n+m}{n} \epsilon(\mu)$. The assertion follows by

$$(H_B^1(S/J))_\mu = \binom{n+m}{n} d^n e^m - \mathrm{HF}_V(\mu).$$

(5) By part (3), $\mathrm{HF}_V(\mu) = \binom{n+m}{n} d^n e^m - \mathrm{HF}_V(\sigma - \mu)$. Because $\mu \notin \Gamma_{-1}$ ($\mu \in \Gamma_{-1}$ yields $\mu \in \Gamma_0$) therefore $\sigma - \mu \notin \Gamma_1$, which by part (2), yields $\mathrm{HF}_V(\sigma - \mu) = \mathrm{HF}_{S/I}(\sigma - \mu)$. On the other hand, since $\mu \notin \cup_{i \geq 2} \Gamma_i$, by part (1), $\mathrm{HF}_{S/I}(\sigma - \mu) = \chi(\sigma - \mu)$. \square

Let $S = [x_0, x_1, x_2, y_0, y_1, y_2]$ and $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Let $I = (f_1, \dots, f_4)$ where $\deg(f_i) = (2, 2)$. Suppose scheme V defined by I is complete intersection. The following picture demonstrate the regions Γ_i in the Theorem 4.3.



Γ_{-1} is the red region, Γ_0 is the blue region and Γ_1 is the green region. By Theorem 4.3, Hilbert function of V at the bidegrees except the intersection of blue and green are independent from the choices of f_i 's and they can be computed via χ and ϵ defined in the Theorem 4.3. On the other hand, the rest do depends upon the f_i 's. Here we computed two example via computer system Macaulay2 [5].

Example 4.4. Let $S = k[x_0, x_1, x_2, y_0, y_1, y_2]$, $I = (x_0^2 y_0^2, x_1^2 y_1^2, x_2^2 y_2^2, (x_0 + x_1 + x_2)^2 (y_0 + y_1 + y_2)^2)$ and V be the complete intersection scheme of points defined by I . For $(0, 0) \leq \mu \leq (7, 7)$ the bigraded Hilbert function $\text{HF}_V(\mu)$ is

24	72	96	96	96	96	96	96
24	72	96	96	96	96	96	96
21	63	86	90	93	95	96	96
15	45	66	78	87	93	96	96
10	30	48	64	78	90	96	96
6	18	32	48	66	86	96	96
3	9	18	30	45	63	72	72
1	3	6	10	15	21	24	24

Where blue corresponds to the bidegrees in Γ_0 , green corresponds to Γ_1 and orange to their intersections. Also red indicate the $\text{HF}_V(\sigma)$. For $\mu \notin \Gamma_0$ by Theorem 4.3 parts (1), (2) and (3) one can compute the bigraded Hilbert function and for blue points by part (4). For the orange points, Theorem 4.3 does not say anything. In addition, any points except the orange ones only depend on the degree of the generator of I , which in this case is $(2, 2)$.

Asking `random` command in Macaulay2 and constructing I' with 4 generators of bidegree $(2, 2)$. Let V' be the complete intersection scheme of points defined by I .

For $(0, 0) \leq \mu \leq (13, 7)$ the bigraded Hilbert function $\text{HF}_{V'}(\mu)$ is

33	84	96	96	96	96	96	96	96	96	96	96	96	96
27	81	96	96	96	96	96	96	96	96	96	96	96	96
21	63	86	90	93	95	96	96	96	96	96	96	96	96
15	45	66	78	87	93	96	96	96	96	96	96	96	96
10	30	48	64	78	90	96	96	96	96	96	96	96	96
6	18	32	48	66	86	96	96	96	96	96	96	96	96
3	9	18	30	45	63	81	84	96	96	96	96	96	96
1	3	6	10	15	21	28	36	45	55	66	78	91	96

The only differences are in the orange spots. In addition, in this case $\text{HF}_{V'}(0, 13) = \text{HF}_{V'}(13, 0) = 96$ which means the natural projection of V' to each \mathbb{P}^n and \mathbb{P}^m is one to one.

Theorem 4.5. *Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring where $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Assume V be a complete intersection scheme of points defined by $I = (f_1, \dots, f_{n+m})$ with $\deg(f_i) = (d, e)$. Let $\pi_1 : V \rightarrow \mathbb{P}^n$ and $\pi_2 : V \rightarrow \mathbb{P}^m$ be the natural projections. Then*

$$\begin{aligned} \max\{\text{reg}(\pi_1^{-1}(p)) \text{ for } p \in \mathbb{P}^n\} &= \min\{a \in \mathbb{N} \mid (a, b) \in \text{reg}_B(V) \text{ for some } b\} \\ \max\{\text{reg}(\pi_2^{-1}(p)) \text{ for } p \in \mathbb{P}^m\} &= \min\{b \in \mathbb{N} \mid (a, b) \in \text{reg}_B(V) \text{ for some } a\}. \end{aligned}$$

Proof. Adopt Notation 2.1 and let J be the saturation of I with respect to B . Since S/J is B -saturated, it is also B_1 and B_2 saturated. Let $\mu \gg 0$ and $\nu \gg 0$, therefore $H_{\mathfrak{m}}^1(S/J)_{\mu, \nu} = H_{\mathfrak{m}}^2(S/J)_{\mu, \nu} = 0$. Suppose $\mu \gg 0$, Hence by Mayer-Vietoris exact sequence

$$H_{B_1}^1(S/J)_{\mu, \nu} \oplus H_{B_2}^1(S/J)_{\mu, \nu} = H_B^1(S/J)_{\mu, \nu}.$$

Since S/J is B_1 -saturated, for $\mu \gg 0$, $H_{B_1}^1(S/J)_{\mu, \nu} = 0$ which means

$$H_{B_2}^1(S/J)_{\mu, \nu} = H_B^1(S/J)_{\mu, \nu}.$$

We state the proof for the first equality, the argument for the second one is the same. There exist $(a_1, b_1), (a_2, b_2) \in \mathbb{N}^2$ such that $H_{\mathfrak{m}}^1(S/J)_{\mu} = H_{\mathfrak{m}}^2(S/J)_{\nu} = 0$ for all $\mu \geq (a_1, b_1)$ and $\nu \geq (a_2, b_2)$. Also for any $a \in \mathbb{N}$, there exists b_a such that for all $\mu \geq b_a$, $(H_{B_2}^1(S/J))_{(a, \mu)} = 0$. Set

$$b := \max_{a \in \mathbb{N}} \{b_1, b_2, b_a\}.$$

The Mayer-Vietoris short exact sequence yields $H_B^1(S/J)_{(\mu, \nu)} = H_{B_1}^1(S/J)_{(\mu, \nu)}$ if $\nu > b$. Therefore

$$\begin{aligned} \min\{a \in \mathbb{N} \mid (a, b) \in \text{reg}_B(V) \text{ for some } b\} &= \min\{\mu \in \mathbb{N} \mid H_{B_1}^1(S/J)_{\mu, \nu} = 0 \text{ for } \nu > b\} \\ &= \min\{\mu \in \mathbb{N} \mid \exists \mathfrak{p} \neq (y_0, y_1, y_2); H_{B_1}^1(S/J \otimes_{k[y_0, y_1, y_2]} k[y_0, y_1, y_2]_{\mathfrak{p}})_{\mu, \nu} = 0 \text{ for } \nu > b\} \end{aligned}$$

□

5. GENERIC COMPLETE INTERSECTION POINTS IN $\mathbb{P}^n \times \mathbb{P}^m$

In this subsection we adopt the Notation 2.1 as well. We focus on the case where ideal I is generated by the generic forms. In this regard, we first need to provide an analogous version of Bertini's theorem in our case.

Proposition 5.1. *Let k be a field of characteristics zero and $V \subset \mathbb{P}_k^n \times \mathbb{P}_k^m$ be a reduced scheme and $(d, e) \in \mathbb{N}^2$ with $d, e \neq 0$. Let f_U be a form of bidegree (d, e) with indeterminate coefficients $U_{\alpha, \beta}$. There exists a non empty open set $\Omega \subset \text{Spec}(k[U_{\alpha, \beta}])$ such that for any $\mathfrak{p} \in \Omega$, the corresponding form $f_{\mathfrak{p}}$ is such that*

$$V \cap Z(f_{\mathfrak{p}}) \subseteq \mathbb{P}_{k_{\mathfrak{p}}}^n \times \mathbb{P}_{k_{\mathfrak{p}}}^m,$$

where $k_{\mathfrak{p}}$ is the residue field of $k[U_{\alpha, \beta}]_{\mathfrak{p}}$, is reduced of dimension equal $\dim V - 1$, unless the $\dim V = 0$ in which case $V \cap Z(f_{\mathfrak{p}}) = \emptyset$.

Proof. Consider the Segre-Veronese map $\Psi : \mathbb{P}^n \times \mathbb{P}^m \rightarrow \mathbb{P}^N$ with $N = \binom{n+d}{d} \times \binom{m+e}{e} - 1$. Let $V' = \Psi(V)$. Under the map Ψ , f_U is mapped to a linear form ℓ_U in $N + 1$ variables and this correspondence is one to one. By Bertini theorem (see [3, Corollary 3.4.9]) there exists Ω as claimed such that for $\mathfrak{p} \in \Omega$, $V' \cap Z(\ell_{\mathfrak{p}}) \cong V \cap Z(f_{\mathfrak{p}})$ is a reduced scheme with the asserted dimension. \square

Definition 5.2. *Forms of bidegree (d, e) are in one to one correspondence with closed points in $\mathbb{A}_k^{\dim_k S(d, e)}$.*

Remark 5.3. *Notice that $\Omega \subset \text{spec } k[U_{\alpha, \beta}] \setminus Z(F)$ for some form $F \neq 0$. In particular, for $c = (c_{\alpha, \beta}) \in k^N$ such that $F(c) \neq 0$, Proposition 5.1 with \mathfrak{p} generated by the elements $u_{\alpha, \beta} - c_{\alpha, \beta}$, shows that $Z(f_c) \cap V$ is reduced.*

Theorem 5.4. *Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring over a field k of characteristic zero where $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$. Let $(d_i, e_i) \geq (1, 1)$ for $i = 1, \dots, n + m$ be bidegrees. Then, there exists a non-empty open set*

$$\Omega \subset \mathbb{A}_k^N := \prod_{i=1}^{n+m} \mathbb{A}_k^{|S(d_i, e_i)|}$$

such that, for any (f_1, \dots, f_{n+m}) corresponding to a point in Ω ,

- (1) $Z := \text{proj}(S/(f_1, \dots, f_{n+m}))$ is reduced of dimension zero.
- (2) The natural projections of Z to the factors \mathbb{P}^n and \mathbb{P}^m are isomorphism.

Proof. Let $V := Z(g_1, \dots, g_{n+m}) \subset \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m$ where $\mathbb{A}_k^N = \text{spec } k[U_{i, \alpha, \beta}]$ for $1 \leq i \leq n + m$, $|\alpha| = d_i$, $|\beta| = e_i$ and $g_i := \sum U_{i, \alpha, \beta} \mathbf{x}^{\alpha} \mathbf{y}^{\beta}$. Set $\pi : \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m \rightarrow \mathbb{P}_k^n \times \mathbb{P}_k^m$ be the natural projection, then V is a vector bundle over $\mathbb{P}_k^n \times \mathbb{P}_k^m$ via π . For any point $p \in \mathbb{P}_k^n \times \mathbb{P}_k^m$ the fiber of p is a linear space of dimension $N - n - m$. Hence V is geometrically irreducible scheme and $\dim(V) = n + m + (N - n - m) = N$.

(1) Set $W := (g_1, \dots, g_{n+m}, \text{Jac}^{n+m}(g_1, \dots, g_{n+m})) \subset V$ where Jac^{n+m} is the Jacobian of order $n + m$. By using Proposition 5.1 inductively, there exist a point $u \in \mathbb{A}_k^N$ and (f_1, \dots, f_{n+m}) corresponding to u such that $\text{Jac}^{n+m}(f_1, \dots, f_{n+m}) \neq 0$ which yields $W \subsetneq V$ and $\dim(W) < \dim(V) = N$. Consider the natural projection $p : \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m \rightarrow \mathbb{A}_k^N$. Since $\dim(W) < N$, $p(W)$ is a non empty closed subset of \mathbb{A}_k^N . Set $\Omega_0 := (\mathbb{A}_k^N \setminus p(W))$. Therefore for any (f_1, \dots, f_{n+m}) corresponding to a point in Ω_0 , $\text{proj}(S/(f_1, \dots, f_{n+m}))$ is geometrically reduced and is of dimension

zero.

(2) Set $p_1 : \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m \rightarrow \mathbb{A}_k^N \times \mathbb{P}_k^n$ and $q_2 : \mathbb{A}_k^N \times \mathbb{P}_k^n \rightarrow \mathbb{A}_k^N$ the natural projections. First we show that it is enough to show that the restriction $p'_1 : V \rightarrow p_1(V)$ is birational. Indeed if p'_1 is birational then there exists a closed subset $Z_1 \subsetneq p_1(V)$ such that setting $W_1 := (p'_1)^{-1}(Z_1)$ then $p'_1|_{V \setminus W_1} : V \setminus W_1 \rightarrow p_1(V) \setminus Z_1$ is an isomorphism. As $\dim(Z_1) < N$, $q_2(Z_1) \subsetneq \mathbb{A}_k^N$. Set $\Omega_1 := \mathbb{A}_k^N \setminus (q_2(Z_1) \cup \pi(W))$. For any point $u \in \Omega_1$ and (f_1, \dots, f_{n+m}) corresponding to u ,

$$q_2^{-1}(u) = p_1(u, Z(f_1, \dots, f_{n+m})) \cong (u, Z(f_1, \dots, f_{n+m}))$$

showing that the natural projection of Z onto \mathbb{P}^n is an isomorphism. By replacing p_1 and q_2 by $p_2 : \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m \rightarrow \mathbb{A}_k^N \times \mathbb{P}_k^m$ and $q_1 : \mathbb{A}_k^N \times \mathbb{P}_k^m \rightarrow \mathbb{A}_k^N$, along the same lines it shows the existence of $Z_2 \in p_2(V)$ and $\Omega_2 = \mathbb{A}_k^N \setminus (q_1(Z_2) \cup \pi(W))$ such that for $u \in \Omega_2$ and (f_1, \dots, f_{n+m}) corresponding to u ,

$$q_1^{-1}(u) = p_2(u, Z(f_1, \dots, f_{n+m})) \cong (u, Z(f_1, \dots, f_{n+m})).$$

Consider the following diagram

$$\begin{array}{ccc} V & \xrightarrow{\subset} & \mathbb{A}_k^N \times \mathbb{P}_k^n \times \mathbb{P}_k^m \\ \downarrow p'_1 & & \downarrow p_1 \\ V_1 := p_1(V) & \xrightarrow{\subset} & \mathbb{A}_k^N \times \mathbb{P}_k^n \\ \downarrow q_2|_{V_1} & & \downarrow q_2 \\ \mathbb{A}_k^N & \xrightarrow{=} & \mathbb{A}_k^N. \end{array}$$

We now show that p'_1 is birational.

Let $I := (g_1, \dots, g_{n+m})$ and $J := I : B^\infty$. By part (1), V_1 is a geometrically irreducible scheme. Hence J and $I_{V_1} = J \cap k[\mathbf{x}, \mathbf{U}]$ are prime and remain prime under any extension of k .

As V_1 is generically smooth, there exists $G \in I_{V_1}$ such that $D := \frac{\partial G}{\partial U_{i,\alpha,\beta}} \notin I_{V_1}$ for some i, α , and β unless $G \in k[\mathbf{y}]$ which is impossible: indeed $I_{V_1} \cap k[\mathbf{y}] = (0)$ because the projection $V \rightarrow \mathbb{P}^m$ is onto.

Set $Z_1 := Z(D) \subset V_1$. By [7, Lemma 4.6.1],

$$\mathbf{x}^\alpha \mathbf{y}^\beta \frac{\partial G}{\partial U_{i,\alpha',\beta'}} - \mathbf{x}^{\alpha'} \mathbf{y}^{\beta'} \frac{\partial G}{\partial U_{i,\alpha,\beta}} \in I_V$$

for any choices of α' and β' . Let $\mathbf{y}^\beta = y_p \mathbf{y}^{\beta_0}$ for some i . For any $0 \leq j \leq m$ put $\alpha' := \alpha$ and $\beta' := y_j \beta_0$. As \mathbf{x}^α and \mathbf{y}^{β_0} are not in I_V it follows that

$$y_p \frac{\partial G}{\partial U_{i,\alpha',\beta'}} - y_j \frac{\partial G}{\partial U_{i,\alpha,\beta}} \in I_V$$

for all $0 \leq j \leq m$. Therefore by localizing at D , we get an isomorphism

$$\phi : \frac{(k[\mathbf{U}, \mathbf{x}]_D[\mathbf{y}])}{I_V} \xrightarrow{\cong} \frac{(k[\mathbf{U}, \mathbf{x}]_D[y_p])}{I_{V_1}}.$$

Therefore the natural maps

$$\left(\frac{k[\mathbf{U}, \mathbf{x}]}{I_{V_1}} \right)_D \xrightarrow{\iota} \frac{(k[\mathbf{U}, \mathbf{x}]_D[\mathbf{y}])}{I_V} \xrightarrow{\phi} \frac{(k[\mathbf{U}, \mathbf{x}]_D[y_p])}{I_{V_1}}$$

are such that $\phi \circ \iota$ induces the identity from $\text{proj}((k[\mathbf{U}, \mathbf{x}]_D[y_p])/I_{V_1})$ to $\text{spec}((k[\mathbf{U}, \mathbf{x}]_D)/I_{V_1})$. Hence ϕ provides the inverse φ

$$V_1 \setminus Z_1 \xrightarrow{\varphi} V \setminus (Z_1 \times \mathbb{P}_k^m) \xrightarrow{p'_1} V_1 \setminus Z_1.$$

□

Theorem 5.5. *Let $S = k[x_0, \dots, x_n, y_0, \dots, y_m]$ be a bigraded polynomial ring over a field k of characteristics zero. If I is generated by $n + m$ generic forms of bidegree (d, e) , then the scheme V defined by I is a set of reduced points and*

$$|\{\mu \in \mathbb{N}^2 \mid \text{HF}_V(\mu) \neq \text{deg}(V)\}| < \infty,$$

which means the natural projections are one-to-one.

Proof. By Theorem 5.4 part (1), V is a set of reduced points and by part (2), $\text{HF}_V(0, i) = \text{HF}_V(j, 0) = \text{deg}(V)$ for any $i, j \gg 0$. □

REFERENCES

- [1] BOTBOL, N., AND CHARDIN, M. Castelnuovo Mumford regularity with respect to multigraded ideals. *Journal of Algebra* 474 (mar 2017), 361–392.
- [2] EISENBUD, D., AND GOTO, S. Linear free resolutions and minimal multiplicity. *Journal of Algebra* 88, 1 (may 1984), 89–133.
- [3] FLENNER, H., O’CARROLL, L., AND VOGEL, W. *Joins and intersections*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1999.
- [4] FULTON, W. *Intersection theory*, second ed., vol. 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1998.
- [5] GRAYSON, D. R., AND STILLMAN, M. E. Macaulay2, a software system for research in algebraic geometry. Available at <https://faculty.math.illinois.edu/Macaulay2/>.
- [6] HOFFMAN, J. W., AND WANG, H. H. Castelnuovo–Mumford regularity in biprojective spaces. *Advances in Geometry* 4, 4 (jan 2004).
- [7] JOUANOLOU, J.-P. Le formalisme du résultant. *Adv. Math.* 90, 2 (1991), 117–263.
- [8] MACLAGAN, D., AND SMITH, G. G. Multigraded Castelnuovo–Mumford regularity. *J. Reine Angew. Math* 14, 1 (2005), 137–164.
- [9] PEEVA, I., AND STILLMAN, M. Open problems on syzygies and hilbert functions. *J. Commut. Algebra* 1, 1 (03 2009), 159–195.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, CNRS & SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

Email address: marc.chardin@imj-prg.fr

UNIVERSITÉ CÔTE D’AZUR, INRIA, 2004 ROUTE DES LUCIOLES, 06902 SOPHIA ANTIPOLIS, FRANCE

Email address: navid.nemati@inria.fr