



First mononuclear organometallics of MoII and MoIII containing terminal hydroxide ligands. X-Ray structure of $[\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{OH})(\text{PMe}_3)_3]\text{BF}_4$

James C Fettinger, Heinz-Bernhard Kraatz, Rinaldo Poli, E. Alessandra Quadrelli

► To cite this version:

James C Fettinger, Heinz-Bernhard Kraatz, Rinaldo Poli, E. Alessandra Quadrelli. First mononuclear organometallics of MoII and MoIII containing terminal hydroxide ligands. X-Ray structure of $[\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{OH})(\text{PMe}_3)_3]\text{BF}_4$. Chemical Communications, 1997, 9, pp.889-890. 10.1039/A701616F . hal-03370824

HAL Id: hal-03370824

<https://hal.science/hal-03370824>

Submitted on 8 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First mononuclear organometallics of Mo^{II} and Mo^{III} containing terminal hydroxide ligands. X-Ray structure of [Mo(η^5 -C₅H₅)(OH)(PMe₃)₃]BF₄

James C. Fetting^a, Heinz-Bernhard Kraatz,^b Rinaldo Poli^{a,c} and E. Alessandra Quadrelli^a

^a Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, USA


^b Steacie Institute for Molecular Science, National Research Council of Canada, 100 Sussex Drive, Ottawa, Canada, K1N 0R6

^c Laboratoire de Synthèse et d'Electrosynthèse Organometallique, Faculté des Sciences "Gabriel", Université de Bourgogne, 6 Boulevard Gabriel, 21100 Dijon, France

The new complexes [Mo(η^5 -C₅H₅)(OH)(PMe₃)₂] and [Mo(η^5 -C₅H₅)(OH)(PMe₃)₃]⁺ are unique examples of electronically unsaturated organometallics of Mo^{II} and Mo^{III} containing terminal hydroxide ligands; they show no tendency to dimerize with formation of hydroxo-bridges.

Hydroxide ligands are ubiquitous of Werner-type coordination chemistry, whereas they are less commonly found in organometallic compounds. The availability of lone pairs on the oxygen atom frequently leads a terminal hydroxide complex (**I**) to lose ligands and afford doubly or triply bridged oligomers, *e.g.* **II** and **III**.^{1–4} Organometallic compounds with terminal hydroxide ligands have rarely been observed for early- and late-transition-metal systems,^{5–10} and for low-oxidation state saturated (18-electron) systems with tightly bonded ancillary ligands, examples of which are [Mo(OH)(NO)(dppe)₂]¹¹ and [W(CO)₅(OH)][–].¹² We consider it therefore quite surprising to find the existence and stability toward dimerization of a mononuclear and paramagnetic organometallic Mo^{II} hydroxo complex with a 16-electron configuration. To the best of our knowledge, there are only two previous examples of formally 16-electron Mo^{II} complexes with terminal OH ligands, *i.e.* [MoL(NO)(O)(OH)] (L = 1,4,7-triisopropyl-1,4,7-triazacyclononane), and its corresponding protonated form, [MoL(NO)(OH)₂]⁺.¹³ These, however, do not contain M–C bonds and are diamagnetic, probably because extra π donation from the oxo lone pair(s) renders the compounds effectively saturated.

The reaction of [Mo(η^5 -C₅H₅)Cl(PMe₃)₃]¹⁴ with dry KOH in thf leads to the exchange of the chloride ion and the formation of paramagnetic [Mo(η^5 -C₅H₅)(OH)(PMe₃)₂] **1** [eqn. (1)].[†] The



same transformation occurs in C₆D₆, allowing the observation of 1 equiv. of free PMe₃ by a ¹H NMR monitoring experiment (Fig. 1). The ¹H NMR spectrum of **1** shows contact-shifted resonances at δ 21.8 and 20.3 in the correct ratio for the PMe₃ and C₅H₅ protons. The ³¹P NMR spectrum shows only the resonance of free PMe₃. The compound is extremely soluble in saturated hydrocarbons, thus preventing its isolation in crystalline form. The formulation of the complex as mononuclear rather than dinuclear is based on a comparison with literature analogues: [Mo(η^5 -C₅Me₅)ClL₂] (L = PMe₃, PMe₂Ph or L₂ = dppe) and [Mo(η^5 -C₅H₅)Cl(PMe₂Ph)₂] are also mononuclear paramagnetic (*S* = 1) complexes,¹⁵ and show contact-

shifted ¹H NMR resonances similar to those of **1**. On the other hand, [Mo(η^5 -C₅H₅)(μ -Br)(η^4 -C₄H₆)]₂ is dinuclear and diamagnetic.¹⁶ Mononuclear 16-electron [Mo(η^5 -C₅H₅)XL₂] complexes also exist with a spin-singlet ground state,¹⁷ whereas there are no paramagnetic 18-electron complexes in this class. A bis(μ -hydroxo) dimeric structure for **1** is therefore ruled out.

It is rather peculiar that **1** is stable not only toward dimerization, but also toward coordination of the PMe₃ ligand [eqn. (1)]. The corresponding chloro parent compound does not have any tendency to lose the third PMe₃. However, sterically more encumbered [Mo(η^5 -C₅Me₅)Cl(PMe₃)₃] and [Mo(η^5 -C₅H₅)Cl(PMe₂Ph)₃] exist in equilibrium with the corresponding 16-electron bis-phosphine complexes.¹⁵ This behaviour is in line with the greater π -donating ability of OH *vs.* Cl. The paramagnetism of **1** parallels that of the chloro analogues, but is in contrast with the diamagnetism of the phosphido or arsenido derivatives [Mo(η^5 -C₅Me₅)(PPh₂)L₂]¹⁷ and [Mo(η^5 -C₅H₅)(As-Bu_t)₂(CO)₂]¹⁸. We have previously rationalized this phenomenon in terms of differences in MO interactions (single-sided *vs.* double-sided π donors) and pairing energy stabilization.^{15,19} Elaborating on those arguments, the paramagnetism of **1** might suggest the involvement of both oxygen lone pairs in two orthogonal two-centre–three electron interactions.²⁰

Further treatment of the **1**–PMe₃ solution in thf with [Fe(η^5 -C₅H₅)₂]BF₄ leads to at least two products. Crystallization from acetone–heptane affords yellow needles of [Mo(η^5 -C₅H₅)(OH)(PMe₃)₃]BF₄ **2** and red cubes of [Mo(η^5 -C₅H₅)O(PMe₃)₂]BF₄ **3**. The structure of **2** (Fig. 2)[‡] shows a terminal mode for the hydroxide ligand in the expected four-legged piano-stool geometry, as found¹⁴ for the related [Mo(η^5 -C₅H₅)Cl(PMe₃)₃]⁺. The structural parameters are normal for this class of compounds,²¹ with the Mo–O bond measuring 2.084(2) Å. To the best of our knowledge, **2** is the first reported organometallic hydroxide of Mo^{III}, and the first complex of Mo^{III} (including Werner-type and clusters) with a terminally bonded hydroxide.^{22–25} A deprotonated form of the Mo^{III} aquo complex [Mo(H₂O)₆]³⁺ is known, but this corresponds to a μ -hydroxo dimer, *i.e.* [Mo(H₂O)₄(μ -OH)]₂⁴⁺.²⁶ **3** has been previously reported.^{27§}

The formation of **2** by oxidation of **1** occurs according to the stoichiometry of eqn. (2). Half-sandwich Mo^{III} complexes are





Fig. 1 Room-temperature ¹H NMR spectrum of the products of reaction (1); solvent C₆D₆






Fig. 2 Top view of the cation in compound **2**. Ellipsoids drawn at 30% probability. Selected bond lengths (Å) and angles (°): Mo–P(1) 2.4530(12), Mo–P(2) 2.4984(12), Mo–P(3) 2.4741(11), Mo–O(1) 2.084(2), Mo–Cp 1.980(4); P(1)–Mo–P(2) 94.19(5), P(1)–Mo–P(3) 93.21(4), P(1)–Mo–O(1) 139.27(10), P(1)–Mo–Cp 108.5(1), P(2)–Mo–P(3) 132.81(4), P(2)–Mo–O(1) 71.53(8), P(2)–Mo–Cp 110.7(1), P(3)–Mo–O(1) 72.75(8), P(3)–Mo–Cp 110.8(1), O(1)–Mo–Cp 112.2(2). Cp is the C₅H₅ ring centroid.

known to prefer the 17-electron configuration.^{21,28} The nature of the observed product **2** shows that a bis(μ -hydroxo)-bridged [**1**₂]²⁺ complex would be unstable relative to the mononuclear 17-electron PMe₃ adduct. The hydroxide proton in complex **2** can be reasonably expected to be acidic. Presumably, the presence of free PMe₃ or other adventitious proton acceptors during the formation of **2** induces a deprotonation equilibrium to afford a neutral Mo^{III} oxo species which is then easily oxidized by [Fe(η^5 -C₅H₅)₂]⁺ to afford the observed product **3** [eqn. (3)].



In conclusion, we have reported here for the first time the existence and stability of terminal hydroxo complexes of paramagnetic organometallics of Mo^{II} and Mo^{III} in defiance of the possible dimerization with formation of hydroxo-bridges. For the dimerization process to occur, **2** would have to lose the stabilization of the Mo^{III}–PMe₃ bond, whereas compound **1** would have to pay the price of pairing the electrons. Terminal hydroxo complexes of Mo^{IV}, on the other hand, have been previously reported.^{2,29–34}

This research was supported by the US Department of Energy Office of Energy Research (grant DEFG0295ER14550) and the National Science Foundation (grant CHE-9508521).

Footnotes

* E-mail: poli@u-bourgogne.fr

† By carrying out all operations under a dinitrogen atmosphere, solid KOH (pre-dried at 110 °C and 6 mmHg for 48 h; ca. 300 mg, ca. 5.3 mmol) was added to a blue solution of [Mo(η^5 -C₅H₅)Cl(PMe₃)₃] (215 mg, 0.507 mmol) in 6 ml of thf. After stirring overnight at room temp., complete evaporation of the solvent from the yellow–brown solution yielded an oily residue, which was extracted into heptane (3 × 5 ml). An aliquot of this solution was dried under reduced pressure and the residue was investigated by ¹H NMR in C₆D₆: δ 21.8 (br, *w*_{1/2} 90 Hz, 18 H, Me), 20.3 (br, *w*_{1/2} 75 Hz, 5 H, C₅H₅). Concentration of the heptane solution and cooling to –80 °C did not separate any solid.

‡ Crystal data for **2**: C₁₄H₃₃BF₄MoOP₃, *M* = 493.06, orthorhombic, space group *Pna*2₁, *a* = 20.525(5), *b* = 13.380(4), *c* = 7.9740(7) Å, *U* = 2189.9(9) Å³, *Z* = 4, *D*_c = 1.496 Mg m^{–3}, $\mu(\text{Mo-K}\alpha)$ = 0.851 mm^{–1},

$2\theta_{\text{max}}$ = 49.96°, ω – 2θ scan mode, λ = 0.71073 Å, *T* = 293(2) K, *R*₁ = 0.0353 (*wR*₂ = 0.0734) for all data, *R*₁ = 0.0267 (*wR*₂ = 0.0567) for 226 parameters and 3339 observed [*I* > 2σ(*I*)] data (3712 collected and independent reflections). Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre (CCDC). See Information for Authors, Issue No. 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 182/429.

§ The ¹H NMR properties match with those reported.²⁷ ³¹P NMR (CD₃COCD₃): 13.8. The identity of this compound is also verified by an X-ray crystallographic study, which will be reported separately.

References

- M. E. Falkley and A. Pidcock, *J. Chem. Soc., Dalton Trans.*, 1977, 1444.
- V. Albano, P. Bellon, G. Cini and M. Manassero, *Chem. Commun.*, 1969, 1242.
- U. Sartorelli, L. Garlaschelli, G. Ciani and G. Bonora, *Inorg. Chim. Acta*, 1971, **5**, 191.
- M. Herberhold, G. Süß, J. Ellermann and H. Gäbelein, *Chem. Ber.*, 1978, **111**, 2931.
- G. L. Hillhouse and J. E. Bercaw, *J. Am. Chem. Soc.*, 1984, **106**, 5472.
- S. A. Al-Jibori, *J. Organomet. Chem.*, 1996, **506**, 119.
- V. V. Grushin and H. Alper, *Organometallics*, 1996, **15**, 5242.
- T. G. Appleton and M. A. Bennett, *Inorg. Chem.*, 1978, **17**, 738.
- D. P. Arnold and M. A. Bennett, *J. Organomet. Chem.*, 1980, **199**, 119.
- D. P. Arnold and M. A. Bennett, *Inorg. Chem.*, 1984, **23**, 2110.
- C. T. Kan, P. B. Hitchcock and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, 1982, 79.
- J. L. Cihonski and R. A. Levenson, *Inorg. Chem.*, 1975, **14**, 1717.
- J. Böhrer, G. Haselhorst, K. Wieghardt and B. Nuber, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1473.
- F. Abugideiri, M. A. Kelland, R. Poli and A. L. Rheingold, *Organometallics*, 1992, **11**, 1303.
- F. Abugideiri, J. C. Fettinger, D. W. Keogh and R. Poli, *Organometallics*, 1996, **15**, 4407.
- L.-S. Wang, J. C. Fettinger and R. Poli, *J. Am. Chem. Soc.*, in the press.
- R. T. Baker, J. C. Calabrese, R. L. Harlow and I. D. Williams, *Organometallics*, 1993, **12**, 830.
- M. Lukasz, S. Kimmer and W. Malisch, *Angew. Chem., Int. Ed. Engl.*, 1983, **22**, 416.
- F. Abugideiri, D. W. Keogh and R. Poli, *J. Chem. Soc., Chem. Commun.*, 1994, 2317.
- R. Poli, *Chem. Rev.*, 1996, **96**, 2135.
- R. Poli, *J. Coord. Chem. B*, 1993, **29**, 121.
- K. Wieghardt, M. Hahn, W. Swiridoff and J. Weiss, *Inorg. Chem.*, 1984, **23**, 94.
- K. Wieghardt, M. Guttman, P. Chaudhuri, W. Gebert, M. Minelli, C. G. Young and J. H. Enemark, *Inorg. Chem.*, 1985, **24**, 3151.
- A. G. Sykes, in *Comprehensive Coordination Chemistry*, ed. G. Wilkinson, R. D. Gillard and J. McCleverty, Pergamon, Oxford, 1987, ch. 36.1.
- K. Wieghardt, U. Bossek, A. Neves, B. Nuber and J. Weiss, *Inorg. Chem.*, 1989, **28**, 432.
- D. T. Richens and A. G. Sykes, *Inorg. Synth.*, 1985, **23**, 130.
- M. Brookhart, K. Cox, F. G. N. Cloke, J. C. Green, M. L. H. Green, P. M. Hare, J. Bashkin, A. E. Derome and P. D. Grebenik, *J. Chem. Soc., Dalton Trans.*, 1985, 423.
- I. Cacelli, D. W. Keogh, R. Poli and A. Rizzo, *New J. Chem.*, 1997, **21**, 133.
- K. Prout, T. S. Cameron, R. A. Forder, S. R. Critchley, B. Denton and G. V. Rees, *Acta Crystallogr., Sect. B*, 1974, **30**, 2290.
- P. R. Robinson, E. O. Schlemper and R. K. Murmann, *Inorg. Chem.*, 1975, **14**, 2035.
- M. R. Churchill and F. J. Rotella, *Inorg. Chem.*, 1978, **17**, 668.
- M. Berry, N. J. Cooper, M. L. H. Green and S. J. Simpson, *J. Chem. Soc., Dalton Trans.*, 1980, 29.
- K. Wieghardt, G. Backes-Dahman, W. Holzbach, W. J. Swiridoff and J. Weiss, *Z. Anorg. Allg. Chem.*, 1983, **499**, 44.
- T. Adachi, D. L. Hughes, S. K. Ibrahim, S. Okamoto, C. J. Pickett, N. Yabanouchi and T. Yoshida, *J. Chem. Soc., Chem. Commun.*, 1995, 1081.