



H, ^{15}N and ^{13}C resonance assignments of the Cterminal domain of Pull, a component of the *Klebsiella oxytoca* type II secretion system

Régine Dazzoni, Aracelys López-Castilla, Florence Cordier, Benjamin Bardiaux,
Michael Nilges, Olivera Francetic, Nadia Izadi-Pruneyre

► To cite this version:

Régine Dazzoni, Aracelys López-Castilla, Florence Cordier, Benjamin Bardiaux, Michael Nilges, et al.. H, ^{15}N and ^{13}C resonance assignments of the Cterminal domain of Pull, a component of the *Klebsiella oxytoca* type II secretion system. 2021. <hal-03371648>

HAL Id: hal-03371648

<https://hal.science/hal-03371648v1>

Preprint submitted on 8 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

^1H , ^{15}N and ^{13}C resonance assignments of the C-terminal domain of Pull, a component of the *Klebsiella oxytoca* type II secretion system

Régine Dazzoni¹, Aracelys López-Castilla¹, Florence Cordier^{1,2}, Benjamin Bardiaux¹, Michael Nilges¹, Olivera Francetic³, Nadia Izadi-Pruneyre^{1*}

¹Structural Bioinformatics Unit, Department of Structural Biology and Chemistry, CNRS UMR3528, Institut Pasteur, 28 rue du Dr Roux, 75724 Paris, France.

²Biological NMR Technological Platform, Institut Pasteur, Department of Structural Biology and Chemistry, CNRS UMR3528, 28 rue du Dr Roux, 75724 Paris, France

³Biochemistry of Macromolecular Interactions Unit, Department of Structural Biology and Chemistry, Institut Pasteur, CNRS UMR3528, 28 rue du Dr Roux, 75724 Paris, France.

* corresponding author: nadia.izadi@pasteur.fr

Keywords: NMR resonance assignment, Type II Secretion System, Pull, *Klebsiella oxytoca*.

Declarations

Keywords: NMR resonance assignment, Type II Secretion System, Pull, *Klebsiella oxytoca*.

Funding

This work was funded by the Institut Pasteur, the Centre National de la Recherche Scientifique (CNRS), the French Agence Nationale de la Recherche (ANR Synergy-T2SS ANR-19-CE11-0020-01), the Fondation pour la Recherche Médicale (Equipe FRM 2017M.DEQ20170839114 to M.N).

Conflicts of interest/Competing interests

The authors declare they have no conflict of interest.

Availability of data and material (data transparency)

The chemical shift values have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu/>) under the accession number 50966.

Abstract

Type II secretion systems (T2SS) allow Gram-negative bacteria to transport toxins and enzymes from the periplasm to the external milieu, and are thus important for the pathogenicity of bacteria. To drive secretion, T2SS assemble filaments called pseudopili closely related to bacterial type IV pili. These filaments are non-covalent polymers of proteins that are assembled by an inner membrane complex called the assembly platform connected to a cytoplasmic ATPase motor. In the *Klebsiella oxytoca* T2SS, the PulL protein from the assembly platform is essential for pseudopilus assembly and protein secretion. However, its role in these processes is not well understood. To decipher the molecular basis of PulL function, we used solution NMR to study its structure and interactions with other components of the machinery. Here as a first step, we report the ^1H , ^{15}N and ^{13}C backbone and side-chain chemical shift assignments of the C-terminal periplasmic domain of PulL and its secondary structure based on NMR data.

Biological context

In Gram-negative bacteria, several complex machineries are dedicated to the secretion of proteins involved in important adaptive processes such as nutrient acquisition, motility, adhesion, respiration and virulence (Nivaskumar and Francetic, 2014). The type II secretion system (T2SS) is one of the most widespread and relevant from biomedical and environmental standpoints (Cianciotto and White, 2017). Together with the closely related type IV pili, T2SSs belong to the versatile superfamily of type IV filament assembly systems (Berry and Pelicic, 2015, Denise *et al.*, 2019). The T2SS is composed of 12-15 proteins organized in four subcomplexes: 1) an outer membrane secretin channel (Hardie *et al.*, 1996); 2) a periplasmic pseudopilus consisting of a non-covalent polymer of proteins (Sauvonnet *et al.*, 2000, López-Castilla *et al.*, 2017); 3) an inner membrane assembly platform (Py *et al.*, 2001); and 4) a cytosolic ATPase (Possot *et al.*, 1997, Camberg and Sandkvist, 2005, Patrick *et al.*, 2011). While the secretin channel allows passage of folded proteins across the outer membrane, the inner membrane subcomplex couples the energy of the ATPase motor to the pseudopilus polymerization, which is thought to drive protein transport (Nivaskumar and Francetic, 2014). Despite decades of studies, the molecular basis of this process remains elusive.

In this study, we focused on the T2SS from *Klebsiella oxytoca* that secretes pullulanase, the enzyme which degrades branched maltotriose polymers allowing their uptake and use as a nutrient (d'Enfert *et al.*, 1987). Pullulanase secretion requires pseudopilus assembly, which is orchestrated by the assembly platform proteins PulF, PulL and PulM and hexameric ATPase PulE (Possot *et al.*, 2000).

PulL protein comprises 398 residues forming an N-terminal cytoplasmic domain, a transmembrane helix and a C-terminal periplasmic domain. In the T2SS of *Vibrio cholerae*, the cytoplasmic N-terminal domain of the PulL homologue EpsL forms a stable complex with the N-terminal subdomain of the ATPase (Abendroth *et al.*, 2005). This interaction anchors the ATPase hexamer to the assembly platform complex. The C-terminal domain (CTD) of PulL interacts with the C-terminal domain of PulM in the periplasm (Possot *et al.*, 2000). In several bacteria these domains have been shown to be each dimeric in solution (Abendroth *et al.*, 2004a, Abendroth *et*

al., 2004b, Fulara *et al.*, 2018) and to interact together as a heterodimer in bacterial two-hybrid and cysteine crosslinking assays (Lallemand *et al.*, 2013).

We report here the backbone and side-chain chemical shift assignments and secondary structure of the C-terminal domain of Pull. This assignment is the first necessary step towards the study of Pull structure and its dimerization. It will be further required for interaction studies with its partners to elucidate the role of Pull in the pseudopilus assembly mechanism. To our knowledge, this is the first NMR study of a protein from the T2SS assembly platform.

Methods and experiments

Production and purification of isotope-labelled Pull_{CTD}

The pMalp2 vector was used for the expression of the C-terminal domain of Pull (Pull_{CTD}) containing 88 residues (residues 312 to 398 of Pull with an additional N-terminal serine). This vector directs production of a N-terminal Maltose Binding Protein (MBP)-tagged protein in the periplasm, followed by a His₆-tag and a TEV cleavage site. Uniformly ¹⁵N labelled and ¹⁵N/¹³C double-labelled Pull_{CTD} were produced in M9 minimal medium by using 1 g/L of ¹⁵NH₄Cl and 4 g/L ¹³C-glucose, as the sole nitrogen and carbon sources, respectively. Gene expression was induced with 1 mM IPTG overnight at 18 °C in *E. coli* BL21 (DE3) cells. The MBP-His₆-tag-Pull_{CTD} labelled protein was purified from the supernatant after the sonication of the whole cells and centrifugation at 16000g, during 1 hour at 4°C. The purification was performed in three steps. The supernatant fraction was loaded onto a HisTrap HP column (*Cytiva*) previously equilibrated with 50 mM Tris-HCl, pH 8, 100 mM NaCl, 10 mM imidazole. Bound proteins were eluted with a linear imidazole gradient going from 10 to 500 mM. After a TEV cleavage overnight at 10°C, proteins were loaded on HisTrap HP column (*Cytiva*) and the flow-through corresponding to Pull_{CTD} without the tag was recovered and concentrated on a 3-kDa cutoff centricon device (*Cytiva*). The concentrated protein fraction was applied on a Sephacryl S-100 column (*Cytiva*) equilibrated with 50 mM HEPES, pH 7, 50 mM NaCl. Fractions containing the purified Pull_{CTD} were pooled and concentrated to 0.45 mM with pH adjusted to 6.5 for NMR data acquisition. Protease inhibitor cocktail, EDTA-free (*Roche*) was added to all buffers during the purification.

NMR spectroscopy

NMR experiments were recorded at 25°C on a 600 MHz Avance III HD (*Bruker Biospin*) spectrometer equipped with a cryogenically cooled triple resonance ¹H [¹³C / ¹⁵N] probe. The ¹H, ¹⁵N and ¹³C backbone and side chain resonance assignments (Cavanagh *et al.*, 1996) was carried out by using standard NMR experiments as implemented in TopSpin 3.6.1 (*Bruker Biospin*) and IBS libraries (Favier and Brutscher, 2019): Backbone and side chain resonances assignment was carried out by using standard NMR experiments: 2D ¹H-¹⁵N-HSQC, ¹H-¹³C-HSQC, and 3D HNCA, HN(CO)CA, HNCACB, CBCA(CO)HN, HNCO, HN(CA)CO, HCCH-TOCSY, C(CO)NH-TOCSY and H(CCO)NH-TOCSY. 2,2-Dimethyl-2-silapentane-5-sulfonate (DSS) signal was taken as 0 ppm for referencing proton chemical shifts and ¹⁵N and ¹³C chemical shifts were indirectly referenced to DSS (Wishart *et al.*, 1995). CcpNmr Analysis (Vranken *et al.*, 2005) was used for NMR data analysis. Analysis of the secondary structure was performed using HN, Hα, Cα, Cβ, CO, and N chemical shifts with the TALOS-N

Extent of assignments and data deposition

4

Secondary structure

Secondary structure elements of Pull_{CTD} were determined based on resonance assignments by calculating the deviation of C α , C β and C' chemical shifts from the corresponding random coil chemical shifts ($\Delta\delta C\alpha$, $\Delta\delta C\beta$ and $\Delta\delta C'$, Figure 2A). Residues with positive or negative values on $\Delta\delta C\alpha$ and $\Delta\delta C'$ are classified as helical or β -strand structures, respectively. For $\Delta\delta C\beta$, positive and negative values correspond to β -sheet and α -helix, respectively (Figure 2A). To complete these data, a TALOS-N analysis was also performed using the HN, H α , C α , C β , C' and N chemical shifts (Shen and Bax, 2013). As shown in Figure 2B, the TALOS prediction agrees with the secondary chemical shifts results. Both predictive approaches revealed that Pull_{CTD} presents two helical segments (α 1: I320-D331 and α 2: S359-A369) and 4 β -strands (β 1: G335-S342, β 2: L350-A356, β 3: F373-K381 and β 4: G386-G395), distributed in the α 1- β 1- β 2- α 2- β 3- β 4 sequential order (Figure 2).




Figure 2: Pull_{CTD} secondary structure predictions based on the NMR assignment data. **A.** Secondary chemical shifts from C α , C β , C'. The same scale (grey line = 2 ppm) is used for the three plots. **B.** TALOS-N secondary structure probability of each residue to be involved in an α -helix (black bars) or in a β -strand (grey bars). The Pull_{CTD} secondary structure, as a consensus of both predictive approaches, is represented between the panels. α -helices and β -strands are displayed as cylinders and arrows, respectively. The sequence of Pull_{CTD} is shown on top.

We observed significant differences in the peak intensities in the ^1H - ^{15}N -HSQC spectrum. Some signals are very weak leading also to weak peaks or absence of peaks in the 3D experiments. As shown in Figure 3, two regions ($\alpha 1$, $\beta 1$ - $\beta 2$) of the protein displayed significantly weaker signal intensity compared to the others. This spectral behaviour can be due to conformational exchange of the residues in these regions. Since there is evidence that Pull_{CTD} and its homologs are dimeric in solution (Fulara *et al.*, 2018, Abendroth *et al.*, 2004a), the regions with low signal intensity might be at the dimer interface or affected by the dimerization. Also, in the HSQC spectrum, no peaks corresponded to the residues 320 and 346, suggesting that these residues can be at the dimer interface. Determination of the 3D structure will be performed as a subsequent step to validate this hypothesis.




Figure 3: ^1H - ^{15}N -HSQC peak intensity (arbitrary unit a.u.) as a function of residue numbers. The two regions with lower peak intensity than the average (grey line) are highlighted with grey boxes.

Acknowledgements:

This work was funded by the Institut Pasteur, the Centre National de la Recherche Scientifique (CNRS), the French Agence Nationale de la Recherche (ANR Synergy-T2SS ANR-19-CE11-0020-01), the Fondation pour la Recherche Médicale (Equipe FRM 2017M.DEQ20170839114). We thank Rémy Le Meur and Bruno Vitorge for their help in NMR experiments.

References:

- ABENDROTH, J., BAGDASARIAN, M., SANDKVIST, M. & HOL, W. G. 2004a. The structure of the cytoplasmic domain of EpsL, an inner membrane component of the type II secretion system of *Vibrio cholerae*: an unusual member of the actin-like ATPase superfamily. *J Mol Biol*, 344, 619-33.
- ABENDROTH, J., MURPHY, P., SANDKVIST, M., BAGDASARIAN, M. & HOL, W. 2005. The X-ray structure of the type II secretion system complex formed by the N-terminal domain of EpsE and the cytoplasmic domain of EpsL of *Vibrio cholerae*. *Journal of molecular biology*, 348.
- ABENDROTH, J., RICE, A. E., MCLUSKEY, K., BAGDASARIAN, M. & HOL, W. G. 2004b. The crystal structure of the periplasmic domain of the type II secretion system protein EpsM from *Vibrio cholerae*: the simplest version of the ferredoxin fold. *J Mol Biol*, 338, 585-96.
- BERRY, J. & PELICIC, V. 2015. Exceptionally widespread nanomachines composed of type IV pilins: the prokaryotic Swiss Army knives. *FEMS microbiology reviews*, 39.

- CAMBERG, J. L. & SANDKVIST, M. 2005. Molecular analysis of the *Vibrio cholerae* type II secretion ATPase EpsE. *Journal of bacteriology*, 187.
- CAVANAGH, J., FAIRBROTHER, W. J., PALMER III, A. G. & SKELTON, N. J. 1996. *Protein NMR Spectroscopy: Principles and Practice*, New York, NY, U.S.A.
- CIANCIOOTTO, N. P. & WHITE, R. C. 2017. Expanding Role of Type II Secretion in Bacterial Pathogenesis and Beyond. *Infection and immunity*, 85.
- D'ENFERT, C., RYTER, A. & PUGSLEY, A. P. 1987. Cloning and expression in *Escherichia coli* of the *Klebsiella pneumoniae* genes for production, surface localization and secretion of the lipoprotein pullulanase. *The EMBO journal*, 6.
- DENISE, R., ABBY, S. & ROCHA, E. 2019. Diversification of the type IV filament superfamily into machines for adhesion, protein secretion, DNA uptake, and motility. *PLoS biology*, 17.
- FAVIER, A. & BRUTSCHER, B. 2019. NMRlib: user-friendly pulse sequence tools for Bruker NMR spectrometers. *Journal of biomolecular NMR*, 73.
- FULARA, A., VANDENBERGHE, I., READ, R. J., DEVREESE, B. & SAVVIDES, S. N. 2018. Structure and oligomerization of the periplasmic domain of GspL from the type II secretion system of *Pseudomonas aeruginosa*. *Sci Rep*, 8, 16760.
- HARDIE, K. R., LORY, S. & PUGSLEY, A. P. 1996. Insertion of an outer membrane protein in *Escherichia coli* requires a chaperone-like protein. *The EMBO journal*, 15.
- LALLEMAND, M., LOGIN, F. H., GUSCHINSKAYA, N., PINEAU, C., EFFANTIN, G., ROBERT, X. & SHEVCHIK, V. E. 2013. Dynamic interplay between the periplasmic and transmembrane domains of GspL and GspM in the type II secretion system. *PLoS One*, 8, e79562.
- LÓPEZ-CASTILLA, A., THOMASSIN, J., BARDIAUX, B., ZHENG, W., NIVASKUMAR, M., X, Y., NILGES, M., EGELMAN, E., IZADI-PRUNEYRE, N. & FRANCETIC, O. 2017. Structure of the calcium-dependent type 2 secretion pseudopilus. *Nature microbiology*, 2.
- NIVASKUMAR, M. & FRANCETIC, O. 2014. Type II secretion system: a magic beanstalk or a protein escalator. *Biochim Biophys Acta*, 1843, 1568-77.
- PATRICK, M., KOROTKOV, K. V., HOL, W. G. & SANDKVIST, M. 2011. Oligomerization of EpsE coordinates residues from multiple subunits to facilitate ATPase activity. *The Journal of biological chemistry*, 286.
- POSSOT, O. M., LETELLIER, L. & PUGSLEY, A. P. 1997. Energy requirement for pullulanase secretion by the main terminal branch of the general secretory pathway. *Molecular microbiology*, 24.
- POSSOT, O. M., VIGNON, G., BOMCHIL, N., EBEL, F. & PUGSLEY, A. P. 2000. Multiple interactions between pullulanase secretion components involved in stabilization and cytoplasmic membrane association of PulE. *J Bacteriol*, 182, 2142-52.
- PY, B., LOISEAU, L. & BARRAS, F. 2001. An inner membrane platform in the type II secretion machinery of Gram-negative bacteria. *EMBO reports*, 2.
- SAUVONNET, N., VIGNON, G., PUGSLEY, A. P. & GOUNON, P. 2000. Pilus formation and protein secretion by the same machinery in *Escherichia coli*. *The EMBO journal*, 19.
- SHEN, Y. & BAX, A. 2013. Protein backbone and sidechain torsion angles predicted from NMR chemical shifts using artificial neural networks. *Journal of biomolecular NMR*, 56.
- VRANKEN, W. F., BOUCHER, W., STEVENS, T. J., FOGH, R. H., PAJON, A., LLINAS, M., ULRICH, E. L., MARKLEY, J. L., IONIDES, J. & LAUE, E. D. 2005. The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins*, 59.
- WISHART, D., BIGAM, C., YAO, J., ABILDGAARD, F., DYSON, H., OLDFIELD, E., MARKLEY, J. & SYKES, B. 1995. ¹H, ¹³C and ¹⁵N chemical shift referencing in biomolecular NMR. *Journal of biomolecular NMR*, 6.