



HAL
open science

Body representation plasticity is altered in Developmental Coordination Disorder

Marie Martel, Véronique Boulenger, Eric Koun, Livio Finos, Alessandro Farnè,
Alice Catherine Roy

► To cite this version:

Marie Martel, Véronique Boulenger, Eric Koun, Livio Finos, Alessandro Farnè, et al.. Body representation plasticity is altered in Developmental Coordination Disorder. 2021. ⟨hal-03373678⟩

HAL Id: hal-03373678

<https://hal.science/hal-03373678v1>

Preprint submitted on 11 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Body representation plasticity is altered in Developmental Coordination Disorder

Marie Martel ^{1,2}, Véronique Boulenger ¹, Eric Koun ², Livio Finos ³, Alessandro Farnè ^{2,4,5}, Alice Catherine Roy ^{1,5}

¹ Laboratoire Dynamique Du Langage, UMR5596, CNRS/University Lyon 2, Lyon, France

² Integrative Multisensory Perception Action & Cognition Team - ImpAct, Lyon Neuroscience Research Center, INSERM U1028, CNRS U5292, University Lyon 1, Lyon, France

³ Department of Statistical Sciences, University of Padua, Italy

⁴ Hospices Civils de Lyon, Mouvement et Handicap, Neuro-immersion, Lyon, France

⁵ Center for Mind/Brain Sciences (CIMeC), University of Trento, Italy.

Corresponding author:

Marie Martel (present address)

mf.martel90@gmail.com

Department of Psychology

Royal Holloway University of London

Egham Hill, Surrey

TW20 0EX Egham (UK)

Abstract

Developmental Coordination Disorder (DCD) is a pathological condition characterized by impaired motor skills. Current theories advance that a deficit of the internal models is mainly responsible for DCD children's altered behavior. Yet, accurate movement execution requires not only correct movement planning, but also integration of sensory feedback into body representation for action to update the state of the body. Here we advance and test the hypothesis that the plasticity of body representations is altered in DCD. To probe Body Representations (BR) plasticity, we submitted a well-established tool-use paradigm to seventeen DCD children, required to reach for an object with their hand before and after tool use, and compared their movement kinematics to that of a control group of Typically Developing (TD) peers. We also asked both groups to provide explicit estimates of their arm length. Results revealed that DCD children explicitly judged their arm shorter after tool use, similarly to their TD peers. Unlike them, though, DCD did not update their implicit BR estimate: kinematics showed that tool-use affected their peak amplitudes, but not their latencies. Remarkably, the kinematics of tool-use showed that the motor control of the tool was comparable between groups, both improving with practice, confirming that motor learning abilities are preserved in DCD. These findings indicate that the update of the BR for action is impaired in DCD, suggesting a novel deficit may contribute to this pathology. Further studies are needed to tease apart the possible role played by motor and body representations deficits in DCD.

Keywords

Body representation, body schema, internal model, tool use, motor control, body image.

Introduction

Developmental Coordination Disorder (DCD) is a neurodevelopmental condition marked by impaired motor skills in the absence of neurological injury, given a child's chronological age and previous opportunities for skill acquisition (DSM-V: American Psychiatric Association, 2013). It affects approximately 5% of school-aged children (Mandich & Polatajko, 2003; range between 1.8% and 8% depending on the selection criteria; for reviews see Biotteau et al., 2020; Gomez & Sirigu, 2015; Zwicker et al., 2012), and persists through adulthood (e.g. Cantell et al., 2003; Cousins & Smyth, 2003; Losse et al., 1991; Rasmussen & Gillberg, 2000), thus considerably impacting academic and life achievements (e.g. Cheng et al., 2011; Geuze, 2005a; Kirby & Sugden, 2007; Smits-Engelsman et al., 2001; Van der Linde et al., 2015).

The etiology of DCD has gained considerable interest over the years, with converging evidence towards an internal modelling deficit (also called predictive control; Adams et al., 2014; Gomez & Sirigu, 2015; Wilson et al., 2013). Within internal models, the inverse model defines the accurate motor command, and the forward model predicts its consequences. Actual sensory feedback are also monitored allowing for movement correction and update of both models to improve motor control (Kawato & Wolpert, 1998; Shadmehr & Krakauer, 2008). Children with DCD can produce some appropriate motor commands when required, even though these appear to be more variable than in typically developing (TD) children (e.g. Roche et al., 2016; Smits-Engelsman et al., 2008; Smits-Engelsman & Wilson, 2013). Accordingly, visuomotor adaptation studies showed that children with DCD are able to update their inverse model, provided they are given more trials and larger error signals (Cantin et al., 2007; Kagerer et al., 2004, 2006; Zoia et al., 2005). This suggests that they need more time to process feedback, and that they might ignore error signals that are not relevant enough. When relevant, however, they can learn from these signals and update their models, as shown by their preserved ability for motor learning (Smits-Engelsman et al., 2015). Furthermore, children with DCD plan and execute a whole movement (i.e., in one time), rather than performing an incomplete movement and updating it online, as this may seem too costly for their motor system (Mon-williams et al., 2005). Hence, in double-step paradigms, children with DCD show difficulties in correcting 3D movement trajectories through rapid online control (Hyde & Wilson, 2011a, 2011b, 2013). Yet, they have no deficit when tasks are easier, namely when more time is allowed to correct the movement, or in tasks that involve 2D movements in the transversal plane (Adams, Lust, et al., 2016; Plumb et al., 2008). Online control performance can be predicted by the ability to represent action, that is motor imagery abilities, both in typical and atypical development (Fuelscher, Williams, Enticott, et al., 2015; Fuelscher, Williams, & Hyde, 2015). Motor imagery studies, which tackle the integrity of the forward model (Kiltner et al., 2018), revealed that DCD participants can imagine movements, but again less consistently, less accurately and less rapidly than their TD peers (e.g. Barhoun et al., 2019, 2021; Deconinck et al., 2009; Noten et al., 2014; Reynolds et al., 2015; Williams et al., 2008; Wilson et al., 2004). Children with DCD can make proper use of instructions to improve their motor imagery abilities (Reynolds et al., 2015; Williams et al., 2008), and motor imagery training has been shown helpful for motor control remediation in DCD (Steenbergen et al., 2020), again attesting that internal models can be updated. Overall, deficits in DCD therefore seem to affect different components of internal models, making it difficult to decipher the core neurocognitive deficit.

A so far relatively neglected aspect of the internal models is the body state estimation. This Body Representation (BR), also referred to as body schema, is defined as an implicit (i.e., unconscious) sensorimotor representation that allows monitoring the position and size of the different effectors (de Vignemont, 2010; Head & Holmes, 1911; Martel et al., 2016; Medina & Coslett, 2010; Schwoebel & Coslett, 2005). This representation it is often opposed to the so-called body image, an action-free,

explicit (i.e., conscious) representation of the body shape and size (de Vignemont, 2010; Gallagher, 2005). Substantial evidence for the plastic monitoring of limb's size in motor control comes from studies on tool use. After using a tool for a few minutes, healthy adults start performing free-hand movements differently, with longer latencies and reduced amplitudes for the acceleration, velocity and deceleration profiles (e.g. Cardinali et al., 2009; for review Martel et al., 2016). This kinematic signature of what has been called tool-incorporation, typically observed also in long-armed vs. short-armed participants (Cardinali et al., 2012; Martel et al., 2019), is indicative of a longer arm estimate after tool use. This body state plasticity has been suggested to allow tools to become extensions of our limbs for action and perception (e.g. Arbib et al., 2009; Jacobs et al., 2009; Maravita & Iriki, 2004; Martel et al., 2016; Miller et al., 2018, 2019; Witt, 2021).

Importantly, this plasticity requires years to develop fully. Martel and colleagues (2021) reported that the plasticity of the body state estimation is not mature in TD children and early adolescents. After using a tool for a few minutes, their free-hand kinematic pattern is actually opposite to what has been consistently observed in healthy adults: the kinematics peaks amplitudes of the wrist increased, and their latencies decreased. Following the rationale from previous studies in adults, this was interpreted as a movement performed with an arm estimated as being shorter (rather than longer in adults) after use of the same tool. Reduction in arm length estimate may result from the need to build new sensorimotor associations for a tool that has never been used before (Ganesh et al., 2014) or from a stronger reliance on visual as compared to proprioceptive guidance in TD children, compared to adults (Martel et al., 2021). Besides, investigating the plasticity of the body estimate in DCD may offer complementary insights to better understand its aetiology (Martel et al., 2017, 2019, 2021; Medendorp & Heed, 2019).

Surprisingly, the role potentially played by an altered body state estimation in DCD's motor deficits has yet been scarcely investigated (Gomez & Sirigu, 2015), although it has been linked indirectly with them. Impairment in multisensory body representations have been suggested after findings of poorer performance in somatosensory localization (Elbasan et al., 2012; Johnston et al., 2017; Schoemaker et al., 2001). Wilson and colleagues (2004) also hypothesized that motor imagery difficulties in DCD could result from inaccurate body estimate: children with DCD would not automatically use motor imagery from their perspective, but would rather rely on visual imagery from an object-based third-person perspective (see also Barhoun et al., 2021).

Here, we aimed at filling this gap by investigating the plasticity of both the implicit body estimate and the explicit body image in DCD, taking advantage of the above-mentioned tool-use paradigm. First, given their relatively preserved ability to update the inverse model and program movements (e.g. Kagerer et al., 2006; Smits-Engelsman et al., 2008, 2015), we predicted that children and early adolescents with DCD should be able to control the tool adequately and to improve with practice. Quantitative difference with their TD peers might however be expected as they sometimes ignore sensory signals, or need more trials to reach the same level. Second and most importantly, we predicted that children with DCD would show a less plastic body estimate than their TD peers. Accordingly, free-hand movements following tool use should display a different kinematic signature than that observed in TD children. Lastly, we also investigated the conscious, explicit representation (or body image) in DCD and TD, asking them to estimate their perceived forearm length before and after tool use. While, to the best of our knowledge, there is no previous investigation of the body image in DCD, we anticipated that it would be preserved (as it is not linked to motor control). Thus, their perceived forearm length should be shorter after tool use, as recently observed in TD children (Martel et al., 2021).

Material and methods

Participants

Children and adolescents with DCD were recruited during the school year, in parallel to the recruitment of TD participants from a previous study using the same paradigm (Martel et al., 2021). A total of 32 DCD participants, aged between 9.5 and 16.5 years old, were referred to our lab by several physical therapists as well as by the health network Dys/10 (specialized in the care of developmental disorders, from diagnosis with a multidisciplinary team to follow-ups with schools). All children were diagnosed with DCD following the DSM-5 (American Psychiatric Association, 2013), and accordingly had no neurological condition nor intellectual disability. They were additionally screened on the French adaptation of the Movement ABC (Henderson & Sugden, 1992; Soppelsa & Albaret, 2004) by a trained physical therapist. As the French version of the M-ABC2 (Marquet Doléac et al., 2016), standardized for adolescents up to 16 years of age, was not available at the time of the study, all the adolescents with DCD older than 12 years were evaluated based on the standardization for the 12-year-olds. To be included in the final sample, participants with DCD could not be born preterm ($n = 2$ excluded) and had to score below the 5th percentile on the M-ABC, or below the 15th percentile but with at least one subcategory under the 5th percentile ($n = 5$ excluded). Comorbidity with other developmental disorders was not an exclusion criterion (see Table 1).

Out of the 25 remaining DCD participants who fulfilled these criteria, we enrolled in the study children and early puberty adolescents only. We assessed their puberty level with the Self-Rating Scale for Pubertal Development (Carskadon & Acebo, 1993; Petersen et al., 1988) including five items standing for five phenomena: growth spurt, body hair, skin changes, deepening of the voice/breast growth and growth of hair on the face/menstruations. For each item, puberty-induced changes are rated on a 4-point scale (“not started yet”, “barely started”, “definitely underway”, “finished”), leading to a global puberty score (ranging between 5 and 20) which considers that children of the same age/sex have not necessarily reached the same pubertal stage. Adolescents who would indicate that each of the phenomena was “definitely underway” would score 15/20, meaning that they reached their puberty peak, while if all of the phenomena had “barely started”, they would score 10/20. We placed the cut-off for inclusion at 13/20 a posteriori, based on the results in typically developing children and adolescents using the same paradigm (Martel et al., 2021), which showed a change in tool-use effects on their body representations at mid-puberty (score ≥ 14). By including only the children and early adolescents with DCD scoring between 5 and 13/20, we minimized the potential role of puberty, and focused on a period where the pattern of results in TD participants is rather stable. This led to a final sample of 17 DCD participants, who were compared to a subsample of a larger cohort of TD participants from a previous study (Martel et al., 2021). The two groups were matched for daily tool practice, age, sex and puberty (see Table 2). TD children and early adolescents had no learning disabilities or delayed psychomotor acquisition according to their parents’ report. DCD and TD participants all had normal or corrected-to-normal vision. We also ensured that DCD and TD groups did not differ in their sport and/or musical practice or in their daily tool-use activities (i.e. activities involving tools such as tennis, ping pong, golf, drums etc. that are regularly practiced outside school). Non-tool users practiced activities without any tool involved (e.g., football, handball, climbing, piano...). Matching DCD and TD participants for tool-use activities ensures that any difference in performance between the groups could not result from different knowledge in such tool-based activities. The main characteristics of the two groups are displayed in Table 2.

Parents or guardians as well as children and early adolescents gave written informed consent to participate in the study, which was approved by the French ethics committee (*Comité de Protection des*

Personnes CPP Sud-Est II) and conformed to the Helsinki declaration. Children and early adolescents were naïve to the purpose of the study and received a board game after completion.

Gesture Imitation proficiency

We first quantified participants' sensorimotor proficiency and compared the performance of DCD and TD groups with a gesture imitation task, initially developed for the assessment of apraxia (De Renzi et al., 1980). Participants were instructed to imitate a set of different gestures performed by the experimenter using their dominant arm/hand (anatomical imitation, not mirrored), while standing. Two training gestures allowed them to familiarize with the task. We also informed them that some gestures would be repeated and emphasized the importance of exact imitation of the gesture (fingers opening, hand orientation etc.). We reminded participants of these instructions several times before the test. The test was composed of 24 gestures: for correct imitation on the first time, participants scored 3 points; 2 points were given if the experimenter had to perform the movement a second time and participants succeeded in imitating it. In case of failure, on the third and last repetition, participants scored 1 for a correct gesture and 0 for inaccurate imitation. We defined the assessment criteria classically used in motor imitation tasks (Rothi et al., 2014). These criteria included the configuration of the arm and/or hand (e.g. extended arm and fist configuration), the limb orientation in space (e.g. palm down) and its target location (e.g. palm on the contralateral shoulder). For sequential gestures, we additionally assessed the correct order and number of occurrences (e.g. three repetitions of fist and hand flat on the table sequence). Any element differing from one of these criteria was considered an incorrect imitation. The maximum score was 72. The same, trained experimenter demonstrated the gestures and evaluated online the imitation for all participants.

Table 1. Characteristics of the group of children and early adolescents with DCD (n=17). The scores on manual dexterity, ball skill and balance are the three subtests of the M-ABC. The scores $\leq 5^{\text{th}}$ percentile are indicated in bold, and those $\leq 15^{\text{th}}$ percentile in italics. Scores for the VCI (Verbal Comprehension Index), PRI (Perceptual Reasoning Index), WMI (Working Memory Index) and PSI (Processing Speed Index) are subtests of the IQ test (Grizzle, 2011). Comorbidity with other developmental disorders such as dyslexia, dyscalculia, developmental Language Disorder (DLD) or Attention Deficit Disorder with (ADHD) or without Hyperactivity (ADD) is indicated.

	Age (years)	Puberty Score (/20)	Sex (Female/Male)	Manual Laterality (Left/Right)	Manual	Ball	Balance	Total M-ABC	VCI	PRI	WMI	PSI	Comorbidity
DCD02	13.8	13	F	RH	11.5	8	1	20.5	103	84	100	78	DYSLEXIA DYSCALCULIA
DCD03	12.0	12	M	LH	13.5	6.5	9	29	152	88	94	69	ADHD DYSLEXIA
DCD07	13.7	12	M	RH	15	8	8	31	130	90	88	86	ADD
DCD09	16.2	13	M	RH	11	4	1	16	126	86	91	100	NONE
DCD10	9.7	5	M	RH	9	3	1.5	<i>13.5</i>	120	99	91	78	NONE
DCD11	14.5	12	M	RH	11	5	8	24	NA	NA	NA	NA	DLD
DCD12	12.6	11	M	RH	4.5	1.5	10	16	118	114	100	88	ADD DYSLEXIA
DCD13	12.1	8	F	RH	6	10	7	23	94	81	103	103	DYSLEXIA
DCD14	12.1	5	M	LH	10.5	9	8	27.5	NA	NA	NA	NA	ADD DYSCALCULIA
DCD15	11.5	10	M	RH	13	1.5	6	20.5	116	90	109	76	DYSLEXIA DYSCALCULIA
DCD16	13.1	7	M	LH	12.5	9	12	33.5	112	94	91	83	ADD DYSLEXIA
DCD18	12.7	8	M	RH	18	8	11	37	94	77	62	73	ADD DLD
DCD19	11.1	6	M	RH	15	0	5	20	79	86	94	83	DYSLEXIA
DCD20	10.6	9	F	RH	14	0	0	14	96	79	70	71	ADD DYSCALCULIA
DCD21	13.4	10	M	RH	11	0	0	11	84	99	94	90	DYSLEXIA
DCD23	15.8	10	F	LH	12	1	10	23	113	104	NA	NA	DYSLEXIA DYSCALCULIA
DCD24	13.5	13	M	RH	12.5	4	6	22.5	148	114	106	90	ADD

Table 2. Characteristics of the Typically Developing and the Developmental Coordination Disorder groups.

	n	Mean Age ± SD (year range)	Mean Puberty score ± SD (score range)	Mean Height ± SD (range in m)	Sex (Female/M ale)	Manual Laterality (Left/Right)	Practice of tool- based activities
TD	17	12.4 ± 1.8 (9.7 – 16.2)	9.3 ± 3.6 (5 – 13)	1.54 ± 0.12 (1.35 – 1.75)	4F / 13M	2LH / 15RH	4 Yes / 13 No
DCD	17	12.8 ± 1.7 (9.9 – 16.2)	9.8 ± 3.6 (5 – 13)	1.57 ± 0.12 (1.35 – 1.78)	4F / 13M	4LH / 13RH	4 Yes / 13 No
<i>t</i> -test		<i>t</i> (32) = .685 <i>p</i> = .498	<i>t</i> (32) = .591 <i>p</i> = .558	<i>t</i> (32) = .639 <i>p</i> = .528			

Apparatus and procedures

The paradigm was identical the one used in a previous study in TD children and adolescents (Martel et al., 2021). This paradigm has consistently been shown to be sensitive to changes in arm length implicit estimate after tool use (see, for review, Martel et al., 2016). In summary, participants were required to reach-to-grasp an object with their free dominant hand before and after performing reach-to-grasp movements towards the same object with a tool lengthening their arm. Comparison of the kinematics of free-hand movements before and after tool use allows to measure the effects of tool use on free-hand movements, and thus to assess the plasticity of the body representation for action (body schema). We also assessed the effect of tool use on the subjective estimate of the forearm length (body image) by asking participants to estimate the length of their forearm before and after performing similar reach-to-grasp movements with the same tool.

Participants were comfortably seated on an adjustable swivel chair, at a fixed distance from a table in front of them. The experiment was divided into three sessions: a pre- and post-tool-use session, separated by a tool use session. During both pre- and post-tool-use sessions, participants performed a Free-Hand Movement task, and a Forearm Length Estimation task, counterbalanced across participants.

Free-Hand Movement. The object to reach for was a wooden parallelepiped (10 × 2.5 × 5cm, weighting 96g) situated 35 cm away from a starting switch on the table, aligned with participants' dominant shoulder along the sagittal axis. Each trial (18 trials in total) started with participants holding their fingers in a pinch grip position on the starting switch. A tone signaled that they could start their movement. Participants were instructed to reach, grasp and lift the object at their natural speed, using their dominant hand, then to put the object down and go back to the starting position until the next tone was presented.

Forearm Length Estimation. In this task, participants were blindfolded. Each trial (18 trials in total) started with them holding their index finger on the starting switch, waiting for a tone to indicate the start of the estimation. Participants were asked to slide their dominant index finger horizontally (i.e., towards the right/left) on the surface of the table from the starting switch to a final position corresponding to the perceived estimation of their forearm (i.e. perceived distance between wrist and elbow). To avoid estimation bias by visual forearm measurement, the experimenter gave task instructions once the participants were blindfolded. The experimenter also named and touched the elbow and the wrist of each participant while giving the instructions, so that they would know exactly which body part they had to estimate.

Tool movement. The tool-use task was composed of four blocks of 12 trials. Each trial started with participants holding the tool “fingers” in a pinch grip position on the starting switch. After a tone indicating the go for the movement, participants had to reach, grasp and lift the same object as previously described at their natural speed, using the tool with their dominant hand. Between each trial, they had to put the object down and go back to their starting position until presentation of the next tone. Each participant underwent a short practice tool-use session at the beginning of the tool session. The tool, based on a commercial grabber (Unger Enterprise Inc, CT, USA), had an ergonomic handle fitted with a lever, a long rigid shaft, and a “hand” with two articulated fingers. It was customized and scaled on children’s height (see Martel et al., 2021 for details). Participants between 123 and 146 cm of height used a 32 cm long tool (DCD group: 4/17; TD group: 5/17). The remaining participants were taller than 147 cm and used a tool of 40 cm long, which is the original length also used for adults (Baccarini et al., 2014; Cardinali et al., 2009, 2011; Martel et al., 2019, 2021), but 100g lighter to prevent fatigue. Participants’ height was collected before the experiment, the lengths of their arm and forearm were measured afterwards.

Kinematic recording system

We recorded the spatial localization of the hand and of the tool using infrared light emitting diodes (IREDs) with an Optotrak 3020 (Northern Digital Inc; sampling rate: 200 Hz; 3D resolution: 0.01 mm at 2.25 m distance). Following previous studies using the same paradigm (Baccarini et al., 2014; Cardinali et al., 2009, 2011, 2012; Martel et al., 2019, 2021), we assessed the grasping component of the hand/tool movements by placing IREDs on the thumb and index finger nails of participants’ dominant hand, as well as on the two “fingers” of the tool. The reaching component was evaluated thanks to IRED located on the dominant wrist (styloid process of the radius and distal part of the tool shaft).

For each free-hand movement, we extracted and analyzed off-line several kinematic parameters with a custom-made Matlab program: latencies and amplitudes of wrist acceleration, wrist velocity and wrist deceleration peaks (reaching component), and latency and amplitude of the maximum thumb-index distance (hereafter MGA for Maximum Grip Aperture) (grasping component). We also measured the overall movement time as the time between the beginning of the movement (velocity ≥ 10 mm/s after switch release) and stabilized grasp on the object (before the lift). Extracted parameters were the same for the tool movement task, except that they involved kinematics of the tool instead of the hand. The arm length was estimated using the marker on participants’ index finger by subtracting the starting position to the final one.

Statistics

We used a Linear Model (LM) on the gesture imitation score of each participant, and Linear Mixed Models (LMM) on individual trials of each kinematic parameter of tool and free-hand movements, implemented in R (v3.6.1; RStudio v1.1.442; R Core Team, 2018) with the package lme4 (Bates et al., 2015). Although similar to LM in terms of interpretation, LMM are more adequate for the present study as they take into consideration the variability between participants (Boisgontier & Cheval, 2016), which is especially high in participants with developmental disorders. Statistics for main effects and interactions were extracted with the package car (Fox & Weisberg, 2011): we report these values based on Type III Wald chi-square tests rather than lme4 output. We used the package emmeans (Russell, 2019) for post-hoc tests, adjusting the *p-values* with the Tukey method for multiplicity adjustment.

Gesture Imitation task. We performed a LM with factor *Group* (TD/DCD) to assess whether DCD participants had poorer sensorimotor abilities than their TD peers, as expected from the literature.

Free-hand movement. We used a LMM with factors *Session* (PRE/POST) and *Group* (TD/DCD) and a random intercept for participants. We were particularly interested in the presence of interactions between the two factors, as they would indicate that tool use modifies free-hand movements of DCD children and early adolescents differently than it does in their TD peers. Additionally, we expected the pattern observed in our subsample of TD participants to conform to the one from the larger sample (Martel et al., 2021), namely that TD children and early adolescents display shorter latencies and higher amplitudes of acceleration, velocity and/or deceleration peaks after tool use.

Forearm Length Estimation. A LMM with factors *Session* (PRE/POST) and *Group* (TD/DCD) and a random intercept for participants was used (note that data were missing for one TD and one DCD participant). Interactions between the two factors would indicate that tool use does not affect the conscious forearm representation of DCD children and early adolescents the same way it does for TD participants. Additionally, for the latter, we expected the conscious forearm representation to be reduced after tool use, as previously reported in the larger sample of TD participants (Martel et al., 2021).

Tool movement. We used a LMM with factors *Block* (FIRST/LAST) and *Group* (TD/DCD) and a random intercept for participants. Note that the second and third blocks of tool use were not analyzed as we were interested in motor learning with practice. Would the two factors interact, this would show that both groups do not control the tool similarly. As before, we also anticipated to observe the motor learning effects found in the larger sample of TD participants (Martel et al., 2021), with children and early adolescents displaying shorter latencies and higher amplitudes in the last compared to the first block.

Results

DCD participants show poorer gesture imitation abilities than TD participants

The DCD group showed a mean score for gesture imitation of 49.2 ± 8.8 . As evidenced by a significant effect of *Group* in a linear model ($F(1, 32) = 10.34, p = .003$), this was significantly lower than the performance of the TD group who scored at 57.6 ± 6.2 (Figure 1). Noteworthy, when considering the cutoffs from the original test in adults (healthy ≥ 62 ; apraxia ≤ 53 ; De Renzi et al., 1980), the two groups did not fall in the same category, underlining their different sensorimotor abilities. Note that this test has been standardized on adults, lower performance are therefore to be expected in the present study (see also Martel et al., 2021), however this can not explain a change in categories between the groups.




Figure 1. Performance in the gesture imitation test in TD and DCD participants. The thick horizontal lines indicate the mean for each group ± 1 SEM. Single dots represent individual performance. Horizontal dashed lines represent the cutoffs in the original task standardized on adults (healthy ≥ 62 ; apraxia ≤ 53 ; De Renzi et al., 1980)

Tool use affects free-hand movements differently in TD and DCD children and early adolescents

Reaching component. The LMM revealed a significant main effect of *Session* on most of the latencies (all $p < .009$; except the acceleration: $\chi^2(1) = 3.48$, $p = .062$), and on all of the amplitudes (all $p < .042$). Of major interest here, significant interactions between *Session* and *Group* were observed for all the latencies (acceleration: $\chi^2(1) = 3.99$, $p = .046$; velocity: $\chi^2(1) = 9.04$, $p = .003$; deceleration: $\chi^2(1) = 11.0$, $p < .001$). Post-hoc comparisons showed that while the TD participants displayed decreased latencies for two out of three parameters when reaching after tool use (velocity: $t = 2.62$, $p = .044$; deceleration: $t = 3.46$, $p = .003$; Figure 2), no latency modulation in subsequent free-hand movements was found in the DCD children and early adolescents (velocity: $t = -1.63$, $p = .364$; deceleration: $t = -1.23$, $p = .611$). Both TD and DCD participants displayed increased amplitudes following tool use, as indicated by the lack of significant *Session* \times *Group* interactions (acceleration: $\chi^2(1) < .001$, $p = .928$; velocity: $\chi^2(1) = .444$, $p = .505$; deceleration: $\chi^2(1) = 3.21$, $p = .073$).




Figure 2. Kinematics of the Reaching component before (blue) and after (orange) tool use (top panel) and individual performance ordered by size (bottom panel) in TD and DCD participants for the latencies and amplitudes of velocity and deceleration. On the bottom panel, positive values indicate that the latency/amplitude increased after tool use, while negative values indicate reduced latency/amplitude. Most of the TD children and early adolescents displayed reduced latencies and increased amplitudes after tool use. DCD children and early adolescents displayed increased amplitudes but no modulation of the latencies. Error bars indicate the means \pm 1 SEM. Asterisks denote significance.

Grasping component. The factor *Session* significantly affected the latency of MGA ($\chi^2(1) = 13.8$, $p < .001$): children and early adolescents opened their fingers earlier after tool use. There was no evidence that DCD participants modulated this parameter differently than their TD peers, with neither a main effect of *Group* ($\chi^2(1) = .499$, $p = .480$), nor an interaction between *Session* and *Group* ($\chi^2(1) = .683$, $p = .409$). Regarding the MGA, the DCD participants opened their fingers larger than the TD participants with a main effect of *Group* ($\chi^2(1) = 9.92$, $p = .002$). There was also a significant main effect of *Session* ($\chi^2(1) = 4.03$, $p = .045$) and a significant *Session* \times *Group* interaction ($\chi^2(1) = 14.2$, $p < .001$). As revealed by post-hoc comparisons, tool use increased subsequent free-hand finger opening in the DCD group ($t = -7.12$, $p < .001$) while it did not modulate grip aperture in TD participants ($t = -2.01$, $p = .186$; Figure 3A).




Figure 3. Kinematics of the Grasping component (A) and the Movement Time (B) before (blue) and after (orange) tool use (top panel) and individual performance ordered by size (bottom panel) in TD and DCD participants. On the bottom panel, positive values indicate that the latency/grip aperture and movement time increased after tool use, while negative values indicate reduced latency/grip aperture and shorter movement time. Note that data for the grasping component is missing for one DCD participant. Most of the TD children and early adolescents displayed reduced MGA latencies, which led to reduced movement time after tool use. DCD children and early adolescents opened their fingers earlier and larger after tool use. They however did not display any modulation of their movement time. Error bars indicate the means \pm 1 SEM. Asterisks denote significance.

Movement Time. The LMM revealed a significant main effect of *Session* ($\chi^2(1) = 11.6, p < .001$) as well as a significant interaction between *Session* and *Group* ($\chi^2(1) = 4.85, p = .028$). No significant main effect of *Group* ($\chi^2(1) < .001, p = .987$) was found. Post-hoc comparisons indicated that free-hand movement time was shorter after tool use in the TD group ($t = 3.41, p = .004$; Figure 3B), while the duration of the movement was not modulated in the DCD group ($t = .291, p = .991$).

In sum, the group of TD participants depicted larger amplitudes and reduced latencies during the reaching component of their movement, resulting in shorter movement times. They also displayed a reduced latency, opening their fingers earlier, but no modulation of the amplitude of the grasping component. Instead, DCD children and early adolescents displayed modulations of the amplitudes of the reaching component only, while their latencies were not affected, which did not lead to a change in movement times. They however modulated their grasping component, opening their fingers both earlier and larger after tool use.

Forearm length estimation is reduced after tool use in both TD and DCD groups

Participants estimated their forearm shorter following tool use, as shown by a significant main effect of *Session* ($\chi^2(1) = 23.5, p < .001$; Figure 4; TD: 114% vs 105%; DCD: 131% vs 122%). However, the two groups did not significantly differ in their estimation (no main effect of *Group*: $\chi^2(1) = 1.91, p = .167$), which was overall slightly overestimated before tool-use. There was no evidence that DCD participants modulated their conscious forearm representation differently than their TD peers as shown by the absence of interaction between *Session* and *Group* ($\chi^2(1) = .528, p = .467$).




Figure 4. Forearm length estimation (in % of the actual forearm length) before (blue) and after (orange) tool use (left panel) and individual performance ordered by size (right panel). On the right panel, positive values indicate that the perceived forearm length increased after tool use, while negative values indicate reduced forearm length representation. Note that data was missing for one TD and one DCD participant. Most of the TD and DCD children and early adolescents estimated their forearm length to be shorter after tool use. Error bars indicate the means \pm 1 SEM. Asterisks denote significance.

TD and DCD children and early adolescents behave similarly during tool use

Reaching component. The LMM revealed a significant main effect of *Block* on all the latencies (all $p < .031$), and most of the amplitudes (all $p < .001$; except deceleration: $\chi^2(1) = 1.30$, $p = .255$). Latencies decreased and amplitudes increased between the first and last blocks of the tool-use session (Figure 5). Crucially, no evidence indicated that the DCD participants controlled the tool differently to reach for the object than the TD group. The main effect of *Group* was not significant neither on the latencies (acceleration: $\chi^2(1) = .003$, $p = .954$; velocity: $\chi^2(1) = .078$, $p = .780$; deceleration: $\chi^2(1) = .129$, $p < .719$) nor on the amplitudes (acceleration: $\chi^2(1) = .524$, $p = .469$; velocity: $\chi^2(1) = 9.04$, $p = .003$; deceleration: $\chi^2(1) = .583$, $p = .445$). Similarly, there was no significant *Block* \times *Group* interaction on the latencies (acceleration: $\chi^2(1) = .946$, $p = .331$; velocity: $\chi^2(1) = .441$, $p = .507$; deceleration: $\chi^2(1) = 1.26$, $p = .203$) and amplitudes (acceleration: $\chi^2(1) = .009$, $p = .925$; velocity: $\chi^2(1) = .069$, $p = .793$; deceleration: $\chi^2(1) = .371$, $p = .543$).




Figure 5. Kinematics of the Reaching component during the first (turquoise green) and the last (turquoise blue) blocks of tool use (top panel) and individual performance ordered by size (bottom panel) in TD and DCD participants for the latencies and amplitudes of velocity and deceleration. On the bottom panel, positive values indicate that the latency/amplitude increased with practice, while negative values indicate reduced latency/amplitude. Most of the TD and DCD children and adolescents reduced their latencies and increased their amplitudes with tool practice. Error bars indicate the means \pm 1 SEM. Asterisks denote significance.

Grasping component. Similar to the reaching component, a significant main effect of *Block* ($\chi^2(1) = 5.33$, $p = .021$) was found on MGA latency, with participants opening their tool fingers earlier with practice. There was no indication of significant modulation by *Group* ($\chi^2(1) = 1.96$, $p = .162$) nor an interaction between *Block* and *Group* ($\chi^2(1) = 1.62$, $p = .204$). As to the tool maximal grip aperture, the LMM did not reveal any significant main effect of *Block* ($\chi^2(1) = .187$, $p = .666$), nor *Group* ($\chi^2(1) =$