



Improving Stereotaxic Neurosurgery Techniques and Procedures Greatly Reduces the Number of Rats Used per Experimental Group-A Practice Report

Barbara Ferry, Damien Gervasoni

► To cite this version:

Barbara Ferry, Damien Gervasoni. Improving Stereotaxic Neurosurgery Techniques and Procedures Greatly Reduces the Number of Rats Used per Experimental Group-A Practice Report. *Animals*, 2021, 11 (9), pp.2662. 10.3390/ani11092662 . hal-03374352

HAL Id: hal-03374352

<https://hal.science/hal-03374352>

Submitted on 14 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improving Stereotaxic Neurosurgery Techniques and Procedures Greatly Reduces the Number of Rats Used per Experimental Group – A Self-Experience Report

Barbara Ferry ^{1,2} and Damien Gervasoni ¹

¹ Centre of Research in Neuroscience Lyon CNRS UMR 5292 - INSERM U 1028 - Université Claude Bernard Lyon 1, Université de Lyon

² Centre Hospitalier Le Vinatier - Bâtiment 462 – Neurocampus, 95 boulevard Pinel, 69675 Bron Cedex, France

* Correspondence: barbara.ferry@cnrs.fr; Tel.: + 33 4 81 10 65 57

Simple Summary: Techniques of stereotaxic surgery are commonly used today in research laboratories by a diversity of students, technicians and researchers. Over the past twenty years, technical and scientific progresses have been made in neurosurgery to meet the evolving requirements imposed by international legislation, and to promote the implementation of « three R » rules. These improvements were motivated by a greater awareness of animal welfare and the necessary effort in the reduction of the number of animals used in experiments. The data presented in the present study show that technical and methodological improvements brought to our surgical procedures from 1992 resulted in reproducible stereotaxic neurosurgeries and in a significant reduction experimental errors and animal morbidity. The decreasing effect of these improvements on the final number of animals used in our experiments, including (1) management of pain during and after surgery, (2) appropriate aseptic techniques and (3) correct stereotaxic surgical approaches are precisely described throughout the text.

Abstract: Techniques of stereotaxic surgery are commonly used today in research laboratories by a diversity of students, technicians and researchers. To meet the evolving requirements imposed by international legislation, and to promote the implementation of 3R rules (replacement, reduction and refinement) by reducing experimental error, animal morbidity and mortality, it is essential that standard operating procedures and proper conduct to follow during such complex surgeries be precisely described and respected. The present report shows how refinements of our own neurosurgical techniques over decades, have significantly reduced the number of animals (rats) used in experiments and improved the animals' well-being during the post-surgical recovery period. The current way of conducting pre-, per- and post-surgical procedures used in our laboratory is detailed. It describes, with high standard levels, the practical aspects of stereotaxic neurosurgery that has been refined in our laboratory since 1992 and that cover various areas including appropriate anaesthesia, pain management during and after surgery, method to determine the stereotaxic coordinates and the best approach to the target brain structure. The application of these optimal surgical methods that combines reliable and reproducible results with an acute awareness of ethics and animal welfare leads to a significant reduction in the number of animals included in experimental research in accordance with ethical and regulatory rules as required by the European Directive on laboratory animal welfare.

Keywords: stereotaxic neurosurgery; refinement; reduction; 3R; well-being; rat model

Citation: Lastname, F.; Lastname, F.; Last-name, F. Title. *Animals* **2021**, *11*, x. <https://doi.org/10.3390/xxxxx>

Received: 28 December 2020

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors.

Submitted for possible open access

publication under the terms and condi-

tions of the Creative Commons Attribu-

tion (CC BY) license

(<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Stereotaxic neurosurgery is used in combination with other diverse techniques necessary for addressing the specific objectives of a proposed neuroscience study. These

techniques include *in vivo* experimental procedures for activating or inactivating brain structures or transmitter systems, permanent selective lesioning, functional neuroanatomy using tracers, and acute or chronic measurements or recordings. With the Directive 2010/63/EU for the protection of laboratory animals 2010,¹ Europe has reinforced the implementation of the Three Rs principle (Replacement, Reduction and Refinement) on the use of animals for experimental and other scientific purposes, originally put forward by Russell and Burch in their “The Principles of Humane Experimental Technique”.² As in the preceding Directive (86/609/EEC), appropriate formation and training are required for all those engaged in the use of stereotaxic neurosurgery in rodents for scientific purposes. A high standard level of formation in stereotaxic neurosurgery implies a good knowledge in various areas of expertise but also a great awareness of ethics and animal welfare.

While legal frameworks applicable to animal research have considerably evolved since the early 1980s, animal care and use guidelines and recommendations have also been provided and regularly updated by expert groups or entities (eg National Research Council 201),³ Canadian Council on Animal Care 2013,⁴ Federation for Laboratory Animal Science Associations, animal welfare bodies, see Guillén & Steckler for review,⁵). Overall, in conjunction with regulatory incentive these helpful recommendations and advices have promoted better practices and improved the day-to-day follow-up and care of laboratory animals. Additionally, with a general public being more and more informed, animal researchers have also become increasingly aware of societal expectations on laboratory animal welfare. Compliance with the 3R principle, implementation of humane endpoints, higher scrutiny in pain recognition and management have now become essential requirements.

Animals, especially rodents, are widely used in neuroscience research in order to study normal and abnormal brain functions.⁶ Since 1992, our research is focusing on the role of the central structures involved in learning and memory. In particular our experiments have addressed the involvement of several temporal lobe structures in the successive memory phases underlying associative learning task in the rat model. For this purpose, we have used different techniques aimed at testing the effect of local activation or inactivation of target structures on these processes. Such approach involves the stereotaxic placement of guide cannulas in adult rats, that allow intracerebral injections (and collections) of fluids.

The present work describes the techniques based on common standard procedures of stereotaxic surgery, during which material can be inserted into the brain in order to microinfuse drugs in behaving animals performing a task. Based on the data published or collected in the laboratory notebooks, reports and theses from master and PhD-students under our supervision, we provide a detailed report on the changes that were introduced over time in our experimental procedures, and we show their effect on the proportion of animals that were excluded from the final experimental groups due to per- and post-surgical complications or intra-brain misposition of the material, throughout our research activities. The description of the general methodology includes the chronological list of the refinements (modification, additional or abandon of a component of the overall procedure) that have been introduced in the stereotaxic procedures during the period that covers our research activity from 1992 to 2018.

2. Material and Methods

2.1. Implementation of Aseptic Techniques

2.1.1. Environment and Material

Aseptic techniques have received a special attention since we started our experiment in 1992 and sterilization of the surgical equipment was a first step implemented in our procedure. To this end, surgical tools (surgical drapes, gowns, compresses, cannulae, electrodes, micrometer screws, obturators, tools used for drilling the skull bone, scalpel,

retractors, scissors and needle holder) were sterilized by placing them for 30 minutes at 170°C.

The ear and incisor bars, the hand piece of the dental drill and the stereotaxic frame were simply cleaned with disinfectant wipes.

From 1994 on, a change was introduced in the procedure and the cannulas were not sterilized by heat anymore. Instead, cannulas were put in a bath of hexamidine solution and the inside was systematically infused by a solution of hexamidine and rinsed with a sterile NaCl solution.

From 2005, we implemented from the beginning to the end of surgery a sequence of steps under the go-forward principle in order to limit contacts between soiled and sterile instruments or materials and therefore to keep a high level of asepsis. The organisation of space was included in this process with the delineation of two distinct areas, one called "dirty" area for the preparation of the animal (see below), and a "clean" zone for the surgery.

2.1.2. Preparation of the Surgeon

From 2005 on, an assistant was always present and helped the experimenter to conform with the go-forward principle and to prepare the surgical set-up. After a thorough surgical handwashing, the surgeon is helped for gowning and gloving, with a sterilized gown and mask as well as sterile gloves prepared in advance. Once the surgeon is equipped, the assistant places the instrument box on a sterile drape next to the stereotaxic frame.

2.1.3. Preparation of the Animal

A clinical exam is made to ensure that each animal has a good health status before proceeding. Importantly, none of the rats was subject to food restriction before surgery. The animals' weight is carefully measured for the adjustment of anaesthesia dosage and kept as reference for the post-surgical follow-up. An animal is first being anesthetized and prepared (surgical shearing) in the "dirty" area. Second, the paws and tail of the animal are gently cleaned with an iodine or hexamidine scrub solution. The animal is then brought by the assistant to the "clean" zone. From 1994 on, a thermostatically controlled heating blanket with a rectal probe was used to ensure an optimal control of internal body temperature. At this point, the surgeon takes charge, checks the depth of anaesthesia and install the animal into the stereotaxic frame by fixing its head between ear- and nose-bars. An ophthalmic ointment is applied on the eyes to protect the cornea from desiccation. The top of the animal's head is then scrubbed with an iodine foaming solution (Vetidine Scrub®), rinsed with sterile water and disinfected with iodine solution (Vetidine Solution®). Alternatively, chlorhexidine-based soap and solution (Hibitane®) were used. The surgeon then let the antiseptic solution dry out and checks the position of the head.

2.2. Anaesthesia and Presurgical Analgesia

From 1992 to 1999, anaesthesia was induced by an intraperitoneal (i.p.) injection of 5 mg/kg diazepam followed by 100 mg/kg ketamine (Ferry et al., 1995; 1996). From 1999 to 2006, anaesthesia was induced by using i.p. injection of sodium pentobarbital (50 mg/kg) and supplemented with given atropine sulfate (0.4 mg/kg, i.p.) to suppress salivation and bronchial secretions.^{7,8,9,10,11,12} From 2006, anaesthesia was induced by i.p. injection of a mixture of 3/4 part of ketamine (Imalgene®; 100 mg/kg) and 1/4 part of the alpha-2 agonist xylazine hydrochloride (Rompun®, 10 mg/kg).^{13,14,15,16,17,18,19,20} Anaesthesia was maintained by reinjections of a half and quarter-dose after an initial phase of 30 to 45 min.

From 1992, the animals received a subcutaneous injection of lidocaine (Xylocaine®) at the incision site once deeply anesthetized (suspension of consciousness, loss of tail- or toe-pinch withdrawal reflex, pain sensitivity and mobility.^{21,22} From 2006, the animals

received also a general analgesic through a subcutaneous injection of an analgesic dose of the mu opioid receptor agonists butorphanol (Dolorex®) 0.4mg/kg and later on buprenorphine (Buprecare©) 0.05 mg/kg in order to reduce pain during surgery.^{23,24}

2.3. Stereotaxic surgery procedure

Incision of the skin and exposure of the cranial surface is made using a scalpel with sterile blade and tissue retractors are placed on each side of the cranial area of interest. The bone surface is gently scraped with a periosteal raspator or a bone scraper. Sterile cotton-tips and saline solution are used to remove bone debris and dry the skull surface.

The cannula holder, or injection needle is installed on the object carrier and its verticality relative to the coronal and median planes is checked. The coordinates of the bregma and lambda landmarks are measured in order to position the skull in a horizontal position ("flat-skull"). The dorso-ventral (DV) coordinates of both landmarks are checked and the inclination of the head is adjusted whenever the difference between both DV coordinates exceeds 0.1 mm. Sterile anchor screws (when needed; for most procedures two or three screws are used) are then inserted to the skull to strengthen the fixation of the implant.

From 1995, the "flat-skull" position was verified once. From 2004, the flat position of the skull was checked twice before and after the insertion of the anchor screws.

Then, the ML and AP coordinates of the target structure are calculated relatively to the appropriate reference (Bregma or Lambda) with support of a reference atlas and its updated versions.^{25,26,27} The needle/cannula is then positioned above the target and its tip used to determine the final DV coordinate of the target from the skull surface. The skull is then drilled, and the dura is hooked using the tip of a small sterile hypodermic needle, and incised in a circular motion.

From 1995, the verification of the vertical position of the holders was also checked after having positioned the carrier on the micromanipulator on the stereotaxic frame.

From 2004, the final DV coordinates of the target were calculated from the dura surface and the cannula, or injection needle, was installed on the object carrier after the dura was resected. This step was repeated in the case of bilateral insertion.

Then, the cannula or injection needle is carefully lowered according to the DV coordinates, with a speed of about 1 millimeter per minute.

In 2012, we introduced a new holder with groves aligned at fixed intervals to maintain two cannulas strictly in parallel for bilateral implantations. This tool allowed the simultaneous insertion of two parallel guide cannulas in a single move of the holder. From this period on, the axial position of the skull was also systematically verified before inserting the material.

When the material has been inserted correctly in the brain, the skull is dried and a self-curing adhesive resin (Super-Bond C&B®, Sun Medical Co., Japan) is applied.²⁸ The guide cannula and anchor screws are then fixed solidly to the skull of the animal using dental acrylic cement (e.g. Unifast Trad® powder and solvent, GC America Inc.). The cement is applied in successive thin layers to protect the brain from the thermal effect of the acrylic polymerisation. The edges of the implant are made as smooth as possible. When the cement has completely dried out, the object carrier is slowly raised leaving the guide cannula in place and visible on the skull. A sterilized obturator (stylet) is then inserted into the guide to avoid clogging during the post-operative recovery period. Depending upon the size of the skin incision and the volume of the implant, the wound is sutured at the end of the procedure. For this purpose, sterile suture material is used. Single stiches are done using resorbable suture wire on a 3/8 curved needle. At the end of the procedure, the wound is cleaned with a compress soaked with a 1:10 dilution of disinfectant in sterile water. Any trace of cement, hair or blood clots or other debris that can interfere with wound healing is removed. Before the animal fully recovers from anaesthesia, the nails of all paws are trimmed to minimize any impact of the animal's nor-

mal scratching and grooming behaviour on the healing process. The return of the palpebral reflex is checked using a cotton-tip applied on the eye's corner.

From 1995, pilot studies were conducted with injections of dye at the target sites and immediate histological verification of the position of dye deposits to determine the correspondence between the theoretical coordinates and actual placement within the target area. Occasional discrepancies between actual and atlas coordinates were thus corrected.²⁹

From 2004, cannulas with reduced diameter were used (from 22 to 23 gauge).

2.4. Monitoring Vital Signs During Surgery

Toe- or tail-pinch withdrawal reflexes, breathing rhythm and amplitude, colour of the mucous membranes, body temperature and possible dehydration of the animal are checked throughout the surgery. This monitoring was applied since 1992. From 1994, in addition to the visual control of the animal's breathing and the colour of its mucous membranes, internal temperature was systematically monitored and maintained throughout the surgery.

2.5. Post-surgical Procedure

At the end of the surgical procedure, the animal is temporarily placed in the recovery zone with an infrared spot above the cage. Importantly, the light is placed such that an animal can move away from the hot spot once recovering its righting reflex. Post-operative care procedures such as manual feeding and facilitated access to drinking source are provided until an animal recovers its autonomy and a complete restoration of normal behaviour. Each animal returns to the housing facilities only after complete awakening from anaesthesia and is placed back in its home cage with its litter mates.

From 1992 to 1994 systematic i.p. injection of 5 ml of NaCl and heating of the animal at the end of surgery have been used. From that period, animal facilities staff were involved in daily post-surgical care and nursing. Individual follow-up sheets were filled for each animal which lists all the substances an animal had received pre- per and post-surgery. In some rare circumstances of severe complication, the involvement of the technical staff enabled rapid euthanasia according to predefined endpoints, thus avoiding unnecessary suffering. Also, from 1992 to 2000, a prophylactic antibiotic treatment (penicillin 0.12 MU/0.4 ml, im) was used in order to complete our aseptic techniques.^{7,30,31,32}

From 2012, i.p. injections of 5 ml of glucose Ringer lactate 5% replaced the use of NaCl at the end of the surgery and all animals received a subcutaneous injection of a nonsteroidal anti-inflammatory drug (carprofen, Rimadyl®). An evaluation of signs of distress and pain was done daily, observations were written on the individual's follow-up sheet.³³ Anti-inflammatory treatment was maintained for at least 48h and prolonged when needed. Contemporaneous records of each animal served to prepare a final report on animal use, and to determine the actual level of pain and distress of the procedures. With the improvement of the aseptic conditions, the antibiotic prophylaxis was discontinued in 2000.

2.6. Pharmacological Experiments

Animals were allowed to completely recover from the surgery for at least two weeks before performing behavioural and pharmacological studies. During this period, the experimenter visited the animals once daily, and check the integrity of the implant and wound, as well as the operability of the guide cannula. For this purpose, each animal was carefully handled and restrained by hand while the experimenter checked the presence of the stylet and the absence of resistance when sliding it up and down. This daily handling contributed to the habituation of the animals to the experimental conditions. The injected drugs were always freshly prepared on the day of the experiment using aseptic

techniques and frozen aliquots of sterile medium (sodium chloride 0.9%, or artificial cerebrospinal fluid solution). In addition, the inside of the tubing was infused with a 70% alcoholic solution before each microinfusion. From 1994 on, the animal care staff was involved in the follow-up of the animals and would inform the experimenter whenever an abnormal or unexpected situation occurred.

2.7. Post-mortem Examination of Animals

After completion of behavioural and pharmacological testing, each rat was euthanized for brain removal and histological examination of the injection sites. Euthanasia was obtained with an overdose of anaesthetic.

2.8. Statistics

All analyses were performed with Prism (GraphPad Software Inc.). For normalization purpose, a ratio of the “total number of animals discarded from the statistical analyses / total number of animals used in experimental research in our group from 1992 to 2018” was calculated and analysed. The Agostino Pearson’s analysis was used to test the normality of the distribution and this was followed by Pearson’s parametric correlation test.

3. Results and Discussion

Data has been obtained from a total of 113 studies and reports in our research group from 1992 to 2018. A total of 3228 animals, weighing 250-300g at surgical period, were involved in the present study (n = 348 wistar; n = 1880 Long-Evans and n = 1000 Sprague-Dawley strains). The numbers of animal finally discarded in each study are compiled and presented as proportions for normalization and inter-studies comparison purposes.

The proportion of animals that were discarded from the final statistical analyses are spread relatively to the category of improvement brought in our experimental procedure. Figure 1 shows the effect of refinements in anaesthesia and control of vital signs on the general recovery of the animals (clinical signs of post-surgical pain and distress); Figure 2 illustrates the effect of improvement of aseptic techniques on the number of infectious complications (internal and external) and Figure 3 shows the effects of stereotaxic methodology improvement influenced the number of animals discarded because 13.




Figure 1. Effect of refinement in anaesthesia, pain management and rehydration on animal recovery. Letters indicate the chronology of various improvements introduced in the surgical procedure. (a): from 1994-96, internal temperature has been monitored and maintained. Systematic i.p. injection of 5 ml of NaCl and heating of the animal at the end of surgery have been used. Involvement of animal facilities staff in daily post-surgical care and nursing. Individual follow-up is filled for each animal which lists all the substances the animal received pre- per and post-surgery and daily evaluation of signs of distress. (b): from 2004-06, diazepam/xylazine or sodium pentobarbital anaesthetics have been gradually replaced by the association of ketamine and xylazine. (c): from 2012, systematic i.p. injections of 5 ml of glucose Ringer lactate 5% at the end of the surgery and a subcutaneous injection of an anti-inflammatory were done at the end of the surgery.




Figure 2. Effect of refinement in the surgical and microinjection procedures on aseptic complications. Letters indicate the chronology of various improvements introduced in the procedures. (d): from 1994, the inside of the cannulas was systematically sterilized. In addition, during the microinfusion procedure, the inside of the tubing, the injection needles and the injected drugs were prepared in careful aseptic conditions. (e): from 2005, we implemented an asepsis chain following a “go-forward” workflow.




Figure 3: Effect of refinement in the stereotaxic procedure on the accuracy of placement of the implanted item(s). LetTable 1995. verification of the flat-skull position before each surgical session. Verification of the vertical position of the holders. Introduction of pilot studies and immediate histological verification of implantation site to check the correspondence between the theoretical coordinates and actual placement within the target area. (g): from 2004, reduction of the cannula diameter, final DV coordinate of the target structure is calculated from the dura; systematic verification of the flat position of the skull after screw position and on each animal (h): from 2012, development of a fixed spaced slide for cannulas (bilateral implantation); systematic verification of the axial position of the skull.

As illustrated in the figures, we observed significant decreases in the number of animals excluded from final analyses following the implementation of technical (in stereotaxic pre-, per- and post-surgery) and welfare (pain and distress managements, anaesthesia) refinements in our procedures from 1992. Statistical analyses confirmed these observation and Pearson rank correlation revealed a significant relationship between the proportion of animals discarded from the analyses and the refinement in the different techniques throughout the successive periods of research ($r = -0.93$, $P < 0.001$ for Figure 1; $r = -0.94$, $P < 0.001$ for Figure 2 and $r = -0.95$, $P < 0.001$ for Figure 3). These refinements concern different areas of expertise that are listed below.

3.1. Pain and Distress

Figure 1 shows how technical improvements in our anaesthesia and pain management procedures positively impacted post-surgical welfare of the animals. While the priority was to monitor the proper depth of anaesthesia during the operation by checking the various reflexes in 1992, gradual increase in the attention to general homeostasis, body temperature and proper hydration maintenance of the animal prompted us to complete our surgical procedure. As shown in Figure1 (event a), improvements brought to the monitoring technique from 1994 lead to a substantial decrease in the post-surgical mortality of the animals. From 2004 (event b), replacement of diazepam/xylazine or sodium pentobarbital by the association of ketamine and xylazine improved the overall quality of surgical anaesthesia, by reducing the time of induction and nociception and increasing myorelaxation.³⁴ Moreover, from 2012, other refinements such as

post-operative management of pain by administration of a NSAID at the end of the procedure, introduction of health follow-up checks involving the animal facility staff and modification of the rehydration procedure resulted in a reduction of the number of animals discarded from the experimental groups.

The welfare of animals included in an experiment is essential to ensure reliable and reproducible the results. From their arrival in the laboratory facilities, animals were kept under a constant supervision in order to detect signs of pain and distress as early as possible. Recognition of these signs can sometimes be difficult especially in prey animals, but helpful information can be found in the literature.^{33,35,36} From 1994, animal caretakers were trained to evaluate post-operative signs of distress and involved in the daily follow-up of animals. This involvement of the animal facility staff improved the level of animal care and nursing.

Minimizing pain and discomfort during and after a surgery strongly relies on the choice of sedation, anaesthetic and analgesic compounds, and an appropriate rehydration procedure when needed, but also on the monitoring procedure. Depth of anaesthesia, and vital signs such as respiratory rhythm and amplitude, should frequently be checked throughout surgery in order to react rapidly. General anaesthesia causes a dose-dependent impairment of thermoregulation, a depression of cardiovascular and respiratory systems,^{22,37} a lowering of metabolic activity and a peripheral vasodilation that can lead in a matter of minutes to a drop in body temperature and a fatal hypothermia under 32°C. Moreover, prolonged hypothermia during a surgery can lead to difficulties in awaking, discomfort and complications from poor blood perfusion, inadvertent coagulation and ischemia. Conversely, hyperthermia might lead to death. Since a faulty heating blanket or probe can cause severe harm, appropriate check and maintenance of heating devices is recommended when preparing the surgery and attention should be paid to its correct functioning throughout the procedure. From a general point of view, a good monitoring procedure can prevent complications occurring during surgery.

3.2. Surgical Asepsis

Surgical asepsis refers to the absence of infectious microorganisms during an invasive procedure. Such organisms are naturally present in the environment, in animals and humans, and on all foreign material that can be introduced in a living tissue (biofilm). Aseptic techniques are thus essential to prevent **surgical site infections**, post-surgical complications and to minimize the risk of mortality.

Figure 2 shows the effect of asepsis-related refinements on the post-surgical welfare of the animals. As shown in the figure, a first noticeable decrease in the number of animals discarded from the analyses was observed at period 1994-96 (event d). Less parenchymal over-reactions were observed post-mortem when surgical tools and the material such as cannulas, anchoring screws, stylets and cannula holders were chemically decontaminated by a bath of antiseptic solution. A special attention was paid to the decontamination of the inside of the cannulas. Stylets that were replaced in the cannulas throughout the experiment were always decontaminated with 70% alcoholic solution before being re-inserted. The second important event that influenced the number of animals discarded from the final experimental groups after the 2004-2006 period (event e) corresponded to the enforcement of the “go-forward” process. With the clear spatial delineation of three zones (i.e. “dirty”, “intermediate” and “clean”), we were able to implement a “go-forward” workflow whatever the type of manipulation, i.e. the surgery itself, single microinfusion or stylets replacement. As shown on the figure, all these refinements lead to a sustained reduction of the number of animals used in the analyses so that today there is no more animals excluded from experimental groups due to contamination event.

3.3. Stereotaxic Coordinates and Angle of Approach During Surgery

When determining stereotaxic coordinates of a target structure one should consider its anatomical and functional characteristics, its spatial coordinates as given by the atlas, the organization of the cerebral vascular system, and the age or state of development of the animal. In order to ensure a high precision and reproducibility of stereotaxic surgeries, the implantation of material such as cannulas, electrodes or microdialysis probes should aggregate all these informations.

In this regard, from 1992, a series of improvements have been brought to our stereotaxic calculation of the brain target coordinates by a systematic verification of the adequacy between the theoretical position of the target structure found in the atlases^{25,26,27} and its actual position in our animal model.²⁹ In addition, the cerebrovascular anatomy, especially the arrangement of the leptomeningeal vessels has been taken into account for the calculation of the angle of approach in order to avoid critical danger of rupturing important vessels such as the transverse sinus during material insertion.^{38,39,40}

As shown in figure 3, the first event observed after the 1994-96 period corresponded to a refinement in the method to verify the angle of insertion of the cannula into the brain. From that period, the flat position of the skull was verified before each session of surgery. To do this, the correspondence between the DV (± 0.1 mm) coordinates measured at the Lambda and Bregma reference points was verified before inserting anchoring screws. From that period, we also systematically verified the vertical alignment of the cannula holders. Both operations significantly contributed to improve the repeatability in the intracerebral A/P placement of the cannula. Also, pilot non-survival surgeries and immediate histological verification of implantation site were conducted at that period on animals previously used in other experiments. This helped to adjust the stereotaxic coordinates indicated in the Paxinos and Watson atlas in Wistar rats (1) to those found in other type of rats (Sprague Dawley, Long Evans) and (2) to those found in rats of a different weight.

The second important improvement was observed during the 2002-2004 period when cannulas of a smaller diameter were introduced (from 23 to 24 gauge); less invasive, thinner cannulas have a better tissue tolerance and allow a more precise positioning into the brain. Also, from that period, due to the imprecision observed in some cases between the theoretical DV coordinate of the target structure and the position of the cannula observed post-mortem (i.e. convex shape of the cranial bone), the final D/V coordinate of the target structure was calculated from the dura. In addition, from 2002, the flat position of the skull was verified before and after the screw anchoring.

The third significant progress in the method appeared in the 2010-12 period and resulted from the development of custom-made holders maintaining a given distance and a strict parallelism between cannulas for bilateral implantations in symmetric areas. Such holders were manufactured with parallel grooves to fit exactly the M/L distance between each structure targeted. Also, the ML axial flat skull position was systematically verified in order to ensure perfect vertical position and right M/L position of the cannulas. To this aim, the D/V coordinates of the point above the target structure was bilaterally checked for correspondence.

As shown on the figure, these refinements lead to a steady decline of the number of animals discarded from the analyses due to misplacement of lesion or injection site. Our method has gained in accuracy and reproducibility so that only 1/166 subject was excluded from the final analysis in the last period.

3.4. Further Refinement Perspectives

The refinements introduced in our research are somewhat diverse and might not apply to any stereotaxic surgery. Stereotaxic surgeries can indeed be used a variety of research projects; while general recommendations can apply, such as the systematic use of aseptic techniques in survival procedures, others need to be considered on a

case-by-case basis. This seems particularly true when considering the nature of anaesthetic, analgesic and other drugs to administer to an animal. Dosage and administration route are obvious parameters, but others such as half-life and potential side-effects also deserve attention especially in neuroscience research. Some drugs acting on the central nervous system can have delayed effects that can impact the animal's behaviour and well-being. For example, compounds like buprenorphine²⁴ or fentanyl offer nice pain relief properties but can also cause nausea⁴¹ and a decrease of intestinal motility.^{42,43} Some anaesthesia methods can cause excessive salivation and bronchial secretion. To minimize such effect, experimenters can consider the use of atropine sulfate. Because of its central parasympathomimetic effect, especially at low dose, atropine sulfate can also induce bradycardia instead of tachycardia. Experimenters can alternatively consider glycopyrrolate (Robinul®) that does not cross the blood-brain barrier or choose another anaesthesia method that does not cause excessive secretions. The choice of a drug should also take into account the potential delayed effect, such as anxiety signs, that may need a specific test to be objectivated.

Regarding anesthesia in our own research, we observed a significant decrease in mortality following the refinement in the method. As an alternative to fixed anaesthesia, we have introduced inhalational anaesthesia, since it has been shown to be safer and more effective than parenteral anaesthetics.^{44,45} Isoflurane, the most used compound⁴⁶ has a rapid induction and awakening due to its low solubility.⁴⁷ It provokes little or no salivation or tracheobronchial secretions. By adjusting the vaporizer, the level of anaesthesia can be rapidly adjusted. Recommended levels for use in the rodent are 2 to 3% for induction, and 0.25 to 2% for maintenance. Moreover, a supplement of oxygen should be used during surgery to avoid hypoxia.⁴⁸ In order to control this, several devices now exist for monitoring blood gases (oximetry, capnography) and arterial pressure in rodents. Although not widely considered as being necessary for operator procedures of relatively short duration, they will be tested for use in the next future in our laboratory during long surgeries.

Because of its poor analgesic properties and its rapid elimination, switching to isoflurane anaesthesia prompted us to reconsider our analgesia protocols. Since pain can be much more difficult to control once the animal has become conscious of it, we systematically use local anaesthetic drugs (lidocaine, bupivacaine), and opioids (butorphanol, buprenorphine) for surgery and a post-operative 48h treatment with NSAIDs (carprofen) to provide good analgesic levels and comfort for the animals.

Any significant change in a procedure should therefore be thoroughly thought before being implemented. A cost/benefit approach might be appropriate in this matter. Such methodology implies a careful assessment of many parameters, not only during the surgery itself, but also during the immediate and long-term periods that follows. These parameters can strongly differ between research studies. When targeting a brain area involved in motor control, the follow-up of animal's well-being should include signs of motor deficits or handicap. Such signs can be extremely subtle, or even somehow compensated, but they should always be anticipated. Animal caretakers are the first-line actors, because of their presence in animal facilities, but refining a complete procedure will also benefit from close interactions between experimenters, the animal welfare body involved in the follow-up of projects and the designated veterinarian. If required by the ethical oversight body or institutional animal care and use committee, a retrospective assessment can also provide guidance and suggestions for future improvements.

Although the proportion of animals excluded from our experiment today has reached a low level, we continue our efforts to further improve our procedure and the welfare of the animals without forgetting the well-being of the experimenter and facility staff, bearing in mind that good working conditions (physical and mental) will contribute to high-quality care to the animals and limit compassion fatigue. For this purpose, we regularly update and enrich our training program for experimenters and caretakers in our animal facility on stereotaxic surgery, nursing skills and ethical requirements.

4. Conclusion

Over the past twenty years, technical and scientific progresses have been made in neurosurgery to meet higher ethical standards. These improvements of pre- per and post-surgical approaches were motivated by a greater awareness of animal welfare and the necessary effort in the reduction of the number of animals used in experiments. We identified three major domains where improvements allowed us to achieve reliable and reproducible stereotaxic neurosurgeries with a lower number of subject rejections. These domains include (1) management of pain during and after surgery, (2) appropriate aseptic techniques and (3) correct determination of stereotaxic coordinates and surgical approaches.

The data presented in the present study show that refinements made on our procedures from 1992 to 2018 resulted in a significant reduction in the number of animals used in experimental research through continuous decrease in experimental errors and animal morbidity over time. These technical and methodological improvements were guided by our growing ethical and animal welfare that prompted us to evaluate and implement new practices and technologies found in the literature and throughout numerous exchanges within the Neuroscience community.

The Authors declares that there is no conflict of interest.

References

1. Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes. Official Journal of the European Union, Strasbourg, <http://data.europa.eu/eli/dir/2010/63/2019-06-26> (2010).
2. Russell, WMS and Burch RL. The Principles of Humane Experimental Technique. London, UK: Methuen, 1959.
3. National Research Council (US) Committee for the Update of the Guide for the Care and Use of Laboratory Animals. Guide for the Care and Use of Laboratory Animals. 8th ed. Washington (DC): National Academies Press (US), 2011.
4. Canadian Council on Animal Care. Guide to the care and use of experimental animals: Chapter XXI (laboratory rats), www.ccac.ca/Documents/Standards/Guidelines/Vol2/rats.pdf (2003).
5. Guillén J, Steckler T. Good Research Practice: Lessons from Animal Care and Use. In: Bernalov A, Michel MC and Steckler T (eds) Handbook of Experimental Pharmacology. Berlin: Springer, 2019, pp.367-382.
6. El Ayache N, Galligan JJ. The rat in neuroscience research. In: Mark A, Suckow F, Hankenson C et al. (eds) American College of Laboratory Animal Medicine, The Laboratory Rat. 3rd ed. Academic Press, 2019, pp. 1003-1022.
7. Ferry B and McGaugh JL. Clenbuterol administration into the basolateral amygdala post-training enhances retention in an inhibitory avoidance task. NLM 1999; 72:8-12.
8. Ferry B, Roozendaal, B and McGaugh JL. Involvement of alpha1-adrenoceptors in the basolateral amygdala in modulation of memory storage. Eur J Pharmacol 1999; 372:9-16.
9. Ferry B, Roozendaal B and McGaugh JL. Basolateral amygdala noradrenergic influences on memory are mediated by an interaction between beta- and alpha1-adrenoceptors. J Neurosci 1999; 19:5119-5123.
10. Roozendaal B, De Quervain D, Ferry B et al. Basolateral amygdala-nucleus accumbens interactions in mediating glucocorticoid enhancement of memory consolidation. J Neurosci 2001; 21:2518-2525.
11. Majchrzak M, Ferry B, Marchand AR et al. Entorhinal cortex lesions disrupt fear conditioning to context but spare fear conditioning to a tone. Hippocampus 2006; 16(2):114-124.
12. Ferry B, Ferreira G, Traissard N et al. Selective involvement of the lateral entorhinal cortex in the control of the olfactory memory trace during conditioned odor aversion in the rat. Behav Neurosci 2006; 120(5):1180-1186.
13. Miranda MI, Rodríguez-García G, Reyes-Lopez JV et al. Blockade of β -adrenergic receptors in basolateral amygdala or insular cortex disrupts appetitive but not aversive taste. NLM 2008; 90(1):54-61.
14. Ferry B, and McGaugh JL. Involvement of the basolateral amygdala α 2-adrenoceptors in memory modulation of inhibitory avoidance in the rat. L and M 2008; 15(4):238-43.
15. Ferry B and Duchamp-Viret P. The orexin component of fasting triggers memory processes underlying food selection in the rat Orexin. L and M 2014; 21:185-189.

16. Boisselier L, Ferry B and Gervais, R. Selective involvement of NMDA transmission in the lateral entorhinal cortex for the formation of cross-modal olfactory-tactile associations in the rat. *Hippocampus* 2014; 24:877-891. 533
17. Ferry B, Parrot S, Marien M et al. Noradrenergic influences in the basolateral amygdala on inhibitory avoidance memory are mediated by an action on $\alpha 2$ -adrenoceptors. *Psychoneuroendocrinol* 2015; 51:68-79. 534
18. Ferry B, Herbeaux K, Javelot H et al. The entorhinal cortex is involved in conditioned odor and context aversions. *Front Neurosci Res* 2015; 9:342. 535
19. Boisselier L, Ferry B and Gervais R. Respective role of the dorsal hippocampus and the entorhinal cortex during the recombination of previously learned olfactory-tactile associations in the rat. *L and M* 2017; 24(1):24-34. 536
20. Estrade L, Duchamp-Viret P, Cassel JC et al. Microdialysis unveils the role of the alpha2-adrenergic system in the basolateral amygdala during acquisition of conditioned odor aversion in the rat. *ACS chem Neurosci* 2019; 10(4):1929-1934. 537
21. Harrison NL. Knockin' on the door of general anesthetic mechanisms: but will US researchers be shut out ? *Anesth Analg* 2003; 97:616-618. 538
22. Liles JH and Flecknell PA. The effects of buprenorphine, nalbuphine and butorphanol alone or following halothane anaesthesia on food and water consumption and locomotor movement in rats. *Lab Anim* 1992; 2:180-189. 539
23. Roughan JV and Flecknell PA. Buprenorphine: a reappraisal of its antinociceptive effects and therapeutic use in alleviating post-operative pain in animals. *Lab Anim* 2002; 36:322-343. 540
22. Flecknell P. Laboratory animals anesthesia. 3rd ed. Academic Press, 2009. 541
24. Paxinos G and Watson C. The Rat Brain in Stereotaxic Coordinates. 2nd ed. Sidney: Academic Press, 1982. 542
25. Paxinos G and Watson C. The Rat Brain in Stereotaxic Coordinates, 3rd ed. San Diego: Academic Press, 1996. 543
26. Paxinos G and Watson C. The rat brain in stereotaxic coordinates. 5th ed. San Diego: Academic Press, 2004. 544
27. Gervasoni D, Darracq L, Fort P et al. Electrophysiological evidence that noradrenergic neurons of the rat locus coeruleus are tonically inhibited by GABA during sleep. *Eur J Neurosci* 1998; 10:964-70. 545
28. Azimi N, Yadollahikhales G, Argenti JP et al. Discrepancies in stereotaxic coordinate publications and improving precision using an animal-specific atlas. *J Neurosci Methods* 2017; 284:15-20. 546
29. Ferry B, Sandner G and Di Scala G. Neuroanatomical and functional specificity of basolateral amygdaloid nucleus in Taste Potentiated Odor Aversion. *NLM* 1995; 64:169-180. 547
30. Ferry B, Oberling P, Jarrard LE et al. Facilitation of conditioned odor aversion learning by entorhinal cortex lesions in the rat. *Behav Neurosci* 1996; 110(3):443-450. 548
31. Wirth S, Ferry B and Di Scala G. Facilitation of olfactory recognition by entorhinal cortex lesions in the rat. *Behav Brain Res* 1998; 91:49-59. 549
32. Ferry B, Gervasoni D and Vogt C. Stereotaxic neurosurgery in laboratory rodent - handbook on best practices. New York: Springer Verlag 2014. 550
33. Stokes EL, Flecknell PA, Richardson CA. Reported analgesic and anaesthetic administration to rodents undergoing experimental surgical procedures. *Lab Anim* 2009; 43:149-154. 551
34. Rehbindler C, Baneux P, Forbes D et al. FELASA recommendations for the health monitoring of mouse, rat, hamster, gerbil, guinea pig and rabbit experimental units. *Lab Anim* 1995; 30:193-208. 552
35. Nicklas W, Baneux P, Boot R et al. Recommendations for the health monitoring of rodent and rabbit colonies in breeding and experimental units. *Lab Anim* 2002; 36:20-42. 553
36. Albrecht M, Henke J, Tacke S et al. Influence of repeated anaesthesia on physiological parameters in male Wistar rats: a telemetric study about isoflurane, ketamine-xylazine and a combination of medetomidine, midazolam and fentanyl. *BMC Vet Res* 2014; 10:310. 554
37. Bär T, Scheck T. The vascular system of the rat cerebral cortex - basic pattern of leptomeningeal vessels and numerical densities of neocortical arteries and veins. In: Auer LM and E Loew E (eds) *The cerebral veins – An experimental and clinical update*. Wien: Springer Verlag, 1983, pp. 65-72. 555
38. Scremin OU. Cerebral Vascular System. In: Paxinos G (ed) *The Rat Nervous System*. New York: Academic Press, 2004, pp.1167-1202. 556
39. Dashwood MR and Loesch A. Endothelin-1 as a neuropeptide: neurotransmitter or vascular effects? *J Cell Commun Signal* 2010; 4:51-62. 557
40. Sarabia-Estrada R, Cowan A, Tyler B et al. Association of nausea with buprenorphine analgesia for rats. *Lab Anim* 2017; 46: 242-244. 558
41. Cowan A, Doxey JC and Harry EJ. The animal pharmacology of buprenorphine, an oripavine analgesic agent. *Br J Pharmacol* 1977; 60:547-554. 559

-
42. Kuo A and Smith MT. In vivo profiling of four centrally administered opioids for antinociception, constipation and respiratory depression: Between-colony differences in Sprague Dawley rats. *Clin Exp Pharmacol Physiol* 2018; 45:1056-1066. 587
588
589
43. Sonner JM, Antognini JF, Dutton RC et al. Inhaled anesthetics and immobility: mechanisms, mysteries and MAC. *Anesth Analg* 2003; 97:718-740. 590
591
44. Jensen EW, Litvan H, Struys M et al. Pitfalls and challenges when assessing the depth of hypnosis during general anesthesia by clinical signs and electronic indices. *Acta Anaesthesiol Scand* 2004; 48:1267-1269. 592
593
45. Messier C, Emond S and Ethier K. New techniques in stereotaxic surgery and anesthesia in the mouse. *Pharmacol Biochem Behav* 1999; 63(2):313-318. 594
595
46. Jurd R, Arras M, Lambert S et al. General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA(A) receptor beta3 subunit. *The Fed Am Soc Exp Biol Journal* 2003; 17:250-252. 596
597
47. Fornari RV, Wichmann R, Atsak P, et al. Rodent stereotaxic surgery and animal welfare outcome improvements for behavioral neuroscience. *J Vis Exp* 2012; 59:e3528. 598
599