



HAL
open science

The cis-enhancers 3'RR and 5'Emu are Independent Motors of IgH Locus Remodelling

Melissa Ferrad, Nour Ghazzaoui, Hussein Issaoui, Alyssa Martin, Jeanne Cook-Moreau, François Boyer, Sandrine Le Noir, Yves Denizot

► **To cite this version:**

Melissa Ferrad, Nour Ghazzaoui, Hussein Issaoui, Alyssa Martin, Jeanne Cook-Moreau, et al.. The cis-enhancers 3'RR and 5'Emu are Independent Motors of IgH Locus Remodelling. Immunology and Cancer Biology, 2021. hal-03374932

HAL Id: hal-03374932

<https://hal.science/hal-03374932>

Submitted on 12 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Book Chapter

The *cis*-enhancers 3'RR and 5'E μ are Independent Motors of IgH Locus Remodelling

Melissa Ferrad[#], Nour Ghazzaoui[#], Hussein Issaoui[#], Ophélie Alyssa Martin, Jeanne Cook-Moreau, François Boyer, Sandrine Le Noir and Yves Denizot*

CNRS UMR 7276, Inserm U1262, Equipe Labellisée LIGUE 2018, Université de Limoges, France

[#]Equal contributors

***Corresponding Author:** Yves Denizot, CNRS UMR 7276, Inserm U1262, Equipe Labellisée LIGUE 2018, Université de Limoges, France

Published **July 05, 2021**

How to cite this book chapter: Melissa Ferrad, Nour Ghazzaoui, Hussein Issaoui, Ophélie Alyssa Martin, Jeanne Cook-Moreau, François Boyer, Sandrine Le Noir, Yves Denizot. The *cis*-enhancers 3'RR and 5'E μ are Independent Motors of IgH Locus Remodelling. In: Hussein Fayyad Kazan, editor. Immunology and Cancer Biology. Hyderabad, India: Vide Leaf. 2021.

© The Author(s) 2021. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author Contributions: MF, HI, NG, FB, OAM, JCM, SLN and YD designed and performed experiments and wrote the manuscript. YD obtained financial grants.

Conflicts of Interest: Authors declare no conflict of interest.

Acknowledgements: This work was supported by grants from Ligue Contre le Cancer (Equipe labellisée LIGUE 2018) and Agence Nationale de la Recherche (ANR: projet EpiSwitch-3'RR 2016). N. Ghazzaoui was supported by a grant from Association de Spécialisation et d'Orientation Scientifique (Lebanon), the municipality of Khiam (Lebanon) and the Société Française d'Hématologie. H. Issaoui is supported by a fellowship of the University of Limoges. F.B. is supported by Fondation Partenariale de l'Université de Limoges and ALURAD. We thank the genomics platform of Nice Sophia Antipolis for RNAseq experiments and the animal facility of University of Limoges for mice housing.

Abstract

B-cell development is spatially and temporally regulated with the Ig heavy chain (IgH) locus as a conductor. Starting with the first steps of B-cell development, IgH DNA remodeling constantly occurs under the control of the two *cis*-regulatory elements 5'E_μ (during early stages of B-cell development) and 3'RR (during late stages of B-cell development). These enhancers are reported to physically interact via IgH DNA loops of still unclear functional significance at the pro-B cell stage. We thus investigated if 5'E_μ and 3'RR were independent drivers of locus remodelling or if their functions were more intimately intermingled during B-cell ontogeny. RNAseq experiments reported that these two *cis*-enhancers are independent motors of IgH locus remodelling with no mutual *cis*-transcriptional interactions within the IgH locus and no *trans*-transcriptional interaction with the Ig light chain kappa (Igκ) locus. These results validate the hypothesis that the two major enhancers of the IgH locus are independent engines of locus remodelling; the activation of one does not depend on the activation of the other (and vice versa).

Keywords

IgH 3' Regulatory Region; 5'E_μ Enhancer; IgL Locus; Transcriptional Enhancer; Knock-out Mice; RAG-Deficient Mice

Introduction

By their impact on nuclear organisation, enhancers are master regulators of cell fate [1]. The immunoglobulin heavy chain (IgH) locus undergoes numerous changes throughout B-cell differentiation. Among them, transcription and accessibility for V(D)J recombination, class switch recombination (CSR) and somatic hypermutation (SHM) are the most notable [2]. The IgH locus carries two potent enhancers separated by 200 kb (Figure 1A). 5'E_μ and the 3' regulatory region (3'RR), located at both ends of the constant gene cluster, control locus remodelling during B-cell differentiation [2]. Several studies have reported long-range interactions between 5'E_μ and 3'RR enhancers during B-cell maturation [3-5]. The question of mutual transcriptional cross-talk between these two enhancer entities remains open. During B-cell development, the heavy and light chain (IgL) loci are poised for their VDJ and VJ rearrangements, respectively [2]. The IgH locus first rearranges with D-J segments at the pro-B-cell stage followed by V-DJ joining at the pre-B-cell stage. The Igk locus is poised for VJ rearrangements at the pre-B cell stage. A transient association (*trans*-mediated by Igk enhancer elements) between IgH and Igk loci has been demonstrated at the pre-B cell stage [6,7]. The question of a *trans*-mediated effect of IgH enhancer elements (5'E_μ and 3'RR) on Igk locus remains open. In this study we examined whether 5'E_μ and 3'RR enhancers were independent motors of locus remodelling or if their functions were more intimately intermingled during B-cell ontogeny. We thus developed ΔE_μ-RAG-deficient and Δ3'RR-RAG-deficient mice to investigate potential transcriptional cross-talk between 5'E_μ and 3'RR enhancers at the immature B-cell stage. We also analysed transcriptomic data in order to identify a putative *trans*-transcriptional effect beyond the Igk locus in IgH enhancer-deficient mice.




Figure 1: 3'RR eRNA in pro-B cells.

A: Schematic representation of the IgH locus (not to scale). V (variable), D (diversity), J (junctional) and C (constant) segments are located as well as the 5'E μ element and the 3'RR. The 3'RR contains 4 transcriptional enhancers. Three of them are encompassed in a 25kb palindromic structure. B: Detection of 3'RR eRNAs. Lower panel: *Wt* B-cell splenocytes were stimulated 2 days with 5 μ g/ml LPS (8-12 weeks old, males and females). RNAseq experiments were done after depletion of rRNA. One representative experiment out of two is reported with 3 mice per sample. Upper and medium panels: Pro-B cells of RAG-deficient (upper panel) and Δ E μ -RAG-deficient (medium panel) mice. One representative experiment out of two is reported with 4 to 5 mice per genotype.

Material and Methods

Mice

RAG-deficient mice (Janvier Labs, France), ΔE_{μ} [8] -RAG-deficient mice and $\Delta 3'RR$ [9] -RAG-deficient mice were housed in the EOPS facility of the University of Limoges (France) and procedures were conducted in agreement with European Directive 2010/63/EU on animals used for scientific purposes. The APAFiS#13855 projet was authorized by the French « Ministère de l'Education Nationale, de l'Enseignement Supérieur et de la Recherche ».

Cells for RNAseq Experiments

Femoral pro-B cells were recovered with the EasySep™ mouse B-cell isolation Kit (STEMCELL Technologies, France) designed to isolate B cells from single-cell suspensions by negative selection. Unwanted cells were targeted for removal with biotinylated antibodies directed against non-B cells and streptavidin-coated magnetic particles (RapidSpheres™). Labeled cells were separated using an EasySep™ magnet without the use of columns. Desired cells were collected into a new tube. Cells of RAG-deficient, ΔE_{μ} -RAG-deficient and $\Delta 3'RR$ -RAG-deficient mice (8-12 weeks old, males and females) were used. In another set of experiments, *wt* B-splenocytes were stimulated with 5µg/ml LPS for two days as a positive control of 3'RR eRNA detection.

RNAseq Experiments

Pro-B cells were obtained from 10 RAG-deficient, 10 $\Delta 3'RR$ -RAG-deficient mice and 10 ΔE_{μ} -RAG-deficient mice. RNA was extracted using miRNeasy kit from QIAGEN, according to the manufacturer's instructions. Two RNA pools (five samples each) were obtained for each genotype. RNA libraries were prepared using TruSeq Stranded Total RNA with Ribo-Zero Gold (Illumina), according to the manufacturer's instructions. Libraries were sequenced on a NextSeq500 sequencer, using NextSeq 500/550 High Output Kit (Illumina). Illumina

NextSeq500 paired-end 2x150nt reads were mapped with STAR release v2.4.0a versus mm10 with a gene model from Ensembl release 77 with default parameters. 10-12 RNAseq experiments were done by the genomics platform of Nice Sophia Antipolis (France). Data were deposited in Gene Expression Omnibus under the accession number GSE117449, GSE169690 and GSE169691.

Results and Discussion

Femoral pro-B cells were isolated from RAG-deficient, ΔE_{μ} -RAG-deficient and $\Delta 3'RR$ -RAG-deficient mice to investigate the potential transcriptional cross-talk between $5'E_{\mu}$ and $3'RR$ enhancers in immature B-cells. A schematic representation of the IgH locus is reported in Figure 1A. Non coding RNAs (ncRNAs) contribute to chromosomal looping.¹³ Among these ncRNAs, enhancer RNAs (eRNAs) are transcribed from DNA sequences of enhancers including the $3'RR$ and contribute to their enhancer function [14,15]. As a positive control of $3'RR$ eRNA detection, LPS-stimulated *wt* B-splenocytes were used (Figure 1B, lower panel). RNAseq experiments did not highlight any $3'RR$ eRNAs in pro-B cells of RAG and ΔE_{μ} -RAG-deficient mice (Figure 1B, upper and middle panel, respectively) confirming results from a previous study with specific RT-QPCR [16]. The absence of $3'RR$ eRNAs in pro-B cells is in agreement with studies reporting that $3'RR$ has no direct role on V(D)J recombination [17,18].

The $5'E_{\mu}$ enhancer is implicated during DJ recombination [8]. A schematic representation of the variable part of the IgH locus is shown in Figure 2A. Genomic deletion of the $5'E_{\mu}$ enhancer abrogated transcription around its location including J_H and C_{μ} transcription as well as peaks of transcription specifically found to originate from the D_{Q52} promoter (D_{4-1}) and the $5'E_{\mu}$ enhancer (known as μo and I_{μ} sense transcripts, respectively) (Figure 2B, upper and middle panels). As previously reported by Braikia and coll, [16] genomic deletion of the $3'RR$ had no effect on both μo and I_{μ} transcripts (Figure 2B, lower panel). A small increase in D antisense transcription was found in $3'RR$ -deficient mice

supporting a 3'RR-mediated transcriptional DJ silencing activity as previously suggested by Braikia and coll, [16] with RT-QPCR analysis. Together with the absence of 3'RR eRNA (Figure 1B), these results strongly suggest no transcriptional interaction of the 3'RR with the 5'E_μ element that transcriptionally control DJ recombination at the pro-B cell stage.

A *trans*-mediated transcriptional effect of the Igκ locus (mediated by Igκ enhancer elements) on the IgH locus has been demonstrated at the pre-B cell stage [6,7]. We next explored the potential transcriptional cross-talk between IgH enhancers on the Igκ locus (schematized in Figure 3A) in pro-B cells. Deletion of 5'E_μ did not affect Igκ locus transcription (Figure 3B, middle panel). Deletion of 3'RR enhancers slightly affected Igκ locus sense and antisense transcription (Figure 3B, lower panel) but this effect was weak and its relevance (if any) remains obscure. In mature B-cells, deletion of 3'RR and 5'E_μ had no *trans* effect on Igκ transcription in LPS-stimulated splenocytes (Figure 3C, middle and lower panels, respectively). These results are expected since no close association between the Igκ and IgH loci has been reported in mature B-cells.




Figure 2: Influence of 5'E $_{\mu}$ and 3'RR deletion on pro-B cell DJ transcription. A: Schematic representation of the IgH locus (not to scale). V, D, J and C $_{\mu}$ segments are located as well as the 5'E $_{\mu}$ element and the 3'RR. B: D, J, 5'E $_{\mu}$, and C $_{\mu}$ sense and antisense transcription in pro-B cells of RAG-deficient, ΔE_{μ} -RAG-deficient and $\Delta 3'RR$ -RAG-deficient mice. One representative experiment out of two is reported with 4 to 5 mice per genotype. Locations of D $_{4-1}$ (also known as D $_{Q52}$) and I $_{\mu}$ promoters are indicated.




Figure 3: Influence of 5'E μ and 3'RR deletion on Ig κ transcription. A: Schematic representation of the Ig κ locus (not to scale). B: Ig κ transcription in pro-B cells of RAG-deficient, $\Delta E\mu$ -RAG-deficient and $\Delta 3'RR$ -RAG-deficient mice. One representative experiment out of two is shown with 4 to 5 mice per genotype. C: Ig κ transcription in B-cell splenocytes from *wt*, $\Delta E\mu$ and $\Delta 3'RR$ mice. B-cell splenocytes were stimulated 2 days with 5 $\mu\text{g/ml}$ LPS (8-12 weeks old, males and female). One representative experiment out of two is shown with 3 mice per sample.

Concluding Remarks

Studies have reported the role of 5'E $_{\mu}$ and 3'RR enhancers during B-cell fate and maturation. Deletion of the 5'E $_{\mu}$ enhancer markedly lowered B-cell V(D)J recombination with no effect on SHM and CSR [8,19]. In contrast, deletion of the 3'RR enhancer affected B-cell fate, [20] SHM, [21] and CSR [9,11,12]. If studies reported the independent roles of 5'E $_{\mu}$ and 3'RR in B-cell maturation, few data were available concerning their synergy, cooperation and transcriptional cross-talk. This study shows that despite strong physical interactions during IgH locus DNA looping at the pro-B cell stage, [4,5] 5'E $_{\mu}$ and 3'RR enhancers are independent drivers of locus remodelling. Their function is not intimately intermingled and their optimal activation does not require physical contact with each other. Analysis of IgH locus transcription in 5'E $_{\mu}$ - and 3'RR-deficient mice reveals unilateral dependence of this pair of enhancers: 5'E $_{\mu}$ and 3'RR autonomously in immature B-cells and mature B-cells, respectively. These results obtained with knock-out (KO) mice are in agreement with previous results obtain with 5'E $_{\mu}$ -GFP, GFP-3'RR and 5'E $_{\mu}$ -GFP-3'RR transgenics [22-24].

References

1. Qian J, Wang Q, Dose M, Pruett N, Kieffer-Kwon KR, et al. B cell super-enhancers and regulatory clusters recruit AID tumorigenic activity. *Cell*. 2014; 159: 1524-1537.
2. Pinaud E, Marquet M, Fiancette R, Péron S, Vincent-Fabert C, et al. The IgH locus 3' regulatory region: pulling the strings from behind. *Adv Immunol*. 2011; 110: 27-70.
3. Wuerffel R, Wang L, Grigera F, Manis J, Selsing E, et al. S-S synapsis during class switch recombination is promoted by distantly located transcriptional elements and activation-induced deaminase. *Immunity*. 2007; 27: 711-722.
4. Medvedovic J, Ebert A, Tagoh H, Tamir IM, Schwickert TA, et al. Flexible long-range loops in the VH gene region of the IgH locus facilitate the generation of a diverse antibody repertoire. *Immunity*. 2013; 39: 229-244.

5. Guo C, Yoon HS, Franklin A, Jain S, Ebert A, et al. CTCF-binding elements mediate control of V(D)J recombination. *Nature*. 2011; 477: 424-430.
6. Xiang Y, Zhou X, Hewitt SL, Skok JA, Garrard WT. A multifunctional element in the mouse Igk locus that specifies repertoire and Ig loci subnuclear location. *J Immunol*. 2011; 186: 5356-5366.
7. Hewitt SL, Farmer D, Marszalek K, Cadera E, Liang HE, et al. Association between the Igk and Igh immunoglobulin loci mediated by the 3' Igk enhancer induces "decontraction" of the Igh locus in pre-B cells. *Nat Immunol*. 2008; 9: 396-404.
8. Marquet M, Garot A, Bender S, Carrion C, Rouaud P, et al. The E μ enhancer region influences H chain expression and B cell fate without impacting IgVH repertoire and immune response in vivo. *J Immunol*. 2014 ; 193: 1171-1183.
9. Vincent-Fabert C, Fiancette R, Pinaud E, Truffinet V, Cogné N, et al. Genomic deletion of the whole IgH 3' regulatory region (hs3a, hs1,2, hs3b, hs4) dramatically affects class switch recombination and Ig secretion to all isotypes. *Blood*. 2010; 116: 1895-1898.
10. Saintamand A, Vincent-Fabert C, Marquet M, Ghazzoui N, Magnone V, et al. E μ and 3'RR IgH enhancers show hierarchic unilateral dependence in mature B-cells. *Sci Rep*. 2017; 7: 442.
11. Saintamand A, Rouaud P, Saad F, Rios G, Cogné M, et al. Elucidation of IgH 3' region regulatory role during class switch recombination via germline deletion. *Nature Commun*. 2015 ; 6: 7084.
12. Saintamand A, Vincent-Fabert C, Garot A, Rouaud P, Oruc Z, et al. Deciphering the importance of the palindromic architecture of the immunoglobulin heavy-chain 3' regulatory region. *Nat Comm*. 2016; 7: 10730.
13. Pefanis E, Wang J, Rothschild G, Lim J, Kazadi D, et al. RNA exosome-regulated long non-coding RNA transcription controls super-enhancer activity. *Cell*. 2015; 161: 774-789.
14. Péron S, Laffleur B, Denis-Lagache N, Cook-Moreau J, Tinguely A, et al. AID-driven deletion causes immunoglobulin heavy chain locus suicide recombination in B cells. *Science*. 2012; 336: 931-934.

15. Lam MT, Li W, Rosenfeld G, Glass CK. Enhancer RNAs and regulated transcriptional programs. *Trends Biochem Sci.* 2014; 39: 170-182.
16. Braikia FZ, Conte C, Moutahir M, Denizot Y, Cogné M, et al. Developmental switch in the transcriptional activity of a long range regulatory element. *Mol Cell Biol.* 2015; 35: 3370-3380.
17. Rouaud P, Vincent-Fabert C, Fiancette R, Cogné M, Pinaud E, et al. Enhancers located in heavy chain regulatory region (hs3a, hs1,2, hs3b and hs4) are dispensable for diversity of VDJ recombination. *J Biol Chem.* 2012; 287: 8356-8360.
18. Ghazzaoui N, Issaoui H, Saintamand A, Oblet C, Carrion C, et al. The immunoglobulin heavy chain 3' regulatory region super-enhancer controls mouse B1 B-cell fate and late VDJ repertoire diversity. *Blood Adv.* 2018; 2: 252-262.
19. Perlot T, Alt FW, Bassing CH, Suh H, Pinaud E. Elucidation of IgH intronic enhancer functions via germ-line deletion. *Proc Natl Acad Sci USA.* 2005; 102: 14362-14367.
20. Saintamand A, Rouaud P, Garot A, Saad F, Carrion C, et al. The IgH 3' regulatory region governs μ chain transcription in mature B lymphocytes and the B cell fate. *Oncotarget.* 2015; 6: 4845-4852.
21. Rouaud P, Vincent-Fabert C, Saintamand A, Fiancette R, Marquet M, et al. The IgH 3' regulatory region controls AID-induced somatic hypermutation in germinal centre B-cells in mice. *J Exp Med.* 2013; 210: 1501-1507.
22. Guglielmi L, Le Bert M, Truffinet V, Cogné M, Denizot Y. Insulators to improve expression of a 3'IgH LCR-driven reporter gene in transgenic mouse models. *Biochem Biophys Res Commun.* 2003; 307: 466-471.
23. Guglielmi L, Le Bert M, Comte I, Dessain ML, Drouet M, et al. Combination of 3' and 5' IgH regulatory elements mimics the B-specific endogenous expression pattern of IgH genes from pro-B cells to mature B cells in a transgenic mouse model. *Biochim Biophys Acta.* 2003; 1642: 181-190.
24. Guglielmi L, Truffinet V, Carrion C, Le Bert M, Cogné N, et al. The 5'HS4 insulator element is an efficient tool to analyse the transient expression of an Em-GFP vector in a transgenic mouse model. *Transgenic Res.* 2005; 14: 361-364.