



HAL
open science

TREK channel activation suppresses migraine pain phenotype

Pablo Ávalos Prado, Arnaud Landra-Willm, Clément Verkest, Aurore Ribera, Anne-Amandine Chassot, Anne Baron, Guillaume Sandoz

► **To cite this version:**

Pablo Ávalos Prado, Arnaud Landra-Willm, Clément Verkest, Aurore Ribera, Anne-Amandine Chassot, et al.. TREK channel activation suppresses migraine pain phenotype. *iScience*, 2021, 24 (9), pp.102961. 10.1016/j.isci.2021.102961 . hal-03375300

HAL Id: hal-03375300

<https://hal.science/hal-03375300>

Submitted on 12 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TREK channel activation suppresses migraine pain phenotype

Pablo Ávalos Prado^{1,2}, Arnaud Landra-Willm^{1,2}, Clément Verkest^{1,2,3}, Anne-Amandine Chassot^{1,2}, Anne Baron^{2,3} and Guillaume Sandoz^{1,2,4} *

Affiliations :

¹Université Côte d'Azur, CNRS, INSERM, iBV, France

²Laboratories of Excellence, Ion Channel Science and Therapeutics, Nice, France

³Universite Côte d'Azur, CNRS, Institut de Pharmacologie Moléculaire et Cellulaire, Valbonne, France

⁴Lead contact

*Correspondence to: sandoz@unice.fr

1 **ABSTRACT**

2
3 Activation and sensitization of trigeminal (TG) ganglia sensory neurons, leading to the release
4 of pro-inflammatory peptides such as calcitonin gene-related peptide (CGRP), are likely a key
5 component in migraine-related headache induction. Reducing TG neuron excitability represents
6 therefore an attractive alternative strategy to relieve migraine pain. Here by using pharmacology
7 and genetic invalidation *ex vivo* and *in vivo*, we demonstrate that activating TREK1- and
8 TREK2- Two-Pore-Domain potassium (K_{2P}) channels inhibits TG neuronal firing sufficiently
9 to fully reverse the migraine-like phenotype induced by NO-donors in rodents. Finally,
10 targeting TREK is as efficient as treatment with CGRP antagonists, which represents one of the
11 most effective migraine therapies. Altogether, our results demonstrate that inhibiting TG
12 excitability by pharmacological activation of TREK channels should be considered as an
13 alternative to the current migraine treatment.

14
15
16 **KEYWORDS**

17
18 Allodynia / Pain / K_{2P} channels / Migraine / Neuronal excitability / Trigeminal neurons.

1 INTRODUCTION

2 Migraine is considered one of the most disabling conditions worldwide, affecting
3 around 15% of the global population (Burch et al., 2015). It manifests as a unilateral throbbing
4 headache and is accompanied by multiple symptoms such as nausea, vomiting, photophobia,
5 phonophobia or cutaneous allodynia. It is generally assumed that migraine attacks start with the
6 activation of sensory neurons in the trigeminal ganglia (TG) (Strassman et al., 1996).
7 Hyperexcitability of TG fibers innervating meninges induces the release of neuropeptides (*e.g.*
8 substance P and the calcitonin gene-related peptide CGRP) that triggers vasodilatation of
9 surrounding blood vessels and local neurogenic inflammation, leading to migraine (Frederiksen
10 et al., 2019; Goadsby et al., 2017; Khan et al., 2019). Targeting the pro-inflammatory peptides
11 secreted by TG neurons such as CGRP constitutes one of the most effective current strategy to
12 treat migraine (Edvinsson et al., 2018). However, about 50% of the patients do not respond to
13 this treatment and suffer from secondary effects (joint pain, dizziness, constipation, flu-like
14 symptoms) and alternative patient care needs to be developed.

15 Expression of several K_{2P} channel subunits has been detected in nociceptive dorsal
16 root ganglions and trigeminal neurons (Alloui et al., 2006; Yamamoto et al., 2009). We
17 recently found that the human mutation of K_{2P} -TRESK channel, “TRESK-MT”, related to
18 migraine, inhibits K_{2P} -TREK1 and TREK2 channels and sufficiently increases sensory TG
19 neuronal excitability to generate a migraine-like phenotype (Royal et al., 2019). These results
20 demonstrate the importance of TREK1 and TREK2 channels in the regulation of TG
21 excitability.

22 As the current most effective drugs against migraine target the neuropeptide CGRP
23 released by TG neurons or its receptor (Edvinsson *et al.*, 2018), it is attractive to act on TG
24 excitability to prevent CGRP release and, consequently, migraine. We therefore made the
25 hypothesis that enhancing TREK1 and TREK2 activity would reduce TG excitability, prevent
26 the release of pro-inflammatory peptides and therefore suppress migraine pain. We show that,
27 at a cellular level, ML67-33, a TREK1/2 agonist (Bagriantsev et al., 2013), reduces TG sensory
28 neuronal excitability, explaining its “anti-migraine” properties. Using behavioral tests in wild-
29 type and double knock-out mice for *Trek1/Trek2* (*Trek1^{-/-}-Trek2^{-/-}* mice), we demonstrate that
30 ML67-33 reverses the NO donor-induced migraine-like phenotype. This reversion exhibits a
31 similar potency to BIBN4096, a selective CGRP antagonist used against migraine (Olesen et
32 al., 2004). Finally, we demonstrate that ML67-33 completely reversed the NO donor-induced

1 facial allodynia in rats, which is a direct index of TG excitability. Therefore, targeting TREK
2 channels to inhibit TG firing and prevent CGRP release underlying migraine should therefore
3 be considered when developing new molecules to treat migraine.

4 5 **RESULTS**

6 **TREK1 and TREK2 activation by ML67-33 reverses the migraine-like cutaneous** 7 **allodynia in mice**

8 To investigate whether targeting TREK1 and TREK2 activity would treat migraine, we
9 first tested whether the TREK agonist ML67-33 (Bagriantsev *et al.*, 2013) specifically activated
10 TREK1 and TREK2 channels. We validated that ML67-33 selectively activated both TREK1
11 homodimer and TRESK-TREK1 heterodimer currents whereas TRESK alone was not activated
12 by ML67-33 (**Figure S1**). This shows that ML67-33, by binding to only one channel moiety,
13 can agonize the TREK channels, making it the molecule of choice for studying TREK1/2 homo-
14 and heterodimer channel function.

15 Once validated, we next monitored the effects of ML67-33 in a well-described model
16 of chronic migraine in mice in which migraine is induced by nitric oxide (NO) donors (Bates
17 *et al.*, 2010) such as isosorbide dinitrate (ISDN) (Dallel *et al.*, 2018; Royal *et al.*, 2019; Verkest
18 *et al.*, 2018). We induced and evaluated cutaneous allodynia, a quantifiable marker of migraine,
19 after four consecutive days of ISDN injections in wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice by
20 measuring paw withdrawal mechanical threshold using a dynamic von Frey aesthesiometer
21 before each injection (**Figure 1A**). We first measured the consequences of ISDN injection on
22 the mechanical threshold. In wild-type animals, we observed a significant decrease of this
23 threshold when we compared the first and last measurements (4.04 ± 0.06 g vs 2.87 ± 0.04 g, P
24 < 0.001). *Trek1^{-/-}-Trek2^{-/-}* mice already exhibited a low mechanical threshold, as previously
25 reported (Royal *et al.*, 2019) (4.04 ± 0.06 g vs 3.11 ± 0.08 g for wild-type and *Trek1^{-/-}-Trek2^{-/-}*
26 mice before ISDN i.p., respectively, $P > 0.001$) (**Figure 1B**).

27 We then compared the effect of ML67-33 with the effect of BIBN4096, a CGRP
28 antagonist used for migraine treatment (Olesen *et al.*, 2004), and challenged the capacity of
29 ML67-33 to increase the lowered mechanical threshold induced by ISDN injections. For this
30 purpose, wild-type or *Trek1^{-/-}-Trek2^{-/-}* mice were classified into three groups (for each),
31 following a series of two different injections spaced four hours in time (Figure 1D, F). The first

1 group was first injected with a vehicle solution (saline + DMSO 0.1%) then with BIBN4096.
2 The second group was first injected with BIBN4096 then with the vehicle solution. The third
3 group was first injected with ML67-33 and then BIBN4096.

4 The efficacy of BIBN4096 was confirmed in both wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice
5 from the first group, since the second injection with BIBN4096 strongly increased their
6 mechanical threshold compared to the first injection with the vehicle solution (2.92 ± 0.10 g vs
7 3.47 ± 0.11 g and 2.79 ± 0.13 g vs 3.28 ± 0.26 g, for wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice,
8 respectively, $P > 0.2$ for both conditions) (**Figures 1D and F**, black graphs). In the second
9 group, the mechanical threshold rapidly increased in both wild-type and *Trek1^{-/-}-Trek2^{-/-}*
10 animals after the first injection with BIBN4096 (2.89 ± 0.07 g vs 3.61 ± 0.11 g, $P < 0.01$; and
11 2.87 ± 0.18 g vs 3.87 ± 0.17 g, $P < 0.001$, for wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice, respectively)
12 (**Figures 1C and E**, blue traces), and exhibited no additive effects following the second
13 injection with the vehicle solution (3.61 ± 0.11 g vs 3.85 ± 0.17 g and 3.87 ± 0.17 g vs $3.82 \pm$
14 0.18 g, for wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice, respectively, $P < 0.01$ for both conditions)
15 (**Figures 1D and F**, blue graphs). Nevertheless, wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice from the
16 third group responded differently: whereas the first injection of ML67-33 increased the
17 mechanical threshold of wild-type animals compared to the control (2.78 ± 0.06 g vs $3.92 \pm$
18 0.18 , $P < 0.001$) (**Figure 1C**, red trace), it had no significant impact on *Trek1^{-/-}-Trek2^{-/-}* mice
19 (2.69 ± 0.11 g vs 3.00 ± 0.16 , $P > 0.1$) (**Figure 1E**, red trace). Interestingly, the second injection
20 of BIBN67-33 had no additive effect on the threshold in wild-type mice after the first injection
21 (3.92 ± 0.18 g vs 3.65 ± 0.19 , $P > 0.1$) (**Figure 1D**, red graph), while it increased the threshold
22 in *Trek1^{-/-}-Trek2^{-/-}* mice (3.00 ± 0.16 g vs 3.61 ± 0.18 , $P < 0.05$) (**Figure 1F**, red graph).

23 Together, these results indicate that either antagonizing CGRP release using BIBN4098
24 or specifically activating TREK1/2 using ML67-33 similarly reverses from NO-donors-induced
25 allodynia related to migraine in mice.

26

27 **ML67-33 increases TREK1 and TREK2 currents in TG sensory neurons**

28 To investigate whether the ML67-33 agonist is specifically regulating TREK1/2
29 currents in sensory neurons to reduce allodynia, we studied its effects on small and medium-
30 sized TG sensory neurons either from wild-type or double KO animals. These neurons are
31 classically divided within two populations, according to their ability to bind the plant lectin

1 isolectin B4 (IB4) from *Griffonia simplicifolia* (Stucky and Lewin, 1999). Both IB4⁺ and IB4⁻
2 neurons are involved in nociception but only IB4⁻ are enriched in neuropeptides, notably CGPR,
3 known for inducing neurogenic inflammation and triggering migraine attacks (Frederiksen *et*
4 *al.*, 2019; Khan *et al.*, 2019).

5 On primary cultured TG neurons from wild-type mice, perfusion of ML67-33 increased
6 the amplitude of a sustained K⁺ current in IB4⁺ and IB4⁻ neurons (3.18 ± 0.46 pA/pF and 7.66
7 ± 1.80 and pA/pF before and after exposure to ML67-33, respectively for IB4⁺ neurons, $P <$
8 0.05 ; 2.19 ± 0.30 pA/pF and 8.12 ± 1.62 pA/pF before and after exposure to ML67-33,
9 respectively for IB4⁻ neurons, $P < 0.01$, at -25 mV) (**Figures 2A and 2B**). However, no
10 significant difference was detected in the currents obtained from *Trek1*^{-/-}-*Trek2*^{-/-} TG neurons
11 before and after treatment with the TREK agonist (4.71 ± 1.75 pA/pF and 5.05 ± 0.76 and
12 pA/pF before and after exposure to ML67-33, respectively for IB4⁺, $P > 0.3$; 2.77 ± 0.31 pA/pF
13 and 3.60 ± 0.29 pA/pF before and after exposure to ML67-33, respectively for IB4⁻, $P > 0.05$,
14 at -25 mV) (**Figures 2C and 2D**). Therefore, ML67-33 specifically activates TREK1 and/or
15 TREK2 currents in TG sensory neurons.

16

17 **TREK current activation reduces intrinsic excitability in TG neurons related to migraine**

18 We next determined the functional consequences of the TREK1-TREK2 current on TG
19 sensory neuron excitability. First, our results showed that *Trek1* and *Trek2* genetic deletion
20 increased both IB4⁺ and IB4⁻ neurons excitability (**Figures 3 A-D**, white points and grey traces;
21 1.70 ± 0.33 Hz vs 5.12 ± 1.52 Hz for IB4⁺ from wild-type and *Trek1*^{-/-}-*Trek2*^{-/-}, respectively, P
22 < 0.05 ; 5.12 ± 1.52 Hz vs 10.89 ± 1.57 Hz for IB4⁻ from wild-type and *Trek1*^{-/-}-*Trek2*^{-/-}
23 respectively, $P < 0.01$, for a 150 pA current injection stimulus), indicating that TREK1/2
24 activation is required to reduce sensory neuron excitability.

25 Second, in wild-type IB4⁺ and IB4⁻ sensory neurons, the TREK current increased by
26 ML67-33 led to a non-significant decrease of the spike frequency for IB4⁺ neurons (1.70 ± 0.33
27 Hz vs 1.20 ± 0.36 Hz, $P > 0.2$) (**Figure 3A**) and an important reduction of the spike frequency
28 in IB4⁻ neurons (4.80 ± 0.90 Hz vs 1.00 ± 0.36 Hz, $P > 0.001$) (**Figure 3B**). The ML67-33-
29 induced firing inhibition might not be detected in IB4⁺ neurons because at rest their firing rate
30 is already low (Guo and Cao, 2014; Guo *et al.*, 2014). The reduction of the IB4⁻ excitability is
31 specific to the activation of TREK channels since ML67-33 was not effective in sensory

1 neurons from *Trek1^{-/-}-Trek2^{-/-}* (IB4⁺: 5.12 ± 1.51 Hz vs 4.50 ± 1.19 Hz before and after ML67-
2 33 perfusion, respectively, P > 0.3; IB4⁻: 10.89 ± 1.57 Hz and 9.67 ± 3.03 Hz before and after
3 ML67-33 perfusion, respectively; P > 0.6) (**Figures 3C and 3D**). Furthermore, the rheobase
4 (*i.e.* the lowest current intensity necessary for triggering action potentials (APs) was
5 significantly enhanced by ML67-33 only in wild-type sensory neurons (**Figure S2**). Together,
6 this indicates that activating TREK1/2 using ML67-33 specifically reduces IB4⁻ neuronal
7 excitability.

8 Excitability is closely linked to the resting membrane potential (RMP). Because K_{2P}
9 channels serve as a hub for RMP maintenance at negative values (Enyedi and Czirják, 2010),
10 we evaluated the effect of ML67-33 on this parameter. We found that ML67-33 induced
11 hyperpolarization of TG neurons from wild-type animals (**Figures 3E and S3**, -51.32 ± 2.40
12 mV and -62.13 ± 2.84 mV, before and after perfusion of ML67-33, respectively for IB4⁺, P <
13 0.01; -53.96 ± 0.91 mV and -61.46 ± 2.21 mV, before and after perfusion of ML67-33,
14 respectively for IB4⁻, P < 0.01) mainly through TREK1 and TREK2 activation since the slight
15 RMP decrease observed for *Trek1^{-/-}-Trek2^{-/-}* TG neurons was not statistically significant (-49.94
16 ± 1,42 mV and -53.94 ± 2.28 mV before and after perfusion of ML67-33, respectively for IB4⁺,
17 P > 0.2; -45.70 ± 1,12 mV and -50.70 ± 1.81 mV before and after perfusion of ML67-33,
18 respectively for IB4⁻, P > 0.06). In addition, we noticed that the basal value of the RMP of
19 peptidergic neurons from *Trek1^{-/-}-Trek2^{-/-}* animals was statistically higher than those from wild-
20 type mice (-45.70 ± 1,12 mV vs -53.96 ± 0.91 mV for *Trek1^{-/-}-Trek2^{-/-}* and wild-type peptidergic
21 neurons, respectively, P < 0.001) (**Figure 3G**). These results confirm that, at rest, TREK1 and
22 TREK2 control the RMP basal value.

23 Altogether, these data show that TREK1 and TREK2 play a crucial role in the
24 excitability of TG neurons, and particularly of peptidergic IB4⁻ nociceptive neurons, by
25 determining the threshold for the generation of APs and the spike frequency and by notably
26 setting up the RMP at values close to the K⁺ equilibrium potential. Thus, ML67-33 increases
27 TREK current to reduce the TG excitability, leading to a decrease of the migraine phenotype
28 observed in wild-type animals treated with NO-donors.

29

30 **Activation of TREK1 and TREK2 abrogates migraine-like facial allodynia in rats**

31 To validate that injection of ML67-33 decreases TG sensory neuronal excitability and
32 suppresses NO-migraine phenotype *in vivo*, we conducted behavioral experiments in rats. This

1 model allows to measure the mechanical allodynia of the face which is related to TG excitability
2 and therefore constitutes one of the most reliable and quantifiable readout of TG excitability
3 underlying migraine phenotype (Harris et al., 2017; Pradhan et al., 2014; Royal *et al.*, 2019).
4 The protocol used to induce chronic migraine was similar to the one used in mice despite the
5 use of the facial sensitivity instead of the paw withdrawal threshold as a readout. After a week
6 of habituation, the animals were tested for facial mechanical sensitivity with von Frey filaments.
7 Migraine-like phenotype was induced by daily intraperitoneally (i.p.) injections of ISDN (10
8 mg/kg) for four consecutive days and facial mechanical threshold was measured before each
9 injection over the four days. At day 5, either ML67-33, BIBN4096 or vehicle was administrated
10 and their effects were followed for 3 hours, every 30 minutes (**Figure 4A**).

11 As previously described (Verkest et al., 2018), ISDN induced a strong significant
12 reduction of the face withdrawal threshold compared to normal sensitivity ($P < 0.001$ for the
13 three groups) (**Figure 4B**). This reduction of the face withdrawal threshold is associated with
14 the development of chronic mechanical allodynia that reflects the induction of trigeminal
15 hyperexcitability and migraine (Royal *et al.*, 2019). As shown in **Figure 4C**, both ML67-33
16 and BIBN4096 treatments increased in a similar manner the mechanical threshold previously
17 lowered by ISDN injections (5.7 ± 0.77 g vs 6.7 ± 0.39 g for rats injected with ML67-33 and
18 BIBN4096 respectively $P > 0.6$) (**Figures 4C and 4D**).

19 Thus, in rodent models of migraine, activation of TREK1 and TREK2 channels reverses
20 from chronic allodynia related to migraine pain as efficiently as antagonizing CGRP with
21 BIBN4096.

22

1 DISCUSSION

2 Migraine is a neurological disease caused by the combination of environmental,
3 hormonal and genetic components. Genome-wide association studies have revealed the
4 importance of genetic mutations in ion channels involved in synaptic transmission in migraine
5 predisposition (Gormley et al., 2016; Nyholt et al., 2008). Particularly, K_{2P} channels such as
6 TRESK were shown to play a fundamental role in this disease. The frameshift mutation
7 TRESK-MT perfectly segregates with migraine phenotype in large pedigree (Andres-Enguix et
8 al., 2012; Lafrenière et al., 2010). The direct causality linking TRESK-MT to migraine was
9 recently demonstrated by showing that using CRISPR-Cas9 to fix the MT mutation restores a
10 normal nociceptor excitability (Pettingill et al., 2019) and that the TRESK-MT mutation
11 generated altered proteins which affect TREK1 and TREK2 channel function. The TRESK-MT
12 fragment binds and inhibits TREK1 and TREK2 in TG sensory neurons, leading to neuronal
13 hyperexcitability and migraine (Royal *et al.*, 2019). This makes TREK1 and TREK2 good
14 candidates for regulating TG excitability and subsequently migraine.

15 To analyze the potential of modulating TREK channel activity as a novel target to
16 treat this disease, we combined a pharmacological approach using the TREK agonist ML67-33
17 with mice models deficient for both *Trek1* and *Trek2*. We found that i) in wild-type mice,
18 ML67-33 reverses from NO-donors-induced allodynia and is as efficient as the CGRP
19 antagonist BIBN4096, and that ii) in *Trek1*^{-/-}-*Trek2*^{-/-} mice, ML67-33 has no effect. This shows
20 that reduction of allodynia induced by ML67-33 is due to TREK activation. Furthermore,
21 BIBN4096 and ML67-33 do not trigger an additive effect in wild-type mice, while BIBN4096
22 alone suppresses the allodynia induced by *Trek1/2* genetic deletion. Together, these observation
23 suggests that deleting *Trek1/2* promotes an increase of CGRP release that generates a migraine-
24 like phenotype whereas that their activation reduces CGRP release and suppresses migraine.

25 We confirmed this model by using a rat model allowing to measure the mechanical
26 allodynia pain threshold of the face, which is directly linked to TG neuronal excitability and
27 represents a good index of migraine state (Harris *et al.*, 2017; Pradhan *et al.*, 2014; Royal *et al.*,
28 2019). Accordingly, rats injected with NO-donors recovered from induced allodynia after either
29 BIBN4096 or ML67-33 treatment in a similar manner confirming the TREK1 and TREK
30 activation may prevent CGRP release to suppress migraine phenotype. At the cellular level,
31 activation of TREK1 and TREK2 dramatically decreased the excitability of TG sensory neurons
32 by notably lowering their RMP value. This reduction of neuronal excitability by ML67-33, only

1 observed in wild-type animals, confirms the role of TREK channels in excitability regulation
2 and consequently in pain transmission.

3 In conclusion, our results provide evidence of the key role of TREK1 and TREK2
4 channels in migraine induction by regulating TG excitability. Whereas their genetic
5 invalidation induces a neuronal hyperexcitability leading to migraine-like phenotype, their
6 activation suppresses NO-donor induced-migraine phenotype as efficiently as current anti-
7 migraine drugs targeting neuropeptide release. Therefore, targeting TREK channel intrinsic
8 activity to reduce TG neuron excitability and CGRP release should be considered as an
9 alternative strategy to treat migraine.

10

11 **ACKNOWLEDGMENTS**

12 We thank Pr Daniel L. Minor for providing us the selective activator of TREK channels ML67-
13 33. We thank the Dr Franck Chatelain and Dr Florian Lesage for providing us the TREK1^{-/-}-
14 TREK2^{-/-} mice. This work was supported by a grant to G.S. by the Fondation pour la Recherche
15 Medicale (Equipe labellisée FRM 2017, FRM DEQ20170336753) the Agence Nationale pour
16 la Recherche (AT2R-TRAAK-Bioanalgesics ANR-17-CE18-0001 and ANR-17-ERC2-0023),
17 the Laboratory of Excellence “Ion Channel Science and Therapeutics” (grant ANR-11-LABX-
18 0015-01) and the French government, through the UCAJEDI Investments in the Future project
19 managed by the National Research Agency (ANR) with the reference number ANR-15-IDEX-
20 01.

21

22

23

24 **DECLARATION OF INTERESTS**

25

26 The authors declare no competing interest

27






28

29

30

1 **FIGURE LEGENDS**

2



3

4

1 **Figure 1. ML67-33 reverses NO induced migraine-like-phenotype through TREK channel**
2 **activation.** (A) Migraine behavioral test timetable. (B) Effect of repeated administration of
3 systemic ISDN NO-donors (10 mg/kg i.p.) on the mechanical threshold (g) and the development
4 of cutaneous allodynia related to migraine in wild-type and *Trek1^{-/-}-Trek2^{-/-}* mice. Two-way
5 ANOVA and Sidak post hoc vs wild-type (* p < 0.5, *** p < 0.001); One-way ANOVA and
6 Dunett's post hoc vs pre-treatment day 1 (# p < 0.5, ### p < 0.001). (C) Time-course of
7 cutaneous mechanical threshold (g) after injection of a vehicle solution (saline + DMSO 0.1%
8 i.p.), BIBN4096 (1 mg/kg i.p.) and ML67-33 (1 mg/kg i.p.) in wild-type mice. Two-way
9 ANOVA and Dunett's post hoc vs vehicle (* p < 0.5, ** p < 0.01, *** p < 0.005); One-way
10 ANOVA and Dunett's post hoc vs time -0.5 h (# p < 0.5, ## p < 0.01, ### p < 0.005). (D) Bar
11 graph representing mechanical threshold (g) of the three groups of animals in this study after
12 2h of the second injection of BIBN4096, vehicle and BIBN4096 respectively. Student's t-test
13 (* p < 0.5, ** p < 0.01). (E-F) Same as C-D for *Trek1^{-/-}-Trek2^{-/-}* mice.

14

15

16

17


18

19

20

21


22



1

2 **Figure 2. ML67-33 potentiates TREK channels in TG neurons.** (A-D) Representative
 3 current traces of IB4⁺ and IB4⁻ neurons of small TG neurons obtained from wild-type (A and
 4 B) and *Trek1*^{-/-}-*Trek2*^{-/-} (C and D) mice, before and during perfusion of ML67-33 (10 μM).
 5 Insets Bar graphs representing current densities at -25 mV (pA/pF). Currents were elicited by
 6 voltage-ramps (from -25 to -135 mV, 300 ms duration). Paired t-test (* p < 0.05, ** p < 0.01).
 7 Mean ± SEM.

8



2

3

4 **Figure 3. ML67-33 reduces TG neuron excitability through TREK channels activation.**


5 (A-D) Input-output plots of spike frequency in response to 1-s depolarizing current injection in
 6 small IB4⁺ and IB4⁻ small TG neurons from wild-type (A and B) and *Trek1^{-/-}-Trek2^{-/-}* (C and
 7 D) mice before and after perfusion with ML67-33 (10 μ M). At right, Representative traces of

1 action potentials generated by incremental depolarizing current injections in small-diameter TG
2 neurons. Two-way ANOVA (* $p < 0.5$, ** $p < 0.01$, *** $p < 0.001$). (E) Representative RMP
3 (mV) recordings from TG neurons exposed to ML67-33. (F-G) Bar graphs showing the basal
4 RMP value of IB4⁺ (F) and IB4⁻ (G) TG neurons. Mann-Whitney test (*** $p < 0.001$). Mean \pm
5 SEM.

6

7


8



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

Figure 4. ML67-33 abrogates mechanical facial allodynia related to migraine. (A) Migraine behavioral test timetable. Green arrows represent the injection of ISDN, a known migraine trigger. Magenta arrows represent the injection of either saline or BIBN4096 or ML67-33 solution injection. Blue arrows represent the measurement of the facial withdrawal mechanical threshold (B) Effect of repeated administration of systemic ISDN (10 mg/kg i.p.) on the occurrence and development of cutaneous chronic allodynia (facial mechanical threshold, g) in rats before treatment with different drugs. One-way ANOVA and Sidak post hoc vs pre-treatment day 1 (* $p < 0.5$, ** $p < 0.01$, *** $p < 0.001$). (C) ML67-33 is as efficient as the CGRP receptor antagonist BIBN4096 to reverse migraine like phenotype: time-evolution of von Frey delta facial withdrawal thresholds in rats injected with a vehicle solution (saline + DMSO 0.1% i.p.), BIBN4096 (1 mg/kg i.p.) and ML67-33 (1 mg/kg i.p.). Two-way ANOVA and Sidak post hoc vs vehicle treatment (* $p < 0.05$, *** $p < 0.001$); One-way ANOVA and Dunett's post hoc vs time - 0.5 h (# $p < 0.5$, ## $p < 0.01$, ### $p < 0.001$). (D) Variation of the face withdrawal threshold before and 3 hours after compound administration. Kruskal-Wallis and Student-Newman-Keuls post hoc (** $p < 0.01$). Mean \pm SEM.


1 **SUPPLEMENTAL FIGURES AND LEGENDS**



2

3 **Figure S1. ML67-33 activates TREK1 homodimers and TRESK/TREK1 heterodimers**
 4 **but not TRESK homodimers, Related to Figure 2 and 3.** (A-C) Representative current traces
 5 obtained from HEK293T cells expressing either TREK1 (A) or TRESK (B) homodimers or
 6 TRESK/TREK1 (C) heterodimers before and after application of ML67-33 (10 μ M). Currents
 7 were elicited by voltage-ramps (from -100 to +100 mV from a holding potential at -80 mV, 1s
 8 duration). Insets show a summary of current densities obtained at 0 mV. Paired t-test (** $p <$
 9 0.01). (D) Bar graph showing the fold increase after perfusion with ML67-33. Kruskal-Wallis
 10 and Student-Newman-Keuls post hoc (***) $p <$ 0.001). Mean \pm SEM.

11




1

2 **Figure S2. ML67-33 increases rheobase of TG neurons through activation of TREK1 and**
 3 **TREK2 channels, Related to Figure 3.** (A-D) Bar graphs showing representing rheobase
 4 obtained from IB4⁺ and IB4⁻ TG neurons obtained from wild-type (A and B) and *Trek1*^{-/-}-*Trek2*^{-/-}
 5 (C and D) mice before and after perfusion with ML67-33 (10 μM). Rheobase was determined
 6 by incremental depolarizing current injections in steps of + 10 pA in small-diameter TG
 7 neurons. Paired t-test (** p < 0.01). Mean ± SEM.

8

9



1
2
3 **Figure S3. ML67-33 hyperpolarizes TG neurons from wild-type mice through activation**
4 **of TREK channels, Related to Figure 3.** (A-B) Bar graphs showing representing the RMP
5 value obtained from small IB4⁺ and IB4⁻ TG neurons before and after perfusion with ML67-33
6 (10 μ M) from wild-type (A) and *Trek1*^{-/-}-*Trek2*^{-/-} (B) mice. Paired t-test (** p < 0.01, *** p <
7 0.001). Mean \pm SEM.

1 **STAR★Methods**

2
3
4 RESOURCE AVAILABILITY

5
6 **Lead Contact**

7 Further information and requests for resources and reagents should be directed to and will be
8 fulfilled by the Lead Contact, Guillaume Sandoz (sandoz@unice.fr).

9
10 **Materials Availability**

11 This study did not generate new unique reagents.

12
13 **Data and Code Availability**

14 Source data for the main and supplemental figures in the paper is available at Mendeley
15 doi:10.17632/4vr86gr5ch.1

16
17
18 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**

19 **Mice**

20 All mouse experiments were conducted according to national and international
21 guidelines and have been approved by the local ethical committee and authorized the French
22 Ministry of Research according to the European Union regulations and the Directive
23 2010/63/EU (APAFIS#21943-2019073017246158). The C57BL/6J breeders were maintained
24 on a 12 h light/dark cycle with constant temperature (23–24 °C), humidity (45%–50%), and
25 food and water *ad libitum* at the animal facility of Institut de Biologie de Valrose. Knock-out
26 mice lacking *Trek1* and *Trek2* were generated as described (Guyon et al., 2009). Null mutations
27 were backcrossed against the C57BL/6J inbred strain for more than 10 generations prior to
28 establishing the breeding cages to generate subjects for this study. Age- and sex-matched
29 C57BL/6J wild-type mice, aged 9-12 weeks, were obtained from Charles River Laboratories
30 (Wilmington, MA). Behavioral experiments were performed on 7 to 13-weeks old male mice
31 weighting 20-30 g.

1 **Rats**

2 Experiments were performed on 6 to 9 weeks old male Sprague Dawley rats (Janvier
3 Labs) weighing 250 to 400 g. Animals were housed in a 12 hours light-dark cycle with food
4 and water available *ad libitum*. Animal procedures were approved by the Institutional Local
5 Ethical Committee and authorized by the French Ministry of Research according to the
6 European Union regulations and the Directive 2010/63/EU (Agreements 01550.03). Animals
7 were sacrificed at experimental end points by CO₂ euthanasia.

8 **Primary culture of mouse TG neurons**

9 Trigeminal ganglion tissues were collected from postnatal day 2-8 mice of both sex in
10 L15 Leibovitz medium (Sigma) and treated with 2 mg/ml collagenase type II (Worthington)
11 and BSA (Euromedex) for 2 hours, followed by 2.5 mg/ml trypsin (Sigma) for 15 min. Neurons
12 were dissociated in DMEM/F12 medium (Gibco) by triturating with fire-polished and coated
13 glass pipettes and seeded on poly-L-lysine (Sigma) coated coverslips. The DMEM-based
14 culture medium contained 10% fetal bovine serum (Dutscher) and 2 mM Glutamine (Gibco).

15 **HEK293T cells**

16 HEK293T (ATCC, #CRL11268) cells were maintained in DMEM (Gibco)
17 supplemented with 10% FBS (Dutscher) in 35 mm dishes. At 70%–80% confluency they were
18 transiently co-transfected using the calcium phosphate method with a total amount of 3.5 µg of
19 DNA and seeded on 18 mm diameter poly-L-lysine (Sigma) coated glass coverslips in 12 well
20 plates.

21

22 **METHOD DETAILS**

23 **Molecular biology and gene expression**

24 Clones used were mTREK1 (RefSeq: NP_001153322), mTREK2 (NP_084187.2) and
25 mTRESK (RefSeq: NP_997144). All channel DNA was used in the pIRES2eGFP vector.
26 Heterodimers between TREK1, TREK2 and TRESK were generated by PCR.

27

28

29

1 **Electrophysiology**

2 HEK293T electrophysiology was performed 24 – 48 h after transfection. For whole-cell
3 patch-clamp experiments, cells were recorded in a bath solution containing (in mM) 145 NaCl,
4 4 KCl, 1 MgCl₂, 2 CaCl₂ and 10 HEPES, pH 7.4. The glass pipettes (2-5 MΩ of resistance)
5 were filled with (in mM) 140 KCl, 3 MgCl₂, 5 EGTA, 10 HEPES, pH 7.3. ML67-33 was
6 perfused at 10 μM for 2-5 min in the bath solution. Whole-cell currents were elicited by voltage-
7 ramps (from -100 to +100 mV, 1 s) holding the cells at -80 mV. Current densities were measured
8 at 0 mV.

9 For TG neuron electrophysiology, medium and small sized cells (< 25 μm) were
10 recorded in an extracellular solution containing (in mM): 135 NaCl, 5 KCl, 2 CaCl₂, 1 MgCl₂,
11 5 HEPES, 10 glucose, pH 7.4. In order to distinguish between peptidergic and non-peptidergic,
12 neurons were first incubated with an isolectine-IB4 marker (Invitrogen) for 30 min at 5 μg/mL.
13 The pipette solution contained the following (in mM): 140 K-gluconate, 10 NaCl, 2 MgCl₂, 5
14 EGTA, 10 HEPES, 2 ATP-Mg, 0.3 GTP-Na, 1 CaCl₂. Neurons were excluded from analysis
15 when the RMP was higher than -40 mV. To test neuronal excitability, neurons were held at
16 RMP in current-clamp mode and injected with 1 s depolarizing current in 10 pA incremental
17 steps. The voltage-clamp ramp protocol for TREK channels consisted in voltage-ramps from -
18 25 to -135 mV during 600 ms, holding the cells at -60 mV. ML67-33 was perfused at 10 μM
19 for 2-5 min in the extracellular solution.

20 All cells were recorded at room temperature using an Axopatch 200B (Molecular
21 Devices) amplifier. Signals were filtered at 10 kHz and digitalized at 20 kHz. Cell recordings,
22 data acquisition and analysis of electrophysiology were performed using pClamp software
23 (Molecular Devices).

24 **Migraine model rodents**

25 The rodent models of NO-induced migraine were induced by *i.p.* injection of ISDN
26 (Sanofi) at 10 mg/kg, a long-lasting NO donor in both rats and mice.

27 The mice hindpaw mechanical sensitivity was evaluated with a dynamic plantar
28 aesthesiometer (Ugo Basile). Unrestrained mice were placed in twelve individual plastic boxes
29 on top of a wire surface. The mouse hindpaw was subjected to an increasing force ramp (0 to
30 7.5 g in 10 s), the paw withdrawal force threshold (g) was measured in quadriplate on each hind

1 paw, and the mean force was calculated. For 2 days, mice were habituated to repeated (every
2 30 min) measurements of hindpaw mechanical sensitivities, and basal values were determined
3 2 days before the experiments. ISDN was injected each day for 4 days, and the mechanical
4 threshold was measured each day before the next ISDN injection to follow the settlement of
5 chronic allodynia. Normal mechanical sensitivity and allodynia thresholds were taken as the
6 measurement values of the first day before treatment with ISDN and the last measurement
7 before the injection of drugs on wild-type mice, respectively. On the fifth day, BIBN4096
8 (Tocris) (1 mg/kg *i.p.*) and ML67-33 (1 mg/kg, *i.p.*) were administrated as well as their vehicle
9 solution: 0.9% saline and 0.1% DMSO (Sigma). The hindpaw mechanical sensitivity was
10 measured every 30 min before (basal value) and every 30 min after the *i.p.* injections.

11 The face mechanical sensitivity of rats was measured using calibrated von Frey
12 filaments (Bioseb, France). Unrestrained rats placed in individual plastic boxes on top of a wire
13 surface were trained over 1 week; a stimulus was applied to the periorbital area, following a
14 progressive protocol, starting with non-noxious filaments during the first days of training. The
15 face withdrawal force threshold (g) was determined by the filament evoking at least three
16 responses over five trials, starting with lower force filaments. ISDN (10 mg/Kg, Sanofi) was
17 injected each day for 4 days, and the mechanical threshold was measured each day before the
18 next ISDN injection to follow the settlement of chronic allodynia. To test the effect of drugs,
19 face mechanical sensitivity was measured every 30 min before (basal value) and for 3 h after
20 compound or vehicle injection.

21

22 QUANTIFICATION AND STATISTICAL ANALYSIS

23 Analysis of whole-cell currents was performed using ClampFit software. For analysis
24 of voltage-ramp traces cursors were set at 0 mV for HEK293T cells, to extract the average
25 current. Cursors were set at 355 ms for TG neurons. Spike frequency was determined as the
26 number of APs triggered in every current step. The rheobase was taken as the minimal current
27 injected to generate a first AP. When neurons did not trigger any AP even for current injections
28 >500 pA, the rheobase value was considered as 0 pA. Signals were filtered at 10 kHz and
29 digitalized at 20 kHz. For electrophysiology and behavioral experiments, statistical details and
30 n values can be found in the figure legends and in the main text.

31

32

1 **KEY RESOURCES TABLE**

2

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Chemicals, Peptides, and Recombinant Proteins		
BIBN4096 4096	Tocris	Cat# 4561
BSA	Euromedex	Cat# 04-100-812-C
Collagen	This paper	N/A
DMEM	Gibco	Cat# 41965-039
DMEM/F12	Sigma-Aldrich	Cat# 6421
DPBS	Gibco	Cat# 14190-169
FBS	Dutscher	Cat# S1900-500
Glutamine	Gibco	Cat# 25030-081
HEPES	Gibco	Cat# 15630-080
ISDN Risordan ®	Sanofi	N/A
Isolectin GS-IB4	Thermofischer	Cat# I21411
L15 Medium (Leibovitz)	Sigma-Aldrich	Cat# L5520
ML67-33	Tocris	Cat# 6886
Penicillin-Streptomycin	Gibco	Cat# 15140-122
Poly-L-lysine	Sigma-Aldrich	Cat# P4707
Trypsin	Sigma-Aldrich	Cat# T1763
Deposited Data		
Raw and analyzed data	This paper	doi:10.17632/4vr86gr5ch.1
Experimental Models: Cell Lines		
HEK 293T cells	ATCC	Cat#CRL11268
Mouse trigeminal ganglia cells	This paper	N/A
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Charles River Laboratories	Strain Code 027
Mouse: TREK1 ^{-/-} -TREK2 ^{-/-}	Guyon et al., 2009	N/A
Rat: Sprague Dawley	Janvier Labs	Strain RjHan:SD
Oligonucleotides		
TREK1 (forward): CATCTTCATCCTGTTTGGCTG	Sigma-Aldrich	Custom order
TREK1 (reverse): ATCATGCTCAGAACAGCTGC	Sigma-Aldrich	Custom order
TRESK-TREK1 (internal forward): TTTCGCTACCTTGGGCGGCCCTGACTTGCTGGAT	Sigma-Aldrich	Custom order
TRESK-TREK2 (internal reverse): GCAAGTCAGGGGCCGCCAAGGTAGCGAACTTCC	Sigma-Aldrich	Custom order
TREK2 (forward) AACAGTGGTTGCCATCTTCG	Sigma-Aldrich	Custom order
TREK2 (reverse) CCAGCAAAGAAGAAGGCACT	Sigma-Aldrich	Custom order
TRESK-TREK2 (internal forward): GTTTCGCTACCTTGGAAATTTCCAATCGAGACG	Sigma-Aldrich	Custom order
TRESK-TREK2 (internal reverse): CGTCTCGATTGGAAATTTCCAAGGTAGCAAAC	Sigma-Aldrich	Custom order
TRESK (forward) CTGCTTCTTTGCTGCCTG	Sigma-Aldrich	Custom order
TRESK (reverse) AAGAAGAGAGCGCTCAGGAA	Sigma-Aldrich	Custom order
Recombinant DNA		
pIRES2eGFP	Clontech	6029-1
mTREK1	Royal et al., 2019	N/A
mTREK2	Royal et al., 2019	N/A
mTRESK	Royal et al., 2019	N/A

Software and Algorithms		
Fiji/ImageJ, v1.8	NIH, Schneider et al., 2012	https://imagej.nih.gov/ij/
pCLAMP 10, pCLAMP 11	Molecular Devices	N/A
SigmaPlot v11	Systat Software Inc.	N/A
Other		
Axioplan 2 Imaging Microscope	Zeiss	https://www.microshop.zeiss.com/?s=16103145829fcb6&l=en&p=us&f=a&i=10027
Axopatch 200B amplifier	Molecular Devices	N/A
Dynamic plantar aesthesiometer	Ugo Basil	Cat#: 37450
Micromanipulator MP225	Sutter Instrument	https://www.wpi-europe.com/products/micromanipulators/motorised-manipulators/mp-225.aspx
Digidata 1550B	Molecular Devices	N/A
Perfusion	ValveLink 8.2	https://www.autom8.com/perfusion-systems-overview/valvelink8-2-controller/
Camera EMCCD iXon	Andor	https://andor.oxinst.com/products/ixon-emccd-cameras
sCMOS camera Zyla 4.2+	Andor	https://andor.oxinst.com/products/scmos-camera-series/zyla-4-2-scmos
Von Frey Filaments	Bioseb	Model: Bio-VF-M

1
2
3
4

1 REFERENCES

- 2 Alloui, A., Zimmermann, K., Mamet, J., Duprat, F., Noël, J., Chemin, J., Guy, N., Blondeau, N., Voilley, N.,
3 Rubat-Coudert, C., et al. (2006). TREK-1, a K⁺ channel involved in polymodal pain perception. *EMBO J* 25,
4 2368-2376. 10.1038/sj.emboj.7601116.
- 5 Andres-Enguix, I., Shang, L., Stansfeld, P.J., Morahan, J.M., Sansom, M.S., Lafrenière, R.G., Roy, B., Griffiths,
6 L.R., Rouleau, G.A., Ebers, G.C., et al. (2012). Functional analysis of missense variants in the TRESK
7 (KCNK18) K channel. *Sci Rep* 2, 237. 10.1038/srep00237.
- 8 Bagriantsev, S.N., Ang, K.H., Gallardo-Godoy, A., Clark, K.A., Arkin, M.R., Renslo, A.R., and Minor, D.L.
9 (2013). A high-throughput functional screen identifies small molecule regulators of temperature- and mechano-
10 sensitive K2P channels. *ACS Chem Biol* 8, 1841-1851. 10.1021/cb400289x.
- 11 Bates, E.A., Nikai, T., Brennan, K.C., Fu, Y.H., Charles, A.C., Basbaum, A.I., Ptáček, L.J., and Ahn, A.H.
12 (2010). Sumatriptan alleviates nitroglycerin-induced mechanical and thermal allodynia in mice. *Cephalalgia* 30,
13 170-178. 10.1111/j.1468-2982.2009.01864.x.
- 14 Burch, R.C., Loder, S., Loder, E., and Smitherman, T.A. (2015). The prevalence and burden of migraine and
15 severe headache in the United States: updated statistics from government health surveillance studies. *Headache*
16 55, 21-34. 10.1111/head.12482.
- 17 Dalle, R., Descheemaeker, A., and Luccarini, P. (2018). Recurrent administration of the nitric oxide donor,
18 isosorbide dinitrate, induces a persistent cephalic cutaneous hypersensitivity: A model for migraine progression.
19 *Cephalalgia* 38, 776-785. 10.1177/0333102417714032.
- 20 Edvinsson, L., Haanes, K.A., Warfvinge, K., and Krause, D.N. (2018). CGRP as the target of new migraine
21 therapies - successful translation from bench to clinic. *Nat Rev Neurol* 14, 338-350. 10.1038/s41582-018-0003-
22 1.
- 23 Enyedi, P., and Czirják, G. (2010). Molecular background of leak K⁺ currents: two-pore domain potassium
24 channels. *Physiol Rev* 90, 559-605. 10.1152/physrev.00029.2009.
- 25 Frederiksen, S.D., Haanes, K.A., Warfvinge, K., and Edvinsson, L. (2019). Perivascular neurotransmitters:
26 Regulation of cerebral blood flow and role in primary headaches. *J Cereb Blood Flow Metab* 39, 610-632.
27 10.1177/0271678X17747188.
- 28 Goadsby, P.J., Holland, P.R., Martins-Oliveira, M., Hoffmann, J., Schankin, C., and Akerman, S. (2017).
29 Pathophysiology of Migraine: A Disorder of Sensory Processing. *Physiol Rev* 97, 553-622.
30 10.1152/physrev.00034.2015.
- 31 Gormley, P., Anttila, V., Winsvold, B.S., Palta, P., Esko, T., Pers, T.H., Farh, K.H., Cuenca-Leon, E., Muona,
32 M., Furlotte, N.A., et al. (2016). Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for
33 migraine. *Nat Genet* 48, 856-866. 10.1038/ng.3598.
- 34 Guo, Z., and Cao, Y.Q. (2014). Over-expression of TRESK K(+) channels reduces the excitability of trigeminal
35 ganglion nociceptors. *PLoS One* 9, e87029. 10.1371/journal.pone.0087029.
- 36 Guo, Z., Liu, P., Ren, F., and Cao, Y.Q. (2014). Nonmigraine-associated TRESK K⁺ channel variant C110R
37 does not increase the excitability of trigeminal ganglion neurons. *J Neurophysiol* 112, 568-579.
38 10.1152/jn.00267.2014.
- 39 Guyon, A., Tardy, M.P., Rovère, C., Nahon, J.L., Barhanin, J., and Lesage, F. (2009). Glucose inhibition persists
40 in hypothalamic neurons lacking tandem-pore K⁺ channels. *J Neurosci* 29, 2528-2533.
41 10.1523/JNEUROSCI.5764-08.2009.
- 42 Harris, H.M., Carpenter, J.M., Black, J.R., Smitherman, T.A., and Sufka, K.J. (2017). The effects of repeated
43 nitroglycerin administrations in rats; modeling migraine-related endpoints and chronification. *J Neurosci*
44 *Methods* 284, 63-70. 10.1016/j.jneumeth.2017.04.010.
- 45 Khan, S., Amin, F.M., Christensen, C.E., Ghanizada, H., Younis, S., Olinger, A.C.R., de Koning, P.J.H.,
46 Larsson, H.B.W., and Ashina, M. (2019). Meningeal contribution to migraine pain: a magnetic resonance
47 angiography study. *Brain* 142, 93-102. 10.1093/brain/awy300.
- 48 Lafrenière, R.G., Cader, M.Z., Poulin, J.F., Andres-Enguix, I., Simoneau, M., Gupta, N., Boisvert, K.,
49 Lafrenière, F., McLaughlan, S., Dubé, M.P., et al. (2010). A dominant-negative mutation in the TRESK
50 potassium channel is linked to familial migraine with aura. *Nat Med* 16, 1157-1160. 10.1038/nm.2216.
- 51 Nyholt, D.R., LaForge, K.S., Kallela, M., Alakurtti, K., Anttila, V., Färkkilä, M., Hämaläinen, E., Kaprio, J.,
52 Kaunisto, M.A., Heath, A.C., et al. (2008). A high-density association screen of 155 ion transport genes for
53 involvement with common migraine. *Hum Mol Genet* 17, 3318-3331. 10.1093/hmg/ddn227.
- 54 Olesen, J., Diener, H.C., Husstedt, I.W., Goadsby, P.J., Hall, D., Meier, U., Pollentier, S., Lesko, L.M., and
55 Group, B.B.C.P.o.C.S. (2004). Calcitonin gene-related peptide receptor antagonist BIBN 4096 BS for the acute
56 treatment of migraine. *N Engl J Med* 350, 1104-1110. 10.1056/NEJMoa030505.
- 57 Pettingill, P., Weir, G.A., Wei, T., Wu, Y., Flower, G., Lalic, T., Handel, A., Duggal, G., Chintawar, S., Cheung,
58 J., et al. (2019). A causal role for TRESK loss of function in migraine mechanisms. *Brain* 142, 3852-3867.
59 10.1093/brain/awz342.

1 Pradhan, A.A., Smith, M.L., McGuire, B., Tarash, I., Evans, C.J., and Charles, A. (2014). Characterization of a
2 novel model of chronic migraine. *Pain 155*, 269-274. 10.1016/j.pain.2013.10.004.
3 Royal, P., Andres-Bilbe, A., Ávalos Prado, P., Verkest, C., Wdziekonski, B., Schaub, S., Baron, A., Lesage, F.,
4 Gasull, X., Levitz, J., and Sandoz, G. (2019). Migraine-Associated TRESK Mutations Increase Neuronal
5 Excitability through Alternative Translation Initiation and Inhibition of TREK. *Neuron 101*, 232-245.e236.
6 10.1016/j.neuron.2018.11.039.
7 Strassman, A.M., Raymond, S.A., and Burstein, R. (1996). Sensitization of meningeal sensory neurons and the
8 origin of headaches. *Nature 384*, 560-564. 10.1038/384560a0.
9 Stucky, C.L., and Lewin, G.R. (1999). Isolectin B(4)-positive and -negative nociceptors are functionally distinct.
10 *J Neurosci 19*, 6497-6505.
11 Verkest, C., Piquet, E., Diochot, S., Dauvois, M., Lanteri-Minet, M., Lingueglia, E., and Baron, A. (2018).
12 Effects of systemic inhibitors of acid-sensing ion channels 1 (ASIC1) against acute and chronic mechanical
13 allodynia in a rodent model of migraine. *Br J Pharmacol 175*, 4154-4166. 10.1111/bph.14462.
14 Yamamoto, Y., Hatakeyama, T., and Taniguchi, K. (2009). Immunohistochemical colocalization of TREK-1,
15 TREK-2 and TRAAK with TRP channels in the trigeminal ganglion cells. *Neurosci Lett 454*, 129-133.
16 10.1016/j.neulet.2009.02.069.
17

18

19

20

21

22

23

24

25

26

27

28