



HAL
open science

Energy method for the Boltzmann equation of monatomic gaseous mixtures

Laurent Boudin, Bérénice Grec, Milana Pavić-Čolić, Srboľjub Simić

► **To cite this version:**

Laurent Boudin, Bérénice Grec, Milana Pavić-Čolić, Srboľjub Simić. Energy method for the Boltzmann equation of monatomic gaseous mixtures. *Communications in Mathematical Sciences*, inPress. hal-03376932v1

HAL Id: hal-03376932

<https://hal.science/hal-03376932v1>

Submitted on 13 Oct 2021 (v1), last revised 16 May 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ENERGY METHOD FOR THE BOLTZMANN EQUATION OF MONATOMIC GASEOUS MIXTURES

LAURENT BOUDIN, BÉRÉNICE GREC, MILANA PAVIĆ-ČOLIĆ, AND SRBOLJUB SIMIĆ

ABSTRACT. In this paper, we present an energy method for the system of Boltzmann equations in the multicomponent mixture case, based on a micro-macro decomposition. More precisely, the perturbation of a solution to the Boltzmann equation around a global equilibrium is decomposed into the sum of a macroscopic and a microscopic part, for which we obtain *a priori* estimates at both lower and higher orders. These estimates are obtained under a suitable smallness assumption. The assumption can be justified *a posteriori* in the higher-order case, leading to the closure of the corresponding estimate.

1. INTRODUCTION

In the last decade or so, the Boltzmann equation for mixtures, which was already mentioned in [10], attracted the attention of many works. The modelling issue, for both monatomic and polyatomic gases, was for instance discussed in [14, 1, 33] (see also the references therein). Many works focused on the analysis of the monatomic case, like [6, 12, 9, 3], which were dedicated to compactness, hypocoercivity-related and stability results. Well-posedness and regularity were investigated in [9, 8, 18, 13], and asymptotics questions were tackled in [7, 22, 5, 4].

Some of the previous papers, for instance [9], rely on the so-called micro-macro decomposition. In the present work, we aim to provide a more detailed insight on that decomposition in the mixture case. Indeed, the micro-macro decomposition of a solution of the linearized Boltzmann equation has a key role in the study of both mathematical and numerical properties of that solution. It was introduced for the monospecies Boltzmann equation in [20, 21] on the one hand, and in [27, 29, 34] on the other hand. The method consists in considering the equilibrium perturbation as the sum of a macroscopic part and a microscopic one. The macroscopic part can be decomposed on a finite-dimensional subspace, where the associate coordinates solve some conservation laws of fluid type, whereas the microscopic one still solves a kinetic equation. Nevertheless, the microscopic part is incorporated in macroscopic conservation laws and fills the gap between the usual Navier-Stokes approximation and the complete kinetic equation [27]. In fact, it brings information which is essential to provide proper estimates of the perturbed solutions of the kinetic equation.

In the monospecies case, the micro-macro decomposition and the underlying energy method were used for hypocoercivity estimates, see [16], for large-time behaviour studies [28, 25, 32] (see also [26] for a binary mixture), for propagation of one-dimensional waves [30], or to obtain Green's function for the Boltzmann equation [31]. In numerical analysis, this decomposition is a major tool to build asymptotic-preserving (AP) schemes, allowing to pass, for instance, from the Boltzmann equation to the Navier-Stokes equations [2], to exactly conserve some physical quantities [17], or to quantify uncertainty in kinetic equations [15]. As far as the mixture case is concerned, note that several attempts relying on a micro-macro decomposition were already performed, with a BGK approximation [24, 23], in the two-species case [11], and in the general case with partial results [9].

This work was supported by the European COST Action CA18232 Mat-Dyn-Net and by the French CNRS PICS08057 and IEA00375 projects. S. S. and M. P.-Č. acknowledge support of the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant No. 451-03-9/2021-14/200125). M. P.-Č. was supported by the Alexander von Humboldt Foundation.

In this paper, we study the micro-macro decomposition and the corresponding energy method in the multicomponent mixture case, by following the strategy of [29]. More precisely, we start from an equilibrium not depending on the time and space variables. The perturbation of this equilibrium is then decomposed into the sum of microscopic and macroscopic parts, for which we obtain lower and higher order estimates, first using relevant smallness assumptions, and exhibiting closure in the one-dimensional (in space) setting.

It is peculiar for the mixture, in contrast to the monospecies case, that the microscopic part contributes to macroscopic equations not only in the momentum and energy conservation laws, but also in the mass conservation law. This effect is crossed with the perturbation of the energy variable, which altogether makes the procedure of finding proper estimates much more involved. This problem is solved by means of introducing a suitable fluid quantity.

The paper is structured as follows. In the next section, we give a preliminary overview describing the framework for the subsequent analysis. Then, in Section 3, we discuss the methodology and main ideas relying on the micro-macro decomposition, and state our main results. They are *a priori* estimates on the perturbation based on the decomposition, whose proofs are exposed in Sections 4 and 5. In particular, we provide very detailed explanations for the lower-order estimate, knowing that, for the higher-order one, the same kind of computations and ideas are developed.

2. PRELIMINARIES

We consider an ideal gas mixture constituted with $I \geq 2$ monatomic species. Each species, indexed by $1 \leq i \leq I$, is described thanks to a distribution function F_i , which is nonnegative, and depends on time $t \in [0, T]$, $T > 0$, space position $x \in \mathbb{R}$ and microscopic velocity $v \in \mathbb{R}^3$. We denote by m_i the atomic mass of species i . We emphasize that we choose to work here in a one-dimensional setting for the space variable x , not only for the sake of simplicity. Indeed, if most computations and results remain true in dimensions 2 and 3, the estimates are closed in this work by introducing the antiderivative of the macroscopic part of the decomposition, which can only be performed in a one-dimensional setting.

To consider the species altogether, we introduce the vector distribution function of the mixture, denoted by $\mathbf{F} = (F_i)_{1 \leq i \leq I}$. It satisfies the system of Boltzmann equations, also written in a vector form,

$$(1) \quad \partial_t \mathbf{F} + v_1 \partial_x \mathbf{F} = \mathbf{Q}(\mathbf{F}, \mathbf{F}),$$

where v_1 is the coordinate of velocity v in a direction of the space variable x , and \mathbf{Q} is the vector collision operator, which only acts on the velocity variable v . The vector collision operator \mathbf{Q} can be defined component-wise. To this end, we first need to recall the microscopic context of the collisions.

We assume that the mixture only involves elastic collisions, without chemical reactions. Consider two colliding molecules, one of species i and another one of species j , with respective pre-collisional velocities v' and v'_* . Those velocities change after collision into post-collisional velocities v and v_* , with both momentum and kinetic energy conserved, *i.e.*

$$(2) \quad m_i v' + m_j v'_* = m_i v + m_j v_*, \quad \frac{1}{2} m_i v'^2 + \frac{1}{2} m_j v'^2_* = \frac{1}{2} m_i v^2 + \frac{1}{2} m_j v_*^2.$$

The previous equalities allow to introduce a parameter $\omega \in \mathbb{S}^2$, enabling to write v' and v'_* in terms of v and v_* as

$$(3) \quad v' = \frac{m_i v + m_j v_*}{m_i + m_j} + \frac{m_j}{m_i + m_j} T_\omega(v - v_*), \quad v'_* = \frac{m_i v + m_j v_*}{m_i + m_j} - \frac{m_i}{m_i + m_j} T_\omega(v - v_*),$$

denoting $T_\omega z = z - 2(\omega \cdot z)\omega$ for any $z \in \mathbb{R}^3$.

Then, for any i, j , we can define the operator Q_{ij} describing the atomic interactions of species i with species j . It only acts on the velocity variable and is given by

$$Q_{ij}(f_i, g_j)(v) = \iint_{\mathbb{R}^3 \times S^2} [f_i(v')g_j(v'_*) - f_i(v)g_j(v_*)] \mathcal{B}_{ij}(v, v_*, \omega) d\omega dv_*,$$

for any species-related real-valued functions f_i, g_j of the velocity variable. The cross-section \mathcal{B}_{ij} allows to classify the way species i and j interact and must satisfy the micro-reversibility property

$$\mathcal{B}_{ij}(v', v'_*, \omega) = \mathcal{B}_{ji}(v_*, v, \omega) = \mathcal{B}_{ij}(v, v_*, \omega) \geq 0.$$

Moreover, in this work, we make the hard-sphere assumption, for any i, j ,

$$(4) \quad \mathcal{B}_{ij}(v, v_*, \omega) = \beta_{ij} |(v - v_*) \cdot \omega|,$$

where $\beta_{ij} > 0$ is given. The assumption is required to ensure needed properties of the collision frequency and to deal with the nonlinearity.

Eventually, we can define the i -th component of \mathbf{Q} , with $\mathbf{f} = (f_j)_{1 \leq j \leq I}$, $\mathbf{g} = (g_j)_{1 \leq j \leq I}$, by

$$Q_i(\mathbf{f}, \mathbf{g}) = \sum_{j=1}^I Q_{ij}(f_i, g_j).$$

Before recalling the main properties of the solutions to (1), let us introduce some very convenient notations. First, we define a component-wise product of two vectors $\mathbf{A} = (A_i)_{1 \leq i \leq I}$, $\mathbf{B} = (B_i)_{1 \leq i \leq I}$ and a vector-valued function of \mathbf{A} , for $\Phi : \mathbb{R} \rightarrow \mathbb{R}$, by

$$\mathbf{A}\mathbf{B} = \begin{pmatrix} A_1 B_1 \\ A_2 B_2 \\ \vdots \\ A_I B_I \end{pmatrix}, \quad \Phi(\mathbf{A}) = \begin{pmatrix} \Phi(A_1) \\ \Phi(A_2) \\ \vdots \\ \Phi(A_I) \end{pmatrix}.$$

This way, we can write, for instance, $\mathbf{A}^{1/2} = (A_i^{1/2})_{1 \leq i \leq I}$, when $A_i \geq 0$. Finally, $L^2(\mathbb{R}^3)^I$ is endowed with its natural scalar product and norm, *i.e.* we set, for any vector functions $\mathbf{f} = (f_i)_{1 \leq i \leq I}$, $\mathbf{g} = (g_i)_{1 \leq i \leq I} \in L^2(\mathbb{R}^3)^I$,

$$\langle \mathbf{f}, \mathbf{g} \rangle_I = \sum_{i=1}^I \int_{\mathbb{R}^3} f_i g_i dv, \quad \|\mathbf{f}\|_I = \langle \mathbf{f}, \mathbf{f} \rangle_I^{1/2}.$$

Conservative properties of the Boltzmann equations are obtained thanks to the weak form of the collision operator that uses some symmetries built in the model. In the mixture setting, the weak form is carefully described, for example, in [14, 7, 5]. We only mention here the final formula. For any functions \mathbf{G} and ψ for which it makes sense, we have

$$\begin{aligned} \langle \mathbf{Q}(\mathbf{G}, \mathbf{G}), \psi \rangle_I &= -\frac{1}{4} \sum_{i,j=1}^I \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times S^2} [G_i(v')G_j(v'_*) - G_i(v)G_j(v_*)] \\ &\quad \times [\psi_i(v') + \psi_j(v'_*) - \psi_i(v) - \psi_j(v_*)] \mathcal{B}_{ij}(v, v_*, \omega) d\omega dv_* dv. \end{aligned}$$

In this paper, we work in a perturbative setting, around a global equilibrium distribution function. Its notion is introduced in the so-called H -theorem, see [14] for instance. Let us first define the entropy production functional

$$D(\mathbf{G}) = \langle \mathbf{Q}(\mathbf{G}, \mathbf{G}), \log \mathbf{G} \rangle_I.$$

The H -theorem reads

Proposition 1. *Assume that all the cross sections are positive almost everywhere and that \mathbf{G} is such that both $\mathbf{Q}(\mathbf{G}, \mathbf{G})$ and $D(\mathbf{G})$ are well defined. Then*

- (a) *The entropy production is non-positive, i.e. $D(\mathbf{G}) \leq 0$.*

(b) Moreover, the three following properties are equivalent:

- i. for any $1 \leq i, j \leq I$, $Q_{ij}(G_i, G_j) = 0$;
- ii. the entropy production vanishes, that is $D(\mathbf{G}) = 0$;
- iii. there exist $T > 0$ and $u \in \mathbb{R}^3$ such that, for any i , there exists $n_i \geq 0$ such that

$$G_i(v) = n_i \left(\frac{m_i}{2\pi kT} \right)^{3/2} e^{-\frac{m_i}{2kT}|v-u|^2}.$$

Choosing $kT = 1$, $u = 0$, \mathbf{n} as a nonnegative constant vector, we obtain the normalized centered Maxwell vector function \mathbf{M} as

$$M_i(v) = \left(\frac{m_i}{2\pi} \right)^{3/2} e^{-\frac{m_i}{2}|v|^2}, \quad 1 \leq i \leq I.$$

Let us then recall the collision invariants in the gas mixtures setting which can be found in [9], for instance. The collision invariants are the velocity-depending functions which make the previous weak form of \mathbf{Q} vanish. They are moreover chosen one-to-one orthogonal and normalized with respect to a L^2 scalar product weighted in terms of \mathbf{nM} (remembering the component-wise multiplication defined above). More precisely, we set

$$(5) \quad \left\{ \begin{array}{l} \chi^1 = \frac{1}{\sqrt{n_1}} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad \chi^2 = \frac{1}{\sqrt{n_2}} \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix}, \quad \dots, \quad \chi^I = \frac{1}{\sqrt{n_I}} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}, \\ \chi^{I+1} = \frac{1}{\sqrt{\sum_{j=1}^I n_j m_j}} \begin{pmatrix} m_1 v_1 \\ m_2 v_1 \\ \vdots \\ m_I v_1 \end{pmatrix}, \quad \dots, \quad \chi^{I+3} = \frac{1}{\sqrt{\sum_{j=1}^I n_j m_j}} \begin{pmatrix} m_1 v_3 \\ m_2 v_3 \\ \vdots \\ m_I v_3 \end{pmatrix}, \\ \chi^{I+4} = \frac{1}{\sqrt{6 \sum_{j=1}^I n_j}} \begin{pmatrix} m_1 |v|^2 - 3 \\ m_2 |v|^2 - 3 \\ \vdots \\ m_I |v|^2 - 3 \end{pmatrix}. \end{array} \right.$$

Then the family $(\chi^k)_{1 \leq k \leq I+4}$ satisfies, for any \mathbf{G} ,

$$(6) \quad \begin{aligned} \langle \mathbf{Q}(\mathbf{G}, \mathbf{G}), \chi^k \rangle_I &= 0, \quad 1 \leq k \leq I+4, \\ \langle (\mathbf{nM})^{1/2} \chi^k, (\mathbf{nM})^{1/2} \chi^\ell \rangle_I &= \delta_{k\ell}, \quad 1 \leq k, \ell \leq I+4. \end{aligned}$$

In this paper, we focus on a perturbation of the global equilibrium distribution function \mathbf{nM} . More precisely, we consider a perturbation carried by a vector-valued function \mathbf{f} , which implies that \mathbf{F} takes the form

$$(7) \quad \mathbf{F} = \mathbf{nM} + (\mathbf{nM})^{1/2} \mathbf{f}.$$

Since \mathbf{nM} does not depend on t and x , we shall carefully study the macroscopic part of the perturbation, which contains the time and space variations of \mathbf{f} and subsequently of \mathbf{F} , as in [29], bringing at the same time some more details about the estimates to handle the mixture case. Note that another possibility would have been to follow [27], in which \mathbf{F} is decomposed into the sum of a local Maxwellian, containing the whole macroscopic part of the distribution function, and the microscopic part. This decomposition induces other difficulties, such as the dependence of $\ker \mathcal{L}$ on x and t .

3. MAIN IDEAS AND RESULTS

Let us now focus our attention on the micro-macro decomposition of the perturbation \mathbf{f} . Straightforwardly, (1) implies that \mathbf{f} satisfies

$$(8) \quad \partial_t \mathbf{f} + v_1 \partial_x \mathbf{f} - \mathcal{L} \mathbf{f} = \mathcal{N}(\mathbf{f}),$$

where \mathcal{L} and \mathcal{N} are respectively the linearized Boltzmann operator and a quadratic operator defined by

$$\begin{aligned} \mathcal{L} \mathbf{f} &= (\mathbf{n}M)^{-1/2} \left(\mathbf{Q}(\mathbf{n}M, (\mathbf{n}M)^{1/2} \mathbf{f}) + \mathbf{Q}((\mathbf{n}M)^{1/2} \mathbf{f}, \mathbf{n}M) \right), \\ \mathcal{N}(\mathbf{f}) &= (\mathbf{n}M)^{-1/2} \mathbf{Q}((\mathbf{n}M)^{1/2} \mathbf{f}, (\mathbf{n}M)^{1/2} \mathbf{f}). \end{aligned}$$

In the remainder of the paper, we shall denote by \mathbb{D} the domain of \mathcal{L} in $L^2(\mathbb{R}^3)^I$.

It is easy to see [9, 6] that the operator \mathcal{L} is a non-positive self-adjoint operator, *i.e.* for any $\mathbf{f}, \mathbf{g} \in \mathbb{D}$,

$$\langle \mathcal{L} \mathbf{f}, \mathbf{g} \rangle_I = \langle \mathbf{f}, \mathcal{L} \mathbf{g} \rangle_I, \quad \langle \mathcal{L} \mathbf{f}, \mathbf{f} \rangle_I \leq 0.$$

The collisions invariants allow to characterize the elements of $\mathbb{P}^0 = \ker \mathcal{L}$, *i.e.*

$$(9) \quad \mathbb{P}^0 = \ker \mathcal{L} = \text{Span} \left\{ (\mathbf{n}M)^{1/2} \boldsymbol{\chi}^k \mid 1 \leq k \leq I + 4 \right\},$$

which is a finite-dimensional subspace of \mathbb{D} with $\dim \mathbb{P}^0 = I + 4$. Let us denote by \mathbb{P}^1 its orthogonal complement in \mathbb{D} with respect to the $\langle \cdot, \cdot \rangle_I$ scalar product, *i.e.* $\mathbb{P}^1 = (\ker \mathcal{L})^\perp = (\mathbb{P}^0)^\perp$.

If we naively proceed by multiplying (8) by \mathbf{f} and integrate with respect to t , x and v , the only term we can hope to upper-bound comes from a spectral gap estimate for \mathcal{L} : it is a square norm of the projection of \mathbf{f} onto \mathbb{P}^1 , which we denote \mathbf{f}^1 . That implies that we fail to control the norm of the projection \mathbf{f}^0 of \mathbf{f} onto \mathbb{P}^0 . To control \mathbf{f}^0 , we use the so-called micro-macro decomposition of \mathbf{f} , as \mathbf{f} is uniquely written as

$$\mathbf{f} = \mathbf{f}^0 + \mathbf{f}^1.$$

We shall get back to it later, but \mathbf{f}^0 is called the macroscopic part of \mathbf{f} , and \mathbf{f}^1 its microscopic part.

The statements below will be accurately justified in the upcoming sections. The projections of \mathbf{f} satisfy, as in [29],

$$\begin{aligned} \partial_t \mathbf{f}^0 + \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1) &= 0, \\ \partial_t \mathbf{f}^1 + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1) - \mathcal{L} \mathbf{f}^1 &= \mathcal{N}(\mathbf{f}). \end{aligned}$$

As we shall see, $\bar{\mathcal{L}} = \mathcal{L}|_{\mathbb{P}^1}$ is invertible, thus we get, from the previous equation on \mathbf{f}^1 ,

$$\mathbf{f}^1 = \bar{\mathcal{L}}^{-1} \left(\partial_t \mathbf{f}^1 + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1) - \mathcal{N}(\mathbf{f}) \right).$$

We now plug this expression of \mathbf{f}^1 in the previous equation on \mathbf{f}^0 , which ensures

$$\partial_t \mathbf{f}^0 + \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^0 \left(v_1 \partial_x \bar{\mathcal{L}}^{-1} \left(\partial_t \mathbf{f}^1 + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1) - \mathcal{N}(\mathbf{f}) \right) \right) = 0.$$

We immediately observe that, when scalarly multiplying by \mathbf{f}^0 , there is no way to exhibit an estimate on a norm of \mathbf{f}^0 itself. There may only be hope to find an estimate of the norm $\|\partial_x \mathbf{f}^0\|_I^2$. Indeed, we recall that \mathbb{P}^0 is a finite-dimensional space, and $\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)$ can be expressed in terms of the space derivatives of the coordinates of \mathbf{f}^0 in the orthonormal basis of \mathbb{P}^0 . Then a subtle combination of arguments, some new and specific to the mixture case, some others coming from [27, 29], leads to an estimate involving $\|\partial_x \mathbf{f}^0\|_I^2$ on the left-hand side. We still do not have any control on $\|\mathbf{f}^0\|_I^2$, but such a control is needed, since the nonlinear term involves \mathbf{f} , and not only \mathbf{f}^1 (this is due to the fact that $\mathcal{N}(\mathbf{f}^0) \neq 0$), and derivatives of the quadratic term $\mathcal{N}(\mathbf{f})$ involve the derivatives of \mathbf{f} and \mathbf{f} itself. We can see that it is possible to do so by introducing the antiderivative

\mathbf{W}^0 of \mathbf{f}^0 with respect to x , which justifies the fact that we are working in a one-dimensional setting. Without loss of generality, as explained below in Remark 1, we shall assume that

$$(10) \quad \int_{\mathbb{R}} \mathbf{f}^0(0, x, v) dx = 0, \quad v \in \mathbb{R}^3,$$

so that

$$\mathbf{W}^0 : (t, x, v) \mapsto \int_{-\infty}^x \mathbf{f}^0(t, y, v) dy$$

can be treated within an L^2 -framework in both variables x and v . This allows to derive an estimate on $\partial_x \mathbf{W}^0 = \mathbf{f}^0$, roughly as we did for $\partial_x \mathbf{f}^0$. Eventually, our global estimate requires the control of norms of the time and space derivatives of \mathbf{f}^1 , which can be obtained after differentiation of the equation on \mathbf{f}^1 with respect to t or x , thanks to the spectral gap of \mathcal{L} . This whole process also requires a mandatory assumption to deal with the nonlinear term $\mathcal{N}(\mathbf{f})$, which implies that \mathbf{W}^0 and \mathbf{f} must remain small in the $\|\cdot\|_I$ norm, pointwise in time and space.

Let us first make the following smallness assumption, where $\varepsilon > 0$ will be chosen afterwards.

Assumption. *The perturbation \mathbf{f} must satisfy*

$$(11) \quad \sup_{\substack{t \geq 0 \\ x \in \mathbb{R}}} \left(\|\mathbf{W}^0\|_I + \|\mathbf{f}^0\|_I + \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I \right) \leq \varepsilon.$$

This assumption is stronger than the one used in [29], where the term $\|\mathbf{f}^0\|_I^2$ was not involved, but also allows to obtain an estimate on the full macroscopic part, which was not the case previously. More precisely, our first main result is an *a priori* estimate on norms of \mathbf{f} , its partial derivatives, and the antiderivative of \mathbf{f}^0 . The generic term denoted by $\mathcal{I}(0)$ which appears below must be understood as a linear combination of square norms of initial data of the pointwise in time integrals of the left-hand side of the estimates, with coefficients only depending on the problem data. Without assuming (10), one must add in $\mathcal{I}(0)$ a term involving the square norm in $L^2(\mathbb{R}^3)^I$ of the element $(v \mapsto \int_{\mathbb{R}} \mathbf{f}^0(0, x, v) dx)$ of \mathbb{P}^0 , see Remark 1.

Proposition 2. *There exists $\varepsilon_0 > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0]$, we can find positive constants $\alpha_0, \alpha_1, \hat{C}_1, \hat{C}_2, \hat{C}_3$ such that, for any solution \mathbf{f} to (8) satisfying the smallness assumption (11), the following lower-order a priori estimate holds*

$$(12) \quad \begin{aligned} & \frac{1}{4} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_{t=T} + \frac{1}{4} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_{t=T} + \alpha_0 \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{\alpha_1}{2} \int_{\mathbb{R}} \|\partial_x \mathbf{f}\|_I^2 dx \Big|_{t=T} \\ & + \frac{\alpha_1}{2} \int_{\mathbb{R}} \|\partial_t \mathbf{f}\|_I^2 dx \Big|_{t=T} + \hat{C}_1 \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt + \hat{C}_1 \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt \\ & + \hat{C}_2 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 dx dt + 2\hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial_x \mathbf{f}^1 \right\|_I^2 dx dt \\ & + 2\hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial_t \mathbf{f}^1 \right\|_I^2 dx dt \leq \mathcal{I}(0). \end{aligned}$$

Unfortunately, estimate (12) is not closed. For the latter property to hold, choosing the initial data such that $\mathcal{I}(0)$ is of order ε^2 would have to imply (11). However, $\|(1 + |v|)^{1/2} \mathbf{f}^1\|_I$ would only lie in $H_{t,x}^1$, which does not continuously inject in $L_{t,x}^\infty$. We thus need a higher-order estimate involving more derivatives with respect to t and x . In order to obtain it, we differentiate the equations satisfied by \mathbf{f}^0 and \mathbf{f} as many times as necessary. To this aim, we introduce the following notation. For any $p = (p_1, p_2) \in \mathbb{N}^2$, we set $|p| = p_1 + p_2$, and, for any function \mathbf{g} of t, x and v ,

$$\partial^p \mathbf{g} = \partial_t^{p_1} \partial_x^{p_2} \mathbf{g}.$$

We emphasize that the multi-index notation does not imply any derivative with respect to v .

At this point, we make the following smallness assumption, where $\varepsilon > 0$ will be chosen afterwards.

Assumption. *The perturbation \mathbf{f} must satisfy*

$$(13) \quad \sup_{\substack{t>0 \\ x \in \mathbb{R}}} \left[\|\mathbf{W}^0\|_I + \max_{|p| \leq 2} \left(\|\partial^p \mathbf{f}^0\|_I + \|(1+|v|)^{1/2} \partial^p \mathbf{f}^1\|_I \right) \right] \leq \varepsilon.$$

Under Assumption (13), we obtain the next theorem. This time, as we explain below, the smallness assumption can be dropped, provided that we assume instead that $\mathcal{I}(0)$ is small.

Theorem 3. *There exists $\varepsilon_1 > 0$ such that, for any $\varepsilon \in (0, \varepsilon_1]$, and any solution \mathbf{f} to (8) such that the corresponding $\mathcal{I}(0)$ is at most of order ε^2 , the following higher-order a priori estimate holds*

$$(14) \quad \begin{aligned} & \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_{t=T} + \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_{t=T} + \sum_{1 \leq |r| \leq 4} \int_{\mathbb{R}} \|\partial^r \mathbf{f}^0\|_I^2 dx \Big|_{t=T} \\ & \quad + \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_{t=T} + \sum_{1 \leq |p| \leq 5} \int_{\mathbb{R}} \|\partial^p \mathbf{f}\|_I^2 dx \Big|_{t=T} \\ & \quad + \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt + \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt + \sum_{1 \leq |r| \leq 4} \int_0^T \int_{\mathbb{R}} \|\partial_x \partial^r \mathbf{f}^0\|_I^2 dx dt \\ & \quad + \int_0^T \int_{\mathbb{R}} \|(1+|v|)^{1/2} \mathbf{f}^1\|_I^2 dx dt + \sum_{1 \leq |p| \leq 5} \int_0^T \int_{\mathbb{R}} \|(1+|v|)^{1/2} \partial^p \mathbf{f}^1\|_I^2 dx dt \leq \mathcal{I}(0). \end{aligned}$$

Estimate (14) is closed because, this time, $\|(1+|v|)^{1/2} \mathbf{f}^1\|_I$ lies in $H_{t,x}^5 \hookrightarrow W_{t,x}^{2,\infty}$, and $\|\mathbf{f}^0\|_I$ lies in $H_{t,x}^4 \hookrightarrow W_{t,x}^{2,\infty}$. Estimate (14) implies that Assumption (13) is automatically satisfied when $\mathcal{I}(0)$ is of order ε^2 . The latter closed estimate (14) appears as a very useful tool. For instance, it allows to obtain the stability of the global Maxwellian function (\mathbf{nM}) in large time, provided that the perturbation at initial time is chosen small enough in the $H^s(L_v^2)$ norm, for $s \geq 5$.

Let us now focus on the proofs of Proposition 2 and Theorem 3, starting with the lower-order estimate.

4. PROOF OF THE LOWER-ORDER ESTIMATE

4.1. Estimates on \mathcal{L} and \mathcal{N} . The linearized operator \mathcal{L} can be written as $\mathcal{L} = \mathbf{K} - \nu$, where \mathbf{K} is compact [6] and ν is a multiplicative operator, called the collision frequency, given by

$$\nu_i(v) = \sum_{j=1}^I n_j \left(\frac{m_j}{2\pi} \right)^{3/2} \iint_{\mathbb{R}^3 \times S^2} e^{-\frac{m_j}{2}|v_*|^2} \mathcal{B}_{ij}(v, v_*, \omega) d\omega dv_*, \quad v \in \mathbb{R}^3, \quad 1 \leq i \leq I.$$

Note that, thanks to the hard-sphere assumption (4) on the cross sections, as in [27], ν satisfies a growth estimate. More precisely, there exist positive constants ν_0 and $\bar{\nu}_0$, such that

$$(15) \quad 0 < \nu_0 \leq \nu_0(1+|v|) \leq \nu_i(v) \leq \bar{\nu}_0(1+|v|), \quad v \in \mathbb{R}^3, \quad 1 \leq i \leq I.$$

Besides, in [9], a constructive spectral gap estimate on \mathcal{L} is proved. In our notation, it means that there exists $\lambda > 0$ such that, for any $\mathbf{h} \in (\ker \mathcal{L})^\perp$,

$$(16) \quad \langle \mathcal{L}\mathbf{h}, \mathbf{h} \rangle_I \leq -\lambda \|\nu^{1/2} \mathbf{h}\|_I^2.$$

The spectral gap estimate (16) on \mathcal{L} and the lower bound ν_0 on ν from (15), together with the Cauchy-Schwarz inequality, yield

$$\nu_0 \|\mathbf{g}\|_I^2 \leq \|\nu^{1/2} \mathbf{g}\|_I^2 \leq -\frac{1}{\lambda} \langle \mathcal{L}\mathbf{g}, \mathbf{g} \rangle_I \leq \frac{1}{\lambda} \|\mathbf{g}\|_I \|\mathcal{L}\mathbf{g}\|_I.$$

Since $\mathbb{P}^1 = (\ker \mathcal{L})^\perp$ and \mathcal{L} is self-adjoint, it is clear that $\mathcal{L}(\mathbb{P}^1) = \mathbb{P}^1$, hence $\bar{\mathcal{L}} = \mathcal{L}|_{\mathbb{P}^1}$ is an invertible operator on \mathbb{P}^1 . Now, for any $\mathbf{h} \in \mathbb{P}^1$, writing $\mathbf{g} = \bar{\mathcal{L}}^{-1}\mathbf{h}$ and setting $C_{\text{inv}} = (\nu_0\lambda)^{-1} > 0$, the previous inequality implies that

$$(17) \quad \|\bar{\mathcal{L}}^{-1}\mathbf{h}\|_I \leq C_{\text{inv}}\|\mathbf{h}\|_I,$$

which ensures the boundedness of $\bar{\mathcal{L}}^{-1}$ on \mathbb{P}^1 .

Eventually, we write an estimate on the nonlinear operator \mathcal{N} . Its i -th component, $1 \leq i \leq I$, is given by

$$\mathcal{N}_i(\mathbf{f}) = M_i^{-1/2} \sum_{j=1}^I n_j^{1/2} Q_{ij}(M_i^{1/2} f_i, M_j^{1/2} f_j).$$

Thanks to Lemma 8 stated in Appendix A, we immediately have

$$(18) \quad \left\| (1 + |v|)^{-1/2} \mathcal{N}(\mathbf{f}) \right\|_I \leq C_\beta \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I^2,$$

$$(19) \quad \left\| (1 + |v|)^{-1/2} \partial_\star \mathcal{N}(\mathbf{f}) \right\|_I \leq 2 C_\beta \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial_\star \mathbf{f} \right\|_I,$$

where ∂_\star denotes either time or space partial differentiation.

4.2. Micro-macro decomposition. The micro-macro decomposition method lies on the orthogonal decomposition of \mathbb{D} onto $\mathbb{P}^0 = \ker \mathcal{L}$ and $\mathbb{P}^1 = (\ker \mathcal{L})^\perp = \text{im } \mathcal{L}$. In order to perform it, let \mathbf{P}^0 and \mathbf{P}^1 respectively denote the orthogonal projections on \mathbb{P}^0 and \mathbb{P}^1 . It is clear that

$$\mathcal{L}\mathbf{P}^0 = \mathbf{P}^0\mathcal{L} = 0, \quad \mathcal{L}\mathbf{P}^1 = \mathbf{P}^1\mathcal{L}.$$

Moreover, thanks to (9), we can write, for any $\mathbf{g} \in \mathbb{D}$,

$$(20) \quad \mathbf{P}^0\mathbf{g} = \sum_{k=1}^{I+4} \left\langle (\mathbf{nM})^{1/2} \boldsymbol{\chi}^k, \mathbf{g} \right\rangle_I (\mathbf{nM})^{1/2} \boldsymbol{\chi}^k.$$

We decompose the perturbation \mathbf{f} following the direct orthogonal sum $\mathbb{P}^0 \oplus \mathbb{P}^1$. Then, $\mathbf{f}^0 = \mathbf{P}^0\mathbf{f}$ and $\mathbf{f}^1 = \mathbf{P}^1\mathbf{f}$ satisfy

$$\mathbf{f} = \mathbf{P}^0\mathbf{f} + \mathbf{P}^1\mathbf{f} = \mathbf{f}^0 + \mathbf{f}^1, \quad \langle \mathbf{f}^0, \mathbf{f}^1 \rangle_I = 0.$$

Functions \mathbf{f}^0 and \mathbf{f}^1 are respectively referred to as the macroscopic (or fluid) and microscopic (or non-fluid) components of \mathbf{f} .

The coordinates of \mathbf{f}^0 in the orthonormal basis $((\mathbf{nM})^{1/2} \boldsymbol{\chi}^k)_{1 \leq k \leq I+4}$ of \mathbb{P}^0 , also known as the fluid quantities, are given by

$$\begin{aligned} \rho_i(t, x) &= \left\langle (\mathbf{nM})^{1/2} \boldsymbol{\chi}^i, \mathbf{f} \right\rangle_I, \quad 1 \leq i \leq I, \\ q^k(t, x) &= \left\langle (\mathbf{nM})^{1/2} \boldsymbol{\chi}^{I+k}, \mathbf{f} \right\rangle_I, \quad k = 1, 2, 3, \\ e(t, x) &= \left\langle (\mathbf{nM})^{1/2} \boldsymbol{\chi}^{I+4}, \mathbf{f} \right\rangle_I, \end{aligned}$$

so that

$$(21) \quad \mathbf{f}^0 = \sum_{i=1}^I \rho_i (\mathbf{nM})^{1/2} \boldsymbol{\chi}^i + \sum_{k=1}^3 q^k (\mathbf{nM})^{1/2} \boldsymbol{\chi}^{I+k} + e (\mathbf{nM})^{1/2} \boldsymbol{\chi}^{I+4}.$$

Using (5) and the more compact (I -sized) vector writing, we obtain

$$(22) \quad \mathbf{f}^0 = \boldsymbol{\rho} \mathbf{M}^{1/2} + \left(\sum_{k=1}^3 \frac{q^k v_k}{\sqrt{\sum_j n_j m_j}} \right) \mathbf{m} (\mathbf{nM})^{1/2} + \frac{e}{\sqrt{6 \sum_j n_j}} (|v|^2 \mathbf{m} - 3\mathbf{E}) (\mathbf{nM})^{1/2},$$

where we set $\mathbf{E} = (1 \dots 1)^\top$.

If we project the equation (8) on the perturbation \mathbf{f} onto \mathbb{P}^0 and \mathbb{P}^1 , respectively, we obtain the following equations respectively satisfied by \mathbf{f}^0 and \mathbf{f}^1 , *i.e.*

$$(23) \quad \partial_t \mathbf{f}^0 + \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1) = 0,$$

$$(24) \quad \partial_t \mathbf{f}^1 + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1) - \mathcal{L} \mathbf{f}^1 = \mathcal{N}(\mathbf{f}).$$

The nonlinear term $\mathbf{P}^0 \mathcal{N}(\mathbf{f})$ vanished in (23) by combining the representation (20) of elements of \mathbb{P}^0 , the definition of \mathcal{N} , and the one (6) of the collision invariants. This means that $\mathcal{N}(\mathbf{f})$ lies in \mathbb{P}^1 , which we take into account, together with the commutation of \mathbf{P}^1 and \mathcal{L} to rewrite (24) as an equality regarding \mathbf{f}^1 , that is

$$(25) \quad \mathbf{f}^1 = \bar{\mathcal{L}}^{-1} (\partial_t \mathbf{f}^1 + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) + \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1) - \mathcal{N}(\mathbf{f})).$$

Of course, (25) does not provide a direct expression of \mathbf{f}^1 , since its right-hand side still depends on the nonlinear term $\mathcal{N}(\mathbf{f}) = \mathcal{N}(\mathbf{f}^0 + \mathbf{f}^1)$ and first-order derivatives of \mathbf{f}^0 and \mathbf{f}^1 . Nevertheless, this equality is crucial for the rest of the paper. Let us also emphasize here that time or space differentiations commute with \mathcal{L} , \mathbf{P}^0 and \mathbf{P}^1 , since these three operators only act on the velocity variable.

Let us now state a result on the fluid quantities, which is important for the further analysis. It comes from the equation (23) satisfied by \mathbf{f}^0 and provides conservation laws for $(\rho_i)_{1 \leq i \leq I}$, $(q^k)_{k \in \{1,2,3\}}$ and e .

Proposition 4. *The fluid quantities of \mathbf{f}^0 satisfy the following conservation laws*

$$(26) \quad \partial_t \rho_i + \frac{n_i}{\sqrt{\sum n_j m_j}} \partial_x q^1 + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), \chi^i (\mathbf{nM})^{1/2} \right\rangle_I = 0, \quad 1 \leq i \leq I,$$

$$(27) \quad \partial_t q^1 + \frac{1}{\sqrt{\sum n_j m_j}} \partial_x \left(\sum_i \sqrt{n_i} \rho_i + \frac{\sqrt{6 \sum n_j}}{3} e \right) + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), \chi^{I+1} (\mathbf{nM})^{1/2} \right\rangle_I = 0,$$

$$(28) \quad \partial_t q^k + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), \chi^{I+k} (\mathbf{nM})^{1/2} \right\rangle_I = 0, \quad k = 2, 3,$$

$$(29) \quad \partial_t e + \frac{1}{3} \sqrt{\frac{6 \sum n_j}{\sum n_j m_j}} \partial_x q^1 + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), \chi^{I+4} (\mathbf{nM})^{1/2} \right\rangle_I = 0.$$

The principle of the proof is very simple (checking the equations), but the computations inside are tedious. The proof is provided in Appendix B for the sake of completeness. Note that the term with \mathbf{f}^1 in (26) is peculiar to the mixture and does not appear in the monospecies case.

In the following, we shall also need to introduce the fluid quantity appearing in (27) with its space derivative, that is

$$(30) \quad \ell = \frac{1}{\sqrt{\sum n_j m_j}} \left(\sum_i \sqrt{n_i} \rho_i + \frac{2\sqrt{\sum n_j}}{\sqrt{6}} e \right).$$

Using (26) and (29), ℓ clearly satisfies

$$(31) \quad \partial_t \ell + \frac{\sum n_j^{3/2} + \frac{2}{3} \sum n_j}{\sum n_j m_j} \partial_x q^1 + \frac{1}{\sqrt{\sum n_j m_j}} \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), \left(\sum_i \sqrt{n_i} \chi^i + \frac{1}{3} \sqrt{6 \sum n_j} \chi^{I+4} \right) (\mathbf{nM})^{1/2} \right\rangle_I = 0,$$

so that q^1 and ℓ can easily be linked through (27) and (31).

To conclude this subsection, note that, thanks to (10), the perturbation \mathbf{f} has zero total macroscopic quantities at initial time, *i.e.*

$$\int_{\mathbb{R}} \rho_i dx \Big|_{t=0} = 0, \quad 1 \leq i \leq I, \quad \int_{\mathbb{R}} q^k dx \Big|_{t=0} = 0, \quad 1 \leq k \leq 3, \quad \int_{\mathbb{R}} e dx \Big|_{t=0} = 0.$$

The antiderivatives (R_i) , (Q^k) and E of (ρ_i) , (q^k) and e with respect to x are also the coordinates of \mathbf{W}^0 in the basis $(\chi^k(\mathbf{nM})^{1/2})$ of \mathbb{P}^0 . Denote by L the antiderivative of ℓ , then we have the straightforward corollary of Proposition 4.

Corollary 5. *The fluid quantities in \mathbf{W}^0 satisfy*

$$(32) \quad \partial_t Q^1 + \ell + \left\langle \mathbf{P}^0(v_1 \mathbf{f}^1), \chi^{I+1}(\mathbf{nM})^{1/2} \right\rangle_I = 0,$$

$$(33) \quad \partial_t L + \frac{\sum n_j^{3/2} + \frac{2}{3} \sum n_j}{\sum n_j m_j} q^1 + \frac{1}{\sqrt{\sum n_j m_j}} \left\langle \mathbf{P}^0(v_1 \mathbf{f}^1), \left(\sum_i \sqrt{n_i} \chi^i + \frac{1}{3} \sqrt{6 \sum n_j} \chi^{I+4} \right) (\mathbf{nM})^{1/2} \right\rangle_I = 0.$$

Remark 1. *As promised, let us comment on Assumption (10) on the initial datum on \mathbf{f}^0 . If it is not satisfied, Proposition 4 ensures that $\int_{\mathbb{R}} \rho_i dx$, $\int_{\mathbb{R}} q^k dx$ and $\int_{\mathbb{R}} e dx$ remain constants with respect to t , one of them, at least, being nonzero. Consequently, one of the antiderivatives of the perturbation macroscopic quantity would not be in $L^2(\mathbb{R})$. To deal with that issue, consider a nonnegative C^∞ -compactly supported function ψ in the variable x satisfying $\int_{\mathbb{R}} \psi(y) dy = 1$, and set*

$$\widetilde{\mathbf{W}}^0(t, x, v) = \mathbf{W}^0(t, x, v) - \left(\int_{-\infty}^x \psi(y) dy \right) \left(\int_{\mathbb{R}} \mathbf{f}^0(0, y, v) dy \right).$$

As a function of v , $\widetilde{\mathbf{W}}^0$ clearly belongs to \mathbb{P}^0 , and it is not restrictive to assume that $\widetilde{\mathbf{W}}^0(0, \cdot, \cdot)$ lies in $L^2(\mathbb{R}; L^2(\mathbb{R}^3)^I)$. Then it is safe to proceed with the proof by using $\widetilde{\mathbf{W}}^0$ instead of \mathbf{W}^0 .

We conclude this subsection by the following lemma, which is useful in the proofs of the upcoming *a priori* estimates. It relies on the fact that \mathbb{P}^0 is finite-dimensional.

Lemma 6. *The norms $\|\cdot\|_I$, $\|v_1 \cdot\|_I$ and $\|(1 + |v|)^{1/2} \cdot\|_I$ are equivalent on \mathbb{P}^0 . Hence, there exists a constant $C_{\text{eq}} > 0$, depending on \mathbf{n} and \mathbf{m} , such that for any $\mathbf{g} \in \mathbb{P}^0$,*

$$(34) \quad \|v_1 \mathbf{g}\|_I \leq C_{\text{eq}} \|\mathbf{g}\|_I \quad \text{and} \quad \|(1 + |v|)^{1/2} \mathbf{g}\|_I \leq C_{\text{eq}} \|\mathbf{g}\|_I.$$

Moreover, from (20), we can deduce that there exists a constant $C_\chi > 0$ only depending on \mathbf{n} and \mathbf{m} , such that for any $\mathbf{f} \in \mathbb{D}$,

$$(35) \quad \|\mathbf{P}^0(v_1 \mathbf{f})\|_I^2 = \sum_{k=1}^{I+4} \left| \left\langle v_1 (\mathbf{nM})^{1/2} \chi^k, \mathbf{f} \right\rangle_I \right|^2 \leq C_\chi \|\mathbf{f}\|_I^2.$$

4.3. Handling the lower-order estimate on \mathbf{f}^1 through \mathbf{f} . Since we deal with \mathbf{f}^0 estimates separately, it is equivalent to treat \mathbf{f} or \mathbf{f}^1 to obtain an estimate on the microscopic part. We choose to proceed with \mathbf{f} . Let us scalarly multiply (8) by \mathbf{f} , integrate with respect to $t \in [0, T]$ and $x \in \mathbb{R}$ to obtain

$$(36) \quad \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_0^T + \int_0^T \int_{\mathbb{R}} \langle v_1 \partial_x \mathbf{f}, \mathbf{f} \rangle_I dx dt - \int_0^T \int_{\mathbb{R}} \langle \mathcal{L} \mathbf{f}, \mathbf{f} \rangle_I dx dt \leq \int_0^T \int_{\mathbb{R}} \langle \mathcal{N}(\mathbf{f}), \mathbf{f} \rangle_I dx dt.$$

The second term on the left-hand side of this equation vanishes by conservativity with respect to the x variable, since

$$\int_0^T \int_{\mathbb{R}} \langle v_1 \partial_x \mathbf{f}, \mathbf{f} \rangle_I dx dt = \frac{1}{2} \int_0^T \int_{\mathbb{R}} \frac{\partial}{\partial x} \left(\sum_{i=1}^I \int_{\mathbb{R}^3} v_1 f_i^2 dv \right) dx dt.$$

The term with \mathcal{L} can be lower-bounded thanks to the \mathcal{L} spectral gap estimate (16) and to (15) in the following way

$$\lambda \nu_0 \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2 \leq \lambda \|\nu^{1/2} \mathbf{f}^1\|_I^2 \leq -\langle \mathcal{L} \mathbf{f}^1, \mathbf{f}^1 \rangle_I = -\langle \mathcal{L} \mathbf{f}, \mathbf{f} \rangle_I.$$

Eventually, we deal with the term involving \mathcal{N} . We first notice that

$$\langle \mathcal{N}(\mathbf{f}), \mathbf{f} \rangle_I = \langle \mathcal{N}(\mathbf{f}), \mathbf{f}^1 \rangle_I = \langle (1 + |v|)^{-1/2} \mathcal{N}(\mathbf{f}), (1 + |v|)^{1/2} \mathbf{f}^1 \rangle_I,$$

since $\mathbf{P}^0 \mathcal{N}(\mathbf{f}) = 0$. Then, using (18), we have

$$(37) \quad |\langle \mathcal{N}(\mathbf{f}), \mathbf{f} \rangle_I| \leq C_\beta \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I^2 \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I \\ \leq 2C_\beta \left(\left\| (1 + |v|)^{1/2} \mathbf{f}^0 \right\|_I^2 + \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 \right) \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I.$$

Using the equivalence of norms (34) on \mathbb{P}^0 , we can find a constant $K_{\text{eq}} > 0$, depending on \mathbf{m} , \mathbf{n} , the cross sections and \mathcal{L} through its null space, such that (36) becomes

$$(38) \quad \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_0^T + \lambda \nu_0 \int_0^T \int_{\mathbb{R}} \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2 dx dt \\ \leq K_{\text{eq}} \int_0^T \int_{\mathbb{R}} \left(\|\mathbf{f}^0\|_I^2 + \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 \right) \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I dx dt.$$

Of course, we can see that, if we choose ε small enough in assumption (11), there will only remain a nonnegative contribution of a norm of \mathbf{f}^0 on the right-hand side of (38). This is why we have to focus now on an estimate on \mathbf{f}^0 .

4.4. Lower-order estimate on \mathbf{f}^0 : first steps. In order to deal with the estimate on \mathbf{f}^0 , we plug the expression (25) of \mathbf{f}^1 into the equation (23) satisfied by \mathbf{f}^0 , then scalarly multiply the new equation by \mathbf{f}^0 , and integrate with respect to t and x , to obtain

$$(39) \quad \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_0^T + L_1 + L_2 + L_3 + L_4 = 0,$$

where the term with $\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0), \mathbf{f}^0 \rangle_I$ vanishes, again by conservativity in x , and where we set

$$L_1 = \int_0^T \int_{\mathbb{R}} \left\langle v_1 \partial_x \bar{\mathcal{L}}^{-1} \partial_t \mathbf{f}^1, \mathbf{f}^0 \right\rangle_I dx dt = - \int_0^T \int_{\mathbb{R}} \left\langle \bar{\mathcal{L}}^{-1} \partial_t \mathbf{f}^1, \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I dx dt, \\ L_2 = \int_0^T \int_{\mathbb{R}} \left\langle v_1 \partial_x \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0), \mathbf{f}^0 \right\rangle_I dx dt = - \int_0^T \int_{\mathbb{R}} \left\langle \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0), \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I dx dt, \\ L_3 = \int_0^T \int_{\mathbb{R}} \left\langle v_1 \partial_x \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^1), \mathbf{f}^0 \right\rangle_I dx dt = - \int_0^T \int_{\mathbb{R}} \left\langle \partial_x \mathbf{f}^1, v_1 \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I dx dt, \\ L_4 = - \int_0^T \int_{\mathbb{R}} \left\langle v_1 \partial_x \bar{\mathcal{L}}^{-1} \mathcal{N}(\mathbf{f}), \mathbf{f}^0 \right\rangle_I dx dt = \int_0^T \int_{\mathbb{R}} \left\langle \mathcal{N}(\mathbf{f}), \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I dx dt.$$

Let us show how to handle these terms by performing the following preliminary computations on L_1, L_2, L_3, L_4 .

4.4.1. *Term L_1 .* Combining the equivalence of norms (34) on \mathbb{P}^0 with the boundedness of $\bar{\mathcal{L}}^{-1}$, and setting $K_1 = C_{\text{inv}}C_{\text{eq}} > 0$, we obtain

$$(40) \quad |L_1| \leq K_1 \int_0^T \int_{\mathbb{R}} \|\partial_t \mathbf{f}^1\|_I \|\partial_x \mathbf{f}^0\|_I \, dx \, dt.$$

4.4.2. *Term L_2 .* It is clear that $V = \mathbf{P}^1(v_1\mathbb{P}^0)$ is a finite-dimensional subspace of \mathbb{P}^1 . Consequently, $\bar{\mathcal{L}}|_V^{-1}$ is a bounded invertible operator, as well as its inverse. Hence, there exists a constant $C_V > 0$, only depending on \mathcal{L} , such that, for any $\mathbf{g} \in \mathbb{P}^0$,

$$\|\mathbf{P}^1(v_1\mathbf{g})\|_I \leq C_V \left\| \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1\mathbf{g}) \right\|_I.$$

Thus, thanks to the previous inequality and to the spectral gap estimate (16) on \mathcal{L} , we get

$$(41) \quad \begin{aligned} L_2 &= - \int_0^T \int_{\mathbb{R}} \left\langle \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0), \mathcal{L} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I \, dx \, dt \\ &\geq \lambda \int_0^T \int_{\mathbb{R}} \left\| \nu^{1/2} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I^2 \, dx \, dt \\ &\geq \lambda_2 \int_0^T \int_{\mathbb{R}} \left\| \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I^2 \, dx \, dt, \end{aligned}$$

where we chose $\lambda_2 = \lambda\nu_0/C_V > 0$.

4.4.3. *Term L_3 .* We apply the Cauchy-Schwarz inequality to yield

$$|L_3| \leq \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I \left\| v_1 \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I \, dx \, dt.$$

Then, introducing $C_{\mathbf{K}} > 0$ as the boundedness constant of the compact operator $\mathbf{K} = \mathcal{L} + \nu$, we notice that

$$\begin{aligned} \left\| v_1 \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I &\leq \frac{1}{\nu_0} \left\| \nu \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I \\ &\leq \frac{1}{\nu_0} \left(\left\| \mathbf{K} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I + \left\| \mathcal{L} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I \right) \leq \frac{C_{\text{eq}}}{\nu_0} (C_{\mathbf{K}} C_{\text{inv}} + 1) \|\partial_x \mathbf{f}^0\|_I, \end{aligned}$$

where we also used the boundedness of $\bar{\mathcal{L}}^{-1}$ and \mathbf{P}^1 (as a projector), and the norm equivalence argument on \mathbb{P}^0 , involving the constant C_{eq} . Setting $K_3 = C_{\text{eq}}(C_{\mathbf{K}}C_{\text{inv}} + 1)/\nu_0 > 0$, we obtain

$$(42) \quad |L_3| \leq K_3 \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I \|\partial_x \mathbf{f}^1\|_I \, dx \, dt,$$

4.4.4. *Term L_4 .* We first have to estimate

$$\begin{aligned} \left\langle \mathcal{N}(\mathbf{f}), \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I &= \left\langle (1 + |v|)^{-1/2} \mathcal{N}(\mathbf{f}), (1 + |v|)^{1/2} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\rangle_I \\ &\leq \left\| (1 + |v|)^{-1/2} \mathcal{N}(\mathbf{f}) \right\|_I \left\| (1 + |v|)^{1/2} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I. \end{aligned}$$

We treat the norm with \mathcal{N} using (18) and the same kind of argument as in (37), including norm equivalence on \mathbb{P}^0 . It ensures that

$$\left\| (1 + |v|)^{-1/2} \mathcal{N}(\mathbf{f}) \right\|_I \leq C_{\text{eq}} \left(\|\mathbf{f}^0\|_I^2 + \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2 \right).$$

Besides, the norm with $\bar{\mathcal{L}}^{-1}$ is treated in the same way as for L_3 , *i.e.*

$$\left\| (1 + |v|)^{1/2} \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I \leq \left\| (1 + |v|) \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \mathbf{f}^0) \right\|_I \leq K_3 \|\partial_x \mathbf{f}^0\|_I.$$

Therefore, setting $K_4 = C_{\text{eq}}K_3 > 0$, we can write

$$(43) \quad |L_4| \leq K_4 \int_0^T \int_{\mathbb{R}} \left(\|\mathbf{f}^0\|_I^2 + \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2 \right) \|\partial_x \mathbf{f}^0\|_I dx dt.$$

4.4.5. *Before we proceed.* Let us explain what the current situation on \mathbf{f}^0 is. Applying Young's inequality in (40) and (42) with a parameter $\delta_0 > 0$ to be chosen later, and plugging (40)–(43) in (39), we get

$$(44) \quad \begin{aligned} & \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_0^T + \lambda_2 \int_0^T \int_{\mathbb{R}} \|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 dx dt \\ & \leq K_1 \int_0^T \int_{\mathbb{R}} \left(\frac{1}{2\delta_0} \|\partial_t \mathbf{f}^1\|_I^2 + \frac{\delta_0}{2} \|\partial_x \mathbf{f}^0\|_I^2 \right) dx dt + K_3 \int_0^T \int_{\mathbb{R}} \left(\frac{\delta_0}{2} \|\partial_x \mathbf{f}^0\|_I^2 + \frac{1}{2\delta_0} \|\partial_x \mathbf{f}^1\|_I^2 \right) dx dt \\ & \quad + K_4 \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I \left(\|\mathbf{f}^0\|_I^2 + \|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2 \right) dx dt. \end{aligned}$$

We intend afterwards to combine (44) with the estimate (38) for \mathbf{f}^1 . All the terms are product of exactly two norms, apart from the ones coming from the nonlinear operator. This is why, in order to proceed, we shall need the same kind of smallness assumption as in [29], namely (11).

The term $\|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2$ appears in the left-hand side of (38), and in the right-hand side of both estimates in the terms coming from \mathcal{N} , with a multiplication by the arbitrary small parameter ε when using the smallness assumption. So they can be put on the left-hand side to obtain a still positive coefficient for $\|(1 + |v|)^{1/2} \mathbf{f}^1\|_I^2$. On the contrary, the time and space derivatives of \mathbf{f}^1 only appear in the right-hand sides of our estimates, and so does $\|\mathbf{f}^0\|_I^2$. We have no way to control them for the time being. Last, the space derivative of \mathbf{f}^0 appears on the right-hand side of (44) as its I -norm, and on the left-hand side in $\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I$, which will provide a helpful contribution.

Consequently, our next two steps are natural: deal with the norms of the time and space derivatives of \mathbf{f}^1 , and $\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I$.

4.5. **Handling the derivatives of \mathbf{f}^1 .** We proceed in the same way as in Subsection 4.3. We denote by ∂_\star any time or space partial differentiation. We differentiate (8) accordingly, scalarly multiply it by $\partial_\star \mathbf{f}$ and integrate with respect to t and x . Then, using the spectral gap (16) of \mathcal{L} , the growth property (15) of ν and the estimate (19) on $\partial_\star \mathcal{N}(\mathbf{f})$, we obtain, similarly to (38), the existence of $C_\star > 0$, depending on \mathbf{m} , \mathbf{n} and β and \mathcal{L} through its null space, such that

$$(45) \quad \begin{aligned} & \frac{1}{2} \int_{\mathbb{R}} \|\partial_\star \mathbf{f}\|_I^2 dx \Big|_0^T + \lambda \nu_0 \int_0^T \int_{\mathbb{R}} \|(1 + |v|)^{1/2} \partial_\star \mathbf{f}^1\|_I^2 dx dt \\ & \leq C_\star \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial_\star \mathbf{f}^1 \right\|_I \left(\|\partial_\star \mathbf{f}^0\|_I + \left\| (1 + |v|)^{1/2} \partial_\star \mathbf{f}^1 \right\|_I \right) dx dt. \end{aligned}$$

Let us deal with the time derivative, since no other term involving $\partial_t \mathbf{f}^0$ will appear in our estimates. In fact, using (23) and (35), we get, using again the Cauchy-Schwarz inequality,

$$\|\partial_t \mathbf{f}^0\|_I^2 = \|\mathbf{P}^0(v_1 \partial_x \mathbf{f})\|_I^2 \leq C_\chi \|\partial_x \mathbf{f}\|_I^2.$$

Consequently, (45) becomes, for the time derivative,

$$(46) \quad \begin{aligned} & \frac{1}{2} \int_{\mathbb{R}} \|\partial_t \mathbf{f}\|_I^2 dx \Big|_0^T + \lambda \nu_0 \int_0^T \int_{\mathbb{R}} \|(1 + |v|)^{1/2} \partial_t \mathbf{f}^1\|_I^2 dx dt \\ & \leq C'_\star \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial_t \mathbf{f}^1 \right\|_I \left(\|\partial_x \mathbf{f}^0\|_I + \left\| (1 + |v|)^{1/2} \partial_t \mathbf{f}^1 \right\|_I \right) dx dt, \end{aligned}$$

where $C'_\star = C_\star \max(C_\chi, 1)$.

We can thus observe that (45) and (46) have exactly the same structure and can be rewritten, for some constant $K_5 > 0$ only depending on \mathbf{n} , \mathbf{m} and β , as

$$(47) \quad \frac{1}{2} \int_{\mathbb{R}} \|\partial_\star \mathbf{f}\|_I^2 dx \Big|_0^T + C_0 \int_0^T \int_{\mathbb{R}} \|(1 + |v|)^{1/2} \partial_\star \mathbf{f}^1\|_I^2 dx dt \\ \leq K_5 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial_\star \mathbf{f}^1 \right\|_I \left(\|\partial_x \mathbf{f}^0\|_I + \left\| (1 + |v|)^{1/2} \partial_\star \mathbf{f}^1 \right\|_I \right) dx dt,$$

where we set $C_0 = \lambda \nu_0 > 0$.

4.6. Lower bound for $\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I$. Let us now focus on the term with $\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I$. The following lemma allows to estimate the L^2 -norm of $\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)$ in terms of $\partial_x \mathbf{f}^0$, up to a contribution in $\partial_x \mathbf{f}^1$ which can be as small as desired. We must emphasize that this result is the main improvement in the mixture case, compared to its monospecies counterparts from [27, estimate (2.22) p.185] and [29, Lemma 3.1 p.139]. Indeed, in the monospecies case, the lower bound of the term $\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I$ did not give any $\partial_x \rho$ contribution, only $\partial_x q$ and $\partial_x e$ parts of $\|\partial_x \mathbf{f}^0\|_I$.

Lemma 7. *There exists $\theta_0 \in (0, 1]$ such that, for any $\mathbf{f} \in \mathbb{D}$ and any $0 < \theta < \theta_0$,*

$$(48) \quad \int_0^T \int_{\mathbb{R}} \|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 dx dt \geq \gamma_\theta \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt \\ - \theta C_2 \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt + 2\theta \int_{\mathbb{R}} q \partial_x \ell dx \Big|_0^T$$

where $\gamma_\theta > 0$ only depends on \mathbf{n} , \mathbf{m} and θ , $C_2 > 0$ on \mathbf{n} and \mathbf{m} .

Proof. By orthogonality, we first have

$$\|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 = \|v_1 \partial_x \mathbf{f}^0\|_I^2 - \|\mathbf{P}^0(v_1 \partial_x \mathbf{f}^0)\|_I^2.$$

The first term of the right-hand side writes

$$\|v_1 \partial_x \mathbf{f}^0\|_I^2 = \sum_i \frac{1}{m_i} (\partial_x \rho_i)^2 + \frac{4}{\sqrt{6} \sum_j n_j} \sum_i \frac{\sqrt{n_i}}{m_i} (\partial_x \rho_i) (\partial_x e) \\ + \frac{7}{3} \frac{1}{\sum_j n_j} \sum_i \frac{n_i}{m_i} (\partial_x e)^2 + \frac{\sum_j n_j}{\sum_j n_j m_j} \left[3 (\partial_x q^1)^2 + (\partial_x q^2)^2 + (\partial_x q^3)^2 \right].$$

It can become a sum of square quantities, as

$$(49) \quad \|v_1 \partial_x \mathbf{f}^0\|_I^2 = \sum_i \frac{1}{m_i} \left(\partial_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6} \sum_j n_j} \partial_x e \right)^2 + \frac{5}{3} \frac{1}{\sum_j n_j} \sum_i \frac{n_i}{m_i} (\partial_x e)^2 \\ + \frac{\sum_j n_j}{\sum_j n_j m_j} \left[3 (\partial_x q^1)^2 + (\partial_x q^2)^2 + (\partial_x q^3)^2 \right].$$

The term with \mathbf{P}^0 decomposes on our basis of \mathbb{P}^0 into

$$\mathbf{P}^0(v_1 \partial_x \mathbf{f}^0) = \partial_x q^1 \sum_i \frac{\sqrt{n_i}}{\sqrt{\sum_j n_j m_j}} \chi^i (\mathbf{nM})^{1/2} + \partial_x q^1 \frac{2\sqrt{\sum_j n_j}}{\sqrt{6} \sum_j n_j m_j} \chi^{I+4} (\mathbf{nM})^{1/2} \\ + \frac{1}{\sqrt{\sum_j n_j m_j}} \left(\sum_i \sqrt{n_i} \partial_x \rho_i + \frac{2\sqrt{\sum_j n_j}}{\sqrt{6}} \partial_x e \right) \chi^{I+1} (\mathbf{nM})^{1/2}.$$

It is thus easy to check that

$$(50) \quad \|\mathbf{P}^0(v_1 \partial_x \mathbf{f}^0)\|_I^2 = (\partial_x \ell)^2 + \frac{5}{3} \frac{\sum n_j}{\sum n_j m_j} (\partial_x q^1)^2.$$

Therefore, we immediately get

$$(51) \quad \|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 = \sum_i \frac{1}{m_i} \left(\partial_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \partial_x e \right)^2 + \frac{5}{3} \frac{1}{\sum n_j} \left(\sum \frac{n_j}{m_j} \right) (\partial_x e)^2 \\ + \frac{\sum n_j}{\sum n_j m_j} \left[\frac{4}{3} (\partial_x q^1)^2 + (\partial_x q^2)^2 + (\partial_x q^3)^2 \right] - (\partial_x \ell)^2.$$

Thanks to the Cauchy-Schwarz inequality, we have

$$\left(\sum_i \sqrt{n_i} \partial_x \rho_i + \frac{2\sqrt{\sum n_j}}{\sqrt{6}} \partial_x e \right)^2 = \left[\sum_i \left(\sqrt{n_i} \partial_x \rho_i + \frac{2n_i}{\sqrt{6 \sum n_j}} \partial_x e \right) \right]^2 \\ \leq \left(\sum_j n_j m_j \right) \sum_i \frac{1}{m_i} \left(\partial_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \partial_x e \right)^2,$$

which implies that

$$(\partial_x \ell)^2 \leq \sum_i \frac{1}{m_i} \left(\partial_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \partial_x e \right)^2.$$

Let $\theta \in (0, 1)$. Then we split $(\partial_x \ell)^2$ in (51) into the sums of itself respectively multiplied by θ and $(1 - \theta)$. Applying the previous Cauchy-Schwarz inequality, it follows from (51) that

$$(52) \quad \|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 \geq \theta \sum_i \frac{1}{m_i} \left(\partial_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \partial_x e \right)^2 + \frac{5}{3} \frac{1}{\sum n_j} \left(\sum \frac{n_j}{m_j} \right) (\partial_x e)^2 \\ + \frac{\sum n_j}{\sum n_j m_j} \left[\frac{4}{3} (\partial_x q^1)^2 + (\partial_x q^2)^2 + (\partial_x q^3)^2 \right] - \theta (\partial_x \ell)^2.$$

The term $-\theta |\partial_x \ell|^2$ appearing in (52) is absorbed thanks to the conservation laws. More precisely, multiplying (27) by $\partial_x \ell$ and integrating with respect to t and x gives

$$\int_0^T \int_{\mathbb{R}} |\partial_x \ell|^2 dx dt = - \int_0^T \int_{\mathbb{R}} \partial_t q^1 \partial_x \ell dx dt - \int_0^T \int_{\mathbb{R}} \left\langle \partial_x \mathbf{f}^1, v_1 \chi^{I+1} (\mathbf{nM})^{1/2} \right\rangle_I \partial_x \ell.$$

Then, using integrations by parts, one with respect to t , one for x , for the first term on the right-hand side, and Cauchy-Schwarz and Young's inequalities, we obtain with (35) that

$$\int_0^T \int_{\mathbb{R}} |\partial_x \ell|^2 dx dt \leq - \int_0^T \int_{\mathbb{R}} \partial_t \ell \partial_x q^1 dx dt - \int_{\mathbb{R}} q^1 \partial_x \ell dx \Big|_0^T \\ + \frac{1}{2} \int_0^T \int_{\mathbb{R}} |\partial_x \ell|^2 dx dt + \frac{C_\chi}{2} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt,$$

ensuring that

$$(53) \quad \frac{1}{2} \int_0^T \int_{\mathbb{R}} |\partial_x \ell|^2 dx dt + \int_{\mathbb{R}} q^1 \partial_x \ell dx \Big|_0^T \leq \left| \int_0^T \int_{\mathbb{R}} \partial_t \ell \partial_x q^1 dx dt \right| + \frac{C_\chi}{2} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt.$$

Further, we multiply (31) by $\partial_x q^1$, integrate with respect to x and t and use again Cauchy-Schwarz and Young's inequalities as well as (35), to get that there exists a some constant $K_\chi > 0$ depending only on \mathbf{n} and \mathbf{m} such that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}} \partial_t \ell \partial_x q^1 dx dt &= \frac{1}{\sum n_j m_j} \left(\sum n_j^{3/2} + \frac{2}{3} \sum n_j \right) \int_0^T \int_{\mathbb{R}} |\partial_x q^1|^2 dx dt \\ &\quad + \frac{1}{2} \int_0^T \int_{\mathbb{R}} |\partial_x q^1|^2 dx dt + \frac{K_\chi}{2} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt. \end{aligned}$$

This estimate, combined with (53), gives, after multiplication by 2θ ,

$$\begin{aligned} (54) \quad &\theta \int_0^T \int_{\mathbb{R}} |\partial_x \ell|^2 dx dt + 2\theta \int_{\mathbb{R}} q^1 \partial_x \ell dx \Big|_0^T \\ &\leq 2\theta \left(\frac{\sum n_j^{3/2} + \frac{2}{3} \sum n_j}{\sum n_j m_j} + \frac{1}{2} \right) \int_0^T \int_{\mathbb{R}} |\partial_x q^1|^2 dx dt + \theta(C_\chi + K_\chi) \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt. \end{aligned}$$

Setting

$$(55) \quad \theta_0 = \min \left\{ 1, \frac{\sum n_j}{3 \sum n_j m_j + 6 \sum n_j^{3/2} + 4 \sum n_j} \right\} \in (0, 1],$$

we can come back to (52) with a fixed $0 < \theta < \theta_0$, so that the term in $\|\partial_x q^1\|^2$ in (54) can be absorbed by the following term, taken in (52),

$$\frac{\sum n_j}{\sum n_j m_j} \int_0^T \int_{\mathbb{R}} \frac{1}{3} (\partial_x q^1)^2 dx dt.$$

Let us now introduce $\mathcal{G} : \mathbb{P}^0 \rightarrow \mathbb{R}_+$,

$$\mathbf{f}^0 \mapsto \theta \sum_i \frac{1}{m_i} \left(\rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} e \right)^2 + \frac{5}{3} \frac{1}{\sum n_j} \left(\sum \frac{n_j}{m_j} \right) e^2 + \frac{\sum n_j}{\sum n_j m_j} \left[(q^1)^2 + (q^2)^2 + (q^3)^2 \right],$$

which defines a positive definite quadratic form on the finite-dimensional subspace \mathbb{P}^0 , thus equivalent to $\|\cdot\|_I^2$ on \mathbb{P}^0 . Therefore, there exists a constant $\gamma_\theta > 0$, only depending on \mathbf{n} , \mathbf{m} and θ , such that, for any $\mathbf{f}^0 \in \mathbb{P}^0$,

$$\mathcal{G}(\mathbf{f}^0) \geq \gamma_\theta \|\mathbf{f}^0\|_I^2.$$

All the previous considerations lead to the following estimate

$$\begin{aligned} (56) \quad &\int_0^T \int_{\mathbb{R}} \|\mathbf{P}^1(v_1 \partial_x \mathbf{f}^0)\|_I^2 dx dt \geq \gamma_\theta \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt \\ &\quad - \theta(C_\chi + K_\chi) \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^1\|_I^2 dx dt + 2\theta \int_{\mathbb{R}} q^1 \partial_x \ell \Big|_0^T. \end{aligned}$$

We obtain the required estimate (48) by setting $C_2 = (C_\chi + K_\chi)$. \square

Remark 2. *Let us emphasize that the previous lemma holds in higher dimensions. We explain in Appendix C how to handle the proof of Lemma 7 in a three-dimensional setting.*

4.7. Estimate using an antiderivative of \mathbf{f}^0 . As we already pointed out at the end of Subsection 4.4, we still fail to control $\|\mathbf{f}^0\|_I$. Since we work in a one-dimensional space setting, following the strategy of [29], we write an estimate on the antiderivative \mathbf{W}^0 of \mathbf{f}^0 .

We first integrate (23) with respect to the space variable between $-\infty$ and $x \in \mathbb{R}$, so that

$$(57) \quad \partial_t \mathbf{W}^0 + \mathbf{P}^0(v_1 \partial_x \mathbf{W}^0) + \mathbf{P}^0(v_1 \mathbf{f}^1) = 0.$$

We then scalarly multiply (57) by \mathbf{W}^0 and integrate with respect to t and x . Since the second term has a conservative form in x and using the expression (25) of \mathbf{f}^1 , we get

$$(58) \quad \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_0^T + \hat{L}_1 + \hat{L}_2 + \hat{L}_3 + \hat{L}_4 = 0,$$

where we set

$$\begin{aligned}\hat{L}_1 &= \int_0^T \int_{\mathbb{R}} \langle \mathbf{P}^0(v_1 \bar{\mathcal{L}}^{-1} \partial_t \mathbf{f}^1), \mathbf{W}^0 \rangle_I dx dt \\ \hat{L}_2 &= - \int_0^T \int_{\mathbb{R}} \langle \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \mathbf{f}^0), \mathbf{P}^1(v_1 \mathbf{f}^0) \rangle_I dx dt \\ \hat{L}_3 &= - \int_0^T \int_{\mathbb{R}} \langle \mathbf{P}^1(v_1 \mathbf{f}^1), \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \mathbf{f}^0) \rangle_I dx dt \\ \hat{L}_4 &= \int_0^T \int_{\mathbb{R}} \langle \mathcal{N}(\mathbf{f}), \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \mathbf{W}^0) \rangle_I dx dt.\end{aligned}$$

The term \hat{L}_2 in (58) is treated in the same way as the corresponding term L_2 in Subsections 4.4.2 and 4.6. Indeed, we can prove a result of the same kind as Lemma 7 involving \mathbf{f}_0 instead of $\partial_x \mathbf{f}^0$, by using, among others properties, (32)–(33). Hence, we can write, for any $\theta \in (0, \theta_0)$,

$$\hat{L}_2 \geq \lambda_2 \gamma \theta \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt - \lambda_2 \theta C_2 \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^1\|_I^2 dx dt + 2\lambda_2 \theta \int_{\mathbb{R}} Q^1 \ell dx \Big|_0^T.$$

The term \hat{L}_3 has the same structure as L_3 in 4.4.3. Consequently, involving the same constant K_3 as in (42), we have

$$|\hat{L}_3| \leq K_3 \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I \|\mathbf{f}^1\|_I dx dt.$$

The term \hat{L}_4 with the nonlinear operator is also treated as L_4 in 4.4.4 to yield

$$|\hat{L}_4| \leq K_4 \int_0^T \int_{\mathbb{R}} \|\mathbf{W}^0\|_I \left(\|\mathbf{f}^0\|_I^2 + \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 \right) dx dt.$$

The main difference with Subsection 4.4 comes from the treatment of \hat{L}_1 . Integrating by parts with respect to t , and then replacing $\partial_t \mathbf{W}^0$ by its expression in (57), yields

$$|\hat{L}_1| \leq \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^1\|_I \left\| \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \mathbf{P}^0(v_1 \mathbf{f})) \right\|_I dx dt + \left| \int_{\mathbb{R}} \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I dx \right|_0^T.$$

Thanks to the boundedness of $\bar{\mathcal{L}}^{-1}$ and the norm equivalence argument on \mathbb{P}^0 (34), and computing directly $\mathbf{P}^0(v_1 \mathbf{f})$ as in (35), we obtain

$$\left\| \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \mathbf{P}^0(v_1 \mathbf{f})) \right\|_I \leq C_{\text{inv}} C_{\text{eq}} C_{\chi} \|\mathbf{f}\|_I.$$

Setting $\hat{K}_1 = C_{\text{inv}} C_{\text{eq}} C_{\chi} > 0$, we get

$$|\hat{L}_1| \leq \hat{K}_1 \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^1\|_I \|\mathbf{f}\|_I dx dt + \left| \int_{\mathbb{R}} \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I dx \right|_0^T.$$

Let us sum up the situation on \mathbf{W}^0 . Taking into account the estimates on \hat{L}_1 , \hat{L}_2 , \hat{L}_3 , \hat{L}_4 in (58), and applying Young's inequality in the estimates on \hat{L}_1 and \hat{L}_3 with a parameter $\delta_1 > 0$ to be chosen later, we get

$$\begin{aligned}(59) \quad & \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_0^T + \lambda_2 \gamma \theta \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt - \theta \lambda_2 C_2 \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^1\|_I^2 dx dt + 2\lambda_2 \theta \int_{\mathbb{R}} Q^1 \ell dx \Big|_0^T \\ & \leq \hat{K}_1 \int_0^T \int_{\mathbb{R}} \left(\frac{1}{2\delta_1} \|\mathbf{f}^1\|_I^2 + \frac{\delta_1}{2} \|\mathbf{f}\|_I^2 \right) dx dt + K_3 \int_0^T \int_{\mathbb{R}} \left(\frac{\delta_1}{2} \|\mathbf{f}^0\|_I^2 + \frac{1}{2\delta_1} \|\mathbf{f}^1\|_I^2 \right) dx dt \\ & \quad + K_4 \int_0^T \int_{\mathbb{R}} \|\mathbf{W}^0\|_I \left(\|\mathbf{f}^0\|_I^2 + \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 \right) dx dt + \left| \int_{\mathbb{R}} \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I dx \right|_0^T.\end{aligned}$$

4.8. Proof of the global lower-order estimate. We now carefully mix all the estimates we got so far, starting with the use of the smallness assumption (11). We also have to treat pointwise in time integrals. Recall that all terms at initial time are put in a generic term denoted by $\mathcal{I}(0)$. Terms at time T must be handled more shrewdly.

First, imposing $\varepsilon \leq C_0/(2K_{\text{eq}})$, (38) becomes

$$(60) \quad \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{C_0}{2} \int_0^T \int_{\mathbb{R}} \|(1+|v|)^{1/2} \mathbf{f}^1\|_I^2 dx dt \leq \varepsilon K_{\text{eq}} \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt + \mathcal{I}(0).$$

In the same way, if $\varepsilon \leq C_0/(3K_5)$, (47) can be rewritten as

$$(61) \quad \frac{1}{2} \int_{\mathbb{R}} \|\partial_\star \mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{C_0}{2} \int_0^T \int_{\mathbb{R}} \|(1+|v|)^{1/2} \partial_\star \mathbf{f}^1\|_I^2 dx dt \leq \frac{\varepsilon K_5}{2} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt + \mathcal{I}(0).$$

Let us now deal with the estimates on elements of \mathbb{P}^0 . In (59), we can find three terms at time T , which are

$$\frac{1}{2} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx + 2\lambda_2 \theta \int_{\mathbb{R}} Q^1 \ell dx - \int_{\mathbb{R}} \left| \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I \right| dx.$$

We then notice that, if we set $\delta_2 = (C_{\text{inv}} C_{\text{eq}})^2$,

$$\int_{\mathbb{R}} \left| \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I \right| dx \leq C_{\text{inv}} C_{\text{eq}} \int_{\mathbb{R}} \|\mathbf{f}^1\|_I \|\mathbf{W}^0\|_I dx \leq \frac{1}{4} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx + \delta_2 \int_{\mathbb{R}} \|\mathbf{f}^1\|_I^2 dx.$$

Moreover, we can write

$$-2 \int_{\mathbb{R}} Q^1 \ell dx \leq \int_{\mathbb{R}} (Q^1)^2 dx + \int_{\mathbb{R}} \ell^2 dx \leq \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx + \int_{\mathbb{R}} \ell^2 dx.$$

Hence, for $\theta \leq (8\lambda_2)^{-1}$, we get

$$-2\theta\lambda_2 \int_{\mathbb{R}} Q^1 \ell dx \leq \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx + \theta\lambda_2 \int_{\mathbb{R}} \ell^2 dx.$$

All in all, the terms at time T in (59) satisfy, for any $0 < \theta \leq (8\lambda_2)^{-1}$,

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx + 2\lambda_2 \theta \int_{\mathbb{R}} Q^1 \ell dx - \int_{\mathbb{R}} \left| \langle \bar{\mathcal{L}}^{-1} \mathbf{f}^1, v_1 \mathbf{W}^0 \rangle_I \right| dx \\ \geq \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx - \delta_2 \int_{\mathbb{R}} \|\mathbf{f}^1\|_I^2 dx - \theta\lambda_2 \int_{\mathbb{R}} |\ell|^2 dx. \end{aligned}$$

Let us set $\hat{C}_1 = \lambda_2 \gamma \theta$, $\delta_1 = \hat{C}_1 (\hat{K}_1 + K_3)^{-1}/2$, $\hat{C}_2 = \theta_0 \lambda_2 C_2 + \hat{K}_1 + \frac{\hat{K}_1 + K_3}{2\delta_1} + K_4$. Using that $\|\mathbf{f}\|_I^2 = \|\mathbf{f}^0\|_I^2 + \|\mathbf{f}^1\|_I^2$ and the previous inequality in (59), we obtain

$$(62) \quad \begin{aligned} \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_{t=T} + \frac{\hat{C}_1}{2} \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt \\ \leq \hat{C}_2 \int_0^T \int_{\mathbb{R}} \left\| (1+|v|)^{1/2} \mathbf{f}^1 \right\|_I^2 dx dt + \theta\lambda_2 \int_{\mathbb{R}} |\ell|^2 dx \Big|_{t=T} + \delta_2 \int_{\mathbb{R}} \|\mathbf{f}^1\|_I^2 dx \Big|_{t=T} + \mathcal{I}(0), \end{aligned}$$

where we imposed $\varepsilon \leq \min(\hat{C}_1/(4K_4), 1)$.

Eventually, we similarly tackle the estimate (44). We apply (48), and choose $\delta_0 = \hat{C}_1 (K_1 + K_3)^{-1}/2$, $\hat{C}_3 = \max(\theta_0 C_2 + K_3/(2\delta_0), K_1/(2\delta_0))$, to obtain

$$(63) \quad \begin{aligned} \frac{1}{4} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_{t=T} + \frac{\hat{C}_1}{2} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt \\ \leq \hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1+|v|)^{1/2} \partial_x \mathbf{f}^1 \right\|_I^2 dx dt + \hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1+|v|)^{1/2} \partial_t \mathbf{f}^1 \right\|_I^2 dx dt \end{aligned}$$

$$+ \varepsilon K_4 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 dx dt + \theta \lambda_2 \int_{\mathbb{R}} |\partial_x \ell|^2 dx \Big|_{t=T} + \mathcal{I}(0),$$

with $\varepsilon \leq \hat{C}_1/4$.

Now, let us set

$$\alpha_0 = \max \left(2\delta_2, \frac{4\hat{C}_2}{C_0} \right) > 0, \quad \alpha_1 = \frac{4\hat{C}_3}{C_0} > 0.$$

The remaining pointwise in time terms are handled by choosing θ small enough, say $\theta \leq \theta_1$, where $\theta_1 > 0$ only depends on the problem data, so that

$$\theta \lambda_2 \int_{\mathbb{R}} |\ell|^2 dx \Big|_{t=T} \leq \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_{t=T}, \quad \theta \lambda_2 \int_{\mathbb{R}} |\partial_x \ell|^2 dx \Big|_{t=T} \leq \frac{\alpha_1}{4} \int_{\mathbb{R}} \|\partial_x \mathbf{f}\|_I^2 dx \Big|_{t=T}.$$

We then multiply (60) by α_0 and both estimates (61) (for derivatives in t and x) by α_1 , to jointly add them to (62)–(63). Assuming moreover that

$$\varepsilon \leq \min \left(\frac{\hat{C}_2}{2K_4}, \frac{\hat{C}_1}{2\alpha_1 K_5}, \frac{\hat{C}_1}{4\alpha_0 K_{\text{eq}}} \right),$$

we can finally write

$$\begin{aligned} & \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{W}^0\|_I^2 dx \Big|_{t=T} + \frac{1}{8} \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx \Big|_{t=T} + \frac{\alpha_0}{4} \int_{\mathbb{R}} \|\mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{\alpha_1}{4} \int_{\mathbb{R}} \|\partial_x \mathbf{f}\|_I^2 dx \Big|_{t=T} \\ & + \frac{\alpha_1}{2} \int_{\mathbb{R}} \|\partial_t \mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{\hat{C}_1}{4} \int_0^T \int_{\mathbb{R}} \|\mathbf{f}^0\|_I^2 dx dt + \frac{\hat{C}_1}{4} \int_0^T \int_{\mathbb{R}} \|\partial_x \mathbf{f}^0\|_I^2 dx dt \\ & + \frac{\hat{C}_2}{2} \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \mathbf{f}^1 \right\|_I^2 dx dt + \hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial_x \mathbf{f}^1 \right\|_I^2 dx dt \\ & + \hat{C}_3 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial_t \mathbf{f}^1 \right\|_I^2 dx dt \leq \mathcal{I}(0), \end{aligned}$$

which yields (12). To summarize on ε , we set

$$\varepsilon_0 = \min \left(\frac{\hat{C}_2}{2K_4}, \frac{\hat{C}_1}{2\alpha_1 K_5}, \frac{\hat{C}_1}{4\alpha_0 K_{\text{eq}}}, \frac{\hat{C}_1}{4}, \frac{\hat{C}_1}{4K_4}, \frac{C_0}{3K_5}, \frac{C_0}{2K_{\text{eq}}}, 1 \right).$$

The previous inequality holds as soon as $\varepsilon \leq \varepsilon_0$. This concludes the proof of Proposition 2.

5. ELEMENTS OF PROOF FOR THE HIGHER-ORDER ESTIMATE

Unlike what we did in the previous section about the lower-order estimate, for the sake of simplicity, we shall now use the corresponding smallness assumption (13) as soon as possible in our computations for the higher-order estimate.

Estimate involving $\partial^p \mathbf{f}^1$, $1 \leq |p| \leq 5$. We take the ∂^p derivative of the linearized Boltzmann equation (8), scalarly multiply it by $\partial^p \mathbf{f}$ and integrate with respect to t and x , to obtain

$$(64) \quad \frac{1}{2} \int_{\mathbb{R}} \|\partial^p \mathbf{f}\|_I^2 dx \Big|_0^T - \int_0^T \int_{\mathbb{R}} \langle \mathcal{L} \partial^p \mathbf{f}, \partial^p \mathbf{f} \rangle_I dx dt = \int_0^T \int_{\mathbb{R}} \langle \partial^p \mathcal{N}(\mathbf{f}), \partial^p \mathbf{f} \rangle_I dx dt.$$

In the second integral, we use the spectral gap property (16) of \mathcal{L} , so that

$$- \int_0^T \int_{\mathbb{R}} \langle \mathcal{L} \partial^p \mathbf{f}, \partial^p \mathbf{f} \rangle_I dx dt \geq C_0 \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial^p \mathbf{f}^1 \right\|_I^2 dx dt.$$

Let us focus on the term with the nonlinear operator \mathcal{N} . We first notice that

$$\partial^p \mathcal{N}(\mathbf{f}) = (\mathbf{nM})^{-1/2} \sum_{0 \leq |p'| \leq |p|} \binom{|p|}{|p'|} \mathbf{Q} \left((\mathbf{nM})^{1/2} \partial^{p'} \mathbf{f}, (\mathbf{nM})^{1/2} \partial^{p-p'} \mathbf{f} \right).$$

Thanks to Lemma 8, we get

$$(65) \quad \left\| (1 + |v|)^{-1/2} \partial^p \mathcal{N}(\mathbf{f}) \right\|_I \leq C_\beta \sum_{0 \leq |p'| \leq |p|} \binom{|p|}{|p'|} \left\| (1 + |v|)^{1/2} \partial^{p'} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial^{p-p'} \mathbf{f} \right\|_I.$$

For $3 \leq |p| \leq 5$, if the sub-index p' of p satisfies $|p'| \leq 2$, the smallness assumption (13) can be used for the term $\|(1 + |v|)^{1/2} \partial^{p'} \mathbf{f}\|_I$, and if $3 \leq |p'| \leq 5$, then $|p - p'| \leq 2$, and the term $\|(1 + |v|)^{1/2} \partial^{p-p'} \mathbf{f}\|_I$ can be handled through (13). Hence, in any case (when $|p| \leq 2$, it is straightforward), we can write, for some constant $C_p > 0$ only depending on \mathbf{m} , \mathbf{n} , β and p ,

$$\left\| (1 + |v|)^{-1/2} \partial^p \mathcal{N}(\mathbf{f}) \right\|_I \leq C_p \varepsilon \sum_{|p'| \leq |p|} \left\| (1 + |v|)^{1/2} \partial^{p'} \mathbf{f} \right\|_I.$$

Consequently, we have, for any p , $1 \leq |p| \leq 5$,

$$\int_0^T \int_{\mathbb{R}} |\langle \partial^p \mathcal{N}(\mathbf{f}), \partial^p \mathbf{f} \rangle_I| dx dt \leq C_p \varepsilon \int_0^T \int_{\mathbb{R}} \sum_{|p'| \leq |p|} \left\| (1 + |v|)^{1/2} \partial^{p'} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \partial^p \mathbf{f} \right\|_I dx dt.$$

All in all, summing (64) for all indices p , $1 \leq |p| \leq 5$, we get, for ε small enough,

$$(66) \quad \frac{1}{2} \sum_{1 \leq |p| \leq 5} \int_{\mathbb{R}} \|\partial^p \mathbf{f}\|_I^2 dx \Big|_{t=T} + \frac{C_0}{2} \sum_{1 \leq |p| \leq 5} \int_0^T \int_{\mathbb{R}} \left\| (1 + |v|)^{1/2} \partial^p \mathbf{f}^1 \right\|_I^2 dx dt \\ \leq C \varepsilon \sum_{1 \leq |p| \leq 5} \int_0^T \int_{\mathbb{R}} \|\partial^p \mathbf{f}^0\|_I^2 dx dt + \mathcal{I}(0),$$

where $C > 0$ only depends on the data of the problem.

Estimate involving $\partial^r \mathbf{f}^0$, $1 \leq |r| \leq 4$. We take the ∂^r derivative of (23) and observe that $\partial^r \mathbf{f}^0$ satisfies exactly the same kind of equation as \mathbf{f}^0 itself. Consequently, (44) also holds for $\partial^r \mathbf{f}^0$ instead of \mathbf{f}^0 , with some changes only on the nonlinear term \hat{L}_4^r ,

$$(67) \quad \frac{1}{2} \int_{\mathbb{R}} \|\partial^r \mathbf{f}^0\|_I^2 dx \Big|_0^T + \lambda_2 \int_0^T \int_{\mathbb{R}} \|\mathbf{P}^1(v_1 \partial_x \partial^r \mathbf{f}^0)\|_I^2 dx dt \\ \leq K_1 \int_0^T \int_{\mathbb{R}} \left(\frac{1}{2\delta_0} \|\partial_t \partial^r \mathbf{f}^1\|_I^2 + \frac{\delta_0}{2} \|\partial_x \partial^r \mathbf{f}^0\|_I^2 \right) dx dt \\ + K_3 \int_0^T \int_{\mathbb{R}} \left(\frac{\delta_0}{2} \|\partial_x \partial^r \mathbf{f}^0\|_I^2 + \frac{1}{2\delta_0} \|\partial_x \partial^r \mathbf{f}^1\|_I^2 \right) dx dt + |\hat{L}_4^r|,$$

where

$$\hat{L}_4^r = - \int_0^T \int_{\mathbb{R}} \left\langle v_1 \partial_x \bar{\mathcal{L}}^{-1} \partial^r \mathcal{N}(\mathbf{f}), \partial^r \mathbf{f}^0 \right\rangle_I dx dt = \int_0^T \int_{\mathbb{R}} \left\langle \partial^r \mathcal{N}(\mathbf{f}), \bar{\mathcal{L}}^{-1} \mathbf{P}^1(v_1 \partial_x \partial^r \mathbf{f}^0) \right\rangle_I dx dt.$$

Using the smallness assumption (13) and (65), we get

$$(68) \quad |\hat{L}_4^r| \leq C_r K_3 \varepsilon \left[\sum_{|p'| \leq |r|} \left\| (1 + |v|)^{1/2} \partial^{p'} \mathbf{f} \right\|_I \right] \|\partial_x \partial^r \mathbf{f}^0\|_I.$$

The second term in (67) is handled thanks to Lemma 7, leading to

$$(69) \quad \int_0^T \int_{\mathbb{R}} \|\mathbf{P}^1(v_1 \partial_x \partial^r \mathbf{f}^0)\|_I^2 dx dt \geq \gamma_\theta \int_0^T \int_{\mathbb{R}} \|\partial_x \partial^r \mathbf{f}^0\|_I^2 dx dt \\ - \theta C_2 \int_0^T \int_{\mathbb{R}} \|\partial_x \partial^r \mathbf{f}^1\|_I^2 dx dt + 2\theta \int_{\mathbb{R}} \partial^r q \partial_x \partial^r \ell dx \Big|_0^T.$$

Proof of Theorem 3. The same reasoning as in the previous section provides, for θ and ε small enough, the estimate (14), by combining the lower-order estimate (12), the estimate (66) on the derivatives of \mathbf{f}^1 , and the estimate on the derivatives of $\partial_x \mathbf{f}^0$, itself obtained from (67)–(69).

APPENDIX A. ESTIMATE ON \mathbf{Q}

In order to deal with the terms involving $\mathcal{N}(\mathbf{f})$, we need the following result. It requires the hard-sphere assumption (4) and its proof is provided here, despite its similarity to the one in [19, Lemma A.1] (see also [29, Lemma B.1]), for the sake of completeness.

Lemma 8. *Assuming that all the cross sections satisfy the hard-sphere assumption (4), there exists $C_\beta > 0$, only depending on \mathbf{m} , \mathbf{n} and the cross sections, such that, for any $\mathbf{f}, \mathbf{g} \in \mathbb{D}$,*

$$(70) \quad \left\| (1 + |v|)^{-1/2} (\mathbf{nM})^{-1/2} \mathbf{Q} \left((\mathbf{nM})^{1/2} \mathbf{f}, (\mathbf{nM})^{1/2} \mathbf{g} \right) \right\|_I \leq C_\beta \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I \left\| (1 + |v|)^{1/2} \mathbf{g} \right\|_I.$$

Proof. Denote by A the left-hand side of (70). We can write, thanks to the Cauchy-Schwarz inequality and the hard-sphere assumption (4) on the cross sections,

$$A^2 = \sum_i \int_{\mathbb{R}^3} (1 + |v|)^{-1} (n_i M_i)^{-1} Q_i \left((\mathbf{nM})^{1/2} \mathbf{f}, (\mathbf{nM})^{1/2} \mathbf{g} \right)^2 dv \\ \leq I \sum_{i,j} n_j \int_{\mathbb{R}^3} (1 + |v|)^{-1} \left[\iint_{\mathbb{R}^3 \times S^2} M_j(v_*)^{1/2} (f_i(v') g_j(v'_*) - f_i(v) g_j(v_*)) \mathcal{B}_{ij} d\omega dv_* \right]^2 dv \\ \leq 2I \sum_{i,j} n_j \int_{\mathbb{R}^3} \frac{\beta_{ij}}{1 + |v|} \left[\left(\iint_{\mathbb{R}^3 \times S^2} M_j(v_*)^{1/2} f_i(v') g_j(v'_*) |(v - v_*) \cdot \omega| d\omega dv_* \right)^2 \right. \\ \left. + \left(\iint_{\mathbb{R}^3 \times S^2} M_j(v_*)^{1/2} f_i(v) g_j(v_*) |(v - v_*) \cdot \omega| d\omega dv_* \right)^2 \right] dv.$$

Denote $\beta = \max \beta_{ij} > 0$. Noticing that

$$|(v - v_*) \cdot \omega| \leq (|v| + |v_*|),$$

and that $v_* \mapsto M_j(v_*)^{1/2} |v_*|$ is a bounded function on \mathbb{R}^3 , there exists a constant $C_{\mathbf{n},\beta} > 0$, only depending on \mathbf{n} and β , such that A is upper-bounded by

$$C_{\mathbf{n},\beta} \sum_{i,j} \left[\iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times S^2} f_i(v')^2 g_j(v'_*)^2 (1 + |v|) d\omega dv_* dv + \iint_{\mathbb{R}^3 \times \mathbb{R}^3} f_i(v)^2 g_j(v_*)^2 (1 + |v|) dv_* dv \right]$$

Let us focus on the first addend, since the second one clearly equals $C_{\mathbf{n},\beta} \left\| (1 + |v|)^{1/2} \mathbf{f} \right\|_I^2 \left\| \mathbf{g} \right\|_I^2$ thanks to the Fubini theorem. We perform the change of variables $(v, v_*, \omega) \mapsto (v', v'_*, \omega)$ in the integral, which becomes

$$\iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times S^2} f_i(v)^2 g_j(v_*)^2 (1 + |v'|) d\omega dv_* dv.$$

The collision rules (3) then ensure that, for some constant $C_{\mathbf{m}} > 0$ only depending on \mathbf{m} (uniform with respect to the indices i and j),

$$|v'| \leq |v| + \frac{2m_j}{m_i + m_j} |v_*| \leq C_{\mathbf{m}} (|v'| + |v'_*|), \quad \forall v, v_* \in \mathbb{R}^3,$$

so that, up to a value change of $C_m > 0$, still only depending on \mathbf{m} ,

$$1 + |v'| \leq C_m(2 + |v| + |v_*|), \quad \forall v, v_* \in \mathbb{R}^3.$$

The previous inequality allows to upper-bound the first addend in the same way as the second one thanks to the Fubini theorem, which concludes the proof. \square

APPENDIX B. CONSERVATION LAWS

In this section, we provide the proof of Proposition 4. We take the projected Boltzmann equation (23) satisfied by \mathbf{f}^0 and scalarly multiply it by each $\chi^i(\mathbf{nM})^{1/2}$, $1 \leq i \leq I + 4$.

We first choose $1 \leq i \leq I$ to obtain the conservation law for ρ_i , that is

$$\partial_t \left\langle \mathbf{f}^0, (\mathbf{nM})^{1/2} \chi^i \right\rangle_I + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0), (\mathbf{nM})^{1/2} \chi^i \right\rangle_I + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), (\mathbf{nM})^{1/2} \chi^i \right\rangle_I = 0.$$

The first term obviously gives $\partial_t \rho_i$. The second term can of course be rewritten as

$$\left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0), (\mathbf{nM})^{1/2} \chi^i \right\rangle_I = \left\langle v_1 \partial_x \mathbf{f}^0, (\mathbf{nM})^{1/2} \chi^i \right\rangle_I,$$

which we can explicitly compute, helped by parity arguments,

$$\left\langle v_1 \partial_x \mathbf{f}^0, (\mathbf{nM})^{1/2} \chi^i \right\rangle_I = \partial_x \left[\int_{\mathbb{R}^3} v_1 f_i^0 (n_i M_i)^{1/2} dv \right] = \frac{n_i \partial_x q^1}{\sqrt{\sum n_j m_j}} \int_{\mathbb{R}^3} m_i v_1^2 M_i dv = \frac{n_i \partial_x q^1}{\sqrt{\sum n_j m_j}}.$$

The third term being unchanged, we recover (26).

To obtain the conservation law on q^1 , we can write

$$\partial_t q^1 + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0), (\mathbf{nM})^{1/2} \chi^{I+1} \right\rangle_I + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), (\mathbf{nM})^{1/2} \chi^{I+1} \right\rangle_I = 0.$$

The second term also simplifies thanks to parity arguments

$$\begin{aligned} \left\langle v_1 \partial_x \mathbf{f}^0, (\mathbf{nM})^{1/2} \chi^{I+1} \right\rangle_I &= \sum_i \partial_x \left[\int_{\mathbb{R}^3} v_1 f_i^0 (n_i M_i)^{1/2} \chi_i^{I+1} dv \right] \\ &= \sum_i \frac{m_i}{\sqrt{\sum n_j m_j}} \partial_x \int_{\mathbb{R}^3} \left(\rho_i v_1^2 \sqrt{n_i} M_i + \frac{e}{\sqrt{6 \sum n_j}} (m_i |v|^2 - 3) v_1^2 n_i M_i \right) dv \\ &= \frac{1}{\sqrt{\sum n_j m_j}} \left(\sum_i \sqrt{n_i} \partial_x \rho_i + \frac{\sqrt{6 \sum n_j}}{3} \partial_x e \right). \end{aligned}$$

This finishes the derivation of the conservation law (27) for q^1 . The one for q^2 and q^3 (28), is then straightforward.

Finally, the conservation law (29) for e reads

$$\partial_t e + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^0), (\mathbf{nM})^{1/2} \chi^{I+4} \right\rangle_I + \left\langle \mathbf{P}^0(v_1 \partial_x \mathbf{f}^1), (\mathbf{nM})^{1/2} \chi^{I+4} \right\rangle_I = 0.$$

Let us compute the second term of the above equation, which gives

$$\begin{aligned} \left\langle v_1 \partial_x \mathbf{f}^0, (\mathbf{nM})^{1/2} \chi^{I+4} \right\rangle_I &= \sum_{i=1}^I \int_{\mathbb{R}^3} v_1 \partial_x f_i^0 (n_i M_i)^{1/2} \chi_i^{I+4} dv \\ &= \frac{1}{\sqrt{6 \sum n_j}} \sum_i n_i \partial_x \left[\int_{\mathbb{R}^3} (m_i |v|^2 - 3) \left(\frac{q^1}{\sqrt{\sum n_j m_j}} m_i v_1 \right) v_1 M_i dv \right] \\ &= \sum_i \frac{2n_i}{\sqrt{6 \sum n_j}} \frac{\partial_x q^1}{\sqrt{\sum n_j m_j}} = \frac{1}{3} \sqrt{\frac{6 \sum n_j}{\sum n_j m_j}} \partial_x q^1. \end{aligned}$$

Summarizing, the conservation law (29) for e is obtained.

APPENDIX C. PROOF OF LEMMA 7 IN THREE DIMENSIONS

In what follows, the one-dimensional notations are straightforwardly extended in three dimensions. The only significant difference with respect to the one-dimensional case is the treatment of $\nabla_x q$, which requires the use of the Korn inequality to conclude. Let us rewrite the main equalities and estimates in the three-dimensional setting. We first note that the conservation laws on the fluid quantities in \mathbf{f}^0 still hold, *i.e.*

$$\begin{aligned} \partial_t \rho_i + \frac{n_i}{\sqrt{\sum n_j m_j}} \nabla_x \cdot q + \left\langle \mathbf{P}^0(v \cdot \nabla_x \mathbf{f}^1), \chi^i (\mathbf{nM})^{1/2} \right\rangle_I &= 0, \quad 1 \leq i \leq I, \\ \partial_t q^k + \nabla_x \ell + \left\langle \mathbf{P}^0(v \cdot \nabla_x \mathbf{f}^1), \chi^{I+1} (\mathbf{nM})^{1/2} \right\rangle_I &= 0, \quad k = 1, 2, 3, \\ \partial_t e + \frac{1}{3} \sqrt{\frac{6 \sum n_j}{\sum n_j m_j}} \nabla_x \cdot q + \left\langle \mathbf{P}^0(v \cdot \nabla_x \mathbf{f}^1), \chi^{I+4} (\mathbf{nM})^{1/2} \right\rangle_I &= 0, \\ \partial_t \ell + \frac{\sum n_j^{3/2} + \frac{2}{3} \sum n_j}{\sum n_j m_j} \nabla_x \cdot q & \\ + \frac{1}{\sqrt{\sum n_j m_j}} \left\langle \mathbf{P}^0(v \cdot \nabla_x \mathbf{f}^1), \left(\sum_i \sqrt{n_i} \chi^i + \frac{1}{3} \sqrt{6 \sum n_j} \chi^{I+4} \right) (\mathbf{nM})^{1/2} \right\rangle_I &= 0, \end{aligned}$$

where ℓ is still defined by (30). Of course, we follow the strategy of Section 4.6. We first compute

$$\begin{aligned} \|v \cdot \nabla_x \mathbf{f}^0\|_I^2 &= \sum_i \frac{1}{m_i} \left| \nabla_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \nabla_x e \right|^2 + \frac{5}{3} \frac{1}{\sum n_j} \sum_i \frac{n_i}{m_i} |\nabla_x e|^2 \\ &+ \frac{\sum n_j}{\sum n_j m_j} \left[(\nabla_x \cdot q)^2 + \sum_{l,m} \left((\partial_{x_l} q^m)^2 + \partial_{x_l} q^m \partial_{x_m} q^l \right) \right]. \end{aligned}$$

In the same way, as in (50), we can write

$$\|\mathbf{P}^0(v \cdot \nabla_x \mathbf{f}^0)\|_I^2 = |\nabla_x \ell|^2 + \frac{5}{3} \frac{\sum n_j}{\sum n_j m_j} (\nabla_x \cdot q)^2.$$

Then, for some $\theta \in (0, 1)$, (52) becomes

$$\begin{aligned} \|\mathbf{P}^1(v \cdot \nabla_x \mathbf{f}^0)\|_I^2 &\geq \theta \sum_i \frac{1}{m_i} \left| \nabla_x \rho_i + \frac{2\sqrt{n_i}}{\sqrt{6 \sum n_j}} \nabla_x e \right|^2 + \frac{5}{3} \frac{1}{\sum n_j} \left(\sum \frac{n_j}{m_j} \right) |\nabla_x e|^2 \\ &+ \frac{\sum n_j}{\sum n_j m_j} \left[-\frac{2}{3} (\nabla_x \cdot q)^2 + \sum_{l,m} \left((\partial_{x_l} q^m)^2 + \partial_{x_l} q^m \partial_{x_m} q^l \right) \right] - \theta |\nabla_x \ell|^2. \end{aligned}$$

Besides, using the three-dimensional laws on q^k and ℓ , (54) can be rewritten into

$$\begin{aligned} \theta \int_0^T \int_{\mathbb{R}^3} |\nabla_x \ell|^2 dx dt + 2\theta \int_{\mathbb{R}^3} q \cdot \nabla_x \ell dx \Big|_0^T \\ \leq 2\theta \left(\frac{\sum n_j^{3/2} + \frac{2}{3} \sum n_j}{\sum n_j m_j} + \frac{1}{2} \right) \int_0^T \int_{\mathbb{R}^3} (\nabla_x \cdot q)^2 dx dt + \theta C_{\mathcal{X}} \int_0^T \int_{\mathbb{R}^3} \|\nabla_x \mathbf{f}^1\|_I^2 dx dt. \end{aligned}$$

Thanks to the Korn inequality, for θ small enough, there exists $C_\theta > 0$, only depending on θ , such that

$$-\left(\frac{2}{3} + \theta\right) \int_{\mathbb{R}^3} (\nabla_x \cdot q)^2 dx + \sum_{l,m} \int_{\mathbb{R}^3} \left((\partial_{x_l} q^m)^2 + \partial_{x_l} q^m \partial_{x_m} q^l \right) dx \geq C_\theta \int_{\mathbb{R}^3} |\nabla_x q|^2 dx.$$

Eventually, in the same way as in Subsection 4.6, the norm equivalence argument on \mathbb{P}^0 (34) allows to obtain

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^3} \|\mathbf{P}^1(v \cdot \nabla_x \mathbf{f}^0)\|_I^2 dx dt &\geq \gamma_\theta \int_0^T \int_{\mathbb{R}^3} \|\nabla_x \mathbf{f}^0\|_I^2 dx dt \\ &\quad - \theta C_2 \int_0^T \int_{\mathbb{R}^3} \|\nabla_x \mathbf{f}^1\|_I^2 dx dt + 2\theta \int_{\mathbb{R}^3} q \cdot \nabla_x \ell dx \Big|_0^T, \end{aligned}$$

for some constants $\gamma_\theta > 0$ depending on \mathbf{n} , \mathbf{m} and θ , and $C_2 > 0$ depending on \mathbf{n} and \mathbf{m} .

REFERENCES

- [1] C. Baranger, M. Bisi, S. Brull, and L. Desvillettes. On the Chapman-Enskog asymptotics for a mixture of monoatomic and polyatomic rarefied gases. *Kinet. Relat. Models*, 11(4):821–858, 2018.
- [2] M. Bennoune, M. Lemou, and L. Mieussens. Uniformly stable numerical schemes for the Boltzmann equation preserving the compressible Navier-Stokes asymptotics. *J. Comput. Phys.*, 227(8):3781–3803, 2008.
- [3] A. Bondesan, L. Boudin, M. Briant, and B. Grec. Stability of the spectral gap for the Boltzmann multi-species operator linearized around non-equilibrium Maxwell distributions. *Commun. Pure Appl. Anal.*, 19(5):2549–2573, 2020.
- [4] A. Bondesan and M. Briant. Stability of the Maxwell-Stefan system in the diffusion asymptotics of the Boltzmann multi-species equation. *Comm. Math. Phys.*, 382(1):381–440, 2021.
- [5] L. Boudin, B. Grec, and V. Pavan. The Maxwell-Stefan diffusion limit for a kinetic model of mixtures with general cross sections. *Nonlinear Anal.*, 159:40–61, 2017.
- [6] L. Boudin, B. Grec, M. Pavić, and F. Salvarani. Diffusion asymptotics of a kinetic model for gaseous mixtures. *Kinet. Relat. Models*, 6(1):137–157, 2013.
- [7] L. Boudin, B. Grec, and F. Salvarani. The Maxwell-Stefan diffusion limit for a kinetic model of mixtures. *Acta Appl. Math.*, 136:79–90, 2015.
- [8] M. Briant. Stability of global equilibrium for the multi-species Boltzmann equation in L^∞ settings. *Discrete Contin. Dyn. Syst.*, 36(12):6669–6688, 2016.
- [9] M. Briant and E. S. Daus. The Boltzmann equation for a multi-species mixture close to global equilibrium. *Arch. Ration. Mech. Anal.*, 222(3):1367–1443, 2016.
- [10] C. Cercignani. *The Boltzmann equation and its applications*, volume 67 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1988.
- [11] A. Crestetto, C. Klingenberg, and M. Pirner. Kinetic/fluid micro-macro numerical scheme for a two component gas mixture. *Multiscale Model. Simul.*, 18(2):970–998, 2020.
- [12] E. S. Daus, A. Jüngel, C. Mouhot, and N. Zamponi. Hypocoercivity for a linearized multispecies Boltzmann system. *SIAM J. Math. Anal.*, 48(1):538–568, 2016.
- [13] E. de la Canal, I. M. Gamba, and M. Pavić-Čolić. Propagation of L^p_β -norm, $1 < p \leq \infty$, for the system of Boltzmann equations for monoatomic gas mixtures. ArXiv e-print 2001.09204, 2020.
- [14] L. Desvillettes, R. Monaco, and F. Salvarani. A kinetic model allowing to obtain the energy law of polytropic gases in the presence of chemical reactions. *Eur. J. Mech. B Fluids*, 24(2):219–236, 2005.
- [15] G. Dimarco and L. Pareschi. Multi-scale control variate methods for uncertainty quantification in kinetic equations. *J. Comput. Phys.*, 388:63–89, 2019.
- [16] J. Dolbeault, C. Mouhot, and C. Schmeiser. Hypocoercivity for linear kinetic equations conserving mass. *Trans. Amer. Math. Soc.*, 367(6):3807–3828, 2015.
- [17] I. M. Gamba, S. Jin, and L. Liu. Micro-macro decomposition based asymptotic-preserving numerical schemes and numerical moments conservation for collisional nonlinear kinetic equations. *J. Comput. Phys.*, 382:264–290, 2019.
- [18] I. M. Gamba and M. Pavić-Čolić. On existence and uniqueness to homogeneous Boltzmann flows of monoatomic gas mixtures. *Arch. Ration. Mech. Anal.*, 235(1):723–781, 2020.
- [19] F. Golse, B. Perthame, and C. Sulem. On a boundary layer problem for the nonlinear Boltzmann equation. *Arch. Rational Mech. Anal.*, 103(1):81–96, 1988.
- [20] Y. Guo. The Vlasov-Maxwell-Boltzmann system near Maxwellians. *Invent. Math.*, 153(3):593–630, 2003.

- [21] Y. Guo. The Boltzmann equation in the whole space. *Indiana Univ. Math. J.*, 53(4):1081–1094, 2004.
- [22] H. Hutridurga and F. Salvarani. Maxwell–Stefan diffusion asymptotics for gas mixtures in non-isothermal setting. *Nonlinear Analysis*, 159:285–297, 2017.
- [23] S. Jin and Q. Li. A BGK-penalization-based asymptotic-preserving scheme for the multispecies Boltzmann equation. *Numer. Methods Partial Differential Equations*, 29(3):1056–1080, 2013.
- [24] S. Jin and Y. Shi. A micro-macro decomposition-based asymptotic-preserving scheme for the multispecies Boltzmann equation. *SIAM J. Sci. Comput.*, 31(6):4580–4606, 2009/10.
- [25] M.-Y. Lee, T.-P. Liu, and S.-H. Yu. Large-time behavior of solutions for the Boltzmann equation with hard potentials. *Comm. Math. Phys.*, 269(1):17–37, 2007.
- [26] S. Liu and H. Liu. Optimal time decay of the Boltzmann system for gas mixtures. *Nonlinear Anal.*, 95:592–606, 2014.
- [27] T.-P. Liu, T. Yang, and S.-H. Yu. Energy method for Boltzmann equation. *Phys. D*, 188(3-4):178–192, 2004.
- [28] T.-P. Liu, T. Yang, S.-H. Yu, and H.-J. Zhao. Nonlinear stability of rarefaction waves for the Boltzmann equation. *Arch. Ration. Mech. Anal.*, 181(2):333–371, 2006.
- [29] T.-P. Liu and S.-H. Yu. Boltzmann equation: micro-macro decompositions and positivity of shock profiles. *Comm. Math. Phys.*, 246(1):133–179, 2004.
- [30] T.-P. Liu and S.-H. Yu. Initial-boundary value problem for one-dimensional wave solutions of the Boltzmann equation. *Comm. Pure Appl. Math.*, 60(3):295–356, 2007.
- [31] T.-P. Liu and S.-H. Yu. Solving Boltzmann equation, Part I: Green’s function. *Bull. Inst. Math. Acad. Sin. (N.S.)*, 6(2):115–243, 2011.
- [32] C. Mouhot and L. Neumann. Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus. *Nonlinearity*, 19(4):969–998, 2006.
- [33] S. Simić, M. Pavić-Čolić, and D. Madjarević. Non-equilibrium mixtures of gases: modelling and computation. *Riv. Math. Univ. Parma (N.S.)*, 6(1):135–214, 2015.
- [34] T. Yang and H.-J. Zhao. A new energy method for the Boltzmann equation. *J. Math. Phys.*, 47(5):053301, 19, 2006.

LB: SORBONNE UNIVERSITÉ, CNRS, UNIVERSITÉ DE PARIS, LABORATOIRE JACQUES-LOUIS LIONS (LJLL), F-75005 PARIS, FRANCE

Email address: laurent.boudin@sorbonne-universite.fr

BG: MAP5, CNRS UMR 8145, UNIVERSITÉ DE PARIS, F-75006 PARIS, FRANCE

Email address: berenice.grec@u-paris.fr

MPC: DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA OBRADOVIĆA 4, 21000 NOVI SAD, SERBIA

Email address: milana.pavic@dmi.uns.ac.rs

SS: DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA OBRADOVIĆA 4, 21000 NOVI SAD, SERBIA

Email address: srboljub.simic@dmi.uns.ac.rs