



HAL
open science

Friction mediated by transient elastic linkages : extension to loads of bounded variation

Samar Allouch, Vuk Milišić

► **To cite this version:**

Samar Allouch, Vuk Milišić. Friction mediated by transient elastic linkages : extension to loads of bounded variation. *Journal of Integral Equations and Applications*, 2022. hal-03378290

HAL Id: hal-03378290

<https://hal.science/hal-03378290>

Submitted on 14 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Friction mediated by transient elastic linkages : extension to loads of bounded variation

Samar Allouch^{*†} Vuk Milišić^{*‡}

October 13, 2021

Abstract

In this work, we are interested in the convergence of a system of integro-differential equations with respect to an asymptotic parameter ε . It appears in the context of cell adhesion modelling [16, 15]. We extend the framework from [12, 13], strongly depending on the hypothesis that the external load f is in $\text{Lip}([0, T])$ to the case where $f \in \text{BV}(0, T)$ only. We show how results presented in [13] naturally extend to this new setting, while only partial results can be obtained following the comparison principle introduced in [12].

1 Introduction

Cell motility plays a central role in several important phenomena in biology : cancer cell migration, neutrophils' extravasation, chemotaxis, etc. The present paper fits in the modelling framework presented in [17, 19, 10]. The adhesive dynamics of actin filaments are at the heart of the project : they contribute to lamellipodium's stabilization and allow the cell to attach to the substrate or the surrounding tissue. This paper contributes to the better mathematical understanding of a minimal model introduced first in [12], its aim is to extend results already obtained in [12, 13] to the case of *stiffer* external loads.

More precisely, we are interested in the motion of a single binding site, linked to a one-dimensional substrate and subjected to an external force f . As in [12, 13], the position of this binding site, denoted z_ε , solves a Volterra integral equation

$$\begin{cases} \frac{1}{\varepsilon} \int_0^\infty (z_\varepsilon(t) - z_\varepsilon(t - \varepsilon a)) \rho_\varepsilon(a, t) da = f(t), & t \geq 0, \\ z_\varepsilon(t) = z_p(t), & t < 0. \end{cases} \quad (1)$$

The kernel ρ_ε above solves a non-local age-structured problem :

$$\begin{cases} \varepsilon \partial_t \rho_\varepsilon + \partial_a \rho_\varepsilon + \zeta_\varepsilon \rho_\varepsilon = 0, & t > 0, a > 0, \\ \rho_\varepsilon(a = 0, t) = \beta_\varepsilon(t) \left(1 - \int_0^\infty \rho_\varepsilon(t, \tilde{a}) d\tilde{a} \right), & t > 0, \\ \rho_\varepsilon(a, t = 0) = \rho_{I, \varepsilon}(a), & a \geq 0, \end{cases} \quad (2)$$

where $\beta_\varepsilon \in \mathbb{R}_+$ (resp. $\zeta_\varepsilon \in \mathbb{R}_+$) is the kinetic on-rate (resp. off-rate) function. These possibly depend on the dimensionless parameter $\varepsilon > 0$. The past positions are stored in the Lipschitz function $z_p(t) \in \mathbb{R}$, prescribed for every $t < 0$.

⁰Acknowledgments : the authors thank the referee for valuable remarks that helped to improve the clarity of this work.

^{*}Laboratoire Analyse, Géométrie & Applications (LAGA), Université Paris 13

[†]e-mail : allouch@math.univ-paris13.fr

[‡]e-mail : milisic@math.univ-paris13.fr

Various mathematical issues related to this system have already been investigated [12, 13, 11]. In [12], the authors have introduced a specific Lyapunov functional in order to study the convergence of (2) when ε goes to 0. Indeed, due to the saturation effect in the non-local boundary condition in (2), neither the Generalized Relative Entropy [18, 5] nor more generic comparison principles [6] do apply. Then, concerning (1), under the assumptions that the force f is Lipschitz on \mathbb{R} , and because the kernel ρ_ε in (1) is non-negative, an extension of Gronwall's Lemma to integral equations, shows convergence of z_ε towards z_0 the solution of (3), the limit equation associated to (1). These two steps show that

$$\|z_\varepsilon - z_0\|_{C^0([0,T])} + \|\rho_\varepsilon - \rho_0\|_{C^0([0,T];L^1(\mathbb{R}_+))} \rightarrow 0.$$

where z_0 is given by

$$\begin{cases} \mu_{1,0}(t)\partial_t z_0(t) = f(t), & t > 0 \\ z_0(0) = z_p(0) & t = 0 \end{cases} \quad (3)$$

where $\mu_{1,0}(t) := \int_0^\infty a\rho_0(a,t)da$, and ρ_0 solves :

$$\begin{cases} \partial_a \rho_0 + \zeta_0(a,t)\rho_0 = 0, & t > 0, a > 0, \\ \rho_0(a=0,t) = \beta_0(t) \left(1 - \int_0^\infty \rho_0(t,\tilde{a}) d\tilde{a}\right), & t > 0. \end{cases} \quad (4)$$

In [13], the authors weakened some assumptions concerning the off-rate ζ_ε , by assuming that ζ_ε is not necessarily non-decreasing passed a certain age a_0 . Then, they introduce a new variable u_ε related to z_ε which transforms (1) into a transport problem with a non-local source term :

$$\begin{cases} \varepsilon\partial_t u_\varepsilon + \partial_a u_\varepsilon = \frac{1}{\mu_{0,\varepsilon}(t)} \left(\varepsilon\partial_t f + \int_0^\infty \zeta_\varepsilon(\tilde{a},t)u_\varepsilon(\tilde{a},t)\rho_\varepsilon(\tilde{a},t) d\tilde{a} \right), & t > 0, a > 0, \\ u_\varepsilon(a=0,t) = 0, & t > 0, \\ u_\varepsilon(a,t=0) = u_{I,\varepsilon}(a) := \frac{z_\varepsilon(0) - z_p(-\varepsilon a)}{\varepsilon}, & a \geq 0, \end{cases} \quad (5)$$

where $\mu_{0,\varepsilon}(t) := \int_0^\infty \rho_\varepsilon(\tilde{a},t) d\tilde{a}$ and according to (1), it holds that

$$z_\varepsilon(0) = \frac{1}{\mu_{0,\varepsilon}(0)} \left(\int_0^\infty z_p(-\varepsilon a)\rho_{I,\varepsilon}(a) da + \varepsilon f(0) \right). \quad (6)$$

If $f \in \text{Lip}(\mathbb{R})$, systems (1) and (5) are equivalent. Nevertheless, (5) admits a stability result that allows to show a weak-* convergence of $u_\varepsilon/(1+a)$ towards $u_0/(1+a)$ in $L^\infty(\mathbb{R}_+ \times (0,T))$, where u_0 is the solution of the limit problem

$$\begin{cases} \partial_a u_0 = \frac{1}{\mu_{0,0}} \int_0^\infty \zeta_0 u_0 \rho_0 d\tilde{a}, & t > 0, a > 0, \\ u_0(a=0,t) = 0, & t > 0, \end{cases} \quad (7)$$

which in turn provides the strong convergence of z_ε in $C([0,T])$ towards z_0 solving (3).

In our analysis, however, when $f \in \text{BV}((0,T))$, the derivative of f is neither a function nor it is bounded, since it is a Radon measure. Therefore, we cannot apply directly results from [12]. Instead, defining f_δ to be a specific regularization of f [20, Section 5.3] provides a *regular* function u_ε^δ solving (5). To do this, we use the framework already established in [13]. Then we show that u_ε^δ satisfies certain *a priori* estimates that are uniform with respect to both δ and ε . These provide necessary compactness in order to pass to the limit with the regularization parameter δ and give existence and uniqueness of a weak solution u_ε associated to (5) with a load $f \in \text{BV}((0,T))$. The *a priori* estimates holding also in this weaker framework, we can consider convergence with respect to ε and prove consistency with the formal limit system. We show that, in the BV framework, the equivalence between (5) and (1) still holds. For the particular case when the kernel ρ_ε is independent

on time and on ε and under suitable hypotheses, we show error estimates to be compared with [12], the comparison principle being applied to the integral of the error's modulus.

In order to clarify the interplay between parameters ε and δ , we make the following remarks : in the previous literature [12, 13], not only existence (and uniqueness) but also convergence results were *strongly* related to the Lipschitz regularity of the load f . This motivates the present work since it is not clear that the convergence occurs with respect to ε in this weaker framework. This explains also why we first regularize the problem with the parameter δ , make δ tend to zero and then consider the convergence with respect to ε .

The outline of the paper is as follows : in Section 2, collecting various results from the literature on BV-functions in one space dimension, we introduce the framework used in the rest of the paper. We make the link with the Riemann-Stieltjes integral, through a careful analysis of different definitions of BV-functions with respect to the boundary of the time domain $(0, T)$. In Section 3, we recall some results concerning (2) already established in [12]. Then in Section 4, we establish uniform (with respect to ε) *a priori* estimates for the regularized system in u_ε . After that, in Section 5, we show the weak convergence of u_ε towards u_0 , the solution of the limit problem. This implies strong convergence of z_ε in $L^\infty(0, T)$ as stated in Theorem 8. We establish, in Section ??, a specific comparison principle for Volterra equations when the density ϱ is constant in time and does not depend on ε .

2 Notations and main assumptions

We denote $L_t^p L_a^q := L^p((0, T); L^q(\mathbb{R}_+))$ for any real $(p, q) \in [1, \infty]^2$ and

$$X_T := \left\{ g \in L_{loc}^\infty((0, T) \times \mathbb{R}_+) ; \sup_{t \in (0, T)} \|g(t, a)w(a)\|_{L_a^\infty} < \infty \right\} \quad (8)$$

where $w(a) := (1 + a)^{-1}$. The space $\text{Lip}(I)$ is the set of Lipschitz functions on the interval I .

Assumptions 1. For any $T > 0$ possibly infinite, we assume that :

i) The past condition z_p is L_{z_p} -Lipschitz on \mathbb{R}_- *i.e.* :

$$|z_p(a_2) - z_p(a_1)| \leq L_{z_p} |a_2 - a_1|, \quad \forall (a_2, a_1) \in \mathbb{R}_- \times \mathbb{R}_-.$$

ii) The function $\beta_\varepsilon(t)$ is in $L^\infty(0, T)$ and $\zeta_\varepsilon(a, t)$ is in $L^\infty(\mathbb{R}_+ \times (0, T))$.

iii) For limit functions $\beta_0 \in L_t^\infty$ and $\zeta_0 \in L_t^\infty L_a^\infty$ it holds that

$$\|\zeta_\varepsilon - \zeta_0\|_{L_{a,t}^\infty} \rightarrow 0 \quad \text{and} \quad \|\beta_\varepsilon - \beta_0\|_{L_t^\infty} \rightarrow 0$$

as $\varepsilon \rightarrow 0$.

iv) There are upper and lower bounds such that

$$0 < \zeta_{\min} \leq \zeta_\varepsilon(a, t) \leq \zeta_{\max} \quad \text{and} \quad \beta_{\min} \leq \beta_\varepsilon(a, t) \leq \beta_{\max},$$

for all $\varepsilon > 0$, $a \geq 0$ and $t > 0$.

Assumptions 2. The initial condition $\rho_{I,\varepsilon} \in L_a^\infty(\mathbb{R}_+)$ satisfies

i) positivity

$$\rho_{I,\varepsilon}(a) \geq 0, \quad \text{a.e. in } \mathbb{R}_+,$$

moreover, one has also that the total initial population satisfies

$$0 < \int_{\mathbb{R}_+} \rho_{I,\varepsilon}(a) da < 1;$$

ii) boundedness of higher moments,

$$0 < \int_{\mathbb{R}_+} a^p \rho_{I,\varepsilon}(a) da < c_p, \quad \text{for } p = 1, 2,$$

where c_p are positive constants depending only on p ;

Next, we introduce definitions of functions with bounded variation in one dimension, as well as some related properties.

Definition 1. Let $f : (0, T) \rightarrow \mathbb{R}$ be a Lebesgue measurable function. The pointwise variation (or Jordan variation) of f on $(0, T)$ is

$$\text{pvar}(f, (0, T)) := \sup_P \text{var}(f, P) \quad (9)$$

where $\text{var}(f, P) := \sum_{k=1}^n |f(t_k) - f(t_{k-1})|$ and $P = \{0 < t_0 < \dots < t_n < T\}$ is a partition of $(0, T)$.

Moreover, we denote $\text{BPV}((0, T)) := \{f \in \mathcal{L}((0, T)), \text{ s.t. } \text{pvar}(f, (0, T)) < +\infty\}$, the space of measurable functions with pointwise bounded variation, see for example, [1, section 3.2], [9, chapter 2] and [7, section 2.2, 2.3]. The pointwise variation of f is clearly dependent on the value of f at each point of the domain, and it differs from one a.e.-representative of f to another. For this reason, for every measurable function f , one defines the essential pointwise variation :

$$\text{epvar}(f, (0, T)) := \inf \{\text{pvar}(g, (0, T)) : f(t) = g(t) \text{ a.e. } t \in (0, T)\} \quad (10)$$

In [9, Chapter 6], another functional space is defined :

Definition 2. Given an open interval $(0, T) \subset \mathbb{R}$, the space of functions with bounded variation $\text{BV}((0, T))$ is defined as the space of all functions $f \in L^1((0, T))$ for which there exists a signed Radon measure μ_f such that

$$\int_{(0, T)} f \phi' dt = - \int_{(0, T)} \phi d\mu_f, \text{ for every } \phi \in C_c^1((0, T)) \quad (11)$$

for all $\phi \in C_c^1((0, T))$. The measure μ_f is called the weak or distributional derivative of f .

Remark 1.

i) We define the total variation of $f \in L^1((0, T))$ by

$$\|Df\|((0, T)) = \sup \left\{ - \int_{(0, T)} f \phi' dt, \phi \in C_c^1((0, T)), |\phi|_\infty \leq 1 \right\}. \quad (12)$$

Moreover, $f \in \text{BV}((0, T))$ if $\|Df\|((0, T)) < +\infty$.

ii) Definitions (9) and (12) are not equivalent. For instance, the Dirichlet indicatrix function $\chi_{\mathbb{Q} \cap [0, 1]}$ is not of pointwise bounded variation in $(0, 1)$ in the sense of Definition 1 but is well defined in the sense of Definition 2. The equivalence between the two definitions holds up to a.e. equality. Moreover every integrable function $f : (0, T) \rightarrow \mathbb{R}$ such that $\text{pvar}(f, (0, T)) < +\infty$, is in $\text{BV}((0, T))$ and $\|Df\|((0, T)) \leq \text{pvar}(f, (0, T))$. On the other hand, if f belongs to $\text{BV}((0, T))$, then f admits a right continuous representative \bar{f} with bounded pointwise variation such that

$$\text{pvar}(\bar{f}, (0, T)) = \|Df\|((0, T)).$$

For more details, see, e.g., [9, theorem 7.3] and [7].

iii) Under the norm

$$\|f\|_{\text{BV}} := \|f\|_{L^1} + \text{epvar}(f, (0, T)) < \infty$$

$\text{BV}((0, T))$ is a Banach space.

Next, we provide existence of the left and right limits of functions with bounded variation [7, Proposition 2.2].

Lemma 1. Let $f \in \text{BV}((0, T))$, Then both the limits

$$f(0^+) = \lim_{s \rightarrow 0, s > 0} f(s) \quad \text{and} \quad f(T^-) = \lim_{s \rightarrow T, s < T} f(s) \quad \text{exist.}$$

Additionally, if f is integrable, the left and right limits are as follows:

Lemma 2. Suppose that $f \in \text{BV}((0, T))$, then

$$f(0^+) = \lim_{\rho \rightarrow 0^+} \frac{1}{\rho} \int_0^\rho f(t) dt, \quad f(T^-) = \lim_{\rho \rightarrow 0^+} \frac{1}{\rho} \int_{T-\rho}^T f(t) dt.$$

Next, we present a result used in the proof of Proposition 3, which relates the pointwise variation to the Lebesgue measure:

$$\lambda(f, h, \Omega) := \int_{\{t \in \Omega : t+h \in \Omega\}} |f(t+h) - f(t)| dt,$$

Lemma 3. If f is in $\text{BPV}((0, T))$, then $\lambda(f, h, (0, T))/|h|$ is bounded. Moreover,

$$\lambda(f, h, (0, T)) \leq |h| \text{pvar}(f, (0, T)).$$

For the proof we can see [9, Theorem 2.20].

Finally, in [7], the authors add a new notion of variation containing the boundary value in order to expand the total variation of f to $[0, T]$. This variation is defined as

$$\text{varw}(f) := \sup_{\substack{\phi \in C_c^1((0, T)) \\ |\phi|_\infty \leq 1}} \left\{ \phi(T)f(T^-) - \phi(0)f(0^+) - \int_{(0, T)} f \phi' dt \right\} \quad (13)$$

Moreover, by summarizing the results of [7, Proposition 2.3, 2.6 and 2.7] all notions of variations coincide:

$$\text{epvar}(f, (0, T)) = \|Df\|((0, T)) = \text{varw}(f)$$

The previous result allows to extend Lemma 3 to $\text{BV}((0, T))$ functions :

Lemma 4. If f is in $\text{BV}((0, T))$, then $\lambda(f, h, (0, T))/|h|$ is bounded. Moreover,

$$\lambda(f, h, (0, T)) \leq |h| \|Df\|((0, T))$$

PROOF. By taking the infimum over almost every equal measurable functions, one has

$$\inf_{f=\tilde{f} \text{ a.e.}} \lambda(\tilde{f}, h, (0, T)) \leq |h| \inf_{f=\tilde{f} \text{ a.e.}} \text{pvar}(\tilde{f}, (0, T)) = |h| \text{epvar}(f, (0, T)) = |h| \|Df\|((0, T))$$

Since the left hand side is a Lebesgue integral one has :

$$\inf_{f=\tilde{f} \text{ a.e.}} \lambda(\tilde{f}, h, (0, T)) = \lambda(f, h, (0, T))$$

which ends the proof. □

2.1 Data regularization

Theorem 1. For every $f \in \text{BV}((0, T))$, there exists a sequence of smooth functions $(f_\delta)_\delta$ in $C^\infty((0, T))$ such that

$$\lim_{\delta \rightarrow 0} \int_{(0, T)} |f_\delta - f| dt = 0 \quad \text{and} \quad \lim_{\delta \rightarrow 0} \int_{(0, T)} |f'_\delta| dt = \|Df\|((0, T)).$$

Although the proof is classical (see for instance [20, Theorem 5.3.3 p.225]), we need the explicit form of f_δ in the rest of the paper. For this reason, we present in Section A.1 the proof of Theorem 1.

Lemma 5. Let $f \in \text{BV}((0, T)) \cap L^\infty((0, T))$. Then the regularization function f_δ defined as (60) is bounded in $(0, T)$.

Next, we compare the left and right limits of f and its' approximation f_δ on the boundary:

Lemma 6. Let $f \in \text{BV}((0, T))$ and f_δ defined as (60), then

$$f_\delta(0^+) = f(0^+) \quad \text{and} \quad f_\delta(T^-) = f(T^-).$$

First we need the following result :

Proposition 1. Let $f \in \text{BV}((0, T))$. For every $\delta > 0$, and $t_0 \in \{0, T\}$,

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau} \int_{I_\tau \cap (0, T)} |f_\delta - f| dt = 0, \quad (14)$$

where $I_\tau = \{t \in \mathbb{R} : |t - t_0| < \tau\}$.

PROOF. For a fixed $t_0 \in \{0, T\}$ and $t \in I_\tau \in (0, T)$, we have

$$f_\delta(t) - f(t) = \sum_{i=0}^{\infty} [\chi_{\delta_i} * (\phi_i f) - \phi_i f]$$

by the definition of $\text{supp } \phi_i$ (see (56)), we have $1/(j_0 + i + 1) < \tau < 1/(j_0 + i - 1)$ then $i > 1/\tau - j_0 - 1$. Since, \mathbb{R} is an archimedean space then

$$\forall \tau > 0, \exists! i_0 := \left\lfloor \frac{1}{\tau} \right\rfloor - j_0 \text{ s.t } i_0 \leq \frac{1}{\tau} < i_0 + 1 \quad (15)$$

which implies by using (58) that

$$\begin{aligned} \int_{I_\tau \cap (0, T)} |f_\delta - f| dt &= \sum_{i=i_0}^{\infty} \int_{I_\tau \cap (0, T)} [\chi_{\delta_i} * (\phi_i f) - \phi_i f] dt \leq \sum_{i=i_0}^{\infty} \delta 2^{-i} \\ &\leq \delta 2^{-i_0} \sum_{i=0}^{\infty} 2^{-i} = 2^{j_0+1} \delta 2^{-\lfloor \frac{1}{\tau} \rfloor} \quad \square \end{aligned}$$

then

$$\frac{1}{\tau} \int_{I_\tau \cap (0, T)} |f_\delta - f| dt \leq C \delta \frac{2^{-\lfloor \frac{1}{\tau} \rfloor}}{\tau}$$

using again (15), we have

$$\frac{2^{-\lfloor \frac{1}{\tau} \rfloor}}{\tau} = \frac{\exp(-\lfloor \frac{1}{\tau} \rfloor \ln 2)}{\tau} \leq \frac{2 \exp(-\frac{1}{\tau} \ln 2)}{\tau}$$

Finally, we conclude that

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau} \int_{I_\tau \cap (0, T)} |f_\delta - f| dt = \lim_{\tau \rightarrow 0^+} \frac{2 \exp(-\frac{1}{\tau} \ln 2)}{\tau} = 0.$$

PROOF OF LEMMA 6. According to the Lemma 2,

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau} \int_0^\tau |f_\delta(t) - f_\delta(0^+)| dt = 0 \quad \text{and} \quad \lim_{\tau \rightarrow 0^+} \frac{1}{\tau} \int_0^\tau |f(t) - f(0^+)| dt = 0.$$

Moreover, we have, thanks to Proposition 1

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau} \int_0^\tau |f_\delta - f| dt = 0.$$

Thus, for all $\varepsilon' > 0$, there exist $\delta' > 0$ such that $0 < \tau < \delta'$ implies

$$\begin{aligned} |f_\delta(0^+) - f(0^+)| &= \frac{1}{\tau} \int_0^\tau |f_\delta(0^+) - f(0^+)| dt \\ &\leq \frac{1}{\tau} \int_0^\tau |f_\delta(0^+) - f_\delta(t)| dt + \frac{1}{\tau} \int_0^\tau |f_\delta(t) - f(t)| dt + \frac{1}{\tau} \int_0^\tau |f(t) - f(0^+)| dt \leq 3\varepsilon' \end{aligned}$$

which proves the required result. Similarly, we can prove that $f_\delta(T^-) = f(T^-)$. \square

In the previous setting, the weak derivative of $f \in \text{BV}((0, T))$ defines a linear continuous form on $C((0, T))$. In the next section, we show how to *extend* this measure on functions in $C([0, T])$.

2.2 Definition and basic properties of Stieltjes integral

The Riemann–Stieltjes integral (RS–integral) is a generalization of the Riemann integral. Let \mathcal{P} a tagged partition of $[0, T]$, defined as

$$\mathcal{P} := \{(\xi_i, [t_{i-1}, t_i]) : 1 \leq i \leq n\} \quad (16)$$

where $0 = t_1 \leq \dots \leq t_n = T$, and on each interval $[t_{i-1}, t_i]$ we choose a single value ξ_i , for $i \in \{1, \dots, n\}$.

Definition 3. For any function $f, g : [0, T] \rightarrow \mathbb{R}$ and a partition \mathcal{P} , we define the Riemann–Stieltjes sum by

$$S(f, dg, \mathcal{P}, [0, T]) := \sum_i f(\xi_i)[g(t_i) - g(t_{i-1})].$$

Moreover, the RS-integral of f with respect to g

$$(RS) \int_{[0, T]} f(t) dg(t)$$

exists and has a value $I \in \mathbb{R}$, if, for every $\varepsilon > 0$, there exists $\delta > 0$, such that the mesh size $\max_i(t_i - t_{i-1}) < \delta$ and for every ξ_i in $[t_i, t_{i+1}]$,

$$|S(f, dg, \mathcal{P}, [0, T]) - I| < \varepsilon.$$

Lemma 7. Suppose that f is continuous on $[0, T]$ and g is of bounded pointwise variation on $[0, T]$, then

$$\left| \int_{[0, T]} f dg \right| \leq \|f\|_\infty \text{pvar}(g, [0, T])$$

In the following Theorem we see that a Riemann–Stieltjes integral can be used to describe any bounded linear functional on $C([0, T])$ (see [3, Theorem 7.1.1] and [8, Theorem 4.4-1] for more details)

Theorem 2. Let $\Gamma_f \in (C([0, T]))'$, then there exist $g \in BPV([0, T])$ such that

$$\Gamma_f(\varphi) = \int_{[0, T]} \varphi dg, \quad \forall \varphi \in C([0, T]).$$

Theorem 3. (Integration by parts). If one of the integrals $\int_{[0, T]} f dg$ and $\int_{[0, T]} g df$ exists, then the other exists as well, and we have

$$\int_{[0, T]} f dg + \int_{[0, T]} g df = [fg(t)]_{t=0}^{t=T}.$$

Moreover, If $f \in C^1([0, T])$ and $g \in BPV([0, T])$, then $df = f' dt$ in the second term of the left hand side.

For the proof cf [14, Theorem 5.52] and [2, Lemma 2].

Lemma 8. Let $f \in \text{BV}((0, T))$. Then there exists $g \in \text{BPV}([0, T])$ s.t

$$[f\varphi]_{t=0^+}^{t=T^-} - \int_{(0, T)} f\varphi' dx = \int_{[0, T]} \varphi dg, \quad \forall \varphi \in C^1([0, T])$$

s.t. $f(t) = g(t)$, a.e. $t \in (0, T)$

PROOF. We regularize $f \in \text{BV}((0, T))$ by $f_\delta \in C^\infty((0, T))$ as in Theorem 1, then we have :

$$\int_{t_k}^{s_k} f'_\delta \varphi dt + \int_{t_k}^{s_k} f_\delta \varphi' dt = [f_\delta \varphi]_{t=t_k}^{t=s_k} =: L_k, \quad \forall \varphi \in C^1([0, T])$$

where $s_k \rightarrow T^-$ and $t_k \rightarrow 0^+$. We define :

$$I_k := \int_{t_k}^{s_k} f'_\delta \varphi dt, \quad J_k := \int_{t_k}^{s_k} f_\delta \varphi' dt.$$

Thanks to Lebesgue's Theorem, one has that

$$\lim_{k \rightarrow \infty} I_k = I := \int_0^T f'_\delta \varphi dt, \quad \lim_{k \rightarrow \infty} J_k = J := \int_0^T f_\delta \varphi' dt$$

and thanks to Lemma 6 and the continuity of φ ,

$$L_k = L := f(T^-)\varphi(T) - f(0^+)\varphi(0), \quad \forall k \in \mathbb{N}$$

So that we have :

$$\int_0^T f'_\delta \varphi dt + \int_0^T f_\delta \varphi' dt = [f\varphi]_{t=0^+}^{t=T^-}$$

If we set

$$\mathcal{I}_{f_\delta}(\varphi) := \int_0^T f'_\delta \varphi dt,$$

it is a linear continuous form on $C([0, T])$, since one has :

$$|\mathcal{I}_{f_\delta}(\varphi)| \leq \|f'_\delta\|_{L^1((0, T))} \|\varphi\|_{L^\infty} \leq \{\|Df\|((0, T)) + \delta\} \|\varphi\|_{L^\infty((0, T))} \quad (17)$$

where we used estimates from the proof of [20, Theorem 5.3.3]. Since it is a continuous linear form on $C([0, T])$, by Theorem 2 there exists $h_\delta \in \text{BPV}([0, T])$ s.t.

$$\mathcal{I}_{f_\delta}(\varphi) = \int_{[0, T]} \varphi dh_\delta, \quad \forall \varphi \in C([0, T])$$

in the Stieljes' sense. But by using the integration by parts from Theorem 3, we have that

$$[f\varphi]_{t=0^+}^{t=T^-} - \int_{(0, T)} f_\delta \varphi' dt = \int_{[0, T]} \varphi dh_\delta = [h_\delta \varphi]_{t=0}^{t=T} - \int_{[0, T]} h_\delta \varphi' dt, \quad \forall \varphi \in C^1([0, T]) \quad (18)$$

which implies that

$$\int_{(0, T)} f_\delta \varphi' dt = \int_{(0, T)} h_\delta \varphi' dt, \quad \forall \varphi \in \mathcal{D}((0, T))$$

and then we can apply [9, Lemma 7.4] and conclude that there exists $c \in \mathbb{R}$, s.t.

$$f_\delta = h_\delta + c.$$

Then setting $g_\delta := h_\delta + c$ provides a function s.t.

$$[f_\delta \varphi]_0^T - \int_0^T f_\delta \varphi' dt = \int_0^T \varphi dg_\delta, \quad \forall \varphi \in C^1([0, T])$$

and s.t.

$$f_\delta(t) = g_\delta(t), \quad \text{a.e. } t \in (0, T).$$

Thanks to (17), \mathcal{I}_{f_δ} is a linear continuous form on $C([0, T])$ uniformly bounded with respect to δ . It can be identified via the Riesz representation theorem as a Radon measure μ_δ on $[0, T]$. Therefore, there exist $\mu \in M^1([0, T])$ and a sub-sequence μ_{δ_k} such that

$$\mu_{\delta_k} \xrightarrow{*} \mu$$

in $\sigma(M^1([0, T]), C([0, T]))$ with respect to the weak-* topology. By Theorem 2, there exists $h \in \text{BPV}([0, T])$ s.t.

$$\mu(\varphi) = \int_0^T \varphi dh$$

where the left side is a Radon measure and the right hand side is the Riemann-Stieltjes integral. Because f_δ tends to f in the $L^1(0, T)$ topology, one has then that

$$[f\varphi]_{t=0^+}^{t=T^-} - \int_{(0, T)} f\varphi' dt = \int_{[0, T]} \varphi dh, \quad \forall \varphi \in C^1([0, T])$$

then using again integration by parts from Theorem 3, one concludes that

$$f(t) = h(t) + \tilde{c}, \quad \text{a.e. } t \in (0, T)$$

and setting $g = h + \tilde{c}$ ends the proof. \square

Corollary 1. There exists a sub-sequence $(f_{\delta_k})_{k \in \mathbb{N}}$, s.t.

$$I_{f_{\delta_k}}(\varphi) := \int_0^T \varphi f_{\delta_k}' dt \rightarrow \int_0^T \varphi dg, \quad \forall \varphi \in C([0, T])$$

when $k \rightarrow \infty$

3 Mathematical background for the linkages' density

We list here some of the results proved in [12] used in the next sections of the paper.

Theorem 4. Let Assumptions 1 and 2 hold, then for every fixed $\varepsilon > 0$ there is a unique solution $\rho_\varepsilon \in C^0(\mathbb{R}_+; L^1(\mathbb{R}_+)) \cap L^\infty(\mathbb{R}_+^2)$ of the problem (2). It satisfies (2) in the sense of characteristics, namely

$$\rho_\varepsilon(a, t) = \begin{cases} \beta_\varepsilon(t - \varepsilon a) \left(1 - \int_{\mathbb{R}_+} \rho_\varepsilon(\tilde{a}, t - \varepsilon a) d\tilde{a}\right) \exp\left(-\int_0^a \zeta_\varepsilon(\tilde{a}, t - \varepsilon(a - \tilde{a})) d\tilde{a}\right), & \forall a < \frac{t}{\varepsilon} \\ \rho_{I, \varepsilon}\left(a - \frac{t}{\varepsilon}\right) \exp\left(-\frac{1}{\varepsilon} \int_0^t \zeta_\varepsilon\left(\frac{\tilde{t}-t}{\varepsilon} + a, \tilde{t}\right) d\tilde{t}\right), & \forall a \geq \frac{t}{\varepsilon} \end{cases} \quad (19)$$

Moreover, it is a weak solution as well since it satisfies (cf [12, Lemma 2.1])

$$\begin{aligned} & \int_0^T \int_0^{+\infty} \rho_\varepsilon(a, t) (\varepsilon \partial_t \varphi + \partial_a \varphi - \zeta_\varepsilon \varphi) da dt - \varepsilon \int_0^{+\infty} \rho_\varepsilon(a, T) \varphi(a, T) da \\ & + \int_0^T \rho_\varepsilon(a=0, t) \varphi(a=0, t) dt + \varepsilon \int_0^{+\infty} \rho_{I, \varepsilon}(a) \varphi(a, t=0) da = 0 \end{aligned} \quad (20)$$

for every $T > 0$ and test function $\varphi \in C^\infty(\mathbb{R}_+^2) \cap L^\infty(\mathbb{R}_+^2)$. Now we define the moments of ρ_ε which we denote by $\mu_{p, \varepsilon}(t) := \int_{\mathbb{R}_+} a^p \rho_\varepsilon(a, t) da$, with $p = 1, 2$.

Lemma 9. Let Assumptions 1 and 2 hold, then the unique solution $\rho_\varepsilon \in C^0(\mathbb{R}_+; L^1(\mathbb{R}_+)) \cap L^\infty(\mathbb{R}_+^2)$ of (2) satisfies

- i) $\rho_\varepsilon(a, t) \geq 0$ for a.e (a, t) in \mathbb{R}_+^2 ,
- ii) $\mu_{0, \min} \leq \mu_{0, \varepsilon}(t) < 1, \forall t \in \mathbb{R}_+$ where $\mu_{0, \min} < \min\left(\mu_{0, \varepsilon}(0), \frac{\beta_{\min}}{\beta_{\min} + \zeta_{\max}}\right)$
- iii) $\mu_{p, \min} \leq \mu_{p, \varepsilon}(t) \leq k$, where $\mu_{p, \min} = \min\left(\mu_{p, \varepsilon}(0), \frac{\mu_{p-1, \min}}{\zeta_{\max}}\right)$

The authors provide a Liapunov functional that reads :

$$\mathcal{H}[u] := \left| \int_0^\infty u(a) da \right| + \int_0^\infty |u(a)| da ,$$

thanks to which they obtain the following convergence result for ρ_ε :

Lemma 10. Let $\zeta_{\min} > 0$ be the lower bound to $\zeta_\varepsilon(a, t)$ according to Assumptions 1, and setting $\hat{\rho}_\varepsilon := \text{one}$ has

$$\mathcal{H}[(\rho_\varepsilon - \rho_0)(\cdot, t)] \leq \mathcal{H}[\rho_{I, \varepsilon} - \rho_0(\cdot, 0)] \exp\left(\frac{-\zeta_{\min} t}{\varepsilon}\right) + \frac{2}{\zeta_{\min}} \left(\|R_\varepsilon\|_{L_a^1(\mathbb{R}_+)} + \|M_\varepsilon\|_{L_t^\infty(\mathbb{R}_+)} \right) \quad (21)$$

with $R_\varepsilon := -\varepsilon \partial_t \rho_0 - \rho_0(\zeta_\varepsilon - \zeta_0)$ and $M_\varepsilon := (\beta_\varepsilon - \beta_0)(1 - \int_0^\infty \rho_0 da)$.

This ensures the convergence of ρ_ε that reads :

Theorem 5. Let ρ_ε the solution of the system (2) and let ρ_0 given by (4), then

$$\rho_\varepsilon \rightarrow \rho_0 \text{ in } C^0((0, \infty); L^1(\mathbb{R}_+)) \text{ as } \varepsilon \rightarrow 0 ,$$

where the convergence with respect to time is meant in the sense of uniform convergence on compact subintervals.

4 Existence, uniqueness and stability

Using the regularized function f_δ introduced in Theorem 1, we consider an approximation of (1) : we denote by $z_\varepsilon^\delta := z_\varepsilon^\delta(t)$ the function solving

$$\begin{cases} \frac{1}{\varepsilon} \int_0^\infty (z_\varepsilon^\delta(t) - z_\varepsilon^\delta(t - \varepsilon a)) \rho_\varepsilon(a, t) da = f_\delta(t), & t \leq 0 \\ z_\varepsilon^\delta(t) = z_p(t), & t < 0 \end{cases} \quad (22)$$

We also define $u_\varepsilon^\delta(a, t)$, an approximation of the elongation variable u_ε , defined as the mild solution of

$$\begin{cases} \varepsilon \partial_t u_\varepsilon^\delta + \partial_a u_\varepsilon^\delta = \frac{1}{\mu_{0,\varepsilon}} \left(\varepsilon f'_\delta + \int_0^\infty \zeta_\varepsilon u_\varepsilon^\delta \rho_\varepsilon da \right), & t > 0, a > 0 \\ u_\varepsilon^\delta(a = 0, t) = 0, & t > 0 \\ u_\varepsilon^\delta(a, t = 0) = u_{I,\varepsilon}^\delta(a), & a \geq 0 \end{cases} \quad (23)$$

where

$$u_{I,\varepsilon}^\delta(a) := \frac{z_\varepsilon^\delta(0^+) - z_p(-\varepsilon a)}{\varepsilon} \quad (24)$$

and

$$z_\varepsilon^\delta(0^+) = \frac{1}{\mu_{0,\varepsilon}(0)} \left(\varepsilon f_\delta(0^+) + \int_0^\infty z_p(-\varepsilon a) \rho_{I,\varepsilon}(a) da \right). \quad (25)$$

More precisely, u_ε^δ is a solution of system (23) in the sense of characteristics, namely

$$u_\varepsilon^\delta(a, t) = \begin{cases} \int_0^a h(t - \varepsilon \tilde{a}) d\tilde{a}, & \text{if } t > \varepsilon a \\ \int_0^{t/\varepsilon} h(t - \varepsilon \tilde{a}) d\tilde{a} + u_{I,\varepsilon}^\delta(a - t/\varepsilon), & \text{if } t \leq \varepsilon a, \end{cases} \quad (26)$$

where

$$h(t) := \frac{1}{\mu_{0,\varepsilon}} \left(\varepsilon \partial_t f_\delta + \int_0^\infty \zeta_\varepsilon u_\varepsilon^\delta \rho_\varepsilon da \right).$$

By arguments similar to [12, Lemma 3], it is as well a weak solution of (23) *i.e.*

$$\begin{aligned} & - \int_0^T \int_0^\infty u_\varepsilon^\delta (\varepsilon \partial_t \varphi + \partial_a \varphi) da dt + \left[\int_0^\infty u_\varepsilon^\delta(s, a) \varphi(s, a) da \right]_{s=0}^{s=T} \\ & = \int_0^T \frac{1}{\mu_{0,\varepsilon}} \left(\varepsilon f'_\delta + \int_0^\infty \zeta_\varepsilon u_\varepsilon^\delta \rho_\varepsilon da \right) \left(\int_0^\infty \varphi(t, \tilde{a}) d\tilde{a} \right) dt \end{aligned} \quad (27)$$

for any function $\varphi \in C_c^\infty([0, T] \times \mathbb{R}_+)$. Although, problem (22) can be defined for weaker data (typically $L^1((0, T))$ or the space of Radon measures $M((0, T))$), the elongation problem (23), requires to give a meaning to the time derivative of f , which is more restrictive. Nevertheless, as we are mainly interested in convergence results, $f \in \text{BV}((0, T))$ seems the weakest possible regularity to our knowledge.

Theorem 6. Let Assumptions 1 hold, and let ρ_ε be the unique solution of (2) then the system (23) has a unique solution $u_\varepsilon^\delta \in X_T$.

We are in the framework of [13, Theorem 6.1], but for sake of self-containness, we recall in an abridged version the proof hereafter.

PROOF. A Banach fixed point Theorem is used to prove this result. We define the mapping $\phi(v) = u$ such that by Duhamel's principle

$$u(a, t) = \begin{cases} \int_0^a G(t - \varepsilon \tilde{a}) d\tilde{a}, & \text{if } t > \varepsilon a \\ \int_0^{t/\varepsilon} G(t - \varepsilon \tilde{a}) d\tilde{a} + u_{I,\varepsilon}^\delta(a - t/\varepsilon), & \text{if } t \leq \varepsilon a. \end{cases} \quad (28)$$

where

$$G(t) := \frac{1}{\mu_{0,\varepsilon}} \left(\varepsilon f'_\delta + \int_0^\infty \zeta_\varepsilon(a, t) v(a, t) \rho_\varepsilon(a, t) da \right)$$

As in [13], a simple computation shows that

$$\|u\|_{X_T} \leq \|G\|_{L^\infty((0, T))} \frac{T}{T + \varepsilon} + \left\| \frac{u_{I,\varepsilon}^\delta(\cdot)}{1 + a} \right\|_{L^\infty(\mathbb{R}_+)}$$

Moreover, since $\partial_t f_\delta \in L^\infty((0, T))$

$$\|G\|_{L^\infty((0, T))} \leq \frac{1}{\mu_{0, \min}} \varepsilon \|f'_\delta\|_{L^\infty((0, T))} + \frac{\zeta_{\max}(1+k)}{\mu_{0, \min}} \|v\|_{X_T}$$

where k is the upper bound of $\mu_{1, \varepsilon}$ proved in Lemma 9. Furthermore, by the same argument we can prove that ϕ is a contraction. Indeed, if $u_i = \phi(v_i)$ for $i \in \{1, 2\}$

$$\|u_2 - u_1\|_{X_T} \leq C \frac{T}{T + \varepsilon} \|v_2 - v_1\|_{X_T}$$

for a constant $C > 0$. Then we can choose $T < \varepsilon/C$ and we obtain the existence of a local solution in time of (23), by Banach-Picard's fixed point theorem. As the contraction time does not depend on the initial data, we shall extend the same result by continuation. This shows existence and uniqueness in X_T for any $T > 0$. \square

Lemma 11. If Assumptions 1 holds, then the solution of system (23) satisfies the uniform a priori estimates

$$\begin{aligned} \int_0^\infty \rho_\varepsilon(a, t) |u_\varepsilon^\delta(a, t)| da &\leq \int_0^t |f'_\delta| d\tilde{t} + \int_{\mathbb{R}_+} \rho_{I, \varepsilon} |u_{I, \varepsilon}^\delta| da \\ &\leq C \left(\|f\|_{\text{BV}((0, T))}, \|(1+a)\rho_I\|_{L^1(\mathbb{R}_+)}, \|z'_p\|_{L^\infty(\mathbb{R}_-)} \right) \end{aligned} \quad (29)$$

where C is independent on ε and on δ .

PROOF. Again, we proceed as in [13, Lemma 5.1], multiplying (23) by $\text{sgn}(u_\varepsilon^\delta)$, testing against ρ_ε , and integrating with respect to a gives :

$$\frac{d}{dt} \int_{\mathbb{R}_+} |u_\varepsilon^\delta| \rho_\varepsilon da + \int_{\mathbb{R}_+} |u_\varepsilon^\delta| \zeta_\varepsilon \rho_\varepsilon da \leq \varepsilon |f'_\delta| + \int_{\mathbb{R}_+} |u_\varepsilon^\delta| \zeta_\varepsilon \rho_\varepsilon da$$

the rigorous proof relies on arguments exposed in [12, Lemma 3.1] and is left to the reader. Finally, after integration with respect to time, we conclude that

$$\int_0^\infty \rho_\varepsilon(a, t) |u_\varepsilon^\delta(a, t)| da \leq \int_0^t |f'_\delta| dt + \int_{\mathbb{R}_+} \rho_{I, \varepsilon} |u_{I, \varepsilon}^\delta| da$$

since

$$\begin{aligned} |u_{I, \varepsilon}^\delta(a)| &\leq \left| \frac{z_\varepsilon^\delta(0^+) - z_p(0)}{\varepsilon} \right| + \left| \frac{z_p(0) - z_p(-\varepsilon a)}{\varepsilon} \right| \\ &\leq \frac{1}{\mu_{0, \varepsilon}(0)} \left| f_\delta(0^+) + \frac{1}{\varepsilon} \int_0^\infty (z_p(-\varepsilon a) - z_p(0)) \rho_{I, \varepsilon}(a) da \right| + \left| \frac{z_p(0) - z_p(-\varepsilon a)}{\varepsilon} \right| \\ &\leq \frac{1}{\mu_{0, \min}} \left(\|z'_p\|_{L^\infty(\mathbb{R}_-)} \mu_{1, \varepsilon}(0) + f_\delta(0^+) \right) + \|z'_p\|_{L^\infty(\mathbb{R}_-)} a \\ &\leq \max \left\{ \frac{1}{\mu_{0, \min}} \|z'_p\|_{L^\infty(\mathbb{R}_-)} \mu_{1, \varepsilon}(0) + f_\delta(0^+), \|z'_p\|_{L^\infty(\mathbb{R}_-)} \right\} (1+a), \end{aligned}$$

the result follows. \square

In order to establish the convergence of u_ε^δ in X_T , for a fixed ε , we introduce an intermediate variable w defined as

$$w(a, t) := u_\varepsilon^\delta(a, t) - \frac{f_\delta(t)}{\mu_{0, \varepsilon}(t)}. \quad (30)$$

It satisfies

$$\begin{cases} \varepsilon \partial_t w + \partial_a w = \frac{1}{\mu_{0,\varepsilon}} \left(\varepsilon \frac{f_\delta \partial_t \mu_{0,\varepsilon}}{\mu_{0,\varepsilon}} + \int_0^\infty \zeta_\varepsilon u_\varepsilon^\delta \rho_\varepsilon da \right), & t > 0, a > 0, \\ w(a=0, t) = \frac{-f_\delta(t)}{\mu_{0,\varepsilon}(t)}, & t > 0, \\ w(a, t=0^+) = u_{I,\varepsilon}^\delta(a) - \frac{f_\delta(0^+)}{\mu_{0,\varepsilon}(0)}, & a \geq 0, \end{cases} \quad (31)$$

The following crucial result holds:

Lemma 12. For a fixed δ and ε , and under the Assumptions 1, the unknowns w and u_ε^δ , are uniformly bounded in X_T with respect to δ and ε .

PROOF. Using arguments from [12, Lemma 2.1], one can show that w defined as

$$w(a, t) := \begin{cases} w(0, t - \varepsilon a) + \int_0^a G_w(t - \varepsilon \tilde{a}) d\tilde{a}, & \text{if } t > \varepsilon a \\ w(a - t/\varepsilon, 0^+) + \int_0^{t/\varepsilon} G_w(t - \varepsilon \tilde{a}) d\tilde{a}, & \text{if } t \leq \varepsilon a. \end{cases} \quad (32)$$

is a weak solution of (31). In the latter definition $G_w(t) := \left\{ \varepsilon \frac{f_\delta \partial_t \mu_{0,\varepsilon}}{\mu_{0,\varepsilon}^2} + \frac{1}{\mu_{0,\varepsilon}} \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon^\delta da \right\}$. A simple computation shows that

$$\|w\|_{X_T} \leq \|G_w\|_{L^\infty((0,T))} + \|w(0, \cdot)\|_{L^\infty((0,T))} + \left\| \frac{w(\cdot, 0)}{1+a} \right\|_{L^\infty(\mathbb{R}_+)}.$$

It remains to estimate $\|G_w\|_{L^\infty((0,T))}$. For every fixed ε , $\mu_{0,\varepsilon}$ is a Lipschitz continuous function. Indeed, $\mu_{0,\varepsilon}$ satisfies

$$\varepsilon \partial_t \mu_{0,\varepsilon} - \beta_\varepsilon (1 - \mu_{0,\varepsilon}) + \int_{\mathbb{R}_+} \zeta_\varepsilon \rho_\varepsilon da = 0$$

and then

$$\|\varepsilon \partial_t \mu_{0,\varepsilon}\|_{L_t^\infty} \leq \|\beta_\varepsilon\|_{L_t^\infty} + \|\zeta_\varepsilon\|_{L_{t,a}^\infty} \|\rho_\varepsilon\|_{L_t^\infty L_a^1},$$

which shows, by using the result of Lemma 11 that G_w , and also w , are uniformly bounded in $L^\infty((0,T))$ with respect to δ and ε . Indeed,

$$\|G_w\|_{L^\infty((0,T))} \leq \frac{\|\varepsilon \partial_t \mu_{0,\varepsilon}\|_{L^\infty((0,T))} \|f_\delta\|_{L^\infty((0,T))}}{\mu_{0,\min}^2} + \frac{\zeta_{\max}}{\mu_{0,\min}} \int_{\mathbb{R}_+} \rho_\varepsilon |u_\varepsilon^\delta| da < +\infty.$$

Finally, we have that

$$\|u_\varepsilon^\delta\|_{X_T} \leq \|w\|_{X_T} + \frac{\|f_\delta\|_{L^\infty((0,T))}}{\mu_{0,\min}} < +\infty$$

which ends the proof. \square

Previous stability estimates allow to show :

Theorem 7. Under Assumption 1, one has for any fixed $\varepsilon > 0$,

$$u_\varepsilon^\delta \rightharpoonup u_\varepsilon \text{ weakly-}^* \text{ in } X_T$$

as $\delta \rightarrow 0$, where u_ε solves the weak problem

$$\begin{aligned} & - \int_0^T \int_0^\infty u_\varepsilon (\varepsilon \partial_t + \partial_a) \varphi \, da \, dt + \varepsilon \left[\int_0^\infty u_\varepsilon(a, s) \varphi(a, s) \, da \right]_{s=0}^{s=T} \\ & = \varepsilon \int_0^T \frac{\int_0^\infty \varphi(\tilde{a}, t) \, d\tilde{a}}{\mu_{0,\varepsilon}} \, dg + \int_0^T \frac{\int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon \, da}{\mu_{0,\varepsilon}} \left(\int_0^\infty \varphi(\tilde{a}, t) \, d\tilde{a} \right) \end{aligned} \quad (33)$$

for any $\varphi \in C_c^1([0, T] \times \mathbb{R}_+)$.

PROOF. The uniform bound on u_ε^δ in X_T , proved in Lemma 12, implies that

$$\frac{u_\varepsilon^\delta}{1+a} \xrightarrow{*} \frac{u_\varepsilon}{1+a}$$

in $L^\infty((0, T) \times \mathbb{R}_+)$ in the weak- $*$ sense and the limit function u_ε belongs X_T . For every $\psi \in L^\infty((0, T) \times \mathbb{R}_+)$, we have $\zeta_\varepsilon(1+a)\rho_\varepsilon\psi \in L^1((0, T) \times \mathbb{R}_+)$ and then

$$\int_0^T \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon^\delta \psi \, da \, dt \rightarrow \int_0^T \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon \psi \, da \, dt.$$

By Corollary 1, the first term of right-hand in (27) tends to $\int_0^T \int_0^\infty \varphi(t, \tilde{a}) \, d\tilde{a} / \mu_{0,\varepsilon} \, dg$ as $\delta \rightarrow 0$, for any $\varphi \in C_c^1([0, T] \times \mathbb{R}_+)$. Regarding the second term of the right hand side in (27), one has that $\zeta_\varepsilon \rho_\varepsilon / \mu_{0,\varepsilon} \in L^1((0, T) \times \mathbb{R}_+)$ and this leads, thanks again to the weak- $*$ convergence above to write :

$$\int_0^t \frac{1}{\mu_{0,\varepsilon}} \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon^\delta \, da \, d\tilde{t} \rightarrow \int_0^t \frac{1}{\mu_{0,\varepsilon}} \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon \, da \, d\tilde{t} \text{ as } \delta \rightarrow 0, \quad (34)$$

which ends the proof. \square

The latter theorem allows to prove a convergence result when returning to the z_ε variable :

Proposition 2. Under the same assumptions as above, it holds that

$$z_\varepsilon^\delta \rightarrow z_\varepsilon \text{ strongly in } L^\infty((0, T)) \text{ as } \delta \rightarrow 0,$$

where z_ε satisfies

$$z_\varepsilon(t) = z_\varepsilon(0^+) + \int_0^t \frac{\varepsilon}{\mu_{0,\varepsilon}} \, dg + \int_0^t \frac{1}{\mu_{0,\varepsilon}} \int_0^\infty \zeta_\varepsilon u_\varepsilon \rho_\varepsilon \, da \, d\tilde{t} \quad (35)$$

which is also a solution of (1).

Before showing this result, we make some comments : if u_ε^δ is a solution of (23) then z_ε^δ defined as

$$z_\varepsilon^\delta(t) := z_\varepsilon^\delta(0^+) + \int_0^t \frac{1}{\mu_{0,\varepsilon}(\tilde{t})} \left(\varepsilon f'_\delta(\tilde{t}) + \int_0^\infty \zeta_\varepsilon u_\varepsilon^\delta \rho_\varepsilon \, da \right) \, d\tilde{t} \quad (36)$$

solves (22). Conversely, if z_ε^δ solves (22) then u_ε^δ , given by

$$u_\varepsilon^\delta(a, t) = \begin{cases} \frac{z_\varepsilon^\delta(t) - z_\varepsilon^\delta(t - \varepsilon a)}{\varepsilon}, & \text{if } t > \varepsilon a \\ \frac{z_\varepsilon^\delta(t) - z_p(t - \varepsilon a)}{\varepsilon}, & \text{if } t \leq \varepsilon a \end{cases} \quad (37)$$

is a solution of (23). For more details see [13, Lemma 4.1 and Lemma 4.2].

PROOF. First, by using (34) in the proof of Theorem 7, and Lemma 6, we have that $f_\delta(0^+) = f(0^+)$ and z_ε^δ given by (36) converge strongly in $L^\infty((0, T))$ to z_ε which verifies (35). Using [13, Lemma 4.2], if z_ε^δ is defined as (36) it solves (22). Multiplying (22) by a test function $\varphi \in L^1(0, T)$ gives :

$$\frac{1}{\varepsilon} \int_0^T \int_0^{+\infty} (z_\varepsilon^{\delta k}(t) - z_\varepsilon^{\delta k}(t - \varepsilon a)) \rho_\varepsilon(a, t) \varphi(t) da dt = \int_0^T f_{\delta k}(t) \varphi(t) dt \quad (38)$$

As $z_\varepsilon^{\delta k}$ converges strongly in $L^\infty(0, T)$ to z_ε , the difference $z_\varepsilon^{\delta k}(t) - z_\varepsilon^{\delta k}(t - \varepsilon a)$ converges almost everywhere for any fixed (a, t) in $\mathbb{R}_+ \times (0, T)$ towards $z_\varepsilon(t) - z_\varepsilon(t - \varepsilon a)$. Thanks to the L^∞ bounds on $u_\varepsilon^{\delta k}/(1+a)$, and the bounds in $L^1(\mathbb{R}_+ \times (0, T))$ on the first moment of ρ_ε , there exists an integrable majorizing function $g(a, t)$ on $\mathbb{R}_+ \times (0, T)$ s.t.

$$|z_\varepsilon^{\delta k}(t) - z_\varepsilon^{\delta k}(t - \varepsilon a)| |\varphi(t)| \rho_\varepsilon(a, t) \leq g(a, t)$$

uniformly for every k . Thus one can apply the Lebesgue's Theorem in the right hand side of (38). Since f_δ converges in $L^1(0, T)$ the convergence occurs in (38) for every $\varphi \in \mathcal{D}(0, T)$ and thus almost everywhere in $(0, T)$ and thus z_ε solves (1). \square

5 Weak convergence when ε goes to zero

Next, we prove the weak convergence of u_ε from which we deduce the strong convergence of z_ε .

Theorem 8. Under the same assumptions as above, one has

$$u_\varepsilon \rightharpoonup u_0 \text{ weakly-}^* \text{ in } X_T$$

as $\varepsilon \rightarrow 0$, where u_0 satisfies (7) and

$$\int_0^\infty u_0(a, t) \rho_0(a, t) da = f(t) \text{ a.e } t \in (0, T).$$

Furthermore, it also holds that

$$z_\varepsilon \rightarrow z_0 \text{ strongly in } L^\infty((0, T)) \text{ as } \varepsilon \rightarrow 0.$$

PROOF. The proof follows the same steps as in [13, Theorem 6.2]. First, by Lemma 12, u_ε^δ is uniformly bounded in X_T with respect to δ and ε , and therefore u_ε is uniformly bounded in X_T with respect to ε , then u_ε is weakly convergent to u_0 in X_T . On the other hand, Theorem 3.2 and Lemma 3.4 imply that

$$(1+a)\rho_\varepsilon \rightarrow (1+a)\rho_0$$

strongly in $L^1((0, T) \times \mathbb{R}_+)$. These arguments justify that for every $\psi \in L^\infty((0, T) \times \mathbb{R}_+)$ one has

$$\int_0^T \int_0^\infty \zeta_\varepsilon \rho_\varepsilon u_\varepsilon \psi da dt \rightarrow \int_0^T \int_0^\infty \zeta_0 \rho_0 u_0 \psi da dt.$$

Indeed, one has

$$\begin{aligned} & \int_0^T \int_0^\infty \{\zeta_\varepsilon \rho_\varepsilon u_\varepsilon - \zeta_0 \rho_0 u_0\} \psi da dt = \int_0^T \int_0^\infty (\zeta_\varepsilon - \zeta_0) \rho_\varepsilon u_\varepsilon \psi da dt \\ & + \int_0^T \int_0^\infty \zeta_0 (\rho_\varepsilon - \rho_0) u_\varepsilon \psi da dt + \int_0^T \int_0^\infty \zeta_0 \rho_0 (u_\varepsilon - u_0) \psi da dt \end{aligned}$$

As $\zeta_\varepsilon \rightarrow \zeta_0$ by Assumptions 1, and thanks to the weak convergence of u_ε , both terms on the right-hand side tend to zero as $\varepsilon \rightarrow 0$. Note that this implies the weak convergence of $\int_{\mathbb{R}_+} \zeta_\varepsilon \rho_\varepsilon u_\varepsilon da$ in $L^1((0, T))$, since we can choose $\psi \in L^\infty((0, T))$. Moreover, thanks to Lemma 7 we obtain that

$$\left| \int_{[0, T]} \frac{\int_0^{+\infty} \varphi da}{\mu_{0, \varepsilon}} dg \right| \leq \frac{C \|\varphi\|_{L_{t, a}^\infty}}{\mu_{0, \min}} \text{pvar}(g, [0, T]) \leq C \|f\|_{\text{BV}((0, T))}.$$

As in [13, Theorem 6.2], passing to the limit in the weak formulation (33) we obtain

$$- \int_0^T \int_0^\infty u_0 \partial_a \psi da dt = \int_0^T \int_0^\infty \frac{\zeta_0 \rho_0 u_0 \psi}{\mu_{0, 0}} da dt$$

which implies that u_0 satisfies

$$\begin{cases} \partial_a u_0 = \frac{1}{\mu_{0, 0}} \int_0^\infty \zeta_0(\tilde{a}, t) u_0(\tilde{a}, t) \rho_0(\tilde{a}, t) d\tilde{a}, & t > 0, a > 0, \\ u_0(a = 0, t) = 0, & t > 0, \end{cases} \quad (39)$$

Similarly, we have the weak convergence of $\int_0^\infty u_\varepsilon(t, a) \rho_\varepsilon(t, a) da$ towards $\int_0^\infty u_0(t, a) \rho_0(t, a) da$ in $L^1((0, T))$. Hence, one concludes that u_0 satisfies also

$$\int_0^\infty u_0(t, a) \rho_0(t, a) da = f(t), \quad \text{a.e. } t \in (0, T).$$

As the the right hand side of (39) does not depend on age, one has that $u_0 = \gamma(t) a$, where in order to satisfy the last compatibility condition implies that

$$\gamma(t) \int_{\mathbb{R}_+} a \rho_0(a, t) da = f(t), \quad \text{a.e. } t \in (0, T).$$

Thus $u_0(a, t) = f(t)/\mu_{1, 0}(t)a$ for almost every $t \in (0, T)$ and every $a \in \mathbb{R}_+$. Using again the weak convergence of $\int_{\mathbb{R}_+} \zeta_\varepsilon \rho_\varepsilon u_\varepsilon da$ in $L^1((0, T))$ combined with the strong convergence of $\mu_{0, \varepsilon}$ allows to pass to the limit in the third term of (35).

Moreover,

$$\begin{aligned} |z_\varepsilon(0^+) - z_p(0)| &= \frac{1}{\mu_{0, \varepsilon}(0)} \left| \int_0^\infty (z_p(-\varepsilon a) - z_p(0)) \rho_{I, \varepsilon}(a) da + \varepsilon f(0^+) \right| \\ &\leq \frac{\varepsilon k \|z_p'\|_{L^\infty(\mathbb{R}_-)}}{\mu_{0, \min}} + \frac{\varepsilon |f(0^+)|}{\mu_{0, \min}} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0 \end{aligned}$$

where k is the constant from Lemma 9.

All together this provides that z_0 solves :

$$z_0(t) = z_p(0) + \int_0^t \frac{f(\tau)}{\mu_{1, 0}(\tau)} \frac{\int_{\mathbb{R}_+} \zeta_0(a, \tau) \rho_0(a, \tau) da}{\mu_{0, 0}(\tau)} d\tau \quad (40)$$

but because ρ_0 solves (4), one has that $a\rho_0$ solves :

$$\partial_a(a\rho_0) - \rho_0 + a\zeta_0(a, t)\rho_0 = 0,$$

which after integration in time shows that

$$\frac{\int_{\mathbb{R}_+} a\zeta_0(a, t)\rho_0(a, t) da}{\mu_{0, 0}(t)} = 1$$

and this shows in turn that (40) reduces to

$$z_0(t) = z_0(0) + \int_0^t \frac{f(\tau)}{\mu_{1, 0}(\tau)} d\tau$$

which is the integrated version of (3). □

6 A comparison principle

In this section, we give error estimates between z_ε and z_0 , the solution of the problem

$$\begin{cases} \frac{1}{\varepsilon} \int_0^\infty (z_\varepsilon(t) - z_\varepsilon(t - \varepsilon a)) \varrho(a) da = f(t), & t \geq 0, \\ z_\varepsilon(t) = z_p(t), & t < 0, \end{cases} \quad (41)$$

where ϱ is constant in time and satisfies

$$\begin{cases} \partial_a \varrho + \zeta(a) \varrho = 0, & a > 0, \\ \varrho(0) = \beta \left(1 - \int_0^\infty \varrho(\tilde{a}) d\tilde{a} \right), \end{cases} \quad (42)$$

where the data of (41) and (42) satisfy

Assumptions 3.

- i) $f \in \text{BV}((0, T))$,
- ii) $z_p \in \text{Lip}(\mathbb{R}_-)$,
- iii) $\beta \in \mathbb{R}_+^*$,
- iv) $\zeta \in L^\infty(\mathbb{R}_+)$ such that

$$0 < \zeta_{\min} \leq \zeta(a) \leq \zeta_{\max}, \quad \text{a.e. } a \in \mathbb{R}_+.$$

Setting $\hat{z}_\varepsilon(t) := z_\varepsilon(t) - z_0(t)$, it solves :

$$\hat{z}_\varepsilon(t) = \frac{1}{\mu_0} \int_0^{\frac{t}{\varepsilon}} \hat{z}_\varepsilon(t - \varepsilon a) \varrho(a) da + \tilde{h}_\varepsilon(t)$$

where

$$\begin{aligned} \tilde{h}_\varepsilon(t) &= \frac{\varepsilon}{\mu_0} \int_0^{t/\varepsilon} \left(\frac{z_0(t) - z_0(t - \varepsilon a)}{\varepsilon} - a \partial_t z_0(t) \right) \varrho(a) da \\ &+ \frac{\varepsilon}{\mu_0} \int_{t/\varepsilon}^{+\infty} \left(\frac{z_0(t) - z_0(0)}{\varepsilon} - a \partial_t z_0(t) \right) \varrho(a) da + \frac{1}{\mu_0} \int_{t/\varepsilon}^{+\infty} (z_p(t - \varepsilon a) - z_p(0)) \varrho(a) da \end{aligned} \quad (43)$$

where $\mu_0 := \int_0^{+\infty} \varrho(a) da$ then

$$|\hat{z}_\varepsilon(t)| \leq \frac{1}{\mu_0} \int_0^{\frac{t}{\varepsilon}} |\hat{z}_\varepsilon(t - \varepsilon a)| \varrho(a) da + |\tilde{h}_\varepsilon(t)| \quad (44)$$

Then integrating in time and setting

$$\hat{Z}_\varepsilon(t) := \int_0^t |\hat{z}_\varepsilon(\tau)| d\tau$$

one has that :

$$\hat{Z}_\varepsilon(t) = \int_0^t |\hat{z}_\varepsilon(\tau)| d\tau \leq \frac{1}{\mu_0} \int_0^t \int_0^{\frac{\tau}{\varepsilon}} |\hat{z}_\varepsilon(\tau - \varepsilon a)| \varrho(a) da d\tau + \int_0^t |\tilde{h}_\varepsilon(\tau)| d\tau$$

then, we change the order of integration and the domain of integration becomes $D' := \{(a, \tau) \in (0, t/\varepsilon) \times (\varepsilon a, t)\}$. We use the change of variable $\tilde{t} = \tau - \varepsilon a$ in order to write :

$$\begin{aligned} \int_0^t \int_0^{\frac{\tau}{\varepsilon}} |\hat{z}_\varepsilon(\tau - \varepsilon a)| \varrho(a) da d\tau &= \int_0^{\frac{t}{\varepsilon}} \int_{\varepsilon a}^t |\hat{z}_\varepsilon(\tau - \varepsilon a)| d\tau \varrho(a) da \\ &= \int_0^{\frac{t}{\varepsilon}} \int_0^{t - \varepsilon a} |\hat{z}_\varepsilon(\tilde{t})| d\tilde{t} \varrho(a) da = \int_0^{\frac{t}{\varepsilon}} \hat{Z}_\varepsilon(t - \varepsilon a) \varrho(a) da \end{aligned}$$

So that finally \hat{Z}_ε solves :

$$\hat{Z}_\varepsilon(t) \leq \frac{1}{\mu_0} \int_0^{\frac{t}{\varepsilon}} \hat{Z}_\varepsilon(t - \varepsilon a) \varrho(a) da + \int_0^t \tilde{h}_\varepsilon(\tau) d\tau = \int_0^t \hat{Z}_\varepsilon(\tilde{a}) K_\varepsilon(t - \tilde{a}) d\tilde{a} + \int_0^t \tilde{h}_\varepsilon(\tau) d\tau$$

where $K_\varepsilon(\tilde{a}) := \frac{1}{\varepsilon\mu_0} \varrho\left(\frac{\tilde{a}}{\varepsilon}\right)$ is the kernel of the integral operator. We use a comparison principle [6, the Generalised Gronwall Lemma 8.2 p. 257] and construct a majorizing function U_ε of the form $U_\varepsilon(t) = \varepsilon(K_0 + K_1 t)$ where K_0 and K_1 are suitably chosen, such that $U_\varepsilon \geq |\hat{Z}_\varepsilon|$ and $U_\varepsilon \sim \varepsilon$. The following two lemmas are required in order to apply this comparison principle :

Lemma 13. The Volterra kernel K_ε satisfies :

$$\|K_\varepsilon\|_{L^\infty(0,T)} := \operatorname{ess\,sup}_{t \in (0,T)} \int_0^t |K_\varepsilon(\tilde{a})| d\tilde{a} < 1$$

PROOF. To prove this result, we need to show that

$$0 \leq \int_0^t |K_\varepsilon(\tilde{a})| d\tilde{a} = \frac{\int_0^{\frac{t}{\varepsilon}} \varrho(a) da}{\int_0^{+\infty} \varrho(a) da} < 1. \quad (45)$$

The kernel ϱ solves (42), thus it can be explicitly computed as

$$\varrho(a) = \frac{\beta}{1 + \beta I} \exp\left(-\int_0^a \zeta(s) ds\right)$$

one has the lower bound :

$$\varrho(a) \geq \frac{\beta}{1 + \beta I} \exp(-\zeta_{\max} a) > 0, \quad \forall a \in \mathbb{R}_+.$$

This in turn shows that

$$\int_{\frac{t}{\varepsilon}}^{\infty} \varrho(a) da > 0$$

which is equivalent to the claim. \square

Lemma 14. Consider the expectation value of a given density ϱ with respect to the tail $a > t/\varepsilon$,

$$A_1[\varrho](t) := \frac{\int_0^{+\infty} a \varrho(a + \frac{t}{\varepsilon}) da}{\int_0^{+\infty} \varrho(a + \frac{t}{\varepsilon}) da} \quad (46)$$

then under Assumptions 3, one has

$$A_1[\varrho](t) \leq \frac{\zeta_{\max}}{\zeta_{\min}^2}.$$

PROOF. Setting

$$q(a, t) := \frac{\varrho(a + \frac{t}{\varepsilon})}{\varrho(\frac{t}{\varepsilon})}, \quad (47)$$

it solves

$$\partial_a q(a, t) + \zeta(a + t/\varepsilon) q(a, t) = 0, \quad q(0, t) = 1.$$

This problem admits an explicit solution of the form

$$q(a, t) = \exp\left(-\int_0^a \zeta(\bar{a} + t/\epsilon) d\bar{a}\right) = \exp\left(-\int_{t/\epsilon}^{a+t/\epsilon} \zeta(\hat{a}) d\hat{a}\right). \quad (48)$$

Then, we shall rewrite (46) as :

$$A_1[\varrho](t) := \frac{\int_0^{+\infty} a q(a, t) da}{\int_0^{+\infty} q(a, t) da}$$

by using hypothesis iv) from Assumptions 3, one has

$$\exp(-\zeta_{\max} a) \leq q(a, t) \leq \exp(-\zeta_{\min} a)$$

This gives :

$$\int_0^{+\infty} q(a, t) da \geq \int_0^{+\infty} \exp(-\zeta_{\max} a) da = \frac{1}{\zeta_{\max}}$$

and

$$\int_0^{+\infty} a q(a, t) da \leq \int_0^{+\infty} a \exp(-\zeta_{\min} a) da = \frac{1}{\zeta_{\min}^2}$$

which shows the final upper bound. \square

Proposition 3. Under Assumptions 3, for $0 < t < T$ one has the estimates :

$$\tilde{H}_\epsilon(t) := \int_0^t |\tilde{h}_\epsilon(\tau)| d\tau \leq \epsilon^2 C_1$$

where C_1 depends on $\mu_2, \mu_1, \|\partial_t z_0\|_{\text{BV}((0, T))}$ and on $\|z_p\|_{\text{Lip}(\mathbb{R}_-)}$ but not on ϵ .

PROOF. Recalling the definition of \tilde{h}_ϵ in (43), we split $\int_0^t |\tilde{h}_\epsilon(\tau)| d\tau$ into three parts. First, we define

$$I_1 := \frac{\epsilon}{\mu_0} \int_0^t \int_0^{\tau/\epsilon} \left| \frac{z_0(\tau) - z_0(\tau - \epsilon a)}{\epsilon} - a \partial_t z_0(\tau) \right| \varrho(a) da d\tau$$

Since $\partial_t z_0 \in \text{BV}((0, T))$, then I_1 can be written in the form

$$I_1 = \frac{1}{\mu_0} \int_0^t \int_0^{\tau/\epsilon} \left| \int_{\tau - \epsilon a}^{\tau} (\partial_t z_0(\tilde{t}) - \partial_t z_0(\tau)) d\tilde{t} \right| \varrho(a) da d\tau$$

by switching the integration order between τ and a , and using the change of variable $\tilde{t} = \tau + h$, we get that

$$\begin{aligned} I_1 &\leq \frac{1}{\mu_0} \int_0^{t/\epsilon} \int_{\epsilon a}^t \int_{\tau - \epsilon a}^{\tau} |\partial_t z_0(\tilde{t}) - \partial_t z_0(\tau)| d\tilde{t} d\tau \varrho(a) da \\ &\leq \frac{1}{\mu_0} \int_0^{t/\epsilon} \int_{\epsilon a}^t \int_{-\epsilon a}^0 |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| dh d\tau \varrho(a) da \\ &\leq \frac{1}{\mu_0} \int_0^{t/\epsilon} \int_{-\epsilon a}^0 \int_{\epsilon a}^t |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| d\tau dh \varrho(a) da \end{aligned} \quad (49)$$

and thus applying Lemma 3, one has the estimate of the inner integral of the latter right hand side :

$$\int_{\epsilon a}^t |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| d\tau \leq |h| \|\partial_t z_0\|_{\text{BV}}, \quad \forall h \in (-\epsilon a, 0)$$

which implies that

$$I_1 \leq \frac{\|\partial_t z_0\|_{BV}}{\mu_0} \int_0^{t/\varepsilon} \int_{-\varepsilon a}^0 |h| dh \varrho(a) da \leq \frac{\varepsilon^2 \|\partial_t z_0\|_{BV}}{2\mu_0} \int_0^{t/\varepsilon} a^2 \varrho(a) da.$$

Next, we set:

$$\begin{aligned} I_2 &= \frac{\varepsilon}{\mu_0} \int_0^t \int_{\tau/\varepsilon}^{+\infty} \left| \frac{z_0(\tau) - z_0(0)}{\varepsilon} - a \partial_t z_0(\tau) \right| \varrho(a) da d\tau \\ &\leq \frac{1}{\mu_0} \int_0^t \int_{\tau/\varepsilon}^{+\infty} \left| \int_0^\tau \partial_t z_0(\tilde{t}) d\tilde{t} - \tau \partial_t z_0(\tau) \right| \varrho(a) da d\tau + \frac{1}{\mu_0} \int_0^t \int_{\tau/\varepsilon}^{+\infty} |\tau - \varepsilon a| |\partial_t z_0(\tau)| \varrho(a) da d\tau \\ &=: I_{2,1} + I_{2,2} \end{aligned}$$

As in the estimates of I_1 , first, one switches the order of integration and then one integrates on

$$D := \{(a, \tau) \in (0, t/\varepsilon) \times (0, \varepsilon a)\} \cup \{(a, \tau) \in (t/\varepsilon, +\infty) \times (0, t)\}, \quad (50)$$

and one makes the change of variable $\tilde{t} = \tau + h$ in order to obtain

$$\begin{aligned} I_{2,1} &= \frac{1}{\mu_0} \int_0^{t/\varepsilon} \int_0^{\varepsilon a} \int_{-\tau}^0 |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| dh d\tau \varrho(a) da \\ &\quad + \frac{1}{\mu_0} \int_{t/\varepsilon}^{+\infty} \int_0^t \int_{-\tau}^0 |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| dh d\tau \varrho(a) da \\ &= \frac{1}{\mu_0} \int_0^{t/\varepsilon} \int_{-\varepsilon a}^0 \int_{-h}^{\varepsilon a} |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| d\tau dh \varrho(a) da \\ &\quad + \frac{1}{\mu_0} \int_{t/\varepsilon}^{+\infty} \int_{-t}^0 \int_{-h}^t |\partial_t z_0(\tau + h) - \partial_t z_0(\tau)| d\tau dh \varrho(a) da \end{aligned}$$

also, by using Lemma 4, we get that

$$\begin{aligned} I_{2,1} &\leq \frac{\|\partial_t z_0\|_{BV}}{\mu_0} \int_0^{t/\varepsilon} \int_{-\varepsilon a}^0 |h| dh \varrho(a) da + \frac{\|\partial_t z_0\|_{BV}}{\mu_0} \int_{t/\varepsilon}^{+\infty} \int_{-t}^0 |h| dh \varrho(a) da \\ &\leq \frac{\varepsilon^2 \|\partial_t z_0\|_{BV}}{2\mu_0} \int_0^{t/\varepsilon} a^2 \varrho(a) da + \frac{\|\partial_t z_0\|_{BV}}{2\mu_0} \int_{t/\varepsilon}^{+\infty} t^2 \varrho(a) da \\ &\leq \frac{\varepsilon^2 \|\partial_t z_0\|_{BV}}{2\mu_0} \int_0^{t/\varepsilon} a^2 \varrho(a) da + \frac{\varepsilon^2 \|\partial_t z_0\|_{BV}}{2\mu_0} \int_{t/\varepsilon}^{+\infty} a^2 \varrho(a) da \\ &\leq \frac{\varepsilon^2 \mu_2 \|\partial_t z_0\|_{BV}}{\mu_0}. \end{aligned}$$

The second term $I_{2,2}$ is estimated in the same way as $I_{2,1}$. We have

$$\begin{aligned} I_{2,2} &= \frac{1}{\mu_0} \int_0^t \int_{\tau/\varepsilon}^{+\infty} |\tau - \varepsilon a| |\partial_t z_0(\tau)| \varrho(a) da d\tau \\ &= \frac{1}{\mu_0} \left\{ \int_0^{t/\varepsilon} \int_0^{\varepsilon a} + \int_{t/\varepsilon}^{+\infty} \int_0^t |\tau - \varepsilon a| |\partial_t z_0(\tau)| d\tau \varrho(a) da \right\} \\ &\leq \frac{\varepsilon^2 \|\partial_t z_0\|_{\infty}}{2\mu_0} \int_0^{t/\varepsilon} a^2 \varrho(a) da + \frac{\varepsilon^2 \|\partial_t z_0\|_{\infty}}{2\mu_0} \int_{t/\varepsilon}^{+\infty} a^2 \varrho(a) da \leq \frac{\varepsilon^2 \mu_2 \|\partial_t z_0\|_{\infty}}{\mu_0}. \quad \square \end{aligned}$$

Finally, by similar computations, one has

$$\frac{1}{\mu_0} \int_0^t \int_{\tau/\varepsilon}^{+\infty} |z_p(\tau - \varepsilon a) - z_p(0)| \varrho(a) da d\tau \leq \varepsilon^2 \|z_p\|_{\text{Lip}(\mathbb{R}_-)} \frac{\mu_2}{\mu_0}$$

Theorem 9. Under Assumptions 3, z_ε tends to z_0 , the solution of (3), strongly in $L^1(0, T)$ as ε goes to zero. Moreover, there exists a generic constant C depending only on the data of the problem but not on ε , such that :

$$\|z_\varepsilon - z_0\|_{L^1(0, T)} \leq \varepsilon C.$$

PROOF. We have proved in Lemma 13 that the Volterra kernel K_ε is non-positive and bounded (with a bound strictly less than one) in the sense of [6, Definition 2.2 p. 227 and Proposition 2.7 p.231] so [6, the Generalised Gronwall Lemma 8.2 p. 257] applies. First observe, by using Proposition 3, that

$$\hat{Z}_\varepsilon(t) - \int_0^t \hat{Z}_\varepsilon(\tilde{a}) K_\varepsilon(\tilde{a}) d\tilde{a} \leq \tilde{h}_{1,\varepsilon}(t) + \tilde{h}_{2,\varepsilon}(t)$$

We construct a function U_ε which satisfies,

$$U_\varepsilon(t) - \int_0^t U_\varepsilon(\tilde{a}) K_\varepsilon(\tilde{a}) d\tilde{a} \geq \tilde{h}_{1,\varepsilon} + \tilde{h}_{2,\varepsilon} \quad (51)$$

We split the integral operator applied to U_ε in two parts

$$\begin{aligned} U_\varepsilon(t) - \int_0^t U_\varepsilon(\tilde{a}) K_\varepsilon(\tilde{a}) d\tilde{a} &= U_\varepsilon(t) - \frac{1}{\mu_0} \int_0^{t/\varepsilon} U_\varepsilon(t - \varepsilon a) \varrho(a) da \\ &= \underbrace{\frac{1}{\mu_0} \int_0^{+\infty} (U_\varepsilon(t) - U_\varepsilon(t - \varepsilon a)) \varrho(a) da}_{:=H_{1,\varepsilon}} + \underbrace{\frac{1}{\mu_0} \int_{t/\varepsilon}^{+\infty} U_\varepsilon(t - \varepsilon a) \varrho(a) da}_{:=H_{2,\varepsilon}} \end{aligned}$$

and we shall specify U_ε such that $H_{1,\varepsilon} \geq \tilde{H}_\varepsilon(t)$ and $H_{2,\varepsilon} \geq 0$. To this end we set

$$U_\varepsilon(t) := \varepsilon(K_0 + K_1 t), \quad \forall t \in \mathbb{R} \quad (52)$$

with constants K_0 and K_1 to be specified. One has obviously that

$$H_{1,\varepsilon}(t) = \frac{\varepsilon^2 K_1 \mu_1}{\mu_0} \geq \varepsilon^2 C_1 \geq \tilde{H}_\varepsilon(t)$$

a.e. on \mathbb{R}_+ , provided that K_1 is chosen as

$$K_1 > \frac{\mu_0}{\mu_1} C_1$$

Using (52) and the change of variable $\tilde{a} = -t/\varepsilon + a$, we obtain that

$$H_{2,\varepsilon} = \int_0^{+\infty} (K_0 - \varepsilon K_1 \tilde{a}) \varrho(\tilde{a} + t/\varepsilon) d\tilde{a} = (K_0 - \varepsilon K_1 A_\varepsilon[\varrho](t)) \int_{\mathbb{R}_+} \varrho\left(\frac{t}{\varepsilon} + a\right) da$$

We are in the hypotheses of Lemma 14 : $A_\varepsilon[\varrho](t)$, the expectation of a given density ϱ with respect to the tail $a > t/\varepsilon$ is bounded by a positive constant A_{\max}

$$A_\varepsilon[\varrho](t) := \frac{\int_{\mathbb{R}_+} a \varrho\left(\frac{t}{\varepsilon} + a\right) da}{\int_{\mathbb{R}_+} \varrho\left(\frac{t}{\varepsilon} + a\right) da} \leq A_{\max}.$$

Therefore it suffices to chose $K_0 > \varepsilon K_1 A_{\max}$ in order to obtain that $H_{2,\varepsilon} \geq 0$. These computations show that U_ε is a super-solution. Then the comparison principle implies that, for all $0 \leq t \leq T$,

$$0 \leq \hat{Z}_\varepsilon(t) = \int_0^t |z_\varepsilon(s) - z_0(s)| ds \leq U_\varepsilon(t) = \varepsilon(K_0 + K_1 t) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0,$$

hence $\hat{Z}_\varepsilon \rightarrow 0$ in $C([0, T])$, which ends the proof since $\|z_\varepsilon - z_0\|_{L^1(0, T)} \leq \|\hat{Z}_\varepsilon\|_{C([0, T])}$. \square

7 A simple example

We construct *by hand* solutions of problems (1) and (3) when the load f is explicitly defined as

$$f(t) := \begin{cases} 1/2 & \text{if } 0 < t \leq \frac{1}{3}, \\ 1 & \text{if } \frac{1}{3} < t \leq \frac{2}{3}, \\ 3/2 & \text{if } \frac{2}{3} < t < 1, \end{cases} \quad (53)$$

and the kernel ϱ is a simple exponential (see more precise statements below). So defined f is of course of bounded variation on $(0, 1)$. The solution z_ε solving (1) and its limit z_0 show different regularities (see Figure 1) : the adhesive approximation is rougher than the limit solution. This is an interesting feature of our approach.




Figure 1: The solutions z_ε and z_0 as a function of time, for a fixed $\varepsilon = 10^{-1}$ on the left y-axis. The load f on the right y-axis.

Assumptions 4.

- i) the load f is defined in (53),
- ii) the on and off rates are constants defined as :

$$\zeta_\varepsilon = \zeta_0 = \zeta, \quad \beta_\varepsilon = \beta_0 = \beta.$$

- iii) the initial condition is at equilibrium :

$$\rho_{I,\varepsilon} = \rho_0 = \frac{\beta\zeta}{\beta + \zeta} e^{-\zeta a}.$$

Lemma 15. Under Assumptions 4, one has that $\mu_{0,\varepsilon} = \mu_{0,0} = \beta/(\beta + \zeta)$, $\mu_{1,\varepsilon} = \mu_{1,0} = \beta/(\zeta(\beta + \zeta))$ and $\rho_0(a) = \mu_{0,0}\zeta e^{-\zeta a}$. Then the solution z_ε of (1) is *BPV*([0, 1]) and it is explicitly given by

$$z_\varepsilon(t) = \int_0^t \frac{f}{\mu_{1,0}} ds + \varepsilon \frac{f(t)}{\mu_{0,0}} + \frac{1}{\mu_{0,0}} \int_0^{+\infty} z_p(-\varepsilon a) \rho_0 da$$

and hence,

$$z_\varepsilon(t) - z_0(t) = \varepsilon \frac{f(t)}{\mu_{0,0}} + \int_{-\infty}^0 z'_p(s) \exp\left(\frac{\zeta s}{\varepsilon}\right) ds$$

with $z_0(t) = z_p(0) + \int_0^t f(s) ds / \mu_{1,0}$ is a continuous functions in $[0, 1]$. Note that the last term is an ε order term according to Assumption 1, indeed it holds that

$$\left| \int_{-\infty}^0 z'_p(s) \exp\left(\frac{\zeta s}{\varepsilon}\right) ds \right| \leq \frac{\varepsilon}{\zeta} \|z_p\|_{W^{1,\infty}(\mathbb{R}_-)}.$$

PROOF. The specific choice of data and kernel allows to rephrase (1) as

$$z_\varepsilon(t) - \frac{\zeta}{\varepsilon} \int_0^t z_\varepsilon(s) \exp\left(-\frac{\zeta(t-s)}{\varepsilon}\right) ds = \varepsilon \frac{f(t)}{\mu_{0,0}} + \frac{\zeta}{\varepsilon} \int_{-\infty}^0 z_p(s) \exp\left(-\frac{\zeta(t-s)}{\varepsilon}\right) ds.$$

Next, setting

$$q_\varepsilon(t) = z_\varepsilon(t) \exp\left(\frac{\zeta t}{\varepsilon}\right), \quad t \geq 0.$$

Then we can rewrite (15) for all $t \geq 0$ as

$$q_\varepsilon(t) - \frac{\zeta}{\varepsilon} \int_0^t q_\varepsilon(s) ds = \varepsilon \exp\left(\frac{\zeta t}{\varepsilon}\right) \frac{f(t)}{\mu_{0,0}} + \frac{\zeta}{\varepsilon} \int_{-\infty}^0 z_p(s) \exp\left(\frac{\zeta s}{\varepsilon}\right) ds. \quad (54)$$

By differentiating (54) in time and using (6), we prove that q_ε solve the equation

$$\begin{cases} \dot{q}_\varepsilon(t) - \frac{\zeta}{\varepsilon} q_\varepsilon(t) = \frac{\exp\left(\frac{\zeta t}{\varepsilon}\right)}{\mu_{0,0}} (\zeta f(t) + \varepsilon f'(t)), & t > 0, \\ q_\varepsilon(0^+) = \varepsilon f(0) / \mu_{0,0} + \frac{\zeta}{\varepsilon} \int_{-\infty}^0 z_p(s) \exp\left(\frac{\zeta s}{\varepsilon}\right) ds \end{cases} \quad (55)$$

and therefore q_ε is explicitly given by

$$q_\varepsilon(t) = \exp\left(\frac{\zeta t}{\varepsilon}\right) \left(\varepsilon \frac{f(t)}{\mu_{0,0}} + \frac{\zeta}{\varepsilon} \int_{-\infty}^0 z_p(s) \exp\left(\frac{\zeta s}{\varepsilon}\right) ds + \int_0^t \frac{f(s)}{\mu_{1,0}} ds \right)$$

which gives the formula of z_ε . Moreover, it is clear that z_ε is of bounded variation since f is it and z_0 is an absolutely continuous function. \square

A Auxiliary proofs

In this appendix, the domain Ω is an open set of \mathbb{R}^n .

A.1 Proof of Theorem 1

As in the proof of [9, Theorem 14.9], the aim is to show that for every $\delta > 0$, there exists a sequence $\{f_\delta\}_\delta$ in $C^\infty(\Omega)$ such that

$$\int_\Omega |f - f_\delta| dt < \delta \quad \text{and} \quad |Df_\delta|(\Omega) < \|Df\|(\Omega) + \delta.$$

Since the total variation $\|Df\|(\Omega)$ is bounded,

$$\lim_{j \rightarrow +\infty} \|Df\|(\Omega \setminus \{t, \text{dist}(t, \partial\Omega) > 1/j, |t| < j\}) = 0$$

then for fixed $\delta > 0$, there exists a $j_0 \in \mathbb{N}$ such that for all $j \geq j_0$,

$$\|Df\|(\Omega \setminus \{t, \text{dist}(t, \partial\Omega) > 1/j, |t| < j\}) \leq \delta.$$

For $i \in \mathbb{N}$, we define the subdomain Ω_i of Ω by

$$\Omega_i := \{t \in \Omega, \text{dist}(t, \partial\Omega) > 1/j_0 + i, |t| < j_0 + i\}$$

such that $\Omega_i \subset\subset \Omega_{i+1}$ and $\bigcup_{i=0}^{\infty} \Omega_i = \Omega$. Let $W_0 = \Omega_1$ and $W_i = \Omega_{i+1} \setminus \bar{\Omega}_{i-1}$, where $\Omega_{-1} = \Omega_0 := \emptyset$, and let $\{\phi_i\}$ be a partition of the unity subordinate to the covering $\{W_i\}_{i \in \mathbb{N}}$

$$\phi_i \in C_0^\infty(W_i), \quad 0 \leq \phi_i \leq 1, \quad \sum_{i=0}^{\infty} \phi_i = 1.$$

For $i \in \mathbb{N}$, the idea is to find $\delta_i > 0$ so small that

$$\text{supp } \chi_{\delta_i} * (\phi_i f) \subset W_i \tag{56}$$

which allows to write

$$\|\chi_{\delta_i} * (\phi_i f) - \phi_i f\|_{L^1(\Omega)} = \|\chi_{\delta_i} * (\phi_i f) - \phi_i f\|_{L^1(W_i)} \tag{57}$$

and then from the local approximation Theorem (see Theorem 2 p.125 in [4]),

$$\|\chi_{\delta_i} * (\phi_i f) - \phi_i f\|_{L^1(W_i)} < C\delta$$

and also by convenience, we can take $C = \frac{1}{2^i}$. Hence we conclude that

$$\int_{\Omega} |\chi_{\delta_i} * (\phi_i f) - \phi_i f| \, dt < \delta 2^{-i} \tag{58}$$

$$\int_{\Omega} |\chi_{\delta_i} * (f \phi_i') - f \phi_i'| \, dt < \delta 2^{-i}. \tag{59}$$

for a positive mollifiers χ_δ defined as

$$\chi_\delta(t) := \frac{1}{\delta} \chi\left(\frac{t}{\delta}\right), \quad \chi(t) = \begin{cases} 1 & \text{if } |t| < 1 \\ 0 & \text{if } |t| > 2 \end{cases}$$

Define

$$f_\delta = \sum_{i=0}^{\infty} \chi_{\delta_i} * (\phi_i f). \tag{60}$$

By the construction of $\{W_i\}$, we have $W_i \cap W_{i+1} \neq \emptyset$ and $W_i \cap W_{i+1} \cap W_{i+2} = \emptyset$ which give that for every $t \in \Omega$,

$$\#\{i \in \mathbb{N} : \chi_{\delta_i} * (\phi_i f)(t) \neq 0\} \leq 2$$

and since the finite sum of infinitely differentiable functions is infinitely differentiable, we conclude that $f_\delta \in C^\infty(\Omega)$ and

$$\int_{\Omega} |f_\delta - f| \, dt \leq \sum_{i=1}^{\infty} \int_{\Omega} |\chi_{\delta_i} * (\phi_i f) - \phi_i f| \, dt < \sum_{i=1}^{\infty} \frac{\delta}{2^i} \leq \delta,$$

since $f = \sum_{i=0}^{\infty} f\phi_i$. Thus $f_\delta \rightarrow f$ in $L^1(\Omega)$ as $\delta \rightarrow 0$. Using the theorem of Lower semi-continuity of variation measure (see chap 5, [4]), we have

$$\|Df\|(\Omega) \leq \liminf_{\delta \rightarrow 0} |Df_\delta|(\Omega). \quad (61)$$

it remains to show that

$$\limsup_{\delta \rightarrow 0} |Df_\delta|(\Omega) \leq \|Df\|(\Omega).$$

Let $\psi \in C_c^\infty(\Omega)$ be such that $|\psi|_\infty \leq 1$. Since $\phi_i f \in \text{BV}(\Omega)$ for every $i \in \mathbb{N}$, we have by Lemma 14.10 in [9] that

$$\int_{\Omega} \chi_{\delta_i} * (\phi_i f) \psi' dt = \int_{\Omega} \phi_i f (\chi_{\delta_i} * \psi)' dt \quad (62)$$

let $\psi_{\delta_i} = \chi_{\delta_i} * \psi$, and using the fact that support of ψ is compact and that the partition of unity is locally finite, we have that

$$\begin{aligned} \int_{\Omega} f_\delta \psi' dt &= \sum_{i=1}^{+\infty} \int_{\Omega} \chi_{\delta_i} * (\phi_i f) \psi' dt \\ &= \sum_{i=1}^{+\infty} \int_{\Omega} \phi_i f \psi'_{\delta_i} dt \\ &= \sum_{i=1}^{+\infty} \int_{\Omega} f((\phi_i \psi_{\delta_i})' - \phi_i' \psi_{\delta_i}) dt := I_1 + I_2 \end{aligned}$$

since $\text{supp}(\phi_i \psi_{\delta_i}) \subset W_i$ and $|\phi_i \psi_{\delta_i}|_\infty \leq 1$, it follows that

$$\begin{aligned} I_1 &= \int_{\Omega} f (\phi_i \psi_{\delta_i})' dt + \sum_{i=2}^{+\infty} \int_{\Omega} f (\phi_i \psi_{\delta_i})' dt \\ &\leq \|Df\|(\Omega) + \sum_{i=2}^{+\infty} \|Df\|(W_i) \\ &\leq \|Df\|(\Omega) + 3\|Df\|(\Omega \setminus \Omega_1) \\ &\leq \|Df\|(\Omega) + 3\delta \end{aligned}$$

since each $t \in \Omega$ belongs at most two of the sets U_i . On the other hand, by Fubini's theorem we have

$$\begin{aligned} I_2 &= - \sum_{i=1}^{+\infty} \int_{\Omega} \chi_{\delta_i} * (f\phi_i') \psi dt \\ &= - \sum_{i=1}^{+\infty} \int_{\Omega} (\chi_{\delta_i} * (f\phi_i') - f\phi_i') \psi dt \end{aligned}$$

by using the fact that $\sum_{i=1}^{+\infty} \phi_i' = 0$. We now use (59) and the fact that $|\psi|_\infty \leq 1$, to conclude that $I_2 \leq \delta$ and then we obtain that

$$\int_{\Omega} f_\delta \psi' dt \leq \|Df\|(\Omega) + 3\delta.$$

By taking the supremum and passing to the limit when $\delta \rightarrow 0$, we obtain that

$$\limsup_{\delta \rightarrow 0} \|Df_\delta\|(\Omega) \leq \|Df\|(\Omega) \quad (63)$$

Finally, (61) and (63) together concludes the proof.

A.2 Proof of Lemma 5:

Let $f \in \text{BV}(\Omega) \cap L^\infty(\Omega)$. By using (60) and (56) we can write for all t in Ω ,

$$\begin{aligned} |f_\delta(t)| &= \left| \sum_{i=0}^{\infty} \chi_{\delta_i} * (f\phi_i)(t) \right| \\ &\leq \sum_{i=0}^{\infty} \|f\|_{L^\infty} \int_{W_i} \phi_i(t-t') \chi_{\delta_i}(t') dt' \\ &\leq \sum_{i=0}^{\infty} \|f\|_{L^\infty} \int_{W_i} \chi_{\delta_i}(t') dt' \end{aligned}$$

since $0 \leq \phi_i \leq 1$. On the other hand, we have $W_i \cap W_{i-1} = \Omega_i \setminus \Omega_{i-1}$ and $W_{i-1} \cap W_i \cap W_{i+1} = \emptyset$ which implies that

$$|f_\delta(t)| \leq \|f\|_{L^\infty} \int_{W_{i-1}} \chi_{\delta_i}(t') dt' + \|f\|_{L^\infty} \int_{W_i} \chi_{\delta_{i-1}}(t') dt' \leq 2\|f\|_{L^\infty(\Omega)} \|\chi\|_{L^1(\mathbb{R})}$$

which ends the proof.

References

- [1] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford: Clarendon Press, 2000.
- [2] D. Anevski. Riemann-stieltjes integrals. *Mathematical Sciences, Lund University, Lund, Sweden*, 2012.
- [3] J. J. Benedetto and W. Czaja. *Integration and modern analysis*. Basel: Birkhäuser, 2009.
- [4] L. C. Evans and R. F. Gariepy. *Measure theory and fine properties of functions*. Boca Raton: CRC Press, 1992.
- [5] P. Gabriel. *Équations structurées en dynamique des populations*. ENS Cachan, Dec. 2013. Licence.
- [6] G. Gripenberg, S.-O. Londen, and O. Staffans. *Volterra integral and functional equations*, volume 34 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1990.
- [7] M. Heida, R. I. A. Patterson, and D. R. M. Renger. Topologies and measures on the space of functions of bounded variation taking values in a Banach or metric space. *J. Evol. Equ.*, 19(1):111–152, 2019.
- [8] E. Kreyszig. *Introductory functional analysis with applications*. New York etc.: John Wiley & Sons. XIV, 688 p. £15.00; \$ 27.00 (1978)., 1978.
- [9] G. Leoni. *A first course in Sobolev spaces*, volume 105. Providence, RI: American Mathematical Society (AMS), 2009.
- [10] A. Manhart, D. Oelz, C. Schmeiser, and N. Sfakianakis. An extended Filament Based Lamellipodium Model produces various moving cell shapes in the presence of chemotactic signals. *J. Theoret. Biol.*, 382:244–258, 2015.
- [11] V. Milišić and D. Oelz. Tear-off versus global existence for a structured model of adhesion mediated by transient elastic linkages. *Commun. Math. Sci.*, 14(5):1353–1372, 2016.

- [12] V. Milišić and D. Oelz. On the asymptotic regime of a model for friction mediated by transient elastic linkages. *J. Math. Pures Appl. (9)*, 96(5):484–501, 2011.
- [13] V. Milišić and D. Oelz. On a structured model for load-dependent reaction kinetics of transient elastic linkages mediating nonlinear friction. *SIAM J. Math. Anal.*, 47(3):2104–2121, 2015.
- [14] G. A. Monteiro, A. Slavík, and M. Tvrdý. *Kurzweil-Stieltjes integral. Theory and applications*, volume 15. Hackensack, NJ: World Scientific, 2019.
- [15] D. Oelz and C. Schmeiser. Derivation of a model for symmetric lamellipodia with instantaneous cross-link turnover. *Arch. Ration. Mech. Anal.*, 198(3):963–980, 2010.
- [16] D. Oelz, C. Schmeiser, and J. Small. Modeling of the actin-cytoskeleton in symmetric lamellipodial fragments. *Cell adhesion & migration*, 2:117–26, 05 2008.
- [17] D. Oelz, C. Schmeiser, and J. V. Small. Modeling of the actin-cytoskeleton in symmetric lamellipodial fragments. *Cell Adhesion & Migration*, 2(2):117–126, 2008.
- [18] B. Perthame. *Transport equations in biology*. Basel: Birkhäuser, 2007.
- [19] N. Sfakianakis, D. Peurichard, A. Brunk, and C. Schmeiser. Modelling cell-cell collision and adhesion with the filament based lamellipodium model. *Biomath*, 7(2):78–91, 2018.
- [20] W. P. Ziemer. *Weakly differentiable functions. Sobolev spaces and functions of bounded variation.*, volume 120. Berlin etc.: Springer-Verlag, 1989.