



HAL
open science

Copper Content and Export in European Vineyard Soils Influenced by Climate and Soil Properties

Boris Droz, Sylvain Payraudeau, José Antonio Rodríguez Martín, Gergely Tóth, Panos Panagos, Luca Montanarella, Pasquale Borrelli, Gwenael Imfeld, José Antonio, Rodríguez Martín

► **To cite this version:**

Boris Droz, Sylvain Payraudeau, José Antonio Rodríguez Martín, Gergely Tóth, Panos Panagos, et al.. Copper Content and Export in European Vineyard Soils Influenced by Climate and Soil Properties. Environmental Science and Technology, 2021, 55 (11), pp.7327-7334. 10.1021/acs.est.0c02093 . hal-03378939

HAL Id: hal-03378939

<https://hal.science/hal-03378939>

Submitted on 18 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Copper Content and Export in European Vineyards
2 Soils Influenced by Climate and Soil Properties

3 *Boris Droz,^{†*} Sylvain Payraudeau,[†] José Antonio Rodríguez Martín,[‡] Gergely Tóth,[§]*
4 *Panos Panagos,^{||} Luca Montanarella,^{||} Pasquale Borrelli^{□, ¥} & Gwenaël Imfeld^{†*}*

5 [†] Institut Terre et Environnement de Strasbourg, Université de Strasbourg, ENGEES,
6 CNRS, UMR 7063, 5 rue Descartes, Strasbourg F-67084, France

7 [‡] Department of the Environment, Instituto Nacional de Investigación y Tecnología
8 Agraria y Alimentaria (INIA), Carretera de la Coruña 7.5, 28040 Madrid, Spain

9 [§] Institute of Advanced Studies, Chernek str. 14., 9730 Kőszeg, Hungary

10 ^{||} European Commission, Joint Research Centre (JRC), Institute for Environment and
11 Sustainability, Via E. Fermi 2749, 21027 Ispra, Italy

12 [□] Department of Earth and Environmental Sciences, University of Pavia, Via Ferrata, 1,
13 27100 Pavia, Italy

14 [¥] Department of Biological Environment, Kangwon National University, Chuncheon
15 24341, Republic of Korea

16

17 ABSTRACT

18 Copper-based fungicides (Cu_f) are used in European (EU) vineyards to prevent fungal
19 diseases. Soil physicochemical properties govern locally the variation of total copper
20 content (Cu_t) in EU vineyards. However, variables controlling Cu_t distribution at larger
21 scale are poorly known. Here machine learning techniques were used to identify
22 governing variables and to predict the Cu_t distribution in EU vineyards. Precipitation,
23 aridity and soil organic carbon are key variables explaining together 45% of Cu_t
24 distribution across EU vineyards. This underlines the effect of both climate and soil
25 properties on Cu_t distribution. The average net export of Cu at the EU scale is 0.29 kg
26 Cu ha^{-1} which is two orders of magnitude less than the net accumulation of Cu (24.8 kg
27 Cu ha^{-1}). Four scenarios of Cu_f application were compared. The current EU regulation
28 with a maximum of $4 \text{ kg Cu ha}^{-1} \text{ year}^{-1}$ may increase by 2% the EU vineyard area
29 exceeding the predicted no-effect concentration (PNEC) in soil in the next 100 years.
30 Overall, our results highlight vineyard areas requiring specific remediation measures
31 and strategies of Cu_f use to manage a trade-off between pest control and soil and water
32 contamination.

33 INTRODUCTION

34 Copper-based fungicides (Cu_f) are intensively used in European vineyards since the
35 end of the 19th century to prevent fungal disease such as 'Downy Mildew'.¹ Cu_f
36 application doses and timing vary across regions depending on practices, regulation,
37 hydro-climatic conditions and vine variety. The European Union (EU 2018/1981)
38 regulation recently decreased from 6 to 4 kg Cu ha⁻¹ year⁻¹ the maximal dose allowed.²
39 However, doses as high as 50 kg Cu ha⁻¹ year⁻¹ were applied frequently for decades in
40 the mid-20th century.³ Today, the total copper content in topsoil (Cu_t) exceeds 100 mg
41 Cu kg⁻¹ in 15% of the European vineyard area. This corresponds to the average
42 proposed threshold values for which soil remediation is needed.⁴⁻⁶ European vineyards
43 represent more than 50% of the total vineyard surface worldwide.⁷ Although vine
44 growing accounts for only 3.3% of the agricultural area in Europe, it uses 86% of total
45 fungicides consumed in Europe.⁸ Hence, vineyard soil is often more contaminated by
46 Cu_f than soils of any other agricultural soils. Cu_f accumulates in soil and can impact soil
47 organisms and plants,⁹ thereby reducing soil fertility¹⁰ and productivity.¹¹ In contrast,
48 leaching¹² and surface runoff^{13,14} export Cu_f from vineyard soil, which can contaminate
49 aquatic ecosystems¹⁵ and drinking water resources.¹⁶

50 Cu_f mobility in topsoil controls Cu_t accumulation and off-site export. Cu mobility in soil
51 is mainly controlled by (i) sorption on organic matter,¹⁷ and to a lesser extent on clay
52 and Fe-, Mn-(hydr)oxide and carbonate,¹⁸ (ii) pH-redox dependency since alkaline soil
53 increase the proportion of available binding sites on organic matter.^{17,19} Cu sorbed on
54 fine particle matter may result in Cu export in runoff.^{13,14} Hydro-climatic and field
55 conditions indirectly control Cu export by (i) affecting soil properties,²⁰ (ii) mediating

56 dissolved and particulate Cu_f transport, and (iii) modifying preferential transport.²¹
57 Accordingly, presumed or established processes governing Cu_t mobility in vineyard soil
58 may co-occur and/or may be interrelated (Table 1). Variables governing Cu_t and
59 dynamics in vineyard soil are, however, examined in laboratory and field studies
60 separately. As a result, the understanding of variables governing Cu_t in soil remains
61 study- or site-specific. The identification of variables governing Cu_t in vineyard soil and
62 leading to Cu accumulation and/or off-site export over time is necessary to evaluate the
63 current and future distribution of Cu_t at the European scale.

64 In this context, a major issue is to predict the spatial variation of Cu_t in the European
65 vineyards, to quantify Cu accumulation and export and to identify priority areas requiring
66 soil remediation. The purpose of this study was (i) to identify key variables potentially
67 governing Cu_t in vineyard topsoil (0 – 30 cm), (ii) to predict average Cu_t at European
68 vineyards scale using machine learning technics, and (iii) to estimate Cu_t accumulation,
69 export and associated toxicity for terrestrial trophic levels, for historical, current and
70 future scenarios of Cu_f application (i.e, 0, 2, 4 or 8 kg Cu ha⁻¹ y⁻¹).

71

72 MATERIALS AND METHODS

73 **Topsoil total Copper Datasets.**

74 Cu_t datasets ($Cu_{t, \text{dat}}$; i.e., soils within 0–30 cm) were obtained from several soil
75 surveys across European vineyards (–9.4° to 42° E; 35° to 53° N; SI, Figure S1)
76 delineated by the Corine Land Cover (CLC 2012; v18.5.1; <http://land.copernicus.eu>). In
77 total, 1,202 $Cu_{t, \text{dat}}$ measurements (average: 27.8 mg Cu kg⁻¹; min–max: 0.54–774 mg
78 Cu kg⁻¹) were sampled between 1992 and 2016 (median = 2008, standard deviation

79 (SD) = 5) and used in our study. Summary of the studied $Cu_{t, dat}$ are provided in Table
80 S1 of the Supporting Information (SI). Although the Cu quantification methods differ
81 among datasets, inter-comparability between datasets has been demonstrated
82 previously for the methods used in our study.²² The impact of topsoil sampling depth
83 (first 20 or 30 cm) and sampling time span on the average prediction of Cu_t (Cu_{pred}) at
84 the European scale remains low (SI, section Influence of heterogeneous sampling
85 depths and time on the Cu prediction). The heterogeneity of topsoil sampling depth and
86 time span did not significantly (within 2σ) impact the average predictions of Cu_t at the
87 European scale.

88

89 **Predictive Variables.**

90 Soil properties, topology, climate and land cover were selected and evaluated as
91 hypothetical predictive variables of Cu_t (Table 1 & SI, Table S2). Geogenic, i.e., from bed
92 rocks,²³ and atmospheric non-agricultural anthropogenic inputs of Cu, i.e., mining,
93 smelter, sewage sludge and vehicle brake pads are sources of Cu contributing from 5 to
94 $100 \text{ g Cu ha}^{-1} \text{ year}^{-1}$ whereas about $8 \text{ kg Cu ha}^{-1} \text{ years}^{-1}$ of Cu_f are applied on
95 vineyards.²⁴ Topsoils surrounding Cu mining or smelter, that may reach up to 6 g Cu kg^{-1}
96 ^{1, 25, 26} were not identified close to vineyard areas. Consequently, geogenic and industrial
97 sources of Cu were not considered in this study.

98

99

100

101

102 **Table 1.** Hypothetic Predictive Variables Governing the Total Copper Content in
 103 Vineyard Topsoil (Cu_t) Leading to Cu_t Accumulation and/or off-site Export.

Variables ^a	Processes
Soil properties	
Soil physicochemical properties affect Cu_t mobility	Cu_t correlates positively to soil organic matter since Cu preferentially sorbs to organic matter in soil, ^{13,17} which affects Cu_t mobility. Higher pH increases sorption sites (e.g. negative charge) for Cu_t on clay, organic matter and sand containing carbonate, ¹⁹ and reduces Cu_t leaching. ¹²
Soil moisture influences soil biogeochemistry	Soil moisture directly affects microbial respiration and activities in soil resulting in biogeochemical gradients. ^{27,28} Cu occurs as Cu(II) in oxic soils. In contrast, Cu(II) may potentially precipitate to less mobile Cu(I) forms in anoxic soils. ^{17,29}
Topography/climate	
Topography and climate conditions drive soil erosion	Greater rainfall depth and higher slope length enhance soil erosion by runoff. ²¹ Cu_t is mostly bound to suspended particles matter (SPM) in topsoil (~85% ^{13, 14}) where SPM-bound Cu can account for 84.4% of the total Cu_t exported by runoff. ¹⁴
Hydro-climatic conditions influence Cu application doses	Intense rainfalls during warm conditions favor the development of 'Downy Mildew'. This increases the frequency and amount of Cu_t use in some areas to control it. ^{30,31}
Hydro-climatic conditions influence (i) soil chemistry	Arid environments (with higher Al values), characterized by higher soil pH and oxidizing soils, ²⁰ favor Cu_t accumulation (see soil physicochemical properties above). In contrast, arid soils tend to lose more soil organic matter and associated Cu_t since Cu is preferentially associated to soil organic matter. ^{13,17}
(ii) Vegetation growth	Optimal conditions for vegetation growth depend on temperature, soil moisture and solar radiation, and vary across plant growth stages. ³² Vegetation growth controls Cu_t mobility indirectly by limiting erosion via surface runoff and thus can be related to Cu_t .

(iii) Plant uptake Cu uptake from soil by plants remains limited: 0.1–7.2% by wild³³ and 0.1–3.8% by the vineyard³⁴ plants of the Cu_t. Hence, plants only marginally decrease Cu_t in soil.

Plant cover

Plant cover affects surface runoff and organic matter content in topsoil

Plant cover, i.e., vine and grass in vineyards, can stabilize the vineyard soil and thus limit erosion via surface runoff.³⁵ In addition, plant decomposition can slightly increase organic matter content in topsoil,³⁶ thereby reducing Cu mobility.

104 ^a Hypothetic predictive variables governing Cu_t in vineyard soil. Variables associated
105 with each process are provided in the SI, Table S2. Processes were identified based on
106 experimental studies in vineyard or related soil.

107

108 Predictive variables were computed or directly obtained on a grid cell of 250x250 m
109 for the study area. Details concerning building spatial variables are provided in the in SI,
110 Table S2 & S3. Cu_{t, dat} coordinates were used to extract values of each predictive
111 variable in the corresponding grid cells. Extracted data were then used to select
112 variables and calibrate the model (details in SI). Extracted data represented adequately
113 the range of predictive variables within the prediction domain (details given in SI; paired
114 t-test; p-value_{average} >0.01) and did not display any spatial correlation (Moran's test;
115 I_{average} = -0.211 ± 0.09; p-value >0.01). The selection of variables was carried out from
116 the 24 original predictive variables (SI, Table S2) to limit collinearity between predictive
117 variables and optimize model predictions as described elsewhere.³⁷ (i) Pearson
118 correlation was used to discard variables providing the similar information and (ii)
119 principal component analysis (PCA) was used to discard noise variables containing the
120 least information.³⁸

121

122

123 **Prediction of Total Copper Content in European Vineyard Topsoil.**

124 The variable selection procedure highlight that the following variables were predictive
125 of the average Cu_t in European vineyards: clay and silt fractions (%), soil organic carbon
126 content (C_{org} ; %), soil pH (unitless), aridity index (AI; unitless), annual average
127 temperature (T_{av} ; °C), annual rainfall (rainfall; mm), soil moisture content (moisture; %),
128 the enhanced vegetation index (EVI; unitless) and slope (degree; SI, Table S3). This
129 selection is consistent with predictive variables reported previously (Table 1). Three
130 independent machine learning techniques, i.e., neural networks (R package nnet v 7.3-
131 12), random forest (R package randomForest v.4.6-12) and bagging tree (R package
132 ipred v.0.9-5), were used in parallel and were averaged into an ensemble model (ENS;
133 details in SI) for the final prediction, as described elsewhere.³⁹ The ENS resulted in
134 fewer false predictions and higher predictive accuracy than a single model. Consistency
135 of Cu_{pred} was evaluated for each grid cell on a SD of 1,000 predictions for each machine
136 learning technique. Model performance was evaluated using a 10-fold cross-validation
137 (CV). Model spatial transferability was evaluated using a spatially constrained 10-fold
138 CV (details in SI). For each model, the averaged relative importance (ARI) of each
139 variable in the model was calculated with an input permutation technique and one-
140 factor-at-a-time (OFAT) sensitivity analysis (details in SI). The complete modelling
141 framework to select variables, calibrate the model and predict Cu_t is available for
142 download at https://github.com/Boris-Droz/ML_soil_predi_conc.

143

144

145

146 **Copper Accumulation and Export.**

147 Cu_t net mass accumulation since the beginning of the Cu_f application was estimated
148 assuming that Cu_t is distributed similarly across a homogenous topsoil layer, following
149 eq. 1,

$$150 \text{ net acc.} = (Cu_{Pred} - Cu_{bgd}) \times BD \times h_{topsoil} \quad (1)$$

151 where *net acc.* stands for the Cu_t net mass accumulation ($kg\ Cu\ ha^{-1}$), Cu_{bgd} the
152 background of geogenic and atmospheric non-agricultural anthropogenic Cu_t ($mg\ Cu$
153 kg^{-1} ; SI, Figure S2), BD is the soil bulk density (<https://soilgrids.org/>, series M_sl1,
154 v18.4.2017; in $kg\ m^{-3}$)⁴⁰ and the average topsoil thickness where Cu mainly
155 accumulates ($h_{topsoil} = 30\ cm$; SI, Table S1). The net mass export of Cu_t (*net exp.*; kg of
156 $Cu\ ha^{-1}$) was assumed to be the sum of Cu_t exported from topsoil via runoff and
157 leaching, and was calculated yearly following eq. 2,^{13, 14}

$$158 \text{ net exp.} = (\dot{q}_{RUSLE} + f_{sol} \times BD \times h_{topsoil}) \times Cu_{Pred} \quad (2)$$

159 where \dot{q}_{RUSLE} is the soil loss rate ($kg\ ha^{-1}$) as defined elsewhere²¹ and f_{sol} the soluble
160 fraction of Cu_{Pred} estimated from soil properties (both details in SI).

161

162 **Scenarios of Copper-Based Fungicides Application Doses.**

163 Four scenarios of Cu_f application doses were compared: (i) an historical scenario with
164 $8\ kg\ Cu\ ha^{-1}\ year^{-1}$ inferred from sold Cu_f amount before the 2002 European
165 regulation,² and/or speculated illegal usage in some areas,⁴¹ (ii) current regulation
166 scenarios, with either (iii) 2 or (iii) 4 $kg\ Cu\ ha^{-1}\ year^{-1}$ as average and maximum doses,

167 respectively,^{2, 42} and (iv) a scenario of Cu_f substitution, i.e., no Cu_f application or 0 kg Cu
168 ha⁻¹ year⁻¹.

169 For the scenario with 8 kg Cu ha⁻¹ year⁻¹ and current regulation scenarios, a stepwise
170 mass balance calculation based on a year t was established (eq. 3 and detail in SI)
171 assuming no temporal variation of the topsoil volume and a homogeneous topsoil layer
172 (0–30 cm),

$$173 \quad Cu_{pred,t+1} = Cu_{pred,t} + (APP_{Cu,f} - Cu_{pred,t} \times net\ exp. - Cu_{input} \times f_{sol}) / (BD \times h_{topsoil}) \quad (3)$$

174 where Cu_{pred} is the average predictions of Cu_t (mg Cu kg⁻¹), $APP_{Cu,f}$ the yearly
175 application dose of Cu_f (kg Cu ha⁻¹ year⁻¹) and f_{sol} the soluble fraction as calculated in
176 eq S3–6. The *net exp.* was previously calculated in eq. 2. $Cu_{pred,t+1}$ accounted for the
177 yearly Cu application dose to which the soluble fraction of Cu_f (f_{sol}) and the *net exp.*
178 were deducted. For each, pixel, the scenario was applied until Cu_{pred} at year $t + 1$
179 reaches for the given pixel the predicted no-effect concentration (PNEC; SI, building
180 spatial variables). In contrast, the time required to export the accumulated Cu_f via runoff
181 and associated suspended solids and reach back Cu_{bgd} , i.e., assuming the existence of
182 alternatives to Cu_f, was estimated by dividing eq. 1 by eq. 2.

183 The uncertainty associated with each scenario was estimated by numerical
184 propagation of uncertainties across the entire modelling framework. Briefly, each
185 variable was associated with lower and an upper uncertainty limits accounting for the
186 analytical or monitoring uncertainty or calculation steps, depending on the variable (SI,
187 Table S4). All combinations of lower and upper uncertainty limits for the selected
188 variables were then included to propagate uncertainties and estimate the lower and
189 upper uncertainty limits associated to each scenario for each pixel independently.

190

191

192 RESULTS AND DISCUSSION

193 **Prediction of Total Copper in Topsoil.**

194 Ten variables governing Cu_t in vineyard topsoil were considered (see above Materials
195 and Methods and SI, Figure S4. b). Using these climate and soil variables, the ENS
196 accurately predicted Cu_t (slope of 1.13, intercept of 0.01; SI, Figure S4. a), and
197 predictions were fairly precise (averaged $R^2 = 0.61$). Comparison between ENS and CV
198 (averaged R^2 -CV = 0.58 ± 0.06) indicates no over-fitting of the ENS. For the three
199 machine learning technics, Cu_{pred} fit $Cu_{t, dat}$ ($SD\ Cu_{pred}/Cu_{t, dat} \times 100 = 13 \pm 2\%$),
200 indicating a good overall agreement of predictions between the three technics. Average
201 uncertainty of the Cu_{pred} falls within $\pm 11\text{ mg Cu kg}^{-1}$ for 75% of the data (RMSE of 0.63
202 ± 0.06). The scattering observed in the predictions (SI, Figure S4) mainly concerns data
203 above 130 mg Cu kg^{-1} , corresponding to 6.3% of the dataset only. ENS predictions
204 failed to reproduce observed Cu_t in the corresponding vineyard areas, which potentially
205 received larger historical Cu_f applications than other areas under similar climatic and
206 soil conditions. Additional model performances and limits are provided in SI.

207

208 **Climate Variables rather than Soil Variables Determine Total Copper in Topsoil.**

209 Climate variables (i.e., rainfall and aridity index) mainly determined Cu_{pred} at the
210 European scale, possibly by controlling physicochemical processes, including Cu
211 leaching and the redox state of soil. Hydro-climatic conditions likely determined regional
212 Cu application practices as 'Downy Mildew's' pressure increases in more humid and

213 warmer areas in early summer.^{30,31} In contrast, soil physicochemical properties (i.e., pH)
214 predominantly governed Cu_t on the local scale and under homogeneous climatic
215 conditions. The averaged relative importance (ARI) from the ENS and the sensitivity
216 analyses confirmed the relationship between climate variables and Cu_{pred} . Cu_{pred}
217 correlated positively with both the annual rainfall (rain; $R^2 = 0.71$) and the aridity index
218 (AI; potential evapotranspiration divided by rainfall; $R^2 = 0.79$), while the combined
219 effect of rainfall and AI accounted for 32.6% of the Cu_{pred} ARI (rainfall = $17.4 \pm 0.9\%$, AI
220 = $15.2 \pm 1.3\%$; SI, Figure S4. b).

221 Although rainfall was negatively correlated with AI, both variables were positively
222 correlated to Cu_{pred} , suggesting that two different mechanisms control Cu_t in vineyard
223 soil. First, rainfall could increase temporally soil water content and reduce oxygen
224 diffusion in water, leading to reducing soil conditions.⁴³ Under anoxic conditions,
225 biological activity may decrease Cu mobility by precipitating soluble Cu(II) to Cu(I),^{44,45}
226 which is strongly complexed and stabilized with reduced organic sulfur.⁴⁶ Additionally,
227 depending on the solutes present in soil, e.g., carbonate, sulphide, biotic and abiotic
228 processes may form minerals and further reduce the Cu_t mobility.^{33, 47, 48}

229 In contrast, higher AI leads to drier soil,⁴⁹ which facilitates the formation of soil
230 macropores favoring oxygen transfer.⁵⁰ As a result, Cu(I) may partly re-oxidize into
231 mobile Cu(II) in oxic soil. However, an opposite trend is observed as AI may indirectly
232 increase the Cu-binding affinity of soil organic matter (ARI of $13.1 \pm 0.1\%$). Indeed, the
233 oxidation of organic matter under oxic conditions increases the contribution of carboxylic
234 and hydroxylic groups as preferential binding partners for Cu in organic matter.⁵¹ In


235 addition, pH (ARI of $7.5 \pm 1.3\%$) directly controls the deprotonation of sorption sites on
236 soil constituent. Hence, higher pH increases available sorption sites for Cu binding.¹⁹

237

238 **Distribution of the Total Copper in Topsoil Across European Vineyards.**

239 Cu_{pred} in European vineyards were successfully made as the datasets covered the
240 range of predictive variables within the domain of prediction. The ENS also displays
241 good model transferability in space evaluated on four equivalent geographical sub-
242 domains (average $R^2 = 0.84 \pm 0.07$, details in SI). This suggests a homogenous
243 influence of predictive variables within the European vineyards. Overall, Cu_{pred} for
244 vineyards is $25.4 \text{ mg Cu kg}^{-1}$ which is 30.5% higher than Cu_{bgd} including geogenic and
245 atmospheric non-agricultural anthropogenic Cu (Figure 1. a–b).

246



247
 248 **Figure 1.** Distribution of the total copper content in vineyard topsoil (Cu_t), accumulation
 249 and net export of Cu in European Vineyards. (a) Observed background Cu_t . (b)
 250 Predicted Cu_t (Cu_{pred}). (c) Estimated net accumulation of Cu_t . (d) Estimated net export of
 251 Cu_t . Dark grey: Area of the European Corine Land Cover (CLC). Pixels depict average
 252 predictive values at a spatial resolution of 20 km^2 . Net accumulation was not computed
 253 for Turkey due to absence of background data. Overall, Cu_{pred} were higher for vineyards

254 in the alpine regions (Figure 1. b; average = 57 mg Cu kg⁻¹) than in other European
255 biogeographic regions (SI, Table S5).

256
257 The threshold of contamination (>100 mg Cu kg⁻¹)⁵ was reached for 2% of Cu_{pred}
258 within the prediction domain. Contaminated areas were mainly located in central
259 Western Europe, i.e., northern Italy, eastern France, Switzerland and Slovenia. In those
260 regions, C_{org} is medium (up to 14%) while Cu_{pred} >100 mg Cu kg⁻¹ accounted for 70% of
261 vineyard soil.⁵² Data points above the contamination threshold in those regions are
262 characterized by high AI, soil moisture and annual rainfall, although only C_{org} was
263 significantly higher (>9.0%) than in other regions of southern Europe (t-test; p-value
264 <0.01). This suggests that higher annual rainfall correlates with higher temperature in
265 central Western Europe. This may favor 'Downy Mildew' emergence and concomitant
266 use of higher doses and more frequent applications of Cu_f. In central Western Europe,
267 soil with higher C_{org} may influence the Cu_{pred} predominantly and locally, although C_{org}
268 accounted in total for 13.1% ARI of the Cu_{pred}.

269

270 **Accumulation and Export of Copper in Topsoil.**

271 The estimated average net accumulation and net export of Cu in topsoil in European
272 vineyards are 24.8 (SD = 42.7) and 0.29 kg Cu ha⁻¹ (SD = 0.64), respectively (Figure 1
273 c–d). This corresponds to 112,400 tons of Cu accumulated over time, and a yearly Cu
274 export from vineyards of 1,466 tons across Europe. The distribution of Cu net export
275 fitted with that of Cu_{pred} (R² = 0.65). On average, predicted loss rate of vineyard soil
276 (9.47 tons ha⁻¹ year⁻¹) is 3.5 times greater than any other European arable land due to


277 low vegetation and sloping area in vineyards. Consequently, net Cu export is
278 particularly large in the Southern Alps due to intense soil erosion. Large soil loss is
279 associated to higher slope ($>25^\circ$) and greater rainfall amounts ($>1,600 \text{ mm year}^{-1}$) in
280 the Southern Alps than in other vineyard areas.²¹

281

282 **Potential Future Copper Accumulation in European Vineyards.**

283 Overall, the proportion of European vineyard area exceeding PNEC values (i.e.,
284 ranged from 30 to 290 mg Cu kg⁻¹ according to soil properties calculate elsewhere,^{53, 54}
285 details in SI) may largely vary in the future depending on the application dose of Cu_f
286 (Figure 2). Cu accumulation due to Cu_f application doses of 2, 4 and 8 kg Cu ha⁻¹ year⁻¹
287 may be balanced with Cu export after 1,200, 750 and 150 years, with 75, 89, 97% of
288 European vineyards areas exceeding PNEC values, respectively. Assuming constant
289 soil properties, we speculate that the current European regulation with a maximum
290 application of 4 kg Cu ha⁻¹ year⁻¹ will increase by only 2% the proportion of vineyard
291 areas exceeding PNEC in the next 100 years.

292 In contrast, 94% of vineyard areas may exceed PNEC in the historical application
293 scenario of 8 kg Cu ha⁻¹ year⁻¹. With an application of 2 kg Cu ha⁻¹ year⁻¹, vineyards
294 areas exceeding PNEC values may increase by less than 0.5% for the next 100 years.
295 In contrast, a substitution of Cu_f (0 kg Cu ha⁻¹ year⁻¹) may decrease the proportion of
296 vineyard areas exceeding the background Cu_{bgd} from 48 to 38% after 100 years. Such
297 long-term indicative estimation for different Cu application dose provides critical
298 temporal trend to support the reduction of Cu application in vineyard.



299

300 **Figure 2.** Proportion of European vineyards area exceeding predicted no-effect
 301 concentration (PNEC) values of total Cu in topsoil (scenarios with Cu_f application dose
 302 of 2, 4, 8 $\text{kg ha}^{-1} \text{ year}^{-1}$) or equal to or lower than the background (bgd) Cu content
 303 (scenario with no Cu_f application). Predictions account for the balance between yearly
 304 Cu_f application at a given application dose and Cu export over time. Dashed lines
 305 indicate the upper and lower uncertainties of predictions.

306

307 Our back-envelope is a first attempt to evaluate Cu accumulation and export at the
 308 European scale and the proposed mass balance approach may lead to inaccurate
 309 estimations of Cu_t export. For instance, freshly applied Cu_f is a soluble Cu pool that can
 310 be directly exchanged with the labile cation present in vineyard soil following Cu_f

311 application. Depending on the time following Cu_f application, Cu_f may be partly mobilized
312 during rainfall-runoff events¹⁴ and thus bias estimations of Cu_t in soil. Since rainfall
313 erosivity is predicted to increase by 18% in Europe over time, Cu_f export may also
314 increase.⁵⁵ In addition, estimations using the semi empirical model account for Cu aging
315 in contaminated soil and rely on realistic desorption experiments, which may
316 underestimate Cu_f export.¹² However, Cu_f accumulation over time may be also
317 underestimated. Climate change models predict an average of 10% increase of soil
318 organic matter content by 2100 in regions of European vineyards,⁵⁶ which may in turn
319 reduce Cu_t mobility and increase the amount of Cu accumulated in the vineyard soil.

320

321 **Implications for Winegrowing Practices.**

322 Cu is an element that cannot be degraded in soil and soil remediation processes,
323 such as phyto-extraction⁵⁷ or enhanced mobilization by plants,⁵⁸ have limited
324 efficiencies. The most efficient strategies to limit the increase of Cu_t is thus to reduce
325 application doses of Cu_f . Reducing strategies consist in applying Cu_f during long humid
326 periods only⁴² and in reducing preventively the use of Cu_f . Forecast models of
327 grapevine 'Downy Mildew'⁵⁹ relying on climate variables can help to improve the
328 efficiency of Cu_f reduction strategies. Alternatives to Cu_f include replacing current grape
329 varieties with new grape varieties with a higher resistance to fungus pests,⁶⁰ using
330 microorganisms for biocontrol,⁶¹ introducing plant defense stimulators⁶² and/or applying
331 plant extracts to substitute Cu_f .⁶³ Alternative scenarios without Cu_f are still debated, but
332 are currently used in some Biodynamic approaches accounting for only ca. 1% of
333 vineyards worldwide (<http://www.demeter.net>). Additional benefits of biodynamic wine

334 growing include the higher soil biodiversity and microbiological activity¹⁰ and potentially
335 greater resilience to climatic and pathogenic threats in vineyards⁶⁴ as opposed to most
336 conventional wine growing.⁶⁵

337 Cu export is mainly driven by soil erosion with concomitant loss of solid-bound Cu,
338 which should be prevented and lessened in regions prone to soil loss to reduce
339 contamination of connected aquatic ecosystems. In fact, 10% of the vineyards are
340 located close to a river (<500 m)⁶⁶ and the associated catchment may act as a point
341 source of contamination. The transport of solid-bound Cu into aquatic ecosystems may
342 also affect the downstream areas due to the progressive release of toxic Cu(II) resulting
343 from a change in pH and/or redox conditions. Anti-erosion soil management strategies
344 should thus be prioritized and designed to retain Cu on the vineyard plots. In particular,
345 traditional soil management strategies, including grass or mulch cover, reduced tillage,
346 contour planting and terraced vineyards are most effective in limiting soil erosion and
347 associated Cu runoff.³⁵ Complementarily, stormwater wetlands may be deployed at the
348 outlet of vineyard catchments. Stormwater wetlands have the potential to retain >68% of
349 the dissolved Cu and >92% of the solid-bound Cu, the principal contributions to Cu
350 export in runoff.¹⁴

351

352 ASSOCIATED CONTENT

353 Supporting Information

354 The Supporting Information is available free of charge on the ACS Publications
355 website at DOI: XXX/XXXXX.

356 Texts, tables, and figures provide detailed methods (total copper in the topsoil dataset,
357 spatial variable building, data pre-processing, variable selection, machine learning
358 building, performance and evaluation, scenario of copper-based fungicide application)
359 and supporting results and discussion (model performance, regional prediction, limit of
360 the approach).

361
362 AUTHOR INFORMATION

363 * Corresponding author. e-mail: bodroz@bluewin.ch; imfeld@unistra.fr phone: + 33 3
364 6885 0474

365 ORCID

366 Boris Droz: 0000-0002-3942-704X

367 Sylvain Payraudeau: 0000-0003-2546-7857

368 José Antonio Rodríguez Martín: 0000-0002-9158-9564

369 Gergely Tóth: 0000-0002-4201-1678

370 Panos Panagos: 0000-0003-1484-2738

371 Luca Montanarella: 0000-0002-4176-3878

372 Pasquale Borrelli: 0000-0002-4767-5115

373 Gwenaël Imfeld: 0000-0002-7054-4666

374

375 Author contributions

376 B.D., S.P. and G.I. designed research, B.D. performed research; S.P., J.A., R.M., G.T.,

377 P.P., L. M. and P.B. contributed to the acquisition of data/soil samples; B.D. and S.P.

378 analyzed data; and B.D., S.P. and G.I. wrote the manuscript with critical review from all
379 authors. The authors declare no conflict of interest.

380

381 ACKNOWLEDGEMENTS

382 Boris Droz was supported by a fellowship of the Region Grand Est and the Rhine-
383 Meuse Water Agency (AERM). Pasquale Borrelli was funded by the EcoSSSoil Project,
384 Korea Environmental Industry & Technology Institute (KEITI), Korea (Grant No.
385 2019002820004). We acknowledge the copper dataset owners for sharing their data:
386 Holger Tülp (Mosel, Germany), Denis Rusjan (Slovenia), Maria Concepcion Ramos
387 (Spain), Nikolai Dinev (Bulgaria), several Swiss state environmental agencies
388 (Schaffhausen, Neuchâtel, Genève, Tecino, Aargau, Valais), Central Institute for
389 Supervising and Testing in Agriculture (ÚKZÚZ, Czech Republic), Bundesanstalt für
390 Geowissenschaften und Rohstoffe (Germany), le Réseau de Mesures de la Qualité des
391 Sols (RMQS) de l'Institut National de Recherche pour l'Agriculture, l'Alimentation et
392 l'Environnement (INRAE, France). We acknowledge Koens Oorts to sharing the
393 toxicological Model. The LUCAS topsoil dataset, CORINE land cover and Copernic data
394 were funded by the European Environment Agency (EEA). MODIS data was supported
395 by the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC),
396 USGS/Earth Resources Observation and Science (EROS). We acknowledge Dimitri
397 Rambourg, Raphaël di Chiara, Marwan Fahs, Gerhard Schäfer and Benoit Masson-
398 Bedeau for enable computing time. We gratefully acknowledge Margaret Johnson for

399 enabling significant improvement of the manuscript. Finally, we thank the two
400 anonymous reviewers and Guillaume Drouin for useful comments.

401 REFERENCES

- 402 1. Borkow, G.; Gabbay, J., Copper, an ancient remedy returning to fight microbial,
403 fungal and viral infections. *Curr. Chem. Biol.* **2009**, 3, (3), 272–278.
- 404 2. Commission Implementing regulation (EU), 2018/1981 renewing the approval of
405 the active substances copper compounds, as candidates for substitution, in
406 accordance with regulation (EC) No 1107/2009 of the european parliament and
407 of the council concerning the placing of plant protection products on the market,
408 and amending the annex to commission implementing regulation (EU) No
409 540/2011. *OJEU* **2018**, 61, 16–20.
- 410 3. Lamichhane, J. R.; Osdaghi, E.; Behlau, F.; Köhl, J.; Jones, J. B.; Aubertot, J.-N.,
411 Thirteen decades of antimicrobial copper compounds applied in agriculture. A
412 review. *Agron. Sustain. Dev.* **2018**, 38, (3), 28–45.
- 413 4. Ballabio, C.; Panagos, P.; Lugato, E.; Huang, J.-H.; Orgiazzi, A.; Jones, A.;
414 Fernández-Ugalde, O.; Borrelli, P.; Montanarella, L., Copper distribution in
415 European topsoils: An assessment based on LUCAS soil survey. *Sci. Total*
416 *Environ.* **2018**, 636, 282–298.
- 417 5. Ministry of the Environment Finland *Government decree on the assessment of*
418 *soil contamination and remediation needs*; Finland, 2007.
- 419 6. SCHER Voluntary risk assessment report on copper and its compounds,
420 environmental part; 2009.

- 421 http://echa.europa.eu/chem_data/transit_measures/vrar_en.asp (14th January
422 2021).
- 423 7. OIV *World vitiviniculture situation, OIV statistical report on world vitiviniculture*;
424 Organisation Internationale de la Vigne et du Vin: Paris, France, 2016; p 16.
- 425 8. Nadin, P.; Eurostat, *The use of plant protection products in the European Union*
426 *data 1992–2003*. Office for Official Publications of the European Communities:
427 Luxembourg, 2007.
- 428 9. Flemming, C. A.; Trevors, J. T., Copper toxicity and chemistry in the
429 environment: a review. *Water, Air, Soil Pollut.* **1989**, *44*, (1), 143–158.
- 430 10. Maeder, P.; Fließbach, A.; Dubois, D.; Gunst, L.; Fried, P.; Niggli, U., Soil fertility
431 and biodiversity in organic farming. *Science* **2002**, *296*, (5573), 1694–1697.
- 432 11. Hendgen, M.; Hoppe, B.; Döring, J.; Friedel, M.; Kauer, R.; Frisch, M.; Dahl, A.;
433 Kellner, H., Effects of different management regimes on microbial biodiversity in
434 vineyard soils. *Sci. Rep.* **2018**, *8*, (1), 9393–9406.
- 435 12. McBride, M.; Sauve, S.; Hendershot, W., Solubility control of Cu, Zn, Cd and Pb
436 in contaminated soils. *Eur. J. Soil Sci.* **1997**, *48*, (2), 337–346.
- 437 13. Xue, H.; Sigg, L.; Gächter, R., Transport of Cu, Zn and Cd in a small agricultural
438 catchment. *Water Res.* **2000**, *34*, (9), 2558–2568.
- 439 14. Babcsányi, I.; Imfeld, G.; Granet, M.; Chabaux, F., Copper stable isotopes to
440 trace copper behavior in wetland systems. *Environ. Sci. Technol.* **2014**, *48*, (10),
441 5520–5529.

- 442 15. Roussiez, V.; Probst, A.; Probst, J.-L., Significance of floods in metal dynamics
443 and export in a small agricultural catchment. *J. Hydrol.* **2013**, *499*, 71–81.
- 444 16. Schwarzenbach, R. P.; Egli, T.; Hofstetter, T. B.; von Gunten, U.; Wehrli, B.,
445 Global water pollution and human health. *Annu. Rev. Env. Resour.* **2010**, *35*,
446 109–136.
- 447 17. Strawn, D. G.; Baker, L. L., Speciation of Cu in a contaminated agricultural soil
448 measured by XAFS, mu-XAFS, and mu-XRF. *Environ. Sci. Technol.* **2008**, *42*,
449 (1), 37–42.
- 450 18. Bradl, H. B., Adsorption of heavy metal ions on soils and soils constituents. *J.*
451 *Colloid Interface Sci.* **2004**, *277*, (1), 1–18.
- 452 19. Sposito, G., *The chemistry of soils*. second ed.; Oxford University Press: 2008.
- 453 20. Silver, W. L.; Lugo, A. E.; Keller, M., Soil oxygen availability and biogeochemistry
454 along rainfall and topographic gradients in upland wet tropical forest soils.
455 *Biogeochemistry* **1999**, *44*, (3), 301–328.
- 456 21. Borrelli, P.; Robinson, D. A.; Fleischer, L. R.; Lugato, E.; Ballabio, C.; Alewell, C.;
457 Meusburger, K.; Modugno, S.; Schütt, B.; Ferro, V.; Bagarello, V.; Oost, K. V.;
458 Montanarella, L.; Panagos, P., An assessment of the global impact of 21st
459 century land use change on soil erosion. *Nat. Commun.* **2017**, *8*, (1), 2013.
- 460 22. Reimann, C.; de Caritat, P., New soil composition data for Europe and Australia:
461 Demonstrating comparability, identifying continental-scale processes and

- 462 learning lessons for global geochemical mapping. *Sci. Total Environ.* **2012**, *416*,
463 239–252.
- 464 23. Ander, E. L.; Johnson, C. C.; Cave, M. R.; Palumbo-Roe, B.; Nathanail, C. P.;
465 Lark, R. M., Methodology for the determination of normal background
466 concentrations of contaminants in English soil. *Sci. Total Environ.* **2013**, *454–*
467 *455*, 604–618.
- 468 24. Panagos, P.; Ballabio, C.; Lugato, E.; Jones, A.; Borrelli, P.; Scarpa, S.; Orgiazzi,
469 A.; Montanarella, L., Potential sources of anthropogenic copper inputs to
470 european agricultural soils. *Sustainability* **2018**, *10*, 2380–2397.
- 471 25. Chopin, E. I. B.; Alloway, B. J., Distribution and mobility of trace elements in soils
472 and vegetation around the mining and smelting areas of Tharsis, Ríotinto and
473 Huelva, Iberian Pyrite Belt, SW Spain. *Water, Air, Soil Pollut.* **2007**, *182*, (1),
474 245–261.
- 475 26. Wilson, B.; Pyatt, F. B., Heavy metal dispersion, persistence, and bioaccumulation
476 around an ancient copper mine situated in Anglesey, UK. *Ecotoxicol. Environ.*
477 *Saf.* **2007**, *66*, (2), 224–231.
- 478 27. Orchard, V. A.; Cook, F. J., Relationship between soil respiration and soil
479 moisture. *Soil Biol. Biochem.* **1983**, *15*, (4), 447–453.
- 480 28. Borch, T.; Kretzschmar, R.; Kappler, A.; Cappellen, P. V.; Ginder-Vogel, M.;
481 Voegelin, A.; Campbell, K., Biogeochemical redox processes and their impact on
482 contaminant dynamics. *Environ. Sci. Technol.* **2010**, *44*, (1), 15–23.

- 483 29. Donner, E.; Howard, D. L.; Jonge, M. D. d.; Paterson, D.; Cheah, M. H.; Naidu,
484 R.; Lombi, E., X-ray absorption and micro X-ray fluorescence spectroscopy
485 investigation of copper and zinc speciation in biosolids. *Environ. Sci. Technol.*
486 **2011**, *45*, (17), 7249–7257.
- 487 30. Besnard, E.; Chenu, C.; Robert, M., Influence of organic amendments on copper
488 distribution among particle-size and density fractions in Champagne vineyard
489 soils. *Environ. Pollut.* **2001**, *112*, (3), 329–337.
- 490 31. Komárek, M.; Čadková, E.; Chrastný, V.; Bordas, F.; Bollinger, J.-C.,
491 Contamination of vineyard soils with fungicides: A review of environmental and
492 toxicological aspects. *Environ. Int.* **2010**, *36*, (1), 138–151.
- 493 32. Mora, C.; Caldwell, I. R.; Caldwell, J. M.; Fisher, M. R.; Genco, B. M.; Running,
494 S. W., Suitable days for plant growth disappear under projected climate change:
495 potential human and biotic vulnerability. *PLoS Biol.* **2015**, *13*, (6), e1002167.
- 496 33. Brun, L. A.; Maillet, J.; Richarte, J.; Herrmann, P.; Remy, J. C., Relationships
497 between extractable copper, soil properties and copper uptake by wild plants in
498 vineyard soils. *Environ. Pollut.* **1998**, *102*, (2–3), 151–161.
- 499 34. Miotto, A.; Ceretta, C. A.; Brunetto, G.; Nicoloso, F. T.; Giroto, E.; Farias, J. G.;
500 Tiecher, T. L.; De Conti, L.; Trentin, G., Copper uptake, accumulation and
501 physiological changes in adult grapevines in response to excess copper in soil.
502 *Plant Soil* **2014**, *374*, (1), 593–610.

- 503 35. Durán Zuazo, V. H.; Rodríguez Pleguezuelo, C. R., Soil-erosion and runoff
504 prevention by plant covers. A review. *Agron. Sustain. Dev.* **2008**, *28*, (1), 65–86.
- 505 36. Schmidt, M. W. I.; Torn, M. S.; Abiven, S.; Dittmar, T.; Guggenberger, G.;
506 Janssens, I. A.; Kleber, M.; Kogel-Knabner, I.; Lehmann, J.; Manning, D. A. C.;
507 Nannipieri, P.; Rasse, D. P.; Weiner, S.; Trumbore, S. E., Persistence of soil
508 organic matter as an ecosystem property. *Nature* **2011**, *478*, (7367), 49–56.
- 509 37. Araujo, M. B.; Guisan, A., Five (or so) challenges for species distribution
510 modelling. *J. Biogeogr.* **2006**, *33*, (10), 1677–1688.
- 511 38. Press, W. H.; Teukolsky, S. A.; Vetterling, W. T.; Flannery, B. P., *Numerical*
512 *recipes in C: The art of scientific computing*. Cambridge University Press: 2002.
- 513 39. Dietterich, T. G., Ensemble methods in machine learning. In *Proceedings of the*
514 *first international workshop on multiple classifier systems*, Springer-Verlag: New
515 York, NY, 2000.
- 516 40. Hengl, T.; Mendes de Jesus, J.; Heuvelink, G. B. M.; Ruiperez Gonzalez, M.;
517 Kilibarda, M.; Blagotić, A.; Shangguan, W.; Wright, M. N.; Geng, X.; Bauer-
518 Marschallinger, B.; Guevara, M. A.; Vargas, R.; MacMillan, R. A.; Batjes, N. H.;
519 Leenaars, J. G. B.; Ribeiro, E.; Wheeler, I.; Mantel, S.; Kempen, B.,
520 SoilGrids250m: Global gridded soil information based on machine learning. *PLoS*
521 *One* **2017**, *12*, (2), e0169748.
- 522 41. Mackie, K. A.; Müller, T.; Kandeler, E., Remediation of copper in vineyards – A
523 mini review. *Environ. Pollut.* **2012**, *167*, 16–26.

- 524 42. Speiser, B.; Mieves, E.; Tamm, L., Utilisation de cuivre par les paysans bio
525 suisses dans différentes cultures. *Rech. Agr. Suisse* **2015**, 6, (4), 160–165.
- 526 43. Liptzin, D.; Silver, W. L.; Detto, M., Temporal dynamics in soil oxygen and
527 greenhouse gases in two humid tropical forests. *Ecosystems* **2011**, 14, (2), 171–
528 182.
- 529 44. Hofacker, A. F.; Behrens, S.; Voegelin, A.; Kaegi, R.; Losekann-Behrens, T.;
530 Kappler, A.; Kretzschmar, R., Clostridium species as metallic copper-forming
531 bacteria in soil under reducing conditions. *Geomicrobiol. J.* **2015**, 32, (2), 130–
532 139.
- 533 45. Weber, S. D.; Hofmann, A.; Pilhofer, M.; Wanner, G.; Agerer, R.; Ludwig, W.;
534 Schleifer, K. H.; Fried, J., The diversity of fungi in aerobic sewage granules
535 assessed by 18S rRNA gene and ITS sequence analyses. *FEMS Microbiol. Ecol.*
536 **2009**, 68, (2), 246–254.
- 537 46. Fulda, B.; Voegelin, A.; Maurer, F.; Christl, I.; Kretzschmar, R., Copper redox
538 transformation and complexation by reduced and oxidized soil humic acid. 1. X-
539 ray absorption spectroscopy study. *Environ. Sci. Technol.* **2013**, 47, (19), 10903–
540 10911.
- 541 47. Ponizovsky, A. A.; Allen, H. E.; Ackerman, A. J., Copper activity in soil solutions
542 of calcareous soils. *Environ. Pollut.* **2007**, 145, (1), 1–6.
- 543 48. Patrick, R. A. D.; Mosselmanns, J. F. W.; Charnock, J. M.; England, K. E. R.;
544 Helz, G. R.; Garner, C. D.; Vaughan, D. J., The structure of amorphous copper


- 545 sulfide precipitates: An X-ray absorption study. *Geochim. Cosmochim. Acta*
546 **1997**, *61*, (10), 2023–2036.
- 547 49. Jarvis, N. J., A review of non-equilibrium water flow and solute transport in soil
548 macropores: principles, controlling factors and consequences for water quality.
549 *Eur. J. Soil Sci.* **2007**, *58*, (3), 523–546.
- 550 50. Katuwal, S.; Arthur, E.; Tuller, M.; Moldrup, P.; de Jonge, L. W., Quantification of
551 soil pore network complexity with X-ray computed tomography and gas transport
552 measurements soil physics & hydrology. *Soil Sci. Soc. Am. J.* **2015**, *79*, (6),
553 1577–1589.
- 554 51. Manceau, A.; Matynia, A., The nature of Cu bonding to natural organic matter.
555 *Geochim. Cosmochim. Acta* **2010**, *74*, (9), 2556–2580.
- 556 52. European commission CO-FREE: Innovative strategies for copper-free low input
557 and organic farming systems; 2016. <https://cordis.europa.eu/project/id/289497>
558 (17 December 2019).
- 559 53. ECHA, Characterisation of dose [concentration]-response for environment. In
560 *Guidance on Information Requirements and Chemical Safety Assessment*,
561 European Chemicals Agency: Helsinki, Finland, 2008.
- 562 54. Smolders, E.; Oorts, K.; Van Sprang, P.; Schoeters, I.; Janssen, C. R.; McGrath,
563 S. P.; McLaughlin, M. J., Toxicity of trace metals in soil as affected by soil type
564 and aging after contamination: Using calibrated bioavailability models to set
565 ecological soil standards. *Environ. Toxicol. Chem.* **2009**, *28*, (8), 1633–1642.

- 566 55. Panagos, P.; Ballabio, C.; Meusburger, K.; Spinoni, J.; Alewell, C.; Borrelli, P.,
567 Towards estimates of future rainfall erosivity in Europe based on REDES and
568 WorldClim datasets. *J. Hydrol.* **2017**, *548*, 251–262.
- 569 56. Gottschalk, P.; Smith, J. U.; Wattenbach, M.; Bellarby, J.; Stehfest, E.; Arnell, N.;
570 Osborn, T. J.; Jones, C.; Smith, P., How will organic carbon stocks in mineral
571 soils evolve under future climate? Global projections using RothC for a range of
572 climate change scenarios. *Biogeosciences* **2012**, *9*, (8), 3151–3171.
- 573 57. Tack, F. M. G.; Meers, E., Assisted phytoextractions helping plants to help us.
574 *Elements* **2010**, *6*, 383–388.
- 575 58. Schmidt, U., Enhancing phytoextraction: The effect of chemical soil manipulation
576 on mobility, plant accumulation, and leaching of heavy metals. *J. Environ. Qual.*
577 **2003**, *32*, (6), 1939–1954.
- 578 59. Thomas, C. S.; Gubler, W. D.; Leavitt, G., Field testing of a powdery mildew
579 disease forecast model on grapes in California. *Phytopathology* **1994**, *84*, 1070.
- 580 60. Pedneault, K.; Provost, C., Fungus resistant grape varieties as a suitable
581 alternative for organic wine production: Benefits, limits, and challenges. *Sci.*
582 *Hortic.* **2016**, *208*, 57–77.
- 583 61. Ongena, M.; Jacques, P., Bacillus lipopeptides: versatile weapons for plant
584 disease biocontrol. *Trends Microbiol.* **2008**, *16*, (3), 115–125.

- 585 62. Harm, A.; Kassemeyer, H.-H.; Seibicke, T.; Regner, F., Evaluation of chemical
586 and natural resistance inducers against Downy Mildew (*Plasmopara viticola*) in
587 grapevine. *Am. J. Enol. Viticult.* **2011**, *62*, (2), 184–192.
- 588 63. Abdu-Allah, G. A. M.; Abo-Elyousr, K. A. M., Effect of certain plant extracts and
589 fungicides against powdery mildew disease of grapevines in upper Egypt. *Arch.*
590 *Phytopathol. Plant Protect.* **2017**, *50*, (19–20), 957–969.
- 591 64. Soustre-Gacougnolle, I.; Lollier, M.; Schmitt, C.; Perrin, M.; Buvens, E.;
592 Lallemand, J. F.; Mermet, M.; Henaux, M.; Thibault-Carpentier, C.; Dembele, D.;
593 Steyer, D.; Clayeux, C.; Moneyron, A.; Masson, J. E., Responses to climatic and
594 pathogen threats differ in biodynamic and conventional vines. *Sci. Rep.* **2018**, *8*,
595 16857–16871.
- 596 65. Karimi, B.; Cahurel, J.-Y.; Gontier, L.; Charlier, L.; Chovelon, M.; Mahé, H.;
597 Ranjard, L., A meta-analysis of the ecotoxicological impact of viticultural
598 practices on soil biodiversity. *Environ. Chem. Lett.* **2020**, *18*, (6), 1947–1966.
- 599 66. Andreadis, K. M.; Schumann, G. J.-P.; Pavelsky, T., A simple global river bankfull
600 width and depth database. *Water Resour. Res.* **2013**, *49*, (10), 7164–7168.

601

602 **FOR TOC ONLY**



603