



**HAL**  
open science

## **Lysyl oxidase is a strong determinant of tumour cell colonization in bone**

Caroline E Reynaud, Laura Ferreras, Paola Di Mauro, Casina Kan, Martine Croset, Edith Bonnelye, Floriane Pez, Clémence Thomas, Géraldine Aimond, Antoine E Karnoub, et al.

► **To cite this version:**

Caroline E Reynaud, Laura Ferreras, Paola Di Mauro, Casina Kan, Martine Croset, et al.. Lysyl oxidase is a strong determinant of tumour cell colonization in bone. *Cancer Research*, 2017, 77 (2), pp.268-278. 10.1158/0008-5472.CAN-15-2621 . hal-03380388

**HAL Id: hal-03380388**

**<https://hal.science/hal-03380388v1>**

Submitted on 15 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Lysyl oxidase is a strong determinant of tumour cell colonization in bone.**

Caroline Reynaud<sup>1,2</sup>, Laura Ferreras<sup>1,2</sup>, Paola Di Mauro<sup>1,2</sup>, Casina Kan<sup>1,2</sup>, Martine Croset<sup>1,2</sup>, Edith Bonnelye<sup>1,2</sup>, Floriane Pez<sup>3</sup>, Clémence Thomas<sup>4</sup>, Géraldine Aimond<sup>3</sup>, Antoine E. Karnoub<sup>4</sup>, Marie Brevet<sup>1,5</sup>, Philippe Clézardin<sup>1,2</sup>.

<sup>1</sup>INSERM, UMR1033, F-69372 Lyon, France

<sup>2</sup> University of Lyon, F-69622 Villeurbanne, France

<sup>3</sup> CNRS, UMR5305, IBCP, F-69367 Lyon, France

<sup>4</sup> Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA

<sup>5</sup> Department of Pathology, Groupement Hospitalier Est, Hospices Civils de Lyon, Lyon, France

**Short Title:** LOX triggers tumour cell colonization in bone

**Keywords:** metastasis; IL-6; osteoclast; osteoblast.

**Disclosure of potential conflict of interests:** All authors state that they have no conflicts of interest.

**Financial support:** CR acknowledges the support of the Ligue Contre le Cancer (Comité du Rhône) and CNRS. PC is supported by INSERM, the University of Lyon, the Grand Prix de la Recherche Ruban Rose and INCa (grant n°: 2014-164). CK is a recipient of a Marie Curie individual fellowship from the Horizon 2020 European Programme under agreement number n°655777. Additionally, this work was supported by the LabEX

DEVweCAN (ANR-10-LABX-61) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

**Corresponding authors:**

P. Clézardin, C. Reynaud, INSERM, UMR1033, Faculté de Médecine Lyon-Est (domaine Laennec), Rue Guillaume Paradin, 69372 Lyon cedex 08, France, Tel: 33 4 78 78 57 38;

Fax: 33 4 78 77 87 72; emails: [philippe.clezardin@inserm.fr](mailto:philippe.clezardin@inserm.fr);

[caroline.reynaud@inserm.fr](mailto:caroline.reynaud@inserm.fr)

## **Abstract**

Lysyl oxidase (LOX) is a secreted copper-dependent amine oxidase whose primary function is to drive collagen crosslinking and extracellular matrix stiffness. LOX in colorectal cancer (CRC) synergizes with hypoxia-inducible factor-1 (HIF-1 $\alpha$ ) to promote tumour progression. Here we investigated whether LOX/HIF1 endows CRC cells with full competence for aggressive colonization in bone. We show that a high *LOX* expression in primary tumours from CRC patients was associated with poor clinical outcome, irrespective of *HIF-1*. Additionally, LOX was expressed by tumour cells in the bone marrow from CRC patients with bone metastases. *In vivo* experimental studies show that *LOX* overexpression in CRC cells or systemic delivery of the conditioned medium from *LOX*-overexpressing CRC cells promoted tumour cell dissemination in the bone marrow and enhanced osteolytic lesion formation, irrespective of *HIF-1*. Conversely, silencing or pharmacological inhibition of LOX activity blocked dissemination of CRC cells in the bone marrow and tumour-driven osteolytic lesion formation. *In vitro*, tumour-secreted LOX supported the attachment and survival of CRC cells to and in the bone matrix, and inhibited osteoblast differentiation. *LOX* overexpression in CRC cells also induced a robust production of IL-6. In turn, both LOX and IL-6 were acting in concert to stimulate osteoclast differentiation, thereby creating an imbalance between bone resorption and bone formation. Collectively, our findings show that LOX supports CRC cell dissemination in the bone marrow and they reveal a novel mechanism through which LOX-driven IL-6 production by CRC cells impairs bone homeostasis.

*Word count (limit.: 250): 239*

## Introduction

The lysyl oxidase (LOX) family of secreted copper-dependent amine oxidases consists of five paralogs: LOX and LOX-like 1-4 (LOXL 1-4). The primary function of the LOX family is to catalyse the covalent crosslinking of collagen and elastin in the extracellular matrix, thereby increasing insoluble matrix deposition and tensile strength (1). Increased expression of *LOX* and *LOXL2* has been consistently reported in various cancer types (colorectal, breast, prostate, lung, bladder) (1-5). LOX and LOXL2 are closely associated with desmoplastic areas at the invasive front of infiltrating tumours (1). They mediate collagen/elastin crosslinking that increases extracellular matrix stiffness, a process which is associated with enhanced integrin-mediated mechanotransduction coupled to increased tumour cell invasion in breast and colorectal cancers (1,3,6). LOX- and LOXL2-mediated collagen crosslinking are also responsible for enhancing outgrowth in metastatic xenograft models of breast cancer (5,7). Additionally, LOX enhances the metastatic trait of breast tumour xenografts in animals by stimulating *TWIST1* transcription in tumour cells (8). In the same vein, hypoxia-inducible factor-1 (HIF-1) induces the expression of several members of the LOX family in breast cancer cells, including *LOX* and *LOXL2*, which then catalyses collagen crosslinking in the lungs, facilitating the recruitment of bone marrow-derived cells and the subsequent colonization of the pulmonary tissue by tumour cells (5, 9-11). Furthermore, we have shown that LOX synergises with HIF-1 in promoting *in vivo* growth of colorectal tumours (2). These experimental results (2, 3, 5-11) probably explain why increased expression of *LOX* and *LOXL2* in primary tumours (colorectal, breast, lung, prostate) is associated with distant relapse and poor survival (1). Indeed, targeting of LOX and LOXL2 with the LOX inhibitor  $\beta$ -aminopropionitrile ( $\beta$ APN) or a function-blocking antibody (AB0023), respectively, is efficacious at reducing metastatic tumour burden in xenograft models of

cancer (7,12). Thus, there is a body of experimental evidence indicating that LOX and LOXL2 facilitate the development of metastases in distant organs, such as the lungs, liver, and brain. Surprisingly, little is known regarding the role of LOX proteins during bone metastasis formation.

In bone metastasis, metastatic cancer cells residing in the bone marrow alter the functions of bone-resorbing (osteoclasts) and bone-forming (osteoblasts) cells and hijack signals coming from the bone matrix (13). By disrupting the physiological balance between bone resorption and bone formation, metastatic cells therefore promote skeletal destruction. The detection rate of disseminated tumour cells (DTC) in the bone marrow from colorectal cancer (CRC) patients receiving curatively intended surgery is 27% and their presence is associated with a poor clinical outcome (14,15). This detection rate is relatively high and comparable to what is observed in bone-tropic cancers such as breast cancer (about 30%) (14), which is in intriguing contrast to the low incidence of overt bone metastases in CRC (16). A current hypothesis gaining ground is that DTCs may be representative of dormant tumour cells with the ability to escape dormancy upon receiving appropriate stimuli from the microenvironment (17). Given the role of LOX in driving collagen crosslinking and extracellular matrix stiffness (1), we hypothesized that LOX in CRC may be one of the factors that endow DTCs with full competence for aggressive colonization in bone. Interestingly, it has been very recently shown during the course of this study that hypoxia-induced LOX in breast cancer cells generates premetastatic osteolytic lesions in animals through the stimulation of NFATc1, a master regulator of osteoclastogenesis (18). Here, we investigated the role of LOX and its regulator HIF-1 $\alpha$  in CRC bone metastases.

## Materials and Methods

**Reagents, human cancer cell lines, and animals.** All chemicals were obtained from Sigma-Aldrich, unless otherwise specified. Primer pairs used for quantitative real-time PCR analysis are listed in Supplementary Table S1. Human Hct116, HT29 and LS174Tr CRC cell lines (ATCC) were tested before their usage for authentication by DNA fingerprinting using short tandem repeat (STR) method in 2010. The mutation in the *ras* proto-oncogene was also tested in 2014. Hct116 cells were maintained in RPMI-1640-Glutamax (Invitrogen) with 10% (v/v) FBS, 10mM HEPES, 1mM sodium pyruvate, and 0.1mM non essential amino acids (Invitrogen) in humidified atmosphere at 37 °C. HT29 and LS174Tr cells were grown in DMEM-glutamax (Invitrogen), 10% (v/v) FBS and 50µg/ml gentamycin. These cells were first transduced with lentiviral shRNA particles targeting *LOX* (shLOX) or a control sequence (shCTL) then co-transduced with particles containing an empty vector (EV) or a *LOX*-expressing vector (LOX+) under the control of a doxycycline (dox)-inducible promoter, as previously described (2). Transduced cell lines initially called shCTL/EV, shLOX/EV (cells silenced for endogenous *LOX*) and shCTL/LOX+ (cells overexpressing *LOX* in an inducible manner) (2) were named, respectively, Ctrl, LOX- and LOX+ for brevity. Additionally, Ctrl and LOX+ Hct116 cells were co-transduced with lentiviral shRNA particles targeting *HIF-1a* (Ctrl/HIF1 $\alpha$ - and LOX+/HIF1 $\alpha$ -, respectively), as previously described (2).

Four-week-old female Balb/c immunocompromised mice were purchased from Janvier (Saint-Berthevin, France). Animals were maintained in a 12-h light-dark cycle and given free access to food and water. All procedures involving animals, including their housing and care, the method by which they were culled, and experimental protocols were conducted in accordance with a code of practice established by the local ethical committee of the University of Lyon.

**Immunohistochemistry.** CRC tissue specimens (primary tumors and bone metastases) were obtained from the Department of Pathology, Hospices Civils de Lyon. Tumour sections (5µm) were de-paraffinized, treated with pepsin for 15-min at 37°C (Zymed, Invitrogen® Carlsbad USA), and processed for immunohistochemical staining using a LOX antibody (1:250 dilution; ab31238 Abcam).

**LOX activity measurement.** LOX activity was measured using a fluorescent assay, according to manufacturer's instructions (ab112139 Abcam).

**ELISA.** Conditioned media from LOX- and LOX+#Hct116 were collected, and IL-6 was measured by ELISA, according to manufacturer's instructions (Clinisciences).

**Protein extraction and Western blotting.** Proteins were extracted in RIPA buffer (Sigma), electrophoresed on a SDS-polyacrylamide gel (Life technologies), then transferred onto nitrocellulose membranes (Millipore, Billerica, MA) and proteins were probed with a primary antibody against STAT3 (HP1001671, 1:300 dilution, Sigma), phosphoSTAT3 (9145, 1:1,000 dilution, Cell Signaling), Src (clone GD11, 1:1,000 dilution, Millipore), Src<sup>Y416</sup> (Clone 916, 1:1,000 dilution, Millipore), FAK (05-537, 1:1,000 dilution, Upstate), phosphoFAK (44-624, 1:1,000 dilution, Biosource), β-actin (A5060, 1:15,000 dilution, Sigma) or tubulin (T5168, 1:2,000 dilution, Sigma). After incubation with primary antibodies, membranes were incubated with horseradish peroxidase (HRP)-conjugated secondary goat anti-mouse (1:2,000 dilution, BioRad) or goat anti-rabbit (1:2,500 dilution, BioRad) antibody. Immunostaining was performed with enhanced chemiluminescence (ECL) detection system (Perkin Elmer).



**Functional cell-based assays.** Ctrl, LOX- and LOX+ Hct116 cells ( $4 \times 10^6$ /ml), treated or not treated with  $350 \mu\text{M}$   $\beta$ APN or  $100 \mu\text{g/ml}$  tocilizumab, were cultured for 48h in complete RPMI medium without phenol red. Conditioned media were then collected, centrifuged and stored as aliquots at  $-80^\circ\text{C}$  until used. LOX production was analyzed by western blotting, as previously described (2).

For cell adhesion assays, experiments were conducted as previously described (19). Briefly, 96-well tissue culture plates were coated with increasing concentrations of type-I collagen or fibronectin ( $0.01 \text{mg/cm}^2$  to  $100 \text{mg/cm}^2$ ) and incubated overnight at  $4^\circ\text{C}$ . Cells ( $4 \times 10^4$  cells/0.1 ml/well) were allowed to adhere for 20 min at  $37^\circ\text{C}$  in a 5%  $\text{CO}_2$  atmosphere. After washing to remove non-adherent cells, adherent cells were fixed and stained with 0.1% (w/v) crystal violet. The dye was eluted with 2% (w/v) SDS and the optical density quantified at 595 nm. Alternatively, 96-well tissue culture plates were coated with type-I collagen ( $15 \mu\text{g/cm}^2$ ) overnight at  $4^\circ\text{C}$ . Following incubation with 1% BSA for 30 min, cells ( $1 \times 10^5$  cells/0.1 ml/well) were incubated for 1h at  $37^\circ\text{C}$  in a 5%  $\text{CO}_2$  atmosphere. After washing, adherent cells were fixed and stained with hematoxylin and fuschin.

For tumour spheroid-formation assays,  $1 \times 10^4$ /0.3 ml Hct116 cells (Ctrl, LOX- and LOX+) were seeded in CytoCapture Chambers, Big Hexagonal Cavities (diameter  $250 \mu\text{m}$ ) (PAA). Cells were grown in suspension in 0.3mL of MammoCult™ basal medium containing 10% (v/v) MammoCult™ proliferation supplement (Stem Cell), 0.0004% (w/v) heparin (Stem Cell),  $1 \mu\text{g/ml}$  hydrocortisone, 1mM glutamine, 50U/mL penicillin, 50 $\mu\text{g/ml}$  streptomycin (Life Technologies), with or without  $350 \mu\text{M}$   $\beta$  APN. Tumour spheroids were imaged under a microscope (Confocal-Leica SP5 X) and quantified using Fiji software. Spheroids of at least  $60 \mu\text{m}$  in diameter were counted.

**Osteoclastogenesis assay.** Experiments were conducted as described previously (20). Briefly, bone marrow cells from 6-week-old OF1 mice were cultured for 7 days in  $\alpha$ -MEM medium (Invitrogen) supplemented with 10% (v/v) fetal calf serum (Invitrogen), 20 ng/ml M-CSF (R&D Systems) and 200 ng/ml RANKL, alone or in combination (from day 1 to day 7) with the conditioned media from transduced Hct116 cells (Ctrl, LOX+ and LOX-) (25  $\mu$ g/ml) or with recombinant LOX (150 ng/ml). After 7 days in culture, mature osteoclasts were enumerated under a microscope on the basis of the number of nuclei (more than three nuclei) and TRAP activity.

**NFATc1 staining.** Cultured osteoclasts on coverslips were fixed for 15 min in 4% PFA then permeabilized with 0.1% Triton X-100 in PBS for 10 min. Saturation was performed with 5% normal goat serum in 0.1% PBS-Tween20 (NGS) for 2h. Monoclonal antibody to NFATc1 (SC-7294, Santa Cruz), diluted 1:100 in NGS, was incubated 1h at room temperature and, after washing, coverslips were further incubated for another 1h with a secondary goat anti-mouse antibody coupled to Alexa Fluor 488 (1:300 dilution in PBS). Coverslips were then mounted in FluorSave Reagent (Calbiochem).

**Osteoblastogenesis assay.** Experiments were conducted as previously described (21). Briefly, calvaria of 3-day-old OF-1 mice were dissected then cells were enzymatically isolated by sequential digestion with collagenase and plated into 24-well plates. After 24-h incubation,  $\alpha$ -MEM medium containing 10% fetal bovine serum (Life technologies, Carlsbad, CA, USA) was changed and supplemented with 50 $\mu$ g/ml ascorbic acid (Sigma-Aldrich, Buchs, Switzerland) and with or without conditioned medium from transduced Hct116 cells (Ctrl, LOX+ and LOX-). Medium was changed every other day for 21 days. 10mM sodium  $\beta$ -glycerophosphate (Sigma-Aldrich, Buchs, Switzerland) was added for

the last week of the experiments. At day 21, bone mineralized nodules were fixed and stained with von Kossa staining.

**Animal studies.** CRC cell lines ( $5 \times 10^5$  cells in 100  $\mu$ l PBS) were inoculated intra-arterially into anaesthetised female Balb/c *nude* mice. When specified, 300  $\mu$ l of conditioned medium from transduced Hct116 cells were intraperitoneally injected daily into mice. For LOX inhibition, animals were treated with 0.2% (w/v)  $\beta$ APN, supplied in the drinking water. For blockade of IL-6 receptor, animals were treated every other day with tocilizumab (50 mg/kg, i.p.). The progression of skeletal tumour burden was monitored by whole-body bioluminescence imaging (NightOwl, Berthold Technologies), following subcutaneous administration of luciferin (100 mg/kg in PBS; Promega) 10 min prior to imaging. The progression of osteolytic lesions in the skeleton of anaesthetised animals was monitored by radiography, using a cabinet X-ray system (MX-20; Faxitron X-ray Corporation). The area of osteolytic lesions was measured using Explora-Nova Morpho Expert software. Animals were sacrificed on day 7 or 35 after tumour cell inoculation, and hind limbs were collected for histology and histomorphometric analyses.

**Bone histology and histomorphometry.** Bone histology and histomorphometric analysis of bone tissue sections were performed on decalcified, 5- $\mu$ m bone-tissue sections stained with Goldner's Trichrome, as previously described (20). Histomorphometric measurements [bone volume (BV)/tissue volume (TV) and tumour volume (TuV)/soft tissue volume (STV) ratios] were performed in a standard zone of the tibial metaphysis, situated at 0.5 mm from the growth plate, including cortical and trabecular bone. The BV/TV ratio represents the percentage of bone tissue. The TB/STV

ratio represents the percentage of tumour tissue. Additionally, osteoclasts within bone tissue sections were stained using a TRAP activity kit assay (Sigma). The resorption surface (Oc.S/BS) was calculated as the ratio of TRAP-positive trabecular bone surface (Oc.S) to the total bone surface (BS) using the image analysis system MorphoExpert (Explora-nova).

**Ex vivo micrometastasis experiments.** Ex-vivo micrometastasis experiments were conducted as previously described (20). Animals were culled on day 7 or day 35 after tumor cell inoculation. Hind limbs were collected and tibiae and femurs were minced then soaked in an enzyme cocktail containing 300U/ml type-I collagenase and 100 U/ml hyaluronidase (StemCell Technologies) in DMEM medium for 2 hours at 37°C. After incubation, bone marrow cell suspensions were seeded in 6-well plates and cultured in complete medium. After 1-day, cultured cells were placed under puromycin selection for 2 weeks, enabling the selective outgrowth of antibiotic-resistant tumour cells. Tumour cell colonies were then fixed, stained with 0.5% (v/v) crystal violet and counted.

**Clinical correlation analyses.** Gene expression data and clinical annotations were downloaded from The Cancer Genome Atlas for colorectal cancer and previously published datasets downloaded from the Gene Expression Omnibus (GSE16125, GSE33113, GSE41258, GSE17536, GSE31595, GSE12945) (see Supplementary Table S2).

**Statistical Analysis.** All experimental data are presented as mean values  $\pm$  SD. Statistical comparisons of values were made using the Mann-Whitney *U*-test. Disease-free survival Kaplan Meier analyses were performed using the log-rank (Mantel-Cox) test. Correlation analyses were performed by the Spearman Rank test and the Pearson

correlation. All tests were two-sided, and  $P$  values less than 0.05 were considered statistically significant.

## Results

### **LOX expression in colorectal cancer is clinically associated with poor prognosis.**

As a first step towards evaluating the role of LOX and its regulator HIF-1 $\alpha$  in CRC bone metastases, we conducted a meta-analysis in a cohort of CRC patients ( $n = 552$ ) and found that high *LOX* expression in primary tumours was associated with poor overall survival ( $p = 0.0109$ ) and poor relapse-free survival ( $p = 0.02$ ) (Supplementary Figs. S1a and S1b). Additionally, high *LOX* expression was a negative determinant of relapse-free survival, irrespective of *HIF-1 $\alpha$*  levels (Supplementary Fig. S1c). Using immunohistochemistry with a anti-LOX antibody, we examined 5 CRC patients with bone metastases, including 3 patients for whom we had pairs of primary tumours and their matching bone metastases. Although all of the matching primary and metastatic tumours expressed LOX, there was a preferential moderate-to-strong staining for LOX associated with the carcinoma cells in bone metastases for all 5 patients (Fig. 1 and Supplementary Fig. S1d).

### **Tumour-secreted LOX in colorectal cancer generates osteolytic lesions in animals.**

We next investigated the role for LOX and HIF-1 $\alpha$  in CRC bone metastasis formation, using human Hct116 cells (Ctrl) previously transduced for HIF-1 $\alpha$  silencing (Ctrl/HIF-1 $\alpha$ -) and/or *LOX* overexpression (LOX+ ; LOX+/HIF-1 $\alpha$ -) (2). Compared to Ctrl and Ctrl/HIF-1 $\alpha$ - tumour-bearing animals, bioluminescent LOX+ Hct116 cells were readily detected in the hind limbs of animals at day 31-post injection, irrespective of

*HIF-1* $\alpha$  expression (Fig. 2a). Additionally, there was a similar effect on osteolytic lesions in LOX+ and LOX+/HIF-1 $\alpha$ - tumour-bearing animals and the extent of lytic lesions was 2.5-fold higher than that of animals bearing Ctrl- or Ctrl/HIF-1 $\alpha$ - tumors (Fig. 2b). Hence, these data indicated that tumour-secreted LOX promotes osteolytic lesion formation *in vivo*, irrespective of *HIF-1* $\alpha$  expression.

To further address the role of LOX in osteolytic lesion formation, we used two additional previously published human CRC cell lines (HT29 and LS174Tr) in which *LOX* expression has been overexpressed or silenced (2). The modulation of *LOX* expression did not affect the expression of other members of the LOX-like family (Supplementary Table S3). We observed that *LOX*-overexpressing Hc116, HT29 and LS174Tr cells caused osteolytic lesions in animals at day 35-post injection (Fig. 2c). The extent of osteolytic lesions in LOX+ tumour-bearing animals was 3- to 5-fold higher than that of animals bearing Ctrl or LOX- tumours (Fig. 2c). Additionally, the treatment of LOX+ Hct116 tumour-bearing animals with the LOX inhibitor  $\beta$ APN for 35 days, prolonged bone metastasis-free survival to a level similar to that observed with animals bearing LOX- or Ctrl-Hc116 tumours (Supplementary Fig. S2). Histomorphometric analysis of hind limbs with metastases from animals bearing Hct116 tumours showed that the BV/TV ratio (a measure of the bone volume) was decreased in LOX+ tumour-bearing animals, compared with LOX- and Ctrl tumour-bearing animals (Fig. 2d and Table 1). This difference was accompanied with a substantial increase in the TRAP staining of bone tissue sections of metastatic legs from LOX+ tumour-bearing animals (indicating a stimulation of active-osteoclast resorption surfaces) (Fig. 2d and Table 1). Additionally, there was a dramatic increase in the TuV/STV ratio (a measure of the skeletal tumour burden) in LOX+ tumour-bearing animals, compared with animals bearing Ctrl- and LOX- tumours (Fig. 2d and Table 1).

To further examine LOX-dependency, *nude* mice were treated with the conditioned medium from LOX+ or LOX- Hct116 cells beginning 1 day (D-1) before intra-arterial inoculation of parental Hct116 cells (D0). The daily treatment with the conditioned medium then continued until day 35 (D35), at which time anaesthetised animals were analysed by radiography (Fig. 3a). Radiographs of animals showed that mice treated with the LOX+ conditioned medium had a 6-fold increase in the extent of osteolytic lesions, compared to that observed with the LOX- medium (Figure 3b). The bone marrow of 3 metastatic mice per group was then collected and placed in culture under antibiotic selection, enabling the selective outgrowth of antibiotic-resistant tumour cells (Fig. 3c). The average number of colonies recovered from the bone marrow of mice treated with the LOX+ conditioned medium was 2.5-fold higher than that recovered from animals treated with the LOX- conditioned medium ( $344 \pm 60$  vs  $121 \pm 25$  colonies/well) ( $P = 0.08$ ) (Fig. 3c). Hence, LOX plays a prominent role in supporting osteolytic lesion formation *in vivo* and tumour cell dissemination *ex vivo*.

### **LOX disrupts the balance between bone resorption and bone formation.**

By disrupting the physiological balance between bone resorption and bone formation, tumour cells promote skeletal destruction (13). To directly test whether *LOX* expression in Hct116 cells could influence osteoclast differentiation, we treated primary mouse bone marrow cell cultures with RANKL and M-CSF together with the conditioned medium from Ctrl, LOX- and LOX+ Hct116 cells. Consistent with *in vivo* data (Fig. 2d), the conditioned medium from LOX+ Hct116 cells stimulated the formation of TRAP-positive multinucleated osteoclasts, compared to conditioned media from Ctrl- and LOX- Hct116 cells (Fig. 4a). As aforementioned,  $\beta$ APN treatment of LOX+ tumour-bearing animals inhibited osteolytic lesion formation (Supplementary Fig. S2), suggesting that LOX-

mediated bone destruction critically depends on LOX's enzymatic activity. In agreement with this suggestion  $\beta$ APN almost completely inhibited LOX enzymatic activity in the conditioned medium from LOX+ Hct116 cells, when compared to that observed with the conditioned medium from LOX- Hct116 cells (Supplementary Fig. S3). Additionally,  $\beta$ APN treatment blocked the stimulatory effect of LOX+ on osteoclastogenesis (Fig. 4a), indicating that LOX-induced osteoclastogenesis critically depends on LOX's enzymatic activity.

*LOX* overexpression induced a robust production of IL-6 by Hct116 cells, whose production was almost totally inhibited by  $\beta$ APN treatment of LOX+ Hct116 cells (Fig. 4b). Tumour-derived IL-6 can act as an autocrine agent and activate the signal transducer and activator of transcription 3 (STAT3) signalling pathway in CRC cells (22,23). In agreement with these findings (22, 23), STAT3 phosphorylation levels were substantially increased in LOX+ Hct116 cells, compared to Ctrl- and LOX- Hct116 cells (Fig. 4c). Additionally, blockade of the IL-6 receptor by tocilizumab abrogated STAT3 phosphorylation in LOX+ Hct116 cells, compared with untreated cells (Fig. 4c). By contrast, the FAK (focal adhesion kinase)/Src tyrosine kinase activities in Hct116 cells were the same whether these cells expressed or did not express *LOX* (Supplementary Fig. S4). Thus, our data revealed a previously unreported functional link between LOX and IL-6, in which LOX-driven IL-6 production by CRC cells promotes IL-6R/STAT3 signalling pathway. This link extended to the clinic, because *LOX* and *IL-6* significantly correlated across 7 publically available databases on CRC, when analysed independently or in combination (Fig. 4d and Supplementary Fig. S5). The addition of IL-6 to the conditioned medium of LOX- Hct116 cells stimulated osteoclast formation to an extent similar to that observed with the conditioned medium from LOX+ Hct116 cells (Fig. 4e). Moreover, tocilizumab inhibited osteoclastogenesis induced by the LOX+ conditioned



medium, and an additive inhibitory effect on osteoclast formation was observed when combining tocilizumab and  $\beta$ APN (Fig. 4f). These results (Fig. 4e-f) suggested that both IL-6 and LOX were effective stimulators of osteoclastogenesis. Indeed, in agreement with previous findings (18), we observed that recombinant LOX (in the presence of RANKL+M-CSF) stimulated the formation of osteoclasts *in vitro*, compared to control (RANKL+M-CSF) (Fig. 5a). Additionally, it promoted a greater nuclear localization of NFATc1, the master regulator of osteoclastogenesis, than RANKL+MCSF alone (85 vs 30% of positive nuclear staining;  $P < 0.01$ ) (Fig. 5b). Hence, tumour-secreted LOX and IL-6 were acting in concert to stimulate the generation of differentiated osteoclasts.

To determine whether *LOX* expression in Hct116 cells could also influence osteoblast differentiation, we treated primary calvarial mouse osteoblasts with the conditioned medium from Ctrl, LOX+ or LOX- Hct116 cells. We found that the conditioned medium from LOX+ Hct116 cells partially inhibited the formation of bone nodules and their mineralization, compared to conditioned media from Ctrl and LOX- Hct116 cells (Fig. 5c-d).

Collectively, our findings suggested that tumour-derived LOX induces an imbalance between bone resorption and bone formation, which leads to bone destruction *in vivo*.

### **LOX primes tumour cells for dissemination to the bone marrow**

To determine the role of LOX in the settlement of tumour cells in the bone marrow, animals injected with parental Hct116 cells were treated daily from D-1 to D7 with the conditioned medium from LOX- or LOX+ Hct116 cells, with or without tocilizumab or  $\beta$ APN treatment. The bone marrow was then collected and placed in culture under antibiotic selection, enabling growth of tumour cells that have disseminated to the bone marrow (Fig. 6a). We recovered colony-forming tumour cells in the bone marrow from

7 out of 10 animals treated with the LOX+ conditioned medium, whereas only 2 out of 8 animals treated with the LOX- conditioned medium had tumour cell colonies in the bone marrow (Fig. 6b). Moreover, the average number of colony-forming tumour cells recovered from the bone marrow of the 7 metastatic mice treated with the LOX+ conditioned medium was 4-fold higher than that recovered from the 2 mice treated with the LOX- conditioned medium (Fig. 6b). The treatment of animals with  $\beta$ APN completely blocked the stimulatory effects of LOX on incidence (1 out of 6 animals) and number of colony-forming tumour cells in the bone marrow (Fig. 6b). In sharp contrast, tocilizumab did not inhibit LOX stimulatory effect on tumour cell dissemination (fig. 6b). Thus, these data indicate that LOX (irrespective of IL-6) is crucially important in the settlement of tumour cells in the bone marrow.

### **LOX promotes tumour cell adhesion and survival**

We have previously reported that *LOX* overexpression in breast cancer cells induces bone metastasis, a phenotype associated with its ability to induce the expression of a major epithelial-to-mesenchymal (EMT)-transcription factor, *TWIST1* (8). Additionally, we have shown that *TWIST1* facilitates breast cancer bone metastasis formation (20). We therefore proceeded to explore whether LOX promoted tumour cell colonization in bone *via* induction of EMT. Hct116 cells expressed detectable levels of genes encoding for EMT-inducing transcription factors, such as *TWIST1* and *SNAI1* (Snail) (Supplementary Fig. S6). *LOX* expression in Hct116 cells did not modify the morphological appearance of these cells and it did not modify *TWIST1*, *SNAI1* or vimentin (*VIM*) expression (supplementary Fig. S6), indicating that LOX in colon cancer cells does not contribute to bone marrow colonization by promoting an EMT.

Early steps of bone metastasis involve the attachment and survival of DTCs to and

in the bone matrix (13), two properties we examined using *in-vitro* assays. Collagens constitute 90% of the total protein content in bone, type-I collagen being the most abundant bone extracellular matrix protein (24). Cell attachment experiments to type-I collagen were therefore performed using Ctrl, LOX- and LOX+ Hct116 cells. As shown in Fig. 6c, there was a significant gain in the attachment of *LOX*-overexpressing tumour cells to collagen, compared to Ctrl Hct116 cells. The attachment of LOX-deficient tumour cells was lower than that observed with Ctrl Hct116 cells (Fig. 6c). In contrast, the extent of attachment of Hct116 tumour cell lines to fibronectin was the same, irrespective of *LOX* expression levels (Fig. 6c). Using Hct116 cells silenced for *LOX* expression (LOX-) as a control to measure baseline cell attachment to type-I collagen, we conducted cell attachment experiments with *LOX*+ Hct116 cells, in the presence or absence of  $\beta$ APN. As shown in Fig. 6c,  $\beta$ APN abrogated the gain of attachment of *LOX*+ Hct116 cells to type-I collagen (*LOX*+  $\beta$ APN), compared to that observed with untreated cells (*LOX*+). Hence, these observations suggested that LOX crosslinking activity is important to mediate CRC cell attachment to type-I collagen.

Once tumour cells seed in the bone marrow, they need to survive in this microenvironment. A relevant *in vitro* feature of CRC cell survival is their ability to grow as colonospheres (25). We thus asked whether Ctrl, LOX-, and LOX+ Hct116 cells could form colonospheres in serum-free medium. Indeed, LOX+ Hct116 cells formed a higher number of colonospheres than the two other Hct116 cell lines, and growth of LOX+ Hct116 cells as colonospheres was significantly inhibited in the presence of  $\beta$ APN (Supplementary Fig. 7 and data not shown). Thus, LOX can provide CRC cells with a survival advantage in the bone marrow.

## **Discussion**

A large body of preclinical and clinical evidence has shown that *LOX* expression in tumours facilitates the progression of several cancers and the development of metastases to distant organs, such as the lungs, liver, and brain (1,2,4,7,9,10). Indeed, *LOX* has been identified as an important regulator of hypoxia-induced tumour progression *via* a HIF-1 $\alpha$ -dependent mechanism in numerous cancer types (1,9-11). Surprisingly, aside from a very recent report that hypoxia-induced *LOX* promotes bone metastasis of breast cancer (18), there was no evidence as to whether this is a generalised function; *i.e.* if tumour-derived *LOX* participates in the development of bone metastasis in other cancer models, such as colon cancer. We have previously shown that *LOX* synergizes with HIF-1 $\alpha$  in promoting *in vivo* growth of colorectal cancers (2). In the present study, we found that *LOX* was expressed by tumour cells in the bone marrow from CRC patients with bone metastases and that, irrespective of *HIF-1* expression, tumour-secreted *LOX* promoted settlement of CRC cells in the bone marrow and induced a robust expression of IL-6 by these tumour cells. In turn, both *LOX* and IL-6 played a prominent role in enhancing differentiation of mature osteoclasts. In breast cancer, it has been shown that *LOX* stimulates the generation of differentiated osteoclasts through the activation of NFATc1 (18). *LOX* secreted from CRC cells also promoted the nuclear translocation of NFATc1 in osteoclasts. Additionally, *LOX* inhibited osteoblast differentiation, thereby creating an imbalance between bone resorption and bone formation. Based on these findings we propose a model in which *LOX* supports CRC cell dissemination in the bone marrow and *LOX*-driven IL-6 production by CRC cells impairs bone homeostasis, thereby promoting osteolytic lesion formation *in vivo* (Fig. 6d).

Having shown that tumour-secreted *LOX* drives osteolytic lesion formation *in vivo*, we then attempted to determine the specific steps of bone colonization to which *LOX* functions are manifested. It has been previously reported that *LOX* enhances the

metastatic trait of breast tumours in animals by stimulating *TWIST1* expression in tumour cells (8). Additionally, LOX overexpression in breast cancer has been associated with EMT of tumour cells (26). Here, *LOX* overexpression in Hct116 cells did not modify the morphological appearance of these cells and it did not modify the expression of transcription factors *TWIST1* and *SNAI1*, indicating that LOX in CRC cells did not contribute to tumour cell dissemination in bone by promoting an EMT. Instead, our results established that LOX directly plays a stimulatory role in the survival of CRC cells in the bone marrow by enhancing their attachment to type-I collagen. These results are in agreement with previous findings (1,27,28), showing that LOX is a major enhancer of extracellular matrix stiffness and that, during metastasis initiation, tumour cells respond to tensile forces imposed by extracellular matrix stiffness with activation of AKT, FAK, Src and/or STAT3 signalling pathways. We previously showed that LOX induces AKT phosphorylation in Hct116 cells (2). Here, FAK and Src activation in CRC cells occurred irrespective of LOX expression, whereas STAT3 was activated by LOX through a IL-6/IL-6R-dependent signalling pathway. Additionally, blockade of the IL-6 receptor by tocilizumab abrogated STAT3 phosphorylation in LOX+ Hct116 cells. However, tocilizumab did not inhibit tumour cell dissemination in the bone marrow *in vivo*, whereas the LOX inhibitor  $\beta$ APN did. Interestingly, extracellular matrix tension can directly drive STAT3 activation in pancreatic ductal carcinoma cells (28). It is conceivable that LOX promotes STAT3 activation in CRC cells by increasing extracellular matrix stiffness. Our findings do not preclude the possibility that other members of the LOX family could contribute to the dissemination of CRC cells in the bone marrow. However, endogenous mRNA levels of *LOXL1*, *LOXL2*, *LOXL3* and *LOXL4* in Hct116 cells were only barely detectable (2), and *LOX* overexpression in these tumour cells did not modify expression levels of other LOX family members (Supplementary Table S3).

Although overt bone metastasis is rare in colon cancer and frequent in breast cancer, the detection rate of DTCs in the bone marrow from CRC patients is comparable to what is observed in breast cancer (14). It seems therefore that colon cancer DTCs should harbor a mechanism to grow out, but this appears to be somehow blocked in colon cancer patients. Only 5 CRC patients with bone metastases were studied here. However, all of them had LOX-positive tumour cells in the bone marrow. Additionally, LOX overexpression endowed CRC cells with full competence for aggressive colonization in bone *in vivo*. Environments that are rich in type-I collagen may be critical for DTCs, promoting their transition from dormancy to metastatic growth (29). Given the role of LOX in driving collagen crosslinking (1), it is conceivable that LOX participates to metastatic outgrowth of DTCs.

In conclusion, our findings collectively show that LOX supports CRC cell dissemination in the bone marrow and they reveal a novel mechanism through which LOX-driven IL-6 production by CRC cells impairs bone homeostasis. We believe this is a crucially important observation, which supports targeting LOX for metastasis prevention.

## **Acknowledgements**

We acknowledge Dr Cyril Confavreux (Lyon, France) for providing tocilizumab and Dr. Sabine Riethdorf (Hamburg, Germany) for conducting immunofluorescence experiments on CRC cells. We also thank Dr Klaus Pantel (Hamburg, Germany) and Dr Catherine Panabières (Montpellier, France) for valuable discussion.

## **References**

- 1 - Barker HE, Cox TR, Erler JT. The rationale for targeting the LOX family in cancer. *Nat Rev Cancer* 2012;12:540-552.
- 2 - Pez F, Dayan F, Durivault J, Kaniewski B, Aimond G, Le Provost GS, et al. The HIF-1-inducible lysyl oxidase activates HIF-1 via the Akt pathway in a positive regulation loop and synergizes with HIF-1 in promoting tumor cell growth. *Cancer Res* 2011;71:1647-1657.
3. - Barker HE, Chang J, Cox TR, Lang G, Bird D, Nicolau M, et al. LOXL2-mediated matrix remodeling in metastasis and mammary gland involution. *Cancer Res* 2011;71:1561-1572.
- 4 - Baker AM, Bird D, Lang G, Cox TR, Erler JT. Lysyl oxidase enzymatic function increases stiffness to drive colorectal cancer progression through FAK. *Oncogene* 2013;32:1863-1868.
- 5 - Cox TR, Bird D, Baker AM, Barker HE, Ho MW, Lang G, et al. LOX-mediated collagen crosslinking is responsible for fibrosis-enhanced metastasis. *Cancer Res* 2013;73:1721-1732.
- 6 - Levental KR, Yu H, Kass L, Lakins JN, Egeblad M, Erler JT, et al. Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell* 2009;139:891-906.

- 7 - Barry-Hamilton V, Spangler R, Marshall D, McCauley S, Rodriguez HM, Oyasu M, et al. Allosteric inhibition of lysyl oxidase-like-2 impedes the development of a pathologic microenvironment. *Nat Med* 2010;16:1009-1017.
- 8 - El-Haibi CP, Bell GW, Zhang J, Collmann AY, Wood D, Scherber CM, et al. Critical role for lysyl oxidase in mesenchymal stem cell-driven breast cancer malignancy. *Proc Natl Acad Sci USA* 2012;109:17460-17465.
- 9 - Erler JT, Bennewith KL, Nicolau M, Dornhöfer N, Kong C, Le QT, et al. Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature* 2006; 440:1222-1226.
- 10 - Erler JT, Bennewith KL, Cox TR, Lang G, Bird D, Koong A, et al. Hypoxia-induced lysyl oxidase is a critical mediator of bone marrow cell recruitment to form the premetastatic niche. *Cancer Cell* 2009;15:35-44.
- 11 - Wong CC, Gilkes DM, Zhang H, Chen J, Wei H, Chaturvedi P, et al. Hypoxia-inducible factor 1 is a master regulator of breast cancer metastatic niche formation. *Proc Natl Acad Sci USA* 2011;108:16369-16374.
- 12 - Bondareva A, Downey CM, Ayres F, Liu W, Boyd SK, Hallgrimsson B, et al. The lysyl oxidase inhibitor b-aminopropionitrile diminishes the metastatic colonization potential of circulating breast cancer cells. *PLOS One* 2009;4:e5620.
- 13 - Weilbaecher KN, Guise TA, McCauley LK. Cancer to bone: a fatal attraction. *Nat Rev Cancer* 2011;11:411-424.
- 14 - Pantel K, Schlimok G, Braun S, Kutter D, Lindemann F, Schaller G, et al. Differential expression of proliferation-associated molecules in individual micrometastatic carcinoma cells. *J Natl Cancer Inst.* 1993;85:1419-1424.
- 15 - Flatmark K, Borgen E, Nesland JM, Rasmussen H, Johannessen HO, Bukholm I, et al. Disseminated tumour cells as a prognostic biomarker in colorectal cancer. *Br J Cancer* 2011;104:1434-1439.



- 16 - Qiu M, Hu J, Yang D, Cosgrove DP, Xu R. Pattern of distant metastases in colorectal cancer: a SEER based study. *Oncotarget* 2015;6:38658-38666.
- 17 - Sosa MS, Bragado P, Aguirre-Ghiso JA. Mechanisms of disseminated cancer cell dormancy: an awakening field. *Nat Rev Cancer*. 2014;14:611-22.
- 18 - Cox TR, Rumney RM, Schoof EM, Perryman L, Høye AM, Agrawal A, et al. The hypoxic cancer secretome induces pre-metastatic bone lesions through lysyl oxidase. *Nature* 2015;522:106-110.
- 19 - Noblesse E, Cenizo V, Bouez C, Borel A, Gleyzal C, Peyrol S, et al. Lysyl oxidase-like and lysyl oxidase are present in the dermis and epidermis of a skin equivalent and in human skin and are associated to elastic fibers. *J Invest Dermatol* 2004;122:621-630.
- 20 - Croset M, Goehrig D, Frackowiak A, Bonnelye E, Ansieau S, Puisieux A, et al. *TWIST1* expression in breast cancer cells facilitates bone metastasis formation. *J Bone Miner Res* 2014;29:1886-1899.
- 21 - Fradet A, Sorel H, Depalle B, Serre CM, Farlay D, Turtoi A, et al. A new murine model of osteoblastic/osteolytic lesions from human androgen-resistant prostate cancer. *PLoS ONE* 2013;8:e75092.
- 22 - Hsu CP, Chen YL, Huang CC, Chou CC, Liu CL, Hung CH, et al. Anti-interleukin-6 receptor antibody inhibits the progression in human colon carcinoma cells. *Eur J Clin Invest* 2011;41:277-284.
- 23 - Lin L, Liu A, Peng Z, Lin HJ, Li PK, Li C, et al. STAT3 is necessary for proliferation and survival in colon cancer-initiating cells. *Cancer Res* 2011;71:7226-7237.
- 24 - Rossert J, De Crombrughe B. Type I collagen: Structure, synthesis and regulation. In: Bilezikian JP, Raisz LA, Rodan GA (eds) *Principles of Bone Biology*, 2<sup>nd</sup> ed., vol. 1. Academic Press, San Diego, CA, USA: 189-210 (2013).

- 25 - Shaheen S, Ahmed M, Lorenzi F, Nateri AS. Spheroid-Formation (Colonosphere) Assay for in Vitro Assessment and Expansion of Stem Cells in Colon Cancer. *Stem Cell Rev.* 2016. [Epub ahead of print]
- 26 - Han Y, Lian S, Cui X, Meng K, Györfy B, Jin T et al. Potential options for managing LOX+ ER- breast cancer patients. *Oncotarget* 2016. [Epub ahead of print]
- 27- Oskarsson T, Batlle E, Massagué J. Metastatic stem cells : sources, niches, and vital pathways. *Cell Stem Cell* 2014 ;14 :306-321.
- 28- Laklai H, Miroshnikova YA, Pickup MW, Collisson EA, Kim GE, Barrett AS, *et al.* Genotype tunes pancreatic ductal adenocarcinoma tissue tension to induce extracellular matrix fibrosis and tumor progression. *Nat Med.* 2016;22:497-505.
- 29 - Barkan D, El Touny LH, Michalowski AM, Smith JA, Chu I, Davis AS, *et al.* Metastatic growth from dormant cells induced by a col-I-enriched fibrotic environment. *Cancer Res.* 2010;70:5706-5716.

## Figure legends

**Figure 1. LOX expression in pairs of human primary colorectal carcinoma and their matching bone metastasis.** Immunohistochemistry was performed using 3 pairs of patients' tumour specimens. A strong immunostaining for LOX was observed in neoplastic cells (arrows) and, to a less extent, in the desmoplastic tumor stroma. In bone (\*), immunostaining for LOX was observed in osteocytic lacunae. Scale bar: 1-mm.

**Figure 2. High LOX expression in colorectal cancer cells promotes osteolytic lesion formation *in vivo*.** (a) Whole body bioluminescence imaging of animals at day 31 after intra-arterial inoculation of luciferase-expressing Hct116 cells. Ctrl: parental Hct116 cells. LOX+: *LOX*-overexpressing Hct116 cells. Ctrl/*HIF-1 $\alpha$* -: shRNA-*HIF1 $\alpha$* -expressing

Hct116 cells. LOX+/HIF-1 $\alpha$ - : LOX-overexpressing Hct116 cells silenced for HIF-1 $\alpha$ . The images shown are examples that best illustrate data obtained for each group. **(b)** Quantification of the area of osteolytic lesions (mm<sup>2</sup>) on radiographs on day 31 after tumour cell inoculation. *Inset* : Radiographs that best illustrate data obtained for each group. Arrows indicate osteolytic lesions. **(c)** *Left-hand panels*: Radiographic analysis of hind limbs from mice bearing Hct116, HT29 or LS174Tr colorectal tumours. Images best illustrate the effect of LOX on osteolytic lesion formation. Arrows indicate osteolytic lesions. Ctrl: parental tumour cells; LOX+: LOX-overexpressing tumour cells; LOX-: shRNA-LOX-expressing tumour cells. *Right-hand panels*: Quantification of the area of osteolytic lesions (mm<sup>2</sup>) on radiographs at day 35 after tumour cell inoculation. **(d)** *Upper panels*: Goldner's trichrome staining of tissue sections of tibial metaphysis from animals inoculated with tumour cells. Bone is stained green whereas bone marrow and tumor cells are stained red. The dotted line delimits the extent of tumour burden. *Lower panels*: TRAP-stained metastatic bone tissue sections. Osteoclasts are stained red (arrows). All images were obtained from different mice on day 35 after tumour cell inoculation. The images shown are examples that best illustrate LOX's effects on bone destruction and tumour burden. Scale bar: 0.1 mm. Data are mean  $\pm$  SD. \*, \*\*, \*\*\*:  $p < 0.05$ , 0.01 or 0.001, respectively. NS: not significant.

**Figure 3. Systemic delivery of tumour-secreted LOX enhances osteolytic lesion formation and tumour cell outgrowth in the bone marrow.** **(a)** Schematic representation of the experimental protocol. Parental Hc116 cells were inoculated intra-arterially to 4-week-old female Balb/c nude mice. Animals received a 5-week daily treatment with the conditioned medium from LOX- or LOX+ Hct116 cells, starting one day before tumour cell inoculation. Tumour-bearing mice were analyzed by radiography

(RX) at day 35 after tumour cell inoculation, then culled and the bone marrow was collected for tumour cell colony assays. **(b)** Quantification of osteolytic lesions (mm<sup>2</sup>) on radiographs. *Inset:* Radiographs of hind limbs with osteolytic lesions (arrows) that best illustrate data obtained for animals treated with the conditioned medium of LOX- or LOX+ Hc116 cells. \*\*:  $p < 0.01$  **(c)** Graphs showing the total number of tumour cell colonies formed in the bone marrow of each mouse treated with the LOX- or LOX+ conditioned medium. Representative images of tumour cell colonies are shown for each group.

**Figure 4. LOX-driven IL-6 production by colorectal cancer cells stimulates osteoclastogenesis.** **(a)** *In vitro* osteoclast differentiation of murine bone marrow cells treated with M-CSF+RANKL in combination with the conditioned medium from Ctrl-, LOX- or LOX+ Hct116 cells. The conditioned medium from LOX+ Hct116 cells treated with  $\beta$ APN was also tested. Mature osteoclasts were quantified as multinucleated (more than 3 nuclei), TRAP-positive cells. **(b)** ELISA measurement of IL-6 in the conditioned medium from Ctrl-, LOX-, LOX+ or LOX+/ $\beta$ APN Hct116 cells. **(c)** Immunoblot analysis of phosphorylated STAT3 in Ctrl-, LOX-, and LOX+ Hc116 cells, treated or not treated with tocilizumab. Graphs show intensities of phosphorylated STAT3 relative to total STAT3. Tubulin was used as a control for equal loading. **(d)** Correlation analysis of *LOX* and *IL-6* expression intensities from 7 databases on colon cancer (n = 1,146 patients). **(e)** Effect of exogenous IL-6 (10 ng/ml) on osteoclastogenesis induced with M-CSF+RANKL in combination with the conditioned medium of LOX- Hct116 cells, compared with that of LOX+ Hct116 cells. **(f)** Effects of the anti-IL6 receptor antibody tocilizumab (100 ng/ml) and of  $\beta$ -APN (350 $\mu$ M) on osteoclastogenesis induced with M-CSF+RANKL in

combination with the conditioned medium of LOX+ Hct116 cells. \*,\*\*:  $p < 0.05$  or  $0.01$ , respectively.


**Figure 5. LOX stimulates osteoclastogenesis and inhibits osteoblastogenesis. (a)**

Recombinant LOX (rLOX; 150 ng/ml) enhanced osteoclast differentiation induced by RANKL+M-CSF. **(b)** Conditioned medium from LOX+ Hct116 cells promoted the nuclear translocation of NFATc1 in osteoclasts, compared to conditioned medium from parental Hct116 cells (Ctrl). Nuclei are stained blue with DAPI. **(c)** and **(d)** *In vitro* differentiation of primary mouse calvaria cells cultured in osteogenic conditions with conditioned media from Ctrl, LOX- and LOX+ Hct116 cells. Bone mineralized nodules (stained black with von Kossa staining) were counted under light microscopy. Data are mean  $\pm$  SD. \*,\*\*:  $P < 0.05$  and  $0.01$ , respectively.

**Figure 6. Systemic delivery of tumour-secreted LOX promotes the settlement of colorectal cancer cells in the bone marrow. (a)**

Schematic representation of the experimental protocol. Parental Hct116 cells were inoculated intra-arterially to *nude* mice, then animals were treated with conditioned media of LOX- or LOX+ Hct116 cells (treated or not treated with  $\beta$ APN or tocilizumab). **(b)** Graphs showing for each group the average number of tumour cell colonies recovered from the bone marrow. Each bar corresponds to a single mouse. Data are mean  $\pm$  SD. \*\*:  $P < 0.01$ . **(c)** Attachment of Ctrl-, LOX- and LOX+ Hct116 cells to increasing concentrations of adhesive substrates (collagen, fibronectin). Data are mean  $\pm$  SD. \*,\*\*:  $P < 0.05$  and  $0.01$ , respectively. **(d)** Schematic representation of the role of LOX during bone metastasis formation. Tumour-secreted LOX enhanced tumour cell attachment to type-I collagen and induced IL-6 production which, in turn, acted as an autocrine agent and activated IL-6R/STAT3

signalling pathway. Additionally, IL-6 and LOX (through the nuclear translocation of NFATc1) activated osteoclastogenesis, leading to osteolytic lesion formation *in vivo*.



Reynaud et al., Fig. 1




Fig.2: Reynaud et al.






**A****B****C**

Fig. 3: Reynaud et al.




Fig. 4: Reynaud et al.




Fig. 5: Reynaud et al.




Fig. 6: Reynaud et al.

**Table 1.** Effects of *LOX* overexpression or silencing in Hct116 colorectal cancer cells on bone metastasis formation in animals.

Cell line	Histomorphometry				TRAP staining	
	BV/TV (%)	<i>p</i>	TuV/STV (%)	<i>p</i>	OC.S/BS (%)	<i>p</i>
Ctrl	26.5 ± 2.3 (n=8)	-	9.8 ± 6.4 (n=8)	-	31 ± 4.1 (n=5)	-
<i>LOX</i> +	13.9 ± 2.2 (n=8)	0.0006	57.6 ± 5.5 (n=8)	0.0016	73 ± 6 (n=7)	0.0017
<i>LOX</i> -	28.3 ± 2.9 (n=7)	0.53	2.6 ± 2.1 (n=7)	0.33	19 ± 8.3 (n=3)	0.09

Values are mean ± SD. Values of *p* (two sided) are for pairwise comparison with the control group (Ctrl) using the Mann-Whitney U test.

BV/TV = bone volume-to-tissue volume ratio; TuV/STV = tumour volume-to-total soft tissue volume ratio; OC.S/BS = active osteoclast-resorption surface per trabecular bone surface.; TRAP = tartrate-resistant acid phosphatase.