



HAL
open science

Electron-impact excitation of the $(4p55s5p)$ $4S_{3/2}$ quasimetastable state in Rb

V Roman, S. Demes

► **To cite this version:**

V Roman, S. Demes. Electron-impact excitation of the $(4p55s5p)$ $4S_{3/2}$ quasimetastable state in Rb. XXXI International Conference on Photonic, Electronic, and Atomic Collisions (ICPEAC2015), Jul 2019, Deauville, France. pp.132012, 10.1088/1742-6596/1412/13/132012 . hal-03383342

HAL Id: hal-03383342

<https://hal.science/hal-03383342>

Submitted on 9 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

PAPER • OPEN ACCESS

Electron-impact excitation of the $(4p^5 5s 5p)^4 S_{3/2}$ quasimetastable state in Rb

To cite this article: V Roman and S Demes 2020 *J. Phys.: Conf. Ser.* **1412** 132012

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Electron-impact excitation of the $(4p^55s5p)^4S_{3/2}$ quasimetastable state in Rb

V Roman^{1*} and S Demes²

¹Institute of Electron Physics, National Academy of Sciences of Ukraine, Uzhgorod, 88017, Ukraine

²Institute for Nuclear Research, Hungarian Academy of Sciences, Debrecen, 4026, Hungary

Synopsis The ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ quasimetastable state in rubidium was experimentally investigated by electron-impact in the energy range from the excitation threshold at 16.64 eV to 100 eV. The obtained data at 0.12 eV energy resolution indicate the presence of negative-ion resonances just above excitation threshold. A comparison is made with the optical excitation function of the $(4p^55s5p)^4S_{3/2} \rightarrow (4p^65p)^2P_{3/2}$ transition ($\lambda = 82.37$ nm), measured earlier with a lower energy resolution of ≈ 1.5 eV.

It is known that in alkali atoms (K, Rb, Cs) there is a subclass of p^6 -core excited quartet states which is due to the mixing with doublet states of the same total electronic angular momentum J possess both radiative and electronic decay channels [1]. As a rule, the radiative decay dominates for such quasimetastable states, while autoionization probability remains small. The latter makes difficult the experimental observation of quasimetastable states in ejected-electron spectra [2]. The optical excitation function for the $(4p^55s5p)^4S_{3/2} \rightarrow (4p^65p)^2P_{3/2}$ transition ($\lambda = 82.37$ nm) in Rb was measured earlier with an incident-electron energy resolution of ≈ 1.5 eV [3]. In the present work we obtained the ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ state with an improved 0.12 eV energy resolution by using energy-selective electron beam and by accurate measurement of the ejected-electron spectra in a broad impact energy range.

The apparatus and the experimental procedure were described in details earlier [4]. The ejected-electron spectra were measured at 54.7° observation angle. Due to the overlapping of the $^4S_{3/2}$ line with the high intensity neighboring line $(4p^54d5s)^4P_{1/2}$ with 16.69 eV excitation threshold, the total relative uncertainty is approximately 40% for the most of our data.

Figure 1 shows the measured ejected-electron excitation function of the $(4p^55s5p)^4S_{3/2}$ state and the optical excitation function for $\lambda = 82.37$ nm [3] in an impact-energy range from the 16.64 eV excitation threshold up to 100 eV. Our comparison shows that there is a fairly good agreement between the two types of data. Due to the better energy resolution of the present measurements, the “fine” structure *a-b* is clearly seen in the ejected-electron function just above the excitation threshold. The broad




Figure 1. The electron-impact excitation functions of the $(4p^55s5p)^4S_{3/2}$ state in rubidium: ● - present data; ○ - optical data [3]. The inset shows the threshold region of the present data. The data [3] and our data are normalized with incident-electron energy of 65 eV.

maximum *c* in our data [3] corresponds to the spin-exchange character of the electronic excitation of the $^4S_{3/2}$ state. The notable scatter of experimental data, which is observed between 20 and 25 eV in the ejected-electron excitation function could reflect the combined effect of several overlapping resonances.

In order to obtain the absolute values of the excitation cross sections for the $(4p^55s5p)^4S_{3/2}$ and other p^6 -core excited states in rubidium, we have performed relativistic distorted-wave calculations.

References

- [1] Mendelson A *et al* 1987 *Phys. Rev. A.* **35** 2095
- [2] Borovik A *et al* 2006 *Phys. Rev. A.* **73** 062701
- [3] Bogachev G 1974 *Ukr. Phys. J.* **12** 2061
- [4] Borovik A *et al* 2005 *J. Phys. B* **38** 1081

* E-mail: viktoriyaroman11@gmail.com