



Berry-Esseen type bounds for the matrix coefficients and the spectral radius of the left random walk on $GL_d(\mathbb{R})$

C Cuny, J Dedecker, F Merlevède, M Peligrad

► To cite this version:

C Cuny, J Dedecker, F Merlevède, M Peligrad. Berry-Esseen type bounds for the matrix coefficients and the spectral radius of the left random walk on $GL_d(\mathbb{R})$. 2021. hal-03388718v1

HAL Id: hal-03388718

<https://hal.science/hal-03388718v1>

Preprint submitted on 20 Oct 2021 (v1), last revised 27 Dec 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Berry-Esseen type bounds for the matrix coefficients and the spectral radius of the left random walk on $GL_d(\mathbb{R})$

C. Cuny*, J. Dedecker† and F. Merlevède ‡M. Peligrad §

October 20, 2021

Abstract

We give rates of convergence in the Central Limit Theorem for the coefficients and the spectral radius of the left random walk on $GL_d(\mathbb{R})$, assuming the existence of an exponential or polynomial moment.

1 Introduction

Let $(\varepsilon_n)_{n \geq 1}$ be independent random matrices taking values in $G = GL_d(\mathbb{R})$, $d \geq 2$ (the group of invertible d -dimensional real matrices) with common distribution μ . Let $\|\cdot\|$ be the euclidean norm on \mathbb{R}^d , and for every $A \in GL_d(\mathbb{R})$, let $\|A\| = \sup_{x, \|x\|=1} \|Ax\|$. Let also $N(g) := \max(\|g\|, \|g^{-1}\|)$. We shall say that μ has an exponential moment if there exists $\alpha > 0$ such that

$$\int_G (N(g))^\alpha d\mu(g) < \infty,$$

We shall say that μ has a polynomial moment of order $p \geq 1$ if

$$\int_G (\log N(g))^p d\mu(g) < \infty.$$

*Christophe Cuny, Université de Brest, LMBA, UMR 6205 CNRS, 6 avenue Victor Le Gorgeu, 29238 Brest

†Jérôme Dedecker, Université de Paris, CNRS, MAP5, UMR 8145, 45 rue des Saints-Pères, F-75006 Paris, France.

‡Florence Merlevède, LAMA, Univ Gustave Eiffel, Univ Paris Est Créteil, UMR 8050 CNRS, F-77454 Marne-La-Vallée, France.

§Magda Peligrad, Department of Mathematical Sciences, University of Cincinnati, PO Box 210025, Cincinnati, Oh 45221-0025, USA.

Let $A_n = \varepsilon_n \cdots \varepsilon_1$, with the convention $A_0 = \text{Id}$. It follows from Furstenberg and Kesten [9] that, if μ admits a moment of order 1 then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|A_n\| = \lambda_\mu \quad \mathbb{P}\text{-a.s.}, \quad (1.1)$$

where $\lambda_\mu := \lim_{n \rightarrow \infty} n^{-1} \mathbb{E} \log \|A_n\|$ is the so-called first Lyapunov exponent.

Let now $X := P(\mathbb{R}^d)$ be the projective space of \mathbb{R}^d and write \bar{x} as the projection of $x \in \mathbb{R}^d - \{0\}$ to X . An element A of $G = GL_d(\mathbb{R})$ acts on the projective space X as follows: $A\bar{x} = \overline{Ax}$. Let Γ_μ be the closed semi-group generated by the support of μ . We say that μ is proximal if Γ_μ contains a matrix that admits a unique (with multiplicity 1) eigenvalue of maximal modulus. We say that μ is strongly irreducible if no proper union of subspaces of \mathbb{R}^d is invariant by Γ_μ . Throughout the paper, we assume that μ is strongly irreducible and proximal. In particular, there exists a unique invariant measure ν on $\mathcal{B}(X)$, meaning that for any bounded measurable function h from X to \mathbb{R} ,

$$\int_X h(x) d\nu(x) = \int_G \int_X h(g \cdot x) d\mu(g) d\nu(x). \quad (1.2)$$

Let W_0 be a random variable with values in the projective space X , independent of $(\varepsilon_n)_{n \geq 1}$ and with distribution ν . By the invariance of ν , we see that the sequence $(W_n := A_n W_0)_{n \geq 1}$ is a strictly stationary Markov chain with values in X . Let now, for any integer $k \geq 1$,

$$X_k := \sigma(\varepsilon_k, W_{k-1}) - \lambda_\mu = \sigma(\varepsilon_k, A_{k-1} W_0) - \lambda_\mu, \quad (1.3)$$

where, for any $g \in G$ and any $\bar{x} \in X$, $\sigma(g, \bar{x}) = \log(\|gx\|/\|x\|)$. Note that σ is an additive cocycle in the sense that $\sigma(g_1 g_2, \bar{x}) = \sigma(g_1, g_2 \bar{x}) + \sigma(g_2, \bar{x})$. Consequently

$$S_n := \sum_{k=1}^n X_k = \log \|A_n V_0\| - n\lambda_\mu,$$

where V_0 is a random variable such that $\|V_0\| = 1$ and $\bar{V}_0 = W_0$.

Benoist and Quint [2] proved that if μ has a moment of order 2, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(S_n^2) \rightarrow s^2 > 0, \quad (1.4)$$

and, for any $t \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \sup_{\|x\|=\|y\|=1} \left| \mathbb{P}(\log |\langle A_n x, y \rangle| - n\lambda_\mu \leq t\sqrt{n}) - \phi(t/s) \right| = 0,$$

where ϕ denotes the cumulative distribution function of the standard normal distribution.

Given a matrix $g \in GL_d(\mathbb{R})$ denote by $\lambda_1(g)$ its spectral radius (the greatest modulus of its eigenvalues). Aoun [1] proved that if μ has a moment of order 2, then, for any $t \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} |\mathbb{P}(\log(\lambda_1(A_n)) - n\lambda_\mu \leq t\sqrt{n}) - \phi(t/s)| = 0.$$

In this paper we provide rates of convergence in these Central Limit Theorems, if μ has either an exponential moment, or a polynomial moment of order $p \geq 3$.

2 The case of matrix coefficients

Theorem 2.1. *Let μ be a proximal and strongly irreducible probability measure on $\mathcal{B}(G)$.*

1. *Assume that μ has an exponential moment, and let $s > 0$ be defined by (1.4). Then there exists a positive constant K such that*

$$\sup_{\|x\|=\|y\|=1} \sup_{t \in \mathbb{R}} |\mathbb{P}(\log |\langle A_n x, y \rangle| - n\lambda_\mu \leq t\sqrt{n}) - \phi(t/s)| \leq \frac{K \log n}{\sqrt{n}}. \quad (2.1)$$

2. *Assume that μ has a polynomial moment of order $p \geq 3$ and let $s > 0$ be defined by (1.4). Then there exists a positive constant K such that*

$$\sup_{\|x\|=\|y\|=1} \sup_{t \in \mathbb{R}} |\mathbb{P}(\log |\langle A_n x, y \rangle| - n\lambda_\mu \leq t\sqrt{n}) - \phi(t/s)| \leq \frac{K}{n^{(p-1)/2p}}. \quad (2.2)$$

Remark 2.1. Very recently Dinh, Kaufmann and Wu [8] obtained the bound (2.1) without the extra logarithmic factor.

The proof of this theorem is based on Berry-Esseen estimates for $\log \|A_n x\| - n\lambda_\mu$ (given in [12] and [7]), and on the following elementary lemma (see lemma 5.1 in [11] for a similar result):

Lemma 2.1. *Let $(T_n)_{n \in \mathbb{N}}$ and $(R_n)_{n \in \mathbb{N}}$ be two sequences of random variables. Assume that there exist three sequences of positive numbers $(a_n)_{n \in \mathbb{N}}$, $(b_n)_{n \in \mathbb{N}}$ and $(c_n)_{n \in \mathbb{N}}$ going to infinity as $n \rightarrow \infty$, and a positive constant s such that*

$$\sup_{t \in \mathbb{R}} |\mathbb{P}(T_n \leq t\sqrt{n}) - \phi(t/s)| \leq \frac{1}{a_n}, \quad \text{and} \quad \mathbb{P}(|R_n| \geq \sqrt{2\pi n s}/b_n) \leq \frac{1}{c_n}.$$

Then,

$$\sup_{t \in \mathbb{R}} |\mathbb{P}(T_n + R_n \leq t\sqrt{n}) - \phi(t/s)| \leq \frac{1}{a_n} + \frac{1}{b_n} + \frac{1}{c_n}.$$

Proof of Lemma 2.1. Recall that ϕ is $1/\sqrt{2\pi}$ -Lipschitz. We have

$$\begin{aligned}\mathbb{P}(T_n + R_n \leq t\sqrt{n}) &\leq \mathbb{P}\left(T_n - \sqrt{2\pi n}s/b_n \leq t\sqrt{n}, -R_n \leq \sqrt{2\pi n}s/b_n\right) + \mathbb{P}\left(-R_n \geq \sqrt{2\pi n}s/b_n\right) \\ &\leq \mathbb{P}\left(T_n - \sqrt{2\pi n}s/b_n \leq t\sqrt{n}\right) + \mathbb{P}\left(-R_n \geq \sqrt{2\pi n}s/b_n\right).\end{aligned}$$

Hence

$$\begin{aligned}\mathbb{P}(T_n + R_n \leq t\sqrt{n}) - \phi(t/s) &\leq \frac{1}{a_n} + |\phi(t/s + \sqrt{2\pi}/b_n) - \phi(t/s)| + \frac{1}{c_n} \\ &\leq \frac{1}{a_n} + \frac{1}{b_n} + \frac{1}{c_n}.\end{aligned}$$

The lower bound may be proved similarly, by noting that

$$\begin{aligned}\mathbb{P}(T_n + \sqrt{2\pi n}s/b_n \leq t\sqrt{n}) - \mathbb{P}(R_n \geq \sqrt{2\pi n}s/b_n) &\leq \mathbb{P}(T_n + R_n \leq t\sqrt{n}, R_n \leq \sqrt{2\pi n}s/b_n) \\ &\leq \mathbb{P}(T_n + R_n \leq t\sqrt{n}).\end{aligned}\quad \square$$

Proof of Item 1 of Theorem 2.1. The proof follows the steps used in Section 8.3 of [5]. We shall need some notations. For every $\bar{x}, \bar{y} \in X$, let

$$d(\bar{x}, \bar{y}) := \frac{\|x \wedge y\|}{\|x\| \|y\|},$$

where \wedge stands for the exterior product, see e.g. [4, page 61], for the definition and some properties. Then, d is a metric on X . Let also

$$\delta(\bar{x}, \bar{y}) := \frac{|\langle x, y \rangle|}{\|x\| \|y\|}. \quad (2.3)$$

Recall that the function δ is linked to the distance d on X by the following: For every $\bar{x}, \bar{y} \in X$,

$$\delta^2(\bar{x}, \bar{y}) = 1 - d^2(\bar{x}, \bar{y}). \quad (2.4)$$

We shall also need the following result due to Guivarc'h [10] (see Theorem 14.1 in [3]):

Proposition 2.1. *Let μ be a proximal and strongly irreducible probability measure on $\mathcal{B}(G)$. Assume that μ has an exponential moment. Then, there exists $\eta > 0$, such that*

$$\sup_{\bar{y} \in X} \int_X \frac{1}{\delta(\bar{x}, \bar{y})^\eta} d\nu(\bar{x}) < \infty.$$

We start with the identity, for $\|x\| = \|y\| = 1$,

$$\begin{aligned}\log |\langle A_n x, y \rangle| &= \log \|A_n x\| + \log \frac{|\langle A_n x, y \rangle|}{\|A_n x\| \|y\|} \\ &= \log \|A_n x\| + \log \delta(A_n \cdot \bar{x}, \bar{y}).\end{aligned}$$

We shall then apply Lemma 2.1 to $T_n = \log \|A_n x\| - n\lambda_\mu$ and $R_n = \log \delta(A_n \cdot \bar{x}, \bar{y})$. Since μ has an exponential moment, we know from [12] that we can take $a_n = C\sqrt{n}$ in Lemma 2.1.

In view of Lemma 2.1, we see that Theorem 2.1 will be proved if we can show that there exist $\tau, K > 0$ such that (recall that $\delta(\cdot, \cdot) \leq 1$)

$$\mathbb{P}(|\log \delta(A_n \cdot \bar{x}, \bar{y})| > \tau \log n) = \mathbb{P}(\delta(A_n \cdot \bar{x}, \bar{y}) < n^{-\tau}) \leq \frac{K}{\sqrt{n}}, \quad (2.5)$$

which means that the sequences $(b_n)_{n \in \mathbb{N}}$ and $(c_n)_{n \in \mathbb{N}}$ are such that $b_n = \sqrt{2\pi n s}/(\tau \log n)$ and $c_n = \sqrt{n}/K$.

Recall the identity (2.4). As in [5], we have, using that $d(\cdot, \cdot) \leq 1$,

$$\begin{aligned}\delta^2(A_n \cdot \bar{x}, \bar{y}) &= 1 - d^2(A_n \cdot \bar{x}, \bar{y}) \geq 1 - (d(A_n \cdot \bar{x}, W_n) + d(W_n, \bar{y}))^2 \\ &\geq \delta^2(W_n, \bar{y}) - d^2(A_n \cdot \bar{x}, W_n) - 2d(A_n \cdot \bar{x}, W_n)d(W_n, \bar{y}) \\ &\geq \delta^2(W_n, \bar{y}) - 3d(A_n \cdot \bar{x}, W_n).\end{aligned} \quad (2.6)$$

Hence, to prove (2.5), it suffices to prove that there exist $\tau, K > 0$ such that,

$$\mathbb{P}(\delta^2(W_n, \bar{y}) < n^{-2\tau} + 3d(A_n \cdot \bar{x}, W_n)) \leq \frac{K}{\sqrt{n}}. \quad (2.7)$$

Now, since μ has a polynomial moment of order 3, by Lemma 6 of [5], there exists $\ell > 0$, such that

$$\mathbb{P}(d(A_n \cdot \bar{x}, W_n) \geq e^{-\ell n}) \leq \frac{C}{n}$$

(in fact this estimate remains true as soon as μ has a polynomial moment of order 2, via a monotonicity argument).

Hence, for n large enough (such that $3e^{-\ell n} \leq n^{-2\tau}$), we have

$$\mathbb{P}(\delta^2(W_n, \bar{y}) < n^{-2\tau} + 3d(A_n \cdot \bar{x}, W_n)) \leq \mathbb{P}(\delta^2(W_n, \bar{y}) < 2n^{-2\tau}) + \frac{C}{n}.$$

On another hand, by Markov's inequality, since W_n has law ν ,

$$\mathbb{P}(\delta^2(W_n, \bar{y}) < 2n^{-2\tau}) = \nu \left\{ \bar{x} \in X : \frac{1}{\delta^2(\bar{x}, \bar{y})} > \frac{n^{2\tau}}{2} \right\} \leq \frac{2^{\eta/2}}{n^{\eta\tau}} \sup_{\bar{y} \in X} \int_X \frac{1}{\delta(\bar{x}, \bar{y})^\eta} d\nu(\bar{x}),$$

and (2.7) follows from Proposition 2.1 by taking $\tau = \frac{1}{2\eta}$. \square

Proof of Item 2 of Theorem 2.1. The proof follows the lines of that of Item 1. Instead of Proposition 2.1, we shall use the following result due to Benoist and Quint (see Proposition 4.5 in [2]):

Proposition 2.2. *Let μ be a proximal and strongly irreducible probability measure on $\mathcal{B}(G)$. Assume that μ has a polynomial moment of order $p > 1$. Then*

$$\sup_{\bar{y} \in X} \int_X |\log \delta(\bar{x}, \bar{y})|^{p-1} d\nu(\bar{x}) < \infty.$$

We shall then apply Lemma 2.1 to $T_n = \log \|A_n x\| - n\lambda_\mu$ and $R_n = \log \delta(A_n \cdot \bar{x}, \bar{y})$. Since μ has a moment of order 3, we know from [7] that we can take $a_n = C\sqrt{n}/\log n$ in Lemma 2.1 (and even $a_n = C\sqrt{n}$ if $p \geq 4$).

In view of Lemma 2.1, we see that Theorem 2.1 will be proved if we can show that there exists $K > 0$ such that

$$\mathbb{P}(|\log \delta(A_n \cdot \bar{x}, \bar{y})| > n^{1/2p}) \leq \frac{K}{n^{(p-1)/2p}}, \quad (2.8)$$

which means that the sequences $(b_n)_{n \in \mathbb{N}}$ and $(c_n)_{n \in \mathbb{N}}$ are such that $b_n = \sqrt{2\pi n s}/n^{1/2p}$ and $c_n = n^{(p-1)/2p}/K$.

Starting again from (2.6), we see that it suffices to prove that there exists $K > 0$ such that,

$$\mathbb{P}\left(\delta^2(W_n, \bar{y}) < e^{-2n^{1/2p}} + 3d(A_n \cdot \bar{x}, W_n)\right) \leq \frac{K}{n^{(p-1)/2p}}. \quad (2.9)$$

Proceeding as in the proof of Theorem 2.1, we deduce that, for n large enough (such that $e^{-\ell n} \leq e^{-2n^{1/2p}}$), we have

$$\mathbb{P}\left(\delta^2(W_n, \bar{y}) < e^{-2n^{1/2p}} + 3d(A_n \cdot \bar{x}, W_n)\right) \leq \mathbb{P}\left(\delta^2(W_n, \bar{y}) < 4e^{-2n^{1/2p}}\right) + \frac{C}{n}.$$

On another hand, by Markov's inequality, since W_n has law ν , and for n large enough,

$$\begin{aligned} \mathbb{P}\left(\delta^2(W_n, \bar{y}) < 4e^{-2n^{1/2p}}\right) &= \mathbb{P}\left(|\log \delta(W_n, \bar{y})| > n^{1/2p} - \log 2\right) \\ &= \nu\left\{\bar{x} \in X : |\log \delta(\bar{x}, \bar{y})| > n^{1/2p} - \log 2\right\} \\ &\leq \frac{1}{(n^{1/2p} - \log 2)^{p-1}} \sup_{\bar{y} \in X} \int_X |\log \delta(\bar{x}, \bar{y})|^{p-1} d\nu(\bar{x}), \end{aligned}$$

and (2.9) follows from Proposition 2.2. \square

3 The case of the spectral radius

We now prove similar results for the spectral radius. Given a matrix $g \in GL_d(\mathbb{R})$ denote by $\lambda_1(g)$ its spectral radius (the greatest modulus of its eigenvalues). The first result (Item 1 of

Theorem 3.1 below), assuming an exponential moment for μ , has been recently proved by Xiao et al. [13] (in fact, a stronger result is proved in [13]). We state it only for completeness.

Theorem 3.1. *Let μ be a proximal and strongly irreducible probability measure on $\mathcal{B}(G)$.*

1. *Assume that μ has an exponential moment, and let $s > 0$ be defined by (1.4). Then there exists a positive constant K such that*

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\log \lambda_1(A_n) - n\lambda_\mu \leq t\sqrt{n} \right) - \phi(t/s) \right| \leq \frac{K \log n}{\sqrt{n}}. \quad (3.1)$$

2. *Assume that μ has a polynomial moment of order $p \geq 3$, and let $s > 0$ be defined by (1.4). Then there exists a positive constant K such that*

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\log \lambda_1(A_n) - n\lambda_\mu \leq t\sqrt{n} \right) - \phi(t/s) \right| \leq \frac{K}{n^{(p-1)/2p}}. \quad (3.2)$$

The proof of Item 2 is based on a Berry-Esseen estimate for $\log \|A_n\| - n\lambda_\mu$ given in [7], and on Lemma 2.1.

A key ingredient in the proof of Item 1 by Xiao et al. [13] is Lemma 14.13 of [3].

To prove Item 2, we shall need a suitable version of Lemma 14.13 of [3]. The proof of Lemma 14.13 relies on Lemma 14.2 of [3] (of geometrical nature) and on large deviations, yielding to Lemma 14.3.

We shall need the following consequence of large deviation estimates of Benoist and Quint [2] (see also [6] for a related results under proximality).

Lemma 3.1. *Let μ be a strongly irreducible probability measure on $\mathcal{B}(G)$. Assume that μ has a polynomial moment of order $p > 1$. Let $\varepsilon > 0$. There exists $C > 0$ such that for every $n \in \mathbb{N}$*

$$\sup_{\|x\|=1} \mathbb{P} \left(\max_{1 \leq k \leq n} |\log \|A_k x\| - k\lambda_\mu| > \varepsilon n \right) \leq \frac{C}{n^{p-1}}, \quad (3.3)$$

$$\mathbb{P} \left(\max_{1 \leq k \leq n} |\log \|A_k\| - n\lambda_\mu| > \varepsilon n \right) \leq \frac{C}{n^{p-1}}; \quad (3.4)$$

$$\mathbb{P} \left(\max_{1 \leq k \leq n} |\log \|\Lambda^2(A_k)\| - k(\lambda_\mu + \gamma_\mu)| > \varepsilon n \right) \leq \frac{C}{n^{p-1}}. \quad (3.5)$$

Remark. Let us recall that, for any $A \in GL_d(\mathbb{R})$, $\Lambda^2(A)$ is the matrix on $\Lambda^2(\mathbb{R}^d)$ defined by $\Lambda^2(A)(x \wedge y) = Ax \wedge Ay$. In addition, in (3.5), γ_μ is the second Lyapunov exponent of μ . With the notations of [3, Section 14], λ_μ is denoted either $\lambda_{1,\mu}$ or λ_1 , while γ_μ is either denoted $\lambda_{2,\mu}$ or λ_2 .

Proof of Lemma 3.1. Let u_n be any of the left-hand side in (3.3), (3.4) or (3.5). It follows from Proposition 4.1 and Corollary 4.2 of [2] that

$$\sum_{n \geq 1} n^{p-2} u_n < \infty. \quad (3.6)$$

In fact, in [2], (3.6) is proved for u_n defined without the maximum over $k \in \{1, \dots, n\}$ under the probability. However, it is easy to see that the maximum over k can be added: it suffices to follow the proof of Theorem 2.2 of [2] with obvious changes, and to use a maximal version of Burkholder's inequality for martingales. Now, once (3.6) has been proven, it is easy to infer (via a monotonicity argument) that (3.3), (3.4) and (3.5) are satisfied. \square

Using Lemma 3.1 one can reproduce the proof of Proposition 14.3 of [3] to prove the following version of it.

Lemma 3.2. *Let μ be a strongly irreducible and proximal probability measure on $\mathcal{B}(G)$. Assume that μ has a polynomial moment of order $p > 1$. Then, the estimates (14.5), (14.6), (14.7) and (14.8) of [3] hold with $1 - \frac{C}{n^{p-1}}$ in the right-hand side instead of $1 - e^{-cn}$.*

Lemma 3.2 implies the next result.

Lemma 3.3. *Let μ be a strongly irreducible probability measure on $\mathcal{B}(G)$. Assume that μ has a polynomial moment of order $p > 1$. For every $\varepsilon > 0$, there exist $C > 0$ and $\ell_0 > 0$ such that for every $\ell_0 \leq \ell \leq n$,*

$$\mathbb{P}(\log(\lambda_1(A_n)) - \log \|A_n\| \geq -\varepsilon \ell) \geq 1 - \frac{C}{\ell^{p-1}}.$$

Proof of Lemma 3.3. The lemma is a version of Lemma 14.13 of [3] with the following difference: Lemma 14.13 holds under an exponential moment while in Lemma 3.3 we assume polynomial moments.

Now, it happens that there is a small gap in the proof of Lemma 14.13 of [3] which can be easily fixed thanks to a slight modification of the original argument. The problem comes from the fact that there is no reason why the property (14.43) should be true except on an exponentially small set, see the end of page 233 of [3] for the definition. We explain below how to fix the proof of the original Lemma 14.13 since the proof of our Lemma 3.3 may be done similarly, using our Lemma 3.2 instead of Lemma 14.3 of [3].

From (14.7) of [3] it follows that, with the notations of [3], for every $n \geq n_0$

$$\mu^{\otimes n} \left(\left\{ (b_1, \dots, b_n) \in G^n : \delta(x_{b_n \dots b_{[n/2]+1}}^M, y_{b_{[n/2]} \dots b_1}^m) \geq e^{-\varepsilon [n/2]} \right\} \right) \geq 1 - e^{-c[n/2]},$$

which yields easily (14.33) for all $[n/2] \leq \ell \leq n$, taking c smaller if necessary.

Hence, it remains to prove (14.33) for all $\ell_0 \leq \ell < [n/2]$, for some ℓ_0 large enough.

Let $\ell_0 \leq \ell < [n/2]$, with $\ell_0 \geq n_0$, where n_0 is such that Lemma 14.3 be true. Using that (14.39), (14.41) and (14.42) are true except on an exponentially small set (see the beginning of the proof of Lemma 14.13 which relies on Lemma 14.3), one sees that Lemma 14.13 will be proved (for that ℓ) if we can prove that

$$\mu^{\otimes n} \left(\left\{ (b_1, \dots, b_n) \in G^n : d(x_{b_n \dots b_{n-\ell}}^M, b_n \dots b_1 x_0) \leq e^{-(\lambda_{1,\mu} - \lambda_{2,\mu} - \varepsilon)\ell} \right\} \right) \geq 1 - e^{-c\ell}, \quad (3.7)$$

$$\mu^{\otimes n} \left(\left\{ (b_1, \dots, b_n) \in G^n : \delta(x_{b_n \dots b_{n-\ell}}^M, y_{b_{[n/2]}^m \dots b_1}) \geq e^{-\varepsilon\ell} \right\} \right) \geq 1 - e^{-c\ell}. \quad (3.8)$$

But (3.7) follows from (14.6) of [3] while (3.8) follows from (14.7) after noticing that $n - \ell \geq [n/2]$, so that the sets of coordinates $\{b_1, \dots, b_{[n/2]}\}$ and $\{b_{n-\ell}, \dots, b_n\}$ do not overlap. \square

Proof of Item 2 of Theorem 3.1. We shall apply Lemma 2.1 to $T_n = \log \|A_n\| - n\lambda_\mu$ and $R_n = \log(\lambda_1(A_n)) - \log \|A_n\|$. Since μ has a moment of order 3, we know from [7] that we can take $a_n = C\sqrt{n}/\log n$ in Lemma 2.1 (and even $a_n = C\sqrt{n}$ if $p \geq 4$).

In view of Lemma 2.1, we see that Item 2 of Theorem 3.1 will be proved if we can show that there exists $K > 0$ such that

$$\mathbb{P} \left(|\log(\lambda_1(A_n)) - \log \|A_n\|| > n^{1/2p} \right) \leq \frac{K}{n^{(p-1)/2p}}, \quad (3.9)$$

which means that the sequences $(b_n)_{n \in \mathbb{N}}$ and $(c_n)_{n \in \mathbb{N}}$ are such that $b_n = \sqrt{2\pi n s}/n^{1/2p}$ and $c_n = n^{(p-1)/2p}/K$.

Recall that $\lambda_1(g) \leq \|g\|$ for every $g \in GL_d(\mathbb{R})$. Hence (3.9) follows from Lemma 3.3 by taking $\varepsilon = 1$ and $\ell = n^{1/2p}$. \square

References

- [1] R. Aoun, *The central limit theorem for eigenvalues*, Proc. Amer. Math. Soc. 149 (2021), no. 2, 859-873
- [2] Benoist, Y. and Quint, J.-F. (2016). Central limit theorem for linear groups, *Ann. Probab.* **44** no. 2, 1308–1340.
- [3] Benoist, and Quint, J.-F. Random walks on reductive groups. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* **62** Springer, Cham, 2016. xi+323 pp.
- [4] Bougerol, P. and Lacroix, J. Products of random matrices with applications to Schrödinger operators. *Progress in Probability and Statistics*, 8. Birkhäuser Boston, Inc., Boston, MA, 1985.

- [5] Cuny, C., Dedecker, J. and Jan, C. (2017). Limit theorems for the left random walk on $GL_d(\mathbb{R})$. *Ann. Inst. H. Poincaré Probab. Statist.* **53**, no. 4, 1839–1865.
- [6] Cuny, C., Dedecker, J. and Merlevède, F. (2017). Large and moderate deviations for the left random walk on $GL_d(\mathbb{R})$. *ALEA Lat. Am. J. Probab. Math. Stat.* **14**, no 1, 503–527.
- [7] Cuny, C., Dedecker, J., Merlevède, F. and Peligrad, M. Berry-Esseen type bounds for the left random walk on $GL_d(\mathbb{R})$ under polynomial moment conditions. <https://hal.archives-ouvertes.fr/hal-03329189/document>
- [8] Dinh, T.-C., Kaufmann, L. and Wu, H. Berry-Esseen bound and local limit theorem for the coefficients of products of random matrices. <https://arxiv.org/pdf/2110.09032.pdf>
- [9] Furstenberg, H. and Kesten, H. (1960). Products of Random Matrices. *Ann. Math. Statist.* **31**, no. 2, 457–469.
- [10] Guivarc’h, Y. (1990). Produits de matrices aléatoires et applications. *Ergod. Theory Dyn. Syst.* **10**, 483–512.
- [11] Hörmann, S. (2009). Berry-Esseen bounds for econometric time series. *ALEA Lat. Am. J. Probab. Math. Stat.* **6** 377–397.
- [12] Le Page, E. (1982). Théorèmes limites pour les produits de matrices aléatoires, Probability measures on groups (Oberwolfach, 1981), pp. 258–303, Lecture Notes in Math., 928, Springer, Berlin-New York.
- [13] Xiao, H. Grama, I. and Liu, Q. Berry Esseen bounds and moderate deviations for the random walk on $GL_d(\mathbb{R})$. <https://hal.archives-ouvertes.fr/hal-02911533/document>