



HAL
open science

Compromised nuclear envelope integrity drives TREX1-dependent DNA damage and tumor cell invasion

Guilherme Pedreira de Freitas Nader, Sonia Agüera-Gonzalez, Fiona Routet, Matthieu Gratia, Mathieu Maurin, Valeria Cancila, Clotilde Cadart, Andrea Palamidessi, Rodrigo Nalio Ramos, Mabel San, et al.

► To cite this version:

Guilherme Pedreira de Freitas Nader, Sonia Agüera-Gonzalez, Fiona Routet, Matthieu Gratia, Mathieu Maurin, et al.. Compromised nuclear envelope integrity drives TREX1-dependent DNA damage and tumor cell invasion. *Cell*, 2021, 10.1016/j.cell.2021.08.035 . hal-03392287

HAL Id: hal-03392287

<https://hal.science/hal-03392287v1>

Submitted on 21 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title: Compromised nuclear envelope integrity drives TREX1-dependent**
2 **DNA damage and tumor cell invasion**

3
4 **Authors:** Guilherme Pedreira de Freitas Nader¹, Sonia Agüera-Gonzalez², Fiona Routet²,
5 Matthieu Gratia³, Mathieu Maurin³, Valeria Cancila⁴, Clotilde Cadart^{1,5}, Andrea Palamidessi¹¹,
6 Rodrigo Nalio Ramos¹², Mabel San Roman³, Matteo Gentili^{3,6}, Ayako Yamada^{7,8}, Alice Williard¹,
7 Catalina Lodillinsky^{9,10}, Emilie Lagoutte², Catherine Villard⁷, Jean-Louis Viovy⁷, Claudio
8 Tripodo⁴, Jérôme Galon¹², Giorgio Scita¹¹, Nicolas Manel^{3§}, Philippe Chavrier^{2§}, Matthieu
9 Piel^{1§&}

10 **Affiliations:**

11 1 Institut Curie and Institut Pierre Gilles de Gennes, PSL Research University, CNRS, UMR
12 144, Paris, France

13 2 Institut Curie, PSL Research University, CNRS, UMR 144, Paris, France

14 3 Institut Curie, PSL Research University, INSERM, U 932, Paris, France.

15 4 Tumor Immunology Unit, University of Palermo, Corso Tukory 211, 90234, Palermo, Italy

16 5 Current address: Molecular and Cell Biology Department, University of California, Berkeley,
17 142 Life Sciences Addition #3200, Berkeley, CA 94720-3200, USA

18 6 Current address: Broad Institute of MIT and Harvard, 415 Main Street, Cambridge, MA
19 02142, USA

20 7 Institut Curie, Université PSL, CNRS, UMR 168, Paris, France

21 8 Current address: PASTEUR, Département de chimie, École normale supérieure, Université
22 PSL, Sorbonne Université, CNRS, Paris, France

23 9 Research Area, Instituto de Oncología Ángel H. Roffo, Universidad de Buenos Aires, Buenos
24 Aires, Argentina

25 10 Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires,
26 Argentina

27 11 IFOM, the FIRC Institute of Molecular Oncology IFOM. Via Adamello 16 20139 Milano &
28 Department of Oncology and Hemato-oncology, University of Milan. IFOM. Via Adamello 16
29 20139 Milano, Italy

30 12 INSERM, Sorbonne Université, Université de Paris, Equipe Labellisée Ligue Contre le
31 Cancer, Centre de Recherche des Cordeliers, Laboratory of Integrative Cancer Immunology,
32 Paris, France.

33
34 § These authors contributed equally to the supervision of the work

35 & **Lead contact:** matthieu.piel@curie.fr

36 **Correspondence** should be sent to: nicolas.manel@curie.fr, philippe.chavrier@curie.fr,
37 matthieu.piel@curie.fr

38

1 Summary

2 While mutations leading to a compromised nuclear envelope cause diseases such as muscular
3 dystrophies or accelerated aging, the consequences of mechanically induced nuclear
4 envelope ruptures are less known. Here we show that nuclear envelope ruptures induce DNA
5 damage which promotes senescence in non-transformed cells, and induces an invasive
6 phenotype in human breast cancer cells. We find that the ER-associated exonuclease TREX1
7 translocates into the nucleus after nuclear envelope rupture and is required to induce DNA
8 damage. Inside the mammary duct, cellular crowding leads to nuclear envelope ruptures which
9 generate TREX1-dependent DNA damage, thereby driving the progression of *in situ* carcinoma
10 to the invasive stage. DNA damage and nuclear envelope rupture markers were also enriched
11 at the invasive edge of human tumors. We propose that DNA damage in mechanically
12 challenged nuclei could affect the pathophysiology of crowded tissues by modulating
13 proliferation and extracellular matrix degradation of normal and transformed cells.

14 Introduction

15 In tissues, extracellular matrix fibers and cell packing limit the available space. Migrating and
16 proliferating cells thus often face situations of crowding and confinement, which can result in
17 strong cell and nucleus deformations. We and others showed that, in the case of migrating
18 immune and cancer cells, strong nuclear deformation can result in transient events of nuclear
19 envelope (NE) rupture with subsequent DNA damage (Denais et al., 2016; Hatch, 2018; Hatch
20 and Hetzer, 2016; Irianto et al., 2017; Raab et al., 2016). In this context, efficient repair
21 responses are essential to ensure cell viability, which involve ESCRT proteins restoring NE
22 integrity, as well as BAF1 and the DNA repair pathways (Lusk and Ader, 2020). NE ruptures
23 can be induced by simply confining cells between two plates, independently of their migratory
24 behavior (Le Berre et al., 2012; Gentili et al., 2019; Hatch and Hetzer, 2016). They can also
25 occur spontaneously in fragile mutant nuclei (Vargas et al., 2012; De vos et al., 2011),
26 associated with a large range of degenerative diseases (Robijns et al., 2018). Recent studies
27 showed that these events also occur *in vivo* such as in muscle tissues which are highly solicited
28 mechanically; in the heart of developing embryos, due to stiffening and beating of the tissue
29 (Cho et al., 2019), as well as in skeletal muscle cells harboring mutations in the lamin A/C
30 gene, due to forces exerted on the nuclei by the microtubule cytoskeleton (Earle et al., 2020;
31 Roman et al., 2017). In these tissues, NE ruptures are associated with senescence and aging
32 phenomena that affect cell proliferation, survival, and ultimately tissue function. NE ruptures
33 are often accompanied by DNA damage (Cho et al., 2019; Denais et al., 2016; Earle et al.,
34 2020; Feng et al., 2018; Raab et al., 2016). Mechanisms leading to senescence and DNA
35 damage upon NE rupture are unclear. Here, we investigated the consequences of nuclear
36 deformation and rupture associated with cell migration in crowded tissues, focusing on
37 mammary carcinoma. These tumors, which grow confined inside the mammary ducts, have
38 been shown to display large strands of collectively moving cells associated with the transition
39 from *in situ* to the invasive stage of the tumor development (Palamidessi et al., 2019).
40 Additionally, the breast neoplastic invasive switch requires the activity of matrix proteases for
41 breaching out of the duct (Feinberg et al., 2018; Lodillinsky et al., 2016).

42 Results

43 We first investigated human samples of breast ductal adenocarcinoma *in situ* with
44 microinvasion stained by immunohistochemistry for the small GTPase RAB5A (involved in
45 recycling of cell/cell adhesions, used as a marker for strands of motile cells (Malinverno et al.,
46 2017) and the DNA damage marker phospho- γ H2AX. We observed that tumor regions positive
47 for RAB5A, often corresponding to micro-invasive foci (see methods and (Kim et al., 2018)),
48 displayed a larger number of deformed and elongated nuclei (Fig. 1A-E). This morphology
49 corresponds to cells squeezing each other as they move (Friedl and Gilmour, 2009;
50 Palamidessi et al., 2019). Nuclei positive for γ H2AX were more abundant in invasive foci than
51 in the tumor bulk (Fig. 1D), and, within invasive foci, γ H2AX-positive nuclei were more often
52 found in cells with higher RAB5A expression (presumably more migratory (Palamidessi et al.,

1 2019), Fig. 1E). This suggests that strands of motile cells at the periphery of the tumor exhibit
2 pronounced nuclear deformation, associated with elevated DNA damage.

3 We then investigated a xenograft mouse model based on the intraductal (intra-nipple) injection
4 of transformed human breast cancer MCF10DCIS.com (DCIS hereafter) cells, which are
5 derived from the non-tumorigenic breast epithelial MCF10A cell line. This model recapitulates
6 the transition from *in situ* to invasive stages of breast cancer progression (Behbod et al., 2009;
7 Lodillinsky et al., 2016). *In situ* tumors are characterized by a myoepithelial layer (stained with
8 smooth muscle actin) surrounding human xenografted cells (identified by a human-specific
9 KU70 antibody, Fig. 1F). Tumor sections were stained with γ H2AX (Fig. 1F-G). Advanced
10 tumors (classified based on the state of the myoepithelial layer) displayed more γ H2AX-
11 positive nuclei (Fig. 1F). Deformed nuclei were mostly found at the periphery of the tumor
12 mass, especially in ducts inflated with large tumors, and enriched in micro-invasive foci (Fig.
13 1G). Most of the γ H2AX-positive nuclei did not correspond to mitotic nor apoptotic figures (Fig.
14 S1A), which were excluded from the analysis. We performed an automated analysis of all
15 human nuclei in the tumor regarding their level of γ H2AX staining, as a function of their
16 distance to the tumor edge (Fig. S1A for the method). This analysis showed an enrichment for
17 nuclei with an elevated γ H2AX staining (Z-score above 2) and even more for nuclei with a very
18 high staining (Z-score above 5) in the region closer to the tumor edge (Fig. S1B and Fig. 1H).
19 Together, these data show that the periphery of mammary *in situ* carcinomatous lesions, both
20 in human and in xenografted mice, often display strongly deformed nuclei with elevated levels
21 of DNA damage. This raises the possibility that NE ruptures could be implicated in the induction
22 of the DNA damage, as suggested by previous work on migrating cells (Denais et al., 2016;
23 Raab et al., 2016).

24 Because NE ruptures of deformed nuclei were difficult to investigate in tumor samples, we
25 developed a 'duct-on-chip' microfluidic assay (inspired by (Blaha et al., 2017)). The assay
26 consists of three large 50 μ m-high channels: a lateral channel filled with culture medium, a
27 central one with DCIS cells, and the third channel with a dense fluorescently-labeled type I
28 collagen fibrillar matrix. The channels are connected by regularly spaced V-shape gates, 10
29 μ m-wide in their smaller section, through which cells can migrate towards the collagen-filled
30 chamber (Fig. 1I and Fig. S1C). Cells invaded efficiently the collagen chamber (Fig. 1I) and
31 visualization of the collagen network suggested that they were able to remodel the collagen
32 fibers over time (Fig. 1I). Using a pan inhibitor of matrix metalloproteinases (MMPs), GM6001,
33 we confirmed that collagen degradation by DCIS cells was required for invasion into the
34 collagen chamber (Fig. S1D, E), similar to what we had previously observed *in vivo* (Lodillinsky
35 et al., 2016). DCIS cells expressing catalytically inactive cGAS-mCherry (icGAS hereafter) to
36 detect NE ruptures (Gentili et al., 2019; Raab et al., 2016), and 53BP1-GFP to detect DNA
37 double-strand break (DSB) foci were grown in the device. The average cell density increased
38 by 50% from 24h to 48h after cell loading (Fig 1I). The device was loaded either with a low or
39 a high density of cells (Fig. S1 F, G, H), so that, after 48 hours, the low-seeding density would
40 remain below the initial high-seeding density. We then quantified the shape of nuclei, the
41 number of 53BP1-GFP foci and the fraction of nuclei showing icGAS-mCherry peripheral
42 accumulation, characteristic of NE rupture events (Fig. S1H, (Raab et al., 2016)). 48 h after
43 seeding at high, but not at low density, nuclear roundness decreased, indicating that nuclei
44 were getting confined and deformed (Fig. 1J), and the fraction of nuclei with NE ruptures and
45 elevated 53BP1 foci (> 3) increased (Fig. 1 K, L). DCIS cells remained motile even at high
46 density (Movie S1, (Malinverno et al., 2017; Palamidessi et al., 2019)). This suggests that,
47 similarly to what was observed for single migrating cells in a dense tissue (Denais et al., 2016;
48 Raab et al., 2016), the motility of DCIS cells in a crowded multicellular context can lead to
49 nuclear deformation, NE rupture and DNA damage.

50 To examine the link between nuclear deformation, NE rupture and DNA damage, we used a
51 2D confiner device that allows precise deformation of cells (and nuclei) between two parallel
52 plates (Le Berre et al., 2014). In addition to DCIS cells, we analyzed the non-tumorigenic cell
53 lines RPE1 (normal human hTERT immortalized cells, in which we have previously

1 characterized NE ruptures and DNA damage in migration assays (Raab et al., 2016) and
2 MCF10A expressing icGAS-mCherry and 53BP1-GFP (Fig. 2A). We found a sharp increase
3 in the fraction of RPE1 cells with ruptured nuclei below 3 μm confinement (Fig. S2A, B),
4 confirming our previous observations on another panel of cell types (Raab et al., 2016). To
5 assess the relationship between NE rupture and DNA damage, we analyzed the number of
6 53BP1-GFP foci in nuclei of cells confined at various heights and displaying NE ruptures or
7 not (Fig. S2A). Foci were counted at each time-point using a semi-automated process (see
8 methods and Fig. S2C). We found that, similarly to cells migrating through constrictions (Raab
9 et al., 2016), only cells with ruptured NE displayed increased number of 53BP1 foci (Fig. 2B-
10 D for RPE1 and S2D for DCIS and MCF10A). Almost all cells with NE rupture (> 98%)
11 displayed more than three 53BP1 foci (Fig. 2E-F), while only a small fraction of cells without
12 NE rupture (< 2%) had this number of foci over the course of the movie (Fig. 2E). Analysis of
13 the intensity of foci and of the background (Fig. S2E) showed that 53BP1 accumulated in an
14 increasing number of foci in nuclei with NE ruptures. Cells exhibiting NE ruptures were also
15 positive for other markers of DNA damage, such as RIF1 and γH2AX (Fig S2F, G). Additionally,
16 the number of 53BP1 foci per nucleus did not depend on the cell cycle (Fig. S2H). Collectively,
17 these results indicate that confinement-induced DNA damage was preceded by NE rupture
18 events in a variety of cells, including RPE1 (in accordance with other reports (Raab et al., 2016;
19 Shah et al., 2021)), MCF10A and DCIS cells.

20 In cells displaying NE ruptures, the number of 53BP1 foci remained high throughout
21 the confinement period (Fig. 2C and S2D), and was associated with a majority of cells showing
22 repeated cycles of NE blebbing, rupture and repair over the entire duration of confinement, in
23 contrast to cells confined at 4 μm (Movie S2). We thus hypothesize that the occurrence of DNA
24 damage associated with deformed nuclei in the human tumor samples could correspond to
25 repeated NE ruptures.

26 Previous work suggested that NE rupture might lead to DNA repair proteins leaking out the
27 nucleus (Irianto et al., 2017). Some cytoplasmic factor(s) might also enter the nucleus and
28 access nuclear DNA, causing DNA damage (Maciejowski et al., 2015, 2019; Shah et al., 2017).
29 Live recording of 53BP1-GFP showed the expected dynamic appearance and disappearance
30 of single foci, indicative of DNA damage and repair events (Fig. 2G and Movie S2). We found
31 that the frequency of appearance of new foci was significantly higher in cells with NE rupture
32 with respect to cells without NE rupture, reaching almost the same rate as upon etoposide
33 treatment (Fig. 2H). Moreover, the number of new foci typically increased after a rupture event
34 (Fig. S2I, J), suggesting a causal relationship between NE rupture and new DNA damage. On
35 the contrary, lifetime of the foci showed no difference between cells with or without NE ruptures
36 (Fig. 2I). Overall, these experiments suggest that DNA damage associated with NE ruptures
37 is mostly due to the generation of new DNA damage following NE rupture events rather than
38 to delayed repair.

39 To investigate whether DNA damage was solely associated to NE rupture events or whether
40 cell and nuclear confinement were also required, we recorded non-confined cells plated on a
41 glass coverslip. In these conditions, although very rare, spontaneous NE rupture events could
42 be observed. Such events displayed a similar signature in terms of icGAS accumulation at a
43 local site on the nuclear periphery and leakage of 53BP1-GFP staining in the cytoplasm before
44 re-import into the nucleus (Fig. 2J and Movie S3). Each NE rupture event was followed, within
45 1 to 2 hours, by an increase in the number of 53BP1-GFP foci in the nucleus (Fig. 2J-L). To
46 investigate a larger population of cells, we fixed the cells and counted the number of foci using
47 53BP1-GFP (Fig. 2M, left), or γH2AX staining (Fig. 2M, right), in cells with or without NE rupture
48 (assessed by the presence of an icGAS perinuclear accumulation). The increase in the number
49 of foci appeared similar to the one observed after confinement-induced NE ruptures. Together
50 these experiments indicate that DNA damage follows NE rupture events even in the absence
51 of confinement.

1 NE ruptures at persistent post-mitotic exit chromatin bridges have been shown to cause DNA
2 damage, reduced upon depletion of TREX1, a cytoplasmic ER membrane-associated
3 exonuclease (Maciejowski et al., 2015). We silenced TREX1 by siRNA-mediated knockdown,
4 or permanently knocked it out by CRISPR technology (in RPE1 and DCIS cells, Fig. S3A, B).
5 We found that, upon TREX1 depletion in the three cell lines (RPE1, MCF10A, DCIS), NE
6 ruptures were no longer accompanied by an increase in 53BP1 foci (Fig. 3A, B, Fig. S3C-D,
7 quantified by an independent method in Fig. S3E). TREX1 depletion also prevented the
8 occurrence of DNA damage upon spontaneous NE rupture events in RPE1 cells (Fig. 3C, S4A,
9 B). 53BP1 foci mean intensity (Fig. S3F), the generation of foci upon etoposide treatment (Fig.
10 S3G) and NE rupture frequency (Fig. S3H) were not affected. Wild-type TREX1, but not of a
11 catalytically-deficient mutant (TREX1-D18N, (Lehtinen et al., 2008)) increased the number of
12 foci in cells with NE ruptures (Fig. 3A, B). These experiments unambiguously demonstrate a
13 role for TREX1 in promoting DNA damage upon NE rupture events in this panel of cell lines.

14 To determine how TREX1 gains access to nuclear DNA upon NE rupture events, we imaged
15 TREX1-GFP expressing cells. In non-confined cells, TREX1-GFP was mostly bound to
16 endomembranes and excluded from the nucleoplasm. Upon confinement, TREX1-GFP foci
17 were also found inside the nucleus (Fig. S3I, Movie S4). Immuno-electron microscopy using
18 gold immunolabelling against mCherry on RPE1 TREX1 KO cells reconstituted with TREX1-
19 mCherry, showed that, in non-confined cells, gold particles were found at ER membranes
20 distributed throughout the cytosol, co-localizing with the ER-resident enzyme, protein disulfide
21 isomerase (PDI) (Fig. 3D and S4C). PDI and TREX1 staining were increased inside the
22 nucleus in 2 μ m confined cells (Fig. 3D-E and S4C). TREX1 was particularly enriched at the
23 inner nuclear membrane of confined cells (Fig. S4C, and Tables S1 and S2, (Mayhew, 2011)).
24 When localized inside the nucleus, TREX1-mCherry appeared associated to ER-like
25 membranes together with PDI (Fig. 3D). The presence of such intranuclear ER-like
26 membranes was almost never observed in non-confined cells. These experiments suggest
27 that, upon strong confinement corresponding to heights at which cells undergo NE rupture
28 events, TREX1 localizes, together with other ER-bound proteins like PDI, from cytosolic ER,
29 to the inner side of the nuclear membrane and the inside of the nucleus, but remains mostly
30 bound to membranes.

31 Because TREX1 is an ER membrane protein, and based on the immuno-electron microscopy
32 results, we hypothesized that TREX1 might access the chromatin by diffusion from the ER
33 membrane to connected intra-nuclear membranes ((Ungricht and Kutay, 2015)). Accordingly
34 while WT TREX1 fully rescued the level of DNA damage induced by NE rupture in TREX1 KO
35 DCIS cells, the DeltaC construct, lacking the C-terminal ER-binding site of the protein, failed
36 to do so (Fig. S3J, K). A construct in which the C-terminus of the protein was replaced by the
37 ER binding site of Sec61 (Mohr et al., 2021) was able to partially rescue the level of DNA
38 damage induced by confinement in TREX1 KO DCIS cells (Fig. S3J,K), similarly to what was
39 observed in micro-nuclei (Mohr et al., 2021). Together, these experiments suggest that, upon
40 NE rupture events, ER bound TREX1 gains access to the interior of the nucleus, causing DNA
41 damage.

42 Having established a mechanism of DNA damage generation after NE rupture, we next sought
43 to examine the functional consequences of this damage. In non-transformed cells, chronic
44 exposure to DNA damage-inducing factors is associated with cellular senescence. RPE1 and
45 MCF10A cells showed a very clear induction of senescence upon treatment with etoposide
46 and doxorubicin (hereafter referred to as doxo) as verified by senescence-associated- β -
47 galactosidase (β -gal)-positive staining. However, senescence was less pronounced in the
48 tumorigenic DCIS cells (Fig. S5A). During confinement, even for long confinement periods,
49 most of the cells had not undergone mitosis nor shown any sign of death (Fig. S5B and C).
50 Following confinement, cells were harvested and replated in order to track single cells that had
51 ruptured their NE or not based on icGAS perinuclear staining/distribution (Fig S5D). After a
52 short period of confinement (2h), de-confined cells resumed proliferation with an almost normal
53 cell cycle duration (Fig. 4A, mean cell cycle duration for non-confined cells: 14.66 h +/- 2.31),

1 regardless of their NE status. In contrast, after a long period (12h), cells that displayed NE
2 ruptures, while showing no sign of death, did not resume proliferation (Fig. 4A, right panel)
3 whereas the few dividing cells showed a much longer cell cycle duration (Fig. 4A). These
4 results suggest that NE rupture and sustained confinement delayed cell cycle progression.

5 To characterize the senescence phenotype associated with strong confinement, RPE1 cells
6 were confined for short or long periods of time, under weak (4 μm) or strong (2 μm)
7 confinement, then harvested and replated for 96 h before testing a panel of senescence
8 markers (β -gal positivity, heterochromatin foci-HP1 γ staining, western blot for lamin B1 and
9 p21(Gorgoulis et al., 2019)). All these markers consistently showed that, while short term or
10 weak (4 μm) confinement did not induce senescence, long term, strong confinement, induced
11 a robust senescence phenotype, in both RPE1 and MCF10A, but not in DCIS cells (Fig 4B-G
12 and S5E-G). TREX1-depleted and TREX1 KO RPE1 cells showed no sign of senescence even
13 after strong and prolonged confinement (Fig. 4B, C, E-G and Fig. S5E) and expression of
14 TREX1 WT in TREX1 KO cells rescued the induction of p21 by confinement in RPE1 cells (Fig.
15 4E). The senescence phenotype was not dependent on the activation of the canonical cGAS
16 pathway ((Gorgoulis et al., 2019)), since RPE1 cells do not express cGAS (Fig S5H (Basit et
17 al., 2020)) and did not show any sign of cGAS enzymatic activity in response to transfected
18 DNA (Fig. S5I). Together, these results show that TREX1-dependent chronic DNA damage
19 induced upon strong confinement triggers the hallmarks of the senescence program in normal
20 RPE1 and MCF10A cells, but not in transformed DCIS cells.

21 To investigate the potential consequences of DNA damage associated with NE ruptures during
22 the development of tumors produced by DCIS cells, we used the duct-on-chip assay described
23 above. Both control- and TREX1-depleted cells reached high density within 48 h after injection
24 in the device, showing that proliferation was not affected by TREX1 depletion. They also
25 displayed comparable levels of NE ruptures (Fig. 5A). After 24 h in the device, there was no
26 appreciable difference in 53BP1 foci numbers between control- and TREX1-depleted cells
27 (Fig. 5A). In contrast, while this number significantly increased at 48 h for control cells, it
28 remained constant in TREX1-depleted cells (Fig. 5A). These results indicate that the increased
29 53BP1 foci levels observed at high density in the duct-on-chip system is a consequence of
30 TREX1-dependent DNA damage following nuclear deformations and ruptures.

31 TREX1-depleted DCIS cells can thus grow at high density in the duct-on-chip device without
32 showing an increase in DNA damage. We found that control cells invaded the collagen
33 chamber earlier and migrated farther than TREX1-depleted cells (Fig. 5B). This suggests that
34 the increased invasive behavior of control cells might be due more elevated DNA damage. To
35 test this hypothesis, differently-labelled control and doxo-treated cells were mixed and
36 introduced in the duct-on-chip device. We observed that 24 h after loading the cells into the
37 device (to ensure that cell density was still low), doxo-treated cells displayed increased
38 invasion potential into the collagen chamber as compared to DMSO-treated cells (Fig 5C).
39 These results indicate that sustained DNA damage can be a driver of the invasive behavior of
40 DCIS cells in a 3D matrix environment. We thus assessed the collagen degradation activity of
41 DCIS cells using an antibody that recognizes the collagenase-cleaved $\frac{3}{4}$ fragment of collagen
42 I (Castagnino et al., 2018; Infante et al., 2018; Lodillinsky et al., 2016). Cells were first treated
43 with drugs, or confined for several hours, and then were embedded in a 3D collagen gel in the
44 absence of any further DNA damaging condition. We first treated DCIS cells with increasing
45 doses of doxo and observed a dose-dependent increase in collagen degradation (Fig. S6A),
46 suggesting that prior exposure to DNA damage promotes the collagen degradation capacity of
47 DCIS cells. We then assayed cells after confinement. Once in the gel, the nuclear area of the
48 cells that had been confined at various heights, was comparable (Fig. S6B), showing no long-
49 lasting nuclear deformation. TREX1-proficient cells that had been confined at 2 μm height, but
50 not TREX1-depleted cells, showed a persistent, highly polarized morphology, increased speed
51 in the collagen network (Fig. S6C, quantified in D and Movie S5), and augmented collagen
52 degradation (Fig. 5D-F). In agreement, overexpressing TREX1-WT augmented collagen
53 degradation, while TREX1-D18N did not (Fig. 5E). Doxo treatment induced collagen

1 degradation in TREX1-depleted cells (Fig. S6E), confirming that TREX1 acts upstream of DNA
2 damage to induce collagen degradation. Taken together, these experiments show that, upon
3 strong confinement, TREX1-dependent DNA damage leads to a sustained increase in collagen
4 degradation by DCIS cells.

5 To further clarify how TREX1-mediated DNA damage induces collagen degradation, cells
6 confined at 2 μ m height were treated with the MMP inhibitor, GM6001, or with the ATM inhibitor
7 KU-55933. Both treatments inhibited collagen degradation (Fig. 5G). DNA damage during
8 interphase might lead to cGAS activation and the establishment of an EMT-like phenotype
9 (Harding et al., 2017). cGAS-depletion showed that, similarly to the senescence phenotype in
10 RPE1 cells, the induction of collagen degradation upon confinement was not cGAS-dependent
11 (Fig. S6F). Increased cellular motility and invasion induced by prolonged DNA damage and
12 ATM activation has been shown to promote a partial EMT state (Peng et al., 2019; Sun et al.,
13 2012). We analyzed the mRNA expression level of a series of canonical EMT genes in cells
14 confined at 10 (control) or 2 μ m, comparing WT and TREX1 KO cells. We found that SNAIL1
15 was specifically upregulated by confinement at 2 μ m and that SNAIL1 mRNA induction did not
16 occur in two independent TREX1 KO clones (Fig. S6G). In the absence of confinement,
17 SNAIL1 was induced by both Doxo and etoposide with a peak at 4 h of treatment (Fig. S6H).
18 Moreover, SNAIL1 silencing abrogated the increase in collagen degradation upon confinement
19 (Fig. S6I). Together, these results suggest that a partial EMT phenotype, including increased
20 matrix degradation and invasive potential, is induced in DCIS cells due to prolonged exposure
21 to DNA damage.

22 Putting these results in the context of our initial observations of nuclear deformation and DNA
23 damage both in human breast ductal carcinoma and intraductal DCIS cell tumor xenografts
24 (Fig. 1), we hypothesized that TREX1 is responsible for DNA damage in these deformed nuclei,
25 favoring a collagen degradation program, ultimately promoting the *in situ*-to-invasive
26 carcinoma transition. To test directly this hypothesis in an *in vivo* murine model, we injected
27 either parental or TREX1 KO DCIS cells inside the mammary ducts of immuno-suppressed
28 mice and allowed tumors to develop for 6 to 8 weeks. Both parental- and TREX1 KO-injected
29 mice formed intraductal *in situ* tumors at a similar rate (Fig. S6J). However, there was a sharp
30 decrease in the tumor area of TREX1 KO-injected mice compared to parental-injected mice
31 (Fig. S6K, L). Strikingly, while all the mice injected with parental DCIS cells showed invasive
32 tumors based on whole-mount carmine and histological staining within 6 weeks after injection,
33 the formation of invasive tumors in TREX1 KO-injected mice was greatly reduced for up to 8
34 weeks (Fig. 6A, B). This phenotype was comparable to the reduced invasive potential of
35 Membrane Type I (MT1)-MMP (MMP14)-deficient DCIS cells that we previously reported using
36 this model (Lodillinsky et al., 2016). The proliferation rates of parental DCIS and TREX1 KO
37 clones were similar *in vitro* (Fig. S6M). Additionally, analysis of parental and TREX1 KO tumor
38 xenografts revealed no differences in the fraction of cells positive for the proliferation marker
39 Ki67 (Fig. 6C). Parental tumors displayed a significantly larger percentage of γ H2AX-positive
40 nuclei than TREX1 KO tumors (Fig. 6 C-D). The fraction of cells with high and very high Z-
41 scores was reduced at the margin of TREX1 KO tumors, but not in the tumor bulk (Fig. 6 E, F
42 and S6N). Overall, these results suggest that the transition from *in situ* to the invasive stage is
43 favored in WT cells by TREX1-dependent DNA damage occurring in deformed and ruptured
44 nuclei as the tumor grows in the confined environment of the mammary ducts.

45 We also asked whether NE ruptures can be seen in the mice xenograft and in the human tumor
46 as well. We first used a combination of Lamin A/C and DAPI staining and looked for chromatin
47 herniations, i.e. interruptions of the Lamin A/C staining with a protrusion of chromatin labelled
48 by DAPI. We were able to find several such instances in the intraductal tumor xenografts (Fig.
49 7A), which correlated with nuclei positive for γ H2AX staining. Second, we used a combination
50 of cGAS and DAPI staining. In mouse xenografts, we found regions enriched with the typical

1 cGAS perinuclear accumulation, corresponding to γ H2AX-positive nuclei (Fig. 7B).
2 Immunohistochemistry provided even clearer examples of NE ruptures, with cGAS staining at
3 the periphery of nuclei (Fig. 7C). Of note, the cGAS antibody was validated *in vitro* on DCIS
4 cells depleted for cGAS (Fig. S7A). Human breast cancer samples displayed heterogenous
5 cGAS labelling and cGAS positive micro-nuclei, but we could also detect numerous NE rupture
6 events (Fig. S7B). A complete proof of the role of NE ruptures and TREX1 in human breast
7 tumors will require deeper investigation of a larger number of patient samples. These
8 observations show that NE rupture events occur *in vivo*, both in DCIS tumor xenografts and in
9 human tumors and that cGAS staining has the potential to be a powerful tool to visualize the
10 occurrence of NE rupture events in tumor samples.

11 Discussion

12 TREX1 is a cytoplasmic exonuclease, mostly found associated with ER membranes (Lee-
13 Kirsch et al., 2007a, 2007b), and shown to protect cells from the accumulation of cytoplasmic
14 DNA, either from viral or endogenous origins (Mazur and Perrino, 2001; Stetson et al., 2008).
15 Although TREX1 has already been proposed to act on nuclear DNA as well (Chowdhury et al.,
16 2006; Maciejowski et al., 2015, 2019), possibly in conjunction with an endonuclease
17 (Chowdhury et al., 2006), to generate DSBs, there is so far no clear mechanism explaining
18 how TREX1 could act on nuclear chromatin. ROS and irradiation have been shown to induce
19 TREX1 nuclear translocation, although the precise mechanisms remain elusive (Chowdhury
20 et al., 2006; Martinvalet et al., 2005; Yang et al., 2007). Based on our imaging and rescue
21 experiments, we speculate that TREX1 translocation could potentially implicate fusion of the
22 outer and inner nuclear membranes upon NE ruptures. This could allow ER proteins, by
23 diffusion on the membrane, to gain access to the inner nuclear membrane and potentially to
24 the poorly characterized intranuclear membrane networks of the nuclear reticulum (Malhas et
25 al., 2011). It is likely that the whole ER-bound SET complex transfers to the inner nuclear
26 membrane upon NE rupture events. The SET complex has previously been implicated in the
27 induction of TREX1-dependent nuclear DNA double strand breaks (Chowdhury et al., 2006).
28 In agreement, we did not observe any transient formation of single strand breaks upon
29 confinement, using RPA-GFP, which displays DNA damage foci upon treatment with
30 hydroxyurea (Fig. S7C). We thus speculate that DNA damage after NE rupture could also
31 implicate other members of the SET complex that, together with the TREX1 exonuclease, gain
32 access to inner nuclear membranes due to fusion of the outer and inner nuclear membranes
33 at sites of rupture.

34 In the model systems we studied here, the ATM-dependent DNA damage response pathway
35 is involved both in the senescence phenotype in RPE1 cells, and in the induction of the
36 collagenolytic response in the senescence-deficient DCIS cell line. This is in contrast to the
37 cGAS-dependent inflammatory pathway, which does not seem to be required (Harding et al.,
38 2017; Liu et al., 2018). A potential effect of ATM activation by DNA damage is a cell-cycle
39 arrest. We indeed found that doxo-treated DCIS cells showed a reduction of cells in S/G2 upon
40 48 hours of treatment (Fig. S7D). Nevertheless, this cell cycle arrest *per se* does not explain
41 the increase in collagen degradation as this response was not increased comparing cells in
42 G1 versus S/G2 (Fig. S7E). Two recent studies pointed to a potential activation of an EMT-like
43 phenotype downstream of ATM activation (Jiang et al., 2019; Peng et al., 2019). Furthermore,
44 several ATM substrates have also been associated with lysosomal activity and thus potentially
45 to collagen degradation (Kang et al., 2017). Our results suggest that upon chronic, sub-lethal
46 levels of DNA damage, cells that have lost the normal ATM-dependent senescence-associated
47 proliferation arrest, could initiate a partial EMT program including an invasive behavior, driving
48 the transition from *in situ* to invasive carcinoma stages.

1 A recent study (Shah et al., 2021) found that DNA damage can be triggered by confinement
2 independently of NE rupture events some cell lines, including MDA-MB-231 cells. We thus
3 assayed these cells and found that DNA damage upon confinement was both NE rupture- and
4 TREX1-independent (Fig. S7G, H). These results suggest that tumors might be differentially
5 sensitive to TREX1 inhibition.

6 Our experiments do not prove that TREX1 is the main driver of DNA damage in the human
7 disease nor that DNA damage associated with nuclear deformation plays a role in the human
8 disease progression. TREX1 has been described to promote pro- or anti-tumoral effects
9 depending on different cellular and tumoral contexts (Benci et al., 2019; Erdal et al., 2017;
10 Vanpouille-Box et al., 2017). To get further insight into this question, we analyzed the
11 association of TREX1 expression levels in breast cancer with overall survival using public
12 datasets. This analysis revealed that high level of TREX1 expression is significantly associated
13 with reduced probability of survival (Fig. S7F). Thus, targeting this exonuclease might be an
14 effective therapeutic strategy to curb tumor progression, while promoting anti-tumoral immune
15 response via cGAS activation (Stetson et al., 2008; Takahashi et al., 2018).

16 In this study, we focused on mammary tumors, but nuclear envelope ruptures could also occur
17 in other tumors. As a proof of concept, we performed cGAS staining on colorectal tumor
18 samples. In 2 patients out of 4 tested, we observed a good correlation between regions highly
19 enriched in nuclei positive for γ H2AX, which corresponded to the invasive front of the tumor,
20 and regions with a high fraction of nuclei with NE rupture events (Fig. S7 I, J and K). A broader
21 study is needed to assess whether our findings can be extended to other tumor types.

22 We and others have shown that the NE is a fragile barrier constantly being challenged in
23 physiology and disease (Denais et al., 2016; Raab et al., 2016; Srivastava et al., 2021). The
24 occurrence of DNA damage associated with nuclear deformations is likely to be a common
25 phenomenon. In normal tissues, this would trigger cellular senescence, contributing to
26 developmental programs and tissue homeostasis, or even participating in tissue aging. In
27 contrast, in tumor cells, in which the senescence checkpoint has been lost, it would lead to an
28 aberrant invasiveness in the absence of a proliferative arrest, promoting tumor growth and
29 invasion.

30 **Limitations of the study**

31 Double labelling with fluorescent 53BP1 and icGAS in live cells unambiguously demonstrated
32 the causal relationship between NE rupture events and DNA damage; But the different
33 dynamics of icGAS entry and exit from the nucleus, and DNA damage foci appearance and
34 repair, generate a more complex set of situations when considering cells at a fixed time point,
35 or fixed cells stained with these two proteins by immunofluorescence. This difficulty is
36 exacerbated in 3D samples in which NE rupture events can happen out of the plan of imaging.
37 In cases in which a direct live observation of the rupture events is difficult, for example in slices
38 of tumor samples or in the duct on chip assay, cells in which cGAS perinuclear accumulation
39 and DNA damage are simultaneously present likely represent only a subset of the cells that
40 have undergone NE rupture dependent DNA damage, making precise quantification difficult.

41 **Acknowledgements:**

42 We thank the flow cytometry and cell sorting facility, the animal facility at the Institut Curie and
43 the microscopy facility of IPGG for equipment and technical assistance. We thank Titia de
44 Lange and John Maciejowski for providing the guideRNA to generate the TREX1 KO CRISPR
45 cell lines, John Maciejowski, Alexis Lomakin and Ana-Maria Lennon-Dumenil, for suggestions
46 during the writing of this manuscript, Nicolas Carpi for lab management, plasmid amplifications
47 and computer technical support, Juan-Manuel Garcia Arcos for help in assembling the figures

1 and drawing the graphical abstract, Aastha Mathur for data analysis of the γ H2AX-positive
2 nuclei relative to the xenograft edge, Zahraa Alraies for the preparation of cDNA samples for
3 the quantitative RT-PCR analysis, Dipanjan Bhattacharya who developed the method to
4 automatically segment nuclei at IFOM, Dr. Ines Ferrara for providing technical support for the
5 human tumor samples. Finally, we also thank the patients and their families who contributed
6 tissue samples to these studies. This study was supported by the following grants: Institut
7 Pierre-Gilles de Gennes-IPGG (Equipement d'Excellence, "Investissements d'avenir",
8 program ANR-10-EQPX-34) and laboratoire d'excellence, "Investissements d'avenir" program
9 ANR-10-IDEX-0001-02 PSL and ANR-10-LABX-31; Laboratory of Excellence (LabEx)
10 CelTisPhyBio ANR 11-LABX-0038 to SAG; Plan Cancer 2018 'Single Cells' (19CS007-00) to
11 FR; Fondation pour la Recherche Médicale (FDT20160435078) to CC; Associazione Italiana
12 per la Ricerca sul Cancro (AIRC-IG#18621 to GS, AIRC-0IG#22145 to CT, and 5XMille #22759
13 to GS and CT); the Italian Ministry of University and Scientific Research (MIUR) to GS (PRIN:
14 Progetti di Ricerca di Rilevante Interesse Nazionale – Bando 2017#2017HWTP2K); Institut
15 National du Cancer INCA_7905 (MetaCLEM) and INCA PLBIO 2019-1-PL BIO-07-ICR-1,
16 INSERM Plan Cancer Single Cell grant 19CS007-00, Fondation ARC pour la Recherche
17 contre le Cancer (PGA1 RF20170205408) and institutional funding from Institut Curie and
18 Centre National de la Recherche Scientifique to PC; ERCadg Cello (FP7-IDEAS- ERC-
19 321107) to JLV; LABEX DCBIOL (ANR-10-IDEX-0001-02 PSL and ANR-11-LABX-0043), ANR
20 ANR-17-CE15-0025-01 and ANR-18-CE92-0022-01 to NM; INCA PLBIO 2019-1-PL BIO-07-
21 ICR-1, INSERM Plan Cancer Single Cell grant 19CS007-00 to MP, NM and PC, Italian
22 Foundation for Cancer Research (AIRC) Accelerator Award ID. 24296 to CT. This work was
23 supported in part by grants from Région Ile-de-France.

24 **Author Contributions**

25 G.P.F.N. designed and performed the experiments, analyzed and interpreted the data and
26 wrote the manuscript. S.A-G. and G.P.F.N. performed the 3D collagen degradation assays.
27 S.A-G, A.Y., C.V., P.C. and J-L.V. designed the duct-on-chip device. F.R. and E.L. performed
28 the mouse xenograft assays and S.A-G., F.R. and G.P.F.N. analyzed the data. M.Gr.
29 performed the sequencing of the TREX1 CRISPR clones and generated some stable cell lines.
30 M.M. wrote Image J macros for data analysis and helped on data interpretation. A.W.
31 performed and quantified the experiments with DCIS cells expressing geminin-mCherry. The
32 R.N.R. and J.G. designed and performed multiplex IF and IHC on mice xenografts samples
33 and IHC on human tumor samples. A.P. performed the quantitative RT-PCR analysis. C.C.
34 analyzed data relative to the cell division cycle. M.G. generated stable cell lines and performed
35 molecular cloning. C.L. provided images of mouse xenografts for analysis. V.C. and C.T.
36 performed and interpreted the assays involving human primary tumors. G.S. designed,
37 supervised and interpreted the assays involving human primary tumors and edited the
38 manuscript. P.C. designed, supervised and interpreted the mouse xenograft assays and edited
39 the manuscript. N.M. conceptualized the role of TREX1, helped designing and interpreting
40 experiments, supervised M.Gr. and edited the manuscript. G.P.F.N., S.A-G., M.P., P.C., N.M.
41 and G.S. actively engaged in discussions throughout the study. M.P. supervised the study,
42 designed the experiments, interpreted the data and wrote the manuscript.

43

44 **Declaration of Interest.**

45 Matteo Gentili, Nicolas Manel and Matthieu Piel are authors of a patent entitled 'Method to
46 monitor and quantify interphase nuclear envelope rupture events ', WO2017140875A1

47 **Main Figures Titles and Legends:**

1 **Figure 1. Human breast tumor and mouse xenograft model of breast cancer progression**
2 **display strong nuclear deformation and DNA damage.** (A) Immunohistochemistry of *in situ*
3 advanced human primary breast tumors. RAB5A (red), γ H2AX (brown). Bar, 80 μ m. (B)
4 Magnified region of *in situ* advanced primary breast tumor with a micro-invasive focus. Yellow
5 arrowheads point to deformed nuclei. Bar, 25 μ m. Box and whisker plot showing the median
6 value and 10-90 percentiles of (C) nuclear roundness index or (D) γ H2AX-positive nuclei. (E)
7 Bars are the mean \pm SD of 6 individual tumor samples where at least 150 nuclei were scored
8 per sample. (A-D) Representative of 7 independent tumor samples where 21 and 27 nuclei
9 were analyzed for tumor bulk and micro-invasive focus, respectively, per sample. (F, G)
10 Immunofluorescence analysis of alpha-smooth muscle actin (green), human-specific KU70
11 (cyan) and γ H2AX (red) in 7 weeks old mouse xenografts at different stages generated by
12 intraductal injection of DCIS cells. (F) The *in situ*-advanced stage is characterized by a
13 discontinuous/broken myoepithelial layer. Graph, box and whisker plot showing the median
14 value and 10-90 percentiles of γ H2AX-positive nuclei measured at *in situ*-early and -advanced
15 stages. Data represents 3 independent experiments where 500 nuclei were scored per tumor
16 stage, per experiment. (G) Invasive stage: SMA-positive and elongated cells represent cancer
17 associated fibroblasts. Graph, box and whisker plot showing the median value and 10-90
18 percentiles of the nuclear roundness index measured at invasive and *In situ* stages (divided
19 into edge-up to 2 cell layers below the myoepithelial layer, and bulk). Data represents 3
20 independent experiments where 7 *In situ* and 13 invasive tumor samples were analyzed (for a
21 total of 140 nuclei analyzed for *In situ*-bulk, 139 nuclei analyzed for *In situ*-edge and 264 nuclei
22 analyzed for invasive tumor samples). (F, G) Bar, 100 μ m, (G) Zoom Bar, 50 μ m; Inset Bars,
23 15 μ m. (H) Classification of Z scores categories (γ H2AX intensities) of nuclei as a function of
24 different distances from the tumor edge (determined by the myoepithelial layer) (for details
25 refer to fig. S1A). Data represents a total of 23 images analyzed for DCIS parental tumors from
26 3 independent mice injections. (I) Duct-on-chip assay. Cell density was measured by counting
27 the number of cells/ μ m² in 5 random fields (20x magnification) at the indicated time points and
28 are expressed in the graph relative to the 24h time point. Data represents the mean \pm SD of 3
29 independent experiments. Bar, 50 μ m; Zoom Bar, 25 μ m. (J-L) Duct-on-chip assay. DCIS cells
30 stably expressing 53BP1-mCherry and cGAS-EGFP were injected into the duct-on-a chip
31 device at low and high density. Nuclear roundness (J), DNA damage (K) and NE rupture (L)
32 were assessed. Graphs: (J) Box and whisker plot showing the median value and 10-90
33 percentiles of the nuclear roundness. (K, L) Bars are the mean \pm SD of 3 independent
34 experiments. *P* values were calculated by unpaired Student's t-test, except for items "E" and
35 "K" (One-way ANOVA with post hoc Tukey test) and item "L" (Chi-square test). ****P* < 0.0001;
36 ***P* < 0.005; **P* < 0.05; ns = not significant. See also Fig.S1 and Movie S1

37 **Figure 2. Strong confinement induces NE rupture.** (A) Cells stably expressing 53BP1-
38 EGFP and catalytically inactive cGAS-mCherry (icGAS) were confined and images were taken
39 after 2 h. Bar, 5 μ m. (B) Single cell curves of 53BP1 foci number as a function of time in cells
40 displaying or not NE rupture while under 2 μ m confinement. (C) Quantification of DNA damage
41 level during confinement. Data represents the mean \pm SD of 4 independent experiments where
42 20 cells per experiment per condition were analyzed. (E) Quantification of the percentage of
43 cells displaying more than 3 53BP1 foci/nucleus while confined at 2 μ m. (F) Frequency
44 distribution of 53BP1 foci number/nucleus in cells displaying NE rupture. Data represents 250
45 nuclei pooled from 3 independent experiments. (G) Illustrative panels depicting the
46 appearance and repair of DNA damage foci in cells stably expressing 53BP1-EGFP while
47 confined at 2 μ m. Bar, 5 μ m. (H, I) Frequency of foci appearance and foci life-time (foci
48 duration) in RPE1 cells confined at 2 μ m or treated with etoposide (25 μ M). Box and whisker
49 plot show the median value and 10-90 percentiles of 3 independent experiments where 20
50 cells per condition per experiment were analyzed. (J) Cells stably expressing 53BP1-EGFP

1 and icGAS were plated on glass bottom dishes and imaged for several hours. Arrows point to
2 NE rupture events. (K) Single cell curves showing the relationship between NE rupture and the
3 appearance of 53BP1 foci. (L) Box and whisker plot showing the median value and 10-90
4 percentiles of the ratio between the number of 53BP1 foci after and before NE rupture. Data
5 are from 2 independent experiments where 10 NE rupture events per experiment were
6 analyzed. (M) Cells stably expressing 53BP1-EGFP and icGAS were confined at 2 μm ,
7 harvested and plated for 20 minutes before fixation for immunostaining with antibodies against
8 GFP (left) or γH2AX (right); box and whisker plot show the median value and 10-90 percentiles.
9 *P* values were calculated by unpaired Student's t-test, except for items "H" and "I" (One-way
10 ANOVA with post hoc Tukey test). ****P* < 0.0001; ***P* < 0.005; ns = not significant. See also
11 Fig.S2 and Movies S2, S3

12 **Figure 3. DNA damage upon NE rupture is TREX1-dependent.** (A, B) A variety of TREX1-
13 deficient or -proficient cells was confined at 2 μm and DNA damage level was assessed after
14 2 h under confinement. Graphs represent the mean \pm SD of 3 independent experiments where
15 20 cells were analyzed per condition per experiment. Western blots: TREX1 depletion 48h
16 post-knockdown; tubulin is the loading control. Bar, 10 μm . (C) Box and whisker plot showing
17 the median value and 10-90 percentiles of the ratio between the number of 53BP1 foci after
18 and before NE rupture in siRNA-treated cells. Data are from 2 independent experiments where
19 15 cells per experiment were analyzed. (D) Immuno-gold electron microscopy images of RPE1
20 TREX1 KO cells reconstituted with TREX1-mCherry, confined at 2 μm or left unconfined. Bar,
21 0.5 μm ; magnified regions bar, 0.1 μm . (E) Number of mCherry (TREX1) immune gold particles
22 at the indicated cellular compartments, see also Fig. S5C. The 'cytoplasm' compartment also
23 includes particles associated to cytoplasmic endomembranes like ER and MVBs. *P* values
24 were calculated by One-way ANOVA with post hoc Tukey test, except for item "C", which was
25 calculated by unpaired Student's t-test. **P* < 0.05; ****P* < 0.0001; ns = not significant. In panel
26 E, statistical significance was calculated by chi-square test; *** does not refer to *P* value but
27 significance according to the chi-square statistics (see Table S1). See also Fig.S3, S4 and
28 Movie S4

29 **Figure 4. Long-term, strong confinement causes TREX1-dependent cell senescence.** (A)
30 Cells stably expressing catalytically inactive cGAS were confined at 2 μm for 3 or 12 h. Cells
31 were then harvested and replated for imaging for cell cycle duration assessment in cells
32 displaying or not NE rupture. Graph, left: box and whisker plot showing the median value and
33 10-90 percentiles of the cell cycle duration. Data for 3 h of confinement represents 3
34 independent experiments where a total of 150 cells with non-ruptured and 200 cells with
35 ruptured nuclei were analyzed. Data for 12 h of confinement represents 3 independent
36 experiments where a total of 23 lineages with non-ruptured and 40 lineages with ruptured
37 nuclei were analyzed. Right: Frequency distribution of the number of cell divisions. (B, C, F,
38 G) siRNA-treated cells were confined, then harvested, replated and cultured for 72 h before
39 fixation for β -gal staining (B, C) or immunostaining of heterochromatin foci (HP1 γ) (F, G). (C)
40 Data represents the mean \pm SD of 3 independent experiments where 200 cells were scored
41 per condition per experiment. (D) Cells were confined for 3 or 12 h at 4 or 2 μm ; cells were
42 then harvested and replated for 72 h before lysis for western blot; tubulin is the loading control.
43 (E) RPE1 TREX1 KO cells or RPE1 TREX1 KO cells reconstituted with TREX1 WT-mCherry
44 were confined at the indicated heights for 12 h. Cells were then harvested, replated and
45 cultured for 72 h before lysis for western blot; tubulin is the loading control. Western blots are
46 representative of 2 independent experiments. Bars, 80 μm (B) and 10 μm (F). *P* values were
47 calculated by unpaired Student's t-test (A) or One-way ANOVA with post hoc Tukey test (C,
48 G). ****P* < 0.0001; ***P* < 0.005; ns = not significant. See also Fig.S5

1 **Figure 5. TREX1-dependent DNA damage leads to collagen degradation and invasion.**
2 (A, B, C) Duct-on-chip assay. (A) DCIS cells stably expressing 53BP1 and catalytically inactive
3 cGAS were transiently depleted for TREX1 and 48 h later injected in the chip device. Graphs,
4 data represents the mean \pm SD of 3 independent experiments where 5 random fields (at 20x
5 magnification) were scored per condition per experiment. Bar, 50 μ m; zoom Bar, 10 μ m. (B)
6 DCIS cells were transiently depleted for TREX1 and 48 h later injected in the same chip device
7 with siControl cells (labeled with a green cell tracker). Graphs, (top) data represents the mean
8 \pm SD of 4 independent experiments where cells invading from 7 random gates were scored
9 per experiment; (bottom) frequency distribution histogram of the data displayed on the bar
10 graph. Bar, 50 μ m. (C) DCIS cells stably expressing 53BP1-EGFP were pre-treated for 12 h
11 with doxorubicin (Doxo, 20 nM) and co-injected with DCIS cells stably expressing 53BP1-
12 mCherry pre-treated for 12 h with DMSO. Data represents the mean \pm SD of 2 independent
13 experiments where cells invading from 10 random gates were scored per experiment. Bar, 40
14 μ m. (D) DCIS cells were confined for 2 h. Following the confinement period, cells were
15 harvested and embedded in 3D type I collagen for 12 h. Degraded collagen (red); DAPI (cyan).
16 (E) A variety of TREX1-deficient or -proficient DCIS cells was confined at the indicated heights.
17 Data represents 3 independent experiments where 50 cells per condition per experiment were
18 analyzed. (F) It was proceeded exactly as described in "D" except that the experiment was
19 performed with TREX1-depleted DCIS cells. Data represents 3 independent experiments
20 where 60 cells per condition per experiment were analyzed. Data was normalized by the mean
21 of control samples. (G) It was proceeded exactly as described in "D" except that cells were
22 treated with ATM inhibitor (KU-55933, 10 μ M) or MMP inhibitor (GM6001, 40 μ M) while under
23 confinement and in 3D collagen. Data represents 3 independent experiments where 60 cells
24 per condition per experiment were analyzed. All box and whisker plots show the median value
25 and 10-90 percentiles of the collagen degradation area. *P* values were calculated by unpaired
26 Student's *t*-test (F), Chi-square test (A-right, B, C) or One-way ANOVA with post hoc Tukey
27 test (E, G). ****P* < 0.0001; ns = not significant. See also Fig.S6 and Movie S5

28 **Figure 6. TREX1-dependent DNA damage promotes tumor invasion in mouse xenograft**
29 **model of breast cancer.** (A) Left: Immunofluorescence analysis of alpha-smooth muscle
30 actin, SMA (green) and DAPI (cyan) in 6-8 weeks old mouse xenografts generated by
31 intraductal injection of either parental DCIS cells or TREX1 KO clones. Bar, 100 μ m. Middle:
32 whole-mount carmine-stained 7 weeks-old glands. Bar, 1 mm. Right: Hematoxylin & Eosin
33 stained 7 weeks-old glands. Bar, 60 μ m. (B) Classification of the tumor xenograft stage 6-8
34 weeks post-intraductal injection of either parental DCIS cells or TREX1 KO clones from 3
35 independent mice injections. Data represents a total of 7 glands scored for DCIS parental
36 tumors and 14 glands for TREX1 KO tumors (TREX1 KO #1 +TREX1 KO #2 clones). (C)
37 Scatter dot plots of γ H2AX- and Ki67-positive nuclei (lines are median) 6-8 weeks post-
38 intraductal injection. A total of 13 parental and 34 TREX1 KO (18 TREX1 KO #1 + 16 TREX1
39 KO #2) images were analyzed for Ki67 scoring; a total of 31 parental and 62 TREX1 KO (35
40 TREX1 KO #1 + 27 TREX1 KO #2) images were analyzed for γ H2AX scoring. (D)
41 Immunofluorescence analysis of γ H2AX (red), alpha-smooth muscle actin (green) and DAPI
42 (cyan) in 6-8 weeks old mouse xenografts generated by intraductal injection of either parental
43 DCIS cells or TREX1 KO clone. Blue arrowheads point to DNA damage- γ H2AX-positive cells;
44 yellow arrowheads point to mitotic or apoptotic figures. Bar, 40 μ m. (E, F) Z score categories
45 (γ H2AX intensities, see Methods and fig. S1A) of nuclei localized in the tumor bulk (E) or in
46 the tumor edge (determined by the myoepithelial layer) (F). Data represents a total of 23
47 images analyzed for DCIS parental tumors and 30 images for TREX1 KO tumors (TREX1 KO
48 #1 +TREX1 KO #2 clones) from 3 independent mice injections. *P* values were calculated by
49 unpaired Student's *t*-test, ****P* < 0.0001; **P* < 0.05; ns = not significant. See also Fig.S6

1 **Figure 7. Mouse tumor xenografts display NE rupture and DNA damage.** (A)
2 Immunofluorescence analysis of laminA/C (green), DAPI (cyan) and γ H2AX (red). Bar, 25 μ m;
3 inset bars, 5 μ m (single channels) and 3 μ m (merged channels). (B) Multiplex
4 immunofluorescence analysis of cGAS (green), DAPI (cyan) and γ H2AX (red). Bar, 60 μ m. (C)
5 cGAS immunohistochemistry (brown) and nuclei (blue). Bar, 25 μ m. All analysis was
6 performed in mouse xenografts at *in situ*-advanced stage generated by intraductal injection of
7 DCIS cells. Yellow arrowheads point to NE rupture events. See also Fig. S7
8

1 STAR Methods

2 RESOURCE AVAILABILITY

3 Lead Contact

4
5 Further information and requests for resources and reagents should be directed to and will
6 be fulfilled by the lead contact, Matthieu Piel (matthieu.piel@curie.fr)

7 Materials Availability

8 All newly generated materials associated with this study (stable cell lines and plasmids,
9 design of microfabricated devices) will be provided upon request to the lead contact.

10 Data and Code Availability

- 11 • All data reported in this paper will be shared by the lead contact upon request
- 12 • All original code (Image J macros) has been deposited at Mendely data and is
13 publicly available as of the date of publication. DOIs are listed in the key resources
14 table.
- 15 • Any additional information required to reanalyze the data reported in this manuscript
16 is available from the lead contact upon request.

17 EXPERIMENTAL MODEL AND SUBJECT DETAILS

18 Mice

19 Animal care and use for this study were performed in accordance with the recommendations
20 of the European Community (2010/63/UE) for the care and use of laboratory animals. All
21 animal experimental procedures were approved by ethics committee of Institut Curie CEEA-
22 IC #118 (Authorization APAFiS #24649-2020031220398710-v1 given by National Authority) in
23 compliance with international guidelines. As research was limited to the mammary gland, all
24 transplanted animals were female mice.

25 *Intraductal transplantation method*

26 Eight to ten week-old virgin female CB17/Icr-Prkdc^{scid}/IcrIcoCrl mice (Charles River
27 Laboratories) were anesthetized by inhalation with isoflurane and intraductally injected into the
28 3rd and the 4th pair of glands with 2 μ l PBS containing 50,000 MCF10DCIS.com cells as
29 previously described (Behbod et al., 2009; Lodillinsky et al., 2016). Five to eight weeks after
30 injection with no additional manipulation, mice were sacrificed and the engrafted mammary
31 glands were harvested and processed for whole-mount and histological and IHC staining on
32 tissue sections.

33 Human tissues

34 Human breast cancer tissue samples were collected according to the Helsinki Declaration and
35 the study was approved by the University of Palermo Ethical Review Board (approval number
36 09/2018). The cases were classified according to the World Health Organization classification
37 criteria of the tumors of the breast. All the tissue samples were from 7 adult female patients
38 with breast ductal adenocarcinoma with an age range between 40 and 88 years old (histotypes
39 *in situ* ductal adenocarcinoma, invasive ductal adenocarcinoma or both).

40 Human Cell lines

1
2 Cell lines are described in the *Key Resources Table*. MCF10DCIS.com and MDA-MB-231 cell
3 lines were kindly provided by the lab of Dr. Philippe Chavrier (Institut Curie, Paris, France).
4 MCF10DCIS.com cells were culture in Advanced DMEM- F12 (GIBCO) medium supplemented
5 with 2 mM glutamine, 5% horse serum and 1% penicillin and streptomycin (GIBCO #15140-
6 122). MDA-MB-231 was maintained in L-15 culture medium (Sigma-Aldrich, St Louis, MO,
7 USA) with 2mM glutamine (GIBCO, Cergy Pontoise, France) and 15% fetal bovine serum
8 (GIBCO). MCF10A cells were culture in DMEM-F12 (GIBCO) supplemented with EGF
9 (20ng/ml) (Peprotech #AF-100-15), hydrocortisone (0.5µg/ml) (SIGMA #H0888), cholera toxin
10 (100ng/ml) (SIGMA #C8052), Insulin (10µg/ml) (SIGMA #I9278), 5% horse serum and 1%
11 penicillin and streptomycin. RPE1 cells were grown in DMEM-F12 Glutamax medium (GIBCO),
12 supplemented with 10% FBS and 1% penicillin and streptomycin (GIBCO). All cells were
13 maintained at 37 °C in 5% CO₂, with the exception of MDA-MB-231 cells, which were
14 maintained at 37 °C in 1% CO₂.

15 16 **METHOD DETAILS**

17 18 **Constructs**

19
20 The following constructs were kindly provided by the laboratory of Dr. Nicolas Manel (Institut
21 Curie, Paris, France): pTRIP-CMV-mCherry-TREX1 WT and D18N were cloned from pEGFP-
22 C1-TREX1 (Addgene #27219). pTRIP-SFFV-EGFP-FLAG-cGAS E225A/D227A and pTRIP-
23 CMV-mCherry-FLAG-cGAS E225A/D227A are described elsewhere (Gentili et al., 2019;
24 Raab et al., 2016). pTRIP-SFFV-EGFP-53BP1 (amino acids 1224-1716 for isoform 1) was
25 obtained by cloning from pTRIP-CMV-mCherry-53BP1 (Gentili et al., 2019). The following
26 constructs were provided byb the laboratory of Dr. John Maciejowski (MSKCC, New York,
27 USA): pLENTI-CMV-GFP-TREX1 deltaC (aa1-235), pLENTI-CMV-GFP-TREX1 deltaC-
28 sec61TM and pQCXIP-GFP-RPA70 (Mohr et al., 2021).

29 30 **Lentiviral particles production in 293FT cells and Lentivector transductions**

31
32 Lentiviral particles were produced as previously described from 293FT cells (Lahaye et al.,
33 2013). Lentiviral viral particles and viral-like particles were produced by transfecting 1µg of
34 psPAX2 and 0.4µg of pCMV-VSV-G together with 1.6µg of a lentiviral vector plasmid per well
35 of a 6-well plate. For cell transduction, 0.5x10⁶ cells were plated in a 6 well plate in 1ml and
36 infected with 2ml of fresh lentivector in the presence of 8µg/ml of Protamine.

37 38 **Quantitative RT-PCR analysis**

39
40 Total RNA was extracted using the RNeasy Mini kit (Qiagen) and quantified by NanoDrop to
41 assess both concentration and quality of the samples. Reverse transcription was performed
42 using the SuperScript VILO cDNA Synthesis kit from Invitrogen. Gene expression was
43 analysed by using the TaqMan Gene expression Assay (Applied Biosystems). 0.1 ng of cDNA
44 was amplified, in triplicate, in a reaction volume of 25 µl with 10 pmol of each gene-specific
45 primer and the SYBR Green PCR MasterMix (Applied Biosystems). Real-time PCR was
46 performed on the 14 ABI/Prism 7700 Sequence Detector System (PerkinElmer/Applied
47 Biosystems) using a pre-PCR step of 10 min at 95 °C, followed by 40 cycles of 15 s at 95 °C
48 and 60 s at 60 °C. Specificity of the amplified products was confirmed by melting curve analysis
49 (Dissociation Curve TM; Perkin Elmer/Applied Biosystems) and by 6% PAGE. Preparations
50 with RNA template without reverse transcription were used as negative controls. Samples were
51 amplified with primers for each gene (for details see the quantitative PCR primer list below)
52 and GAPDH as a housekeeping gene. The cycle threshold C_t values were normalized to the
53 GAPDH curve. PCR experiments were performed in triplicate and standard deviations
54 calculated and displayed as error bars.

cGAMP extraction with methanol/water

Cells were seeded and treated the day after with HT-DNA (1µg/ml) for 7h. 1.5×10^6 cells were recovered and washed with PBS. Cell pellets were resuspended in 80% methanol and frozen at -80°C. To perform cGAMP extraction, pellets were frozen/thawed five times in liquid nitrogen (with vortexing steps between each cycle). Then the samples were centrifuged at 16000g for 20 min at 4°C. Supernatants were recovered and dried in a speed vacuum drying in Savant DNA Speed Vac DNA 110 for 3 h at 65°C. Pellets were resuspended in 150µl of DNase/RNase sterile water. cGAMP concentration was obtained with the 2'3'-cGAMP ELISA Kit (Cayman Chemical) according to the manufacturer instruction. cGAMP quantities were normalized to the number of cells extracted.

Generation of TREX1 CRISPR clones

The guideRNA (gRNA) targeting TREX1 was kindly provided by the lab of Titia De Lange (Rockefeller University, New York, USA): gTREX1, 5'-GAGCCCCCCCCACCTCTC-(PAM)-3'. gRNA plasmid was co-transfected into target cells with an hCas9 expression plasmid (Addgene) by nucleofection (Lonza apparatus). 1×10^6 cells were mixed with electroporation buffer (freshly mixed 125 mM Na₂HPO₄, 12.5 mM KCl, 55 mM MgCl₂ pH 7.75), 2 µg Cas9 plasmid, and 2 µg gRNA plasmid, transferred to an electroporation cuvette (BTX), and electroporated with program U-017 for RPE-1 cells or program D-023 for MCF10DCIS.com cells. Cells were then allowed to recover for 48 h before single cell FACS sorting into single wells of a 96 wells plate for single clone amplification and selection. Successful CRISPR/Cas9 editing was confirmed by both Western blotting of the amplified cell colonies and analysis of indel size (using the ICE analysis toolbox (<https://www.synthego.com/products/bioinformatics/crispr-analysis>)).

Duct-on-a-chip assay

Unlabeled type I collagen at 2.4 mg/ml (BD Biosciences, Cat. # 354236) was injected in the "collagen chamber" of the microfabricated duct-on-a-chip and incubated for 1h at 37°C. In the meantime cells were resuspended in matrigel (SIGMA #E1270) at a concentration of 0.7×10^6 cells/µl and then 3 µl were injected in the "cells chamber". Images were acquired at the desired time points after injection to assess cell invasion into the "collagen chamber".

Human primary tumor samples

Sections 2.5/3 micron-thick were cut from paraffin blocks, dried, de-waxed and rehydrated. The antigen unmasking technique was performed using Target Retrieval Solutions pH6 at 98°C for 30 min. After neutralization of the endogenous peroxidase with 3% H₂O₂ and Fc blocking by a specific protein block, double-marker immunohistochemistry was carried out by incubation for 90 min at RT with the primary antibodies RAB5A (Abcam, 1:100, pH6) and γH2AX (Abcam, 1:1000, pH6). Staining was revealed using Novolink Polymer Detection Systems (Novocastra) and SuperSensitive Link-Label IHC Detection System Alkaline Phosphatase (Biogenex). DAB (3,3'-diaminobenzidine) and Vulcan Fast Red were used as substrate chromogens. The slides were counterstained with Harris hematoxylin (Novocastra). All the sections were analyzed under a Zeiss AXIO Scope.A1 microscope (Zeiss, Germany) and microphotographs were collected using a Zeiss Axiocam 503 Color digital camera using the Zen2 imaging software.

Immunohistochemistry and spectral multiplex (processed using Bond RX automated-Leica)

1 Antibody reference: cGAS (D1D3G) Rabbit mAb #15102; Cell Signaling, Dilution: 1/50. Antigen
2 retrieval: BOND Epitope Retrieval Solution 2 (Leica, AR9961; pH9) for 20 min; protein blocking:
3 5 min (protein block serum-free Dako #X0909); incubation: 1h at room temperature

4 For DAB staining, once immunostaining finished, slides were counterstained with hematoxylin
5 before mounting with glycergel medium (Dako). Images were acquired using Digital Pathology
6 slide scanner (Nanozoomer 2.5, Hamamatsu). For multiplex IF, immunostaining was
7 processed with Opal™ 7-Color IHC Kits (Akoya 23 Biosciences, NEL821001KT). HRP labelled
8 Rabbit Polymer and TSA opal fluorophores were incubated for 10 min each. Subsequently,
9 slides were submitted to DAPI staining for 10 min, washed and coverslipped with Prolong™
10 Diamond Antifade Mountant (ThermoFisher) before image acquisition.

11 **Histological and immunofluorescence analysis of mouse tissue sections**

12
13 Whole-mount carmine and hematoxylin and eosin staining were performed as described
14 (Teuliere, 2005). To retrieve antigens on paraffin-embedded tissue samples, sections were
15 incubated for 20 min in 10mM sodium citrate buffer, pH 6.0 at 90°C. Then, after 1h incubation
16 in 5% fetal calf serum, sections were incubated overnight with diluted primary antibodies,
17 washed and further incubated for 2h at room temperature with appropriate secondary
18 antibodies.

19

20 **Indirect immunofluorescence microscopy**

21
22 Samples were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton X-100 for 5
23 min and then incubated for 1h at room temperature with primary antibodies. Following three
24 washes with PBS, samples were incubated with the appropriate secondary antibodies for 1h
25 at room temperature. Samples were again washed three times with PBS and mounted with
26 Fluoromount G (Molecular probes).

27

28 **3D type I collagen degradation assay**

29
30 For collagen degradation assessment followed by immunofluorescence analysis, glass bottom
31 dishes (MatTek Corporation) were layered with 15 µl of a solution of 5 mg/ml unlabeled type I
32 collagen (IBIDI Cat. # 50201) mixed with 1/20-40 volume of Alexa Fluor 647-labeled collagen.
33 Polymerization was induced at 37°C for 3 min and complete medium was added. Cells were
34 seeded onto the polymerized collagen layer and incubated for 1h at 37 °C. The medium was
35 then gently removed and two drops of a mix of Alexa Fluor 647-labeled type I collagen (10%
36 final) and unlabeled type I collagen at 2.4 mg/ml (BD Biosciences, Cat. # 354236) were added
37 on top of the cells (top layer). After polymerization at 37 °C for 1.5h, 1 ml of medium added to
38 the MatTek dishes.

39

40 **Transfection procedure and siRNA oligonucleotides**

41
42 For RNA interference experiments, cells were transfected with siRNA oligonucleotides
43 (Dharmacon) using Lipofectamine® RNAiMAX reagent (Invitrogen) (RPE1 cells), or with
44 Lullaby (OZ Biosciences, France) (MCF10A cells, MCF10DCIS.com cells and MDA-MB-231
45 cells) according to manufacturer's protocol. The following SMARTpool siRNAs were used:
46 human TREX1 (Dharmacon, cat. # L-013239-02-0005), human MB21D1 (cGAS) (Dharmacon,
47 cat. # L-015607-02-0005) and validated non-targeting siRNAs (Dharmacon, cat. # D-001810-
48 10-20). Cells were analyzed 72 h post-transfection using standard Western blot or
49 immunofluorescent analysis protocols.

50

51 **Drug treatments**

52
53 The following pharmacological inhibitors and chemical compounds were used: Etoposide
54 (DNA damaging agent, topoisomerase-II inhibitor; Cell Signaling, #2200), Doxorubicin (DNA

1 damaging agent, topoisomerase-II inhibitor; Cell Signaling, #5927), GM6001 (pan-inhibitor of
2 matrix metalloproteinases; Millipore, # CC1010), KU-55933 (ATM kinase inhibitor; Abcam,
3 #120637).

4 5 **Western blotting**

6
7 Cells were collected and resuspended in Laemmli buffer. Proteins were separated using
8 sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE, 4-20%) and
9 transferred onto nitrocellulose or PVDF membranes. Membranes were blocked with 5% BSA
10 in Tris-buffered saline containing 0.05% Tween (TBS-Tween) for 1 h at room temperature,
11 after which primary antibodies were added in TBS-Tween 5% BSA and incubated over night
12 at 4°C. The membranes were subsequently washed in TBS-Tween after which secondary
13 horseradish peroxidase conjugated antibodies (Jackson ImmunoResearch) were added in 5%
14 BSA in TBS-Tween and incubated for 1 h at room temperature. After extensive washing in
15 TBS-Tween ECL signal was recorded on a ChemiDoc Touch Biorad Imager.

16 17 **Microfabrication-based confinement**

18 19 *Confinement using the “6 well confiner”:*

20 To obtain large quantities of confined cells for cell population or biochemical studies, cell
21 confinement was performed using a version of the cell confiner adapted to multi-well plates
22 (Liu et al., 2015). To make the polydimethylsiloxane (PDMS, RTV615, GE) microspacers
23 (micropillars) at the desired height, 12 mm glass coverslips were plasma treated and then
24 placed on top of a PDMS mixture (10/1 w/w PDMS A / crosslinker B) on the wafer molds
25 (containing holes/micropillars, fabricated following standard photolithography procedures).
26 The height of the micropillars determines the height for spatial confinement of cells between
27 the coverslip and the substrate. The surface of the confining side was always treated with non-
28 adhesive pLL-PEG (SuSoS, PLL(20)-g[3.5]-PEG(2)). After baking at 95°C for 15 min,
29 coverslips with PDMS pillars were carefully removed from the wafers under isopropanol. They
30 were then cleaned with isopropanol, well-dried, treated with plasma for 1 min, and treated with
31 0.5 mg/mL pLL-PEG in 10 mM pH 7.4 HEPES buffer for 1h at room temperature. Coverslips
32 with PDMS pillars were rinsed and incubated in medium for at least 2 hours before confining
33 the cells. The modified cover-lid of a multi-well plate was used to apply confining slides to cells.
34 In this case, large PDMS pillars were stuck on the cover-lid of the multi-well plate to hold the
35 confining slides containing a layer of microfabricated micropillars. These large PDMS pillars
36 push the confining slides from the top of the plastic 6 well cover-lid to confine the cells in 6 well
37 glass/plastic bottom plates. The process of fabrication for these large pillars attached to the 6
38 well plate lid is as follows: the large PDMS pillars were fabricated by pouring a PDMS mixture
39 (A:B = 35:1) into a custom-made metallic mold, removing bubbles under vacuum, then baking
40 overnight at 80°C, and getting the pillars out of the mold with the help of small amount of
41 isopropanol.

42 43 **Live-cell imaging**

44
45 Time-lapse recordings were acquired with 20x (NA 0.75) or 40x (NA 0.85) dry objectives using
46 either an Eclipse Ti inverted microscope (Nikon) equipped with a Coolsnap HQ2 camera
47 (Photometrics) or with an spinning-disc confocal microscope with a Yokogawa CSU-X1
48 spinning-disc head on a DMI-8 Leica inverted microscope equipped with a Hamamatsu
49 OrcaFlash 4.0 Camera, a NanoScanZ piezo focusing stage (Prior Scientific) and a motorized
50 scanning stage (Marzhauser). Both microscopes were controlled by MetaMorph software
51 (Molecular Devices). All microscopes were equipped with an on-stage incubation chamber
52 which maintained the temperature at 37°C and CO₂ concentration at 5% at all times. Image
53 analysis was performed using ImageJ/Fiji software (NIH, <http://rsb.info.nih.gov/ij/index.html>) or
54 MetaMorph software (Universal Imaging).

1 **Measurements of γ H2AX intensity relative to tumor edge in mice xenografts**

2
3 For semi-automated analysis of nuclear γ H2AX intensity in tissues as a function of distance
4 from tumor edge, a macro was written and run on Fiji software. First, individual nuclear regions
5 were obtained after thresholding using the DAPI channel. Subsequently, the tumor edge was
6 manually defined on the image using the myoepithelial layer as the reference. Next, a map of
7 the distance to the tumor edge was obtained. Finally, γ H2AX intensity and the distance to the
8 edge was measured for each nuclei of the tumor xenograft. Cells with strongly condensed
9 chromatin (as assessed by DAPI staining) were excluded from the analysis as they could be
10 either mitotic or apoptotic cells. Due to intensity variation between experiments, γ H2AX
11 intensities were normalized using Z scores. For each image, the distribution of the intensities
12 of nuclei inside the tumor ($>30\mu\text{m}$ from the myoepithelial layer) and at the tumor edge ($<30\mu\text{m}$
13 from the myoepithelial layer) were fit using a Gaussian function to obtain an average nuclear
14 intensity and a standard deviation. Thus, individual nuclei intensities were normalized as: Z
15 score = (Nuclear Intensity - Average Nuclear Intensity) / Standard Deviation, to compare all
16 experiments.
17

18 **Electron microscopy**

19 Sample preparation, ultrathin cryo-sectioning and immunolabelling were performed as
20 previously described (Slot and Geuze, 2007). Briefly, RPE1 TREX1 KO + TREX1-mCherry
21 were confined at $2\mu\text{m}$ for 30 min. Next, the confinement lid was removed (in order to have
22 enough number of cells for sample preparation, a total of twelve 6well plates were confined,
23 cells were harvested and pooled together). Cells were harvested and processed for fixation for
24 2h with 2% formaldehyde + 0.125% Glutaraldehyde in 0.1 M phosphate buffer pH 7.4. The
25 fixative was the replaced by PFA 1% in 0.1 M phosphate buffer and sample was stored at 4°C
26 in a sealed tube. For the preparation of the blocks cells were rinsed with PBS, the blocks were
27 embedded in 12% gelatin, cryoprotected with 2.3 M sucrose, and frozen in liquid nitrogen.
28 Ultrathin cryo-sections were cut on a Leica ultracut UC7 cryo-microtome and picked up in a
29 freshly prepared 1:1 mixture of 2.3 M sucrose and 1.8% methylcellulose, thawed and collected
30 on formvar-coated grids. After washing with PBS containing 0.02 M glycine, sections were
31 incubated with primary antibody (anti-mCherry; GeneTex #GTX 128508 used at 1:250 dilution
32 and anti-PDI; EnzoLife Sciences ADI-SPA-891) and protein A-gold conjugates (PAG) (Utrecht
33 University, The Netherlands). Sections were examined using a Tecnai Spirit electron
34 microscope (FEI Company) equipped with a digital camera Quemesa (SIS). Quantitative
35 Immuno EM analysis was used as described by (Mayhew, 2011). *Approach 1: Testing for shifts*
36 *in compartment labelling in different groups of cells.* This approach is used to test whether the
37 distribution of gold particles across compartments alters between not confined and $2\mu\text{m}$
38 confined cells. The observed numerical frequency distributions of raw gold counts in different
39 groups of cells are compared directly by contingency table analysis. For a particular
40 compartment in a given cell, the number of expected gold particles is calculated by multiplying
41 the corresponding column sum by the corresponding row sum and then dividing by the grand
42 row sum (see Table S1). *Approach 2: Relative labelling index analysis, testing for differences*
43 *in specific-labelling enrichment of TREX1-mCherry gold particles between cellular*
44 *compartments.* Labelling density values relate numbers of gold particles to the sizes of
45 compartments. For the relative labelling index analysis, a total of four compartments, including
46 "nucleus", "peripheral ER", "inner nuclear membrane" and "outer nuclear membrane" were
47 selected to analyze the intracellular distribution of TREX1-gold particles. The expected
48 distribution was obtained by superimposing to pictures an array of points that was generated
49 digitally, and points (P) were counted in the selected compartments. Pictures were taken
50 randomly with the only criterion of a well-preserved morphology. Gold particles on all sampled
51 fields were counted and named as observed gold particles (N_{go}). For each compartment, N_{go}
52 was compared with the expected numbers of gold particles (N_{ge} , derived from the observed
53 frequencies of points P). Labelling density (LD) is calculated as the number of gold particles
54 per test point ($LD = N_{go}/P$). For each compartment, relative labelling index $RLI = LD$

1 *compartment/LD cell*. $RLI = 1$ indicates random labeling while $RLI > 1$ indicates when
2 compartments are preferentially labeled. By means of a two-sample Chi-squared (χ^2) analysis
3 with two columns (N_{go}) and (N_{ge}) and c compartments (arranged in rows), two distributions
4 were compared, the total and partial χ^2 values were calculated, and whether to accept or reject
5 the null hypothesis was decided (of no difference between distributions) for $c-1$ degrees of
6 freedom. For any given compartment, the partial χ^2 is calculated as $(N_{go}-N_{ge})^2/N_{ge}$. If the
7 observed and expected distributions are different, examining the partial χ^2 values will identify
8 those compartments that are mainly responsible for that difference. A convenient arbitrary cut-
9 off is a partial χ^2 value accounting for 10% or more of total (Mayhew, 2011).

10

11 **QUANTIFICATION AND STATISTICAL ANALYSIS**

12

13 Statistical analyses for all experiments were performed in Prism (GraphPad) v5.02. Statistical
14 data are presented as median or mean \pm SEM, SD or 10-90 percentiles. Sample size (n),
15 statistical tests used and p-values are specified in the figure legends. Samples in most cases
16 were defined as the number of cells counted/examined within multiple different fields of view
17 on the same dish/slide, and thus represent data from a single sample within a single
18 experiment that are representative of at least three additional independently conducted
19 experiments.

20

1 Supplementary Figures Titles and Legends

2 **Figure S1. The “duct-on-chip” device: reconstitution of the mammary duct using**
3 **microfabrication (related to Figure 1)** (A) Schematic showing the nuclei segmentation and
4 Z score transformation for the normalization of γ H2AX intensities as a function of the distance
5 from the tumor xenograft edge (defined by the myoepithelial layer). Note that the great majority
6 of nuclei with Z score > 2 are detected within the first 30 μm from the tumor edge/myoepithelial
7 layer. Cells with strongly condensed chromatin (as assessed by DAPI staining) were excluded
8 from the analysis as they could be either mitotic or apoptotic cells. Zoom bars, 5 μm . (B)
9 Density scatter plot of Z scores (γ H2AX intensities) of DCIS parental tumors. Data represents
10 a total of 23 images analyzed from 3 independent experiments (mice injections). (C) Diagram
11 illustrating the duct-on-chip device. A mixture of cells and matrigel is injected in the “cell
12 chamber” and type I collagen is injected in the collagen chamber. Subsequently, culture
13 medium is injected in the “medium chamber”. Note that cells invade towards the collagen
14 chamber through 10 μm wide gates that recapitulate the breaches in the myoepithelial layer
15 during *in vivo* of cell invasion. Bar, 80 μm . (D) DCIS cells were injected in the duct-on-chip in
16 the presence of DMSO (vehicle) or MMP pan inhibitor (GM6001, 40 μM). Cell invasion was
17 measured as distance from the gates. (E) Frequency distribution of invading cells as a function
18 of the distance invaded from the gate. Data represents 3 independent experiments where cells
19 invading from 10 random gates (at 20x magnification) were analyzed per condition per
20 experiment. Bar, 60 μm . (G-I) Duct-on-chip assay. DCIS cells stably expressing 53BP1-
21 mCherry and cGAS-EGFP were injected in the chip device at low (G) and high (H, I) densities.
22 Images were acquired at the indicated time points. (I) Nuclei displaying both NE rupture events
23 (discrete cGAS accumulation at the nuclear border) and augmented 53BP1 foci are shown.
24 Bars, 50 μm (F, G); 5 μm (H); zoom bars 20 μm (F), 10 μm (G). *P* value was calculated by Chi-
25 square test in “E”.

26

27 **Figure S2. Characterization of nuclear envelope rupture under different confinement**
28 **heights and of DNA damage following nuclear envelope rupture (related to Figure 2).** (A)
29 RPE1 cells stably expressing catalytically inactive cGAS-EGFP (icGAS) were confined at the
30 indicated heights and images were acquired immediately after (and while cells were under
31 confinement). Bar, 15 μm . (B) Quantification of NE rupture events as assessed by cGAS
32 perinuclear localization. Data represents the mean \pm SD of 3 independent experiments where
33 50 cells per experiment per height were analyzed. (C) Frequency distribution of 53BP1-EGFP
34 (RPE1 cells, left) and 53BP1-mCherry (DCIS cells, right) mean intensities. (D) DCIS and
35 MCF10A cells stably expressing 53BP1-mCherry and catalytically inactive cGAS-EGFP were
36 confined at 2 μm . Top: examples of cells displaying or not NE rupture events (note perinuclear
37 cGAS accumulation on cells displaying NE rupture). Bar, 10 μm . Bottom: quantification of DNA
38 damage levels following 2 μm confinement as assessed by the number of 53BP1 foci in cells
39 displaying or not NE rupture. Data represents the mean of 3 independent experiments
40 (MCF10A cells) or 2 independent experiments (DCIS cells) where 20 cells per experiment per
41 condition were analyzed. (E) Top: curves showing the 53BP1-EGFP mean intensity of RPE1
42 cells while under confinement at the indicated heights; data represents 4 independent
43 experiments where 20 cells per experiment and per condition were analyzed. Bottom: curves
44 showing the 53BP1-EGFP total intensity (blue) and the average nuclear background intensity
45 (red) of RPE1 cells while under confined at 2 μm . Data are pooled from 3 independent
46 experiments where a total of 50 cells per condition were analyzed. Bars are SD. (F) RPE1 cells
47 stably expressing catalytically inactive cGAS-EGFP (icGAS) were confined at 2 μm for 2 h;
48 subsequently, the confinement lid was removed, cells were harvested, replated for 20 minutes
49 and fixed for immunostaining with the DNA damage markers RIF1 and γ H2AX (red) and DAPI

1 (blue). Arrowheads point to cells with NE rupture, which are also positive for the DNA damage
2 markers analyzed. Bar, 20 μm . (G) Quantification of DNA damage foci at the indicated
3 conditions (DMSO, etoposide 25 μM and 2 μM , NE rupture). Images are representative of 2
4 independent experiments where 30 cells per experiment per condition were analyzed. Bars
5 represent the mean \pm SD. (H) DCIS cells stably expressing Geminin-mCherry and 53BP1-
6 EGFP were confined at 2 μm and the number of 53BP1 foci was assessed; box and whisker
7 plot shows the median value and 10-90 percentiles of 3 independent experiments where 20
8 cells per experiment per condition were analyzed. (I) RPE1 cells stably expressing 53BP1-
9 EGFP were confined at 2 μm and imaged under spinning disc microscopy. White arrowhead
10 points to a non-ruptured NE bleb while the yellow arrowhead points to a bleb bursting event
11 that is followed by the appearance of DNA damage foci. (J) Graph showing the cumulative
12 number of new 53BP1 foci (red curve) following a bleb bursting event (assessed by an increase
13 in the cytosolic intensity of the probe 53BP1-EGFP, which leaks out of the nucleus upon NE
14 rupture, blue curve). Blue arrow indicates the instant of NE rupture/bleb bursting. *P* value was
15 calculated by unpaired Student's t-test in "H"; ns = not significant.

16

17 **Figure S3. Characterization of DNA damage following nuclear envelope rupture in**
18 **different cell lines and of TREX1 KO clones (related to Figure 3).** (A) Diagrams illustrating
19 the Cas9/gRNA targeting sequence in the *TREX1* gene and the primers used for the
20 sequencing of the TREX1 KO clones generated by CRISPR technology for RPE1 and DCIS
21 cells. (B) Western blot of DCIS/RPE1 TREX1 WT and DCIS/RPE1 TREX1 KO clones; tubulin
22 is a loading control. (C) Single cell curves of 53BP1 foci number as a function of time in RPE1
23 cells (top) or DCIS cells (bottom) displaying NE rupture events while under 2 μm confinement.
24 (D) Top: MCF10A cells stably expressing 53BP1-EGFP and catalytically inactive cGAS-
25 mCherry (icGAS) were transiently depleted for TREX1 with siRNA and 48 h later cells were
26 confined at 2 μm . Bottom: Quantification of DNA damage levels during 2 μm confinement as
27 assessed by the number of 53BP1 foci in siTREX1 MCF10A cells displaying or not NE rupture.
28 Western blot shows TREX1 depletion 48 h post-knockdown; tubulin is the loading control. Data
29 represents the mean of 3 independent experiments where 20 cells per experiment per
30 condition were analyzed. (E) Mean values of the ratio between 53BP1 foci SD mean intensity
31 and 53BP1 foci mean intensity (coefficient of variation of 53BP1 foci intensity) of RPE1 and
32 DCIS cells transiently depleted for TREX1 and confined at 2 μm . This is an alternative way of
33 quantifying the increase in the number of 53BP1 foci in the nucleus, without counting the foci.
34 This globally measures the coefficient of variation of the signal, which increases when the
35 signal becomes less homogeneous. (F) Curves showing the 53BP1-EGFP mean intensity of
36 RPE1 and DCIS cells transiently depleted for TREX1 and confined at 2 μm . (E, F) Graphs
37 represent data pooled from 3 independent experiments for approximately 50 cells analyzed
38 per condition. (G) RPE1 cells stably expressing 53BP1-EGFP were transiently depleted for
39 TREX1 using siRNA and treated with etoposide (25 μM) for 2 h. Subsequently cells fixed and
40 stained for endogenous TREX1. Bar, 25 μm . (H) Frequency distribution of the number of
41 nuclear blebs in RPE1 TREX1 KO clones and in parental RPE1 cells. Data represents 3
42 independent experiments where 60 cells per experiment were analyzed. (I) RPE1 cells
43 transiently transfected with TREX1-EGFP WT were confined at 2 μm using a pressure-
44 controlled dynamic confiner. Images represent a single plane Z slice through the middle section
45 of the nucleus. Bars, 5 μm and 1 μm (magnification). Images are representative of 2
46 independent experiments. (J) Quantification of DNA damage levels (number of 53BP1 foci) in
47 DCIS wt cells and DCIS TREX1 KO #1 cells stably expressing 53BP1-mCherry reconstituted
48 with the indicated TREX1 constructs fused to GFP and confined at 2 μm . Quantification was
49 performed in cells displaying NE rupture. Data represents the mean \pm SD of 3 independent
50 experiments where 25 cells per condition per experiment were analyzed. (K) Images showing

1 the distribution of TREX1 deltaC-GFP and TREX1 deltaC-Sec61TM-GFP in DCIS cells
2 confined at 2 μm (note the diffuse distribution of TREX1 deltaC-GFP). Bar, 5 μm .

3

4 **Figure S4. Characterization of DNA damage following spontaneous nuclear envelope**
5 **rupture and TREX1 localization by electron microscopy (related to Figure 3).** (A, B) Left:
6 RPE1 cells stably expressing 53BP1-EGFP and catalytically inactive cGAS-mCherry were
7 transiently transfected with a non-coding siRNA (A) or with a siRNA targeting TREX1 (B),
8 plated on glass bottom dishes and imaged for several hours. Arrows point to NE rupture events
9 (note the accumulation of the cGAS signal at the rupture site). Right: curves showing the
10 relationship between NE rupture and the appearance of 53BP1 foci (note that in TREX1-
11 depleted cells 53BP1 foci number remains low following NE rupture events). Bars, 5 μm . (C)
12 Immuno-gold electron microscopy images of non-confined (top panels) and confined (2 μm ,
13 middle panels) RPE1 TREX1 KO cells reconstituted with TREX1-mCherry. Bottom panels:
14 non-confined RPE1 WT cells. Following confinement, cells were harvested and labeled with
15 an antibody against mCherry to detect either transduced TREX1 (top and middle panels) or
16 non-specific labeling (negative control, bottom panels). Graph shows the quantification of the
17 relative labelling index (see methods for detailed explanation) of mCherry (TREX1) immune
18 gold particles at the indicated cellular compartments. Bar, 0.5 μm ; magnified regions bar, 0.1
19 μm . For Panel "C", refer to methods and Table S2 for the complete set of numbers and
20 statistics, *** does not stand for P value but significance according to the chi-square statistics.
21 A total of 6 pictures from not confined (260 gold particles) and 9 pictures from 2 μm confined
22 cells (303 gold particles) were used for quantification. For other panels, P values were
23 calculated by One-way ANOVA with post hoc Tukey test in "J". **P < 0.005; *P < 0.05; ns =
24 not significant. See also Table S2 and Movie S4.

25

26 **Figure S5. Long-term, strong confinement causes cell senescence in non-transformed**
27 **cells but not in cancer cells (related to Figure 4).** (A) RPE1, MCF10A and DCIS cells were
28 treated with doxorubicin (40 nM), etoposide (25 μM) or DMSO (vehicle) for the indicated
29 lengths of time. Cells were then fixed and processed for β -gal staining. The percentage of β -
30 gal-positive cells was plotted. Graph, data represents mean \pm SD of 3 independent
31 experiments where 200 cells were scored per condition per experiment. (B) Frequency
32 distribution of the designated categories for cells under the indicated confinement heights.
33 Data represents 3 independent experiments where 37 non-confined cells, 40 confined cells
34 with non-ruptured nuclei and 60 confined cells with ruptured nuclei were scored per condition
35 per experiment. (C) RPE1 cells were confined at 2 μm in the presence of Propidium Iodide and
36 imaged overnight. Graph, data represents the mean \pm SD of 3 independent experiments where
37 5 random fields (at 20x magnification) were scored per time point per experiment. Bar, 50 μm .
38 (D) Epifluorescence and phase images of RPE1 cells stably expressing catalytically inactive
39 cGAS-EGFP (icGAS) harvested from 2 μm confinement and replated for cell cycle duration
40 measurement over a 72 h period. Bar, 25 μm . Inset shows a cell with ruptured NE, as
41 evidenced by cGAS perinuclear accumulation. (E) Graph: box and whisker plot showing the
42 median value and 10-90 percentiles of the ratio between the SD of HP1 γ intensity and HP1 γ
43 mean intensity (coefficient of variation). This is an alternative way of quantifying the increase
44 in the number of HP1 γ foci in the nucleus, without counting the foci. This globally measures
45 the coefficient of variation of the signal, which increases when the signal becomes less
46 homogeneous. Data represents 3 independent experiments where 70 cells were scored per
47 condition per experiment. (F) MCF10A cells or (G) DCIS cells were confined for 12 h at 10 or
48 2 μm , harvested from confinement and replated for 72 h before fixation for immunostaining
49 with heterochromatin foci (HP1 γ) or lysis for western blot analysis of lamin B1 and p21. GAPDH

1 is the loading control. Western blot images are representative of 2 independent experiments.
2 Graph, box and whisker plot showing the median value and 10-90 percentiles of the number
3 of HP1 γ foci per nucleus. Data represents 3 independent experiments where 100 cells per
4 condition per experiment were analyzed. Bars, 10 μ m. (H) Western blot analysis of whole cell
5 extracts of human fibroblasts, RPE1 cells and HeLa cells. GAPDH is the loading control. (I)
6 cGAMP measurements for the indicated cells lines. Bars represent 1 experiment that was
7 performed in duplicate. *P* values were calculated by One-way ANOVA with post hoc Tukey
8 test (A) or unpaired Student's t-test (F, G). ****P* < 0.0001; **P* < 0.05; ns = not significant.

9

10 **Figure S6. Pharmacologically induced DNA damage promotes collagen degradation**
11 **independently of TREX1 and cGAS and strong confinement induces cell polarization**
12 **and collagen degradation in a cGAS-independent fashion (related to Figure 5) TREX1**
13 **KO clones display normal proliferation both *in vitro* and *in vivo* (related to Figure 6).** (A)
14 DCIS cells were treated for 12 h with the indicated concentrations of doxorubicin or DMSO
15 (vehicle). Following the incubation period, cells were harvested and embedded in 3D type I
16 collagen for 12 h to assess their collagen degradation activity by immunostaining with an
17 antibody that specifically recognizes the collagenase-cleaved $\frac{3}{4}$ fragment of collagen I. Graph,
18 box and whisker plot showing the median value and 10-90 percentiles of the collagen
19 degradation area. Data was normalized by the mean of control samples. Data represents 3
20 independent experiments where 50 cells per condition per experiment were analyzed. (B)
21 Quantification of nuclear area of DCIS cells embedded into 3D collagen following the indicated
22 confinement heights. Graph, box and whisker plot showing the median value and 10-90
23 percentiles of nuclear projected area. Data represents 3 independent experiments where 70
24 cells per condition per experiment were analyzed. (C) DCIS cells stably expressing life actin-
25 EGFP were confined for 2 h at the indicated heights. Following the confinement period, cells
26 were harvested and embedded in 3D type I collagen and imaged for 15 h. Bar, 20 μ m. (D)
27 Scatter dot plots of different measured parameters while cells were embedded into 3D
28 collagen (lines are median). Data represents 2 independent experiments where 25 cells per
29 condition per experiment were analyzed. (E) DCIS cells were transiently depleted for TREX1
30 or cGAS using siRNA and 48 h later cells were pre-treated for 12 h with doxorubicin (Doxo, 80
31 nM) or DMSO (vehicle). Following the treatment period, cells were harvested and embedded
32 in 3D type I collagen for 12 h to assess their collagen degradation activity by immunostaining
33 with an antibody that specifically recognizes the collagenase-cleaved $\frac{3}{4}$ fragment of collagen
34 I. Graph, box and whisker plot showing the median value and 10-90 percentiles of the collagen
35 degradation area. Data was normalized by the mean of control samples. Data represents 3
36 independent experiments where 40 cells per condition per experiment were analyzed. (F)
37 DCIS cells were transiently depleted for cGAS using siRNA and 48 h later cells were confined
38 at 2 μ m for 2 h. Following the confinement period, cells were harvested and embedded in 3D
39 type I collagen for 12 h to assess their collagen degradation activity by immunostaining with
40 an antibody that specifically recognizes the collagenase-cleaved $\frac{3}{4}$ fragment of collagen I
41 (red); DAPI (cyan). Bar, 20 μ m. Graph, box and whisker plot showing the median value and
42 10-90 percentiles of the collagen degradation area. Data was normalized by the mean of
43 control samples. Data represents 3 independent experiments where 60 cells per condition per
44 experiment were analyzed. Far right: western blot analysis of cGAS depletion (48 h post-
45 knockdown); tubulin is a loading control. (G) Quantitative Real Time-PCR analysis. DCIS wt or
46 TREX1 KO clones were confined at the indicated heights for 2 h. DCIS cells were harvested,
47 lysed and processed for analysis of the expression of mRNAs of EMT markers. (H) DCIS wt
48 cells were treated for the indicated lengths of time with DMSO (vehicle), Doxorubicin (80 nM)
49 or Etoposide (100 μ M). Cells were harvested, lysed and processed for analysis of the
50 expression of mRNAs of EMT markers. Data represents the mean \pm SD of 3 independent

1 experiments with 3 replicates per experiment. (I) DCIS wt cells were transiently depleted for
2 SNAIL1 and confined at the indicated heights for 2 h. Following the confinement period, cells
3 were harvested and embedded in 3D type I collagen for 12 h to assess their collagen
4 degradation activity by immunostaining with an antibody that specifically recognizes the
5 collagenase-cleaved $\frac{3}{4}$ fragment of collagen I. Graph, box and whisker plot showing the
6 median value and 10-90 percentiles of the collagen degradation area. Data was normalized by
7 the mean of control samples. Data represents 3 independent experiments where 60 cells per
8 condition per experiment were analyzed. Western blot shows SNAIL1 depletion 48 h post-
9 knockdown; tubulin is the loading control. (J) Graph showing the rate of *in situ* tumor burden
10 (period of 6-8 weeks post-intraductal injection). Data represents the mean \pm SEM. A total of 4
11 parental and 11 TREX1 KO glands were analyzed. (K) Graph, box and whisker plot showing
12 the median value and 10-90 percentiles of the tumor area generated by DCIS parental or
13 TREX1 KO clones (6-8 weeks post-intraductal injection). (L) Graph: box and whisker plot
14 showing the median value and 10-90 percentiles of the ratio between tumor area and gland
15 area generated by DCIS parental or TREX1 KO clones (6-8 weeks post-intraductal injection).
16 Data represents a total of 7 glands scored for DCIS parental tumors and 14 glands for TREX1
17 KO tumors. (M) Proliferation curve of DCIS parental and clones TREX1 KO#1 and TREX1
18 KO#2. Graph, data represents the mean \pm SEM of 25 random fields scored at 20x
19 magnification. (N) Graphs showing the proportion of nuclei as a function of distance from the
20 tumor edge (left) or Z score (γ H2AX intensities) (right) for both TREX1 KO and TREX1 WT
21 mice xenografts. Data represents a total of 23 images analyzed for DCIS parental tumors and
22 30 images for TREX1 KO tumors (TREX1 KO #1 + TREX1 KO #2 clones) from 3 independent
23 mice injections. *P* values were calculated by unpaired Student's *t*-test (B, D, F, I, J-L), One-
24 way ANOVA with post hoc Tukey test (A, E). ****P* < 0.0001; **P* < 0.05; ns = not significant. See
25 also Movie S5.

26 **Figure S7. Additional results commented in the discussion of the article, including**
27 **analysis of TREX1 expression that reveals poor prognosis in breast cancer and cGAS**
28 **staining in human tumor samples (related to Figure 7).** (A) cGAS antibody validation by
29 immunohistochemistry (IHC). (Top) DCIS were transiently transfected with siRNA control or
30 siRNA cGAS and 72 hour later cells were paraffin embedded and formalin fixed for IHC
31 processing. Bar, 40 μ m. (Bottom) Quantitative RT-PCR showing cGAS mRNA levels (relative
32 to siControl-treated cells). (B) cGAS immunohistochemistry (brown) and nuclei (blue) in overtly
33 invasive human breast ductal carcinoma. Insets 1 and 2: note cGAS accumulation at the tip of
34 ruptured nuclei. Red arrowheads point to NE rupture events. Left panels: bar, 25 μ m and inset,
35 3 μ m; middle panels: bar, 8 μ m; right panels: bar, 3 μ m. (C) RPE1 cells were transiently
36 transfected with RPA-GFP and either treated with hydroxyurea 2mM or confined at 3 μ m and
37 imaged for several minutes. Yellow arrowhead points to a nuclear bleb that eventually bursts.
38 Bar, 10 μ m. (D) DCIS cells stably expressing Geminin-mCherry were treated with DMSO
39 (vehicle control) or doxorubicin (40 nM) and the percentage of cells in S/G2 was assessed at
40 0 h and 48 h of treatment; box and whisker plot shows the median value and 10-90 percentiles
41 of 2 independent experiments where at least 400 cells per experiment per condition were
42 analyzed. (E) DCIS cells stably expressing Geminin-mCherry were embedded in 3D type I
43 collagen for 12 h to assess their collagen degradation activity by immunostaining with an
44 antibody that specifically recognizes the collagenase-cleaved $\frac{3}{4}$ fragment of collagen I; box
45 and whisker plot shows the median value and 10-90 percentiles of 2 independent experiments
46 where at least 100 cells per experiment were analyzed. (F) Kaplan Meier survival plot for
47 patients expressing low (black line) and high (red line) levels of TREX1 mRNA obtained by
48 RNAseq. Data was obtained from The Cancer Genome Atlas (TCGA), <https://kmplot.com>. (G)
49 MDA-MB-231 cells stably expressing 53BP1-EGFP and catalytically inactive cGAS-mCherry
50 (icGAS) were transiently depleted for TREX1 and confined at 2 μ m. Bar, 10 μ m. (H)

1 Quantification of DNA damage levels following 2 μ m confinement as assessed by the number
2 of 53BP1 foci in MDA-MB-231 cells displaying or not NE rupture (siControl and siTREX1). Data
3 represents the mean of 2 independent experiments where 20 cells per experiment per
4 condition were analyzed. Western blot shows TREX1 depletion 48 h post-knockdown; tubulin
5 is the loading control. (I) Overview of immunohistochemistry analysis of moderately
6 differentiated human colon rectal adenocarcinoma, infiltrating until muscle layer with perineural
7 invasion (classification: pT2). γ H2AX (brown) and nuclei (blue). Bar, 1mm. (J) Magnified
8 regions of the invasive margin shown in "I". (Left) Immunohistochemistry of cGAS (brown) and
9 nuclei (blue). Yellow arrowheads point to NE rupture events. (Right) Immunohistochemistry of
10 γ H2AX (brown) and nuclei (blue). Bar, 20 μ m. (K) Magnified regions of the healthy tissue from
11 the same patient. Bar, 20 μ m. *P* values were calculated by unpaired Student's *t*-test, ****P* <
12 0.0001; ns = not significant.

13

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Supplementary Movies Legends

Movie S1. DCIS cells moving inside the duct on chip device (related to Figure 1). DCIS cells stably expressing 53BP1-mCherry were injected in the duct on chip and imaged using spinning disc microscopy (20x magnification).

Movie S2. RPE1 cells confined at 4 and 2 μm . (related to Figure 2). RPE1 cells stably expressing 53BP1-EGFP and catalytically inactive cGAS-mCherry were confined at 4 (first part of the movie) and 2 μm (second part of the movie) and imaged under 40x magnification using spinning disc microscopy. Cells confined at 4 μm exhibit no NE rupture and cGAS remains cytoplasmic while 53BP1 remains homogenous in the nucleus. At 2 μm sequential nuclear blebbing events are observed, leading to formation of cGAS cluster at the periphery of the nucleus, at the site of blebbing, and accumulation of 53BP1 in multiple foci in the nucleus.

Movie S3. Examples of non-confined RPE1 cells with spontaneous NE rupture and subsequent DNA damage (related to Figure 2). RPE1 cells stably expressing 53BP1-EGFP and catalytically inactive cGAS-mCherry were seeded on a glass bottom dish and imaged under 40x magnification using wide field fluorescence microscopy.

Movie S4. TREX1 nuclear localization upon confinement (related to Figure 3). RPE1 cells were transiently transfected with TREX1-EGFP and 1 μm thickness Z slices were acquired before and after 2 μm confinement. Cells were imaged under 40x magnification using spinning disc microscopy.

Movie S5. DCIS cells subjected to strong confinement exhibit exploratory behavior in 3D collagen (related to Figure 5). DCIS cells stably expressing life-actin-EGFP were confined at 10 μm or 2 μm for 2h. Subsequently, cells were harvested and embedded in a 3D collagen matrix for imaging under 40x magnification using spinning disc microscopy.

References:

Basit, A., Cho, M.G., Kim, E.Y., Kwon, D., Kang, S.J., and Lee, J.H. (2020). The cGAS/STING/TBK1/IRF3 innate immunity pathway maintains chromosomal stability through regulation of p21 levels. *Exp. Mol. Med.*

Behbod, F., Kittrell, F.S., LaMarca, H., Edwards, D., Kerbawy, S., Heestand, J.C., Young, E., Mukhopadhyay, P., Yeh, H.W., Allred, D.C., et al. (2009). An intraductal human-in-mouse transplantation model mimics the subtypes of ductal carcinoma in situ. *Breast Cancer Res.*

Benci, J.L., Johnson, L.R., Choa, R., Xu, Y., Qiu, J., Zhou, Z., Xu, B., Ye, D., Nathanson, K.L., June, C.H., et al. (2019). Opposing Functions of Interferon Coordinate Adaptive and Innate Immune Responses to Cancer Immune Checkpoint Blockade. *Cell.*

Le Berre, M., Aubertin, J., and Piel, M. (2012). Fine control of nuclear confinement identifies a threshold deformation leading to lamina rupture and induction of specific genes. *Integr. Biol.* 4, 1406.

Le Berre, M., Zlotek-Zlotkiewicz, E., Bonazzi, D., Lautenschlaeger, F., and Piel, M. (2014). Methods for Two-Dimensional Cell Confinement. In *Methods in Cell Biology*, pp. 213–229.

Blaha, L., Zhang, C., Cabodi, M., and Wong, J.Y. (2017). A microfluidic platform for modeling metastatic cancer cell matrix invasion. *Biofabrication.*

Castagnino, A., Castro-Castro, A., Irondelle, M., Guichard, A., Lodillinsky, C., Fuhrmann, L., Vacher, S., Agüera-González, S., Zagryazhskaya-Masson, A., Romao, M., et al. (2018). Coronin 1C promotes triple-negative breast cancer invasiveness through regulation of MT1-MMP traffic and invadopodia function. *Oncogene.*

1 Cho, S., Vashisth, M., Abbas, A., Majkut, S., Vogel, K., Xia, Y., Ivanovska, I.L., Irianto, J.,
2 Tewari, M., Zhu, K., et al. (2019). Mechanosensing by the Lamina Protects against Nuclear
3 Rupture, DNA Damage, and Cell-Cycle Arrest. *Dev. Cell*.

4 Chowdhury, D., Beresford, P.J., Zhu, P., Zhang, D., Sung, J.S., Demple, B., Perrino, F.W.,
5 and Lieberman, J. (2006). The Exonuclease TREX1 Is in the SET Complex and Acts in
6 Concert with NM23-H1 to Degrade DNA during Granzyme A-Mediated Cell Death. *Mol. Cell*.

7 Denais, C.M., Gilbert, R.M., Isermann, P., McGregor, A.L., te Lindert, M., Weigelin, B.,
8 Davidson, P.M., Friedl, P., Wolf, K., and Lammerding, J. (2016). Nuclear envelope rupture
9 and repair during cancer cell migration. *Science* (80-.). 352, 353–358.

10 Earle, A.J., Kirby, T.J., Fedorchak, G.R., Isermann, P., Patel, J., Iruvanti, S., Moore, S.A.,
11 Bonne, G., Wallrath, L.L., and Lammerding, J. (2020). Mutant lamins cause nuclear envelope
12 rupture and DNA damage in skeletal muscle cells. *Nat. Mater.*

13 Erdal, E., Haider, S., Rehwinkel, J., Harris, A.L., and McHugh, P.J. (2017). A prosurvival
14 DNA damage-induced cytoplasmic interferon response is mediated by end resection factors
15 and is limited by Trex1. *Genes Dev.*

16 Feinberg, T.Y., Zheng, H., Liu, R., Wicha, M.S., Yu, S.M., and Weiss, S.J. (2018). Divergent
17 Matrix-Remodeling Strategies Distinguish Developmental from Neoplastic Mammary
18 Epithelial Cell Invasion Programs. *Dev. Cell*.

19 Feng, C., Yang, M., Zhang, Y., Lan, M., Huang, B., Liu, H., and Zhou, Y. (2018). Cyclic
20 mechanical tension reinforces DNA damage and activates the p53-p21-Rb pathway to
21 induce premature senescence of nucleus pulposus cells. *Int. J. Mol. Med.*

22 Friedl, P., and Gilmour, D. (2009). Collective cell migration in morphogenesis, regeneration
23 and cancer. *Nat. Rev. Mol. Cell Biol.*

24 Gentili, M., Lahaye, X., Nadalin, F., Nader, G.P.F., Lombardi, E.P., Herve, S., De Silva, N.S.,
25 Rookhuizen, D.C., Zueva, E., Goudot, C., et al. (2019). The N-Terminal Domain of cGAS
26 Determines Preferential Association with Centromeric DNA and Innate Immune Activation in
27 the Nucleus. *Cell Rep.*

28 Gorgoulis, V., Adams, P.D., Alimonti, A., Bennett, D.C., Bischof, O., Bishop, C., Campisi, J.,
29 Collado, M., Evangelou, K., Ferbeyre, G., et al. (2019). Cellular Senescence: Defining a Path
30 Forward. *Cell*.

31 Harding, S.M., Benci, J.L., Irianto, J., Discher, D.E., Minn, A.J., and Greenberg, R.A. (2017).
32 Mitotic progression following DNA damage enables pattern recognition within micronuclei.
33 *Nature*.

34 Hatch, E.M. (2018). Nuclear envelope rupture: little holes, big openings. *Curr. Opin. Cell Biol.*

35 Hatch, E.M., and Hetzer, M.W. (2016). Nuclear envelope rupture is induced by actin-based
36 nucleus confinement. *J. Cell Biol.*

37 Infante, E., Castagnino, A., Ferrari, R., Monteiro, P., Agüera-González, S., Paul-Gilloteaux,
38 P., Domingues, M.J., Maiuri, P., Raab, M., Shanahan, C.M., et al. (2018). LINC complex-Lis1
39 interplay controls MT1-MMP matrix digest-on-demand response for confined tumor cell
40 migration. *Nat. Commun.*

41 Irianto, J., Xia, Y., Pfeifer, C.R., Athirasala, A., Ji, J., Alvey, C., Tewari, M., Bennett, R.R.,
42 Harding, S.M., Liu, A.J., et al. (2017). DNA Damage Follows Repair Factor Depletion and
43 Portends Genome Variation in Cancer Cells after Pore Migration. *Curr. Biol.*

44 Jiang, Y.N., Ni, X.Y., Yan, H.Q., Shi, L., Lu, N.N., Wang, Y.N., Li, Q., and Gao, F.G. (2019).
45 Interleukin 6-triggered ataxia-telangiectasia mutated kinase activation facilitates epithelial-to-

1 mesenchymal transition in lung cancer by upregulating vimentin expression. *Exp. Cell Res.*

2 Kang, H.T., Park, J.T., Choi, K., Kim, Y., Choi, H.J.C., Jung, C.W., Lee, Y.S., and Park, S.C.
3 (2017). Chemical screening identifies ATM as a target for alleviating senescence. *Nat.*
4 *Chem. Biol.*

5 Kim, M., Kim, H.J., Chung, Y.R., Kang, E., Kim, E.K., Kim, S.H., Kim, Y.J., Kim, J.H., Kim,
6 I.A., and Park, S.Y. (2018). Microinvasive carcinoma versus ductal carcinoma in situ: A
7 comparison of clinicopathological features and clinical outcomes. *J. Breast Cancer.*

8 Lahaye, X., Satoh, T., Gentili, M., Cerboni, S., Conrad, C., Hurbain, I., El Marjou, A.,
9 Lacabaratz, C., Lelièvre, J.-D., and Manel, N. (2013). The Capsids of HIV-1 and HIV-2
10 Determine Immune Detection of the Viral cDNA by the Innate Sensor cGAS in Dendritic
11 Cells. *Immunity* 39, 1132–1142.

12 Lee-Kirsch, M.A., Chowdhury, D., Harvey, S., Gong, M., Senenko, L., Engel, K., Pfeiffer, C.,
13 Hollis, T., Gahr, M., Perrino, F.W., et al. (2007a). A mutation in TREX1 that impairs
14 susceptibility to granzyme A-mediated cell death underlies familial chilblain lupus. *J. Mol.*
15 *Med.*

16 Lee-Kirsch, M.A., Gong, M., Chowdhury, D., Senenko, L., Engel, K., Lee, Y.A., De Silva, U.,
17 Bailey, S.L., Witte, T., Vyse, T.J., et al. (2007b). Mutations in the gene encoding the 3'-5'
18 DNA exonuclease TREX1 are associated with systemic lupus erythematosus. *Nat. Genet.*

19 Lehtinen, D.A., Harvey, S., Mulcahy, M.J., Hollis, T., and Perrino, F.W. (2008). The TREX1
20 double-stranded DNA degradation activity is defective in dominant mutations associated with
21 autoimmune disease. *J. Biol. Chem.*

22 Liu, H., Zhang, H., Wu, X., Ma, D., Wu, J., Wang, L., Jiang, Y., Fei, Y., Zhu, C., Tan, R., et al.
23 (2018). Nuclear cGAS suppresses DNA repair and promotes tumorigenesis. *Nature.*

24 Liu, Y.-J., Le Berre, M., Lautenschlaeger, F., Maiuri, P., Callan-Jones, A., Heuzé, M., Takaki,
25 T., Voituriez, R., and Piel, M. (2015). Confinement and Low Adhesion Induce Fast Amoeboid
26 Migration of Slow Mesenchymal Cells. *Cell* 160, 659–672.

27 Lodillinsky, C., Infante, E., Guichard, A., Chaligné, R., Fuhrmann, L., Cyrta, J., Irondelle, M.,
28 Lagoutte, E., Vacher, S., Bonsang-Kitzis, H., et al. (2016). p63/MT1-MMP axis is required for
29 in situ to invasive transition in basal-like breast cancer. *Oncogene.*

30 Lusk, C.P., and Ader, N.R. (2020). CHMPions of repair: Emerging perspectives on sensing
31 and repairing the nuclear envelope barrier. *Curr. Opin. Cell Biol.* 64, 25–33.

32 Maciejowski, J., Li, Y., Bosco, N., Campbell, P.J., and De Lange, T. (2015). Chromothripsis
33 and Kataegis Induced by Telomere Crisis. *Cell.*

34 Maciejowski, J., Chatzipli, A., Dananberg, A., Lange, T. de, and Campbell, P. (2019).
35 APOBEC3B-dependent kataegis and TREX1-driven chromothripsis in telomere crisis.
36 *BioRxiv.*

37 Malhas, A., Goulbourne, C., and Vaux, D.J. (2011). The nucleoplasmic reticulum: Form and
38 function. *Trends Cell Biol.*

39 Malinverno, C., Corallino, S., Giavazzi, F., Bergert, M., Li, Q., Leoni, M., Disanza, A., Frittoli,
40 E., Oldani, A., Martini, E., et al. (2017). Endocytic reawakening of motility in jammed
41 epithelia. *Nat. Mater.*

42 Martinvalet, D., Zhu, P., and Lieberman, J. (2005). Granzyme A induces caspase-
43 independent mitochondrial damage, a required first step for apoptosis. *Immunity.*

44 Mayhew, T.M. (2011). Mapping the distributions and quantifying the labelling intensities of
45 cell compartments by immunoelectron microscopy: Progress towards a coherent set of

1 methods. *J. Anat.* 219, 647–660.

2 Mazur, D.J., and Perrino, F.W. (2001). Excision of 3' termini by the Trex1 and TREX2 3'→5'
3 exonucleases. Characterization of the recombinant proteins. *J. Biol. Chem.*

4 Mohr, L., Toufektchan, E., von Morgen, P., Chu, K., Kapoor, A., and Maciejowski, J. (2021).
5 ER-directed TREX1 limits cGAS activation at micronuclei. *Mol. Cell* 81, 724-738.e9.

6 Palamidessi, A., Malinverno, C., Frittoli, E., Corallino, S., Barbieri, E., Sigismund, S.,
7 Beznoussenko, G. V., Martini, E., Garre, M., Ferrara, I., et al. (2019). Unjamming overcomes
8 kinetic and proliferation arrest in terminally differentiated cells and promotes collective
9 motility of carcinoma. *Nat. Mater.*

10 Peng, B., Ortega, J., Gu, L., Chang, Z., and Li, G.M. (2019). Phosphorylation of proliferating
11 cell nuclear antigen promotes cancer progression by activating the ATM/Akt/GSK3β/Snail
12 signaling pathway. *J. Biol. Chem.*

13 Raab, M., Gentili, M., de Belly, H., Thiam, H.-R., Vargas, P., Jimenez, A.J., Lautenschlaeger,
14 F., Voituriez, R., Lennon-Dumenil, A.-M., Manel, N., et al. (2016). ESCRT III repairs nuclear
15 envelope ruptures during cell migration to limit DNA damage and cell death. *Science* (80-).
16 352, 359–362.

17 Robijns, J., Houthaeve, G., Braeckmans, K., and De Vos, W.H. (2018). Loss of Nuclear
18 Envelope Integrity in Aging and Disease. In *International Review of Cell and Molecular*
19 *Biology*, p.

20 Roman, W., Martins, J.P., Carvalho, F.A., Voituriez, R., Abella, J.V.G., Santos, N.C., Cadot,
21 B., Way, M., and Gomes, E.R. (2017). Myofibril contraction and crosslinking drive nuclear
22 movement to the periphery of skeletal muscle. *Nat. Cell Biol.*

23 Shah, P., Wolf, K., and Lammerding, J. (2017). Bursting the Bubble – Nuclear Envelope
24 Rupture as a Path to Genomic Instability? *Trends Cell Biol.*

25 Shah, P., Hobson, C.M., Cheng, S., Colville, M.J., Paszek, M.J., Superfine, R., and
26 Lammerding, J. (2021). Nuclear Deformation Causes DNA Damage by Increasing
27 Replication Stress. *Curr. Biol.* 31, 753-765.e6.

28 Slot, J.W., and Geuze, H.J. (2007). Cryosectioning and immunolabeling. *Nat. Protoc.* 2,
29 2480–2491.

30 Srivastava, N., de Freitas Nader, G.P., Williard, A., Rollin, R., Cuvelier, D., Lomakin, A., and
31 Piel, M. (2021). Nuclear fragility, blaming the blebs. *Curr. Opin. Cell Biol.* 70, 100–108.

32 Stetson, D.B., Ko, J.S., Heidmann, T., and Medzhitov, R. (2008). Trex1 Prevents Cell-
33 Intrinsic Initiation of Autoimmunity. *Cell.*

34 Sun, M., Guo, X., Qian, X., Wang, H., Yang, C., Brinkman, K.L., Serrano-Gonzalez, M., Jope,
35 R.S., Zhou, B., Engler, D.A., et al. (2012). Activation of the ATM-Snail pathway promotes
36 breast cancer metastasis. *J. Mol. Cell Biol.* 4, 304–315.

37 Takahashi, A., Loo, T.M., Okada, R., Kamachi, F., Watanabe, Y., Wakita, M., Watanabe, S.,
38 Kawamoto, S., Miyata, K., Barber, G.N., et al. (2018). Downregulation of cytoplasmic
39 DNases is implicated in cytoplasmic DNA accumulation and SASP in senescent cells. *Nat.*
40 *Commun.*

41 Teuliere, J. (2005). Targeted activation of -catenin signaling in basal mammary epithelial
42 cells affects mammary development and leads to hyperplasia. *Development* 132, 267–277.

43 Ungricht, R., and Kutay, U. (2015). Establishment of NE asymmetry-targeting of membrane
44 proteins to the inner nuclear membrane. *Curr. Opin. Cell Biol.* 34, 135–141.

- 1 Vanpouille-Box, C., Alard, A., Aryankalayil, M.J., Sarfraz, Y., Diamond, J.M., Schneider, R.J.,
2 Inghirami, G., Coleman, C.N., Formenti, S.C., and Demaria, S. (2017). DNA exonuclease
3 Trex1 regulates radiotherapy-induced tumour immunogenicity. *Nat. Commun.*
- 4 Vargas, J.D., Hatch, E.M., Anderson, D.J., and Hetzer, M.W. (2012). Transient nuclear
5 envelope rupturing during interphase in human cancer cells. *Nucleus.*
- 6 De vos, W.H., Houben, F., Kamps, M., Malhas, A., Verheyen, F., Cox, J., Manders, E.M.M.,
7 Verstraeten, V.L.R.M., Van steensel, M.A.M., Marcelis, C.L.M., et al. (2011). Repetitive
8 disruptions of the nuclear envelope invoke temporary loss of cellular compartmentalization in
9 laminopathies. *Hum. Mol. Genet.*
- 10 Yang, Y.G., Lindahl, T., and Barnes, D.E. (2007). Trex1 Exonuclease Degrades ssDNA to
11 Prevent Chronic Checkpoint Activation and Autoimmune Disease. *Cell.*
- 12


KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
RAB5A	Abcam	Cat# ab109534
γ H2AX	Abcam	Cat# ab11174
Rabbit monoclonal anti-Ku70 (clone EPR4026)	Abcam	Cat# ab108604; RRID: AB_10861861
Rabbit monoclonal anti-p21 (clone ERF3993)	Abcam	Cat# ab109199; RRID: AB_10861551
Mouse monoclonal anti-Ki67 (clone MIB-1)	Agilent	Cat# M7240; RRID: AB_2142367
Mouse monoclonal anti-alpha SMA (clone 1A4)	Agilent	Cat# M0851; RRID: AB_2223500
Rabbit polyclonal anti-RIF1	Bethyl Laboratories	Cat# A300-569A
Rabbit monoclonal anti-cGAS (clone D1D3G)	Cell Signaling Technology	Cat# 15102; RRID: AB_2732795
Rabbit monoclonal anti-TREX1 (clone D8E2O) (discontinued)	Cell Signaling Technology	Cat# 15107; RRID: AB_2798710
Rabbit monoclonal anti-GAPDH (clone 14C10)	Cell Signaling Technology	Cat# 2118; RRID: AB_561053
Rabbit monoclonal anti-Snail1 (clone C15D3)	Cell Signaling Technology	Cat# 3879; RRID: AB_2255011
Mouse monoclonal anti-PDI (clone 1D3)	Enzo LifeSciences	Cat# ADI-SPA-891, RRID: AB_10615355
Rabbit polyclonal anti-mCherry	GeneTex	Cat# GTX128508; RRID: AB_2721247
Rabbit polyclonal anti-Col1-3/4C (collagenase-cleaved 3/4 fragment of collagen I)	ImmunoGlobe GmbH	Cat# #0217-050
Mouse monoclonal anti-phospho-Histone H2A.X (Ser139) (clone JBW301)	Millipore	Cat# 05-636; RRID: AB_309864
Mouse monoclonal anti-HP1 γ (clone 14D3.1)	Millipore	Cat# MABE656; RRID: AB_2722628
Mouse monoclonal anti-TREX1 (clone E-6)	Santa Cruz Biotechnology	Cat# sc-271870; RRID: AB_10708266
Mouse monoclonal anti-laminB1 (clone A-11)	Santa Cruz Biotechnology	Cat# sc-377000; RRID: AB_2861346
Mouse monoclonal anti- α tubulin (clone DM1A)	Sigma-Aldrich	Cat# T9026; RRID: AB_477593
Biological samples		
Human Breast Ductal Adenocarcinoma tissue sections	Archival anonymized tumor tissue collection; Tumor Immunology Unit, Department of Health Sciences, University of Palermo School of Medicine	claudio.tripodo@unipa.it
Chemicals, peptides, and recombinant proteins		
KU-55933 (ATM kinase inhibitor)	Abcam	Cat# 120637; CAS: 587871-26-9
Unlabeled type I collagen	BD Biosciences	Cat# 354236

Etoposide	Cell Signaling Technology	Cat# 2200; CAS: 33419-42-0
Doxorubicin	Cell Signaling Technology	Cat# 5927; CAS: 25316-40-9
Penicillin Streptomycin	Gibco	Cat# 15140-122
Unlabeled type I collagen, 5mg/ml	IBIDI	Cat# 50201
GM6001 (matrix metalloproteinases pan-inhibitor)	Millipore	Cat# CC1010
Human EGF	Peptrotech	Cat# AF-100-15; CAS: 62253-63-8
Hydrocortisone	Sigma-Aldrich	Cat# H0888; CAS: 50-23-7
Cholera toxin	Sigma-Aldrich	Cat# C8052; CAS: 9012-63-9
Insulin	Sigma-Aldrich	Cat# I9278; CAS: 11061-68-0
HT-DNA Deoxyribonucleic acid sodium salt from herring testes	Sigma-Aldrich	Cat# D6898 ; CAS: 438545-06-3
Matrigel	Sigma-Aldrich	Cat# E1270
Protein A gold conjugates	Utrecht University, The Netherlands	N/A
Harris's Hematoxylin	Bio-Optica	Cat# 05-06004/L
pLL-g-PEG	SuSoS	Cat# PLL(20)-g[3.5]-PEG(2)
Critical commercial assays		
2'3'-cGAMP ELISA kit	Interchim	Cat# 501700
DAB Chromogen Kit	BioCare Medical	Cat# DB801L
Vulcan fast red Chromogen Kit	BioCare Medical	Cat# FR805M
Opal 7-Color IHC Kit	Akoya 23 Biosciences	Cat# NEL821001KT
Experimental models: Cell lines		
Human, female: RPE1	Lab of Dr. Matthieu Piel, Institut Curie, Paris	Originally ATCC CRL-4000
Human, female: RPE1 stably expressing icGAS-GFP	This paper	N/A
Human, female: RPE1 stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Human, female: RPE1 TREX1 KO	This paper	N/A
Human, female: RPE1 TREX1 KO stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Human, female: MCF10A	Lab of Dr. Philippe Chavrier, Institut Curie, Paris	Originally ATCC CRL-10317
Human, female: MCF10A stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Human, female: MCF10DCIS.com	Lab of Dr. Philippe Chavrier, Institut Curie, Paris ; Behbod et al. 2009	Originally from Asterand (Detroit, MI, USA)

Human, female: MCF10DCIS.com stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Human, female: MCF10DCIS.com TREX1 KO	This paper	N/A
Human, female: MCF10DCIS.com TREX1 KO stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Human, female: MDA-MB-231	Lab of Dr. Philippe Chavrier, Institut Curie, Paris	Originally ATCC CRM-HTB-26
Human, female: MDA-MB-231 stably expressing icGAS-mcherry and 53BP1-GFP	This paper	N/A
Experimental models: Organisms/strains		
Mouse: CB17/Icr-Prkdc ^{scid} /IcrIcoCrI	Charles River Laboratories	Strain code #236
Oligonucleotides		
ON-TARGETplus SMARTpool siTREX1	Dharmacon	Cat# L-013239-02-0005
ON-TARGETplus SMARTpool sicGAS (MB21D1)	Dharmacon	Cat# L-015607-02-0005
ON-TARGETplus SMARTpool siSNAI1	Dharmacon	Cat# L-010847-01-0005
TREX1 guide RNA to generate CRISPR KO clones: 5'-GAGCCCCCCCACCTCTC-(PAM)-3'	Maciejowski et al., 2015	Lab of Titia De Lange, Rockefeller University, New York, USA
GAPDH: NM_001289746.1 (sequence copyright protected)	This paper	Thermo Fischer
CDH2: NM_001308176.1 (sequence copyright protected)	This paper	Thermo Fischer
VIM: NM_003380.3 (sequence copyright protected)	This paper	Thermo Fischer
SNAI1: NM_005985.3 (sequence copyright protected)	This paper	Thermo Fischer
SNAI2: NM_003068.4 (sequence copyright protected)	This paper	Thermo Fischer
TWIST1: NM_000474.3 (sequence copyright protected)	This paper	Thermo Fischer
ZEB1: NM_001128128.2 (sequence copyright protected)	This paper	Thermo Fischer
Recombinant DNA		
pTRIP-CMV-mCherry-TREX1 WT	Lab of Dr. Nicolas Manel, Institut Curie, Paris	N/A
pTRIP-CMV-mCherry-TREX1 D18N	Lab of Dr. Nicolas Manel, Institut Curie, Paris	N/A
pLENTI-CMV-GFP-TREX1	Mohr et al., 2021	Addgene #164228
pTRIP-SFFV-EGFP-FLAG-cGAS E225A/D227A	Raab et al., 2016; Gentili et al., 2019	Addgene #127661
pTRIP-CMV-mCherry-FLAG-cGAS E225A/D227A	Gentili et al., 2019	Addgene #127657
pTRIP-SFFV-EGFP-53BP1	Lab of Dr. Nicolas Manel, Institut Curie, Paris	N/A
pLENTI-CMV-GFP-TREX1 deltaC (aa1-235)	Mohr et al., 2021	Addgene #164229
pLENTI-CMV-GFP-TREX1 deltaC-sec61TM	Lab of Dr. John Maciejowski, MSKCC, NY	N/A
pQCXIP-GFP-RPA70	Mohr et al., 2021	Addgene #164231
Software and algorithms		

GraphPad Prism version 5.02	GraphPad	https://www.graphpad.com/
Fiji ImageJ 1.52p	ImageJ	https://fiji.sc/
Image J macros	This paper	Mendeley Data doi: 10.17632/hbhv5brnxj. 1
Metamorph version 7.7.11.0	Molecular Devices, LLC	https://www.moleculardevices.com/






Figure 3




Figure 4






Figure 5


Figure 5

[Click here to access/download;Figure;Fig 5.pdf](#)



A




B




C



E



F



D






Figure 7


A





B





C




















Table S1 (relative to figure 3D). Testing for differences in the observed distributions of TREX1-mCherry labelled by gold particles between two groups of cells (not confined and 2µm confined), estimating observed and expected numbers of gold particles in a given compartment. For a particular compartment in a given cell, the number of expected gold particles was calculated by multiplying the corresponding column sum by the corresponding row sum and then dividing by the grand row sum. The compartment "Cytoplasm" also includes the gold particles associated to membranes such as ER and multi-vesicular bodies. Chi-square values accounting for 10% or more of total chi-square are considered to be significant. Refer to methods for a detailed explanation of this analysis (approach 1).

Compartment	Not confined		Confined		row total	X ² (Chi-square)		X ² as %	
	observed	expected	observed	expected		Not confined	Confined	Not confined	Confined
Nucleus	30	62.8	106	73.2	136	17.1361454	14.7042832	37.3566136	32.05518
Outer nuclear membrane	18	20.8	27	24.2	45	0.37229676	0.31946257	0.81160295	0.69642498
Cytoplasm	212	176.4	170	205.6	382	7.17921468	6.16038223	15.6506111	13.4295673
total	260		303		563	45.8717849		100	

Table S2 (relative to figures 3D and S4C). Testing for differences in specific-labelling enrichment of TREX1-mCherry gold particles between compartments in non-confined and 2µm confined RPE1 cells. The relative labelling index is estimated from observed and expected numbers of gold particles. LD=labelling density; RLI=Relative labelling index; Ngo=observed Nano gold particles; Nge=expected nano gold particles; Residuum= all the other cellular compartments. The expected distribution was obtained by superimposing to pictures an array of points that was generated digitally, and points (P) were counted in the selected compartments. For each compartment, Ngo was compared with the expected numbers of gold particles (Nge, derived from the observed frequencies of points P). Chi-square values accounting for 10% or more of total chi-square are considered to be significant and thus to contribute to the enrichment in the indicated compartments. Refer to methods for a detailed explanation of this analysis (approach 2).

Compartment	Non confined							Confined						
	Ngo	P	Nge	LD (=Ngo/P)	RLI (=LD/LD cell)	X ² (Chi-square)	X ² as %	Ngo	P	Nge	LD (=Ngo/P)	RLI (=LD/LD cell)	X ² (Chi-square)	X ² as %
Nucleus	26	932	116.56	0.03	0.22	70.36	18.8	86	1560	176.11	0.06	0.49	46.11	16.3
Inner nuclear membrane	4	45	5.63	0.09	0.71	0.47	0.1	20	54	6.1	0.37	3.28	31.71	11.2
Outer nuclear membrane	18	19	2.38	0.95	7.58	102.73	27.5	27	44	4.97	0.61	5.44	97.73	34.5
Peripheral ER	29	45	5.63	0.64	5.15	97.07	26	20	27	3.05	0.74	6.56	94.28	33.2
Residuum	155	995	124.43	0.16	1.25	7.51	2	150	1719	194.06	0.09	0.77	10	3.5
Column sum	260	2079	260	0.13	1	373.3	100	303	2684	303	0.11	1	283.67	100