



**HAL**  
open science

## A unified strategy for observability of waves in an annulus with various boundary conditions

Lucie Baudouin, Jérémi Dardé, Sylvain Ervedoza, Alberto Mercado

### ► To cite this version:

Lucie Baudouin, Jérémi Dardé, Sylvain Ervedoza, Alberto Mercado. A unified strategy for observability of waves in an annulus with various boundary conditions. 2021. hal-03401646v1

**HAL Id: hal-03401646**

**<https://hal.science/hal-03401646v1>**

Preprint submitted on 25 Oct 2021 (v1), last revised 25 Jan 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A unified strategy for observability of waves in an annulus with various boundary conditions\*

Lucie Baudouin<sup>†</sup>    Jérémi Dardé<sup>‡</sup>    Sylvain Ervedoza<sup>§</sup>    Alberto Mercado<sup>¶</sup>

October 22, 2021

*Dedicated to Marius Tucsnak with friendship*

## Abstract

In this work, we develop a unified strategy which allows to derive observability results for waves in an annulus when the observation is done on the external boundary and under various boundary conditions at the internal boundary. Our approach is based on the fact that the observability of a linear abstract conservative system is equivalent to a suitable resolvent estimate. We develop a full machinery to derive suitable resolvent estimates under some weak assumptions on the boundary condition on the internal sphere, which are shown to be close to sharp. In fact, our strategy allows to exclude the existence of trapped rays close to the internal sphere when the boundary conditions satisfy a suitable assumption. As an application, we apply our results to several wave models.

## 1 Introduction

The goal of this article is to provide a robust strategy to address the observability of the wave equation in a bounded domain with various boundary conditions. In particular, we would like to exhibit sufficient conditions on the boundary conditions which guarantee the observability of the equation.

In order to describe our strategy and to avoid difficulties linked with the geometry, we consider the case of an annulus of  $\mathbb{R}^2$ , i.e.  $A(R_0, R_1) = B(R_1) \setminus \overline{B(R_0)}$ , where  $0 < R_0 < R_1$  and  $B(R)$  denotes the open ball of  $\mathbb{R}^2$  with center at the origin and radius  $R > 0$ , which is observed through the whole external boundary  $S(R_1)$  ( $S(R)$  denotes the sphere of  $\mathbb{R}^2$  of radius  $R$ ).

We will assume that the boundary conditions are homogeneous Dirichlet boundary conditions on  $S(R_1)$  since this does not create any specific difficulty as the observation is done there.

Our goal is to give a functional setting allowing different types of boundary conditions on the internal boundary  $S(R_0)$ , including in particular those given below, and to obtain observability results at once for all those cases.

### 1.1 Examples

Let us mention below several examples of interest on which our method will apply.

---

\*Jérémi Dardé and Sylvain Ervedoza are partially supported by the Project TRECOS ANR-20-CE40-0009 funded by the ANR (2021–2024). Part of this work was done while Alberto Mercado was visiting the Institut de Mathématiques de Toulouse within the Labex CIMI funded by the ANR grant 11-LABX-0040, partially supported by Fondecyt 1211292 and ANID Millennium Science Initiative Program, Code NCN19-161.

<sup>†</sup>LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France, [baudouin@laas.fr](mailto:baudouin@laas.fr).

<sup>‡</sup>Institut de Mathématiques de Toulouse UMR 5219, Université de Toulouse, CNRS, UPS, F-31062 Toulouse Cedex 9, France, [jeremi.darde@math.univ-toulouse.fr](mailto:jeremi.darde@math.univ-toulouse.fr)

<sup>§</sup>Univ. Bordeaux, CNRS, Bordeaux INP, IMB, UMR 5251, F-33400 Talence, France, [sylvain.ervedoza@math.u-bordeaux.fr](mailto:sylvain.ervedoza@math.u-bordeaux.fr).

<sup>¶</sup>Departamento de Matemática, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile, [alberto.mercado@usm.cl](mailto:alberto.mercado@usm.cl).

**Fourier Boundary Conditions.** Let  $\alpha \geq 0$  and consider the wave equation with Fourier boundary conditions on  $S(R_0)$ :

$$\begin{cases} \partial_{tt}y(t, x) - \Delta y(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ y(t, x) = 0, & \text{on } (0, T) \times S(R_1), \\ \partial_\nu y(t, x) + \alpha y(t, x) = 0, & \text{on } (0, T) \times S(R_0), \\ (y(0, \cdot), \partial_t y(0, \cdot)) = (y^0, y^1), & \text{in } A(R_0, R_1), \end{cases} \quad (1.1)$$

where  $\partial_\nu$  denotes the normal derivative in the outward direction, i.e.  $\partial_\nu = -\partial_r$  on  $S(R_0)$ , and  $\partial_\nu = \partial_r$  on  $S(R_1)$ .

The question of observability for (1.1) through the boundary  $S(R_1)$  we shall address asks the existence of a time  $T > 0$  and a constant  $C > 0$  such that for all  $(y^0, y^1) \in H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))$  with  $y^0 = 0$  on  $S(R_1)$ , the solution  $y$  of (1.1) satisfies

$$\|(y^0, y^1)\|_{H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))} \leq C \|\partial_\nu y\|_{L^2(0, T; L^2(S(R_1)))}. \quad (1.2)$$

For system (1.1), the observability inequality (1.2) is known to hold for  $T$  large enough, namely  $T > 2\sqrt{R_1^2 - R_0^2}$  (see [3]).

**Waves on the boundary.** Likewise, we shall consider the case in which a wave equation also takes place on the internal boundary  $S(R_0)$ , also known as dynamic Wentzell boundary conditions. Namely, for  $\alpha > 0$  and  $\beta > 0$ , we consider the equation

$$\begin{cases} \partial_{tt}y(t, x) - \Delta y(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ y(t, x) = 0, & \text{on } (0, T) \times S(R_1), \\ z(t, x) = y(t, x), & \text{on } (0, T) \times S(R_0), \\ \partial_\nu y(t, x) + \alpha \partial_{tt}z(t, x) - \beta \Delta_{S(R_0)}z = 0, & \text{on } (0, T) \times S(R_0), \\ (y(0, \cdot), \partial_t y(0, \cdot)) = (y^0, y^1), & \text{in } A(R_0, R_1), \\ (z(0, \cdot), \partial_t z(0, \cdot)) = (z^0, z^1), & \text{on } S(R_0). \end{cases} \quad (1.3)$$

Here, the operator  $\Delta_{S(R_0)}$  simply coincides with  $\partial_{\theta\theta}/R_0^2$ .

The relevant observability inequality for (1.3) through  $S(R_1)$  then asks the existence of a time  $T > 0$  and a constant  $C > 0$  such that for all  $(y^0, y^1, z^0, z^1) \in H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1)) \times H^1(S(R_0)) \times L^2(S(R_0))$  with  $y^0|_{S(R_0)} = z^0$  and  $y^0|_{S(R_1)} = 0$ , the solution  $(y, z)$  of (1.3) satisfies

$$\|(y^0, y^1, z^0, z^1)\|_{H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1)) \times H^1(S(R_0)) \times L^2(S(R_0))} \leq C \|\partial_\nu y\|_{L^2(0, T; L^2(S(R_1)))}. \quad (1.4)$$

Up to our knowledge, the results we will obtain on the observability properties of (1.3) are new (see Section 2.2). We refer to [5, 13] for controllability results on closely related models.

**Fractional Laplacian on the boundary.** We will also study the case in which the internal boundary condition is given through a non-local boundary condition expressed in terms of powers of the Laplacian on the sphere, given for  $s \in (0, 1]$  as follows:

$$\begin{cases} \partial_{tt}y(t, x) - \Delta y(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ y(t, x) = 0, & \text{on } (0, T) \times S(R_1), \\ \partial_\nu y(t, x) + (-\Delta_{S(R_0)})^s y(t, x) = 0, & \text{on } (0, T) \times S(R_0), \\ (y(0, \cdot), \partial_t y(0, \cdot)) = (y^0, y^1), & \text{in } A(R_0, R_1), \end{cases} \quad (1.5)$$

Note in particular that, for  $s = 1/2$ , using the classical interpretation of the square root of the Laplace operator (see e.g. [8])  $(-\Delta_{S(R_0)})^{1/2}$  corresponds to the Dirichlet to Neumann map for the Laplacian in the ball  $B(0, R_0)$ , see also Remark 2.5 for more precisions.

We shall then investigate the existence of a time  $T > 0$  and a constant  $C > 0$  such that for all  $(y^0, y^1) \in H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))$  with  $y^0 = 0$  on  $S(R_1)$ , the corresponding solution of (1.5) satisfies:

$$\|(y^0, y^1)\|_{H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))} \leq C \|\partial_\nu y\|_{L^2(0, T; L^2(S(R_1)))}. \quad (1.6)$$

We are not aware of any result on that system in the literature.

**A simplified fluid structure model.** One might also consider non-local operators given as follows:

$$\begin{cases} \partial_{tt}y(t, x) - \Delta y(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ y(t, x) = 0, & \text{on } (0, T) \times S(R_1), \\ \partial_\nu y(t, x) = s'(t) \cdot \vec{\nu}, & \text{on } (0, T) \times S(R_0), \\ s''(t) + s(t) = - \int_{S(R_0)} \partial_t y(t, x) \vec{\nu} d\sigma, & \text{in } (0, T), \\ (y(0, \cdot), \partial_t y(0, \cdot), s(0), s'(0)) = (y^0, y^1, s^0, s^1), & \text{in } A(R_0, R_1) \times \mathbb{R}^2 \times \mathbb{R}^2, \end{cases} \quad (1.7)$$

where  $\vec{\nu}$  is the outward normal to  $A(R_0, R_1)$ . This models a fluid-structure interaction problem, see [9], for which  $y$  is the velocity potential of the fluid, and the function  $s$  corresponds to the displacement of an oscillator located in the ball  $B(0, R_0)$ .

Here, the relevant observability inequality reads as follows: there exist a time  $T > 0$  and a constant  $C > 0$  such that for all  $(y^0, y^1, s^0, s^1) \in H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1)) \times \mathbb{R}^2 \times \mathbb{R}^2$  with  $y^0 = 0$  on  $S(R_1)$ , the corresponding solution of (1.7) satisfies:

$$\|(y^0, y^1, s^0, s^1)\|_{H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1)) \times \mathbb{R}^2 \times \mathbb{R}^2} \leq C \|\partial_\nu y\|_{L^2(0, T; L^2(S(R_1)))}. \quad (1.8)$$

We refer to [19, 20, 11] for observability results on that model. In particular, it has been proved in [11] that this model is observable in any time  $T > 2 \sqrt{R_1^2 - R_0^2}$ .

## 1.2 Main results

To address observability issues for all the models presented above, our strategy relies on the fact that all the above examples can be written under an abstract form as

$$Y'(t) = AY(t), \quad t \geq 0, \quad Y(0) = Y^0, \quad (1.9)$$

where  $A$  is a skew-adjoint operator defined on a Hilbert space  $H$  and with domain  $\mathcal{D}(A)$ , and each one of the corresponding observability inequalities reads as follows: Does there exist a constant  $C > 0$  and a time  $T > 0$  such that for all  $Y^0 \in \mathcal{D}(A)$ , the solution  $Y$  of (1.9) satisfies

$$\|Y^0\|_H \leq C \|B^*Y\|_{L^2(0, T; U)}, \quad (1.10)$$

where  $B^*Y$  will correspond to  $\partial_\nu y|_{S(R_1)}$ , and  $U = L^2(S(R_1))$ .

In all the above cases, as we will justify later, the observation operator  $B^*$  is admissible (in the sense of [21, Section 4.3]) and satisfies  $B^* \in \mathcal{L}(\mathcal{D}(A), L^2(S(R_1)))$ . We shall keep the notation  $B^*$  as this observation operator can be thought as the dual of the control operator corresponding to non-homogeneous Dirichlet boundary conditions on the sphere  $L^2(S(R_1))$ , see [21, Section 11.2].

We shall then use a classical result (see for instance [7, 18]) which states that, under this setting, the observability inequality (1.10) is equivalent to the following resolvent estimate:

**Theorem 1.1** ([7, 18]). *Let  $A$  be an unbounded skew-adjoint operator on a Hilbert space  $H$  with domain  $\mathcal{D}(A)$ , and  $B^* \in \mathcal{L}(\mathcal{D}(A), U)$  be an admissible observation operator (in the sense of [21, Section 4.3]) for the group  $(e^{tA})_{t \in \mathbb{R}}$ .*

*Then there exist a time  $T > 0$  and a constant  $C > 0$  such that the observability inequality (1.10) holds for all solutions  $Y$  of (1.9) with initial datum in  $\mathcal{D}(A)$  if and only if there exists  $C > 0$  such that*

$$\|Y\|_H^2 \leq C^2 \|(A - i\omega)Y\|_H^2 + C^2 \|B^*Y\|_U^2, \quad (1.11)$$

for all  $\omega \in \mathbb{R}$ , and for all  $Y \in \mathcal{D}(A)$ .

In fact, our main results below give sufficient conditions on the boundary conditions to ensure that the resolvent estimate (1.11) holds.

In order to express those conditions, due to the radial symmetry of the problems we are considering, it is interesting to decompose  $Y$  using spherical harmonics, indexed by  $k \in \mathbb{Z}$ :

$$Y_k(r) = \frac{1}{2\pi r} \int_0^{2\pi} Y(r, \theta) e^{-ik\theta} r d\theta.$$

Then, as it will be made precise in Section 2, the resolvent condition (1.11) can be reduced to show estimates on the function  $y_k$  solution of

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = i\omega f_{1,k} + f_{2,k}, & \text{in } (R_0, R_1), \\ y_k(R_1) = 0, \\ R_0 \partial_r y_k(R_0) = \rho(\omega, k) y_k(R_0) + g_k, \end{cases} \quad (1.12)$$

where

$$\rho = \rho(\omega, k) : \mathbb{R} \times \mathbb{Z} \rightarrow \mathbb{R}, \quad (1.13)$$

is a kernel attached to the boundary conditions under considerations, and  $f_{1,k} \in H^1(R_0, R_1)$  with  $f_{1,k}(R_1) = 0$ ,  $f_{2,k} \in L^2(R_0, R_1)$  and  $g_k \in \mathbb{C}$ .

We will show in Section 2 that, in the examples listed in Section 1.1, the kernel  $\rho$  used within the corresponding resolvent estimates can be given explicitly:

- For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ ,  $\rho(\omega, k) = R_0 \alpha$  for the wave equation (1.1) with Fourier boundary conditions;
- For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ ,  $\rho(\omega, k) = R_0 \left( -\alpha \omega^2 + \beta \frac{k^2}{R_0^2} \right)$  for the wave equation (1.3) with waves on the boundary;
- For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ ,  $\rho(\omega, k) = R_0^{1-2s} |k|^{2s}$  for the wave equation (1.5) with fractional Laplacian on the boundary;
- For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z} \setminus \{(\pm 1, \pm 1), (\pm 1, \mp 1)\}$ ,  $\rho(\omega, k) = \frac{\pi R_0 \omega^2}{\omega^2 - 1} 1_{|k|=1}$  for the simplified fluid structure model (1.7). (Note that, in this last example,  $\rho$  is singular when  $|k| = |\omega| = 1$ .)

We are now in position to state our main result :

**Theorem 1.2.** *Assume that there exists  $M \geq 0$  such that*

- $\rho = \rho(\omega, k)$  is well-defined for  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying  $\omega^2 R_0^2 + k^2 \geq M^2$  and  $\omega^2 R_0^2 \leq k^2$  with values in  $\mathbb{R}$ ;
- there exist  $\varepsilon \in (0, 1)$ ,  $\mathbf{r} > 0$ , and  $\delta > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2 \geq M^2$  and

$$\log \left( \frac{|k|}{|\omega| R_0} \right) \geq \frac{\mathbf{r}}{|k|^{2/3}}, \quad (1.14)$$

we have

$$I_\varepsilon(\omega, k)^2 \geq \delta \left( (k^2 - \omega^2 R_0^2) - \rho(\omega, k)^2 - \rho(\omega, k) \right), \quad (1.15)$$

where  $I_\varepsilon(\omega, k)$  is defined as

$$I_\varepsilon(\omega, k) = \inf_{c \in [1-\varepsilon, 1]} \left| \rho(\omega, k) + c \sqrt{k^2 - \omega^2 R_0^2} \right|. \quad (1.16)$$

Then there exists a constant  $C > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  and any solution  $y_k$  of

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = i\omega f_{1,k} + f_{2,k}, & \text{in } (R_0, R_1), \\ y_k(R_1) = 0, \\ R_0 \partial_r y_k(R_0) = \rho(\omega, k) y_k(R_0) + g_k, & \text{if } \omega^2 R_0^2 \leq k^2 \text{ with } \omega^2 R_0^2 + k^2 \geq M^2, \end{cases} \quad (1.17)$$

with  $f_{1,k} \in H^1(R_0, R_1)$  satisfying  $f_{1,k}(R_1) = 0$ ,  $f_{2,k} \in L^2(R_0, R_1)$  and  $g_k \in \mathbb{C}$ , it holds

$$\begin{aligned} & \|\partial_r y_k\|_{L^2(R_0, R_1; r dr)} + |k| \|y_k/r\|_{L^2(R_0, R_1; r dr)} + |\omega| \|y_k\|_{L^2(R_0, R_1; r dr)} \\ & \leq C \left( \|\partial_r f_{1,k}\|_{L^2(R_0, R_1; r dr)} + |k| \|f_{1,k}/r\|_{L^2(R_0, R_1; r dr)} + \|f_{2,k}\|_{L^2(R_0, R_1; r dr)} \right. \\ & \quad \left. + |g_k| 1_{\omega^2 R_0^2 \leq k^2} + |\partial_r y_k(R_1)| \right). \end{aligned} \quad (1.18)$$

Furthermore, there exist constants  $C > 0$  and  $A > 0$  such that

$$\begin{aligned}
& 1_{\omega^2 R_0^2 + k^2 \leq M^2} (|\partial_r y_k(R_0)|^2 + |y_k(R_0)|^2) \\
& + 1_{\substack{\omega^2 R_0^2 + k^2 \geq M^2 \\ \omega^2 R_0^2 \geq k^2}} \left( |\partial_r y_k(R_0)|^2 + \max \left\{ \sqrt{k^2 + \omega^2 R_0^2}, \omega^2 R_0^2 - k^2 \right\} |y_k(R_0)|^2 \right) \\
& + 1_{\substack{\omega^2 R_0^2 + k^2 \geq M^2 \\ \omega^2 R_0^2 \leq k^2}} \left( I_\varepsilon(\omega, k)^2 1_{\log\left(\frac{|k|}{|\omega|R_0}\right) \geq \frac{A}{|k|^{2/3}}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} 1_{\log\left(\frac{|k|}{|\omega|R_0}\right) \leq \frac{A}{|k|^{2/3}}} \right) |y_k(R_0)|^2 \quad (1.19) \\
& \leq C \left( \|\partial_r f_{1,k}\|_{L^2(R_0, R_1; r dr)} + |k| \|f_{1,k}/r\|_{L^2(R_0, R_1; r dr)} + \|f_{2,k}\|_{L^2(R_0, R_1; r dr)} \right. \\
& \qquad \qquad \qquad \left. + |g_k| 1_{\omega^2 R_0^2 \leq k^2} + |\partial_r y_k(R_1)| \right)^2,
\end{aligned}$$

where, by analogy with (1.16),  $I_1 = I_1(\omega, k)$  is defined for  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 \leq k^2$  by

$$I_1(\omega, k) = \inf_{c \in [0,1]} \left| \rho(\omega, k) + c \sqrt{k^2 - \omega^2 R_0^2} \right|. \quad (1.20)$$

Condition (1.15) is a sufficient condition to get estimate (1.18), and of course it can be relaxed into a slightly more explicit formulation, proved in Appendix A:

**Corollary 1.3.** *With the same notations as in Theorem 1.2, all solutions  $y_k$  of (1.17) with  $f_{1,k} \in H^1(R_0, R_1)$  satisfying  $f_{1,k}(R_1) = 0$ ,  $f_{2,k} \in L^2(R_0, R_1)$  and  $g_k \in \mathbb{C}$  satisfy the estimates (1.18) and (1.19) if there exist  $\gamma > 0$ ,  $r > 0$  and  $M \geq 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying  $\omega^2 R_0^2 + k^2 \geq M^2$  and (1.14),  $\rho(\omega, k)$  satisfies*

$$\rho(\omega, k) \leq -\sqrt{k^2 - \omega^2 R_0^2} - \frac{1 + \gamma}{2} \quad \text{or} \quad \rho(\omega, k) \geq (-1 + \gamma) \sqrt{k^2 - \omega^2 R_0^2}. \quad (1.21)$$

In particular, if  $\rho(\omega, k) \geq 0$  for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying (1.14), estimates (1.18) and (1.19) hold for solutions of (1.17).

The condition (1.15) is very likely not optimal, but we believe that it is close to be, in the sense that Corollary 1.3 indicates that the resolvent estimate (1.18) holds as soon as we can guarantee that  $\rho(\omega, k)$  is far from  $-\sqrt{k^2 - \omega^2 R_0^2}$  for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2$  large and  $\omega^2 R_0^2 \leq k^2$ .

In fact, we show below the following necessary conditions for (3.3), which underlines that  $|\rho(\omega, k) + \sqrt{k^2 - \omega^2 R_0^2}|$  should be sufficiently large for (1.18) to hold:

**Theorem 1.4.** *Assume that there exists  $C > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying (1.14), any solution  $y_k$  of (1.17) with  $f_{1,k} \in H^1(R_0, R_1)$  satisfying  $f_{1,k}(R_1) = 0$ ,  $f_{2,k} \in L^2(R_0, R_1)$  and  $g_k \in \mathbb{C}$  satisfies the resolvent estimate (1.18).*

*Then, for all  $C_0 > 0$ , there exists  $C > 0$  such that for any sequence  $(\omega_n, k_n)_{n \in \mathbb{N}}$  of elements in  $\mathbb{R} \times \mathbb{Z}$  such that*

$$\lim_{n \rightarrow \infty} (\omega_n^2 R_0^2 + k_n^2) = \infty, \quad \lim_{n \rightarrow \infty} \frac{|k_n|^{2/3}}{\log^{2/3}(|k_n|)} \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) = \infty, \quad (1.22)$$

and

$$\forall n \in \mathbb{N}, \quad \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \leq C_0, \quad (1.23)$$

we have

$$\frac{k_n^2}{\sqrt{k_n^2 - \omega_n^2 R_0^2}} \leq C \left( \rho(\omega_n, k_n) + \sqrt{k_n^2 - \omega_n^2 R_0^2} \right)^2. \quad (1.24)$$

**Remark 1.5.** *Although it is not completely obvious at first glance, tedious computations show that, as expected, condition (1.15) implies that the sufficient condition of Theorem 1.4 is satisfied.*

### 1.3 Related results

There are many works on the observability of wave type equations, mainly triggered by the fact that this is dual to the controllability of such systems, see e.g. [17, 16], and of course we cannot give here a complete account of the theory, but only a rapid overview.

Among the several methods employed for the study of wave systems, the first one was probably the multiplier technique, due to [14] and popularized by [16], which was used to study wave equations with homogeneous Dirichlet boundary conditions in many situations, proving in particular observability in sufficiently large time when the observation set is a part of the boundary in the shadow of a light point, i.e. there exists  $x_0 \in \mathbb{R}^d$  such that the observation set contains  $\{x \in \partial\Omega, (x - x_0) \cdot \nu > 0\}$ . Such condition is usually called the multiplier condition or the  $\Gamma$ -condition of Lions. Note that this condition is satisfied in our setting, since the boundary  $S(R_1)$  is in the shadow from the light point  $x_0 = 0$ . Still, the multiplier condition is quite restrictive on the boundary conditions required at the sphere  $S(R_0)$ , and it does not allow to keep track precisely of what happens on the illuminated boundary  $S(R_0)$ . In particular, it is not clear at all how it can be applied with the generality we are considering here.

Another approach was then developed based on the analysis of the propagation of singularities and on microlocal analysis, yielding to the works [2, 3, 6], and giving a sharp geometric condition for the observation and control of waves, now known under the name of geometric control condition, and asserting that all the rays of geometric optics meet the boundary at a non-diffractive point. However, here again, the analysis close to the part of the boundary on which the control does not act strongly depends on the boundary conditions at hand, and the literature is mainly developed only in the case of Dirichlet or Neumann boundary conditions. In particular, one should exclude the possibility of getting rays captured within the internal sphere  $S(R_0)$ , i.e. solutions whose energy is localized in a neighborhood of  $S(R_0)$ , see also [10]. Besides, microlocal analysis is by construction limited to local operators, while some of the models we present above are not local - even if they become local after having been decomposed in Fourier components under the form (1.12).

Therefore, we believe that our approach is original and allows to attack the observability problem in a somewhat wider generality than what is known so far.

### 1.4 Outline

In Section 2, we explain how observability results for the wave models (1.1), (1.3), (1.5), (1.7) can be deduced from Theorem 1.1, Theorem 1.2, Corollary 1.3 and Theorem 1.4. Section 3 then focuses on the proof of Theorem 1.2. Theorem 1.4 is then proved in Section 4. Section 5 provides further comments. Finally, the proof of Corollary 1.3 is given in Appendix A.

## 2 Observability and non-observability of several wave models

The goal of this section is to explain the functional setting allowing to recast each of the wave equations (1.1), (1.3), (1.5), (1.7) into the abstract form (1.9), prove then that the corresponding resolvent estimate (1.11) reduces to the proof of estimates of the form (1.18) for a suitable kernel function  $\rho$ , and to deduce from them observability and non-observability results for each of the wave models presented in (1.1), (1.3), (1.5), (1.7).

### 2.1 The wave equation (1.1) with Fourier boundary conditions

In this subsection,  $\alpha$  is a fixed non-negative real number.

**Abstract form.** The wave equation (1.1) with Fourier boundary conditions writes as in (1.9) by setting

$$Y(t) = \begin{pmatrix} y(t) \\ \partial_t y(t) \end{pmatrix}, \quad A = \begin{pmatrix} 0 & I \\ -A_0 & 0 \end{pmatrix} \quad (2.1)$$

where  $A_0$  is the Laplace operator defined on  $L^2(\Omega)$  with domain  $\mathcal{D}(A_0) = \{y \in H^2(A(R_0, R_1)), \partial_\nu y + \alpha y = 0 \text{ on } S(R_0), y = 0 \text{ on } S(R_1)\}$  by  $A_0 y = -\Delta y$ , and  $A$  is defined on

$$H = \left\{ \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \text{ with } y_1 \in H^1(A(R_0, R_1)), y_2 \in L^2(A(R_0, R_1)) \text{ and } y_1 = 0 \text{ on } S(R_1) \right\},$$

with domain

$$\mathcal{D}(A) = \left\{ \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \text{ with } y_1 \in \mathcal{D}(A_0), y_2 \in H^1(A(R_0, R_1)) \text{ and } y_2 = 0 \text{ on } S(R_1) \right\}.$$

One then easily checks that  $A$  is skew adjoint on  $H$  provided  $H$  is endowed with the scalar production

$$\left\langle \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \begin{pmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{pmatrix} \right\rangle_H = \int_{A(R_0, R_1)} (\nabla y_1 \cdot \overline{\nabla \tilde{y}_1} + y_2 \overline{\tilde{y}_2}) dx + \alpha \int_{S(R_0)} y_1 \overline{\tilde{y}_1} d\sigma.$$

We then define  $B^*$  as follows:

$$B^* \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \partial_\nu y_1|_{S(R_1)}. \quad (2.2)$$

With this choice, we clearly have that  $B^* \in \mathcal{L}(\mathcal{D}(A), U)$ , with  $U = L^2(S(R_1))$ .

The fact that  $B^*$  is an admissible observation operator in the sense of [21, Section 4.3] follows immediately from classical multiplier arguments as in [16, Section I.4.1].

**Resolvent estimate.** In view of the above setting, the resolvent estimate (1.11) reads as follows. There exists a constant  $C > 0$  such that if  $(y_1, y_2)$  solves

$$\begin{cases} -i\omega y_1 + y_2 = f_1, & \text{in } A(R_0, R_1), \\ \Delta y_1 - i\omega y_2 = f_2, & \text{in } A(R_0, R_1), \\ y_1 = y_2 = 0, & \text{in } S(R_1), \\ \partial_r y_1 = \alpha y_1, & \text{on } S(R_0), \end{cases} \quad (2.3)$$

for some  $(f_1, f_2)^{\text{tr}} \in H$  and  $\omega \in \mathbb{R}$ , then

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y_1|^2 + |y_2|^2) dx + \alpha \int_{S(R_0)} |y_1|^2 d\sigma \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + \alpha \int_{S(R_0)} |f_1|^2 d\sigma \right) + C^2 \|\partial_r y_1\|_{L^2(S(R_1))}^2. \end{aligned} \quad (2.4)$$

Note that for  $(f_1, f_2)^{\text{tr}} \in H$ , using the boundary condition  $f_1 = 0$  on  $S(R_1)$ , Poincaré's estimate gives that

$$\int_{A(R_0, R_1)} |f_1|^2 dx \leq C \int_{A(R_0, R_1)} |\nabla f_1|^2 dx.$$

Accordingly, the above resolvent estimate is equivalent to show the existence of a constant  $C > 0$  such that if  $y$  (corresponding to  $y_1$ ) solves

$$\begin{cases} \omega^2 y + \Delta y = i\omega f_1 + f_2, & \text{in } A(R_0, R_1), \\ y = 0, & \text{in } S(R_1), \\ \partial_r y = \alpha y, & \text{on } S(R_0), \end{cases} \quad (2.5)$$

for some  $(f_1, f_2)^{\text{tr}} \in H$  and  $\omega \in \mathbb{R}$ , then

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y|^2 + |\omega|^2 |y|^2) dx + \alpha \int_{S(R_0)} |y|^2 d\sigma \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + \alpha \int_{S(R_0)} |f_1|^2 d\sigma \right) + C^2 \|\partial_r y\|_{L^2(S(R_1))}^2. \end{aligned} \quad (2.6)$$

Writing the equations (2.5) using the spherical harmonics decomposition, we immediately have that the above resolvent estimate is equivalent to show that there exists a constant  $C > 0$  such that for all  $k \in \mathbb{Z}$ , if  $y_k$  solves

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = i\omega f_{1,k} + f_{2,k}, & \text{in } (R_0, R_1), \\ y_k(R_1) = 0, \\ \partial_r y_k(R_0) = \alpha y_k(R_0), \end{cases} \quad (2.7)$$



for some  $(f_{1,k}, f_{2,k}) \in H^1(R_0, R_1) \times L^2(R_0, R_1)$  with  $f_{1,k}(R_1) = 0$  and  $\omega \in \mathbb{R}$ , then

$$\begin{aligned} & \int_{R_0}^{R_1} \left( |\partial_r y_k|^2 + \left( \frac{k^2}{r^2} + |\omega|^2 \right) |y_k|^2 \right) r dr + \alpha R_0 |y_k(R_0)|^2 \\ & \leq C^2 \left( \int_{R_0}^{R_1} \left( |\partial_r f_1|^2 + \frac{k^2}{r^2} |f_1|^2 + |f_2|^2 \right) r dr + \alpha R_0 |f_{1,k}(R_0)|^2 \right) + C^2 |\partial_r y_k(R_1)|^2. \end{aligned} \quad (2.8)$$

The equation satisfied by  $y_k$  in (2.7) thus corresponds to the equation (1.12) with the choice  $\rho(\omega, k) = R_0 \alpha$  as claimed in Section 1.2.

**Application of Theorem 1.2.** Since we assumed  $\alpha \geq 0$ , the function  $\rho$  is non-negative and thus the condition (1.15) is automatically satisfied. In particular, taking  $g_k = 0$  in (1.12), Theorem 1.2 implies that there exists a constant  $C > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  and any solution  $y_k$  of (2.7) with  $f_{1,k} \in H^1(R_0, R_1)$  satisfying  $f_{1,k}(R_1) = 0$  and  $f_{2,k} \in L^2(R_0, R_1)$ ,

$$\begin{aligned} & \|\partial_r y_k\|_{L^2(R_0, R_1; r dr)} + |k| \|y_k/r\|_{L^2(R_0, R_1; r dr)} + |\omega| \|y_k\|_{L^2(R_0, R_1; r dr)} \\ & \leq C \left( \|\partial_r f_{1,k}\|_{L^2(R_0, R_1; r dr)} + |k| \|f_{1,k}/r\|_{L^2(R_0, R_1; r dr)} + \|f_{2,k}\|_{L^2(R_0, R_1; r dr)} \right) + C |\partial_r y_k(R_1)|. \end{aligned} \quad (2.9)$$

Poincaré estimates then easily implies the resolvent estimate (2.8). Accordingly, as a corollary of Theorem 1.2 and Corollary 1.3, we get the following result:

**Theorem 2.1.** *Let  $\alpha \geq 0$ . Then the wave equation (1.1) with Fourier boundary conditions is observable in some time  $T > 0$ , i.e. there exists a time  $T > 0$  such that the observability inequality (1.2) is true for solutions of (1.1).*

## 2.2 The wave equation (1.3) with waves on the boundary

In this subsection,  $\alpha$  and  $\beta$  are fixed positive numbers.

**Abstract form.** Let us consider the functional space

$$H = \left\{ \begin{pmatrix} y_1 \\ y_2 \\ z_1 \\ z_2 \end{pmatrix}, y_1 \in H_1, y_2 \in L^2(A(R_0, R_1)), z_1 \in H^1(S(R_0)), z_2 \in L^2(S(R_0)), y_1|_{S(R_0)} = z_1 \right\},$$

where

$$H_1 = \{y \in H^1(A(R_0, R_1)), y|_{S(R_1)} = 0\},$$

endowed with the scalar product

$$\left\langle \begin{pmatrix} y_1 \\ y_2 \\ z_1 \\ z_2 \end{pmatrix}, \begin{pmatrix} \tilde{y}_1 \\ \tilde{y}_2 \\ \tilde{z}_1 \\ \tilde{z}_2 \end{pmatrix} \right\rangle_H = \int_{A(R_0, R_1)} (\nabla y_1 \cdot \overline{\nabla \tilde{y}_1} + y_2 \overline{\tilde{y}_2}) dx + \beta \int_{S(R_0)} \nabla_{S(R_0)} z_1 \cdot \overline{\nabla_{S(R_0)} \tilde{z}_1} d\sigma + \alpha \int_{S(R_0)} z_2 \overline{\tilde{z}_2} d\sigma,$$

where  $\nabla_{S(R_0)} = \frac{\partial_\theta}{R_0}$  is the gradient operator on the sphere  $S(R_0)$ . One easily checks that  $H$  is a Hilbert space.

We then set the operator

$$A : \begin{pmatrix} y_1 \\ y_2 \\ z_1 \\ z_2 \end{pmatrix} \in \mathcal{D}(A) \subset H \mapsto \begin{pmatrix} y_2 \\ \Delta y_1 \\ z_2 \\ \frac{\beta}{\alpha} \Delta_{S(R_0)} z_1 - \frac{1}{\alpha} \partial_\nu y_1 \end{pmatrix} \in H,$$

with domain  $\mathcal{D}(A)$  corresponding to the set

$$\left\{ \begin{pmatrix} y_1 \\ y_2 \\ z_1 \\ z_2 \end{pmatrix} \in H, \Delta y_1 \in L^2(A(R_0, R_1)), y_2 \in H_1, z_1 \in H^2(S(R_0)), z_2 \in H^1(S(R_0)), y_2|_{S(R_0)} = z_2 \right\}.$$

Note that for  $(y_1, y_2, z_1, z_2)^{\text{tr}} \in \mathcal{D}(A)$ , as  $y_1$  belongs to  $H^1(A(R_0, R_1))$ ,  $\Delta y_1$  is in  $L^2(A(R_0, R_1))$ , and  $y_1 = z_1 \in H^2(S(R_0))$ , by elliptic regularity  $\partial_\nu y_1$  belongs to  $L^2(S(R_0))$ , hence  $A$  is well-defined. Furthermore,  $A$  is skew-adjoint.

Then, if we consider the unknown

$$Y(t) = \begin{pmatrix} y(t) \\ \partial_t y(t) \\ z(t) \\ \partial_t z(t) \end{pmatrix},$$

Problem (1.3) rewrites in the abstract form  $Y' = AY$ , that is in the form of (1.9).

We can then define  $B^*$  as follows:

$$B^* \begin{pmatrix} y_1 \\ y_2 \\ z_1 \\ z_2 \end{pmatrix} = \partial_\nu y_1|_{S(R_1)}.$$

With this choice, we clearly have that  $B^* \in \mathcal{L}(\mathcal{D}(A), U)$ , with  $U = L^2(S(R_1))$ , and again,  $B^*$  is an admissible observation operator in the sense of [21, Section 4.3] from classical multiplier arguments (see [16, Section I.4.1]).

**Resolvent estimate.** In our context, the resolvent estimate (1.11) reads as follows. There exists a constant  $C > 0$  such that if  $(y_1, y_2, z_1, z_2)^{\text{tr}} \in \mathcal{D}(A)$  verifies

$$\begin{cases} -i\omega y_1 + y_2 = f_1 & \text{in } A(R_0, R_1), \\ \Delta y_1 - i\omega y_2 = f_2 & \text{in } A(R_0, R_1), \\ -i\omega z_1 + z_2 = g_1 & \text{on } S(R_0), \\ \frac{\beta}{\alpha} \Delta_{S(R_0)} z_1 - \frac{1}{\alpha} \partial_\nu y_1 - i\omega z_2 = g_2 & \text{on } S(R_0), \end{cases} \quad (2.10)$$

with  $(f_1, f_2, g_1, g_2)^{\text{tr}} \in H$  and  $\omega \in \mathbb{R}$ , one has

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y_1|^2 + |y_2|^2) dx + \beta \int_{S(R_0)} |\nabla_{S(R_0)} z_1|^2 d\sigma + \alpha \int_{S(R_0)} |z_2|^2 d\sigma \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + \beta \int_{S(R_0)} |\nabla_{S(R_0)} g_1|^2 d\sigma + \alpha \int_{S(R_0)} |g_2|^2 d\sigma \right) \\ & \quad + C^2 \int_{S(R_1)} |\partial_\nu y_1|^2 d\sigma. \end{aligned}$$

Now, using that by definition of  $H$ ,  $y_1 = z_1$  on  $S(R_0)$ , system (2.10) reduces to, with  $y \in H_1$  (corresponding to  $y_1$ ),

$$\begin{cases} \Delta y + \omega^2 y = f_2 + i\omega f_1 & \text{in } A(R_0, R_1), \\ \partial_\nu y - \beta \Delta_{S(R_0)} y - \alpha \omega^2 y = \alpha(g_2 + i\omega g_1) & \text{on } S(R_0), \end{cases} \quad (2.11)$$

and the desired resolvent estimate is equivalent to the existence of a constant  $C > 0$  such that

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y|^2 + |\omega|^2 |y|^2) dx + \beta \int_{S(R_0)} |\nabla_{S(R_0)} y|^2 d\sigma + \alpha \int_{S(R_0)} |\omega|^2 |y|^2 d\sigma \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + \beta \int_{S(R_0)} |\nabla_{S(R_0)} g_1|^2 d\sigma + \alpha \int_{S(R_0)} |g_2|^2 d\sigma \right) \\ & \quad + C^2 \int_{S(R_1)} |\partial_\nu y|^2 d\sigma. \end{aligned}$$

Decomposing our new problem on spherical harmonics leads to the following system, with  $f_{1,k} \in H^1(R_0, R_1)$  such that  $f_{1,k}(R_1) = 0$ , and  $f_{2,k} \in L^2(R_0, R_1)$ :

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = f_{2,k} + i\omega f_{1,k} & \text{in } (R_0, R_1), \\ y_k(R_1) = 0 \\ \partial_r y_k(R_0) = \left( \beta \frac{k^2}{R_0^2} - \alpha \omega^2 \right) y_k(R_0) + \alpha (g_{2,k} + i\omega g_{1,k}), \end{cases} \quad (2.12)$$

that is equation (1.12) with the choice

$$\rho(\omega, k) = R_0 \left( \beta \frac{k^2}{R_0^2} - \alpha \omega^2 \right)$$

as claimed in Section 1.2. Hence, our goal is to find a constant  $C > 0$  such that for all  $k \in \mathbb{Z}$ , all  $\omega \in \mathbb{R}$ , each solution of (2.12) satisfies the estimate

$$\begin{aligned} & \int_{R_0}^{R_1} \left( |\partial_r y_k|^2 + \left( \frac{k^2}{r^2} + \omega^2 \right) |y_k|^2 \right) r \, dr + \left( \beta \frac{k^2}{R_0^2} + \alpha \omega^2 \right) |y_k(R_0)|^2 \\ & \leq C^2 \left( \int_{R_0}^{R_1} \left( |\partial_r f_{1,k}|^2 + \frac{k^2}{r^2} |f_{1,k}|^2 + |f_{2,k}|^2 \right) r \, dr + \beta \frac{k^2}{R_0^2} |g_{1,k}|^2 + \alpha |g_{2,k}|^2 \right) + C^2 |\partial_r y_k(R_1)|^2. \end{aligned} \quad (2.13)$$

**Application of Theorem 1.2: The case  $0 < \alpha < \beta$ .** As an application of Theorem 1.2 and Corollary 1.3, we get the following positive result:

**Theorem 2.2.** *Let  $\alpha$  and  $\beta$  be positive constants with  $\alpha < \beta$ . Then the wave equation (1.3) satisfies the observability inequality (1.4) in some time  $T > 0$ .*

*Proof.* For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 \leq k^2$  we have

$$\rho(\omega, k) = \frac{1}{R_0} (\beta k^2 - \alpha \omega^2 R_0^2) \geq \frac{\beta k^2}{R_0} \left( 1 - \frac{\alpha}{\beta} \right), \quad (2.14)$$

Therefore, for  $\alpha < \beta$ , for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2 \geq M^2$  and  $\omega^2 R_0^2 \leq k^2$ ,  $\rho(\omega, k) \geq 0$ . Corollary 1.3 then applies and the resolvent estimates (1.18)–(1.19) hold, which imply in particular the existence of  $C > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ ,

$$\begin{aligned} & \int_{R_0}^{R_1} \left( |\partial_r y_k|^2 + \left( \frac{k^2}{r^2} + \omega^2 \right) |y_k|^2 \right) r \, dr \\ & \leq C^2 \left( \int_{R_0}^{R_1} \left( |\partial_r f_{1,k}|^2 + \frac{k^2}{r^2} |f_{1,k}|^2 + |f_{2,k}|^2 \right) r \, dr + \beta \frac{k^2}{R_0^2} |g_{1,k}|^2 + \alpha |g_{2,k}|^2 \right) + C^2 |\partial_r y_k(R_1)|^2. \end{aligned} \quad (2.15)$$

It remains to check that the boundary estimates (1.19) indeed yields that the boundary term

$$\left( \beta \frac{k^2}{R_0^2} + \alpha \omega^2 \right) |y_k(R_0)|^2 \quad (2.16)$$

in the left hand-side of (2.13) is bounded by the right hand-side of (2.13), which is of course delicate only for  $\omega^2 R_0^2 + k^2$  large enough.

For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 \leq k^2$ , due to (2.14),

$$I_\varepsilon(\omega, k)^2 \geq I_1(\omega, k)^2 \geq \frac{\beta^2 k^4}{R_0^2} \left( 1 - \frac{\alpha}{\beta} \right)^2.$$

On the other hand, for  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 \geq k^2$  and  $\omega^2 R_0^2 \geq 1$ , taking into account the boundary condition on  $R_0$  of system (2.12), we have

$$\left( \beta \frac{k^2}{R_0^2} - \alpha \omega^2 \right)^2 |y_k(R_0)|^2 \leq 2 |\partial_r y_k(R_0)|^2 + 2 |\alpha (g_{2,k} + i\omega g_{1,k})|^2,$$

so that

$$\frac{k^2}{\omega^2} \left( \beta \frac{k^2}{R_0^2} - \alpha \omega^2 \right)^2 |y_k(R_0)|^2 \leq C |\partial_r y_k(R_0)|^2 + C |g_{2,k}|^2 + C k^2 |g_{1,k}|^2,$$

Accordingly, the estimate on the boundary terms at  $R_0$  in (1.19) corresponding to the case  $\omega^2 R_0^2 \geq k^2$  and  $\omega^2 R_0^2 + k^2 \geq 2R_0^2$  (implying  $\omega^2 \geq 1$ ) allows to estimate

$$\begin{aligned} & \left( \frac{k^2}{\omega^2} \left( \beta \frac{k^2}{R_0^2} - \alpha \omega^2 \right)^2 + \max \left\{ \sqrt{k^2 + \omega^2 R_0^2}, \omega^2 R_0^2 - k^2 \right\} \right) |y_k(R_0)|^2 \\ & \geq \omega^2 R_0^2 \left( \alpha^2 \frac{k^2}{\omega^2 R_0^2} \left( 1 - \frac{\beta}{\alpha} \frac{k^2}{\omega^2 R_0^2} \right)^2 + 1 - \frac{k^2}{\omega^2 R_0^2} \right) |y_k(R_0)|^2 \\ & \geq \omega^2 R_0^2 \inf_{\tau \in [0,1]} \left\{ \alpha^2 \tau^2 \left( 1 - \frac{\beta}{\alpha} \tau^2 \right)^2 + 1 - \tau^2 \right\} |y_k(R_0)|^2 \geq c_* \omega^2 R_0^2 |y_k(R_0)|^2, \end{aligned}$$

for some  $c_* > 0$ .

The previous inequalities and the boundary estimates (1.19) show that the boundary term (2.16) is bounded by the right hand-side of (2.13) with a constant independent of  $(\omega, k)$ .  $\square$

**Remark 2.3.** When  $\alpha = \beta$ , Theorem 1.2 still applies and the above arguments immediately yield (2.15). However, the boundary estimates (1.19) fail to give estimate on (2.16) by the right hand-side of (2.13), in particular in the range  $\omega^2 R_0^2 \simeq k^2$ .

**Application of Theorem 1.4: The case  $0 < \beta < \alpha$ .** As a consequence of Theorem 1.4, we show the following result:

**Theorem 2.4.** Let  $\alpha$  and  $\beta$  be positive constants with  $\alpha > \beta$ . Then the wave equation (1.3) does not satisfy the observability inequality (1.4) in any time  $T > 0$ .

*Proof.* We use Theorem 1.4 and construct a sequence  $(\omega_n, k_n) \in \mathbb{R} \times \mathbb{Z}$  satisfying (1.22)–(1.23) and for which (1.24) cannot hold.

We set  $A = \beta/R_0$  and  $\gamma = \alpha/\beta$ , which satisfies  $\gamma > 1$  from the assumption  $\alpha > \beta$ . With these notations  $\rho$  writes  $\rho(\omega, k) = A(k^2 - \gamma \omega^2 R_0^2)$ .

Then, for  $n \in \mathbb{N}$ , we set

$$k_n = n, \quad \omega_n = \frac{1}{R_0} \sqrt{n^2 - \frac{1}{4\gamma^2 A^2} \left( \sqrt{1 + 4n^2 \gamma (\gamma - 1) A^2} - 1 \right)^2}.$$

This choice is done to guarantee that

$$\forall n \in \mathbb{N}, \quad \rho(\omega_n, k_n) + \sqrt{k_n^2 - \omega_n^2 R_0^2} = 0. \quad (2.17)$$

It is also easy to check that

$$k_n^2 - \omega_n^2 R_0^2 = \frac{1}{4\gamma^2 A^2} \left( \sqrt{1 + 4n^2 \gamma (\gamma - 1) A^2} - 1 \right)^2 \leq \frac{\gamma - 1}{\gamma} k_n^2,$$

hence that

$$\forall n \in \mathbb{N}, \quad 1 \leq \frac{|k_n|}{|\omega_n| R_0} \leq \sqrt{\gamma}, \quad \lim_{n \rightarrow \infty} \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) = \frac{1}{2} \log(\gamma) > 0,$$

and  $\lim_{n \rightarrow \infty} \frac{\sqrt{k_n^2 - \omega_n^2 R_0^2}}{|k_n|} = \sqrt{1 - \frac{1}{\gamma}}.$

Accordingly, we easily get that the sequence  $(\omega_n, k_n)_{n \in \mathbb{N}}$  satisfies (1.22)–(1.23), while the relation (2.17) disproves estimate (1.24). This concludes the proof of Theorem 2.4.  $\square$

### 2.3 The wave equation (1.5) with a fractional laplacian

In this subsection,  $s$  denotes a positive constant in  $(0, 1]$ .

**Abstract form.** Let  $f \in L^2(S(R_0))$ . We denote  $f_k$  its  $k$ -th Fourier coefficient, that is

$$f_k = \frac{1}{2\pi} \int_0^{2\pi} f(\theta) e^{-ik\theta} d\theta,$$

which implies the expansion  $f(\theta) = \sum_{k \in \mathbb{Z}} f_k e^{ik\theta}$ ,

For  $s \in (0, 1]$ , we define the usual fractional Laplace-Beltrami operator

$$(-\Delta_{S(R_0)})^s : f \in \mathcal{D}((-\Delta_{S(R_0)})^s) \subset L^2(S(R_0)) \mapsto (-\Delta_{S(R_0)})^s f \in L^2(S(R_0)),$$

by

$$\mathcal{D}((-\Delta_{S(R_0)})^s) = \left\{ f \in L^2(S(R_0)), \sum_{k \in \mathbb{Z}} |k|^{4s} |f_k|^2 < \infty \right\},$$

and,

$$((-\Delta_{S(R_0)})^s f)(\theta) = \frac{1}{R_0^{2s}} \sum_{k \in \mathbb{Z}} |k|^{2s} f_k e^{ik\theta}.$$

We then define

$$H = \left\{ \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, y_1 \in H^1(A(R_0, R_1)), y_1|_{S(R_0)} \in \mathcal{D}((-\Delta_{S(R_0)})^{s/2}), y_1(R_1) = 0, y_2 \in L^2(A(R_0, R_1)) \right\},$$

which is an Hilbert space when equipped with the scalar product

$$\left\langle \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \begin{pmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{pmatrix} \right\rangle_H = \int_{\Omega} (\nabla y_1 \cdot \overline{\nabla \tilde{y}_1} + y_2 \overline{\tilde{y}_2}) dx + 2\pi R_0^{1-2s} \sum_{k \in \mathbb{Z}} |k|^{2s} y_1(R_0)_k \overline{\tilde{y}_1(R_0)_k}.$$

We also define

$$A : \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \in \mathcal{D}(A) \subset H \mapsto \begin{pmatrix} y_2 \\ \Delta y_1 \end{pmatrix} \in H,$$

with

$$\mathcal{D}(A) := \left\{ \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \in H, \Delta y_1 \in L^2(A(R_0, R_1)), \partial_\nu y_1 + (-\Delta_{S(R_0)})^s(y_1|_{S(R_0)}) = 0 \text{ on } S(R_0) \right. \\ \left. y_2|_{S(R_0)} \in \mathcal{D}((-\Delta_{S(R_0)})^{s/2}), y_2 \in H^1(A(R_0, R_1)), y_2(R_1) = 0 \right\}.$$

It is not difficult to check that the operator  $A$  is skew-adjoint on  $H$ . Therefore, if we consider the unknown  $Y(t) = \begin{pmatrix} y(t) \\ y_t(t) \end{pmatrix}$ , Problem (1.5) rewrites in the abstract form  $Y' = AY$ , that is in the form of (1.9).

The observation operator  $B^*$  can then be deduced on  $H$  by the same formula as in (2.2), and again,  $B^* \in \mathcal{L}(\mathcal{D}(A), U)$ , with  $U = L^2(S(R_1))$ , and  $B^*$  is an admissible observation operator in the sense of [21, Section 4.3] due to classical multiplier arguments as in [16, Section I.4.1].

**Resolvent estimate.** In view of the above setting, the resolvent estimate (1.11) reads as follows. There exists a constant  $C > 0$  such that if  $(y_1, y_2)$  solves

$$\begin{cases} -i\omega y_1 + y_2 = f_1, & \text{in } A(R_0, R_1), \\ \Delta y_1 - i\omega y_2 = f_2, & \text{in } A(R_0, R_1), \\ y_1 = y_2 = 0, & \text{in } S(R_1), \\ \partial_r y_1 = -(-\Delta_{S(R_0)})^s(y_1|_{S(R_0)}), & \text{on } S(R_0), \end{cases} \quad (2.18)$$

for some  $(f_1, f_2)^{\text{tr}} \in H$  and  $\omega \in \mathbb{R}$ , then

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y_1|^2 + |y_2|^2) dx + 2\pi R_0^{1-2s} \sum_{k \in \mathbb{Z}} |k|^{2s} |y_1(R_0)_k|^2 \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + 2\pi R_0^{1-2s} \sum_{k \in \mathbb{Z}} |k|^{2s} |f_1(R_0)_k|^2 \right) + C^2 \|\partial_r y_1\|_{L^2(S(R_1))}^2. \end{aligned} \quad (2.19)$$

Eliminating  $y_2$ , and writing  $y$  for  $y_1$  leads to the equivalent simplified system

$$\begin{cases} \Delta y + \omega^2 y = f_2 + i\omega f_1, & \text{in } A(R_0, R_1), \\ y = 0, & \text{in } S(R_1), \\ \partial_r y = -(-\Delta_{S(R_0)})^s (y|_{S(R_0)}), & \text{on } S(R_0), \end{cases}$$

for which we aim to prove the following equivalent estimate

$$\begin{aligned} & \int_{A(R_0, R_1)} (|\nabla y|^2 + |\omega|^2 |y|^2) dx + 2\pi R_0^{1-2s} \sum_{k \in \mathbb{Z}} |k|^{2s} |y(R_0)_k|^2 \\ & \leq C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) dx + 2\pi R_0^{1-2s} \sum_{k \in \mathbb{Z}} |k|^{2s} |f_1(R_0)_k|^2 \right) + C^2 \|\partial_r y\|_{L^2(S(R_1))}^2. \end{aligned} \quad (2.20)$$

As usual, we now decompose our problem on spherical harmonics. This leads to the following problem

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = f_{2,k} + i\omega f_{1,k} & \text{in } (R_0, R_1), \\ y_k(R_1) = 0, \\ \partial_r y_k(R_0) = \frac{|k|^{2s}}{R_0^{2s}} y_k(R_0), \end{cases} \quad (2.21)$$

with  $f_{1,k} \in H^1(R_0, R_1)$ ,  $f_{1,k}(R_1) = 0$ ,  $f_{2,k} \in L^2(R_0, R_1)$ , for which we aim to prove the existence of  $C > 0$  independent of  $k$  and  $\omega$  such that

$$\begin{aligned} & \int_{R_0}^{R_1} \left( |\partial_r y_k|^2 + \left( \frac{k^2}{r^2} + \omega^2 \right) |y_k|^2 \right) r dr + |k|^{2s} |y_k(R_0)|^2 \\ & \leq C^2 \left( \int_{R_0}^{R_1} \left( |\partial_r f_{1,k}|^2 + \frac{k^2}{r^2} |f_{1,k}|^2 + |f_{2,k}|^2 \right) r dr + |k|^{2s} |f_{1,k}(R_0)|^2 \right) + C^2 |\partial_r y_k(R_1)|^2. \end{aligned} \quad (2.22)$$

In other words, we are back to the study of (1.12) with

$$\rho(\omega, k) = R_0^{1-2s} |k|^{2s}.$$

**Remark 2.5** (The case  $s = 1/2$ ). The operator  $(-\Delta_{S(R_0)})^{1/2}$  actually corresponds to a Dirichlet to Neumann map for the Laplacian in  $B(0, R_0)$ . Namely, if  $h \in \mathcal{D}((-\Delta_{S(R_0)})^{1/2})$ , the solution  $y$  of

$$\Delta y = 0 \text{ in } B(0, R_0), \text{ with } y = h \text{ on } S(R_0),$$

corresponding to  $h(\theta) = \sum_k h_k e^{ik\theta}$ , is explicitly given by

$$y(r, \theta) = \sum_{k \in \mathbb{Z}} h_k \left( \frac{r}{R_0} \right)^{|k|} e^{ik\theta}, \quad \text{in } B(0, R_0),$$

so that  $\partial_r y|_{S(R_0)} = (-\Delta_{S(R_0)})^{1/2} h$ .

## Application of Theorem 1.2.

**Theorem 2.6.** *Let  $s \in (0, 1]$ . The wave equation (1.5) is observable in some time  $T > 0$ .*

*Proof.* Since  $\rho(\omega, k) \geq 0$  for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ , we can apply Corollary 1.3 and deduce the resolvent estimates (1.18) and (1.19) for solutions of (2.21).

Therefore, the only thing to check is that the strength of the boundary terms in (1.19) indeed allows to estimate  $|k|^{2s}|y_k(R_0)|^2$  by the right hand-side of (2.22). As before, this is delicate only when  $\omega^2 R_0^2 + k^2$  is large enough. Then, for  $\omega^2 R_0^2 \geq k^2$ , we simply use that

$$|\partial_r y_k(R_0)|^2 = R_0^{-4s} |k|^{4s} |y_k(R_0)|^2.$$

For  $\omega^2 R_0^2 \leq k^2$ , we use that, since  $\rho(\omega, k) = R_0^{1-2s} |k|^{2s} \geq 0$ ,

$$I_\varepsilon(\omega, k)^2 \geq I_1(\omega, k)^2 \geq R_0^{2-4s} |k|^{4s}.$$

According to the two above estimates, we obviously get the estimate on  $|k|^{2s}|y_k(R_0)|^2$  by the right hand-side of (2.22).  $\square$

## 2.4 The simplified fluid structure model (1.7)

**Abstract form.** We consider the functional space

$$H = \left\{ \begin{pmatrix} y_1 \\ y_2 \\ s_1 \\ s_2 \end{pmatrix}, s_1 \in \mathbb{C}^2, s_2 \in \mathbb{C}^2, y_1 \in H^1(A(R_0, R_1)), y_1|_{S(R_1)} = 0, y_2 \in L^2(A(R_0, R_2)) \right\},$$

endowed with the scalar product

$$\left\langle \begin{pmatrix} y_1 \\ y_2 \\ s_1 \\ s_2 \end{pmatrix}, \begin{pmatrix} \tilde{y}_1 \\ \tilde{y}_2 \\ \tilde{s}_1 \\ \tilde{s}_2 \end{pmatrix} \right\rangle_H = \int_{A(R_0, R_1)} (\nabla y_1 \cdot \overline{\nabla \tilde{y}_1} + y_2 \overline{\tilde{y}_2}) dx + s_1 \cdot \overline{\tilde{s}_1} + s_2 \cdot \overline{\tilde{s}_2},$$

which obviously is a Hilbert space.

We then set the operator

$$A : \begin{pmatrix} y_1 \\ y_2 \\ s_1 \\ s_2 \end{pmatrix} \in \mathcal{D}(A) \subset H \mapsto \left( y_2, \Delta y_1, s_2, - \left( \int_{S(R_0)} y_2 \nu d\sigma \right) - s_1 \right) \in H,$$

with domain  $\mathcal{D}(A)$  corresponding to the set

$$\left\{ \begin{pmatrix} y_1 \\ y_2 \\ s_1 \\ s_2 \end{pmatrix} \in H, \Delta y_1 \in L^2(A(R_0, R_1)), y_2 \in H^1(A(R_0, R_1)), y_2|_{S(R_1)} = 0, \partial_\nu y_1 = s_2 \cdot \nu \text{ on } S(R_0) \right\}.$$

Then, if we consider the unknown  $Y(t) = (y, y_t, s, s_t)^{\text{tr}}$ , problem (1.7) rewrites in the abstract form  $Y' = AY$ , that is in the form of (1.9),  $A$  being a skew-adjoint operator on  $H$ .

Similarly as before, we define the operator

$$B^* \begin{pmatrix} y_1 \\ y_2 \\ s_1 \\ s_2 \end{pmatrix} = \partial_\nu y_1|_{S(R_1)}.$$

With this choice, we clearly have that  $B^* \in \mathcal{L}(\mathcal{D}(A), U)$ , with  $U = L^2(S(R_1))$ , and that  $B^*$  is an admissible observation operator in the sense of [21, Section 4.3] according to classical multiplier arguments as in [16, Section I.4.1].

**Resolvent estimate.** In our context, the resolvent estimate (1.11) reads as follows. There exists a constant  $C > 0$  such that if  $(y_1, y_2, s_1, s_2)^{\text{tr}}$  in  $\mathcal{D}(A)$  verifies

$$\begin{cases} -i\omega y_1 + y_2 = f_1 & \text{in } A(R_0, R_1), \\ \Delta y_1 - i\omega y_2 = f_2 & \text{in } A(R_0, R_1), \\ -i\omega s_1 + s_2 = g_1 & \text{on } S(R_0), \\ -s_1 - \left( \int_{S(R_0)} y_2 \nu \, d\sigma \right) - i\omega s_2 = g_2 & \text{on } S(R_0), \end{cases} \quad (2.23)$$

with  $(f_1, f_2, g_1, g_2)^{\text{tr}}$  in  $H$  and  $\omega$  in  $\mathbb{R}$ , one has

$$\begin{aligned} \int_{A(R_0, R_1)} (|\nabla y_1|^2 + |y_2|^2) \, dx + |s_1|^2 + |s_2|^2 &\leq C^2 \int_{S(R_1)} |\partial_\nu y_1|^2 \, d\sigma \\ &+ C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) \, dx + |g_1|^2 + |g_2|^2 \right). \end{aligned}$$

We already note that the condition  $\partial_\nu y_1 = s_2 \cdot \nu$  on  $S(R_0)$ , when projected on the cylindrical harmonics, immediately gives that  $\partial_r y_{1,k}|_{S(R_0)} = 0$  for all  $k$  but  $-1$  and  $1$ . Conversely, it is not difficult to see that if  $w \in L^2(S(R_0))$  verifies  $w_k = 0$  for all  $k$  but  $-1$  and  $1$ , then there exists  $s \in \mathbb{C}^2$  such that  $w = s \cdot \nu$ . Accordingly  $|s_2|$  is of the order of the  $L^2(S(R_0))$ -norm of  $\partial_\nu y_1|_{S(R_0)}$ .

Similarly, the term

$$\left| \int_{S(R_0)} y_2 \nu \, d\sigma \right|$$

is of the order of  $|y_{2,1}(R_0)| + |y_{2,-1}(R_0)|$ . From the third and fourth lines of (2.23), we can thus get

$$|s_1| \leq |s_2| + |g_1| \quad \text{if } |\omega| \geq 1, \quad |s_1| \leq |s_2| + |g_2| + C|y_{2,1}(R_0)| \quad \text{if } |\omega| \leq 1.$$

Eliminating all unknowns but  $y = y_1$  in system (2.23), we obtain the following reduced system

$$\begin{cases} \Delta y + \omega^2 y = f_2 + i\omega f_1 & \text{in } A(R_0, R_1), \\ (1 - \omega^2) \partial_\nu y - \omega^2 \left( \int_{S(R_0)} y \nu \, d\sigma \right) \cdot \nu = (g_1 - i\omega g_2) \cdot \nu - i\omega \left( \int_{S(R_0)} f_1 \nu \, d\sigma \right) \cdot \nu & \text{on } S(R_0). \end{cases} \quad (2.24)$$

and the desired resolvent estimate is thus equivalent to the existence of a constant  $C > 0$  independent of  $\omega$  such that

$$\begin{aligned} \int_{A(R_0, R_1)} (|\nabla y|^2 + |\omega|^2 |y|^2) \, dx + \int_{S(R_0)} |\partial_\nu y|^2 \, d\sigma + |y_1(R_0)|^2 + |y_{-1}(R_0)|^2 \\ \leq C^2 \int_{S(R_1)} |\partial_\nu y|^2 \, d\sigma + C^2 \left( \int_{A(R_0, R_1)} (|\nabla f_1|^2 + |f_2|^2) \, dx + |g_1|^2 + |g_2|^2 \right). \end{aligned} \quad (2.25)$$

Decomposing our new problem on spherical harmonics leads to the following system, with  $f_{1,k} \in H^1(R_0, R_1)$  such that  $f_{1,k}(R_1) = 0$ , and  $f_{2,k} \in L^2(R_0, R_1)$ :

$$\begin{cases} \omega^2 y_k + \frac{1}{r} \partial_r (r \partial_r y_k) - \frac{k^2}{r^2} y_k = f_{2,k} + i\omega f_{1,k} & \text{in } (R_0, R_1), \\ y_k(R_1) = 0, \\ \partial_r y_k(R_0) = 0 & \text{if } k \text{ in } \mathbb{Z} \setminus \{-1, 1\}, \\ (\omega^2 - 1) \partial_r y_1(R_0) - \pi R_0 \omega^2 y_1(R_0) = \frac{1}{2} (i\omega g_2 - g_1) \cdot \begin{pmatrix} 1 \\ -i \end{pmatrix} - \pi i R_0 f_{1,1}(R_0), & \text{if } k = 1, \\ (\omega^2 - 1) \partial_r y_{-1}(R_0) - \pi R_0 \omega^2 y_{-1}(R_0) = \frac{1}{2} (i\omega g_2 - g_1) \cdot \begin{pmatrix} 1 \\ i \end{pmatrix} - \pi i R_0 f_{1,-1}(R_0), & \text{if } k = -1, \end{cases} \quad (2.26)$$

that is system equation (1.12) with the choice

$$\rho(\omega, k) = \pi R_0 \frac{\omega^2}{\omega^2 - 1} \delta_{k=\pm 1}$$



as claimed in Section 1.2. Hence, our goal is to find a constant  $C > 0$  such that for all  $\omega \in \mathbb{R}$  and all  $k \in \mathbb{Z}$ , each solution of (2.26) satisfies the estimate

$$\begin{aligned} & \int_{R_0}^{R_1} \left( |\partial_r y_k|^2 + \left( \frac{k^2}{r^2} + \omega^2 \right) |y_k|^2 \right) r \, dr + \mathbf{1}_{k=\pm 1} \left( |\partial_r y_k(R_0)|^2 + |y_k(R_0)|^2 \right) \\ & \leq C^2 \left( \int_{R_0}^{R_1} \left( |\partial_r f_{1,k}|^2 + \frac{k^2}{r^2} |f_{1,k}|^2 + |f_{2,k}|^2 \right) r \, dr + |g_1|^2 + |g_2|^2 \right) + C^2 |\partial_r y_k(R_1)|^2. \end{aligned} \quad (2.27)$$

**Application of Theorem 1.2.** In this section, we prove the following result:

**Theorem 2.7.** *There exists a time  $T$  such that the equation (1.7) is observable in time  $T > 0$ .*

*Proof.* For  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2 \geq 2(R_0^2 + 1)$ , we necessarily have  $\rho(\omega, k) \geq 0$  and that  $\rho(\omega, k)$  is bounded, since either  $|k| > 1$  or  $|\omega| \geq \sqrt{(2R_0^2 + 1)}/R_0 > 1$ . Using Corollary 1.3, Theorem 1.2 applies. Estimate (1.18) immediately yields the estimates on the integrated terms, and it thus remains to check that the estimates (1.19) yields the accurate estimates on the boundary terms. This concerns only the case  $|k| = 1$ . Thus, the only delicate case is when  $\omega$  is large, for which estimate (1.19) immediately yields the result. This concludes the proof of the resolvent estimate (2.27).  $\square$

### 3 Proof of Theorem 1.2

Theorem 1.2 relies on several strategies depending on the value of the resolvent parameter  $\omega \in \mathbb{R}$  and the frequency parameter  $k \in \mathbb{Z}$ .

To study the equation (1.12), we first remove the metric from the Laplacian, which can be done by the change of variable  $x = \log(r)$ , so that  $z_k(x) = y_k(r)$  and  $F_{1,k}(x) = r^2 f_{1,k}(r)$ ,  $F_{2,k}(x) = r^2 f_{2,k}(r)$ , for which (1.12) can be rewritten as follows:

$$\begin{cases} \partial_x^2 z_k - (k^2 - \omega^2 e^{2x}) z_k = i\omega F_{1,k} + F_{2,k}, & \text{in } (a_0, a_1). \\ z_k(a_1) = 0, \\ \partial_x z_k(a_0) = \rho(\omega, k) z_k(a_0) + g_k, \end{cases} \quad (3.1)$$

where we have set  $a_0 = \log(R_0)$ ,  $a_1 = \log(R_1)$  and where

$$F_{1,k} \in H^1(a_0, a_1), \text{ with } F_{1,k}(a_1) = 0, \quad \text{and} \quad F_{2,k} \in L^2(a_0, a_1). \quad (3.2)$$

The resolvent estimate we aim at proving, corresponding to (1.18), then reads as follows:

$$\begin{aligned} & \|\partial_x z_k\|_{L^2(a_0, a_1)} + |k| \|z_k\|_{L^2(a_0, a_1)} + |\omega| \|z_k e^x\|_{L^2(a_0, a_1)} \\ & \leq C \left( \|\partial_x F_{1,k}\|_{L^2(a_0, a_1)} + |k| \|F_{1,k}\|_{L^2(a_0, a_1)} + \|F_{2,k} e^x\|_{L^2(a_0, a_1)} + |g_k| \mathbf{1}_{\omega^2 R_0^2 \leq k^2} + |\partial_x z_k(a_1)| \right), \end{aligned} \quad (3.3)$$

and, corresponding to (1.19),

$$\begin{aligned} & \mathbf{1}_{|\omega|^2 R_0^2 + |k|^2 \leq M^2} \left( |\partial_x z_k(a_0)|^2 + |z_k(a_0)|^2 \right) \\ & + \mathbf{1}_{\substack{|\omega|^2 R_0^2 + |k|^2 \geq M^2 \\ \omega^2 R_0^2 \geq |k|^2}} \left( |\partial_x z_k(a_0)|^2 + \max \left\{ \sqrt{k^2 + \omega^2 R_0^2}, \omega^2 R_0^2 - k^2 \right\} |z_k(a_0)|^2 \right) \\ & + \mathbf{1}_{\substack{|\omega|^2 R_0^2 + |k|^2 \geq M^2 \\ \omega^2 R_0^2 \leq |k|^2}} \left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{\log\left(\frac{|k|}{|\omega|R_0}\right) \geq \frac{A}{|k|^{2/3}}} |z_k(a_0)|^2 \right. \\ & \quad \left. + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{\log\left(\frac{|k|}{|\omega|R_0}\right) \leq \frac{A}{|k|^{2/3}}} |z_k(a_0)|^2 \right) \\ & \leq C \left( \|\partial_x F_{1,k}\|_{L^2(a_0, a_1)}^2 + |k|^2 \|F_{1,k}\|_{L^2(a_0, a_1)}^2 + \|F_{2,k}\|_{L^2(a_0, a_1)}^2 + |g_k|^2 \mathbf{1}_{\omega^2 R_0^2 \leq k^2} + |\partial_x z_k(a_1)|^2 \right). \end{aligned} \quad (3.4)$$

We will now focus on the proof of the resolvent estimate (3.3) for solutions of (3.1). In order to do that, we will distinguish several cases:

- The case of bounded frequency parameters  $\omega^2 R_0^2 + k^2 \leq M^2$ , see Section 3.1 and Lemma 3.2, which can be handled using classical Carleman estimates.
- The case  $|\omega| \gg |k|$ , see Section 3.2, which we deal with using multiplier techniques.
- The case  $|\omega| \ll |k|$ , see Section 3.3, in which we use a factorization technique.

Actually, we will do a slightly more subtle analysis, that allows to consider  $|\omega|$  and  $|k|$  of the same order in Section 3.2 and in Section 3.3, and to conclude the resolvent estimate (3.3) for any  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ . To be more precise, for  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ , we introduce the function

$$H_{\omega,k}(x) := k^2 - \omega^2 e^{2x}, \quad x \in \mathbb{R}. \quad (3.5)$$

and we denote by  $x_0^*(\omega, k)$  its zero:

$$x_0^*(\omega, k) = \log \left( \frac{|k|}{|\omega|} \right). \quad (3.6)$$

Since  $H_{\omega,k}$  is a decreasing function of  $x$  and  $H_{\omega,k}(x_0^*(\omega, k)) = 0$ , it is positive for  $x < x_0^*(\omega, k)$  and negative for  $x > x_0^*(\omega, k)$ . We shall thus do as follows:

- When  $x \in [\max\{a_0, x_0^*(\omega, k)\}, a_1]$ , corresponding to  $|\omega|e^x \geq |k|$ , we use a multiplier method, see Section 3.2 and Corollary 3.6.
- When  $x \in [a_0, \min\{x_0^*(\omega, k), a_1\}]$ , corresponding to  $|\omega|e^x \leq |k|$ , we use a factorization argument to get an estimate on  $z_k(a_0)$ , see Section 3.3 and Lemma 3.11.
- And actually, when  $x_0^*(\omega, k) > a_0$ , we also apply the multiplier method in the whole interval  $(a_0, a_1)$  to conclude the resolvent estimate (3.3) more directly, see Theorem 3.8.

**Remark 3.1.** *Neglecting the boundary, the symbol of the operator appearing in the right hand-side of equation (3.1) is of the form  $p(x, \omega, k, \xi) = -\xi^2 - k^2 + \omega^2 e^{2x}$ , where  $x \in \mathbb{R}$ ,  $\omega \in \mathbb{R}$ ,  $k \in \mathbb{Z}$  and  $\xi \in \mathbb{R}$ , where  $\xi$  is the dual variable of  $x \in \mathbb{R}$ , and the above mentioned cases can be clearly understood at this level:*

- When  $x \geq x_0^*(\omega, k)$ , that is  $|\omega|e^x \geq |k|$ , the polynomial  $\xi \mapsto p(x, \omega, k, \xi)$  has two distinct real roots. The equation is thus hyperbolic there.
- When  $x \leq x_0^*(\omega, k)$ , that is  $|\omega|e^x \leq |k|$ , the polynomial  $\xi \mapsto p(x, \omega, k, \xi)$  has two distinct imaginary roots. The equation is thus elliptic there.

*It then appears clearly that our model is in fact closely related to the Friedlander model (see [12]) analyzed in details with homogeneous Dirichlet boundary conditions, and indeed shares some of its features.*

### 3.1 Bounded frequency parameters

The first estimate we give here is rather classical and focuses on the case where  $|\omega| + |k|$  is bounded. It is based on the following result:

**Lemma 3.2.** *Let  $M > 0$ . There exists  $C$  such that for all  $q \in L^\infty(a_0, a_1)$  with  $\|q\|_{L^\infty} \leq M$  and for all  $z$  in  $H^2(a_0, a_1)$  satisfying  $z(a_1) = 0$ ,*

$$\|\partial_x z\|_{L^2(a_0, a_1)} + \|z\|_{L^2(a_0, a_1)} + |\partial_x z(a_0)| + |z(a_0)| \leq C \left\| -\partial_x^2 z + qz \right\|_{L^2(a_0, a_1)} + C |\partial_x z(a_1)|. \quad (3.7)$$

*In particular, for all  $M > 0$ , estimates (3.3)–(3.4), for  $z \in H^2(a_0, a_1)$  solution of (3.1), holds uniformly in the ball*

$$\{(\omega, k) \in \mathbb{R} \times \mathbb{Z}, |k|^2 + |\omega|^2 R_0^2 \leq M^2\}.$$

*Proof.* We first deal with the case  $q = 0$ . For any  $s > 0$  and  $z \in H^2(a_0, a_1)$  with  $z(a_1) = 0$ , we set  $w = e^{sx} z$ , and we consider the conjugate operator

$$P_s w := e^{sx} \partial_x^2 (e^{-sx} w) = (\partial_x - s)^2 w, \quad (3.8)$$

so that  $P_s w = e^{sx} \partial_x^2 z$ .

Setting

$$v := (\partial_x - s)w, \quad (3.9)$$

we have

$$P_s w := (\partial_x - s)v. \quad (3.10)$$

Noting that by (3.9) one has

$$|v|^2 = |\partial_x w|^2 + s^2 |w|^2 - s \partial_x (|w|^2),$$

and integrating over  $(a_0, a_1)$ , we get

$$s|w(a_0)|^2 + \int_{a_0}^{a_1} (|\partial_x w|^2 + s^2 |w|^2) dx = \int_{a_0}^{a_1} |v|^2 dx + s|w(a_1)|^2. \quad (3.11)$$

From  $|(\partial_x - s)v|^2 = |\partial_x v|^2 + s^2 |v|^2 - s \partial_x (|v|^2)$ , we get similarly

$$s|v(a_0)|^2 + s^2 \int_{a_0}^{a_1} |v|^2 dx \leq \int_{a_0}^{a_1} |P_s w|^2 dx + s|v(a_1)|^2. \quad (3.12)$$

It stems from these estimates that

$$s^3 |w(a_0)|^2 + s|v(a_0)|^2 + s^4 \|w\|_{L^2(a_0, a_1)}^2 + s^2 \|\partial_x w\|_{L^2(a_0, a_1)}^2 \leq \|P_s w\|_{L^2(a_0, a_1)}^2 + s|v(a_1)|^2 + s^3 |w(a_1)|^2.$$

Since  $w = e^{sx} z$ , recalling  $z(a_1) = 0$ , we have  $w(a_1) = 0$  and  $v(a_1) = (\partial_x - s)(e^{sx} z)(a_1) = e^{sa_1} \partial_x z(a_1)$ . Similarly,  $w(a_0) = z(a_0) e^{sa_0}$ , and  $v(a_0) = (\partial_x z(a_0) - s z(a_0)) e^{sa_0}$ . Accordingly, recalling also that  $P_s w = e^{sx} \partial_x^2 z$ , there exists a constant  $C > 0$  such that for all  $s \geq 1$ ,

$$\begin{aligned} e^{2sa_0} (s^3 |z(a_0)|^2 + s |\partial_x z(a_0)|^2) + s^4 \|e^{sx} z\|_{L^2(a_0, a_1)}^2 + s^2 \|e^{sx} \partial_x z\|_{L^2(a_0, a_1)}^2 \\ \leq C^2 \left( \|e^{sx} \partial_x^2 z\|_{L^2(a_0, a_1)}^2 + s e^{2sa_1} |\partial_x z(a_1)|^2 \right). \end{aligned} \quad (3.13)$$

Now, since we assume  $\|q\|_{L^\infty} \leq M$ , taking  $s \geq 2\sqrt{CM}$  in (3.13) we get

$$\begin{aligned} e^{2sa_0} (s^3 |z(a_0)|^2 + s |\partial_x z(a_0)|^2) + s^4 \|e^{sx} z\|_{L^2(a_0, a_1)}^2 + s^2 \|e^{sx} \partial_x z\|_{L^2(a_0, a_1)}^2 \\ \leq 4C^2 \left( \|e^{sx} (-\partial_x^2 + q) z\|_{L^2(a_0, a_1)}^2 + s e^{2sa_1} |\partial_x z(a_1)|^2 \right). \end{aligned} \quad (3.14)$$

Finally, taking lower and upper bounds for the weights  $e^{sx}$  and fixing  $s$ , we easily deduce Lemma 3.2.  $\square$

### 3.2 High-frequency estimate $|\omega| \gg |k|$ : the multiplier method

As we said previously, here,  $|\omega| \gg |k|$  means that  $x_0^*(\omega, k)$  defined in (3.6) is such that  $x_0^*(\omega, k) < a_1$ . As we shall see, our arguments below will be used mainly on the interval  $[\max\{a_0, x_0^*(\omega, k)\}, a_1]$ .

**Lemma 3.3.** *For any  $x_0 \in [a_0, a_1]$ , for any  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ , any solution  $z_k$  of equation (3.1) satisfies*

$$\begin{aligned} |\partial_x z_k(x_0)|^2 + \Re(\bar{z}_k(x_0) \partial_x z_k(x_0)) + \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \\ = |\partial_x z_k(a_1)|^2 - \Re \left( \int_{x_0}^{a_1} (2\partial_x \bar{z}_k + \bar{z}_k) (i\omega F_{1,k} + F_{2,k}) dx \right) + H_{\omega, k}(x_0) |z_k(x_0)|^2. \end{aligned} \quad (3.15)$$

**Remark 3.4.** *Note that, when considering Dirichlet boundary conditions  $z_k(a_0) = 0$ , Lemma 3.3 applied to  $x_0 = a_0$  proves the resolvent estimate at once, provided we suitably estimate the terms involving  $F_{1,k}$  and  $F_{2,k}$ , which can of course be done (see for instance the proof of estimate (3.25) afterwards, which can be easily adapted to the case  $z_k(a_0) = 0$ ). In fact, this is the usual multiplier method.*

*Proof.* Multiplying equation (3.1) by  $(2\partial_x \bar{z}_k + \bar{z}_k)$  and integrating on  $(x_0, a_1)$  easily leads to

$$\begin{aligned} |\partial_x z_k(x_0)|^2 - (k^2 - \omega^2 e^{2x}) |z_k(x_0)|^2 + \Re(\bar{z}_k(x_0) \partial_x z_k(x_0)) + \int_{x_0}^{a_1} (|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2) dx \\ = |\partial_x z_k(a_1)|^2 - \Re\left(\int_{x_0}^{a_1} (2\partial_x \bar{z}_k + \bar{z}_k)(i\omega F_{1,k} + F_{2,k}) dx\right) \end{aligned}$$

since  $z_k(a_1) = 0$ . Of course, this coincides with the multiplier identity (3.15).  $\square$

**Remark 3.5.** Another proof of Lemma 3.3 can be obtained by considering the quantity

$$E_k(x) = |\partial_x z_k(x)|^2 - (k^2 - \omega^2 e^{2x}) |z_k(x)|^2 + \Re(\bar{z}_k(x) \partial_x z_k(x)), \quad (x \in [a_0, a_1]),$$

and compute

$$\frac{d}{dx} E_k(x) + |\partial_x z_k(x)|^2 + (k^2 + \omega^2 e^{2x}) |z_k(x)|^2 = \Re((2\partial_x \bar{z}_k(x) + \bar{z}_k(x))(i\omega F_{1,k}(x) + F_{2,k}(x))).$$

Of course, this approach yields the same identity as in (3.15), but the above computation can be interpreted as a version of the so-called lateral propagation of the energy in 1-space dimension for the resolvent equation corresponding to the wave equation.

An immediate corollary of Lemma 3.3 is the following one:

**Corollary 3.6.** Let  $M \geq 2$ . There exists a constant  $C > 0$  such that for any  $z_k$  satisfying (3.1) for  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2 \geq M^2$  with  $x_0^*(\omega, k) < a_1$  and  $(F_{1,k}, F_{2,k})$  as in (3.2), for all  $x_0 \in [\max\{a_0, x_0^*(\omega, k)\}, a_1]$ ,

$$\begin{aligned} |\partial_x z_k(x_0)|^2 + \max\left\{\sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2\right\} |z_k(x_0)|^2 \\ + \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \\ \leq C |\partial_x z_k(a_1)|^2 + C \int_{x_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \end{aligned} \quad (3.16)$$

In particular, if  $\omega^2 R_0^2 + k^2 \geq M^2$  and  $x_0^*(\omega, k) \leq a_0$  (i.e.  $\omega^2 R_0^2 \geq k^2$ ), taking  $x_0 = a_0$  in the above estimate yields the resolvent estimate

$$\begin{aligned} |\partial_x z_k(a_0)|^2 + \max\left\{\sqrt{k^2 + \omega^2 R_0^2}, \omega^2 R_0^2 - k^2\right\} |z_k(a_0)|^2 + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \\ \leq C |\partial_x z_k(a_1)|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx, \end{aligned} \quad (3.17)$$

that is estimates (3.3)–(3.4).

*Proof.* We start from (3.15), and we notice that for  $x_0 \geq x_0^*(\omega, k)$ ,  $H_{\omega,k}(x_0) = k^2 - \omega^2 e^{2x_0} \leq 0$ . Thus, we deduce

$$\begin{aligned} |\partial_x z_k(x_0)|^2 + \Re(\bar{z}_k(x_0) \partial_x z_k(x_0)) + (\omega^2 e^{2x_0} - k^2) |z_k(x_0)|^2 + \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \\ = |\partial_x z_k(a_1)|^2 + 2\Re\left(\int_{x_0}^{a_1} \left(\partial_x \bar{z}_k + \frac{\bar{z}_k}{2}\right) (i\omega F_{1,k} + F_{2,k}) dx\right). \end{aligned} \quad (3.18)$$

Now, using  $z_k(a_1) = 0$ , we get

$$\begin{aligned} \sqrt{k^2 + \omega^2 e^{2x_0}} |z_k(x_0)|^2 = -2\sqrt{k^2 + \omega^2 e^{2x_0}} \Re\left(\int_{x_0}^{a_1} (\partial_x z_k) \bar{z}_k dx\right) \\ \leq 2 \|\partial_x z_k\|_{L^2(x_0, a_1)} \left\| \sqrt{k^2 + \omega^2 e^{2x_0}} z_k \right\|_{L^2(a_0, a_1)} \leq \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx, \end{aligned} \quad (3.19)$$

which immediately implies that

$$\begin{aligned}
& |\partial_x z_k(x_0)|^2 + \Re(\bar{z}_k(x_0)\partial_x z_k(x_0)) + ((\omega^2 e^{2x_0} - k^2) + \frac{1}{2}\sqrt{k^2 + \omega^2 e^{2x_0}})|z_k(x_0)|^2 \\
& \quad + \frac{1}{2} \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x})|z_k|^2] dx \\
& \leq |\partial_x z_k(a_1)|^2 + 2 \left| \int_{x_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) dx \right|. \quad (3.20)
\end{aligned}$$

Then, from  $\omega^2 R_0^2 + k^2 \geq 4$ , i.e.  $\sqrt{k^2 + \omega^2 e^{2x_0}} \geq 2$ , we obtain

$$|\bar{z}_k(x_0)\partial_x z_k(x_0)| \leq \frac{1}{2}|\partial_x z_k(x_0)|^2 + \frac{1}{2}|z_k(x_0)|^2 \leq \frac{1}{2}|\partial_x z_k(x_0)|^2 + \frac{1}{4}\sqrt{k^2 + \omega^2 e^{2x_0}}|z_k(x_0)|^2,$$

and

$$\begin{aligned}
& \frac{1}{2}|\partial_x z_k(x_0)|^2 + ((\omega^2 e^{2x_0} - k^2) + \frac{1}{4}\sqrt{k^2 + \omega^2 e^{2x_0}})|z_k(x_0)|^2 \\
& \quad + \frac{1}{2} \int_{x_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x})|z_k|^2] dx \\
& \leq |\partial_x z_k(a_1)|^2 + 2 \left| \int_{x_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) dx \right|. \quad (3.21)
\end{aligned}$$

We now write that

$$\begin{aligned}
& \int_{x_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) dx = \bar{z}_k(a_1)i\omega F_{1,k}(a_1) - \bar{z}_k(x_0)i\omega F_{1,k}(x_0) \\
& \quad + \int_{x_0}^{a_1} \left( -i\omega \bar{z}_k \partial_x F_{1,k} + \partial_x \bar{z}_k F_{2,k} + \frac{i\omega \bar{z}_k}{2} F_{1,k} + \frac{\bar{z}_k}{2} F_{2,k} \right) dx. \quad (3.22)
\end{aligned}$$

Using  $F_{1,k}(a_1) = 0$ , and the fact that, similarly as in (3.19),

$$|k||F_{1,k}(x_0)|^2 \leq \int_{x_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2|F_{1,k}|^2) dx, \quad (3.23)$$

and that, for  $x_0 \geq x_0^*(\omega, k)$ ,  $\omega^2 e^{2x_0} - k^2 \geq 0$ , the boundary terms in (3.22) can be estimated as follows:

$$\begin{aligned}
& |\bar{z}_k(a_1)i\omega F_{1,k}(a_1) - \bar{z}_k(x_0)i\omega F_{1,k}(x_0)| \leq |\omega||z_k(x_0)||F_{1,k}(x_0)| \\
& \leq \frac{1}{16} \max \left\{ \sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2 \right\} |z_k(x_0)|^2 + C \frac{\omega^2}{\max \left\{ \sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2 \right\}} |F_{1,k}(x_0)|^2 \\
& \leq \frac{1}{16} \max \left\{ \sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2 \right\} |z_k(x_0)|^2 + C|k||F_{1,k}(x_0)|^2 \\
& \leq \frac{1}{16} \max \left\{ \sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2 \right\} |z_k(x_0)|^2 + C \int_{x_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2|F_{1,k}|^2) dx.
\end{aligned}$$

On the other hand, we easily get

$$\begin{aligned}
& \left| \int_{x_0}^{a_1} \left( -i\omega \bar{z}_k \partial_x F_{1,k} + \partial_x \bar{z}_k F_{2,k} + \frac{i\omega \bar{z}_k}{2} F_{1,k} + \frac{\bar{z}_k}{2} F_{2,k} \right) dx \right| \\
& \leq \|\omega z_k\|_{L^2(x_0, a_1)} \|\partial_x F_{1,k}\|_{L^2(x_0, a_1)} + \|\omega z_k\|_{L^2(x_0, a_1)} \|F_{1,k}\|_{L^2(x_0, a_1)} \\
& \quad + (\|\partial_x z_k\|_{L^2(x_0, a_1)} + \|\omega z_k\|_{L^2(x_0, a_1)}) \|F_{2,k}\|_{L^2(x_0, a_1)} \\
& \leq \frac{1}{8} \left( \|\partial_x z_k\|_{L^2(x_0, a_1)}^2 + \|\omega e^x z_k\|_{L^2(x_0, a_1)}^2 \right) \\
& \quad + C \int_{x_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2|F_{1,k}|^2 + |F_{2,k}|^2) dx. \quad (3.24)
\end{aligned}$$

Combining the last two estimates with (3.22), we obtain

$$\begin{aligned} & \left| \int_{x_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) \, dx \right| \\ & \leq \frac{1}{16} \max \left\{ \sqrt{k^2 + \omega^2 e^{2x_0}}, \omega^2 e^{2x_0} - k^2 \right\} |z_k(x_0)|^2 + \frac{1}{8} \left( \|\partial_x z_k\|_{L^2(x_0, a_1)}^2 + \|\omega e^x z_k\|_{L^2(x_0, a_1)}^2 \right) \\ & \quad + C \int_{x_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) \, dx. \end{aligned} \quad (3.25)$$

for some  $C$  independent of  $(\omega, k)$ . Plugging this last estimate in (3.21), we obtain (3.16).

The fact that the resolvent estimate (3.16) yields the resolvent estimate (3.3) when  $(\omega, k)$  satisfy  $\omega^2 R_0^2 + k^2 \geq 4$  and  $x_0^*(\omega, k) \leq a_0$  easily follows.  $\square$

Another interesting straightforward corollary of the identity (3.15) is the following one:

**Corollary 3.7.** *Let  $M \geq 4$ . For any  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $\omega^2 R_0^2 + k^2 \geq M^2$  and  $\omega^2 R_0^2 \leq k^2$ , any solution  $z_k$  of equation (3.1) satisfies, for some  $C$  independent of  $k$ ,*

$$\begin{aligned} & \sqrt{k^2 + \omega^2 e^{2a_0}} |z_k(a_0)|^2 + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] \, dx \leq C |\partial_x z(a_1)|^2 \\ & + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) \, dx + C (H_{\omega, k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k)) |z_k(a_0)|^2. \end{aligned} \quad (3.26)$$

Indeed, this corollary is interesting since it implies that if we know how to estimate  $(H_{\omega, k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k)) |z_k(a_0)|^2$  by the squares of the right hand side of (3.3), we immediately derive the resolvent estimate (3.3). This is in fact the strategy we follow hereafter in Section 3.3.

*Proof.* We start from the fact that, coming from (3.20) applied to  $x_0 = a_0$  and using the fact that  $\partial_x z_k(a_0) = \rho(\omega, k) z_k(a_0)$ , that for any  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$ , any solution  $z_k$  of equation (3.1) satisfies

$$\begin{aligned} & \left( \rho(\omega, k)^2 + \rho(\omega, k) + (\omega^2 e^{2a_0} - k^2) + \frac{1}{2} \sqrt{k^2 + \omega^2 e^{2a_0}} \right) |z_k(a_0)|^2 \\ & + \frac{1}{2} \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] \, dx \\ & \leq |\partial_x z_k(a_1)|^2 + 2 \left| \int_{a_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) \, dx \right|. \end{aligned} \quad (3.27)$$

To bound the last term, we use again identity (3.22). Recalling  $z_k(a_1) = 0$ , the pointwise estimate (3.23) applied to  $x_0 = a_0$ , and using  $\omega^2 R_0^2 \leq k^2$ , we get, for all  $\alpha > 0$ , (with  $C_\alpha = 1/(4\alpha)$ ):

$$\begin{aligned} & |\bar{z}_k(a_1) i\omega F_{1,k}(a_1) - \bar{z}_k(a_0) i\omega F_{1,k}(a_0)| \leq |\omega| |z_k(a_0)| |F_{1,k}(a_0)| \\ & \leq \alpha \frac{|\omega|^2}{|k|} |z_k(a_0)|^2 + C_\alpha |k| |F_{1,k}(a_0)|^2 \\ & \leq \alpha \frac{|k|}{R_0} |z_k(a_0)|^2 + C_\alpha \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2) \, dx. \end{aligned} \quad (3.28)$$

Similarly as in (3.24), we also get, for all  $\alpha > 0$ , the existence of a constant  $C_\alpha > 0$  such that

$$\begin{aligned} & \left| \int_{a_0}^{a_1} \left( -i\omega \bar{z}_k \partial_x F_{1,k} + \partial_x \bar{z}_k F_{2,k} + \frac{i\omega \bar{z}_k}{2} F_{1,k} + \frac{\bar{z}_k}{2} F_{2,k} \right) \, dx \right| \\ & \leq \alpha \left( \|\partial_x z_k\|_{L^2(a_0, a_1)}^2 + \|\omega e^x z_k\|_{L^2(a_0, a_1)}^2 \right) + C_\alpha \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) \, dx. \end{aligned} \quad (3.29)$$

Choosing  $\alpha > 0$  small enough in the above estimates, we get that there exists  $C > 0$  such that

$$2 \left| \int_{a_0}^{a_1} \left( \partial_x \bar{z}_k + \frac{\bar{z}_k}{2} \right) (i\omega F_{1,k} + F_{2,k}) dx \right| \leq \frac{1}{4} \sqrt{k^2 + \omega^2 e^{2a_0}} |z_k(a_0)|^2 \\ + \frac{1}{4} \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx.$$

Combined with (3.27), we easily deduce (3.26).  $\square$

### 3.3 High-frequency estimate $|\omega| \lll |k|$ : a factorization argument

When  $x_0^*(\omega, k) > a_0$ , i.e. when  $k^2 > \omega^2 R_0^2$ , the multiplier identity (3.15) does not allow to conclude immediately since then  $H_{\omega,k}(a_0) > 0$ .

In this section, we shall always assume that  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfy

$$k^2 \geq \omega^2 R_0^2, \quad (\text{equivalently, } x_0^*(\omega, k) \geq a_0), \quad \text{and} \quad k^2 + \omega^2 R_0^2 \geq M^2, \quad (3.30)$$

where  $M$  is the constant appearing in Theorem 1.2.

The main result of this section is the following:

**Theorem 3.8.** *Assume that  $\rho = \rho(\omega, k)$  is as in (1.13) and satisfies the condition (1.15). Then there exist  $C > 0$  and  $A > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying (3.30) and any solution  $z_k$  of (3.1) with  $F_{1,k} \in H^1(a_0, a_1)$  satisfying  $F_{1,k}(a_1) = 0$ ,  $F_{2,k} \in L^2(a_0, a_1)$  and  $g_k \in \mathbb{C}$ ,*

$$\left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A|k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A|k|^{-2/3}} \right) |z_k(a_0)|^2 \\ + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \leq C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx \\ + C|g_k|^2 + C|\partial_x z_k(a_1)|^2. \quad (3.31)$$

*Proof.* Below, we design a factorization technique to get the resolvent estimate (3.31), which consists in writing the differential operator in (3.1) under the form

$$\partial_x^2 - H_{\omega,k}(x) = (\partial_x - X_{\omega,k})(\partial_x + X_{\omega,k}), \quad (3.32)$$

where  $X_{\omega,k}$  will be assumed to be a non-negative function of  $x$  such that

$$\begin{cases} \frac{d}{dx} X_{\omega,k}(x) = X_{\omega,k}^2(x) - H_{\omega,k}(x), & x \in [a_0, x^*], \\ X_{\omega,k}(x^*) = X^*, \end{cases} \quad (3.33)$$

where  $x^* > a_0$  and  $X^*$  are real numbers to be chosen later.

Indeed, this will allow us to get an estimate on  $|z_k(a_0)|$  according to the following lemma:

**Lemma 3.9.** *Let us assume that there exists  $x^* \in (a_0, a_1]$  and  $X^* \geq 0$  such that the ODE (3.33) has a solution  $X_{\omega,k}$  on the interval  $[a_0, x^*]$  satisfying  $X_{\omega,k}(x) \in [0, |k|]$  for all  $x \in [a_0, x^*]$ .*

*Then any solution  $z_k$  of equation (3.1) satisfies the following estimate: for all  $\alpha > 0$ , there exists  $C$  independent of  $(\omega, k)$  such that*

$$(\rho(\omega, k) + X_{\omega,k}(a_0))^2 |z_k(a_0)|^2 \leq \alpha \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx \\ + C|\partial_x z_k(x^*)| + C|X^*|^2 |z_k(x^*)|^2 + C|g_k|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \quad (3.34)$$

*Proof.* Setting  $w_k = \partial_x z_k + X_{\omega,k} z_k$  on the set  $(a_0, x^*)$ , where  $z_k$  solves (3.1), we get that

$$\begin{cases} (\partial_x - X_{\omega,k})w_k = i\omega F_{1,k} + F_{2,k}, & \text{in } (a_0, x^*) \\ w_k(x^*) = \partial_x z_k(x^*) + X_{\omega,k}(x^*)z_k(x^*) = \partial_x z_k(x^*) + X^* z_k(x^*). \end{cases} \quad (3.35)$$

Multiplying the equation by  $\bar{w}_k$  and integrating between  $a_0$  and  $x^*$ , we get

$$\frac{1}{2} (|w_k(x^*)|^2 - |w_k(a_0)|^2) - \int_{a_0}^{x^*} X_{\omega,k} |w_k|^2 dx = \Re \left( \int_{a_0}^{x^*} (i\omega F_{1,k} + F_{2,k}) \bar{w}_k dx \right).$$

Recalling that  $w_k = \partial_x z_k + X_{\omega,k} z_k$ ,  $X_{\omega,k}$  is assumed to be non-negative and that  $w_k(a_0) = \partial_x z_k(a_0) + X_{\omega,k}(a_0) z_k(a_0) = (\rho(\omega, k) + X_{\omega,k}(a_0)) z_k(a_0) + g_k$ , we easily deduce:

$$\begin{aligned} (\rho(\omega, k) + X_{\omega,k}(a_0))^2 |z_k(a_0)|^2 &\leq C \left| \int_{a_0}^{x^*} (i\omega F_{1,k} + F_{2,k}) (\partial_x \bar{z}_k + X_{\omega,k} \bar{z}_k) dx \right| \\ &\quad + C |\partial_x z_k(x^*)|^2 + C |X^*|^2 |z_k(x^*)|^2 + C |g_k|^2. \end{aligned} \quad (3.36)$$

Then, similarly as in the proofs of Corollary 3.6 and Corollary 3.7, we estimate

$$\begin{aligned} \int_{a_0}^{x^*} (i\omega F_{1,k} + F_{2,k}) (\partial_x \bar{z}_k + X_{\omega,k} \bar{z}_k) dx &= i\omega F_{1,k}(x^*) \bar{z}_k(x^*) - i\omega F_{1,k}(a_0) \bar{z}_k(a_0) \\ &\quad + \int_{a_0}^{x^*} (-i\omega \partial_x F_{1,k} \bar{z}_k + F_{2,k} \partial_x \bar{z}_k + X_{\omega,k} F_{1,k} i\omega \bar{z}_k + F_{2,k} X_{\omega,k} \bar{z}_k) dx. \end{aligned} \quad (3.37)$$

Since  $|\omega| \leq |k|/R_0$ , we have from (3.23) that, for all  $\alpha > 0$ , there exists  $C_\alpha > 0$  such that

$$\begin{aligned} |i\omega F_{1,k}(a_0) \bar{z}_k(a_0)| + |i\omega F_{1,k}(x^*) \bar{z}_k(x^*)| &\leq C |k| (|F_{1,k}(a_0)| |z_k(a_0)| + |F_{1,k}(x^*)| |z_k(x^*)|) \\ &\leq C_\alpha \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2) dx + \frac{\alpha}{4} |k| (|z_k(a_0)|^2 + |z_k(x^*)|^2) \\ &\leq C_\alpha \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2) dx + \frac{\alpha}{2} \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx. \end{aligned} \quad (3.38)$$

Using then that  $|X_{\omega,k}| \leq k$ , and that  $|\omega| \leq |k|/R_0$ ,

$$\begin{aligned} &\left| \int_{a_0}^{x^*} (-i\omega \partial_x F_{1,k} \bar{z}_k + F_{2,k} \partial_x \bar{z}_k + X_{\omega,k} F_{1,k} i\omega \bar{z}_k + F_{2,k} X_{\omega,k} \bar{z}_k) dx \right| \\ &\leq C \|\partial_x F_{1,k}\|_{L^2(a_0, a_1)} \| |k| z_k \|_{L^2(a_0, a_1)} + \|F_{2,k}\|_{L^2(a_0, a_1)} \|\partial_x z_k\|_{L^2(a_0, a_1)} \\ &\quad + C \| |k| F_{1,k} \|_{L^2(a_0, a_1)} \| |k| z_k \|_{L^2(a_0, a_1)} + C \|F_{2,k}\|_{L^2(a_0, a_1)} \| |k| z_k \|_{L^2(a_0, a_1)}. \end{aligned}$$

Thus, for all  $\alpha > 0$ , there exists  $C_\alpha > 0$  such that

$$\begin{aligned} &\left| \int_{a_0}^{x^*} (-i\omega \partial_x F_{1,k} \bar{z}_k + F_{2,k} \partial_x \bar{z}_k + X_{\omega,k} F_{1,k} i\omega \bar{z}_k + F_{2,k} X_{\omega,k} \bar{z}_k) dx \right| \\ &\leq C_\alpha \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx + \frac{\alpha}{2} \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx. \end{aligned} \quad (3.39)$$

Plugging this last estimate and (3.38) in the identity (3.37), we easily deduce (3.34) from the estimate (3.36).  $\square$

Lemma 3.9 indicates that, provided we get non-negative solutions  $X_{\omega,k}$  of (3.33) on an interval of the form  $[a_0, x^*]$ , one can get an estimate on  $|\rho + X_{\omega,k}(a_0)| |z_k(a_0)|$  from an estimate on  $|\partial_x z_k(x^*)| + |X^*| |z_{\omega,k}(x^*)|$ , and other terms involving the source terms  $F_{1,k}$ ,  $F_{2,k}$  and  $g_k$  of (3.1).

Our goal now is to combine estimate (3.34) with Corollary 3.6. To do so, we aim to choose  $x^*$  in  $[x_0^*(\omega, k), a_1]$  and  $|X^*|^2 = \theta^2 (\omega^2 e^{2x^*} - k^2)$  for some positive constant  $\theta$  independent of  $\omega$  and  $k$ . We see that  $X^*$  becomes larger as  $x^*$  is chosen further from  $x_0^*(\omega, k)$ , and as a consequence the estimate becomes better. However, we can prove the existence of  $X_{\omega,k}$  solving (3.33) on  $[a_0, x^*]$  only for  $x^*$  close enough to  $x_0^*(\omega, k)$ . Hence a compromise has to be made.

More precisely, for  $x^* = x_0^*(\omega, k) + c/|k|^\alpha$  with  $c \geq 0$  and  $\alpha > 0$ , we have  $X^* = \theta \sqrt{2c} |k|^{1-\alpha/2}$  for some positive constant  $\theta$ . In the following, we make two specific choices, the first one being  $x^* = x_0^*(\omega, k)$  and  $X^* = 0$  (that is  $c = 0$ ), and the second one  $x^* = x_0^*(\omega, k) + 1/(2|k|^{2/3})$  and  $X^* = 2|k|^{2/3}$  (that is  $c = 1/2$ ,  $\theta = 2$ , and  $\alpha = 2/3$ ). The solutions corresponding to these choices enjoy the following properties:



**Lemma 3.10.** *Let  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  satisfying (3.30).*

*The solution  $X_{0,\omega,k}$  of*

$$\begin{cases} \frac{d}{dx} X_{0,\omega,k}(x) = X_{0,\omega,k}^2(x) - H_{\omega,k}(x), & x \in [a_0, x_0^*(\omega, k)], \\ X_{0,\omega,k}(x_0^*) = 0, \end{cases} \quad (3.40)$$

*is well-defined on  $[a_0, x_0^*(\omega, k)]$  and non-negative there.*

*Set*

$$x_1^*(\omega, k) = x_0^*(\omega, k) + \frac{1}{2|k|^{2/3}}. \quad (3.41)$$

*Then the solution  $X_{1,\omega,k}$  of*

$$\begin{cases} \frac{d}{dx} X_{1,\omega,k}(x) = X_{1,\omega,k}^2(x) - H_{\omega,k}(x), & x \in [a_0, x_1^*(\omega, k)], \\ X_{1,\omega,k}(x_1^*(\omega, k)) = 2|k|^{2/3}, \end{cases} \quad (3.42)$$

*is well-defined on  $[a_0, x_1^*(\omega, k)]$  and non-negative there.*

*Besides, we have the following estimates: For all  $\varepsilon > 0$ , there exists  $A_\varepsilon > 0$  independent of  $(\omega, k)$  such that*

$$\forall x < x_0^*(\omega, k) - A_\varepsilon k^{-2/3}, \quad (1 - \varepsilon) \sqrt{H_{\omega,k}(x)} \leq X_{0,\omega,k}(x) \leq X_{1,\omega,k}(x) \leq \sqrt{H_{\omega,k}(x)} \leq |k|. \quad (3.43)$$

*For all  $A > 0$ , there exists  $\delta_A > 0$  independent of  $(\omega, k)$  such that*

$$\forall x \in [x_0^*(\omega, k) - A|k|^{-2/3}, x_0^*(\omega, k)], \quad X_{1,\omega,k}(x) - X_{0,\omega,k}(x) \geq \delta_A k^{2/3}. \quad (3.44)$$

Lemma 3.10 is one of the delicate points of our analysis, and its proof is postponed to Section 3.4. Using then Lemma 3.9, we can prove the resolvent estimates (3.3)–(3.4) for solutions  $z_k$  of (3.1) for  $(\omega, k)$  satisfying (3.30).

Indeed, using Lemma 3.9 with  $X_{0,\omega,k}$  on  $[a_0, \tilde{x}_0^*]$  with  $\tilde{x}_0^* = \min\{x_0^*(\omega, k), a_1\}$  and  $X_{1,\omega,k}$  on  $[a_0, \tilde{x}_1^*]$  with  $\tilde{x}_1^* = \min\{x_1^*(\omega, k), a_1\}$  and summing the two estimates, we obtain the following: for all  $\alpha > 0$ , there exists  $C > 0$  such that

$$\begin{aligned} & ((\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 + (\rho(\omega, k) + X_{1,\omega,k}(a_0))^2) |z_k(a_0)|^2 \\ & \leq \alpha \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx + C |\partial_x z_k(x_0^*)|^2 + C |\partial_x z_k(x_1^*)|^2 + C |k|^{4/3} |z_k(x_1^*)|^2 \\ & \quad + C |g_k|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \end{aligned} \quad (3.45)$$

We shall now bound from below the left hand-side of (3.45):

**Lemma 3.11.** *Let  $\varepsilon > 0$  and  $M \geq 1$  and assume the condition (1.15). Then there exist a constant  $\delta_0 > 0$  and  $A_\varepsilon > 0$  such that for all  $(\omega, k) \in \mathbb{R} \times \mathbb{Z}$  with  $k^2 + \omega^2 R_0^2 \geq M^2$  and  $k^2 \geq \omega^2 R_0^2$ , (equivalently,  $x_0^*(\omega, k) \geq a_0$ ),*

$$\begin{aligned} & ((\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 + (\rho(\omega, k) + X_{1,\omega,k}(a_0))^2) \\ & \geq \delta_0 (I_\varepsilon(\omega, k))^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} \\ & \quad + H_{\omega,k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k). \end{aligned} \quad (3.46)$$

*Proof.* First note that

$$\begin{aligned} & (\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 + (\rho(\omega, k) + X_{1,\omega,k}(a_0))^2 \\ & = 2 \left( \left( \rho(\omega, k) + \frac{X_{0,\omega,k}(a_0) + X_{1,\omega,k}(a_0)}{2} \right)^2 + \left( \frac{X_{1,\omega,k}(a_0) - X_{0,\omega,k}(a_0)}{2} \right)^2 \right). \end{aligned} \quad (3.47)$$

Now, let  $\varepsilon > 0$  and  $M \geq 1$  and assume condition (1.15). Using Lemma 3.9, there exists  $A_\varepsilon > \mathbf{r}$  ( $\mathbf{r}$  is the constant in (1.14)) such that (3.43) holds for all  $x < x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}$ , and there exists  $\delta_\varepsilon > 0$  such that (3.44) holds for  $x \in [x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}, x_0^*(\omega, k)]$ . Consequently, we shall discuss how  $a_0$  compares to  $x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}$ .

If  $a_0 < x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}$ , i.e. if  $\log(|k|/|\omega|R_0) > A_\varepsilon |k|^{-2/3}$ , using the condition (1.15) and the fact that

$$(1 - \varepsilon)\sqrt{H_{\omega,k}(a_0)} \leq X_{0,\omega,k}(a_0) \leq X_{1,\omega,k}(a_0) \leq \sqrt{H_{\omega,k}(a_0)},$$

hence that there exists  $c \in [1 - \varepsilon, 1]$  such that

$$\frac{X_{0,\omega,k}(a_0) + X_{1,\omega,k}(a_0)}{2} = c\sqrt{H_{\omega,k}(a_0)},$$

we easily deduce by bounding from below the first term of the right hand side of (3.47) that

$$(\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 + (\rho(\omega, k) + X_{1,\omega,k}(a_0))^2 \geq 2\delta(H_{\omega,k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k)).$$

If  $a_0 \in [x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}, x_0^*(\omega, k)]$ , i.e. if  $\log(|k|/|\omega|R_0) > A_\varepsilon |k|^{-2/3}$ , we simply use (3.44) and the second term in the right hand side of (3.47) to deduce that

$$(\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 + (\rho(\omega, k) + X_{1,\omega,k}(a_0))^2 \geq 2\delta_\varepsilon |k|^{4/3}.$$

On the other hand, since  $X_{0,\omega,k}(a_0) \in [0, \sqrt{H_{\omega,k}(a_0)}]$ , we easily have

$$(\rho(\omega, k) + X_{0,\omega,k}(a_0))^2 \geq I_1(\omega, k)^2.$$

To end the proof of Lemma 3.11, it suffices to notice that there exists a constant  $C > 0$  such that for all  $a_0 \in [x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}, x_0^*(\omega, k)]$ ,

$$H_{\omega,k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k) \leq H_{\omega,k}(x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}) + 1 \leq C|k|^{4/3}.$$

This last estimate can be easily checked by looking at the asymptotic as  $|k| \rightarrow \infty$  of

$$H_{\omega,k}(x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}) = k^2 - \omega^2 e^{2x_0^*(\omega, k) - 2A_\varepsilon |k|^{-2/3}} = k^2 \left(1 - e^{-2A_\varepsilon |k|^{-2/3}}\right) \underset{|k| \rightarrow \infty}{\simeq} 2A_\varepsilon k^{4/3}.$$

We then easily conclude Lemma 3.11.  $\square$

We then show how Lemma 3.11 and estimate (3.45) imply the resolvent estimate (3.3) for solutions  $z_k$  of (3.1) when  $(\omega, k)$  are as in (3.30).

Using Lemma 3.11, we obtain that for all  $\alpha > 0$ , there exists  $C$  independent of  $(\omega, k)$  satisfying (3.30) such that

$$\begin{aligned} & \left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} \right) |z_k(a_0)|^2 \\ & + (H_{\omega,k}(a_0) - \rho(\omega, k)^2 - \rho(\omega, k)) |z_k(a_0)|^2 \leq \alpha \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx + C |\partial_x z_k(x_0^*)|^2 \\ & + C |\partial_x z_k(x_1^*)|^2 + C |k|^{4/3} |z_k(x_1^*)|^2 + C |g_k|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \end{aligned} \quad (3.48)$$

We then use Corollary 3.7 to deduce

$$\begin{aligned} & \left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} \right) |z_k(a_0)|^2 \\ & + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \leq \alpha \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx + C |\partial_x z_k(x_0^*)|^2 \\ & + C |\partial_x z_k(x_1^*)|^2 + C |k|^{4/3} |z_k(x_1^*)|^2 + C |g_k|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \end{aligned} \quad (3.49)$$

Now, using Corollary 3.6 at  $x_0 = \tilde{x}_0^*(\omega, k)$  and  $x_0 = \tilde{x}_1^*(\omega, k)$ , and the fact that

$$\omega^2 e^{2x_1^*(\omega, k)} - k^2 = k^2 \left( e^{2(x_1^*(\omega, k) - x_0^*(\omega, k))} - 1 \right) \underset{|k| \rightarrow \infty}{\simeq} k^{4/3},$$

we have

$$\begin{aligned} & \sqrt{k^2 + \omega^2 e^{2\tilde{x}_0^*(\omega, k)}} |\partial_x z_k(\tilde{x}_0^*(\omega, k))|^2 + |\partial_x z_k(\tilde{x}_1^*(\omega, k))|^2 + k^{4/3} |z_k(\tilde{x}_1^*(\omega, k))|^2 \\ & \leq C |\partial_x z_k(a_1)|^2 + C \int_{\tilde{x}_0^*}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \end{aligned}$$

Estimate (3.49) thus yields:

$$\begin{aligned} & \left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} \right) |z_k(a_0)|^2 \\ & + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \leq \alpha \int_{a_0}^{a_1} [|\partial_x z_k|^2 + k^2 |z_k|^2] dx \\ & + C |\partial_x z_k(a_1)|^2 + C |g_k|^2 + C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx. \quad (3.50) \end{aligned}$$

Taking  $\alpha > 0$  small enough (independent of  $(\omega, k)$ ), we conclude from (3.15) that

$$\begin{aligned} & \left( I_\varepsilon(\omega, k)^2 \mathbf{1}_{a_0 \leq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} + \max\{|k|^{4/3}, I_1(\omega, k)^2\} \mathbf{1}_{a_0 \geq x_0^*(\omega, k) - A_\varepsilon |k|^{-2/3}} \right) |z_k(a_0)|^2 \\ & + \int_{a_0}^{a_1} [|\partial_x z_k|^2 + (k^2 + \omega^2 e^{2x}) |z_k|^2] dx \leq C \int_{a_0}^{a_1} (|\partial_x F_{1,k}|^2 + k^2 |F_{1,k}|^2 + |F_{2,k}|^2) dx \\ & + C |g_k|^2 + C |\partial_x z_k(a_1)|^2. \end{aligned}$$

which concludes the proof of Theorem 3.8.  $\square$

**Remark 3.12.** *The starting point of the proof of Theorem 3.8 is the factorization of the differential operator  $\partial_x^2 - H_{\omega, k}(x)$  in the set where  $H_{\omega, k}(x) = k^2 - \omega^2 e^{2x}$  is positive.*

*Note that a natural factorization strategy would be to write*

$$\partial_x^2 - H_{\omega, k}(x) = \left( \partial_x - \sqrt{H_{\omega, k}(x)} \right) \left( \partial_x + \sqrt{H_{\omega, k}(x)} \right) + \text{remainder},$$

*but the remainder, which equals to  $-\partial_x \left( \sqrt{H_{\omega, k}(x)} \right)$ , is too singular for this approach close to the zero of  $H_{\omega, k}$ .*

*Our approach thus relies on the exact factorization of the operator (3.32), and avoids the difficulty of handling such remainder terms.*

### 3.4 Proof of Lemma 3.10

With the notations of Lemma 3.10, in this section we study the solutions  $X_{0, \omega, k}$  of (3.40) and  $X_{1, \omega, k}$  of (3.42). In order to do it, we perform the change of variable and unknowns

$$t = |k|^{2/3} (x_0^*(\omega, k) - x), \quad f_{m, k}(t) = |k|^{-2/3} X_{m, \omega, k}(x), \quad m \in \{0, 1\}.$$

Then both  $f_{0, k}$  and  $f_{1, k}$  satisfy the ODE

$$f_k' = H_k(t) - f_k^2, \quad (3.51)$$

where

$$H_k : t \in \mathbb{R} \mapsto |k|^{2/3} \left( 1 - e^{-\frac{2t}{|k|^{2/3}}} \right),$$

plus the initial condition

$$f_{0, k}(0) = 0, \quad f_{1, k} \left( -\frac{1}{2} \right) = 2. \quad (3.52)$$

Note that this equation is now independent of  $\omega$ , thus justifying the fact that we omit the dependence in  $\omega$  and simply use the notation  $f_{0,k}$ ,  $f_{1,k}$ .

Cauchy-Lipschitz theorem implies that equations (3.51) and (3.52) uniquely define  $f_{0,k}$  and  $f_{1,k}$  on some maximal intervals denoted  $I_{0,k}$  and  $I_{1,k}$ .

Then, Lemma 3.10 is implied by the following result.

**Lemma 3.13.** *There exists  $k_0 \geq 0$  such that for all  $|k| \geq k_0$ , the following statements are true:*

*The function  $f_{0,k}$  is well-defined and non-negative on  $[0, \infty)$ . The function  $f_{1,k}$  is well-defined and non-negative on  $[-\frac{1}{2}, \infty)$ . Furthermore,*

- for all  $A > 0$ , there exists  $\delta_A$  independent of  $k$  such that for all  $t \in [0, A]$ ,

$$f_{1,k}(t) - f_{0,k}(t) \geq \delta_A.$$

- for all  $\varepsilon > 0$ , there exists  $A_\varepsilon \geq 0$  independent of  $k$  such that for all  $t \geq A_\varepsilon$ ,

$$(1 - \varepsilon)\sqrt{H_k(t)} \leq f_{0,k}(t) \leq f_{1,k}(t) \leq \sqrt{H_k(t)} \leq |k|^{\frac{2}{3}}.$$

We divide the proof of Lemma 3.13 into several steps. We first concentrate on the function  $f_{0,k}$ .

**Lemma 3.14.** *The function  $f_{0,k}$  is well-defined on  $[0, \infty)$ , and verifies, for all  $t > 0$ ,*

$$0 < f_{0,k}(t) < \sqrt{H_k(t)}.$$

As a consequence,  $f_{0,k}$  is increasing on  $[0, \infty)$ .

*Proof.* A direct computation shows that  $f_{0,k}(0) = f'_{0,k}(0) = 0$  and  $f''_{0,k}(0) > 0$ , so there exists  $\eta_{0,k} > 0$  such that for all  $t \in (0, \eta_{0,k})$ ,  $f_{0,k}(t) > 0$ . Denote

$$T_{0,k} = \sup \{ \eta \in I_{0,k} \cap (0, \infty), \forall t \in (0, \eta), f_{0,k}(t) > 0 \},$$

such that  $\eta_{0,k} \leq T_{0,k} \leq \sup I_{0,k}$ . Suppose  $T_{0,k} \in I_{0,k}$ , then  $f_{0,k}(T_{0,k}) = 0$  which implies  $f'_{0,k}(T_{0,k}) = H_k(T_{0,k}) > 0$ , leading to an immediate contradiction. Hence  $T_{0,k} = \sup I_{0,k}$ , and  $f$  is positive on  $I_{0,k} \cap (0, \infty)$ .

Consider now  $g_{0,k} : t \in I_{0,k} \mapsto f_{0,k}(t)^2 - H_k(t)$ . Using (3.51), we easily obtain that for all  $t$  in  $I_{0,k}$ ,

$$g'_{0,k}(t) + 2f_{0,k}(t)g_{0,k}(t) = -H'_k(t) < 0,$$

from which we infer that for all  $t \in I_{0,k} \cap (0, \infty)$ , denoting  $F_{0,k} : t \in I_{0,k} \mapsto \int_0^t f_{0,k}(s) ds$ ,

$$g_{0,k}(t)e^{2F_{0,k}(t)} < g_{0,k}(0)e^{2F_{0,k}(0)} = 0,$$

hence  $f_{0,k}^2 < H_k$  on  $I_{0,k} \cap (0, \infty)$ . This in turn implies  $\sup I_{0,k} = \infty$ , which ends the proof.  $\square$

**Lemma 3.15.** *There exist  $k_0 > 0$  and a positive constant  $C$  such that for  $|k| \geq k_0$ ,*

- For all  $T \in [2, |k|^{2/3}]$ ,

$$\frac{f_{0,k}(T)}{\sqrt{H_k(T)}} \geq 1 - \frac{C \log(T)}{T\sqrt{T}}. \quad (3.53)$$

- For all  $T \in [|k|^{2/3}, \exp(|k|^{2/3})]$ ,

$$\frac{f_{0,k}(T)}{\sqrt{H_k(T)}} \geq 1 - C \frac{\log(T)}{T}. \quad (3.54)$$

- For all  $T \geq |k|^{2/3}$ ,

$$\frac{f_{0,k}(T)}{\sqrt{H_k(T)}} \geq 1 - C \exp(-T/|k|^{2/3}). \quad (3.55)$$

In particular, for  $T \geq |k|^{4/3}$ , we have

$$\frac{f_{0,k}(T)}{\sqrt{H_k(T)}} \geq 1 - C \exp(-\sqrt{T}). \quad (3.56)$$

*Proof.* Let  $T \geq 2$ . For  $A \in [0, T]$ , we introduce  $f_A$  the solution of  $f'_{A,k} = H_k(A) - f_{A,k}^2$  on  $(A, \infty)$  starting from  $f_{A,k}(A) = 0$ . Easy comparison arguments show that  $f_{0,k}(T) \geq f_{A,k}(T)$ . We then compute  $f_{A,k}(T)$ :

$$f_{A,k}(T) = \sqrt{H_k(A)} \frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}}.$$

Accordingly,

$$\begin{aligned} \frac{f_{0,k}(T)}{\sqrt{H_k(T)}} &\geq \sup_{A \in [0, T]} \left\{ \frac{\sqrt{H_k(A)} (1 - e^{-2\sqrt{H_k(A)}(T-A)})}{\sqrt{H_k(T)} (1 + e^{-2\sqrt{H_k(A)}(T-A)})} \right\} \\ &\geq \sup_{A \in [0, T]} \left\{ \sqrt{\frac{1 - e^{-2A/|k|^{2/3}}}{1 - e^{-2T/|k|^{2/3}}} \frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}}} \right\}. \end{aligned}$$

Let us first focus on the case  $2 \leq T \leq |k|^{2/3}$ . On one hand, we have, for some constants  $C$  independent of  $k$ ,  $A$  and  $T$  and which may change from line to line, that,

$$\begin{aligned} \sqrt{\frac{1 - e^{-2A/|k|^{2/3}}}{1 - e^{-2T/|k|^{2/3}}}} &= \sqrt{1 - e^{-2T/|k|^{2/3}} \frac{e^{2(T-A)/|k|^{2/3}} - 1}{1 - e^{-2T/|k|^{2/3}}}} \\ &\geq 1 - C e^{-2T/|k|^{2/3}} \frac{e^{2(T-A)/|k|^{2/3}} - 1}{1 - e^{-2T/|k|^{2/3}}} \geq 1 - C \left( \frac{T-A}{T} \right). \end{aligned}$$

On the other hand, assuming  $A \geq T/2$ , we get, for some positive constants  $c > 0$  independent of  $k$ ,  $A$ , and  $T$ , which may change from line to line,

$$H_k(A) = |k|^{2/3} \left( 1 - e^{-2A/|k|^{2/3}} \right) \geq cA \geq cT,$$

so that

$$\frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}} \geq 1 - 2e^{-2\sqrt{H_k(A)}(T-A)} \geq 1 - 2 \exp\left(-c\sqrt{T}(T-A)\right).$$

Let us now fix  $\beta = \frac{3}{2c}$  where  $c$  is the constant appearing in the exponential in the last estimate. Then the choice  $A = T - \beta \ln(T)/\sqrt{T}$  satisfies  $A \geq T/2$  provided that  $T$  is greater than some fix constant  $T_0$  depending on  $\beta$  only. With that choice, we obtain on one hand,

$$\sqrt{\frac{1 - e^{-2A/|k|^{2/3}}}{1 - e^{-2T/|k|^{2/3}}}} \geq 1 - C \frac{\log(T)}{T\sqrt{T}},$$

and on the other hand,

$$\frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}} \leq 1 - \frac{2}{T\sqrt{T}}.$$

Combining these last two estimates, and using that  $T \mapsto \frac{1}{T\sqrt{T}}$  is negligible in front of  $\log(T)/(T\sqrt{T})$  as  $T$  goes to infinity, we easily obtain (3.53).

For  $T$  satisfying  $T \geq |k|^{2/3}$ , and  $T - A \leq |k|^{2/3}$ , on one hand, we have

$$\begin{aligned} \sqrt{\frac{1 - e^{-2A/|k|^{2/3}}}{1 - e^{-2T/|k|^{2/3}}}} &= \sqrt{1 - e^{-2T/|k|^{2/3}} \frac{e^{2(T-A)/|k|^{2/3}} - 1}{1 - e^{-2T/|k|^{2/3}}}} \geq 1 - C e^{-2T/|k|^{2/3}} \frac{e^{2(T-A)/|k|^{2/3}} - 1}{1 - e^{-2T/|k|^{2/3}}} \\ &\geq 1 - C e^{-2T/|k|^{2/3}} \left( \frac{T-A}{|k|^{2/3}} \right) \geq 1 - C e^{-2T/|k|^{2/3}} \frac{T}{|k|^{2/3}} \left( \frac{T-A}{T} \right). \end{aligned}$$

On the other hand, for  $T \geq |k|^{2/3}$ , and  $A \geq |k|^{2/3}/2$ ,  $H_k(A) \geq c|k|^{2/3}$  for some  $c$  independent of  $k$ , and thus

$$\frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}} \geq 1 - 2e^{-2\sqrt{H_k(A)}(T-A)} \geq 1 - 2 \exp\left(-c|k|^{1/3}(T-A)\right).$$

Therefore, for  $T$  satisfying  $T \in [|k|^{2/3}, \exp(|k|^{2/3})]$ , setting  $A = T - \log T$ , which indeed satisfies  $A \geq |k|^{2/3}/2$  for  $k$  large enough, and  $T - A \leq |k|^{2/3}$ , we obtain

$$\begin{aligned} \frac{f_{0,k}(T)}{\sqrt{H_k(T)}} &\geq \left(1 - Ce^{-2T/|k|^{2/3}} \frac{T}{|k|^{2/3}} \frac{\log(T)}{T}\right) \left(1 - 2 \exp\left(-c|k|^{1/3} \log(T)\right)\right) \\ &\geq 1 - Ce^{-2T/|k|^{2/3}} \frac{T}{|k|^{2/3}} \frac{\log(T)}{T} - \frac{C}{T^c |k|^{1/3}}. \end{aligned}$$

This last estimate easily implies (3.54), since  $\tau \mapsto \tau \exp(-\tau)$  is bounded on  $[1, \infty]$ , and  $1/T^c |k|^{1/3}$  is negligible in front of  $\log(T)/T$  as  $T \rightarrow \infty$  for  $k$  large enough.

For  $T \geq |k|^{2/3}$ , we can also choose  $A = T/2$ , so that

$$\sqrt{\frac{1 - e^{-2A/|k|^{2/3}}}{1 - e^{-2T/|k|^{2/3}}}} \geq \sqrt{1 - e^{-2A/|k|^{2/3}}} \geq 1 - Ce^{-T/|k|^{2/3}},$$

while

$$\frac{1 - e^{-2\sqrt{H_k(A)}(T-A)}}{1 + e^{-2\sqrt{H_k(A)}(T-A)}} \geq 1 - 2 \exp\left(-c|k|^{1/3}T\right).$$

These two estimates easily yield (3.55) for  $k$  large enough.  $\square$

**Remark 3.16.** *It is not difficult to check that the proof of (3.53) can be adapted to obtain the following result: For all  $C_0 > 0$ , there exist  $k_0$  and  $C$  such that (3.53) holds for all  $k$  with  $|k| \geq k_0$  and  $T \in [2, C_0|k|^{2/3}]$ . This result will be useful later on.*

**Lemma 3.17.** *Let  $k_0$  as in Lemma 3.15. Let  $\varepsilon > 0$ . There exists  $A_\varepsilon \geq 0$  such that, for all  $t \geq A_\varepsilon$  and  $|k| \geq k_0$ ,  $(1 - \varepsilon)\sqrt{H_k(t)} \leq f_{0,k}(t)$ .*

*Proof.* Lemma 3.17 is a simple corollary of Lemma 3.15 by noting that

$$\lim_{T \rightarrow \infty} \left(1 - \frac{C \log(T)}{T\sqrt{T}}\right) = 1, \quad \lim_{T \rightarrow \infty} \left(1 - C \frac{\log(T)}{T}\right) = 1, \quad \lim_{T \rightarrow \infty} \left(1 - \exp(-\sqrt{T})\right) = 1,$$

and using the estimates (3.53) for  $T \in [1, |k|^{2/3}]$ , (3.54) for  $T \in [|k|^{2/3}, |k|^{4/3}]$  and (3.56) for  $T \geq |k|^{4/3}$ .  $\square$

We now study the function  $f_{1,k}$ , starting with the following lemma.

**Lemma 3.18.** *There exists  $k_0 \in \mathbb{N}$  such that, for all  $|k| \geq k_0$ ,  $[-1/2, 0] \subset I_{1,k}$  and*

$$0 < \tan\left(\frac{1}{2}\right) \leq f_{1,k}(0) \leq 2.$$

*Proof.* We denote  $t_0 = -\frac{1}{2}$ . First, as for all  $t \in [t_0, 0]$ , we have  $H_k(t) \leq 0$ , we have  $f'_{1,k} \leq -f_{1,k}^2$  on  $[t_0, 0] \cap I_{1,k}$ , hence  $f_{1,k}(t) \leq f_{1,k}(t_0) = 2$  for all  $t \in [t_0, 0] \cap I_{1,k}$ .

Now, since  $H_k$  is increasing, we have  $f'_{1,k} \geq H_k(t_0) - f_{1,k}^2 = -(|H_k(t_0)| + f_{1,k}^2)$  in  $I$ , which implies

$$\arctan\left(\frac{f_{1,k}}{\sqrt{|H_k(t_0)|}}\right)' \geq -\sqrt{|H_k(t_0)|}. \quad (3.57)$$

For each  $t \in I_{1,k}$  with  $t > t_0$ , we integrate (3.57) in  $[t_0, t]$ , and we get

$$\arctan\left(\frac{f_{1,k}(t)}{\sqrt{|H_k(t_0)|}}\right) \geq \arctan\left(\frac{2}{\sqrt{|H_k(t_0)|}}\right) - \sqrt{|H_k(t_0)|}(t - t_0)$$

Seeing that

$$1 \leq \sqrt{|H_k(t_0)|} \leq \sqrt{1 + \frac{1}{|k|^{2/3}}},$$

we have, for all  $t \in [t_0, 0] \cap I_{1,k}$  such that  $f_{1,k}(t) \geq 0$ ,

$$\begin{aligned} \arctan(f_{1,k}(t)) &\geq \arctan\left(\frac{f_{1,k}(t)}{\sqrt{|H_k(t_0)|}}\right) \\ &\geq \arctan\left(\frac{2}{\sqrt{1 + \frac{1}{|k|^{2/3}}}}\right) - \sqrt{1 + \frac{1}{k^{2/3}}}\left(t + \frac{1}{2}\right) \geq \arctan\left(\frac{2}{\sqrt{1 + \frac{1}{|k|^{2/3}}}}\right) - \frac{\sqrt{1 + \frac{1}{k^{2/3}}}}{2}. \end{aligned}$$

Since

$$\lim_{|k| \rightarrow \infty} \left( \arctan\left(\frac{2}{\sqrt{1 + \frac{1}{|k|^{2/3}}}}\right) - \frac{\sqrt{1 + \frac{1}{|k|^{2/3}}}}{2} \right) = \arctan(2) - \frac{1}{2} > \frac{1}{2},$$

we deduce that for all  $|k|$  sufficiently large,  $f_{1,k} > \tan\left(\frac{1}{2}\right) > 0$  in  $[t_0, 0] \cap I_{1,k}$ .

We hence have obtained that  $\tan\left(\frac{1}{2}\right) \leq f_{1,k} \leq 2$  on  $[t_0, 0] \cap I_{1,k}$ , immediately implying that  $[t_0, 0] \cap I_{1,k} = [t_0, 0]$ , which ends the proof.  $\square$

**Lemma 3.19.** *Under the assumptions of Lemma 3.18, we have  $I_{1,k} \cap [0, \infty) = [0, \infty)$ , and for all  $t \geq 0$ ,  $f_{0,k}(t) < f_{1,k}(t)$ .*

*Proof.* As both  $f_{0,k}$  and  $f_{1,k}$  satisfy (3.51), and  $f_{0,k}(0) < f_{1,k}(0)$ , by Cauchy-Lipschitz theorem, we clearly have  $f_{0,k}(t) < f_{1,k}(t)$  for all  $t \in I_{1,k} \cap [0, \infty)$ . Then, for all  $t \in I_{1,k}$ ,  $t \geq 0$ , we have

$$f'_{1,k}(t) = H_k(t) - f_{1,k}(t)^2 < H_k(t) - f_{0,k}(t)^2 = f'_{0,k}(t),$$

Since  $f_{0,k}$  is bounded by  $|k|^{2/3}$  for all  $t \geq 0$  by Lemma 3.14, we deduce  $I_{1,k} \cap [0, \infty) = [0, \infty)$ .  $\square$

**Lemma 3.20.** *There exists  $k_0 \in \mathbb{N}$  such that for each  $|k| \geq k_0$ , there exists  $\tau_k \in (0, 3]$  such that*

- $f_{1,k}$  is strictly decreasing on  $[-1/2, \tau_k)$ ,
- $f_{1,k}$  is strictly increasing on  $(\tau_k, \infty[$ .

As a consequence,  $f_{1,k}(t) \leq \sqrt{H_k(t)}$  for all  $t \geq 3$ .

*Proof.* Directly from the equation we get that  $f'_{1,k}(t) < 0$  for all  $t \in [-1/2, 0]$ . For  $t \geq 0$ , we define the set

$$\mathcal{E}_k = \{T \geq 0 : f'_{1,k}(t) < 0 \forall t \in [0, T]\}.$$

Since  $f'_{1,k}(0) = -f_{1,k}(0)^2 < 0$ , one has that  $\mathcal{E}_k$  is nonempty and  $\tau_k := \sup \mathcal{E}_k > 0$ . By definition we have  $f'_{1,k}(t) < 0$  for all  $t$  in  $[0, \tau)$ . Therefore, for all  $t \in [0, \tau)$ ,  $f_{1,k}(t) \leq f_{1,k}(0) \leq 2$  and then

$$f'_{1,k}(t) \geq H_k(t) - 4 \tag{3.58}$$

for all  $t < \tau$ . Note that the right hand side of (3.58) vanishes exactly at

$$\tau_{\star,k} := -\frac{|k|^{2/3}}{2} \ln\left(1 - \frac{4}{|k|^{2/3}}\right) \xrightarrow{|k| \rightarrow \infty} 2.$$

In particular,  $\tau_{\star,k} \in [0, 3]$  and then  $\tau_k \leq \tau_{\star,k} \leq 3$  for all  $|k|$  large enough.

On the other hand, given any  $\tilde{\tau} > 0$  such that  $f'_{1,k}(\tilde{\tau}) = 0$ , we have  $f''_{1,k}(\tilde{\tau}) = (H_k - f_{1,k}^2)'(\tilde{\tau}) = H'_k(\tilde{\tau}) > 0$ . Then, a very simple reductio ad absurdum shows that  $f'_{1,k}(t) > 0$  on  $(\tilde{\tau}, \infty)$ , which ends the proof.  $\square$

**Lemma 3.21.** *Let  $k_0$  be fixed as in previous Lemma, and  $A > 0$ . There exists  $\delta_A > 0$  such that for all  $|k| \geq k_0$  and all  $t \in [0, A]$ ,  $f_{1,k}(t) - f_{0,k}(t) \geq \delta_A$ .*

*Proof.* We observe that, for all  $t \geq 0$  and all  $k \geq 1$ ,  $H_k(t) \leq 2t$ . So, combining the results from Lemmas 3.19 and 3.20, we obtain that for all  $t \in [0, A]$  and all  $|k|$  greater than  $k_0$ ,

$$f_{0,k}(t) \leq f_{1,k}(t) \leq 2 + \sqrt{H_k(t)} \leq 2 + \sqrt{2t} \leq 2 + \sqrt{2A}.$$

Setting, for all  $t \in [0, A]$ ,  $g_k(t) = f_{1,k}(t) - f_{0,k}(t)$ , we easily check that  $g_k$  satisfies the equation  $g_k'(t) + (f_{1,k}(t) + f_{0,k}(t))g_k(t) = 0$  on  $[0, A]$ . According to the above estimate, we obtain

$$f_{1,k}(t) - f_{0,k}(t) = g_k(0) \exp\left(-\int_0^t (f_{1,k}(s) + f_{0,k}(s)) ds\right) \geq \tan\left(\frac{1}{2}\right) \exp\left(-\int_0^A (f_{1,k}(s) + f_{0,k}(s)) ds\right).$$

The results follows, with

$$\delta_A = \tan\left(\frac{1}{2}\right) \exp\left(-A(4 + 2\sqrt{2A})\right).$$

□

Combining the results of Lemmas 3.14, 3.17, 3.18, 3.19, 3.20 and 3.21, we obtain Lemma 3.13.

## 4 A necessary condition for the resolvent estimate: Proof of Theorem 1.4

Let  $(\omega_n, k_n)_{n \in \mathbb{N}}$  be a sequence of elements in  $\mathbb{R} \times \mathbb{Z}$  satisfying (1.22) and (1.23). For all  $n \in \mathbb{N}$ , we set

$$\begin{aligned} A_n &= |k_n|^{2/3} \log\left(\frac{|k_n|}{|\omega_n| R_0}\right), & x_{0,n}^* &= \log\left(\frac{|k_n|}{|\omega_n|}\right), & \tilde{x}_{0,n}^* &= \min\{x_{0,n}^*, a_1\}, \\ x_{-1,n}^* &= \frac{1}{3}a_0 + \frac{2}{3}\tilde{x}_{0,n}^*, & x_{-2,n}^* &= \frac{2}{3}a_0 + \frac{1}{3}\tilde{x}_{0,n}^*, & \delta_n &= \min\left\{\frac{A_n}{3|k_n|^{2/3}}, \frac{a_1 - a_0}{3}\right\}, \end{aligned}$$

and we note that  $a_0 < x_{-2,n}^* < x_{-1,n}^* < \tilde{x}_{0,n}^* \leq a_1$  with

$$\tilde{x}_{0,n}^* - x_{-1,n}^* = x_{-1,n}^* - x_{-2,n}^* = x_{-2,n}^* - a_0 = \delta_n.$$

Also note that, due to (1.23),  $\delta_n$  is always of the order of  $A_n/|k_n|^{2/3}$ .

We then set, for  $x \in [a_0, \tilde{x}_{0,n}^*]$ ,

$$\tilde{z}_{k_n}(x) = \exp\left(-\int_{a_0}^x X_{0,\omega_n,k_n}(y) dy\right), \quad (4.1)$$

where  $X_{0,\omega_n,k_n}$  is defined in (3.40), and

$$z_{k_n}(x) = \tilde{z}_{k_n}(x)\eta_n(x) \quad \text{where} \quad \eta_n(x) = \eta\left(\frac{x - a_0}{\tilde{x}_{0,n}^* - a_0}\right), \quad (4.2)$$

where  $\eta = \eta(s)$  is a smooth cut-off function taking value 1 for  $s \leq 1/3$ , vanishing for  $s \geq 2/3$ , and taking values in  $[0, 1]$ .

One then easily check that  $z_{k_n}$ , extended by 0 for  $x \in [\tilde{x}_{0,n}^*, a_1]$ , satisfies (3.1) with

$$\begin{aligned} F_{2,k_n} &= 2\partial_x \eta_n \partial_x \tilde{z}_{k_n} + \partial_{xx} \eta_n \tilde{z}_{k_n}, \\ g_{k_n} &= -(\rho(\omega_n, k_n) + X_{0,\omega_n,k_n}(a_0)), \\ z_{k_n} &= 0 \text{ for } x \geq x_{-1,n}, \\ z_{k_n} &= \tilde{z}_{k_n} \text{ for } x \leq x_{-2,n}, \end{aligned}$$

Our next goal is to check that the resolvent condition (3.3) applied to  $z_{k_n}$  necessarily implies some condition. In order to do so, it is convenient to recall from Lemma 3.10 that, for  $n$  large enough, since  $A_n \rightarrow \infty$  due to (1.22), we have

$$\forall x \in [a_0, x_{-1,n}^*], \quad \frac{1}{2}\sqrt{H_{\omega_n,k_n}(x)} \leq X_{0,\omega_n,k_n}(x) \leq \sqrt{H_{\omega_n,k_n}(x)}. \quad (4.3)$$



We then check that  $\sqrt{H_{\omega_n, k_n}(a_0)}$ ,  $\sqrt{H_{\omega_n, k_n}(x_{-2, n}^*)}$  and  $\sqrt{H_{\omega_n, k_n}(x_{-1, n}^*)}$  are all of the same order. Indeed,

$$H_{\omega_n, k_n}(x) = k_n^2 \left( 1 - e^{-2(x_{0, n}^* - x)} \right).$$

In view of the choice of  $x_{-1, n}^*$  and the fact that  $H_{\omega_n, k_n}$  is decreasing, it is thus clear that there exists a constant  $C > 0$  independent of  $n \in \mathbb{N}$  such that

$$\begin{aligned} \frac{1}{C} \sqrt{H_{\omega_n, k_n}(a_0)} &\leq \sqrt{H_{\omega_n, k_n}(x_{-1, n}^*)} \leq \inf_{[a_0, x_{-1, n}^*]} \sqrt{H_{\omega_n, k_n}(x)} \\ &\leq \sup_{[a_0, x_{-1, n}^*]} \sqrt{H_{\omega_n, k_n}(x)} \leq \sqrt{H_{\omega_n, k_n}(a_0)} \end{aligned} \quad (4.4)$$

and

$$\frac{1}{C} \leq \frac{H_{\omega_n, k_n}(a_0)}{|k_n|^2 \delta_n} \leq C, \quad (4.5)$$

where we used (1.23) to guarantee that  $x_{0, n}^* - a_0$  is bounded uniformly in  $n \in \mathbb{N}$ .

Using the above estimate, we immediately have constants  $C > 0$  independent of  $n$  such that

$$\begin{aligned} \|z_{k_n}\|_{L^2(a_0, a_1)}^2 &\geq \|\tilde{z}_{k_n}\|_{L^2(a_0, x_{-2, n}^*)}^2 \geq \frac{1}{C|k_n|\sqrt{\delta_n}}, \\ \|\partial_x z_{k_n}\|_{L^2(a_0, a_1)}^2 &\geq \|\partial_x \tilde{z}_{k_n}\|_{L^2(a_0, x_{-2, n}^*)}^2 \geq \frac{1}{C} |k_n| \sqrt{\delta_n}, \\ \|\partial_x \eta_n \partial_x z_{k_n}\|_{L^2(a_0, a_1)}^2 &\leq \|\partial_x \eta_n\|_{L^\infty(a_0, a_1)}^2 \|\partial_x z_{k_n}\|_{L^2(x_{-2, n}^*, x_{-1, n}^*)}^2 \leq \frac{C}{|\delta_n|^2} e^{-|k_n| \delta_n^{3/2}/C} |k_n| \sqrt{\delta_n}, \\ \|\partial_{xx} \eta_n z_{k_n}\|_{L^2(a_0, a_1)}^2 &\leq \|\partial_{xx} \eta_n\|_{L^\infty(a_0, a_1)}^2 \|z_{k_n}\|_{L^2(x_{-2, n}^*, x_{-1, n}^*)}^2 \leq \frac{C}{|\delta_n|^4} e^{-|k_n| \delta_n^{3/2}/C} \frac{1}{|k_n| \sqrt{\delta_n}}. \end{aligned}$$

In particular, the estimate (3.3) applied to  $z_{k_n}$  yields, for all  $n \in \mathbb{N}$ ,

$$|k_n| \sqrt{\delta_n} + \frac{|k_n|}{\sqrt{\delta_n}} \leq C e^{-|k_n| \delta_n^{3/2}/C} \left( \frac{|k_n|}{\delta_n^{3/2}} + \frac{1}{|k_n| \delta_n^{9/2}} \right) + C (\rho(\omega_n, k_n) + X_{0, \omega_n, k_n}(a_0))^2.$$

Moreover, since  $\delta_n \leq (a_1 - a_0)/3$ ,

$$\frac{|k_n|}{\sqrt{\delta_n}} \geq C |k_n| \sqrt{\delta_n},$$

while, since  $\delta_n \geq A_n/|k_n|^{2/3}$ ,

$$\frac{|k_n|}{\delta_n^{3/2}} \geq C \frac{1}{|k_n| \delta_n^{9/2}}.$$

We then deduce that for all  $n \in \mathbb{N}$ ,

$$\frac{|k_n|}{\sqrt{\delta_n}} \leq C e^{-|k_n| \delta_n^{3/2}/C} \frac{|k_n|}{\delta_n^{3/2}} + C (\rho(\omega_n, k_n) + X_{0, \omega_n, k_n}(a_0))^2.$$

Then, it is not difficult to check that, since  $|k_n| \rightarrow \infty$  and  $A_n \gg \log^{2/3}(|k_n|)$  by (1.22), and  $\delta_n$  is of the order of  $A_n/(|k_n|^{2/3})$  due to (1.23), we get

$$\liminf_{n \rightarrow \infty} \left( \frac{|k_n|}{\sqrt{\delta_n}} \frac{\delta_n^{3/2}}{|k_n| e^{-|k_n| \delta_n^{3/2}/C}} \right) = \liminf_{n \rightarrow \infty} \left( \delta_n e^{|k_n| \delta_n^{3/2}/C} \right) \geq \lim_{n \rightarrow \infty} \left( \frac{A_n}{|k_n|^{2/3}} e^{A_n^{3/2}/C} \right) = \infty,$$

so that the last condition reduces to

$$\frac{|k_n|}{\sqrt{\delta_n}} \leq C (\rho(\omega_n, k_n) + X_{0, \omega_n, k_n}(a_0))^2,$$

for  $n$  large enough, thus showing that there exists  $c_n \in [0, 1]$  such that

$$\frac{|k_n|^2}{\sqrt{H_{\omega_n, k_n}(a_0)}} \leq C \left( \rho(\omega_n, k_n) + c_n \sqrt{H_{\omega_n, k_n}(a_0)} \right)^2. \quad (4.6)$$

The convergence  $c_n \rightarrow 1$  immediately comes from (3.43) and that  $A_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

To be more precise on the rate at which  $c_n$  goes to 1, we specify that  $(\omega_n, k_n)$  satisfies (1.23). This means

$$|x_{0,n}^* - a_0| = \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \leq C_0.$$

We can thus use Lemma 3.15 and Remark 3.16 to get the existence of  $C$  independent of  $n$  such that

$$1 - \frac{C}{|k_n| \log^{3/2} \left( \frac{|k_n|}{|\omega_n| R_0} \right)} \log \left( |k_n|^{2/3} \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \right) \leq \frac{X_{0, \omega_n, k_n}}{\sqrt{H_{\omega_n, k_n}(a_0)}} \leq 1,$$

i.e. the bound

$$1 - \frac{C}{|k_n| \log^{3/2} \left( \frac{|k_n|}{|\omega_n| R_0} \right)} \log \left( |k_n|^{2/3} \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \right) \leq c_n \leq 1. \quad (4.7)$$

on  $c_n = X_{0, \omega_n, k_n}(a_0) / \sqrt{H_{\omega_n, k_n}(a_0)}$ .

Then, due to (1.23), there exists  $C > 0$  such that for all  $n$ ,

$$\frac{1}{C} |k_n| \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \leq \sqrt{H_{\omega_n, k_n}(a_0)} \leq C |k_n| \log \left( \frac{|k_n|}{|\omega_n| R_0} \right).$$

Accordingly, the right hand-side of (4.6) can be bounded by

$$\begin{aligned} & \left( \rho(\omega_n, k_n) + c_n \sqrt{H_{\omega_n, k_n}(a_0)} \right)^2 \\ & \leq 2 \left( \rho(\omega_n, k_n) + \sqrt{H_{\omega_n, k_n}(a_0)} \right)^2 + \frac{C}{\log \left( \frac{|k_n|}{|\omega_n| R_0} \right)} \log^2 \left( |k_n|^{2/3} \log \left( \frac{|k_n|}{|\omega_n| R_0} \right) \right). \end{aligned} \quad (4.8)$$

We now simply remark that the left hand-side of (4.6) satisfies:

$$\frac{|k_n|^2}{\sqrt{H_{\omega_n, k_n}(a_0)}} \geq \frac{1}{C} \frac{|k_n|}{\log \left( \frac{|k_n|}{|\omega_n| R_0} \right)}.$$

Therefore, in the range (1.23), the second term in (4.8) is negligible compared to the left hand-side of (4.6). Accordingly, if (4.6) holds for  $c_n$  satisfying (4.7), we necessarily have (1.24). This concludes the proof of Theorem 1.4.

## 5 Comments

### 5.1 Estimates on the time of observability

Although the approach we proposed here allows to deal with several models at once, we should underline that the time of observability provided by this approach is not known or badly estimated. In fact, it is known that if the resolvent estimate (1.11) holds, then we can derive a time estimate for the observability inequality (1.10) depending on the constants in (1.11), see [18]. However, this estimate on the time required for observability is, in general, far from optimal.

This remark suggests the development of approaches able to track down the time required for observability, maybe based on suitable weighted estimates in the spirit of Carleman estimates for the waves,

see e.g. [15], able to precisely track down the boundary conditions, similarly to what has been done recently in the context of elliptic equations in [4] for instance.

Semiclassical approaches with detailed analysis of the boundary data could also be adapted to deal with general boundary conditions given by a kernel  $\rho$ , similarly to what has been developed in [2, 3]. This would probably be the first step to deal with more general geometries as well, and we refer also to the forthcoming work [10] for results in this direction.

## 5.2 More general kernels $\rho$

Our approach does not truly require the kernel  $\rho = \rho(\omega, k)$  in (1.12) to be real valued, and the above proofs can easily be adapted to deal with cases in which  $\rho$  is complex valued. This remark can be useful to deal with wave models containing dissipative terms.

For instance, to our knowledge, the null-controllability of the damped wave equation with absorbing boundary conditions on the sphere  $S(R_0)$  and a control function  $u \in L^2((0, T) \times S(R_1))$  is still an open problem:

$$\begin{cases} \partial_{tt}z(t, x) - \Delta z(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ z(t, x) = u(t, x), & \text{on } (0, T) \times S(R_1), \\ \partial_\nu z(t, x) + \partial_t z(t, x) = 0, & \text{on } (0, T) \times S(R_0), \\ (z(0, \cdot), \partial_t z(0, \cdot)) = (z^0, z^1), & \text{in } A(R_0, R_1). \end{cases} \quad (5.1)$$

By duality, see e.g. [17, 21], such property would be equivalent to the observability inequality

$$\|(y(T), \partial_t y(T))\|_{H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))} \leq C \|\partial_\nu y|_{S(R_1)}\|_{L^2((0, T) \times S(R_1))} \quad (5.2)$$

for all solutions  $y$  of

$$\begin{cases} \partial_{tt}y(t, x) - \Delta y(t, x) = 0, & \text{in } (0, T) \times A(R_0, R_1), \\ y(t, x) = 0, & \text{on } (0, T) \times S(R_1), \\ \partial_\nu y(t, x) + \partial_t y(t, x) = 0, & \text{on } (0, T) \times S(R_0), \\ (y(0, \cdot), \partial_t y(0, \cdot)) = (y^0, y^1), & \text{in } A(R_0, R_1), \end{cases} \quad (5.3)$$

with  $(y^0, y^1) \in H^1(A(R_0, R_1)) \times L^2(A(R_0, R_1))$  with  $y^0|_{S(R_1)} = 0$ .

It is then clear that resolvent estimates could be derived from our approach, with in this case,  $\rho(\omega, k) = iR_0\omega$ . However, we do not know if the resolvent estimates we would get this way would yield (5.2) in some time  $T$ .

Note that for this model, the propagation of regularity and of semi-classical measures has been studied and analyzed precisely in the literature, see [3, 1]. Despite of this, the usual compactness-uniqueness arguments to establish observability inequality do not apply since the wave equation (5.3) cannot be solved backward in time, so that to our knowledge, the observability inequality (5.2) for solutions of (5.3) is open.

**Acknowledgments.** The authors are indebted to Belhassen Dehman and Enrique Zuazua for fruitful discussions and interesting comments, and for pointing to our attention the related work [10].

## A Proof of Corollary 1.3

The condition (1.15) clearly holds when  $\rho(\omega, k) \geq 0$  (with  $\delta = (1 - \varepsilon)^2$  for instance), and we can thus focus on the case  $\rho(\omega, k) \leq 0$ .

Now, the right hand-side of (1.15) is negative if  $|\rho + 1/2| \geq \sqrt{k^2 - \omega^2 R_0^2 + 1/4}$ , and in particular if

$$\rho(\omega, k) \leq -\sqrt{k^2 - \omega^2 R_0^2 + \frac{1}{4}} - \frac{1}{2}. \quad (A.1)$$

Now, since in the range (1.14),  $\omega^2 R_0^2 \geq k^2$  and  $k^2 - \omega^2 R_0^2$  is of the order of  $2\mathbf{r}|k|^{4/3}$  at least, when  $\omega^2 R_0^2 + k^2$  is large enough, the first condition in (1.21) implies (A.1) and thus (1.15) obviously holds.

Finally, if  $\rho(\omega, k) \leq 0$  and the second condition in (1.21) holds, then, necessarily  $\gamma \leq 1$ , and taking  $\varepsilon = \gamma/2$ ,  $I_\varepsilon(\omega, k)^2 \geq \varepsilon^2(k^2 - \omega^2 R_0^2)$ , while  $k^2 - \omega^2 R_0^2 - \rho(\omega, k)^2 - \rho(\omega, k) \leq (k^2 - \omega^2 R_0^2) + 1/4$ . Since in the range (1.14),  $\omega^2 R_0^2 \leq k^2$  and  $k^2 - \omega^2 R_0^2$  is of the order of  $2\mathbf{r}|k|^{4/3}$  at least, it is clear that (1.15) holds when  $\omega^2 R_0^2 + k^2$  is large enough with  $\delta = \varepsilon^2/2$ .

## References

- [1] L. Aloui. Stabilisation Neumann pour l'équation des ondes dans un domaine extérieur. *J. Math. Pures Appl. (9)*, 81(11):1113–1134, 2002.
- [2] C. Bardos, G. Lebeau, and J. Rauch. Un exemple d'utilisation des notions de propagation pour le contrôle et la stabilisation de problèmes hyperboliques. *Rend. Sem. Mat. Univ. Politec. Torino*, (Special Issue):11–31 (1989), 1988. Nonlinear hyperbolic equations in applied sciences.
- [3] C. Bardos, G. Lebeau, and J. Rauch. Sharp sufficient conditions for the observation, control and stabilization of waves from the boundary. *SIAM J. Control and Optim.*, 30(5):1024–1065, 1992.
- [4] M. Bellassoued and J. Le Rousseau. Carleman estimates for elliptic operators with complex coefficients. Part I: Boundary value problems. *J. Math. Pures Appl. (9)*, 104(4):657–728, 2015.
- [5] I. Benabbas and D. E. Teniou. Observability of wave equation with Ventcel dynamic condition. *Evol. Equ. Control Theory*, 7(4):545–570, 2018.
- [6] N. Burq and P. Gérard. Condition nécessaire et suffisante pour la contrôlabilité exacte des ondes. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(7):749–752, 1997.
- [7] N. Burq and M. Zworski. Control theory and high energy eigenfunctions. In *Journées “Équations aux Dérivées Partielles”*, pages Exp. No. XIII, 10. École Polytech., Palaiseau, 2004.
- [8] L. Caffarelli and L. Silvestre. An extension problem related to the fractional Laplacian. *Comm. Partial Differential Equations*, 32(7-9):1245–1260, 2007.
- [9] C. Conca, J. Planchard, and M. Vanninathan. *Fluids and periodic structures*, volume 38 of *RAM: Research in Applied Mathematics*. John Wiley & Sons Ltd., Chichester, 1995.
- [10] B. Dehman and E. Zuazua. Sidewise energy estimates for the wave equation, in preparation.
- [11] S. Ervedoza and M. Vanninathan. Controllability of a simplified model of fluid-structure interaction. *ESAIM: Control, Optimisation and Calculus of Variations*, 20:547–575, 4 2014.
- [12] F. G. Friedlander. The wave front set of the solution of a simple initial-boundary value problem with glancing rays. *Math. Proc. Cambridge Philos. Soc.*, 79(1):145–159, 1976.
- [13] C. G. Gal and L. Tebou. Carleman inequalities for wave equations with oscillatory boundary conditions and application. *SIAM J. Control Optim.*, 55(1):324–364, 2017.
- [14] L. F. Ho. Observabilité frontière de l'équation des ondes. *C. R. Acad. Sci. Paris Sér. I Math.*, 302(12):443–446, 1986.
- [15] V. Isakov. *Inverse problems for partial differential equations*, volume 127 of *Applied Mathematical Sciences*. Springer, New York, second edition, 2006.
- [16] J.-L. Lions. *Contrôlabilité exacte, Stabilisation et Perturbations de Systèmes Distribués. Tome 1. Contrôlabilité exacte*, volume RMA 8. Masson, 1988.
- [17] J.-L. Lions. Exact controllability, stabilization and perturbations for distributed systems. *SIAM Rev.*, 30(1):1–68, 1988.
- [18] L. Miller. Controllability cost of conservative systems: resolvent condition and transmutation. *J. Funct. Anal.*, 218(2):425–444, 2005.
- [19] J.-P. Raymond and M. Vanninathan. Exact controllability in fluid-solid structure: the Helmholtz model. *ESAIM Control Optim. Calc. Var.*, 11(2):180–203 (electronic), 2005.
- [20] M. Tucsnak and M. Vanninathan. Locally distributed control for a model of fluid-structure interaction. *Systems Control Lett.*, 58(8):547–552, 2009.
- [21] M. Tucsnak and G. Weiss. *Observation and control for operator semigroups*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2009.