



Diversity, Functions and Antibiotic Resistance of Sediment Microbial Communities From Lake Geneva Are Driven by the Spatial Distribution of Anthropogenic Contamination

Emilie Lyautey, Chloé Bonnineau, Patrick Billard, Jean-Luc Loizeau, Emmanuel Naffrechoux, Ahmed Tlili, Edward Topp, Benoît J.D. Ferrari, Stéphane Pesce

► To cite this version:

Emilie Lyautey, Chloé Bonnineau, Patrick Billard, Jean-Luc Loizeau, Emmanuel Naffrechoux, et al.. Diversity, Functions and Antibiotic Resistance of Sediment Microbial Communities From Lake Geneva Are Driven by the Spatial Distribution of Anthropogenic Contamination. *Frontiers in Microbiology*, 2021, 12, <10.3389/fmicb.2021.738629>. <hal-03403564>

HAL Id: hal-03403564

<https://hal.science/hal-03403564v1>

Submitted on 10 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Diversity, Functions and Antibiotic Resistance of Sediment Microbial Communities From Lake Geneva Are Driven by the Spatial Distribution of Anthropogenic Contamination

Emilie Lyautey^{1,2}, Chloé Bonnineau¹, Patrick Billard³, Jean-Luc Loizeau⁴, Emmanuel Naffrechoux⁵, Ahmed Tlili⁶, Edward Topp^{7,8}, Benoît J.D. Ferrari⁹ and Stéphane Pesce^{1*}

OPEN ACCESS

Edited by:

Carmen Palacios,
Université de Perpignan Via Domitia,
France

Reviewed by:

Kevin Thomas Finneran,
Clemson University, United States
Naresh Singhal,
The University of Auckland,
New Zealand

*Correspondence:

Stéphane Pesce
stephane.pesce@inrae.fr

Specialty section:

This article was submitted to
Microbiotechnology,
a section of the journal
Frontiers in Microbiology

Received: 09 July 2021

Accepted: 28 September 2021

Published: 18 October 2021

Citation:

Lyautey E, Bonnineau C, Billard P,
Loizeau J-L, Naffrechoux E, Tlili A,
Topp E, Ferrari BJD and Pesce S
(2021) Diversity, Functions and
Antibiotic Resistance of Sediment
Microbial Communities From Lake
Geneva Are Driven by the Spatial
Distribution of Anthropogenic
Contamination.
Front. Microbiol. 12:738629.
doi: 10.3389/fmicb.2021.738629

¹ INRAE UR RiverLy, Villeurbanne, France, ² INRAE, Université Savoie Mont Blanc, CARRTEL, Thonon-les-Bains, France, ³ Université de Lorraine, CNRS, LIEC, Nancy, France, ⁴ Department F.A. Forel for Environmental and Aquatic Sciences, University of Geneva, Geneva, Switzerland, ⁵ EDYTEM, CNRS, Université Savoie Mont Blanc, Chambéry, France, ⁶ Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland, ⁷ Agriculture and Agri-Food Canada, London, ON, Canada, ⁸ Department of Biology, University of Western Ontario, London, ON, Canada, ⁹ Swiss Centre for Applied Ecotoxicology (Ecotox Centre), Lausanne, Switzerland

Lake sediments are natural receptors for a wide range of anthropogenic contaminants including organic matter and toxicants such as trace metals, polycyclic aromatic hydrocarbons, polychlorinated biphenyls that accumulate over time. This contamination can impact benthic communities, including microorganisms which play a crucial role in biogeochemical cycling and food-webs. The present survey aimed at exploring whether anthropogenic contamination, at a large lake scale, can influence the diversity, structure and functions of microbial communities associated to surface sediment, as well as their genetic potential for resistance to metals and antibiotics. Changes in the characteristics of these communities were assessed in surface sediments collected in Lake Geneva from eight sampling sites in October 2017 and May 2018. These sampling sites were characterized by a large concentration range of metal and organic compound contamination. Variation between the two sampling periods were very limited for all sampling sites and measured microbial parameters. In contrast, spatial variations were observed, with two sites being distinct from each other, and from the other six sites. Benthic communities from the most contaminated sampling site (Vidy Bay, near the city of Lausanne) were characterized by the lowest bacterial and archaeal diversity, a distinct community composition, the highest abundance of antibiotic resistance genes and functional (respiration, denitrification, methanogenesis, phosphatase, and beta-glucosidase) activity levels. The second sampling site which is highly influenced by inputs from the Rhône River, exhibited low levels of diversity, a distinct community composition, high abundance of antibiotic resistance genes and the highest bacterial abundance. Overall, our results suggest that local anthropogenic contamination, including organic

matter and toxicants, is a major driver of the diversity and functioning of sediment-microbial communities in Lake Geneva. This highlights the need to consider benthic microbial communities and a suite of complementary ecotoxicological endpoints for more effective environmental risk assessments of contaminants in lake sediments.

Keywords: benthic communities, microbial ecotoxicology, metals, organic matter, PCB, PAH, resistance genes, urban contamination

INTRODUCTION

The sediment compartment in lakes can act as a sink for contaminants, including not only nutrients and organic matter but also metals, persistent organic pollutants [e.g., polycyclic aromatic hydrocarbons, polychlorinated biphenyls (PAHs); polychlorinated biphenyls, polychlorinated biphenyls (PCBs)] and other substances such as pesticides or pollutants of emerging concerns (e.g., pharmaceuticals including antibiotics) (Xu et al., 2017; Codling et al., 2018; Golovko et al., 2020). Due to their hydrophobic nature, once deposited most of these contaminants associate strongly to the sediment (Sedláček et al., 2020). Therefore, the vertical depth profiles of contaminants in sediment cores generally reflect the temporal history of contamination, with increasing depth being associated with older contamination events (Lécrivain et al., 2018). On the other hand, horizontal gradients of contamination depend mainly on the distance from the contamination sources. The contamination of surface sediments (0–5 cm layer) in the littoral and pelagic zones of lakes can thus exhibit substantial spatial heterogeneity according to the distribution of point- and diffuse sources of pollution in the surrounding watershed, and contaminant fluxes carried into the lake by tributaries (Marvin et al., 2004; Lécrivain et al., 2019). Therefore, benthic communities in lake sediments can experience different exposures to contaminants including nutrients and organic matter according to their location. However, the resulting effects on these communities still under-investigated (Pesce et al., 2018) and little is known about the potential relationships between the spatial contamination heterogeneity in lake surface sediment, and the consequent spatial alterations of benthic community diversity, structure and functions (Haller et al., 2011; Sauvain et al., 2014; Jin et al., 2019).

In lake surface sediments, microorganisms are highly abundant and characterized by a large taxonomic and functional diversity, making them key players in a multitude of ecosystem processes (Schallenberg and Kalff, 1993; Pearman et al., 2020). These processes include biogeochemical cycling of nitrogen, the recycling of autochthonous and allochthonous organic matter, and the dissipation of organic contaminants (Vadeboncoeur et al., 2002; Haglund et al., 2003; Bedard, 2008; Schultz and Urban, 2008). Several laboratory studies have shown that environmental concentrations of metals (Mahamoud Ahmed et al., 2018, 2020) and organic contaminants such as pesticides (Widenfalk et al., 2004, 2008), PCBs (Diepens et al., 2015) and PAHs (Zhang et al., 2014) can affect the diversity, structure and functional potential of exposed sediment microbial communities. The structural and functional properties of microbial communities in lake sediments can vary horizontally at

a relatively limited geographical scale, according to several local environmental factors (Krzmarzick et al., 2013; Pearman et al., 2020). However, the role of anthropogenic contamination in the spatial structuration of those microbial assemblages and the resulting effects on the functional properties of lake benthic ecosystems remain largely unknown.

The contamination of Lake Geneva has been monitored with a focus on various groups of contaminants such as PAHs, PCBs, metals, pharmaceuticals, and pesticides in surface water and sediments (e.g., Poté et al., 2008; Perazzolo et al., 2010; Hoerger et al., 2014; Larras et al., 2016). For example, a large sampling survey undertaken in 2015 mapped the contamination of the surface sediments at the whole lake scale (Loizeau et al., 2017). Sediments in the Vidy Bay near the city of Lausanne were the most contaminated by metals, organic pollutants and organic matter (Loizeau et al., 2017). The high contamination of sediments in this area has been documented for decades (Grandjean et al., 1990; Poté et al., 2008; Loizeau et al., 2013; Gascon Diez et al., 2014; Masson and Tercier-Waeber, 2014). This is mainly due to historic inputs from urban sewage effluents (Gascón Díez et al., 2017), which is also reflected by the prevalence of fecal-indicator bacteria (Poté et al., 2008) and antibiotic-resistance genes (Czekalski et al., 2014). The contamination has been suggested as an important driver of microbial community structure (Haller et al., 2011). Besides this hotspot of contamination, there is relative spatial heterogeneity in the contamination at the whole Lake scale with a strong influence of the Rhone River, which is the main Lake tributary, and of the lakeshore (Loizeau et al., 2017).

In this context, we performed two sampling campaigns to investigate the spatio-temporal variations in benthic microbial communities collected in surface sediments from eight sampling sites in Lake Geneva. These sites, which are widely geographically distributed at the lake scale, cover a large range of contamination levels by metals and organic toxicants (Loizeau et al., 2017). We included one site located in the Vidy Bay and one site located near the Rhone River mouth into the lake. The objective was to assess whether spatial and/or temporal changes in the structural and functional characteristics of the studied community are driven by anthropogenic contamination. For this purpose, microbial communities in sediment samples were characterized for bacterial and archaeal population diversity, composition and structure, bacterial abundance, functional (respiration, denitrification, methanogenesis, leucine aminopeptidase, phosphatase, and beta-glucosidase) potential activities and the abundance of gene targets associated with resistance to selected metals and antibiotics. We expected the microbial community structure from these eight sites

to exhibit spatial and temporal dynamics, with functional properties linked both to community composition and to the environmental conditions prevailing in the sediment compartment on the sampling dates. Contamination levels by metals and organic toxicants were expected to select for more tolerant populations at the sampling site scale, leading to change in community structure and to increasing genetic potential of resistance, with possible impacts on ecosystem functions.

MATERIALS AND METHODS

Sampling Sites and Sample Collection

Lake Geneva is the largest natural lake in western Europe (580 km², 72 km long and 14 km wide), with a maximum depth of 309 m, and a watershed surface area of 7,999 km². The lake is located between France and Switzerland, and its major tributary and outflow is the Rhône River. The water residence time is 11.3 years (CIPEL, 2018). Lake Geneva is a deep monomictic lake with irregular complete winter overturns occurring on average every 5 years (Schwefel et al., 2016).

Sample collection was carried out on October 2017 (26th) and May 2018 (23rd, 24th, and 25th) at eight sampling sites. Sampling sites were referred as sites number 5, 6, 21, 29, 32, 36, 53, and 78 (**Figure 1**), according to the map of sampling sites of Loizeau et al. (2017). The sites were chosen to cover a large range of contamination levels by metals and organic toxicants, as reported by Loizeau et al. (2017). Sites 29, 32, 53, and 78 were sampled both years whereas sites 5, 6, 21, and 36 were only sampled in 2018 due to weather conditions in 2017. Sampling site descriptions, sampling date, geographical coordinates, and depth are provided in **Table 1**.

Surface sediment sampling was carried out using an Eckman grab sampler, and the top 15–25 cm of sediment were grabbed. Samples for subsequent analyses were collected in the Eckman grab, from the first 2-cm of sediment directly in the field, and stored for further analyses. For each site, three independent samples were collected. Each sample was sub-sampled directly on the boat, and samples were stored at 4°C during transport to the laboratory with arrival within 6 h.

Sediment Chemical Characterization

The analyses of the physico-chemical properties of sediment from sites 29, 32, 53, and 78 were carried out on samples collected on October 2017, and from sites 5, 6, 21, and 36 on samples collected on May 2018. These analyses included concentrations of trace metals (Ti, Cr, Co, Ni, Cu, Zn, As, Mo, Ag, Cd, Sn, Pb, and Hg), 12 PCBs, 16 PAHs, and ancillary parameters [grain size distribution, organic matter (OM), carbonate and total organic carbon content (TOC), total phosphorus content (P_{tot})]. Ti was used as a proxy of the Rhône River influence on Lake Geneva, as suspended particle load from this river mainly originate from the erosion of Alpine rocks richer in Ti-bearing minerals than rocks from the calcareous Jura mountains drained by the others tributaries, or calcite particles directly precipitated from the water column.

Sediment grain-size distribution was determined on wet sediments using a laser diffraction analyzer (Coulter LS-100, Beckman-Coulter, United States), following the procedure described by Loizeau et al. (1994).


Samples were freeze-dried in a CHRIST BETA 1–8 K freeze drying unit (−54°C, 6 Pa) for a minimum of 48 h. Organic matter and calcium carbonate (CaCO₃) contents in sediments were estimated by Loss on Ignition. Samples were heated to 550°C for 30 min to estimate the OM mass loss and then heated to 1000°C for another 30 min to estimate the CaCO₃ content (Dean, 1974). The CaCO₃ content was calculated by multiplying the mass loss at 1000°C by 2.2742, the molar mass ratio of calcite to carbon dioxide.

Total phosphorus (TP) was measured with a spectrophotometer (Helios Gamma UV-Vis Thermo Electroporation, Thermo scientific, United States) at 850 nm. The sediment sample preparation for TP analysis was performed as follow; 50 mg of ignited sediment (after heating at 550°C for 1 h) were diluted in 5 mL HCl 1 N and introduced in centrifuge tubes. The mixture was ultrasonicated (at ambient temperature) during 16 h and centrifuged (4000 × rpm) during 20 min. The TP concentration was determined by measuring the absorbance of the blue complex obtained after reduction of molybdophosphoric acid according to previously described method (Murphy and Riley, 1962; Harwood et al., 1969; Burrus et al., 1990).

Quadrupole based inductively coupled plasma mass spectrometry (ICPMS, model 7700 series, Agilent) was performed following sediment digestion in Teflon bombs heated to 150°C in analytical grade 2 M HNO₃ according to the Swiss Soil Protection Ordinance (OSol, Office Fédéral de l'environnement [OFEV], 1998). Multi-element standard solutions of different concentrations (0, 0.02, 1, 5, 20, 100, and 200 mg L^{−1}) were used for calibration. The total variation coefficients for triplicate sample measurements were less than 10%.

Total Hg was analyzed by Cold Vapor Atomic Absorption 187 Spectrophotometry (CV-AAS) using an Advance Mercury Analyzer (Model AMA 254, Altec, Czech Republic) through dry mineralization and pre-concentration of Hg and amalgamation on a gold trap (Szakova et al., 2004). The detection limit and working range were 0.01 ng and 0.05–600 ng, respectively. The relative error was usually ±5% and always less than 10% (Roos-Barraclough et al., 2002).

Polychlorinated biphenyls and PAHs were quantified on freeze-dried sediments. PCBs and PAHs were characterized according to the method described in Mwanamoki et al. (2014). Briefly, 5 g of dry sediment were extracted with a mixture of acetone (20% of the volume) and hexane (80% of the volume) for 4 h with a Soxhlet apparatus. At this point, interfering sulfur compounds were removed by adding activated copper to the extract. The organic extract was then concentrated to 1 mL in a vacuum rotary evaporator. The extract was further subject to fractionation and clean-up over a chromatographic column containing 3 g of Silica gel. Three separated fractions were collected: first with 16 mL of hexane, then 35 mL of hexane, and finally 50 mL of hexane:dichloromethane (v/v, 1:1). The



first fraction should contain PCBs while PAHs are distributed in the three fractions. Following reduction of the volume, chemicals were measured by gas chromatography with triple mass spectrometry detection (GC–MS/MS, Thermo Scientific, TSQ Quantum XLS Ultra, Waltham, MA, United States).

Bacterial Abundance and Functional Potential of the Microbial Community

Bacterial abundance was estimated by flow cytometry as described in Frossard et al. (2012). Briefly, 3 g of wet sediment were suspended in 10 mL of 2% formalin containing 0.1% pyrophosphate. After an ultrasonic treatment for 1 min (Branson Digital Sonifier 250, Germany) and homogenization, 1 mL subsample of each suspension was placed on top of 0.5 mL Histodenz solution (Sigma-Aldrich, Buchs, Switzerland) and centrifuged at $17,135 \times g$ at 4°C for 90 min. The upper layer, containing bacterial cells, was recovered and stained with 0.1 mL mL⁻¹ of SYBER® Green I (Promega, Switzerland) in anhydrous dimethylsulfoxide and incubated during 15 min in the dark. A known concentration of fluorescent beads (Beckman Coulter, Switzerland) was added to the samples as a standard to determine the cell concentration. Samples were analysed with a Gallios flow cytometer (Beckman Coulter, Switzerland).

The functional potential of the microbial community was assessed through the measurement of three enzymatic activities: β -glucosidase, β -glu, EC 3.2.1.21; leucine aminopeptidase, LAP, EC 3.4.11.1; and phosphatase, Pase, EC 3.1.3.1; and three metabolic activities: aerobic respiration, denitrification and methanogenesis.

Briefly, to measure enzymatic activities, 1.2 g of wet sediment were incubated for 30 min with pre-determined optimal concentrations of the fluorogenic substrates 4-methylumbelliferyl- β -D-glucopyranoside (MUF-Glu, CAS No. 18997-57-4) for β -glu, L-leucine-7-amido-4-methylcoumarin

hydrochloride (Leu-AMC, CAS No. 62480-44-8) for LAP, and 4-methylumbelliferyl phosphatase (MUF-P, CAS No. 3368-04-5) for Pase (Mahamoud Ahmed et al., 2018). After the activities were stopped with glycine buffer (0.05 M glycine, 0.2 M NH₄OH, pH 10.4), the samples were centrifuged and the fluorescence in the supernatant was measured using a microplate reader (Exc: 360 nm/Em: 460 nm, Synergy HT BioTek Instruments). Results are expressed as nmol of fluorochrome (MUF or MCA) per g of sediment dw⁻¹ h⁻¹.

Potential rates of aerobic respiration, denitrification and methanogenesis were measured with 10 g of wet sediment according to the protocols described by Foulquier et al. (2013) and slightly adapted by Mahamoud Ahmed et al. (2018). Sediment was mixed with 10 mL of distilled water under aerobic conditions (aerobic respiration), or 10 mL of a KNO₃ (2.16 g L⁻¹) solution under anaerobic conditions (denitrification), or 10 mL of distilled water under anaerobic conditions (methanogenesis) in 150-mL glass flasks with rubber stoppers. Incubation flasks assigned to denitrification and methanogenesis measurements were purged three times with He to achieve anaerobiosis, and the internal pressure was then adjusted to atmospheric. For denitrification measurements, 15 mL of acetylene (C₂H₂, 10% v/v final volume) was added to inhibit N₂O reductase. All samples were incubated at 20°C in the dark with gentle shaking. After 2 h and 5 h, headspace gasses were sampled and analyzed by gas chromatography (Agilent 490 MICRO GC). Aerobic respiration, denitrification and methanogenesis activities are expressed as ng of gaseous product (CO₂ or N₂O or CH₄, respectively) per g of sediment dw⁻¹ h⁻¹.

Microbial Community Structure

DNA Extraction

Microbial sediment DNA was extracted from 0.5 g of wet sediment using a NucleoSpin Soil Kit (Macherey-Nagel EURL)

TABLE 1 | Presentations of the sampling site characteristics: sampling date (in bold, those when the presented physico-chemical parameters were measured), geographical coordinates, depth, and physico-chemical sediment characteristics: organic matter (OM) content (\pm SD), carbonate content (\pm SD), total organic carbon (TOC) content, total phosphorus (Ptot), and Ti concentrations, median grain size (\pm SD), and the sums of sediment concentrations of the following trace metals (Cr, Co, Ni, Cu, Zn, As, Mo, Ag, Cd, Sn, Pb, and Hg), of the seven indicator PCBs, and of PAHs.

Site	Date	Coordinates		Depth (m)	Organic matter		Carbonate	TOC	P _{tot}	Median grain size		Ti	Σ other metals		Σ7PCBi	ΣPAH
		Longitude	Latitude		± SD (%)	± SD (%)				± SD (μm)	(mg kg ⁻¹ dw)		(mg kg ⁻¹ dw)	(μg kg ⁻¹ dw)		
5	May 18	E6.18008	N46.26927	47	8.64 ± 0.35	48.14 ± 0.83	4.35	848.6		14.77 ± 3.26	37.3 ± 1.0	206.6	5.47		1120.8	
6	May 18	E6.19270	N46.29642	55	8.95 ± 0.10	44.33 ± 0.26	4.59	834.8		7.73 ± 2.73	37.6 ± 1.5	294.5	10.53		1372.4	
21	May 18	E6.33359	N46.37907	50	7.18 ± 0.33	42.09 ± 0.76	2.93	750.4		7.98 ± 2.79	75 ± 1.6	267.1	9.81		1139.4	
29	May 18, Oct 17	E6.38462	N46.45147	95	6.38 ± 0.13	31.21 ± 0.22	2.76	872.0		10.57 ± 3.01	405.5 ± 1.8	321.8	7.40		1922.2	
32	May 18, Oct 17	E6.42316	N46.46618	22	5.27 ± 0.21	36.61 ± 0.36	2.08	670.1		14.95 ± 3.90	177.7 ± 4.5	212.1	7.69		2262.5	
36	May 18	E6.45097	N. 46.37102	32	4.79 ± 0.04	48.87 ± 0.35	2.67	549.3		19.07 ± 5.31	42.5 ± 2.4	169.8	5.20		13581.1	
53	May 18, Oct 17	E6.58810	N46.51014	44	13.65 ± 0.09	19.70 ± 0.23	6.96	3900.0		34.82 ± 5.00	170.7 ± 6.2	1193.4	386.55		25261.9	
78	May 18, Oct 17	E6.89261	N46.40998	60	2.73 ± 0.30	14.40 ± 0.32	0.68	659.4		11.12 ± 2.52	1735.6 ± 30.2	293.0	0.89		774.3	

Detailed metals, PCB and HAP concentrations are presented in **Supplementary Tables 1–3**.

following the manufacturer's instructions with SL1 lysis buffer and additive Enhancer SX buffer. The extracted DNA was quantified fluorometrically after staining with QuantiFluor dsDNA Dye (QuantiFluor dsDNA System, Promega) using a Plate ChameleonTM fluorometer (Hidex; excitation 485 nm, emission 590 nm).

High-Throughput 16S rRNA Gene Sequencing and Bioinformatic Analysis

PCR amplification for high-throughput 16S rRNA gene amplicon sequencing was carried out with the universal primer pair 515F and 909R targeting the V4–V5 hypervariable region of the 16S rRNA gene (Wang and Qian, 2009). Indexes were integrated to both primers following the dual-indexing procedure described by Kozich et al. (2013). Triplicate PCR amplification for each sample was carried out with a total amount of \sim 5 ng of DNA per reaction. Amplicon products were quantified using the Picogreen assay (Life Technologies, Carlsbad, United States) and pooled equimolarly. The final pool was purified with CleanPCR beads (CleanNA). Sequencing was done by Fasteris (Geneva, Switzerland) on an Illumina HiSeq system with 2×300 bp. The analysis yielded 3.7 Gb of sequences with an average error rate of 1.480%, and average Q30 of 85.0%. Adapters were removed using Trimmomatic (Bolger et al., 2014) and barcodes sorted using a Fasteris internal script. Sequences were then processed using the FROGS (Find Rapidly OTUs with Galaxy Solution) Galaxy-supported pipeline (Escudié et al., 2018). Paired-end reads were joined using FLASH (Magoë and Salzberg, 2011) and a quality check was performed using FastQC. Sequences with primers having no mismatch were kept. They were then filtered by size (350–500 bp) and those containing N bases were discarded. The 16S rRNA gene sequences were then denoised and clustered using the Swarm method (Mahé et al., 2015) with a three-base maximum difference, deletion of clusters with less than 0.005% abundance and cluster occurrence in a minimum of two samples of the total library. Chimeras were removed using vchime of vsearch package (Rognes et al., 2016). Affiliation was done using the Silva SSU database 123 (Quast et al., 2013) through BLAST (Altschul et al., 1990) with allowed multiple affiliation and manual curation. All analyses were done on the Galaxy instance of the INRA MIGALE bioinformatics platform¹. Sequences are available at GenBank under accession number PRJNA742879.

Microbial Community Genetic Resistance Potential

Real-time PCR quantification of bacterial 16S rRNA gene abundance was carried out using the primers 968F and 1401R according to Cébron et al. (2008). For the quantification of *cusA* (encoding a copper efflux system inner membrane protein), the primer pair *cusF1* and *cusR2* (Mahamoud Ahmed et al., 2020) was used, and for the quantification of *czcA* (inner membrane protein contributing to cobalt, zinc, and cadmium efflux) the primer pair *czcF3* (5'-CCCTGGACTTCGGCATCATYGTBGAYGG-3') and *czcR1* (5'-GGCCATGGGGTGGAACATYTTNCC-3') was used. The previously described reactions and cycling conditions

¹<http://migale.jouy.inra.fr>

(Mahamoud Ahmed et al., 2020) were applied to both genes. To construct standard plasmids, *czcA* and *cusA* fragments were amplified from *Burkholderia* sp. S911 (accession no. KP081480) and *Cupriavidus metallidurans* CH34, respectively, and cloned into the pGEM-T vector. Gene copy numbers were established using standard curves prepared with 10-fold serial dilutions of standard plasmids. All runs were performed in an iCycler iQ system (Bio-Rad) associated with iCycler Optical System Interface software (version 2.3, Bio-Rad).

The abundance of selected antibiotic resistance gene targets (*sul1*, *strA*, *strB*, *ermB*, *ermF*, *aadA*, and *bla_{OXA-20}*) and the class 1 integron (*intI1*) in sediment DNA was quantified by qPCR with SYBR green or Taqman probe chemistry. All procedures, primers, quality control and standards for quantification were exactly as described in Lau et al. (2020). The identities of the quantified gene targets were verified on the basis of hybridization when using TaqMan chemistry, or melting behavior when using SYBR green. Gene abundances were expressed as a ratio with the total 16S rRNA gene copies.

Data Analyses

All statistical analyses were performed with the R free software (version 3.4.3, R Core Team, 2018). After confirming normality of the residuals (Shapiro-Wilk test; Royston, 1982) and data homoscedasticity (Fligner-Killeen test; Conover et al., 1981), significant differences between sites and dates in the enzymatic activities, metabolic potentials, bacterial abundance, OTU richness and Shannon diversity index and genetic resistance potential were sought by analysis of variance (ANOVA) and further analyzed with a *post hoc* Tukey test. Differences were considered statistically significant when the *p*-value was below 0.05.

For bacteria and archaeal diversity, Bray-Curtis similarities (BC) between samples were calculated, and distances between samples were represented using a Principal Coordinates Analysis (PCoA) with R software (Vegan package). Functional-based community parameters (enzymatic and metabolic activities, metals and antibiotic resistance genes) were analyzed using a Principal Component Analysis (PCA). The significance of the sample groupings on the ordinations was tested using Analysis Of SIMilarity (ANOSIM). The relationship between community composition and functional-based structures (based on PCoA and PCA 2-dimension projections, respectively) were evaluated using a Procrustean analysis (using R software, Vegan package). Procrustean analyses superimpose, scale and rotate one data matrix upon the other until an optimal fit is found (Jackson and Harvey, 1993; Peres-Neto and Jackson, 2001; Heino et al., 2004). Estimated residuals between original values and the derived best fit solution give the *m*² statistic, with a low *m*² statistic indicating a good level of correspondence between data matrices (Paavola et al., 2006). The PROTEST permutation procedure (999 permutations) was used to assess the statistical significance of the Procrustean fit between the two matrices (Peres-Neto and Jackson, 2001; Paavola et al., 2006). Vectors of environmental variable data were fitted onto the procrustean analysis plot using the *envfit* function of the *vegan* package. This function calculates the goodness-of-fit values (*R*²) for environmental

variables onto the procrustean analysis plot and the significance of each correlation was tested based on 999 random permutations (Oksanen et al., 2020).


RESULTS

Environmental Conditions and Sediment Contamination at the Sampling Sites

The main sampling site characteristics are presented in Table 1. Surface sediments were collected at a sampling depth ranging from 22 m (site 32) to 95 m (site 29), and the median grain size varied from 8 μ m (sites 6 and 21) to 35 μ m (site 53). Sediments from the sites 53 and 78 had the highest and lowest organic matter content (13.6 and 2.7%, respectively) and total organic carbon (7.0 and 0.7%, respectively). These two sites were also characterized by lower carbonate content (about 14.4 and 19.7%, respectively) than the six other sites (i.e., 5, 6, 21, 29, 32, and 36) where it varied between 31.2% (site 29) and 48.9% (site 36). Total phosphorus concentrations were 4.5- to 7-fold higher in the site 53 (3900 mg kg⁻¹ dw) than in the seven other sites (659 to 849 mg kg⁻¹ dw). Sediments from the site 53 contained high total concentrations of metals (about 1193 mg kg⁻¹ dw) compared to the seven other ones, where it varied from 170 mg kg⁻¹ dw (site 36) to 322 mg kg⁻¹ dw (site 29). The high total concentration of metals in the site 53 was mainly due to zinc and copper (Zn and Cu, 645 and 281 mg kg⁻¹ dw, respectively; Supplementary Table 1). The sediments from site 78 contained high concentrations of titanium (Ti; 1736 mg kg⁻¹ dw) while for the other sites Ti concentrations ranged between 37 and 405 mg kg⁻¹ dw. The site 53 was also characterized by higher values for the sum of the seven indicator PCB (Σ 7PCBi) (387 μ g Σ 7PCBi kg⁻¹ dw): more than 400-fold higher than those observed in the site 78 (0.9 μ g Σ 7PCBi kg⁻¹ dw) and 36- to 75-fold higher than those observed in the six other sites (5.2–10.5 μ g Σ 7PCBi kg⁻¹ dw) (Supplementary Table 2). Sediments from sites 53 and 36 contained higher concentrations of PAH (25262 μ g kg⁻¹ dw and 13581 μ g kg⁻¹ dw, respectively) compared to the six other ones, where it varied from 774 μ g kg⁻¹ dw (site 78) to 2263 μ g kg⁻¹ dw (site 32) (Supplementary Table 3).

Functional Potential

A set of potential microbial activities involved in the C cycle (β -glu, respiration, and methanogenesis), the N cycle (LAP and denitrification) and the P cycle (Pase) was measured (Figure 2). The results reflected a high spatial heterogeneity between the sampling sites. With the exception of the LAP activity (Figure 2B), sediment microbial communities from the site 53 exhibited the highest functional potential, both in October 2017 and May 2018. Methanogenesis activity (Figure 2F) was only detected in this site and β -glu (Figure 2A), Pase (Figure 2C), aerobic respiration (Figure 2D) and denitrification (Figure 2E) were also significantly higher than in the other sampling sites (*p* < 0.05). Significant but less marked spatial differences were also observed between the seven other sites (i.e., 5, 6, 21, 29, 32, 36, and 78) for the three enzymatic activities, which were



generally the highest in sites 5 and 6 and the lowest in sites 21 and 36 (**Figures 2A–C**). In contrast, no significant difference was observed between these seven sites for both aerobic respiration (**Figure 2D**) and denitrification (**Figure 2E**).

Seasonal variations between the two sampling periods were less pronounced in the four sites that have been sampled in 2017 and 2018 (i.e., 29, 32, 53, and 78). Significant differences ($p < 0.05$) between October 2017 and May 2018 were only observed in site 53 for denitrification (**Figure 2D**) and methanogenesis (**Figure 2E**), with the highest values measured in October 2017.

Microbial Community Structure

The mean bacterial abundance in the sample sediments ranged between 0.55×10^6 to 8.03×10^6 cells g^{-1} dw (**Figure 3A**). A high spatial variability was observed in October 2017, when the bacterial community from site 78 exhibited the highest abundance (about 8.0×10^6 cells g^{-1} dw), which was about twice higher than the one measured in site 53, and threefold higher than those measured in sites 29 and 32. In these four sites, the bacterial abundance was reduced by about two- to fourfold between October 2017 and May 2018. In May 2018, abundances were more stable between the different sites, ranging from 0.55×10^6 at site 36 to 2.80×10^6 cells g^{-1} dw at site 78.

High-throughput amplicon sequencing of the 16S rRNA gene revealed no temporal difference in the number of OTUs affiliated to the archaea and bacteria domains (**Figure 3B**) and the resulting Shannon-Wiener diversity index (**Figure 3C**) between October

2017 and May 2018. Based on these two parameters, communities from site 53, and to a lesser extent, from site 78, exhibited the lowest diversity, whereas these two parameters were relatively similar between the six other sites. According to the sampling date and the sample replicate, the number of OTUs varied from 503 to 626 in site 53, from 947 to 1048 in site 78, and from 1067 to 1311 in the six other sites. The mean Shannon-Wiener diversity index was close to 4.0 (± 0.2) in site 53, 4.9 (± 0.1) in site 78, and between 5.4 and 5.8 in the six other sites.

At the Phylum (or Class for Proteobacteria) level, the microbial community was dominated by members of the Gammaproteobacteria ($25.2 \pm 7.0\%$ of the reads), followed by members of the Bacteroidetes ($19.4 \pm 6.9\%$), Deltaproteobacteria ($13.1 \pm 5.4\%$), Nitrospirae ($9.3 \pm 5.9\%$), and Acidobacteria ($6.8 \pm 1.9\%$) (**Figure 4**). Community composition at site 53 was especially dominated by members of the Bacteroidetes (32%), Deltaproteobacteria (22%) and Gammaproteobacteria (20%), whereas at site 78, community composition was dominated by members of the Gammaproteobacteria (27%), Cyanobacteria (24%) and Bacteroidetes (23%). Site 5 community composition was characterized by the most important proportion of members of the Gammaproteobacteria (42%). Members of the Nitrospirae were present at low abundances at sites 53 (0.30%), 78 (4.23%) and 5 (8.05%), whereas they ranged between 11% (sites 29 and 32) and 18% (site 36). Among the less abundant groups, Actinobacteria, Chloroflexi, Nitrospirae, and other Archaea and Bacteria did not exhibit any temporal or spatial trends. For the main archaeal Phylum, Crenarchaeota peaked up to 4% at site




FIGURE 3 | Dot plots of the (A) bacterial abundance, (B) observed OTU richness, and (C) Shannon diversity index for the triplicate samples from the 8 studied sites in Lake Geneva in October 2017 (open symbols) and May 2018 (closed symbols). For each sampling site (and date if applicable – October 2017, italic letter and May 2018, roman letter), different letters indicate significant differences between treatments ($P < 0.05$, Tukey test).




FIGURE 4 | Microbial community composition for the triplicate samples from the 8 studied sites in Lake Geneva in October 2017 and May 2018. The 15 more abundant phylum (or class for Proteobacteria) are represented and the remaining phylum (or class for Proteobacteria) are grouped under “Other Archaea” or “Other Bacteria”. Data is presented are relative abundance of the read counts.


36, but remained below 1.5% at other sites, Euryarchaeota were generally below 2% except at site 53 where they represented 7% of the community in October 2017 and 3% in May 2018, and Thaumarchaeota represented less than 0.5% of the community at sites 5 and 53 but between 2 and 10% elsewhere. Finally, Verrucomicrobia were below 1% in every samples but at site 53 where they accounted for 5 and 6% of the community in October 2017 and May 2018, respectively.

Metal and Antibiotic Resistance Potential

Two distinct sets of quantitative PCRs were performed to, respectively, quantify two genes involved in microbial resistance

to metals (*cusA* and *czcA*, **Figures 5A,B**) and eight genes conferring antibiotic resistance (*sul1*, *strA*, *strB*, *ermB*, *ermF*, *aadA*, and *bla_{OXA20}*), and the class 1 integron integrase *intI1* (**Figures 5C–J**).

Both metal resistance genes were detected in every site and the two sampling dates (**Figures 5A,B**). The *cusA* gene had the highest relative abundance with mean *cusA* to 16S rRNA gene relative abundances varying between about 0.49×10^{-3} to 210×10^{-3} depending on the sampling site and date (**Figure 5A**). No significant temporal variation was observed for this gene between the two sampling dates with the exception of sites 29 and 32 where mean values were about twofold higher ($p < 0.05$).



in October 2017 than in May 2018. Spatial differences among the sites varied according to the sampling date. In October 2017, the relative abundance of *cusA* was significantly higher in sites 29 and 32 than in sites 53 and 78 ($p < 0.05$). In May 2018, the highest relative abundance was observed in site 5 ($204 \times 10^{-3} \pm 14 \times 10^{-3}$) with values significantly higher than those recorded in sites 21, 32, 36, 53, and 78 ($p < 0.05$). The mean relative abundances of *czcA* to 16S rRNA varied between 1.42×10^{-3} and 15.3×10^{-3} with relatively high variations among replicates and no significant difference between sampling sites or between sampling dates (Figure 5B).

The eight genes for antibiotic resistance (Figures 5C–J) were less represented than *cusA* and *czcA*. The class 1 integron marker *int1* was detected in all samples with the highest relative abundances with mean values varying between 98.7×10^{-6} and 11.2×10^{-3} (Figure 5F). The genes *sul1*, *strA*, and *aadA* (Figures 5C,D,I) were also detected in all the samples but their highest relative abundance value was lower than 1×10^{-3} . The detection of *strB*, *ermB*, *ermF*, and *bla_{OXA-20}* (Figures 5E,G,H,J) genes was variable according to sampling site, date and sample replicate, with mean relative abundances always lower than 1×10^{-4} . In addition to the difference among the measured genes, the results showed a high spatial heterogeneity among the sampling sites. The general tendency showed that sediment microbial communities from site 53, and to a lesser extent those from site 78, were characterized by higher relative abundances of the eight genes than the communities from the six other sites (i.e., 5, 6, 21, 29, 32, and 36). Indeed, sediment microbial communities collected in site 53 were characterized by the highest relative abundances of seven of the eight measured genes. The only exception was for the gene *ermF*, which was about threefold more represented in site 78 than in site 53 (Figure 5H). This gene was under the detection limit in the other sampling sites. At site 53, the other seven genes relative abundances were about 3- to 28-fold more abundant than in site 78 and about 30- to

2000-fold more abundant than at the 6 other sites. Moreover, no significant difference in the relative abundance of the eight genes conferring antibiotic resistance was observed in those sites (i.e., 5, 6, 21, 29, 32, and 36) (Figures 5C–J). No temporal variation was observed between the two sampling dates with the exception of site 53 where relative abundances were significantly higher in October 2017 than in May 2018 for the genes *sul1* (Figure 5C), *strA* (Figure 5D), *strB* (Figure 5E), *int1* (Figure 5F), and *aadA* (Figure 5I).

Environmental Drivers of Microbial Structure and Functional-Based Parameters

The 2-dimensional representation of the procrustean analysis of 16S rRNA gene-based bacterial community structure (PCoA), the first 2 axes explained 74% of the variance) and functional-based community parameters (PCA, the first 2 axes explained 76% of the variance) (Figure 6) showed that sites 53 and 78 were distinct from each other, and from the six other sites (ANOSIM; $p < 0.0001$ for PCoA, and $p < 0.01$ for PCA), with very limited differences between the two sampling dates. Among these six sites, communities from site 5 were also slightly distinct both according to their structure and their functional parameters. The phylogenetic structure and the functional and resistance pattern of the communities displayed a highly significant level of concordance (Procrustean analysis, residual sum of squares = 0.340; $m12 = 0.812$; $p < 0.001$), as indicated by the relatively limited length of the colored arrows on the 2-dimension projection.

The sediment physico-chemical characteristics that were correlated with the microbial community ordinations (Figure 4 and Supplementary Table 4) were organic toxicants (PCB and PAH), phosphorus concentrations and median grain size, which were higher at site 53, some metals, which were high at sites 78




FIGURE 6 | Two-dimensional representation of the procrustean analysis of 16S rRNA gene-based bacterial community structure (PCoA analysis of Bray-Curtis distances, origin of colored arrows) and functional-based community parameters (PCA analysis of microbial activities, metals and antibiotic resistance gene relative abundances, arrival of colored arrows) for the triplicate samples from the 8 studied sites in Lake Geneva in October 2017 (empty symbols) and May 2018 (plain symbols). Black arrows are environmental variables showing significant correlations with the sample 2-dimensional projections.

(As and Cr) and 53 (sum of Co, Ni, Cu, Zn, Mo, Ag, Cd, Sn, Pb, and Hg concentrations) and carbonate, which was lower at sites 53 and 78 (Table 1). Site 78 microbial community ordination was also significantly correlated to Ti concentration.

DISCUSSION

This study investigated *in situ* the possible influence of the anthropogenic contamination of Lake Geneva surface sediments on benthic microbial communities by assessing spatiotemporal changes in their structure, their functional potential and their genetic potential for resistance to selected metals and antibiotics. Our results provide evidence that the multiple contaminants in sediments of Lake Geneva alter the diversity of microbial communities with consequences for the important ecological functions they provide. The magnitude of these effects depends on the types and the concentrations of contaminants in the sediments. All contaminant concentrations measured in the present study are in agreement with previous findings (Haller et al., 2011; Gascon Diez et al., 2014; Masson and Tercier-Waeber, 2014; Gascón Díez et al., 2017; Loizeau et al., 2017). In addition, our results also revealed high concentrations of PAH ($>13.5 \mu\text{g kg}^{-1} \text{ dw}$) in sediments collected at site 36. Such a contamination was not reported in Loizeau et al. (2017) because organic toxicants (PAH and PCB) were not analyzed in this study site (and also not in sites 5 and 21). Our results suggest that the multi-contamination (PAH, PCB, metals, phosphorus, organic matter, and possibly microbial and genetic material) of

sediments observed in the Vidy Bay (site 53) is a strong driver of sediment microbial community structure and diversity, the latter being significantly lower than that at other sites. Some phyla were significantly less represented in the Vidy Bay, such as Planctomycetes, the *Nitrospira* genus from the phylum of Nitrospirae, whereas Euryarchaeota were mainly detected in this highly contaminated site. This archaeal phylum includes both the archaeal sulfide reducers as well as the methanogens (Chrencik and Marsh, 2012). Potential methanogenesis activity was only detected at this site, consistent with the observed abundance of methanogenic archaea, belonging to the *Methanobacterium*, *Methanolinea*, *Methanoregula*, *Methanosaeta*, *Methanosarcina* genera and to the *Methanomassiliicoccaceae* family.

In the present study, the Verrucomicrobia phylum was also more represented in the Vidy Bay than in the other sites, in contrast to previous findings (Haller et al., 2011). However, this is in line with other observations where this phylum was more abundant in very urbanized rivers than in suburban rivers (Lin et al., 2019). The urban contamination is also probably responsible for the observed higher relative abundance of members of the Bacteroidetes phylum, which includes bacteria from the human and animal gastrointestinal tract, frequently detected in wastewater, and generally considered as fecal indicators in the environment (Gomez-Donate et al., 2016; Su et al., 2017; Niestępski et al., 2020). In the Vidy Bay, fecal indicator bacteria and other activated sludge bacteria from local WWTP inputs were previously detected (Haller et al., 2009; Sauvain et al., 2014). Bacteroidetes are known to harbor a large variety of metal- or antibiotic-resistance genes that can favor their establishment and development in contaminated sediments (Chen et al., 2018; Niestępski et al., 2020). The higher abundance of gene markers associated with resistance to various classes of antibiotics (*sulI*, *strA*, *strB*, *ermB*, *ermF*, and *bla_{OXA-20}*) and of the class 1 integron (*intI1*) detected in the Vidy Bay sediment is in accordance with the high occurrence of multiresistant bacteria and antibiotic-resistance genes previously reported at this site (Czekalski et al., 2012, 2014). Reporting in 2011 a 200-fold increase of the relative abundances of six antibiotic-resistance genes (including *sulI*) as compared to sediment sampled in the center of the lake, Czekalski et al. (2014) hypothesized that the increasing abundance of antibiotic-resistance genes was mainly due to direct inputs from wastewater effluents rather than *in situ* development of antibiotic-resistant bacteria in the lake sediment following their chronic exposure to contaminants. In the present study, given the high metal concentrations observed in the surface sediment from the Vidy Bay, the hypothesis of co-selection of antibiotic resistance by this kind of contaminant cannot be excluded (Poole, 2017; Silva et al., 2021). Although pharmaceutical residues in the collected sediments were not analyzed it is more than probable that site 53 sediment were contaminated by antibiotics (Morasch et al., 2010; Bonvin et al., 2011, 2013), potentially favoring the development of antibiotic resistance in sediment communities in response to antibiotic selection pressure (Bengtsson-Palme and Larsson, 2016). In contrast, the analysis of the metal-resistance genes *cusA* (involved in copper resistance) and *czcA* (involved in cobalt-zinc-cadmium resistance) did not show higher relative

abundances in the Vidy Bay, even if this site had the highest concentrations of Cu (about 281 mg kg⁻¹ dw), Zn (about 645 mg kg⁻¹ dw) and Cd (about 1.5 mg kg⁻¹ dw), which were 3- to 12-fold higher than the contamination with these three metals in the seven other stations. Significant increases in the relative abundance of the *cusA* and *czcA* genes in contaminated sediments compared to uncontaminated ones has been previously reported by Besaury et al. (2013) and Roosa et al. (2014). In both cases, however, the level of metal contamination was very high (total Cu > 1400 mg kg⁻¹, Besaury et al., 2013; total Zn > 3000 mg kg⁻¹ and Cd > 35 mg kg⁻¹, Roosa et al., 2014), as was the potential metal bioavailability. Altogether, these results questions whether the qPCR quantification of these metal resistance genes is sensitive enough when contamination gradients are weak, or whether metal bioavailability in the Vidy Bay is poor and causes limited selection pressure.

More surprisingly, a notable and sometimes significant increase in the relative abundance of antibiotic-resistance genes was observed in the site 78 in comparison to the other sites other than the Vidy Bay one. This was especially the case for the *ermB* and *ermF* genes, responsible for erythromycin resistance, which were only (*ermF*) or mainly (*ermB*) detected in site 53 and 78. It cannot be excluded that the surface sediment from the site 78 has been directly contaminated by antibiotic-resistant bacteria and/or antibiotics coming from the upper section of the Rhône River, which has been shown to be an important source of contamination of Lake Geneva with pharmaceutical substances (Loizeau et al., 2013). In this site, benthic microbial richness and diversity index was impacted, being 10 to 20% lower than those observed in the other sites (excluding 53) and community composition was also affected. The relative abundance of Bacteroidetes members could also confirm inputs from sewage water located in the upper Rhône section (see related discussion above for site 53). Cyanobacteria were also significantly more represented at site 78 than the other sites, showing that this kind of microorganisms was particularly adapted to the local site characteristics. Given the water column height at this site and associated limited light penetration, it is unlikely that these microorganisms are part of a microbial mat developing at the sediment – water interface, but nutrients flowing in from the Rhône River might promote their planktonic growth following which they are deposited into sediment. Previous studies (Loizeau et al., 2012; Silva et al., 2019) found that sediment accumulation was 5- to 10-fold more rapid at this site compared to other areas of the lake.

Despite their geographical distance and their notable differences in depth and other physico-chemical characteristics, such as total organic carbon or grain size, the sediment communities collected in the less contaminated sites (i.e., 5, 6, 21, 29, and 32) and at site 36, only contaminated by high concentrations of PAH, exhibited relatively comparable structural and functional properties. All these sampling sites (as well as sites 53 and 78) had a depth below 100 m out of the 309 m of maximal depth for Lake Geneva. Extrapolation of the probe sensor data collected for the environmental survey of Lake Geneva at the deepest site indicated that water temperature at the sediment interface varied between 6 and 14°C in October

2017 and was about 6–7°C in Mai 2018, and that dissolved oxygen concentrations increased from between 6 and 8 mg L⁻¹ in October 2017 and between 9 and 12 mg L⁻¹ in May 2018 (Rimet et al., 2020), because of the first 201 m water column partial turnover on March 6, 2018 (CIPEL, 2019). Even if this data should be interpreted with care since it might not perfectly reflect the local conditions prevailing at sampling sites, whatever the measured parameter, no significant difference was observed between the sediment microbial communities collected in the shallowest (22 m; site 32) and deepest (95 m, site 29) sites. As revealed by the multivariate analysis (Figure 6), it confirms that depth, and indirectly related environmental parameters such as dissolved oxygen, light intensity, temperature, and nutrient availability (Böer et al., 2009), was not a major driver of benthic microbial community structure and functions in Lake Geneva, in contrast to other studies that linked microbial community structure changes with water depth (Edlund et al., 2006; Hewson et al., 2007; Zhang et al., 2014; Ding et al., 2015; Ruuskanen et al., 2018; Wu et al., 2019). The only observed differences between sites 5, 6, 21, 29, 32, and 36 concerned the enzymatic activities (β -glu, LAP, and Pase), which were higher at site 5, and to a lesser extent, at site 6. This difference could be explained, at least partly, by a higher amount of available substrates, as suggested by the higher organic matter and total organic carbon contents measured in these samples. Based on these results, it can be argued that the contamination of the sediment with high concentrations of PAH at site 36 (>13.5 mg Σ PAH kg⁻¹ dw) exerts no or very limited effects on the functional potential, the structure and the composition of the microbial communities, which were almost similar to those from sites characterized by 6- to 12-fold lower PAH concentrations. The limited effects of chronic PAH contamination on sediment microbial community diversity and functions has likewise been observed in coastal ecosystems (Hernandez-Raquet et al., 2006; Païssé et al., 2008; Zhang et al., 2008; Jeanbille et al., 2016), in contrast to the more important influence of other contaminants such as metals (Sun et al., 2012, 2013). The small effect of PAHs could be due to their limited toxicity (Païssé et al., 2010) and/or their limited bioavailability because of their ready sorption to sediments (Kraaij et al., 2002; Ghosh et al., 2003).

Despite the evident negative effects of the anthropogenic contamination on the diversity of sediment microbial communities of sites 53 and 78, no inhibition in their functional potential was observed. On the contrary, communities from site 53 exhibited the highest potential for most of the measured activities (i.e., β -glu, Pase, aerobic respiration, denitrification and methanogenesis). It suggests that the observed shifts in their community composition reflects a long-term selection of benthic microbial species that are strongly adapted to their environment, allowing them to benefit from the regular inputs of nutrients and organic substrates that can be used as energy sources, thus making them key players in benthic biogeochemical cycles (Huang et al., 2019; Steinsberger et al., 2019). This long-term adaptation hypothesis can be supported by the very limited temporal variations that were observed in the present study between the two sampling dates. Indeed, at the four sites sampled twice (i.e., 29, 32, 53, and 78), sediment community diversity and

composition were very similar in October 2017 and May 2018, as illustrated in the multivariate analysis (Figure 6). It suggests, that, the structural characteristics of the surface sediment microbial communities of Lake Geneva are relatively stable over time, unlike the case for shallower lakes (Peng et al., 2017; Wan et al., 2017; Zhang et al., 2019).

CONCLUSION

Using a whole lake approach, the present study highlighted that surface sediment microbial communities were impacted by anthropogenic load of contaminants including toxicants, organic matter, nutrients, and microorganisms in wastewater discharge affecting their diversity, enzymatic and metabolic functional potential as well as their genetic potential resistance to antimicrobial agents. These forcing factors appeared to yield to a more important structuration of the microbial parameters studied than the classical environmental factors generally controlling microbial diversity in lakes.

Given the ecological role of microbial communities in lake sediments, further research to better understand the consequences of anthropogenic contamination on microbial communities should focus on larger scale approaches, including more sites to cover a finer gradient of contamination in order to distinguish the relative importance of the different type of contaminants, integrating the full range of water depths from the littoral to the deeper areas, and considering a more important depth of the sediment compartment to take into account the full range of metabolic activities carried out. Together this would improve the ecotoxicological assessment of sediments in lake ecosystems to understand the consequences of multiple stress at the community level on microbial key activities and functions.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in the NCBI at <https://www.ncbi.nlm.nih.gov/sra/PRJNA742879>.

REFERENCES

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403–410. doi: 10.1016/S0022-2836(05)80360-2
- Bedard, D. L. (2008). A case study for microbial biodegradation: anaerobic bacterial reductive dechlorination of polychlorinated biphenyls – from sediment to defined medium. *Annu. Rev. Microbiol.* 62, 253–270. doi: 10.1146/annurev.micro.62.081307.162733
- Bengtsson-Palme, J., and Larsson, D. G. (2016). Concentrations of antibiotics predicted to select for resistant bacteria: proposed limits for environmental regulation. *Environ. Int.* 86, 140–149. doi: 10.1016/j.envint.2015.10.015
- Besaury, L., Bodilis, J., Delgas, F., Andrade, S., De la Iglesia, R., Ouddane, B., et al. (2013). Abundance and diversity of copper resistance genes *cusA* and *copA* in microbial communities in relation to the impact of copper on Chilean marine sediments. *Mar. Pollut. Bull.* 67, 16–25. doi: 10.1016/j.marpolbul.2012.12.007

AUTHOR CONTRIBUTIONS

BF and SP conceived and designed the study. EL, J-LL, BF, and SP organized and performed the sampling. EL, CB, PB, AT, and ET were responsible for the microbial analyses. J-LL, EN, and BF were responsible for the physico-chemical analyses. EL conducted the bioinformatic and biostatistical analyses. EL and SP drafted the first version of the manuscript. All co-authors analyzed and interpreted the data, contributed to subsequent revisions to the manuscript, and approved its final submitted version.

FUNDING

This study was partly funded by a grant of the Auvergne-Rhône-Alpes Region (CMIRA COOPERA 2016–2018 Project) and by the European Cross-Border Cooperation Program (Interreg France-Switzerland 2014–2020, SYNAQUA).

ACKNOWLEDGMENTS

We are grateful to all the staff who have contributed to the sampling campaigns: the boat skippers, Philippe Arpagaus and Jean-Christophe Hustache, and the colleagues who collected and prepared the samples, Anaïs Charton, Christina Lüthi, Bernard Motte, Amélie Roinat, Régis Vivien, and Sandrine Vix. We thank all the persons who realized microbial and physico-chemical analyses: Anaïs Charton, Bernard Motte, Christophe Rosy, Bernadette Volat, Bettina Wagner, and Yuan-Ching Tien. We also thank OLA-IS, AnaEE-France, INRAE of Thonon-les-Bains, CIPEL for providing probe data at site SHL2.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.738629/full#supplementary-material>

- Böer, S. I., Hedtkamp, S. I. C., Van Beusekom, J. E. E., Fuhrman, J. A., Boetius, A., and Ramette, A. (2009). Time- and sediment depth-related variations in bacterial diversity and community structure in subtidal sands. *ISME J.* 3, 780–791. doi: 10.1038/ismej.2009.29
- Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120. doi: 10.1093/bioinformatics/btu170
- Bonvin, F., Razmi, A. M., Barry, D. A., and Kohn, T. (2013). A coupled hydrodynamic-photolysis model to simulate the fate of micropollutants in Vidy Bay. *Environ. Sci. Technol.* 47, 9207–9216. doi: 10.1021/es401294c
- Bonvin, F., Rutler, R., Chèvre, N., Halder, J., and Kohn, T. (2011). Spatial and temporal presence of a wastewater-derived micropollutant plume in Lake Geneva. *Environ. Sci. Technol.* 45, 4702–4709. doi: 10.1021/es2003588
- Burrus, D., Thomas, R. L., Dominik, B., Vernet, J.-P., and Dominik, J. (1990). Characteristics of suspended sediment in the upper Rhone River, Switzerland, including the particulate forms of phosphorus. *Hydrol. Process.* 4, 85–92. doi: 10.1002/hyp.3360040108

- Cébron, A., Norini, M. P., Beguistain, T., and Leyval, C. (2008). Real-Time PCR quantification of PAH-ring hydroxylating dioxygenase (PAH-RHD α) genes from Gram positive and Gram negative bacteria in soil and sediment samples. *J. Microbiol. Methods* 73, 148–159. doi: 10.1016/j.mimet.2008.01.009
- Chen, Y., Jiang, Y., Huang, H., and Mou, L. (2018). Long-term and high-concentration heavy-metal contamination strongly influences the microbiome and functional genes in Yellow River sediments. *Sci. Total Environ.* 637–638, 1400–1412. doi: 10.1016/j.scitotenv.2018.05.109
- Chrencik, B. J., and Marsh, T. L. (2012). “Contributions of methanogenic Archaeobacteria in community-driven anaerobic chromate reduction by Yellowstone National Park hot spring microorganisms,” in *Microbes in Applied Research: Current Advances and Challenges*, ed. A. Méndez-Vilas (Singapore: World Scientific), 60–64. doi: 10.1142/9789814405041_0012
- CIPEL (2018). *Rapports sur les études et Recherches Entreprises dans le Bassin Lémanique, Campagne 2017*. Nyon: Commission Internationale Pour la Protection des Eaux du Léman, 196.
- CIPEL (2019). *Rapports sur les études et Recherches Entreprises dans le Bassin Lémanique, Campagne 2018*. Nyon: Commission internationale pour la protection des eaux du Léman, 305.
- Codling, G., Sturchio, N. C., Rockne, K. J., Li, A., Peng, H., Tse, T. J., et al. (2018). Spatial and temporal trends in poly- and per-fluorinated compounds in the Laurentian Great Lakes Erie, Ontario and St. Clair. *Environ. Pollut.* 237, 396–405. doi: 10.1016/j.envpol.2018.02.013
- Conover, W. J., Johnson, M. E., and Johnson, M. M. (1981). A comparative study of tests for homogeneity of variances, with applications to the outer continental shelf bidding data. *Am. Stat.* 35, 351–361. doi: 10.2307/1268225
- Czekalski, N., Berthold, T., Caucci, S., Egli, A., and Bürgmann, H. (2012). Increased levels of multiresistant bacteria and resistance genes after wastewater treatment and their dissemination into Lake Geneva, Switzerland. *Front. Microbiol.* 3:106. doi: 10.3389/fmicb.2012.00106
- Czekalski, N., Gascón Díez, E., and Bürgmann, H. (2014). Wastewater as a point source of antibiotic-resistance genes in the sediment of a freshwater lake. *ISME J.* 8, 1381–1390. doi: 10.1038/ismej.2014.8
- Dean, W. E. (1974). Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition; comparison with other methods. *J. Sediment. Res.* 44, 242–248. doi: 10.1306/74D729D2-2B21-11D7-8648000102C1865D
- Diepens, N. J., Dimitrov, M. R., Koelmans, A. A., and Smidt, H. (2015). Molecular assessment of bacterial community dynamics and functional end points during sediment bioaccumulation tests. *Environ. Sci. Technol.* 49, 13586–13595. doi: 10.1021/acs.est.5b02992
- Ding, X., Peng, X. J., Jin, B. S., Xiao, M., Chen, J. K., Li, B., et al. (2015). Spatial distribution of bacterial communities driven by multiple environmental factors in a beach wetland of the largest freshwater lake in China. *Front. Microbiol.* 6:129. doi: 10.3389/fmicb.2015.00129
- Edlund, A., Soule, T., Sjöling, S., and Jansson, J. K. (2006). Microbial community structure in polluted Baltic Sea sediments. *Environ. Microbiol.* 8, 223–232. doi: 10.1111/j.1462-2920.2005.00887.x
- Escudé, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., et al. (2018). FROGS: find, rapidly, OTUs with Galaxy solution. *Bioinformatics* 34, 1287–1294. doi: 10.1093/bioinformatics/btx791
- Foulquier, A., Volat, B., Neyra, M., Bornette, G., and Montuelle, B. (2013). Long-term impact of hydrological regime on structure and functions of microbial communities in riverine wetland sediments. *FEMS Microbiol. Ecol.* 85, 211–226. doi: 10.1111/1574-6941.12112
- Frossard, A., Gerull, L., Mutz, M., and Gessner, M. O. (2012). Disconnect of microbial structure and function: enzyme activities and bacterial communities in nascent stream corridors. *ISME J.* 6, 680–691. doi: 10.1038/ismej.2011.134
- Gascon Díez, E., Bravo, A. G., à Porta, N., Masson, M., Graham, N. D., Stoll, S., et al. (2014). Influence of a wastewater treatment plant on mercury contamination and sediment characteristics in Vidy Bay (Lake Geneva, Switzerland). *Aquat. Sci.* 76, 21–32. doi: 10.1007/s00027-013-0325-4
- Gascón Díez, E., Corella, J. P., Adatte, T., Thevenon, F., and Loizeau, J.-L. (2017). High-resolution reconstruction of the 20th century history of trace metals, major elements, and organic matter in sediments in a contaminated area of Lake Geneva, Switzerland. *Appl. Geochem.* 78, 1–11. doi: 10.1016/j.apgeochem.2016.12.007
- Ghosh, U., Zimmerman, J. R., and Luthy, R. G. (2003). PCB and PAH speciation among particle types in contaminated harbor sediments and effects on PAH bioavailability. *Environ. Sci. Technol.* 37, 2209–2217. doi: 10.1021/es020833k
- Golovko, O., Rehrl, A. L., Köhler, S., and Ahrens, L. (2020). Organic micropollutants in water and sediment from Lake Mälaren, Sweden. *Chemosphere* 258:127293. doi: 10.1016/j.chemosphere.2020.127293
- Gomez-Donate, M., Casanovas-Massana, A., Muniesa, M., and Blanch, A. R. (2016). Development of new host-specific *Bacteroides* qPCRs for the identification of fecal contamination sources in water. *Microbiologyopen* 5, 83–94. doi: 10.1002/mbo3.313
- Grandjean, D., De Alencastro, L. F., and Tarradellas, J. (1990). Micropollutants organiques dans les effluents des stations d'épuration, les eaux du Rhône et les eaux du Léman. *Rapp. Comm. int. prot. eaux Léman contre pollut. Campagne* 1989, 199–212.
- Haglund, A. L., Lantz, P., Törnblom, E., and Tranvik, L. (2003). Depth distribution of active bacteria and bacterial activity in lake sediment. *FEMS Microbiol. Ecol.* 46, 31–38. doi: 10.1016/S0168-6496(03)00190-9
- Haller, L., Poté, J., Loizeau, J.-L., and Wildi, W. (2009). Distribution and survival of faecal indicator bacteria in the sediments of the Bay of Vidy, Lake Geneva, Switzerland. *Ecol. Indic.* 9, 540–547. doi: 10.1016/j.ecolind.2008.08.001
- Haller, L., Tonolla, M., Zopfi, J., Peduzzi, R., Wildi, W., and Poté, J. (2011). Composition of bacterial and archaeal communities in freshwater sediments with different contamination levels (Lake Geneva, Switzerland). *Water Res.* 45, 1213–1228. doi: 10.1016/j.watres.2010.11.018
- Harwood, J. E., Van Steenderen, R. A., and Kühn, A. L. (1969). A comparison of some methods for total phosphate analyses. *Water Res.* 3, 425–432. doi: 10.1016/0043-1354(69)90004-9
- Heino, J., Louhi, P., and Muotka, T. (2004). Identifying the scales of variability in stream macroinvertebrate abundance, functional composition and assemblage structure. *Freshw. Biol.* 49, 1230–1239. doi: 10.1111/j.1365-2427.2004.01259.x
- Hernandez-Raquet, G., Budzinski, H., Caumette, P., Dabert, P., Le Ménach, K., Muyzer, G., et al. (2006). Molecular diversity studies of bacterial communities of oil polluted microbial mats from the Etang de Berre (France). *FEMS Microbiol. Ecol.* 58, 550–562. doi: 10.1111/j.1574-6941.2006.00187.x
- Hewson, I., Jacobson-Meyers, M. E., and Fuhrman, J. A. (2007). Diversity and biogeography of bacterial assemblages in surface sediments across the San Pedro Basin, Southern California Borderlands. *Environ. Microbiol.* 9, 923–933. doi: 10.1111/j.1462-2920.2006.01214.x
- Hoerger, C. C., Akhtman, Y., Martelletti, L., Rutler, R., Bonvin, F., Grange, A., et al. (2014). Spatial extent and ecotoxicological risk assessment of a micropollutant-contaminated wastewater plume in Lake Geneva. *Aquat. Sci.* 76, 7–19. doi: 10.1007/s00027-013-0315-6
- Huang, C., Chen, Z., Gao, Y., Luo, Y., Huang, T., Zhu, A., et al. (2019). Enhanced mineralization of sedimentary organic carbon induced by excess carbon from phytoplankton in a eutrophic plateau lake. *J. Soils Sediments* 19, 2613–2623. doi: 10.1007/s11368-019-02261-2
- Jackson, D., and Harvey, H. (1993). Fish and benthic invertebrates: community concordance and community–environment relationships. *Can. J. Fish. Aquat. Sci.* 50, 2641–2651. doi: 10.1139/f93-287
- Jeanbille, M., Gury, J., Duran, R., Tronczynski, J., Ghiglione, J. F., Agogue, H., et al. (2016). Chronic polyaromatic hydrocarbon (PAH) contamination is a marginal driver for community diversity and prokaryotic predicted functioning in coastal sediments. *Front. Microbiol.* 7:1303. doi: 10.3389/fmicb.2016.01303
- Jin, X., Ma, Y., Kong, Z., Kou, W., and Wu, L. (2019). The variation of sediment bacterial community in response to anthropogenic disturbances of Poyang Lake, China. *Wetlands* 39, 63–73. doi: 10.1007/s13157-017-0909-1
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D. (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Appl. Environ. Microbiol.* 79, 5112–5120. doi: 10.1128/AEM.01043-13
- Kraaij, R., Seinen, W., Tolls, J., Cornelissen, G., and Belfroid, A. C. (2002). Direct evidence of sequestration in sediments affecting the bioavailability of hydrophobic organic chemicals to benthic deposit-feeders. *Environ. Sci. Technol.* 36, 3525–3529. doi: 10.1021/es0102787
- Krzmarzick, M. J., McNamara, P. J., Crary, B. B., and Novak, P. J. (2013). Abundance and diversity of organohalide-respiring bacteria in lake sediments across a geographical sulfur gradient. *FEMS Microbiol. Ecol.* 84, 248–258. doi: 10.1111/1574-6941.12059

- Larras, F., Rimet, F., Gregorio, V., Bérard, A., Leboulanger, C., Montuelle, B., et al. (2016). Pollution-induced community tolerance (PICT) as a tool for monitoring Lake Geneva long-term in situ ecotoxic restoration from herbicide contamination. *Environ. Sci. Pollut. Res.* 23, 4301–4311. doi: 10.1007/s11356-015-5302-0
- Lau, C. H., Tien, Y. C., Stedtfeld, R. D., and Topp, E. (2020). Impacts of multi-year field exposure of agricultural soil to macrolide antibiotics on the abundance of antibiotic resistance genes and selected mobile genetic elements. *Sci. Total Environ.* 727:138520. doi: 10.1016/j.scitotenv.2020.138520
- Lécrivain, N., Aurenche, V., Cottin, N., Frossard, V., and Clément, B. (2018). Multi-contamination (heavy metals, polychlorinated biphenyls and polycyclic aromatic hydrocarbons) of littoral sediments and the associated ecological risk assessment in a large lake in France (Lake Bourget). *Sci. Total Environ.* 619–620, 854–865. doi: 10.1016/j.scitotenv.2017.11.151
- Lécrivain, N., Frossard, V., Naffrechoux, E., and Clément, B. (2019). Looking at organic pollutants (OPs) signatures in littoral sediments to assess the influence of a local urban source at the whole lake scale. *Polycycl. Aromat. Compd.* 41, 950–962. doi: 10.1080/10406638.2019.1631195
- Lin, X., Gao, D., Lu, K., and Li, X. (2019). Bacterial community shifts driven by nitrogen pollution in river sediments of a highly urbanized city. *Int. J. Environ. Res. Public Health* 16:3794. doi: 10.3390/ijerph16203794
- Loizeau, J.-L., Arbouille, D., Santiago, S., and Vernet, J.-P. (1994). Evaluation of a wide-range laser diffraction grain-size analyzer for use with sediments. *Sedimentology* 41, 353–361. doi: 10.1111/j.1365-3091.1994.tb01410.x
- Loizeau, J.-L., Edder, P., De Alencastro, L. F., Corvi, C., and Ramseier Gentile, S. (2013). Lake Geneva pollution by micropollutants – a review of 40 years of study [La contamination du Léman par les micropolluants Revue de 40 ans d'études]. *Arch. Sci.* 66, 117–136.
- Loizeau, J.-L., Girardclos, S., and Dominik, J. (2012). Taux d'accumulation de sédiments récents et bilan de la matière particulaire dans le Léman (Suisse-France). *Arch. Sci.* 65, 81–92.
- Loizeau, J. L., Makri, S., Arpagaus, P., Ferrari, B., Casado-Martinez, C., Benejam, T., et al. (2017). Micropolluants métalliques et organiques dans les sédiments superficiel du Léman. Metallic and organic micropollutants in the surface sediments of Lake Geneva. *Rapp. Comm. Int. Prot. Eaux Léman contre Pollut. Campagne* 2016, 153–207. pdf.
- Magoë, T., and Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27, 2957–2963. doi: 10.1093/bioinformatics/btr507
- Mahamoud Ahmed, A., Lyautey, E., Bonnineau, C., Dabrin, A., and Pesce, S. (2018). Environmental concentrations of copper, alone or in mixture with arsenic, can impact river sediment microbial community structure and functions. *Front. Microbiol.* 9:1852. doi: 10.3389/fmicb.2018.01852
- Mahamoud Ahmed, A., Tardy, V., Bonnineau, C., Billard, P., Pesce, S., and Lyautey, E. (2020). Changes in sediment microbial diversity following chronic copper-exposure induce community copper-tolerance without increasing sensitivity to arsenic. *J. Hazard. Mater.* 391:122197. doi: 10.1016/j.jhazmat.2020.122197
- Mahé, F., Rognes, T., Quince, C., de Vargas, C., and Dunthorn, M. (2015). Swarm v2: highly scalable and high-resolution amplicon clustering. *Peer J.* 3:e1420. doi: 10.7717/peerj.1420
- Marvin, C., Painter, S., Williams, D., Richardson, V., Rossmann, R., and Van Hoof, P. (2004). Spatial and temporal trends in surface water and sediment contamination in the Laurentian Great Lakes. *Environ. Pollut.* 129, 131–144. doi: 10.1016/j.envpol.2003.09.029
- Masson, M., and Tercier-Waeber, M.-L. (2014). Trace metal speciation at the sediment-water interface of Vidy Bay: influence of contrasting sediment characteristics. *Aquat. Sci.* 76, 47–58. doi: 10.1007/s00027-013-0323-6
- Morasch, B., Bonvin, F., Reiser, H., Grandjean, D., de Alencastro, L. F., Perazzolo, C., et al. (2010). Occurrence and fate of micropollutants in the Vidy Bay of Lake Geneva, Switzerland. Part II: micropollutant removal between wastewater and raw drinking water. *Environ. Toxicol. Chem.* 29, 1658–1668. doi: 10.1002/etc.222
- Murphy, J., and Riley, J. P. (1962). A modified single solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta* 27, 31–36. doi: 10.1016/S0003-2670(00)88444-5
- Mwanamoki, P., Devarajan, N., Thevenon, F., Birane, N., De Alencastro, L., Grandjean, D., et al. (2014). Trace metals and persistent organic pollutants in sediments from river-reservoir systems in Democratic Republic of Congo (DRC): spatial distribution and potential ecotoxicological effects. *Chemosphere* 111, 485–492. doi: 10.1016/j.chemosphere.2014.04.083
- Niestępski, S., Harnisz, M., Ciesielski, S., Korzeniewska, E., and Osińska, A. (2020). Environmental fate of Bacteroidetes, with particular emphasis on *Bacteroides fragilis* group bacteria and their specific antibiotic resistance genes, in activated sludge wastewater treatment plants. *J. Hazard. Mater.* 394:122544. doi: 10.1016/j.jhazmat.2020.122544
- Office Fédéral de l'environnement [OFEV] (1998). *Ordonnance du 1er Juillet 1998 sur les Atteintes Portées aux Sols (OSol)*. N° RS 814.12. Berne: OFEV.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., McGlin, D., Minchin, P. R., et al. (2020). *vegan: Community Ecology Package*. R Package Version 2.5-7. Available online at: <https://CRAN.R-project.org/package=vegan>
- Paavola, R., Muotka, T., Virtanen, R., Heino, J., Jackson, D., and Mäki-Petäys, A. (2006). Spatial scale affects community concordance among fishes, benthic macroinvertebrates, and bryophytes in streams. *Ecol. Appl.* 16, 368–379. doi: 10.1890/03-5410
- Paissé, S., Coulon, F., Goñi-Urriza, M., Peperzak, L., McGenity, T. J., and Duran, R. (2008). Structure of bacterial communities along a hydrocarbon contamination gradient in a coastal sediment. *FEMS Microbiol. Ecol.* 66, 295–305. doi: 10.1111/j.1574-6941.2008.00589.x
- Paissé, S., Goñi-Urriza, M., Coulon, F., and Duran, R. (2010). How a bacterial community originating from a contaminated coastal sediment responds to an oil input. *Microb. Ecol.* 60, 394–405. doi: 10.1007/s00248-010-9721-7
- Pearman, J. K., Biessy, L., Thomson-Laing, G., Waters, S., Vandergoes, M. J., Howarth, J. D., et al. (2020). Local factors drive bacterial and microeukaryotic community composition in lake surface sediment collected across an altitudinal gradient. *FEMS Microbiol. Ecol.* 96:fiaa070. doi: 10.1093/femsec/fiaa070
- Peng, Y., Yue, D., Xiao, L., and Qian, X. (2017). Temporal variation and co-occurrence patterns of bacterial communities in eutrophic Lake Taihu, China. *Geomicrobiol. J.* 35, 186–197. doi: 10.1080/01490451.2017.1348406
- Perazzolo, C., Morasch, B., Kohn, T., Magnet, A., Thonney, D., and Chèvre, N. (2010). Occurrence and fate of micropollutants in the Vidy Bay of Lake Geneva, Switzerland. Part I: priority list for environmental risk assessment of pharmaceuticals. *Environ. Toxicol. Chem.* 29, 1649–1657. doi: 10.1002/etc.221
- Peres-Neto, P. R., and Jackson, D. A. (2001). How well do multivariate data sets match? Evaluating the association of multivariate biological data sets: comparing the robustness of Mantel test and a Procrustean superimposition approach. *Oecologia* 129, 169–178. doi: 10.1007/s004420100720
- Pesce, S., Perceval, O., Bonnineau, C., Casado-Martinez, C., Dabrin, A., Lyautey, E., et al. (2018). Looking at biological community level to improve ecotoxicological assessment of freshwater sediments: report on a first French-Swiss workshop. *Environ. Sci. Pollut. Res.* 25, 970–974. doi: 10.1007/s11356-017-0620-z
- Poole, K. (2017). At the nexus of antibiotics and metals: the impact of Cu and Zn on antibiotic activity and resistance. *Trends Microbiol.* 25, 820–832. doi: 10.1016/j.tim.2017.04.010
- Poté, J., Haller, L., Loizeau, J. L., Bravo, A. G., Sastre, V., and Wildi, W. (2008). Effects of a sewage treatment plant outlet pipe extension on the distribution of contaminants in the sediments of the Bay of Vidy, Lake Geneva, Switzerland. *Biores. Technol.* 99, 7122–7131. doi: 10.1016/j.biortech.2007.12.075
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA ribosomal RNA gene database project: improved data processing and webbased tools. *Nucleic Acids Res.* 41, 590–596.
- R Core Team (2018). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Rimet, F., Anneville, O., Barbet, D., Chardon, C., Crépin, L., Domaizon, I., et al. (2020). The Observatory on LAKes (OLA) database: sixty years of environmental data accessible to the public: the Observatory on LAKes (OLA) database. *J. Limnol.* 79, 164–178. doi: 10.4081/jlimnol.2020.1944
- Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. (2016). VSEARCH: a versatile open source tool for metagenomics. *Peer J.* 4:e2584. doi: 10.7717/peerj.2584
- Roosa, S., Wattiez, R., Prygiel, E., Lesven, L., Billon, G., and Gillan, D. C. (2014). Bacterial metal resistance genes and metal bioavailability in contaminated sediments. *Environ. Pollut.* 189, 143–151. doi: 10.1016/j.envpol.2014.02.031
- Roos-Barraclough, F., Givélet, N., Martinez-Cortizas, A., Goodsite, M. E., Biester, H., and Shotyk, W. (2002). An analytical protocol for the determination of total

- mercury concentrations in solid peat samples. *Sci. Total Environ.* 292, 129–139. doi: 10.1016/S0048-9697(02)00035-9
- Royston, J. P. (1982). An extension of Shapiro and Wilk's W test for normality to large samples. *J. R. Stat. Soc. C Appl.* 31, 115–124. doi: 10.2307/2347973
- Ruuskanen, M. O., St Pierre, K. A., St Louis, V. L., Aris-Brosou, S., and Poulain, A. J. (2018). Physicochemical drivers of microbial community structure in sediments of Lake Hazen, Nunavut, Canada. *Front. Microbiol.* 9:1138. doi: 10.3389/fmicb.2018.01138
- Sauvain, L., Bueche, M., Junier, T., Masson, M., Wunderlin, T., Kohler-Milleret, R., et al. (2014). Bacterial communities in trace metal contaminated lake sediments are dominated by endospore-forming bacteria. *Aquat. Sci.* 76, 33–46. doi: 10.1007/s00027-013-0313-8
- Schallenberg, M., and Kalf, J. (1993). The ecology of sediment bacteria in lakes and comparisons with other aquatic ecosystems. *Ecology* 74, 919–934. doi: 10.2307/1940816
- Schultz, P., and Urban, N. R. (2008). Effects of bacterial dynamics on organic matter decomposition and nutrient release from sediments: a modeling study. *Ecol. Modell.* 210, 1–14. doi: 10.1016/j.ecolmodel.2007.06.026
- Schwefel, R., Gaudard, A., Wüest, A., and Bouffard, D. (2016). Effects of climate change on deepwater oxygen and winter mixing in a deep lake (Lake Geneva): comparing observational findings and modeling. *Water Resour. Res.* 52, 8811–8826. doi: 10.1002/2016WR019194
- Sedláček, J., Tolaszová, J., Kříženecká, S., Bábek, O., and Zimová, K. (2020). Regional contamination history revealed in coal-mining-impacted Oxbow Lake sediments. *Water Air Soil Pollut.* 231:208. doi: 10.1007/s11270-020-04583-1
- Silva, I., Tacão, M., and Henriques, I. (2021). Selection of antibiotic resistance by metals in a riverine bacterial community. *Chemosphere* 263:127936. doi: 10.1016/j.chemosphere.2020.127936
- Silva, T. A., Stutenbecker, L., Bakker, M., Costa, A., Schlunegger, F., Lane, S., et al. (2019). The sediment budget and dynamics of a delta-canyon-lobe system over the Anthropocene timescale: the Rhone River Delta, Lake Geneva (Switzerland/France). *Sedimentology* 66, 838–858. doi: 10.1111/sed.12519
- Steinsberger, T., Schmid, M., Wüest, A., Schwefel, R., Wehrli, B., and Müller, B. (2019). Organic carbon mass accumulation rate regulates the flux of reduced substances from the sediments of deep lakes. *Biogeosciences* 14, 3275–3285. doi: 10.5194/bg-14-3275-2017
- Su, J. Q., An, X. L., Li, B., Chen, Q. L., Gillings, M. R., Chen, H., et al. (2017). Metagenomics of urban sewage identifies an extensively shared antibiotic resistome in China. *Microbiome* 5:84. doi: 10.1186/s40168-017-0298-y
- Sun, M. Y., Dafforn, K. A., Brown, M. V., and Johnston, E. L. (2012). Bacterial communities are sensitive indicators of contaminant stress. *Mar. Pollut. Bull.* 64, 1029–1038. doi: 10.1016/j.marpolbul.2012.01.035
- Sun, M. Y., Dafforn, K. A., Johnston, E. L., and Brown, M. V. (2013). Core sediment bacteria drive community response to anthropogenic contamination over multiple environmental gradients. *Environ. Microbiol.* 15, 2517–2531. doi: 10.1111/1462-2920.12133
- Szakova, I., Kolišova, D., Miholova, D., and Mader, P. (2004). Single-purpose atomic absorption spectrometer AMA-254 for mercury determination and its performance in analysis of agricultural and environmental materials. *Chem. Pap.* 58, 311–315.
- Vadeboncoeur, Y., Vander Zanden, M. J., and Lodge, D. M. (2002). Putting the lake back together: reintegrating benthic pathways into lake food web models: lake ecologists tend to focus their research on pelagic energy pathways, but, from algae to fish, benthic organisms form an integral part of lake food webs. *BioScience* 52, 44–54. doi: 10.1641/0006-3568
- Wan, Y., Bai, Y., He, J., Zhang, Y., Li, R., and Ruan, X. (2017). Temporal and spatial variations of aquatic environmental characteristics and sediment bacterial community in five regions of Lake Taihu. *Aquat. Ecol.* 51, 343–358. doi: 10.1007/s10452-017-9621-8
- Wang, Y., and Qian, P. Y. (2009). Conservative fragments in bacterial 16S rRNA genes and primer design for 16S ribosomal DNA amplicons in metagenomic studies. *PLoS One* 4:e7401. doi: 10.1371/journal.pone.0007401
- Widenfalk, A., Bertilsson, S., Sundh, I., and Goedkoop, W. (2008). Effects of pesticides on community composition and activity of sediment microbes—responses at various levels of microbial community organization. *Environ. Pollut.* 152, 576–584. doi: 10.1016/j.envpol.2007.07.003
- Widenfalk, A., Svensson, J. M., and Goedkoop, W. (2004). Effects of the pesticides captan, deltamethrin, isoproturon, and pirimicarb on the microbial community of a freshwater sediment. *Environ. Toxicol. Chem.* 23, 1920–1927. doi: 10.1897/03-345
- Wu, K., Zhao, W., Wang, Q., Yang, X., Zhu, L., Shen, J., et al. (2019). The relative abundance of benthic bacterial phyla along a water-depth gradient in a plateau lake: physical, chemical, and biotic drivers. *Front. Microbiol.* 10:1521. doi: 10.3389/fmicb.2019.01521
- Xu, Y., Wu, Y., Han, J., and Li, P. (2017). The current status of heavy metal in lake sediments from China: pollution and ecological risk assessment. *Ecol. Evol.* 7, 5454–5466. doi: 10.1002/ece3.3124
- Zhang, L., Cheng, Y., Gao, G., and Jiang, J. (2019). Spatial-temporal variation of bacterial communities in sediments in Lake Chaohu, a large, shallow eutrophic lake in China. *Int. J. Environ. Res. Public Health* 16:3966. doi: 10.3390/ijerph16203966
- Zhang, W., Ki, J.-S., and Qian, P.-Y. (2008). Microbial diversity in polluted harbor sediments I: bacterial community assessment based on four clone libraries of 16S rDNA. *Estuar. Coast. Shelf Sci.* 76, 668–681. doi: 10.1016/j.ecss.2007.07.040
- Zhang, X. Z., Xie, J. J., and Sun, F. L. (2014). Effects of three polycyclic aromatic hydrocarbons on sediment bacterial community. *Curr. Microbiol.* 68, 756–762. doi: 10.1007/s00284-014-0535-6

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Lyautey, Bonninaeu, Billard, Loizeau, Naffrechoux, Tlili, Topp, Ferrari and Pesce. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.