

# Enantioselective Au( i )-catalyzed dearomatization of 1-naphthols with allenamides through Tethered Counterion-Directed Catalysis

Yunliang Yu, Zhenhao Zhang, Arnaud Voituriez, Nicolas Rabasso, Gilles Frison, Angela Marinetti, Xavier Guinchard

### ▶ To cite this version:

Yunliang Yu, Zhenhao Zhang, Arnaud Voituriez, Nicolas Rabasso, Gilles Frison, et al.. Enantioselective Au( i )-catalyzed dearomatization of 1-naphthols with allenamides through Tethered Counterion-Directed Catalysis. Chemical Communications, 2021, 57 (82), pp.10779-10782. 10.1039/D1CC04088J. hal-03405405

## HAL Id: hal-03405405 https://hal.science/hal-03405405

Submitted on 5 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Enantioselective Au(I)-Catalyzed Dearomatization of 1-Naphthols with Allenamides through Tethered Counterion-Directed Catalysis

Yunliang Yu,<sup>a,b</sup> Zhenhao Zhang,<sup>a,c</sup> Arnaud Voituriez,<sup>a</sup> Nicolas Rabasso,<sup>b</sup> Gilles Frison,<sup>c,d</sup> Angela Marinetti,<sup>a,\*</sup> Xavier Guinchard<sup>a,\*</sup>

The Tethered Counterion-Directed Catalysis approach has been applied to the enantioselective Au(I) catalyzed dearomatizations of 1-naphthols with allenamides. Stereocontrol is ensured by the intramolecular ion-pairing between the chiral gold-tethered phosphate and an iminium unit, that provides a rigid, well-defined chiral environment to the key electrophilic intermediate.

The impressive potential of gold(I) catalysis for the synthesis of complex molecular scaffolds<sup>1</sup> has largely motivated the design of chiral gold(I) complexes that enable enantioselective variants.<sup>2</sup> Beyond classical approaches based on chiral ligands, the Asymmetric Counterion-Directed Catalysis (ACDC) approach<sup>3</sup> that takes advantage from chiral phosphate counterions as the only source of stereocontrol, proved highly successful in this field, despite being mainly applied to hydrofunctionalizations of unsaturated bonds.<sup>4</sup> As a complement to the ACDC approach, we have disclosed recently<sup>5</sup> that tethering of a phosphate counterion to gold itself via a phosphorus ligand, as in the (CPAPhos<sup>A</sup>)AuCl complex 1 (Scheme 1a), may create additional conformational and steric constraints, that contribute to overcome some of the limitations of ACDC, notably in terms of enantioselectivity. This new method has been named 'Tethered Counterion Directed Catalysis' (TCDC). Thus, in the tandem cycloisomerisation/nucleophilic addition reactions of 2-alkynyl-enones shown in Scheme 1b, (CPAPhos<sup>4</sup>)AuCl 1 provided bicyclic furans with up to 97% ee, at an unusually low catalyst loading (0.2 mol%). This reaction is postulated to proceed via the key carbocationic intermediate I that forms a tight intramolecular ion pair with the tethered phosphate counterion, triggering high enantioselectivity in the addition of the nucleophile.

In this context, with the aim to further expand the scope of the TCDC approach, we have investigated other classes of reactions. Especially, we have postulated that the gold tethered phosphate might form tight ion pairs also with iminium ions that are generated from allenamides under gold activation<sup>6</sup> (*II* in Scheme 1c). Therefore, enantioselective nucleophilic addition reactions involving allenamides might benefit, in terms of stereochemical control, from such intramolecular ion-pairing effects.

a) Bifunctional catalysts (CPAPhos<sup>A</sup>)AuCl (ref 5)



#### Stereochemical control via

b) ion pairing with a remote carbocation (ref 5)



c) ion pairing with iminium ions and application in naphthol dearomatization (this work) OH



Scheme 1: Models for the stereochemical control in reactions involving pre-catalyst 1

In this paper we report preliminary results that validate this working hypothesis, by showing that catalyst **1** gives high enantioselectivity in the catalytic dearomatization of 1-naphthols with allenamides. Generally speaking, catalytic dearomatization has grown as a convenient and reliable strategy for the synthesis of chiral compounds, as it converts flat, achiral derivatives into three dimensional scaffolds decorated by a variety of functional groups<sup>7,8</sup> Naphthols are privileged substrates, since they produce partially dearomatized naphthalenone units often encountered in natural and biologically active products.<sup>9</sup> Despite the extensive development of transition metal catalysts<sup>10</sup> and organocatalysts<sup>11</sup> for these reactions, some substrate combinations remain barely investigated. This is the case notably for the intermolecular dearomatization of naphthols with allenamides that has been reported recently by Shao<sup>11f</sup> and Bandini<sup>10i, 10j</sup> using phosphoric acids and gold phosphates as catalysts, respectively. Bandini's method required highly hindered 3,3'-polyacene-BINOL-based catalysts to achieve high levels of enantioselectivity and was applied only to 2-naphthols. Overall, this state of art that shows only a few sparse reactions involving allenamides, has motivated our attempts to fill this gap by taking advantage of catalyst **1**.

The dearomatization of 2-allylnaphthol **2a** with *N*-tosylallenamide **3a** has been selected as a model catalytic reaction. After a few optimization studies (see the supporting information for full details), the reaction could be carried out in the presence of 5 mol% of (CPAPhos<sup>A</sup>)AuCl **1** as the pre-catalyst (Table 1). Complex **1** was activated *in situ* with Ag<sub>2</sub>CO<sub>3</sub>, in order to remove the chloride atom from gold and concomitantly generate a phosphate counterion. The reaction was carried out at 0 °C in 1,2-dichloroethane for 12h and, gratifyingly, the expected benzocyclohexenone **4a** was produced in good yield (68%) and high enantioselectivity (88% ee, entry 1). The absolute (*R*)-configuration of **4a** has been assigned from literature data.<sup>11f</sup>

Beyond this extremely positive preliminary result, our aim was to substantiate the working hypothesis of the key role of the phosphate-tethered ligand in this catalytic process. Therefore, a few control experiments have been carried out to get more insights on the nature of the catalytically active species. The results are the following:

(a) The non-activated gold complex **1** promoted the reaction with low conversion rate (19% yield) and provided **4a** as a racemic product (entry 2). This demonstrated that the catalytic activity of **1** cannot be due to the phosphoric acid function alone.

(b) CPA1, the gold-free analog of **(S)-1** was catalytically inactive in the presence (entry 3) or absence (entry 4) of silver carbonate. The results obtained with **(S)-1** (entry 1) hence cannot be attributed to silver phosphate nor to an organocatalytic process.

(c) The non-tethered (triphenylphosphine)Au(phosphate) complex **5**, gave **4a** in a very low yield (5%), with a moderate 55% ee, under the same reaction conditions (entry 5). This shows that a non-tethered chiral gold phosphate operating via the ACDC approach does not compete with **1** in terms of both catalytic activity and enantioselectivity, even when the well-known TRIP phosphate is used as the chiral counterion.

(d) The phosphoric acid TRIP  $CPA_2$  promoted the reaction with only moderate yields and ee (38% yield, 57% ee, entry 6).

Overall, these experiments support the synergistic effects between the gold center and the tethered phosphate counterion of **1** that finally produce an effective catalyst.

Based on these encouraging outcomes, we have investigated the scope of this dearomatization reaction promoted by (CPAPhos<sup>A</sup>)AuCl **1**. Results are shown in Scheme 2. The trifluorophenyl substituent of allenamide **3a** could be replaced by a phenyl group, leading to **4b** in 88% ee. The Ts protecting groups of allenamides could then be replaced by either *p*-Ns or *m*-Ns groups, giving **4c**, **4d** and **4e** in 88%, 92% and 91% ee respectively. This set of results expands, to some extent, the scope of the dearomatization reaction, since the established method using chiral phosphoric acids ensures high enantioselectivities only with  $Ts/p-CF_3C_6H_4$  substituted allenamide.

**Table 1:** Dearomatization of 1-naphthol **2a** with allenamide **3a** promoted by activated **1** (entry 1) and control experiments (entries 2-4).

	OH TS PCF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> TS H - I 2a 3a	Catalyst (5 mol%) DCE, 12 h	0 4a	│ N_PCF Ts	<sub>3</sub> -C <sub>6</sub> H <sub>4</sub>
	Catalyst		additiv	yiel	ee
			е	d	
1			Ag <sub>2</sub> CO	68	88
			3	%	%
	O P OH				
2	(S)-1		none	19	0%
				%	
	O II			<b>0</b> (	
3	Ph <sub>2</sub> P		Ag <sub>2</sub> CO	0%	-
			3		
л			<b>no</b> no	~F	
4	O OH		none	<5 %	-
				/0	



A drop of the enantiomeric excess to 68% was observed for  $R^3$  = methallyl (4f), indicating that the reaction is sensitive to the global bulkiness of the substituent  $R^3$ . This was further confirmed by the reaction with 3-*i*-Bu-2naphthol, that resulted in a further decrease of the ee to 62% (4g). On the contrary, naphthols with propyl or methyl groups at the  $R^3$  position maintained a good level of enantioselection, leading to 4h and 4i in 77 and 81% ee, respectively. The reaction with the sterically hindered 3-*t*-Bu-naphthol ( $R^3$ = *t*-Bu) did not proceed, while the reaction with 2-phenyl-1-naphthol led to the *O*-alkylation product in 84% yield (See the supporting information). Naphthols with substituents on their position  $R^4$  were tolerated also: methoxy and ethyl substituted naphthols led to 4j and 4k in 83% and 81% ee respectively. A phenyl substituent in this position however led to a drop of ee to 36% (4l).  $\alpha$ -Amino allenylphosphonates are known to be useful synthons for the synthesis of diverse functionalized molecules.<sup>12</sup> Thus, a range of  $\alpha$ -amino allenylphosphonates have been engaged in the reaction but unfortunately

did not lead to any conversion. Remarkably, none of these reactions afforded products resulting from para addition of allenamides on the naphthols.

For comparison, a few experiments have been carried out on 2-naphthols. Under the same conditions, 1,3dimethyl-2-naphthol delivered naphthalenone **4m** in 75% yield and 65% ee, while 1-methyl-3-benzyl-2-naphthol afforded **4n** in 64% yield and 59% ee. These results indicated that 2-naphthols are suitable substrates in these reactions, but lead to globally lower enantioselectivities.



**Scheme 2:** Scope of the enantioselective dearomatization of 1-and 2-naphthols with allenamides, using pre-catalyst (CPAPhos<sup>4</sup>)AuCl, **1**.

The experimental results in Table 1 and Scheme 2 demonstrate the beneficial effect of tethering the counterion to the gold center, in a catalytic reaction radically different from those disclosed in our previous study (Scheme 1).<sup>5</sup> Thus, to give further rationale to the TCDC concept, we have investigated in more depth the mechanism and stereochemical outcome of this reaction by theoretical methods (see the SI for computational details).

The postulated mechanism (Scheme 3) involves activation of the allenamide by the Au(I) complex **6** to form **7**, association of 1-naphthol to **7** through H-bonding (**8**) and addition of naphthol to the complexed allenamide to produce the  $\sigma$ -complex **9**. Protodeauration will be the final step releasing catalyst **6** and product **4**. Computational studies have been carried out at the DFT level for the key reaction intermediates in Scheme 3.

The most stable conformations of the postulated catalytically active species **6** had been determined in our previous studies.<sup>5</sup> We hence started here by considering the interaction of **6** with allenamide **3b**. The  $\sigma$ -bonded intermediates **7** proved slightly more stable than the corresponding  $\eta^2$ -complexes (see SI, conformers **7a-7y**). A thorough exploration of the possible conformations allowed to identify **7a** as the lowest energy structure (Figure 1). Unexpectedly, **7a** exhibits only a long-range iminium-phosphate intramolecular electrostatic interaction (P-O<sup>---</sup>C=N<sup>+</sup> distance = 4.68 Å). In **7b**, the P-O<sup>---</sup>C=N<sup>+</sup> ion-pair distance is reduced to 3.93 Å but this conformation is higher in energy relative to **7a** (+7.9 kJ/mol) due to the geometrical constraints created by the phosphate group. The formation of a covalent C-O bond as in **7c**, as previously proposed in organocatalytic approaches,<sup>11f</sup> is also energetically demanding.



**Scheme 3:** Putative mechanistic pathways for the enantioselective dearomatization of 1-naphthols with allenamides.



Figure 1: Selected optimized geometries of intermediates 7. Relative Gibbs free energy (in kJ/mol).



**Figure 2:** Optimized geometry of transition states connecting intermediate **8** and **9**. Relative Gibbs free energy (in kJ/mol). For clarity, a wire frame representation of some aryl groups is used.

The most stable conformation 7a allows the 2-methyl-1-naphtol 2d to approach the terminal  $CH_2$  group of the allenamide while enabling its activation through hydrogen bonding with the phosphate group.

The potential energy surface for the addition of 1-naphtol **2d** to **7a** has been computed at the DFT level. The lowest relative energy transition state leads to an (R) configured product **4a**, in agreement with experiments, whereas the (S)-product is obtained through a TS located 9.2 kJ/mol higher in Gibbs free energy (Figure 2). The higher energy of (S)-TS is likely to result from steric hindrance between the naphtol backbone and the phosphoric acid moiety, as revealed by its distortion from planarity (dihedral angle -178.1° vs 173.6°).

Overall, calculations provide an accurate stereochemical model for these reactions. They confirm that the rigidity, bulkiness and three-dimensional arrangement of the intramolecular ion pair **7**, together with hydrogen bonding of 1-naphthol with the phosphate P(O) function, direct the addition of 1-naphthol to the activated allenamide in an enantioselective manner.

In conclusion, we have demonstrated here a second application of the asymmetric Tethered Counterion-Directed Catalysis strategy: the Au(I)-catalyzed dearomatization of 1-naphthols with allenamides. In this reaction, the bifunctional nature of the (CPAPhos<sup>A</sup>)AuCl complex **1** fully reveals its potential, since the phosphate not only contribute to a rigid ion paring within the allenamide-gold adduct, but also directs the addition of the naphthol via H-bonding. Further relevant applications of the TCDC approach will be reported shortly.

We thank the CSC (China Scholarship Council) for PhD grants to Y.Y. and Z.Z..

This paper is dedicated to Prof. Christian Bruneau for his sound contributions to homogeneous catalysis.

#### References

- a) J. Guillem Mayans, H. Armengol-Relats, P. Calleja and A. M. Echavarren, *Isr. J. Chem.*, 2018, 58, 639; b) P.
  Y. Toullec and V. Michelet, *Isr. J. Chem.*, 2018, 58, 578; c) J. L. Mascareñas, I. Varela and F. López, *Acc. Chem. Res.*, 2019, 52, 465.
- a) J.-J. Jiang and M.-K. Wong, *Chem. Asian J.*, 2021, 16, 364; b) W. Zi and D. F. Toste, *Chem. Soc. Rev.*, 2016, 45, 4567.
- 3 a) M. Mahlau and B. List, *Angew. Chem. Int. Ed.*, 2013, **52**, 518; b) K. Brak and E. N. Jacobsen, *Angew. Chem. Int. Ed.*, 2013, **52**, 534.
- a) G. L. Hamilton, E. J. Kang, M. Mba and F. D. Toste, *Science*, 2007, **317**, 496; b) R. LaLonde, Z. Wang, M. Mba, A. Lackner and F. Toste, *Angew. Chem. Int. Ed.*, 2010, **49**, 598; c) W. Zi and F. D. Toste, *Angew. Chem. Int. Ed.*, 2015, **54**, 14447; d) S. Handa, D. J. Lippincott, D. H. Aue and B. H. Lipshutz, *Angew. Chem. Int. Ed.*, 2014, **53**, 10658; e) A. K. Mourad, J. Leutzow and C. Czekelius, *Angew. Chem. Int. Ed.*, 2012, **51**, 11149.
- 5 Z. Zhang, V. Smal, P. Retailleau, A. Voituriez, G. Frison, A. Marinetti and X. Guinchard, *J. Am. Chem. Soc.*, 2020, **142**, 3797.
- 6 For reviews, see: a) E. Manoni and M. Bandini, *Eur. J. Org. Chem.*, 2016, 3135; b)D. Campeau, D. F. León Rayo, A. Mansour, K. Muratov and F. Gagosz, *Chem. Rev.*, 2020, DOI 10.1021/acs.chemrev.0c00788.
- For reviews, see: a) Z.-L. Xia, Q.-F. Xu-Xu, C. Zheng and S.-L. You, *Chem. Soc. Rev.*, 2020, 49, 286; b) H. Abou-Hamdan, C. Kouklovsky and G. Vincent, *Synlett*, 2020, 31, 1775; c) S. Park and S. Chang, *Angew. Chem. Int. Ed.*, 2017, 56, 7720; d) C. Zheng and S.-L. You, *Chem*, 2016, 1, 830; e) C.-X. Zhuo, C. Zheng and S.-L. You, *Acc. Chem. Res.*, 2014, 47, 2558; f) C.-X. Zhuo, W. Zhang and S.-L. You, *Angew. Chem. Int. Ed.*, 2012, 51, 12662; g) S. P. Roche and J. A. Porco, *Angew. Chem. Int. Ed.*, 2011, 50, 4068.
- 8 Previous work from our group: a) N. Sabat, F. Soualmia, P. Retailleau, A. Benjdia, O. Berteau and X. Guinchard, *Org. Lett.*, 2020, 22, 4344; b) V. Magné, Y. Sanogo, C. S. Demmer, P. Retailleau, A. Marinetti, X. Guinchard and A. Voituriez, *ACS Catal.*, 2020, 10, 8141; c) N. Glinsky-Olivier, P. Retailleau and X. Guinchard, *Eur. J. Org. Chem.*, 2018, 2018, 5823; d) V. Magné, A. Marinetti, V. Gandon, A. Voituriez and X. Guinchard, *Adv. Synth. Catal.*, 2017, 359, 4036; e) V. Magné, F. Blanchard, A. Marinetti, A. Voituriez and X. Guinchard, *Adv. Synth. Catal.*, 2016, 358, 3355
- 9 J. An and M. Bandini, Eur. J. Org. Chem., 2020, 4087.

- 10 a) R.-Q. Xu, P. Yang and S.-L. You, *Chem. Commun.*, 2017, **53**, 7553; b) J. Zheng, S.-B. Wang, C. Zheng and S.-L. You, *J. Am. Chem. Soc.*, 2015, **137**, 4880; c) Q. Cheng, Y. Wang and S.-L. You, *Angew. Chem. Int. Ed.*, 2016, **55**, 3496; d) D. Shen, Q. Chen, P. Yan, X. Zeng and G. Zhong, *Angew. Chem. Int. Ed.*, 2017, **56**, 3242; e) J. Nan, J. Liu, H. Zheng, Z. Zuo, L. Hou, H. Hu, Y. Wang and X. Luan, *Angew. Chem. Int. Ed.*, 2015, **54**, 2356; f) T. Oguma and T. Katsuki, *J. Am. Chem. Soc.*, 2012, **134**, 20017; g) J. An, A. Parodi, M. Monari, M. Castineira Reis, C. Silva Lopez and M. Bandini, *Chem. Eur. J.*, 2017, **23**, 17473; h) W.-T. Wu, R.-Q. Xu, L. Zhang and S.-L. You, *Chem. Sci.*, 2016, **7**, 3427; i) J. An, L. Lombardi, S. Grilli and M. Bandini, *Org. Lett.*, 2018, **20**, 7380; j) R. Pedrazzani, J. An, M. Monari and M. Bandini, *Eur. J. Org. Chem.*, 2021, **2021**, 1732.
- 11 a) G. Zhu, G. Bao, Y. Li, J. Yang, W. Sun, J. Li, L. Hong and R. Wang, *Org. Lett.*, 2016, **18**, 5288; b) Z.-L. Xia, C. Zheng, R.-Q. Xu and S.-L. You, *Nature Commun.*, 2019, **10**; c) Q. Yin, S.-G. Wang, X.-W. Liang, D.-W. Gao, J. Zheng and S.-L. You, *Chem. Sci.*, 2015, **6**, 4179; d) M. Uyanik, T. Yasui and K. Ishihara, *Angew. Chem. Int. Ed.*, 2010, **49**, 2175; e) T. Dohi, N. Takenaga, T. Nakae, Y. Toyoda, M. Yamasaki, M. Shiro, H. Fujioka, A. Maruyama and Y. Kita, *J. Am. Chem. Soc.*, 2013, **135**, 4558; f) B. Yang, X. Zhai, S. Feng, D. Hu, Y. Deng and Z. Shao, *Org. Lett.*, 2019, **21**, 330.
- 12 a) P. Adler, A. Fadel and N. Rabasso, *Chem. Commun.*, 2015, **51**, 3612; b) P. Adler, F. Gomes, A. Fadel and N. Rabasso, *Eur. J. Org. Chem.*, 2013, **2013**, 7546.