



**HAL**  
open science

## A general meta-ecosystem model to predict ecosystem function at landscape extents

Eric Harvey, Justin N Marleau, Isabelle Gounand, Shawn J Leroux, Carina R Firkowski, Florian Altermatt, F Guillaume Blanchet, Kevin Cazelles, Cindy Chu, Cassidy d'Aloia, et al.

### ► To cite this version:

Eric Harvey, Justin N Marleau, Isabelle Gounand, Shawn J Leroux, Carina R Firkowski, et al.. A general meta-ecosystem model to predict ecosystem function at landscape extents. 2021. hal-03407501

**HAL Id: hal-03407501**

**<https://hal.science/hal-03407501v1>**

Preprint submitted on 28 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title:** A general meta-ecosystem model to predict ecosystem function at landscape extents

2 **Running title:** Metaecosystem model for landscape of function

3

4 **Type of article:** Perspective

5

6 **Authors:** Eric Harvey<sup>1\*</sup>, Justin N. Marleau<sup>2\*</sup>, Isabelle Gounand<sup>3</sup>, Shawn J. Leroux<sup>4</sup>, Carina R.

7 Firkowski<sup>5</sup>, Florian Altermatt<sup>6a,b,c</sup>, F. Guillaume Blanchet<sup>7a,b,c</sup>, Kevin Cazelles<sup>8</sup>, Cindy Chu<sup>9</sup>,

8 Cassidy D'Aloia<sup>10</sup>, Louis Donelle<sup>5</sup>, Dominique Gravel<sup>7a</sup>, Frédéric Guichard<sup>2</sup>, Kevin McCann<sup>8</sup>,

9 Jonathan L.W. Ruppert<sup>5,11</sup>, Colette Ward<sup>9</sup>, Marie-Josée Fortin<sup>5</sup>

10 Authors marked with an asterisk\* have contributed equally to this work

11 **Corresponding Author:** eric.harvey@umontreal.ca

12

13 **Affiliations:**

14 <sup>1</sup> Département de sciences biologiques, Université de Montréal, Montréal, H2V0B3, Canada.

15 <sup>2</sup> Department of Biology, McGill University, 1205 Docteur Penfield, Montreal, Quebec,

16 H3A1B1, Canada

17 <sup>3</sup> Sorbonne Université, CNRS, UPEC, CNRS, IRD, INRA, Institut d'écologie et des sciences de

18 l'environnement, IEES, F-75005 Paris, France

19 <sup>4</sup> Department of Biology, Memorial University of Newfoundland and Labrador, St. John's,

20 Newfoundland and Labrador, Canada, A1B 3X9.

21 <sup>5</sup> Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario,

22 Canada, M5S 3B2

23 <sup>6a</sup> Eawag: Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic

24 Ecology, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland.

25 <sup>6b</sup> Department of Evolutionary Biology and Environmental Studies, University of Zurich,  
26 Winterthurerstr. 190, CH-8057 Zürich, Switzerland.

27 <sup>6c</sup> Research Priority Programme Global Change and Biodiversity (URPP-GCB), University of  
28 Zurich, Winterthurerstr. 190, CH-8057 Zürich, Switzerland.

29 <sup>7a</sup> Département de biologie, Université de Sherbrooke, 2500 Boulevard Université, Sherbrooke,  
30 Québec, Canada, J1K 2R1

31 <sup>7b</sup> Département de mathématique, Université de Sherbrooke, 2500 Boulevard Université,  
32 Sherbrooke, Québec, Canada, J1K 2R1

33 <sup>7c</sup> Département des sciences de la santé communautaire, Université de Sherbrooke, 3001 12e  
34 Avenue Nord, Sherbrooke, Québec, Canada, J1H 5N4

35 <sup>8</sup> Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1, Canada;

36 <sup>9</sup> Great Lakes Laboratory for Fisheries and Aquatic Sciences, Fisheries and Oceans Canada, 867  
37 Lakeshore Road, Burlington, ON L7S 1A1, Canada

38 <sup>10</sup> Department of Biological Sciences, University of New Brunswick Saint John, Saint John, NB  
39 E2L 4L5, Canada

40 <sup>11</sup> Ecosystem and Climate Science, Toronto and Region Conservation Authority, 101 Exchange  
41 Avenue, Vaughan, ON, L4K 5R6

42

43 **Statement of authorship:** EH and MJF organized the working group, all authors participated in  
44 designing the research during the working group, EH wrote the first draft of the manuscript,  
45 JM;SL;IG;EH;MJF designed the theoretical framework, JM coded and ran the simulations, IG  
46 and CRF designed the illustrations, all authors contributed to further editions of the manuscript.

47 **Data statement:** No data was produced in this study. However, all code necessary to reproduce  
48 the model and results will be made publicly available upon publication of the manuscript.

49

50 **Keywords:** metaecosystem, metacommunity, cross-ecosystem subsidy, spatial networks,  
51 aquatic-terrestrial linkages, ecosystem function, landscape scale

52

53 **Number of words abstract:** 200

54 **Number of references:** 60

55 **Number of words main text:** 5547

56 **Number of figures and tables:** 5 Figures and 0 Table

57

58 **Abstract**

59 The integration of meta-ecosystem processes over large spatial extent is critical to predicting  
60 whether and how global changes might impact biodiversity and ecosystem functions. Yet, there  
61 remains an important gap in meta-ecosystem models to predict multiple ecosystem functions  
62 (e.g., carbon sequestration, elemental cycling, trophic efficiency) across different ecosystem  
63 types (e.g., terrestrial-aquatic, benthic-pelagic). We derive a generic meta-ecosystem model to  
64 predict ecosystem function at landscape extents by integrating the spatial dimension of natural  
65 systems as spatial networks of different habitat types connected by cross-ecosystem flows of  
66 materials and organisms. This model partitions the physical connectedness of ecosystems from  
67 the spatial flow rates of materials and organisms, allowing the representation of all types of  
68 connectivity across ecosystem boundaries as well as the interaction(s) between them. The model  
69 predicts that cross-ecosystem flows maximize the realization of multiple functions at landscape  
70 extent. Spatial flows, even the ones that significantly reduce the overall amount of nutrients in  
71 the meta-ecosystem, can reallocate nutrients to more efficient ecosystems, leading to greater  
72 levels of productivity at both local and regional scales. This '*cross-ecosystem efficiency*  
73 *hypothesis*' is a general and testable hypothesis emphasizing the complementarity and  
74 interconnectedness among ecosystems and the importance of addressing ecosystem diversity for  
75 meta-ecosystem function.

76           **Context: Ecosystem function(s) at the landscape scale**

77           The flows of resources, materials, and species can connect different types of ecosystems  
78 within a landscape (Polis *et al.* 1997; Loreau *et al.* 2003; Massol *et al.* 2011). Meta-ecosystem  
79 theory has been proposed to study these spatial flows across coupled ecosystems, while stressing  
80 how spatial and temporal changes in biodiversity within each ecosystem can affect ecosystem  
81 functions at the meta-ecosystem scale (Loreau *et al.* 2003; Gravel *et al.* 2010; Gounand *et al.*  
82 2014; Marleau *et al.* 2014). Meta-ecosystem theory, however, has recently been criticised for  
83 lack of connection to empirical research (Massol *et al.* 2011; Harvey *et al.* 2016; Gounand *et al.*  
84 2018a) and there is a current push to develop empirically motivated meta-ecosystem models.

85           Classic meta-ecosystem theory tends to focus on measuring how flows influence the  
86 stability of meta-ecosystems (Marleau *et al.* 2010; Gounand *et al.* 2014; McCann *et al.* 2021).  
87 Yet, there remains an important gap in meta-ecosystem models to predict how ecosystem  
88 connections influence functions other than stability (e.g., production, carbon sequestration,  
89 elemental cycling, and trophic efficiency) at both local and landscape scales. To address this gap,  
90 meta-ecosystem theory expanded the two-patch ecosystem approach (Gravel *et al.* 2010;  
91 Marleau *et al.* 2010) to multi-patch systems (Marleau *et al.* 2014; McCann *et al.* 2021) where all  
92 patches are of the same ecosystem type (i.e., terrestrial-terrestrial, aquatic-aquatic, but see  
93 Leroux & Loreau 2012). However, there is no current theoretical model investigating the spatial  
94 flow of both abiotic (i.e., resources, nutrients) and biotic (i.e., organisms) compartments across  
95 different ecosystem types (e.g., terrestrial-aquatic), in multi-patch systems (Massol *et al.* 2017;  
96 Gounand *et al.* 2018a). The theoretical and empirical integration of meta-ecosystem processes at  
97 a broad spatial extent is critical to predicting and therefore understanding whether and how  
98 global changes may impact biodiversity and ecosystem functions at the landscape scale.

99 Empirical examples of spatial flows of energy, materials, or organisms coupling different  
100 ecosystems abound and have recently been reviewed (see reviews in Gounand *et al.* 2018b;  
101 Montagano *et al.* 2019; Peller *et al.* 2020). Several of these studies focus on how cross-  
102 ecosystem exchanges or allochthonous flows affect dynamics at the ecotone (Richardson & Sato  
103 2015). What is missing are studies investigating the functional implications of meta-ecosystem  
104 dynamics at broader multi-patch spatial extents than the ecotone itself (but see Iwata *et al.* 2003;  
105 Largaespada *et al.* 2012). Yet, the effects of material and organismal flows are likely to  
106 propagate or even accumulate across landscapes driving regional variation in ecosystem  
107 function. In watersheds, for instance, different cross-ecosystem flows (e.g., litterfall, fish  
108 migration) will operate at different spatial scales and thus contribute to ecosystem functions (e.g.,  
109 primary and secondary production) at multiple spatial extents (Figure 1). The combined effects  
110 of those flows of abiotic and biotic compartments, however, should predict functioning at the  
111 whole landscape scale (Figure 1).

112 Here, we derive a meta-ecosystem model to predict ecosystem function(s) at landscape  
113 extents by integrating the spatial dimension of ecosystems as spatial networks of different habitat  
114 types connected by cross-ecosystem flows of materials and organisms. This meta-ecosystem  
115 model partitions the physical connectedness of ecosystems from the spatial flow rates of  
116 materials and organisms allowing the representation of all types of connectivity across ecosystem  
117 boundaries as well as the interaction(s) between these two properties. We use this model to make  
118 testable predictions on ecosystem function at landscape extents, using watersheds as an example,  
119 and to investigate the impacts of perturbations on cross-ecosystem flows and corresponding  
120 functions.

121

## 122 **Empirical meta-ecosystem – from the interface between ecosystems to the landscape**

123 Watersheds are a classic and suitable example to illustrate the potential of our proposed  
124 integrated meta-ecosystem approach because they are mosaics of terrestrial and aquatic  
125 ecosystems interconnected by spatial flows of materials, energy, and organisms (Hynes 1975).  
126 Moreover, because of their relevance as a functional unit for conservation and resource  
127 management, watersheds have been extensively studied and spatial flows of materials as well as  
128 organisms have been quantified in many watersheds (Figure 2 and Table S1 for an extensive  
129 review).

130 Previous studies have shown that inputs of terrestrial detritus to aquatic ecosystems are  
131 very common (Gounand *et al.* 2018b, 2020), and they can limit benthic invertebrate production  
132 and contribute to fish diet (Richardson 1991; Kawaguchi *et al.* 2003; Wallace *et al.* 2015) (see  
133 Figure 2a arrow A). Conversely, emerging aquatic insects contribute to the diets of terrestrial  
134 consumers (Nakano & Murakami 2001; Sabo & Power 2002; Iwata *et al.* 2003; Baxter *et al.*  
135 2005; Marczak & Richardson 2007; Bultman *et al.* 2014) (see Figure 2a arrow B). Movements of  
136 organisms, organic matter, and nutrients also occur within ecosystems either passively following  
137 directional flows along the dendritic network (upstream to downstream e.g., particulate organic  
138 matter, see Figure 2a arrow C) or actively via organismal movement (downstream to upstream  
139 e.g., fish migrations, Figure 2a,b arrows D and H). Biomass and resources can also be exchanged  
140 vertically between benthic and pelagic lake zones via the sinking and resuspension of plankton  
141 and organic matter (Jyväsjärvi *et al.* 2013; Matisoff *et al.* 2017) (see Figure 2a arrows E and F).

142 The large body of empirical research on flows of materials and organisms in watersheds  
143 highlight how different types of spatial flows have been studied mostly in isolation, and ignoring  
144 their bi-directional property (Schindler & Smits 2017, but see review in Marcarelli *et al.* 2020).



145 Taken as a whole, however, the data clearly demonstrate that multiple abiotic and biotic flows  
146 interact and flow reciprocally across different ecosystems in watersheds. The different flows can  
147 be separated into three broad categories: (1) **trophic flows within each ecosystem patch** (e.g.,  
148 biomass transfer along the food chain at one location), (2) **spatial flows among patches of the**  
149 **same ecosystem type** (e.g., ungulates foraging across different forest patches), and (3) **spatial**  
150 **flows across patches of different ecosystem types** (flows at the ecotone of two different  
151 ecosystem types, e.g., forest-lake). We surmise that by integrating these three types of flows into  
152 meta-ecosystem theory, we can predict the drivers of variations in ecosystem function across  
153 landscapes (Figure 1). The theory we derive in the next section can be reduced to models  
154 integrating various combinations of the three individual components listed above, but the full  
155 strength of our novel approach is in the integration of these three flow types.

156 Using watersheds as a case-study allows us to highlight (1) the biotic linkages that can  
157 emerge between ecosystems of different types (here terrestrial-aquatic), and (2) how cross-  
158 ecosystem biotic linkages at the ecotone interface are indirectly linked to the whole watershed  
159 via the connectivity structure of the landscape. Although we use watersheds to illustrate the  
160 usefulness of our model, the landscape perspective that we propose is relevant for any system for  
161 which spatial flows within ecosystem types (e.g., seagrass leaves decaying and flowing to an  
162 adjacent seagrass bed) and spatial flows across different ecosystem types (e.g., nutrients leaching  
163 from islands to the seagrass beds) are expected to interact and affect dynamics and functions at  
164 broader scales: marine-island, marine-freshwater, pelagic-benthic, and even, less intuitively,  
165 forest-grassland connections where behavioral movements within and across the two similar  
166 ecosystems can play an important role in driving divergence in trophic dynamics and

167 productivity (Abbas *et al.* 2012; Leroux *et al.* 2017; Gounand *et al.* 2018b; García-Callejas *et al.*  
168 2019).

169         Meta-ecosystem dynamics of spatial flows across different ecosystems involve spatial  
170 couplings where a specific trophic level contributes to different trophic levels in the connected  
171 ecosystems (Leroux & Loreau 2012; Montagano *et al.* 2019). Often, this occurs through the  
172 conversion of living to dead organic matter and eventually inorganic matter. For example,  
173 terrestrial herbivore insects falling in water can subsidize aquatic top-predators and decomposers  
174 at the same time, and also affect aquatic herbivores through indirect interactions by relaxing  
175 predation pressure via an alternative food source (Baxter *et al.* 2005; Allen & Wesner 2016;  
176 Montagano *et al.* 2019). Those indirect cross-ecosystem biotic interactions illustrate the  
177 permeability between ecosystems and the complexity of predicting how human actions in one  
178 ecosystem might affect coupled ecosystems (Leroux & Loreau 2012; Massol *et al.* 2017;  
179 Montagano *et al.* 2019).

180         Cross-ecosystem interactions also constitute a dominant mechanism by which changes in  
181 the processes in one locality can impact processes at a different location, even in the absence of  
182 dispersal (i.e., ‘spatial cascade’, see Gounand *et al.* 2017; García-Callejas *et al.* 2019). For  
183 instance, it has been shown that upstream forest cover contributes ~70% of all dissolved organic  
184 carbon loadings to watersheds of the North American Adirondack mountains (Canham *et al.*  
185 2004), and the spatial configuration of forest patches in watersheds is a direct driver of leaf litter  
186 availability in headwater streams (Little & Altermatt 2018). Cascading effects in space can also  
187 occur through the active movement of organisms subsidized by terrestrial resources along the  
188 connectivity structure of the river network. This is exemplified by the movement of aquatic  
189 invertebrates subsidized by red alder detritus (itself favored by human forest harvesting of other

190 species) from upstream reaches that will, in turn, subsidize downstream fish habitats (Wipfli &  
191 Musslewhite 2004).

192 The magnitude of any spatial cascade across the landscape could be controlled by three  
193 main factors: (1) the level of biotic movement (dispersal or regular foraging movements within a  
194 habitat) of organisms acting as consumers at multiple locations (McCann *et al.* 2005), (2) the  
195 passive abiotic movement of altered nutrient or decaying detritus (*sensus* Vannote *et al.* 1980),  
196 and (3) the constraints imposed by landscape configuration on these processes (Harvey &  
197 Altermatt 2019; McLeod & Leroux 2021). These factors need to be explicitly integrated to  
198 achieve the scaling up of ecosystem function from local to landscape extent.

### 199 **A meta-ecosystem model for landscape ecosystem function**

200 Here, we derive a meta-ecosystem model which integrates trophic flows within ecosystem  
201 patches (Figure 3), spatial flows among ecosystems of the same type (Figure 2 arrows C and D),  
202 and spatial flows across different ecosystem types (Figure 2 arrows A and B) in what would be  
203 by default a multi-patch system (e.g., a leaf falling from the terrestrial ecosystem into a lake that  
204 then decomposes as it moves between connected aquatic ecosystems). Mathematical tractability  
205 of such complexity is challenging and likely a reason why there is a gap in meta-ecosystem  
206 models that have patches that do not share the same ecosystem compartments. To overcome the  
207 complexity involved in integrating the three components above, we generalized a matrix  
208 approach to spatial networks (Figure 3).

209 We derived a modified spatial network formulation that generalises earlier work to allow  
210 us to handle less commonly modelled but empirically abundant spatial flows across different  
211 ecosystems (Figure 3) (Othmer & Scriven 1971; Jansen & Lloyd 2000; Kouvaris *et al.* 2015). In

212 the equation presented in Figure 3a,  $\mathbf{x}$  is a vector containing elements  $x_{ki}$  representing the stock  
 213 (i.e., mass of limiting nutrient) of ecosystem compartment  $k$  in ecosystem patch  $i$ , where  $k$  ranges  
 214 from 1 to the total number of ecosystem compartments  $m$ , and where  $i$  ranges from 1 to the  
 215 number of different ecosystem patches  $n$  (which could be of the same or of different ecosystem  
 216 types). How  $\mathbf{x}$  varies over time  $t$  can be represented by a system of ordinary differential  
 217 equations  $d\mathbf{x}/dt = \mathbf{G}(\mathbf{x})$ , where  $\mathbf{G}$  is a vector-valued function describing rates of change of each  
 218 ecosystem compartment. We decompose  $\mathbf{G}$  in two parts: *flows in local ecosystems* and *between*  
 219 *ecosystem spatial flows*. *Flows in local ecosystems* are collected in the vector-valued function  
 220  $\mathbf{F}(\mathbf{x})$ , while *between ecosystem spatial flows* (i.e., both spatial flows between discrete patches  
 221 within the same ecosystem type and across different ecosystems types) are the result of the  $\mathbf{QCx}$   
 222 matrix multiplication where  $\mathbf{Q}$  is the matrix characterizing the *spatial flow rates* and  $\mathbf{C}$  is the  
 223 matrix of *ecosystem physical connectedness*, both of which are  $nm \times nm$  matrices because  
 224 connectedness and flow rates are both species and ecosystem-specific. Similarly, the vector-  
 225 valued function  $\mathbf{F}$  is composed of  $nm$  functions,  $\mathbf{F}(\mathbf{x}) = [f_{1,1}(\mathbf{x}_1), f_{2,1}(\mathbf{x}_2), \dots, f_{i,1}(\mathbf{x}_i), \dots, f_{n,1}(\mathbf{x}_n),$   
 226  $f_{1,2}(\mathbf{x}_1), \dots, f_{n,2}(\mathbf{x}_n), \dots, f_{i,k}(\mathbf{x}_i), \dots, f_{n,m}(\mathbf{x}_n)]^T$  where  $\mathbf{x}_i = (x_{i,1}, \dots, x_{i,k}, \dots, x_{i,m})$  describe the local  
 227 flows to and from ecosystem compartment  $k$  in ecosystem patch  $i$ . To be more concrete, the  
 228 trophic flow of energy and nutrients in an ecosystem would be categorized in  $\mathbf{F}(\mathbf{x})$ , while the  
 229 biomass of a herbivore that moves from one ecosystem to another divided by the time measured  
 230 would be categorized in  $\mathbf{Q}$  and the direction of flows across the landscape would be categorized  
 231 in  $\mathbf{C}$ .

232 Assuming that the spatial flow rate of an ecosystem compartment does not change between  
 233 ecosystems, the matrix of spatial flow rates  $\mathbf{Q}$  is a diagonal matrix constructed by taking the

234 Kronecker tensor product of the traditional diagonal  $m \times m$  flow (or diffusion or movement) rate  
 235 matrix, which we denote as  $\mathbf{Q}'$  (Marleau *et al.* 2010, 2014, 2015; Marleau & Guichard 2019)  
 236 with an  $n \times n$  identity matrix ( $\mathbf{I}_{(n,n)}$ ):

$$\begin{aligned}
 237 \quad \mathbf{Q}' &= \begin{pmatrix} q_1 & 0 & \dots & 0 \\ 0 & q_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & q_m \end{pmatrix} \mathbf{I}_{(n,n)} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \\
 238 \quad \mathbf{Q} &= \mathbf{Q}' \otimes \mathbf{I}_{(n,n)} = \begin{pmatrix} q_1 \mathbf{I}_{(n,n)} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & q_2 \mathbf{I}_{(n,n)} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & q_m \mathbf{I}_{(n,n)} \end{pmatrix},
 \end{aligned}$$

239 where  $q_k$  is the between ecosystem spatial flow rate for ecosystem compartment  $k$  and  $\mathbf{0}$  is  
 240 an  $n \times n$  zero matrix. Note that if the spatial flow of one compartment is affected by the stock of  
 241 another compartment, then  $\mathbf{Q}'$  (and therefore  $\mathbf{Q}$ ) is no longer diagonal (for example, if a parasite  
 242 is completely dependent on its host for its movement across the landscape). Furthermore, if there  
 243 are ecosystem specific differences in spatial flow rates (for example, certain  
 244 genotypes/phenotypes in an ecosystem disperse more readily than those found in another  
 245 ecosystem), then we can replace the identity matrix with a weighted diagonal matrix instead.

246 The matrix of ecosystem physical connectedness  $\mathbf{C}$  is a block diagonal matrix derived from  
 247 the direct sum of the transpose of each ecosystem compartment  $k$ 's matrix of possible  
 248 connections,  $\mathbf{C}_k$ , which is an  $n \times n$  matrix whose elements,  $c_{ijk}$ , indicate if compartment  $k$  in  
 249 ecosystem  $i$  is physically capable of sending a spatial flow to ecosystem  $j$  (Jansen & Lloyd  
 250 2000):

251

$$\mathbf{C}_k = \begin{pmatrix} c_{11k} & \cdots & c_{1nk} \\ \vdots & \ddots & \vdots \\ c_{n1k} & \cdots & c_{nnk} \end{pmatrix}$$

252

$$\mathbf{C} = \bigoplus_{k=1}^m (\mathbf{C}_k)^T = \begin{pmatrix} (\mathbf{C}_1)^T & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & (\mathbf{C}_2)^T & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & (\mathbf{C}_m)^T \end{pmatrix}.$$

253 The diagonal entries of the  $\mathbf{C}_k$  matrices are negative to indicate the export of organisms and

254 materials from the focal ecosystem, while the off-diagonal entries are positive and represent the

255 arrival of organisms and materials from other ecosystems. Note that we need to transpose the

256 matrices due to the  $\mathbf{C}$  matrix being on the left-hand side of  $\mathbf{x}$  rather than on the right-hand side in257 other derivations (Jansen & Lloyd 2000). Unlike previous work (Marleau *et al.* 2010, 2014,258 2015), we do not require  $\mathbf{C}_k$  matrices to be symmetric and  $c_{iik}$  does not need to equal to the259 negative row sum of its other elements (i.e.  $c_{iik} \neq \sum_{j=1}^n c_{ij}$ ). This means that the flows

260 between ecosystems can be unidirectional or bidirectional (reciprocal) and they can leave the

261 meta-ecosystem partially or entirely.

262 This meta-ecosystem mathematical model allows for the incorporation of many types of

263 flows and thus a better application of the theory to empirical meta-ecosystems. First, there can be

264 different local dynamics (trophic flows) within different ecosystems (or ecosystem patches) for

265 each ecosystem compartment. For example, a terrestrial herbivore (e.g., grasshopper

266 (*Trimerotropis verruculata*)) will interact differently in the local river system, where it dies and

267 becomes detritus, than in the local forest system, where it is a herbivore. This feature of the

268 model makes it possible to appropriately model spatial flows across different ecosystem types,

269 because these types of flows often result in the material or organism moving across different

270 compartments in the donor and recipient ecosystem (in our above-mentioned example, the  
271 flowing material is a living herbivore in the donor ecosystem while it is dead detritus in the  
272 recipient ecosystem). Second, each ecosystem compartment can have unique physical  
273 connectedness (see Figure 3d), which is likely to happen for species that differ in preferred  
274 habitat or foraging areas (McLeod & Leroux 2021). For example, an aquatic-terrestrial boundary  
275 may be more permeable for a terrestrial avian predator (e.g., osprey (*Pandion haliaetus*)) than a  
276 small terrestrial herbivore (e.g., snowshoe hare (*Lepus americanus*)). Third, we can separate the  
277 effects of physical connectedness among ecosystems from the rate of spatial flows (flow  
278 intensity), which are normally measured separately from one another for both organisms and  
279 materials. The combination of physical connectedness of ecosystems and movement or flow  
280 potential leads to realized connectivity. With this model, we are now able to predict impacts of  
281 spatial flows in a simplified watershed meta-ecosystem.

## 282 **Model application: a simulated forest-lake-stream meta-ecosystem**

283 We apply the model to a simulated watershed composed of two aquatic (i.e., stream and  
284 lake) and one riparian forest ecosystem (Figure 3d). Each ecosystem has its own local flows or  
285 internal dynamics of material transfer among its inorganic nutrients, autotrophic, and  
286 heterotrophic components (Figure 3a-b). The local ecosystem components can potentially flow  
287 across boundaries, such that an aquatic herbivore may enter into a terrestrial environment, e.g.,  
288 when aquatic insect larvae mature into flying insects on land. The majority of current meta-  
289 ecosystem theory models flows among ecosystems as diffusion, and therefore implicitly assume  
290 that the material is of the same trophic level and composition in all patches and that it flows with  
291 the gradient in resources (i.e., from high to low). Other studies modelled direct flows from one  
292 ecosystem to another across trophic levels, i.e. a consumer in an ecosystem consumes a resource

293 in another (McCann *et al.* 2005; García-Callejas *et al.* 2019). However, such a flow implicitly  
294 assumes that there is instantaneous movement between ecosystems for either the consumer  
295 and/or resource, and therefore tight coupling between consumption and movement. An  
296 alternative is to explicitly model the dynamics of a non-local compartment in its non-local  
297 ecosystem type (see Figure 3b; Leroux & Loreau 2012). While this approach creates more  
298 variables to keep track of, it also helps us generalize our methods to more diverse situations and  
299 allows for cleaner mathematical treatment.

300 The model units are stocks expressed in mass of a limiting nutrient, such as nitrogen, in  
301 each model compartment (e.g., the mass of nitrogen in autotrophs). Here, vector  $\mathbf{x}$  keeps track in  
302 each ecosystem patch  $i$  of the amount of inorganic nutrients ( $N_i$ ), detritus ( $D_i$ ), aquatic and  
303 terrestrial autotroph stocks ( $A_{A,i}$  and  $A_{T,i}$ ), herbivore stocks ( $H_{A,i}$  and  $H_{T,i}$ ) and predator stocks ( $P_{A,i}$   
304 and  $P_{T,i}$ ) ( $m=8$ ) across a forest, a lake and a stream each composed of a single patch ( $n=3$ ).  
305 Vector  $\mathbf{x}$  has therefore a size of  $mn=24$ .

306 In this simulation, we consider a forest that surrounds a lake and a stream that flows out of  
307 the lake (Figure 3) and common flows among these ecosystems (see Figure 2). Live plant  
308 biomass (e.g., leaves, branches), dead organic matter (e.g., dead leaves, top soil) and inorganic  
309 nutrients can fall into and runoff in the lake, while aquatic insect herbivores (e.g., caddisfly  
310 (*Trichoptera sp.*)) can emerge and enter the forest (Figure 3d). Nutrients, dead organic matter  
311 (detritus), alive terrestrial plant biomass (e.g., white birch (*Betula papyrifera*)) and  
312 phytoplankton flow passively downstream from the lake to the stream, while aquatic herbivores  
313 and carnivores (e.g., dragonfly (*Odonata sp.*)) can move actively between the stream and lake  
314 based on diffusive movements (Figure 3). Thus, while the ecosystems are all physically  
315 connected, the realized biotic connectivity (as defined by the **QC** matrix) is limited and much of



316 the abiotic connectivity is unidirectional. We simulated our system in this way to highlight the  
317 effects of meta-ecosystem connectivity.

318 To model flows in local ecosystems, we assumed a linear food chain for the biotic  
319 ecosystem compartments with Lotka-Volterra functional responses when they are in their local  
320 or donor ecosystem type (Figure 3). When biotic compartments flow from terrestrial to aquatic  
321 or from aquatic to terrestrial ecosystems, the biotic compartments considered here simply  
322 become dead organic material at a given rate as they can only survive a limited time in the  
323 recipient ecosystems (Figure 3).

#### 324 *Simulation scenarios*

325 We chose parameters to produce realistic local flow hierarchy, such that the forest  
326 ecosystem has the greatest primary production, while the aquatic ecosystems are more efficient  
327 in the transfer of biomass between trophic levels and have faster mineralization (Gounand *et al.*  
328 2020). We used empirical parameter ranges for spatial flow rates to explore relevant parameter  
329 space as examples (see Table S1). Furthermore, our analysis focused on functions and  
330 parameters that ensured a stable equilibrium in all ecosystems over the range of parameter values  
331 investigated. For this study, we ran one thousand simulations where we varied the nutrient inputs  
332 to the local ecosystems to examine the impacts of nutrient supply on relative ecosystem  
333 functioning (see Supplementary materials for full case study model equations and details on  
334 model parameterization).

335 To highlight the importance of spatial flows across different ecosystems, we developed  
336 four scenarios where (i) the meta-ecosystem had no spatial flows at all, (ii) the meta-ecosystem  
337 was fully connected (bidirectional flows of biotic and abiotic material between terrestrial and

338 aquatic ecosystems), (iii) scenario without a biotic terrestrial to aquatic flow, and (iv) scenario  
339 without a biotic aquatic to terrestrial flow.

340 Scenario (i) can be viewed as a baseline for compartment stocks (i.e. N, D, A, H, P) and  
341 ecosystem function (primary, herbivore and predator production) through local processes only at  
342 the meta-ecosystem and local levels (Figures 4 and S1). As this scenario has less nutrient losses  
343 given the reduced flow out of the meta-ecosystem, we see higher nutrient stocks (Figure S1) and  
344 higher primary production (Figure 4D) than the scenarios that consider spatial flows, though this  
345 effect is reduced or absent for secondary production for spatial flow scenarios (ii) and (iv)  
346 (Figures 4H and 4L). The key spatial flow to understanding these phenomena is the terrestrial  
347 primary producer flow to the lake. Without the terrestrial primary producer flow (scenario (iii)),  
348 nutrient stocks are relatively high in the meta-ecosystem as terrestrial primary producers keep  
349 their stocks 'on land', leading to higher forest primary and secondary productivity (Figures 4A, E  
350 and I). The spatial flow from the terrestrial to the aquatic ecosystems is then solely composed of  
351 inorganic nutrients and detritus. The detritus and inorganic nutrient flows are relatively small  
352 compared to potential primary producer spatial flows, and the flux into the lake is less than the  
353 flux out of the lake, leading to a decline in lake productivity (Figures 4B, F and J), while the  
354 stream benefits from the larger flows into it compared to those out of it (Figures 4C, G and K).

355 When nutrients were released by the terrestrial primary producers into the aquatic  
356 ecosystems, however, they stimulated aquatic primary producers significantly, which led to  
357 higher secondary production in the stream (Figure 4). This aquatic secondary production is high  
358 enough to compensate for the losses in the terrestrial ecosystem, despite significant losses of  
359 nutrients out of the meta-ecosystem (Figure S1). Furthermore, if aquatic herbivores do not send  
360 individuals to the terrestrial ecosystem, thus breaking reciprocal links between the terrestrial and

361 aquatic ecosystems, predator production greatly increases in the lake, leading to greater meta-  
362 ecosystem predator production overall (Figures 4I-L).

363         These simulation results show how spatial flows between different ecosystems can lead to  
364 complex responses at both local and meta-ecosystem scales. Spatial flows, even the ones that  
365 significantly reduce the overall amount of nutrients in the meta-ecosystem, can reallocate  
366 nutrients to more efficient ecosystems, leading to greater levels of productivity at local and even  
367 regional scales. In other words, cross-ecosystem flows maximize the realization of multiple  
368 functions at landscape extent. We termed this finding the ‘*cross-ecosystem efficiency*  
369 *hypothesis*’. This general hypothesis emphasizes the complementarity and interconnectedness  
370 among ecosystems in the landscape and the importance of addressing ecosystem diversity for  
371 meta-ecosystem function. Therefore, while the application of our model is relatively simple, it  
372 provides a realistic scenario as it generated predictions that were not possible with previous  
373 meta-ecosystem theory. Thus, by utilizing tools to better integrate real world ecosystems into  
374 theory, we have expanded the possibilities of theory and can motivate empirical tests in the  
375 future.

## 376 **Perspectives for predicting ecosystem functions across landscapes**

### 377 **Landscape function(s)**

378         The meta-ecosystem framework we developed highlights the interdependence among the  
379 different ecosystems that can be found alongside one another at the landscape scale. Terrestrial  
380 ecosystems drive primary production but are less efficient than aquatic at transferring the energy  
381 produced up the food chain, which leads to higher nutrient stock accumulation. Conversely,  
382 aquatic systems tend to be more heterotrophic with higher trophic efficiency, which leads to

383 higher secondary production. This ‘spatial trade-off’ can lead to co-dependencies between  
384 systems that share limiting resources through spatial flows (Gounand *et al.* 2017). With our  
385 model, we showed that this trade-off also means that accounting for spatial flows across different  
386 ecosystem types can maximize multiple functions related to biomass accumulation versus  
387 production at the landscape scale (Figures 4 and S1). When flows are accounted for, the energy  
388 lost by the terrestrial to the aquatic system is well compensated at the meta-ecosystem level by  
389 the increase in herbivore and predator production in the aquatic system (Figure 4). Thus, the  
390 landscape can be perceived as an assembly line where each ecosystem type has its own  
391 ‘expertise’ (e.g., biomass accumulation vs. production at different levels) and only by accounting  
392 for energy flows across those systems can we maximize the landscape of functions (hence the  
393 ‘cross-ecosystem efficiency hypothesis’, Figure 5).

394 By pairing our model with realistic landscape perturbation scenarios, we surmise that the  
395 conceptual approach we developed can be used to predict how ecosystem functions may be  
396 altered by different types of perturbations (Figure 5). Watersheds not only constitute relatively  
397 good enclosed functional ecological units, but they also correspond to a human functional unit.  
398 In that sense, watersheds include where we find and manage natural resources (wood, mine),  
399 where we grow our food, where we go for recreational purposes (camping, fishing) and where  
400 we live (urban centers) (Dudgeon *et al.* 2006; Abell *et al.* 2007). The spatial configuration of  
401 these land-use patterns is essential to understanding feedback effects between local ecosystems  
402 and the whole landscape. The conceptual approach we developed here, emphasizes the  
403 importance of considering the mesoscale (watershed, landscape) as a scale of reference for  
404 understanding changes in ecosystem functions that are relevant for human societies.

405           The flow perturbation results also suggest that those cross-scale interactions generally  
406 maximize functions at the landscape scale whereas each of the three ecosystems tend to be more  
407 dynamic in production. Those dynamical variations in production at each ecosystem scale tend to  
408 cancel or compensate each other out at higher spatial scales. Indeed, one could imagine that  
409 some scenarios could lead to the amplification of perturbations in space (across the three  
410 connected ecosystems) and across spatial scales (see McCann *et al.* 2021). In a context where  
411 conservation and restoration sciences are increasingly concerned with landscape management (as  
412 opposed to local-scale habitat management), our approach could lend a direct contribution to  
413 understanding and predicting how a particular perturbation in a given location of the landscape  
414 might affect the whole landscape function in terms of production and standing stocks.

#### 415           **Linking meta-ecosystem theory and empirical studies**

416           We propose a meta-ecosystem model with three major components. First, the model  
417 integrates flows in local ecosystems, spatial flows within the same ecosystem, and spatial flows  
418 across different ecosystems. Empirical studies showed that flows at all three levels are common  
419 (Figure 2, Table S1; see reviews in Allen & Wesner 2016; Gounand *et al.* 2018a; Montagano *et*  
420 *al.* 2019). Yet, existing theory usually focuses on only one of these components. Second, the  
421 framework we propose is flexible enough to incorporate abiotic and biotic flows at different  
422 scales. Empirical studies highlight that the spatial and temporal scales of abiotic and biotic flows  
423 may differ and that there are important interactions between abiotic and biotic flows (see review  
424 in McLeod & Leroux 2021), yet existing theory rarely captures these dynamics - especially in  
425 multi-patch models (Table S1, Figure 1). Third, our framework partitions the physical  
426 connectedness of ecosystems from the movement or flow potential (rate) of a compartment. For  
427 a flow to occur, there needs to be both physical connection and movement potential. This

428 partitioning has three benefits; (i) it allows for a mathematically tractable way to model complex  
429 connectivity scenarios (i.e., K tensor product), (ii) it makes it possible to allow for variable flow  
430 scenarios across different local compartments, for instance in terms of directionality and  
431 differences of connectivity among trophic levels depending on species mobility, and this  
432 flexibility matches with empirical variability in ecosystem connections, and (iii) it provides a  
433 model framework to make predictions based on metrics that are often empirically measured or  
434 can be measured - for example landscape permeability (e.g., terrain ruggedness, Chetkiewicz &  
435 Boyce 200) and animal movement (e.g., movebank, Kranstauber *et al.* 2011). The model could  
436 also be used to determine most important fluxes in and across ecosystems to focus future  
437 monitoring and research efforts. With a case study, we illustrate how this model can be fit to a  
438 specific meta-ecosystem and how it can be used to provide testable predictions in specific  
439 systems. In our forest-lake-stream meta-ecosystem case study, we predict that removing key  
440 flows (e.g., terrestrial plants via forest harvesting) can cascade to impact stocks and productions  
441 at local and landscape scales (Figures 1, 4 and 5), while emphasizing how complementarity in  
442 functions among ecosystem types can maximize ecosystem function in the landscape (*'cross-*  
443 *ecosystem efficiency hypothesis'*).

444 Overall, we anticipate that our framework could be used to develop a suite of predictions  
445 for different ecosystems pertaining to how different flows mediate diverse ecosystem functions.  
446 For example, the model could explore how the demonstrated decline in Pacific salmon  
447 (*Onchorynchus spp.*, e.g. Oke *et al.* 2020) can impact primary and secondary production of natal  
448 streams and riparian forests in the Pacific Northwest of North America. More broadly, habitat  
449 fragmentation and land-use changes are the main factors contributing to the current global  
450 biodiversity loss (IPBES 2019). In that context, we need to better integrate connectivity loss

451 across trophic levels to make testable predictions about the effects of reduced connectivity on  
452 ecosystem function at the mesoscale.

453 From a theory perspective, the model we propose is flexible enough to recover many  
454 existing meta-ecosystem model formulations. For example, by assuming that spatial flows only  
455 occur in the same compartment (i.e., herbivores flow to herbivores), our model can be simplified  
456 to study only spatial flows within the same ecosystem. The use of matrices in our framework  
457 makes for a good match between model predictions and empirical ecological data which are  
458 often readily presented as matrices (e.g., community, connectivity; Gravel *et al.* 2016). In  
459 addition, we advance our framework as a call for theoretical and empirical spatial ecologists to  
460 work together to study landscape scale ecosystem functions. Much of the underlying theory  
461 focuses on stability as a key function but other functions such as production and elemental  
462 cycling are also critical and more commonly measured in natural systems. Recent advances in  
463 spatial stoichiometry provide the statistical methods to map empirical patterns in limiting  
464 nutrients across a landscape (Collins *et al.* 2017; Leroux *et al.* 2017; Soranno *et al.* 2019). These  
465 spatially explicit predictions of elemental surfaces can be used to partially parameterize meta-  
466 ecosystem models such as the one we propose here. Predictions can then be made on current and  
467 future functions.

468 The three-patch implementation of the model we propose here illustrates how to make  
469 predictions on cross-ecosystem exchanges and their regional scale impacts, but it assumes that  
470 each ecosystem is homogeneous in resource distribution. Future use of our framework could also  
471 include multiple patches within each ecosystem so that flows within the same ecosystem and  
472 among different ecosystems would be fully included. This would lead to a more precise

473 quantification of heterogeneity in ecosystem functions (as shown in Figure 1), for instance  
474 allowing for spatial gradients in ecosystem functions to emerge from ecotones.

475 Resource flows from one ecosystem to another are also known to vary at different time  
476 scales, from within a year to inter-annually (Spencer *et al.* 2005). Observational measurements of  
477 those flows could be established as a natural baseline against which flows following a  
478 perturbation could be simulated to analyze changes in the structural stability of the matrix or  
479 resilience (time of return to the natural baseline). This approach could lend interesting insights  
480 on how to offset human impacts, urban development and land conversion, on cross-ecosystem  
481 flows, by providing information such as the amount/configuration of natural cover in riparian  
482 zones to maintain underlying process, especially in the context of a well-connected system like a  
483 river where effects can spread across the watershed. Thus, our approach can be useful to develop  
484 formal tests of landscape implications of local perturbations propagated via spatial cascades.

485 Finally, our framework can also be parametrized with empirical data which could help to  
486 address questions about the functioning of natural systems in the face of perturbations. For  
487 instance, our approach could potentially shed new light on carbon sequestration at the landscape  
488 scale. Most carbon sequestration models assume homogeneous landscapes and ignore animals  
489 (Schmitz *et al.* 2018), but it is not clear how accounting for abiotic and biotic spatial flows in  
490 carbon might affect those predictions. Previous work has shown that carbon exchanges between  
491 ecosystems at large spatial scales can be highly significant (Gounand *et al.* 2018b). In that  
492 context, human induced perturbations such as climate change, but also land use change and  
493 habitat fragmentation could potentially alter carbon flows among ecosystems (Leroux *et al.*  
494 2017), thus influencing carbon sequestration at regional and landscape extents. Yet, much  
495 research is needed to make the link between different types of perturbations and their impacts on



496 spatial flows, and the cumulative effects of different types of perturbations on ecosystem  
497 functions in the landscape.

498

499 **Acknowledgements**

500 We thank the Canadian Institute for Ecology and Evolution for funding the working group that  
501 led to this publication.

502

503 **References**

504

505 Abbas, F., Merlet, J., Morellet, N., Verheyden, H., Hewison, A.J.M., Cargnelutti, B., *et al.*  
506 (2012). Roe deer may markedly alter forest nitrogen and phosphorus budgets across  
507 Europe. *Oikos*, 121, 1271–1278.

508 Abell, R., Allan, J.D. & Lehner, B. (2007). Unlocking the potential of protected areas for  
509 freshwaters. *Biol. Conserv.*, 134, 48–63.

510 Allen, D.C. & Wesner, J.S. (2016). Synthesis: comparing effects of resource and consumer  
511 fluxes into recipient food webs using meta-analysis. *Ecology*, 97, 594–604.

512 Baxter, C.V., Fausch, K.D. & Carl Saunders, W. (2005). Tangled webs: reciprocal flows of  
513 invertebrate prey link streams and riparian zones. *Freshw. Biol.*, 50, 201–220.

514 Bultman, H., Hoekman, D., Dreyer, J. & Gratton, C. (2014). Terrestrial deposition of aquatic  
515 insects increases plant quality for insect herbivores and herbivore density. *Ecol.*  
516 *Entomol.*, 39, 419–426.

517 Canham, C.D., Pace, M.L., Papaik, M.J., Primack, A.G.B., Roy, K.M., Maranger, R.J., *et al.*  
518 (2004). A Spatially Explicit Watershed-Scale Analysis of Dissolved Organic Carbon in  
519 Adirondack Lakes. *Ecol. Appl.*, 14, 839–854.

520 Chetkiewicz, C.-L.B. & Boyce, M.S. (2009). Use of resource selection functions to identify  
521 conservation corridors. *J. Appl. Ecol.*, 46, 1036–1047.

522 Collins, S.M., Oliver, S.K., Lapierre, J.-F., Stanley, E.H., Jones, J.R., Wagner, T., *et al.* (2017).  
523 Lake nutrient stoichiometry is less predictable than nutrient concentrations at regional  
524 and sub-continental scales. *Ecol. Appl.*, 27, 1529–1540.

525 Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.-I., Knowler, D.J., Lévêque, C., *et*  
526 *al.* (2006). Freshwater biodiversity: importance, threats, status and conservation  
527 challenges. *Biol. Rev.*, 81, 163–182.

528 García-Callejas, D., Molowny-Horas, R., Araújo, M.B. & Gravel, D. (2019). Spatial trophic  
529 cascades in communities connected by dispersal and foraging. *Ecology*, 100, e02820.

530 Gounand, I., Harvey, E., Ganesanandamoorthy, P. & Altermatt, F. (2017). Subsidies mediate  
531 interactions between communities across space. *Oikos*, 126, 972–979.

532 Gounand, I., Harvey, E., Little, C.J. & Altermatt, F. (2018a). Meta-Ecosystems 2.0: Rooting the  
533 Theory into the Field. *Trends Ecol. Evol.*, 33, 36–46.

534 Gounand, I., Little, C.J., Harvey, E. & Altermatt, F. (2018b). Cross-ecosystem carbon flows  
535 connecting ecosystems worldwide. *Nat. Commun.*, 9, 4825.

536 Gounand, I., Little, C.J., Harvey, E. & Altermatt, F. (2020). Global quantitative synthesis of  
537 ecosystem functioning across climatic zones and ecosystem types. *Glob. Ecol. Biogeogr.*,  
538 29, 1139–1176.

539 Gounand, I., Mouquet, N., Canard, E., Guichard, F., Hauzy, Céline & Gravel, D. (2014). The  
540 Paradox of Enrichment in Metaecosystems. *Am. Nat.*, 184, 752–763.

541 Gravel, D., Guichard, F., Loreau, M. & Mouquet, N. (2010). Source and sink dynamics in meta-  
542 ecosystems. *Ecology*, 91, 2172–2184.

543 Gravel, D., Massol, F. & Leibold, M.A. (2016). Stability and complexity in model meta-

- 544 ecosystems. *Nat. Commun.*, 7, 12457.
- 545 Harvey, E. & Altermatt, F. (2019). Regulation of the functional structure of aquatic communities  
546 across spatial scales in a major river network. *Ecology*, 100, e02633.
- 547 Harvey, E., Gounand, I., Ward, C.L. & Altermatt, F. (2016). Bridging ecology and conservation:  
548 from ecological networks to ecosystem function. *J. Appl. Ecol.*, 54, 371–379.
- 549 Hynes, H.B.N. (1975). The stream and its valley. *SIL Proc. 1922-2010*, 19, 1–15.
- 550 IPBES. (2019). *Global assessment report on biodiversity and ecosystem services of the*  
551 *Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*.
- 552 Iwata, T., Nakano, S. & Murakami, M. (2003). Stream meanders increase insectivorous bird  
553 abundance in riparian deciduous forests. *Ecography*, 26, 325–337.
- 554 Jansen, V.A.A. & Lloyd, A.L. (2000). Local stability analysis of spatially homogeneous  
555 solutions of multi-patch systems. *J. Math. Biol.*, 41, 232–252.
- 556 Jyväsjärvi, J., Boros, G., Jones, R.I. & Hämäläinen, H. (2013). The importance of sedimenting  
557 organic matter, relative to oxygen and temperature, in structuring lake profundal  
558 macroinvertebrate assemblages. *Hydrobiologia*, 709, 55–72.
- 559 Kawaguchi, Y., Taniguchi, Y. & Nakano, S. (2003). Terrestrial invertebrate inputs determine the  
560 local abundance of stream fishes in a forested stream. *Ecology*, 84, 701–708.
- 561 Kouvaris, N.E., Hata, S. & Guilera, A.D.-. (2015). Pattern formation in multiplex networks. *Sci.*  
562 *Rep.*, 5, 10840.
- 563 Kranstauber, B., Cameron, A., Weinzerl, R., Fountain, T., Tilak, S., Wikelski, M., *et al.* (2011).  
564 The Movebank data model for animal tracking. *Environ. Model. Softw.*, 26, 834–835.
- 565 Largaespada, C., Guichard, F. & Archambault, P. (2012). Meta-ecosystem engineering: Nutrient  
566 fluxes reveal intraspecific and interspecific feedbacks in fragmented mussel beds.  
567 *Ecology*, 93, 324–333.
- 568 Leroux, S.J. & Loreau, M. (2012). Dynamics of Reciprocal Pulsed Subsidies in Local and Meta.  
569 *Ecosystems*, 15, 48–59.
- 570 Leroux, S.J., Wal, E.V., Wiersma, Y.F., Charron, L., Ebel, J.D., Ellis, N.M., *et al.* (2017).  
571 Stoichiometric distribution models: ecological stoichiometry at the landscape extent.  
572 *Ecol. Lett.*, 20, 1495–1506.
- 573 Little, C.J. & Altermatt, F. (2018). Landscape configuration alters spatial arrangement of  
574 terrestrial-aquatic subsidies in headwater streams. *Landsc. Ecol.*, 1–13.
- 575 Loreau, M., Mouquet, N. & Holt, R.D. (2003). Meta-ecosystems: a theoretical framework for a  
576 spatial ecosystem ecology. *Ecol. Lett.*, 6, 673–679.
- 577 Marcarelli, A.M., Baxter, C.V., Benjamin, J.R., Miyake, Y., Murakami, M., Fausch, K.D., *et al.*  
578 (2020). Magnitude and direction of stream–forest community interactions change with  
579 timescale. *Ecology*, 101, e03064.
- 580 Marczak, L.B. & Richardson, J.S. (2007). Spiders and subsidies: results from the riparian zone of  
581 a coastal temperate rainforest. *J. Anim. Ecol.*, 76, 687–694.
- 582 Marleau, J.N. & Guichard, F. (2019). Meta-ecosystem processes alter ecosystem function and  
583 can promote herbivore-mediated coexistence. *Ecology*, 100, e02699.

- 584 Marleau, J.N., Guichard, F. & Loreau, M. (2014). Meta-ecosystem dynamics and functioning on  
585 finite spatial networks. *Proc. R. Soc. Lond. B Biol. Sci.*, 281, 20132094.
- 586 Marleau, J.N., Guichard, F. & Loreau, M. (2015). Emergence of nutrient co-limitation through  
587 movement in stoichiometric meta-ecosystems. *Ecol. Lett.*, 18, 1163–1173.
- 588 Marleau, J.N., Guichard, F., Mallard, F. & Loreau, M. (2010). Nutrient flows between  
589 ecosystems can destabilize simple food chains. *J. Theor. Biol.*, 266, 162–174.
- 590 Massol, F., Altermatt, F., Gounand, I., Gravel, D., Leibold, M.A. & Mouquet, N. (2017). How  
591 life-history traits affect ecosystem properties: effects of dispersal in meta-ecosystems.  
592 *Oikos*, 126, 532–546.
- 593 Massol, F., Gravel, D., Mouquet, N., Cadotte, M.W., Fukami, T. & Leibold, M.A. (2011).  
594 Linking community and ecosystem dynamics through spatial ecology. *Ecol. Lett.*, 14,  
595 313–323.
- 596 Matisoff, G., Watson, S.B., Guo, J., Duewiger, A. & Steely, R. (2017). Sediment and nutrient  
597 distribution and resuspension in Lake Winnipeg. *Sci. Total Environ.*, 575, 173–186.
- 598 McCann, K.S., Cazelles, K., MacDougall, A.S., Fussmann, G.F., Bieg, C., Cristescu, M., *et al.*  
599 (2021). Landscape modification and nutrient-driven instability at a distance. *Ecol. Lett.*,  
600 24, 398–414.
- 601 McCann, K.S., Rasmussen, J.B. & Umbanhowar, J. (2005). The dynamics of spatially coupled  
602 food webs. *Ecol. Lett.*, 8, 513–523.
- 603 McLeod, A.M. & Leroux, S.J. (2021). Incorporating abiotic controls on animal movements in  
604 metacommunities. *Ecology*, 102, e03365.
- 605 Montagano, L., Leroux, S.J., Giroux, M.-A. & Lecomte, N. (2019). The strength of ecological  
606 subsidies across ecosystems: a latitudinal gradient of direct and indirect impacts on food  
607 webs. *Ecol. Lett.*, 22, 265–274.
- 608 Nakano, S. & Murakami, M. (2001). Reciprocal subsidies: Dynamic interdependence between  
609 terrestrial and aquatic food webs. *Proc. Natl. Acad. Sci.*, 98, 166–170.
- 610 Oke, K.B., Cunningham, C.J., Westley, P. a. H., Baskett, M.L., Carlson, S.M., Clark, J., *et al.*  
611 (2020). Recent declines in salmon body size impact ecosystems and fisheries. *Nat.*  
612 *Commun.*, 11, 4155.
- 613 Othmer, H.G. & Scriven, L.E. (1971). Instability and dynamic pattern in cellular networks. *J.*  
614 *Theor. Biol.*, 32, 507–537.
- 615 Peller, T., Andrews, S., Leroux, S.J. & Guichard, F. (2020). From Marine Metacommunities to  
616 Meta-ecosystems: Examining the Nature, Scale and Significance of Resource Flows in  
617 Benthic Marine Environments. *Ecosystems*.
- 618 Polis, G.A., Anderson, W.B. & Holt, R.D. (1997). Toward an integration of landscape and food  
619 web ecology: The Dynamics of Spatially Subsidized Food Webs. *Annu. Rev. Ecol. Syst.*,  
620 28, 289–316.
- 621 Richardson, J.S. (1991). Seasonal Food Limitation of Detritivores in a Montane Stream: An  
622 Experimental Test. *Ecology*, 72, 873–887.
- 623 Richardson, J.S. & Sato, T. (2015). Resource subsidy flows across freshwater–terrestrial  
624 boundaries and influence on processes linking adjacent ecosystems. *Ecohydrology*, 8,

- 625 406–415.
- 626 Sabo, J.L. & Power, M.E. (2002). Numerical Response of Lizards to Aquatic Insects and Short-  
627 Term Consequences for Terrestrial Prey. *Ecology*, 83, 3023–3036.
- 628 Schindler, D.E. & Smits, A.P. (2017). Subsidies of Aquatic Resources in Terrestrial Ecosystems.  
629 *Ecosystems*, 20, 78–93.
- 630 Schmitz, O.J., Wilmers, C.C., Leroux, S.J., Doughty, C.E., Atwood, T.B., Galetti, M., *et al.*  
631 (2018). Animals and the zoogeochemistry of the carbon cycle. *Science*, 362, eaar3213.
- 632 Soranno, P.A., Wagner, T., Collins, S.M., Lapierre, J.-F., Lottig, N.R. & Oliver, S.K. (2019).  
633 Spatial and temporal variation of ecosystem properties at macroscales. *Ecol. Lett.*, 22,  
634 1587–1598.
- 635 Spencer, D.F., Ksander, G.G. & Whitehand, L.C. (2005). Spatial and temporal variation in RGR  
636 and leaf quality of a clonal riparian plant: *Arundo donax*. *Aquat. Bot.*, 81, 27–36.
- 637 Vannote, R.L., Minshall, G.W., Cummins, K.W., Sedell, J.R. & Cushing, C.E. (1980). The River  
638 Continuum Concept. *Can. J. Fish. Aquat. Sci.*, 37, 130–137.
- 639 Wallace, J.B., Eggert, S.L., Meyer, J.L. & Webster, J.R. (2015). Stream invertebrate productivity  
640 linked to forest subsidies: 37 stream-years of reference and experimental data. *Ecology*,  
641 96, 1213–1228.
- 642 Wipfli, M.S. & Musslewhite, J. (2004). Density of red alder (*Alnus rubra*) in headwaters  
643 influences invertebrate and detritus subsidies to downstream fish habitats in Alaska.  
644 *Hydrobiologia*, 520, 153–163.
- 645
- 646

647 **Figure Legends**

648 **Figure 1.** Conceptual diagram showing how different components of a meta-ecosystem  
649 contribute to function(s). Top right panels focus on one specific process each (arrows). Bottom  
650 right panels show an example of how the associated spatial flows would influence secondary  
651 production in a rasterized representation of the landscape (darker colours have more influence).  
652 This can be understood as a log response ratio of an experiment where the flow is removed  
653 (response = secondary production with flow / secondary production without flow). The leftmost  
654 bottom panel presents the sum of flow effects. We propose a novel mathematical model to  
655 integrate the combined effect of those different types of flows at landscape scale.

656 **Figure 2.** Spatial flows in watersheds. a) Illustration and b) schematic diagram of flows of  
657 material and organisms connecting the different habitats of a watershed. We provide one hundred  
658 references quantifying these flows (identified by the numbers on the right panel), all available in  
659 Table S1, providing flow quantifications for watersheds in temperate and cold climates (i.e.,  
660 alpine, subarctic, arctic). The material of quantified flows are: A) Terrestrial detritus, leaves and  
661 insects, eggs deposition of amphibians, leached nitrogen; B) Emergent insects and amphibians,  
662 fish carcasses caught by terrestrial consumers; C) Detritus, sediment DOC, invertebrates drifting,  
663 fish and insects migrating downstream; D) Fish and insects migrating upstream; E) Plankton  
664 sinking, organic matter; F) Resuspension of particles by wind, recycling of benthic phosphorus  
665 by fish; G) Sediment, particulate organic matter, nitrogen flowing downstream, phosphorus  
666 transported by salmon juveniles migrating downstream; H) Spawning salmon migrating  
667 upstream.

668 **Figure 3.** Overview of a meta-ecosystem model that integrates local trophic flows, spatial  
669 flows within the same ecosystem and/or across different ecosystem types (here illustrated for a

670 boreal watershed used as a case-study in our simulations (see Fig. 4). **(a)** All eight ecosystem  
671 compartments included in the landscape, consisting of five trophic levels (detritus (D), inorganic  
672 nutrients (N), autotrophs (A), herbivores (H), and predators (P), with terrestrial and aquatic  
673 specific biotic compartments highlighted in green and blue color, respectively). **(b)** Example of  
674 local forest dynamics describing within ecosystem trophic fluxes among ecosystem  
675 compartments including consumption dynamics, production of detritus by organisms, and  
676 recycling into nutrients. Dotted arrows represent the leaking of nutrients due to the relative lack  
677 of efficiency of trophic interactions. Transparency of aquatic compartments highlights that these  
678 stocks are decaying into detritus in the terrestrial ecosystem without any demographic dynamics.  
679 **(c)** Landscape representation with spatial dynamics decomposed between physical connectedness  
680 among ecosystem patches (**C**) for each ecosystem compartment between each ecosystem (heads  
681 and tails of the arrows), and spatial flow rates (**Q**) (the styles of the body of the arrow). **(d)**  
682 Mathematical representation of the meta-ecosystem. See text for full model description.


683 **Figure 4.** Effects of meta-ecosystem spatial flows on primary production (**A, B, C** and **D**),  
684 herbivore production (**E, F, G** and **H**) and predator production (**I, J, K** and **L**) in the forest (**A, E**  
685 and **I**), lake (**B, F, and J**), stream (**C, G** and **K**) ecosystems and in the whole meta-ecosystem (**D,**  
686 **H** and **L**). The ‘no flows’ scenario is a baseline result with no spatial flows between ecosystems.  
687 The ‘all flows’ scenario is the meta-ecosystem with all the flows specified in Figure 3. The ‘no  
688  $P_T$  flow’ and ‘no  $H_A$  flow’ scenarios have the same spatial flows as the ‘all flows’ scenario,  
689 except for the terrestrial primary producer flow and the aquatic herbivore flow, respectively.  
690 Lines in the middle of the boxes represent median values of 1000 simulations, with the top and  
691 bottom of the boxes representing 75th and 25th percentiles, the whiskers approximate the  
692 expected non-outlier maximum and minimum values. Outliers are defined as being more than the

693 interquartile range multiplied by 1.5, and are represented by red crosses (see Supplementary  
694 Material for parameters and fuller description of the model).

695 **Figure 5.** Landscape-matrix representation of secondary production for the different  
696 scenarios of spatial flows shown on the left column, in our simulated boreal watershed. From top  
697 to bottom: (i) scenario with no spatial flows, (ii) all spatial flows, (iii) only aquatic herbivores  
698 crossing the forest-lake interface, or (iv) only leaf litter. The grey values are proportional to the  
699 median values (opacity of black is median value \* 10) provided in Figure 4 of herbivore and  
700 predator productions (middle columns) in each ecosystem type, which contrasts spatial  
701 distributions of the production among spatial flow scenarios. The right column provides total  
702 secondary production by superimposing the two landscapes. The numbers on the right are the  
703 sum of productions of the three ecosystems, giving total nitrogen used for annual secondary  
704 production at the landscape-scale. Note that here, cases of the matrix are not individual patches  
705 within ecosystems but just a coarse-grain way to represent the landscape.




706 **Figure 1**



707

708


709 **Figure 2**



710


711

712 **Figure 3**



713


714 **Figure 4**



715

716

717 **Figure 5**



718