



Fine morphokinetic motions in ASD children: effect of writing amplitude on the isochrony principle

Aurélie Benchekri, Laetitia Fradet, Romain Tisserand, C. Bidet-Ildei

► To cite this version:

Aurélie Benchekri, Laetitia Fradet, Romain Tisserand, C. Bidet-Ildei. Fine morphokinetic motions in ASD children: effect of writing amplitude on the isochrony principle. 19ème congrès de l'association des chercheurs en activités physiques et sportives (ACAPS), Oct 2021, Montpellier, France. hal-03410750

HAL Id: hal-03410750

<https://hal.science/hal-03410750>

Submitted on 1 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fine morphokinetic motions in ASD children: effect of writing amplitude on the isochrony principle


Aurélie Benchekri ^{1,2}, Laetitia Fradet ², Romain Tisserand ^{1,2}, Christel Bidet-Ildei ¹

¹ : Centre de Recherches sur la Cognition et l'Apprentissage, Université de Poitiers, Université de Tours, CNRS - UMR7295

² : Equipe RoBioSS, Institut PPRIME, Université de Poitiers, CNRS - UPR 3346

Introduction

Autism Spectrum Disorder (ASD) is a neurodevelopmental psychiatric condition characterized by 1) difficulties with social interactions and communication and 2) repeated and stereotypical movements or restricted interests.





Altered motor behaviours were recently observed in animal ASD models and related to **cerebellum disorders** (Jaber, 2017). The cerebellum is particularly involved in the control of **handwriting** (Palmis et al., 2017).


→ Is the writing movement control perturbed in ASD children?

Method

- 7 TD children (2 boys, mean age= 9 years) vs 3 ASD children (1 boy, mean age = 9 years).
- 5 "e" letters of five different sizes in cursive mode.




For each "e", metric coordinates were recorded using a digitizer and the Eye and Pen 3 software



- Movement times and amplitudes were extracted and the isochrony coefficient (γ) and baseline time (T_0) were computed for each participant.

Results



Discussion / Conclusion

2 ASD children/3 are out the typical results for γ and T_0
=> the isochrony principle could be perturbed in ASD

This suggests that the cerebellum disorder could disturb the isochrony principle in the ASD deficit.

Future inclusions are necessary to conclude

References

- Jaber, M. (2017). Le cervelet comme acteur majeur dans les troubles moteurs des syndromes autistiques. *L'Encéphale*, 43(2), 170-175.
Palmis, S., Danna, J., Velay, J.-L., & Longcamp, M. (2017). Motor control of handwriting in the developing brain : A review. *Cognitive Neuropsychology*, 34(3-4), 187-204.

