

Indolizy Carbene Ligand. Evaluation of Electronic Properties and Applications in Asymmetric Gold(I) Catalysis

Thibaut Martinez, Avassaya Vanitcha, Claire Troufflard, Nicolas Vanthuyne, Jérémy Forté, Geoffrey Gontard, Gilles Lemière, Virginie Mouriès-Mansuy,

Louis Fensterbank

▶ To cite this version:

Thibaut Martinez, Avassaya Vanitcha, Claire Troufflard, Nicolas Vanthuyne, Jérémy Forté, et al.. Indolizy Carbene Ligand. Evaluation of Electronic Properties and Applications in Asymmetric Gold(I) Catalysis. Angewandte Chemie International Edition, 2021, 63 (36), pp.19879-19888. 10.1002/anie.202106142. hal-03417325

HAL Id: hal-03417325 https://hal.science/hal-03417325

Submitted on 5 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Indolizy Carbene Ligand. Evaluation of Electronic Properties and Applications in Asymmetric Gold(I) Catalysis

Thibaut Martinez,^[a] Avassaya Vanitcha,^[a] Claire Troufflard,^[a] Nicolas Vanthuyne,^[b] Jérémy Forté,^[a] Geoffrey Gontard,^[a] Gilles Lemière,^[a]* Virginie Mouriès-Mansuy^[a]* and Louis Fensterbank^[a]*

 [a] Dr. T. Martinez, Dr. A. Vanitcha, C. Troufflard, J. Forté, G. Gontard, Dr. G. Lemière, Dr. V. Mouriès-Mansuy, Prof. L. Fensterbank Institut Parisien de Chimie Moléculaire Sorbonne Université, CNRS 4 Place Jussieu, CC 229, F-75252 Paris Cedex 05 (France) E-mail: gilles.lemiere@sorbonne-universite.fr, virginie-mansuy@sorbonne-universite.fr, louis.fensterbank@sorbonne-universite.fr
 [b] Dr. N. Vanthuyne Aix Marseille Univ, CNRS, Centrale Marseille, iSm2 F-13397 Marseille Cedex 20 (France)

Supporting information for this article is given via a link at the end of the document.((Please delete this text if not appropriate))

Abstract: We report herein a new family of carbene ligands based on an indolizine-ylidene (Indolizy) moiety. The corresponding gold(I) complexes are easily obtained from the gold(I)-promoted cyclization of allenylpyridine precursors. Evaluation of the electronic properties by experimental methods and also by DFT calculations confirm strong σ -donating and π -accepting properties of these ligands. Cationization of the gold(I) complexes generates catalytic species that trigger diverse reactions of (poly)unsaturated precursors. When armed with a methylene phosphine oxide moiety on the stereogenic center adjacent to the nitrogen atom, the corresponding bifunctional carbene ligands give rise to highly enantioselective heterocyclizations. DFT calculations brought some rationalization and highlighted the critical roles played by the phosphine oxide group and the tosylate anion in the asymmetric cyclization of yallenols.

Introduction

The last decades have witnessed a strong expansion in homogeneous gold catalysis promoting a broad range of organic transformations.^[1] The high π -Lewis acidity of cationic gold(I) complexes allow them to activate various unsaturations such as alkynes, allenes and also alkenes towards intra- or intermolecular nucleophilic addition.^[2] During the process, the ligands carried by the gold(I) center play a major role in terms of stability, allowing low catalyst loadings, modulation of activity, and selectivity.^[3] Furthermore, the linear coordination of gold(I) complexes that positions the ancillary ligand opposite to the substrate and thus to the outer-sphere attack of the nucleophile renders asymmetric gold catalysis a challenging goal.^[4] The role of the chiral ligand is therefore crucial notably by constraining the degrees of freedom of the system. In this context, C-ligands such as *N*-heterocyclic carbenes (NHCs A)^[5] have exhibited high potential. More accessible than a lot of phosphine ligands that are tedious to prepare and prone to oxidation, they have the ability to form stable complexes thanks to strong σ -bonding with the gold metallic center while π -backbonding to the carbene will be highly dependent from its structural features, the interplay of both effects leading to specific reactivities.^[6] In 2005, the group of Bertrand^[7] developed a new class of carbenes, the cyclic (alkyl)(amino)carbenes (CAACs **B**) in which a nitrogen of a classical NHC diaminocarbene was replaced by a carbon atom. Those are more σ -donating and π -accepting than the parent NHCs.^[8] The corresponding gold complexes^[9] have led to the first examples of hydroamination of alkynes and allenes using ammonia or hydrazine.^[10] The replacement of the alkyl substituent of CAACs by an aryl leads to the formation of new cyclic (amino)(aryl)carbenes (CAArCs **C**),^[11] that still feature high nucleophilicity but also enhanced electrophilicity resulting in a smaller single-triplet gap than for CAACs.

The implementation of asymmetric gold-catalyzed processes relying on monodentate C-ligands has initially consisted in the introduction of stereogenic centers on the imidazolium backbone or on the N-ligated substituents.^[12,13] Tomioka and coll.^[14] synthesized the first chiral C2-symmetric dihydroimidazol-2ylidene gold(I) complexes and applied them in a moderately enantioselective methoxycyclization of 1,6-enynes (up to 59% ee). Improved enantioselectivities (above 70%) for similar reactions were observed using C2-symmetric NHC complexes bearing bulky stereogenic N-substituents.[15,16] Still, these systems suffer from the fact that the nitrogen substituents are too remote from the gold center to fix an adequate steric environment and several strategies for the introduction of higher performance elements of chiral discrimination have been proposed to render carbenes the ligands of choice for asymmetric gold-catalyzed transformations.^[17] Notably, NHCtype ligands based on an imidazo[1,5-a]pyridine-3-ylidene (ImPy D) skeleton, initially disclosed by the groups of Lassaletta^[18] and Glorius^[19] have allowed the synthesis of new families of carbene ligands by the introduction of substituents at the C5 position. The latter are in close proximity to the metal-center and can be designed to convey properties to the gold complexes.^[20] For instance, the π -acceptor properties of ImPy ligands were enhanced by the introduction of a cyclophane scaffold and found applications in the catalysis of the cycloisomerization of eneallenes.^[21] The substitution at the C5 position constitutes also an opportunity to provide axially chiral complexes.^[22] Zhang further refined this approach by providing chiral bifunctional NHC ligands flanked with a tetrahydroisoquinoline moiety at the C5 position.^[23] The corresponding gold complexes proved to be competent in a series of asymmetric transformations (Figure 1).

Regarding CAACs, the presence of a quaternary carbon center adjacent to the ylidene carbon is a good lever to introduce stereogenicity close to the active site. Bertrand and coll. initially prepared an enantiopure (-)-menthol derived CAAC ligand^[7a] but no asymmetric catalysis in a copper-catalyzed asymmetric conjugated borylation reaction was recorded, [24] this being certainly due to a conformational inversion of the menthyl group. In contrast, a more rigid CAAC with extended chirality derived from a 5α -cholestan-3-one steroid backbone was recently synthesized and the corresponding copper complex ^{Cholest}CAACCuCl, promoted encouraging enantioselectivity (55% ee) in the same borylation reaction.^[24,25] It should also be noted that racemic chiral CAACAuCI complexes have been prepared by Kumar and Waldmann via a Me₂SAuCl-promoted cyclizationrearrangement sequence of 1,7-enynes and two examples of catalysis were given.[26]



Figure 1. Milestones in the development of $\mathsf{Au}(\mathsf{I})\text{-}\mathsf{carbene}$ complexes and presentation of the work

As part of our ongoing interest in the design of new chiral ligands for gold complexes^[27-29] we have devised the new family of Indolizy *C*-ligands **E**, possibly bifunctional^[30] (Figure 1). While direct isolation of these ligands has not been accomplished so far, we found more convenient to directly access to the corresponding gold complexes **2** through an efficient gold-promoted cyclization of allenepyridines of type **1** that we previously worked out.^[31,32] Two resonance forms can be written for complexes of type **2**, vinylgold vs carbene, but the very short Au-C bond distance (1.993(4) Å) on **2a**,^[33] reminiscent of other gold carbenes,^[34] suggested a carbenic character favorable for catalytic applications. Overall, the ligands **E** connected to the gold center of complexes **2** could be considered as the first

examples of vinylogous CAACs^[7] that also include chiral congeners.

Prompted by a recent report from the Muñoz group on a related use of bis-pyridyl allenes **1** ($R_1 = Me$, $R_2 = 2$ -methylpyridine) as precursors of gold carbenes and their preliminary catalytic activity,^[35] we herein extend the family of gold(I) complexes **2**, evaluate the electronic properties conferred by ligands **E** and study their catalytic applications notably in enantioselective gold(I) catalysis.

Results and Discussion

Preparation of a family of gold(I) complexes 2. We synthesized a wide range of gold(I) complexes 2 including achiral complex 2b and chiral complexes 2a and 2c-k. As depicted in Scheme 1, they were obtained in two steps from propargyl acetates 3 and 3'. S_N2' reactions on the latter provided various pyridyl-allenes 1 that were engaged in the previously mentioned cyclization reaction using 1 equiv. of Me₂SAuCl to smoothly deliver complexes 2.[31] Through this route, we could control strategic positions on chiral complexes (±)-2: i) the nature of the stereogenic center, either devoid of an additional coordinating group (2c) or flanked with a β -phosphorus group (phosphine oxide for 2a and 2d-g featuring an increasing steric demand, phosphine sulfide for 2h and phosphonate for 2i), ii) the vinyl position close to the gold(I) center by varying the aryl scaffold through the introduction of a 4,6-dimethylphenyl moiety (complexes 2d and 2f). The obtained cyclization yields for complexes 2, from quantitative for 2a-c to 34% for 2f were relatively consistent with the steric bulk imposed by the substituents of the allene moieties and gave sufficient material for catalytic trials. X-ray diffraction (XRD) analyses were obtained for complexes 2c-2f.[33] The Au-C bond lengths lie in the same range from 1.990(6) Å for 2d to 2.002(1) Å for 2f. As shown also on Scheme 1, for complexes 2 bearing a phosphine oxide, a stabilising dipole-dipole interaction between the oxygen of the phosphine oxide and the nitrogen atom of the Ncontaining 6-membered ring seems to operate with an average O-N distance of 2.95 Å.

To obtain enantiomerically pure complexes 2, we relied on a total chirality transfer in the gold-promoted cyclization.^[31] This could be accomplished from allene (-)-1a to give precatalyst catalyst (-)-(S)-2a whose absolute configuration was determined by XRD.^[36] Nevertheless, we found more convenient to obtain each enantiomer of complexes 2a and 2c-i by preparative chiral HPLC.^[37] It should be noted that thiophosphine complex 2h could not be resolved and proved unstable in the HPLC conditions. We were also able to directly obtain enantiopure indolizy-gold(I) complexes 2j and 2k using inexpensive (-)menthone from the chiral pool.^[7] In three steps including initially the addition of ethynylmagnesium bromide onto (-)-menthone,^[38] a mixture of two separable diastereoisomeric propargylic acetates 3j and 3k was obtained. Subsequent cuprate addition led to the formation of tetrasubstituted allenes as single diastereomers in an excellent yield of 97% for both (aS)-1j and (aR)-1k. Finally, the corresponding chiral gold(I) complexes 2j and 2k were obtained in respectively 52 % and 99% yield, by stereospecific reaction with Me₂SAuCI. The structure of 2k was confirmed by XRD analysis (see SI).[39]



^a See SI for details; ^bsolvent: CH₂Cl₂; ^csolvent: DCE

Scheme 1. Indolizy Au(I) precatalysts 2

Evaluation of the Electronic properties. The modification of the NHC backbones can dramatically change their electronic features.^[40] Based on this fact, we have first determined the geometries and energies of the frontier orbitals for the free ligand **Indolizy-b** of complex **2b** (see also Figure 2) by means of

DFT calculations at the B3LYP/def2-TZVP level of theory. It was found that the HOMO of **Indolizy-b** (-4.97 eV) lies considerably higher than those calculated for relevant CAACs considered as very good σ -donor ligands. This data highlights the tendency of the ylidene carbon to increase its electron-donating ability when decreasing the number of electronegative nitrogen atoms connected to it.^[41]



Figure 2. a) Relevant HOMOs and LUMOs (present on the ylidene carbon) energy levels and singlet-triplet transition energy of a series of relevant NHC including Indolizy-b calculated at the B3LYP/def2-TZVP level of theory b) Vizualisation of the frontier orbitals of Indolizy-b plotted with Gaussview at an isosurface value of 0.05 au

Intriguingly, Indolizy-b exhibits a LUMO lying at a very low energy level that is at first glance suggestive of an exceptional πacidity for this ligand, comparable to the coumaraz-2-one-4ylidene carbene.^[42] Nevertheless, π-acceptor capabilities correlate with the energies of the relevant π^* -orbitals and not necessarily the LUMO so that the sole consideration of the LUMO energy can be misleading.^[43] Visualization of the shape of the frontier orbitals allows for a qualitative estimation of their geometrical appropriateness for a possible overlap with the metals d-orbitals. While the HOMO of Indolizy-b is mainly located on the ylidene carbon, the LUMO is rather opposite to this site and mainly delocalized over the N-containing 6membered ring (Figure 2b). Interestingly, the LUMO+1 that lies slightly higher than the one calculated for a highly π -acidic CAArC^[11] (-1.30 vs -1.47), is much present on the ylidene carbon and therefore can also account for the overall π -acidity of the ligand. Finally, Indolizy-b presents a very small singlet-triplet gap (24.3 kcal/mol) smaller than the one calculated for the coumaraz-2-one-4-ylidene carbene (26.7 kcal/mol), which presages good bonding abilities.[44]

To experimentally corroborate these calculations, we first gauged the donating ability of the Indolizy ligand following the Huynh Electronic Parameter (HEP). This method, initially based on the ¹³C NMR spectroscopy analysis of trans-[PdBr₂(*i*Pr₂bimy)L] (*i*Pr2-bimy = 1,3-diisopropylbenzimidazolin-2-ylidene) complexes on which L is a NHC ligand, has also been transposed to [Au(*i*Pr₂-bimy)L] gold(I) complexes.^[45,46] While the HEP method has never been used for the evaluation of the $\sigma\text{-}$ donation of CAACs or analogues, we nevertheless synthesized the [Au(iPr2-bimy)(Indolizy-b)]BF4 complex 5 in good yield (84%) through a one-pot reaction by mixing iPr_2 -bimyH⁺BF₄⁻ (4) in the presence of K₂CO₃ and gold complex **2b** (Scheme 2). The structure of 5 was secured by XRD analysis.[47] As expected, a linear coordination pattern in which the central gold(I) atom is both coordinated to the iPr2-bimy and Indolizy-b ligands was observed. The isopropyl C-H protons of complex 5 appeared by ¹H NMR as a multiplet at 5.05 ppm, falling in the range of the expected values for this type of complexes, and which indicated a free rotation around the Au-C_{carbene} bonds.^[45] Moreover, the ¹³C NMR spectrum of 5 exhibited two carbene resonances at 193.1 ppm, presumably for the iPr2-bimy carbene and more downfield at 204.5 ppm for the Indolizy ligand. A two dimensional HMBC NMR analysis confirmed these assignments (see SI). Based on the linear correlation between the $^{13}C_{carbene}$ (*i*Pr₂-bimy) values for the Pd(II) and Au(I) complexes, a HEP value of 184.7 ppm, corresponding to the highest HEP value ever found for a carbene ligand, and much higher than the HEP of IPr (177.5 ppm), was determined for the Indolizy ligand. It therefore appears that Indolizy ligand features strong σ -donating properties which fits with the DFT calculations.



Scheme 2. Synthesis of complex 5 and its XRD structure (hydrogen atoms are omitted for clarity). Selected bond lengths (Å) and angle (deg): Au1–C27 2.035(3), Au1–C1 2.033(3), C1–Au1–C27 174.5(1)°

Next, the π -acceptor character of the Indolizy ligand was assessed by measuring the ⁷⁷Se NMR chemical shift of the selenium derivative **6**,^[48-49] prepared in 85% yield by mixing in acetonitrile elemental selenium with allene **1b** at 130°C in a sealed tube for 22h (Scheme 3).^[50] The structure was ascertained by a single crystal XRD analysis.^[51]



Scheme 3. Synthesis of Indolizy selenide derivative 6

With a δ^{77} Se NMR value of 566.7 ppm in acetone- d_6 and 528.1 ppm in CDCI₃, Indolizy ligand appears far more π -accepting than IPr (δ^{77} Se: 87.0 ppm in acetone- d_6)^[48] but slightly less π -accepting than the CAArC of Figure 2 (δ^{77} Se: 601 ppm).^[11] Interestingly this last finding is also consistent with the above-presented DFT study.

We also examined the impact of the electronic properties of the Indolizy ligand by using the gold(I) precatalyst 2a in a diagnostic reaction consisting in the cycloisomerization of allenediene 7 (Scheme 4). It was shown that in this transformation the π -activation of the allene by a cationic gold(I) catalyst triggers the generation of a cationic intermediate 8. The latter can evolve following two pathways depending on the influence of the ancillary ligand: an alkyl shift or a 1,2-hydride shift giving respectively the [4+2] cycloadduct 9 or the [4+3] cycloadduct 10.^[52] Bicyclic product 10 is a marker of a strong donor ligand which favors a carbene reactivity of intermediate 8. For instance IPr or IMes ligand will provide almost exclusively 10.[52] In contrast, hydrindane product 9 suggests a cationic pathway because of the reduced donation of the ancillary ligand. This is what happens when using π-accepting cyclophane-NHCs which yields a 72:28 ratio of 9 vs. 10. [21] In the presence of AgSbF₆ and 2a, a 74:26 mixture of cycloadducts 9 and 10 was obtained in 67% yield. While this type of ratio has to be taken cautiously since steric effects can also be at play,^[52e] the convergence between the ⁷⁷Se NMR value and this experiment suggests significant π -acidic properties of the Indolizy ligand.



Scheme 4. Diagnostic reaction using the allenediene precursor 7

Applications in asymmetric catalysis. Besides the cyloisomerization of 7, we checked the reactivity of precatalyst 2a in a few diagnostic reactions (see SI). Consistent results with literature were obtained. We also checked the configurational stability of (-)-(S)-2d in thermal conditions (80°C for 24 h, see SI). All these positive data drove us to test asymmetric catalysis with Indolizy-Au complexes 2 (except 2h)[53] We first studied the methoxycyclization of enyne 11 that has served as benchmark reaction in numerous studies.^[29] Using typical conditions (AgSbF₆ as cationizing agent in MeOH at rt), low enantioselectivities were recorded, the best resulted of 28% ee being obtained with complex (+)-(S)-2f (Table 1, entry 5). Interestingly, a precatalyst bearing a gem-dialkyl group ((+)-2c) with no additional functionality provided a racemic mixture (entry 2) while the other complexes bearing a phosphine oxide arm allowed some tangible substituted enantiomeric discrimination.

Table 1. Methoxycyclization reaction					
$Z [Au] (3 \text{ mol}\%), AgSbF_{6} (4.5 \text{ mol}\%) \\ MeOH, 25 ^{\circ}C, 20 \text{ h} \\ 11 \\ Z = C(CO_2Me)_2 \\ U = 12 $					
Entry	[Au]	Yield of 12 (%)	e.r. ^[a] (R:S) ^[b]		
1	(-)-(S)- 2a	92 ^[c]	59:41		
2	(+)- 2c	97 ^[c]	50:50		
3	(–)-(<i>S</i>)- 2d	76	54:46		
4	(-)-(<i>R</i>)- 2e	76	38:62		
5	(+)-(S)- 2f	95	64:36		

[a] Enantiomeric ratios were determined by chiral HPLC; [b] The configurations were assigned by comparison with literature.^[29]; [c] NMR yields with trimethoxybenzene as internal standard.

Based on these promising findings, we examined the purely intramolecular gold-catalyzed hydroalkoxylation of 4,5-hexadiene-1-ols **13a** and **13b** and the results are summarized in Table 2. Precatalysts **2** were first activated with a silver salt before the addition of substrates **13**. We started our studies with gold(I) complex (–)-(S)-**2a** and initially checked the influence of the silver salt as cationizing agent on the enantioselectivity of the reaction. At 25°C with AgSbF₆ in toluene (entry 1), low yield of **14a** (16%) was obtained with almost no enantioselectivity (e.r. = 45:55). Increasing the temperature to 50°C (entry 2) resulted in a yield increase (52%) but a decrease of enantioselectivity.

Keeping this temperature with other silver salts (entries 3-4), low yields of **14a** in racemic form were observed.

Table 2. Intramolecular	r hydroalkoxylation.
-------------------------	----------------------

	H	[Au] (3 mol	%), [Ag]	(4.5 mol%	%) _0 <u>*</u>	
$R_1 \rightarrow R_1$	R_2	tol	uene, T, t		$ \begin{array}{c} \xrightarrow{} R_1 \xrightarrow{} \\ R_1 \\ R_1 \\ R_2 \end{array} $	R ₂
R_2 13a , R_1 = Ph, R_2 = H 14a , R_1 = Ph, R_2 = H 13b , R_1 = H, R_2 = -C ₅ H ₁₀ - 14b , R_1 = H, R_2 = -C ₅ H ₁₀ -						
Entry	[Au]	[Ag]	T (°C)	t (h)	Yield of 14a (%)	e.r. ^[a] (<i>S:R</i>) ^[b]
1	(–)-(S)- 2a	AgSbF ₆	25	30	16	45:55
2	(+)-(<i>R</i>)- 2a	AgSbF ₆	50	16	52	49:51
3	(-)-(S)- 2a	AgOTf	50	16	28	51:49
4	(+)-(<i>R</i>)- 2a	$AgNTf_2$	50	26	28	50:50
5	(-)-(S)- 2a	AgOTs	25	5	quant.	70:30
6	(-)-(S)- 2a	AgOTs	-20	19	-	-
7	(-)-(S)- 2a	AgOTs	0	5	-	-
8	(+)-(<i>R</i>)- 2a	AgOTs	10	7	34	41:59
9	2j	AgOTs	25	2	quant.	50:50
10	2k	AgOTs	25	2	quant.	51:49
11	(+)- 2c	AgOTs	25	2	quant.	52:48
12	(+)-(<i>R</i>)- 2d	AgOTs	25	3.5	quant.	27:73
13	(-)-(<i>R</i>)- 2e	AgOTs	25	2.5	quant.	12:88
14	(-)-(<i>R</i>)- 2e	AgOTs	-20	5 j	quant.	24:76
15	(-)-(<i>R</i>)- 2e	AgOTs	50	1.5	quant.	13:87
16	(-)-(<i>R</i>)- 2f	AgOTs	25	1.5	quant.	8:92
17	(R)- 2g	AgOTs	25	1.5	quant.	27:73
18	(–)- 2i	AgOTs	25	22	quant.	57:43
Entry	[Au]	[Ag]	T (°C)	t (h)	Yield of 14b (%)	e.r. ^[a]
19 ^[c]	(-)-(S)- 2a	AgSbF ₆	25	16	81%	49:51
20	(+)-(<i>R</i>)- 2a	AgOTs	25	16	quant.	60:40
21	(+)-(<i>R</i>)- 2d	AgOTs	25	1.5	quant.	58:42
22	(-)-(<i>R</i>)- 2e	AgOTs	25	1.5	quant.	74:26
23	(+)-(S)- 2f	ΔαΟTs	25	15	quant	23.77

[a] Enantiomeric ratios were determined by chiral HPLC. [b] The configurations were assigned by comparison with literature.^[55] [c] The reaction was run in DCM.

Interestingly, with AgOTs at 25°C,^[54] tetrahydrofuran **14a** was obtained quantitatively with a promising enantiomeric ratio of 70:30 (entry 5). The absolute configuration of the (S)-major enantiomer was assigned by comparison with previous chiral HPLC data and optical rotation.^[55] From a second set of

experiments with AgOTs, we showed that the reaction does not work at temperatures below 10°C and that no enantioselectivity was observed (entries 6-8). From all these experiments, we converged to catalytic reactions at 25°C with AgOTs as chloride scavenger as in entry 5. Screening of the gold complexes brought intriguing findings. We confirmed that for complexes 2k, 2j and 2c that do not bear a phosphine oxide moiety, the yields were excellent but quasi-racemic mixtures were obtained (entries 9-11). In contrast, for complex (+)-(R)-2d bearing a phosphine oxide moiety, enantioselectivity was restored (e.r. = 27:73, entry 12). This slight improvement compared to (-)-(S)-2a could be ascribed to the larger steric hindrance in vicinity of the gold(I) center brought by the 3,5-dimethylphenyl group on the vinyl carbon. The introduction of 3,5-dimethylphenyl groups on the phosphorus atom as for (-)-(R)-2e significantly increased the enantiomeric ratio to 12:88 (entry 13). As previously shown, the decrease or increase in temperature did not improve the enantioselectivities (entries 14 and 15). Gratifyingly, combining both previous steric constrains by using (-)-(R)-2f yielded the best e.r. value of 8:92 (entry 16). This also corresponds to the best e.r. obtained on 14a with a carbene gold complex.[56],[17] Using gold(I) complex (R)-2g bearing two 3,5-di-tert-butylphenyl groups on the phosphorus atom led however to a lower enantioselectivity (entry 17), suggesting that the steric hindrance around the phosphorus atom does not seem to be the sole factor to influence the enantioinduction. Finally, substitution of the phosphine oxide moiety by a phosphonate ((-)-2i) moiety proved to be highly detrimental to the enantioselectivity (entry 18) giving a e.r. of 57:43. Allenol 13b gave similar results with each catalyst 2, the highest enantioselectivity being recorded with (+)-(S)-2f (entry 23).

The asymmetric intramolecular hydroamination of sulfonamide **15** was next examined. As shown in Table 3, pyrrolidine derivative **16** was obtained in moderate to good yields in 24 h hours at 50°C (Table 2, entries 1-6). The enantiomeric ratios followed the same trend as described for γ -allenols **13** with the same catalysts. As for **13b**, the use of precatalysts (–)-(*R*)-**2e** and (–)-(*R*)-**2f** did not make real difference in terms of enantioselectivity (entries 3 and 4).

Table 3. Intramolecular hydroamination.						
SO ₂ Mes NH 15	[Au] (3 m	bl%), AgOTs (4.5 mol%) ∽ bluene, 50°C, 24h	SO ₂ Mes			
Entry	[Au]	Yield of 16 (%)	e.r. ^[a]			
1	(+)-(<i>R</i>)- 2a	71	28:72			
2	(+)-(<i>R</i>)- 2d	74	25:75			
3	(-)-(<i>R</i>)- 2e	66	15:85			
4	(–)-(<i>R</i>)- 2f	66	17:83			
5	(S)- 2g	66	70:30			
6	(–)- 2i	48	55:45			

[a] Enantiomeric ratios were determined by chiral HPLC.

Mechanistic aspects. As discussed above, analysis of the XRD structures of most of the neutral chlorinated gold complexes 2 revealed that the privileged conformation is the one where the phosphine oxide moiety is directly pointing towards the N-containing heterocyclic ring (Scheme 1). However, the gold catalysts are effective under their cationic form after abstraction of the chloride ligand. We attempted to crystallize the cationic form of the gold complex (±)-2a (as a racemic mixture) after anion metathesis using a stoichiometric amount of silver triflimidate.^[57] The expected cationic complex (±)-17 was formed and its XRD structure^[58] revealed a dimeric complex with a Au-Au distance of 3.0255(4) Å, which is characteristic of an aurophilic interaction (Scheme 5).^[59] An intermolecular interaction between the phosphine oxide group of one monomer and the gold cationic center of the other one was observed, featuring a distance of 2.091(1) Å between the Au and O atoms. Thus, this XRD structure confirmed that the pending arm bearing the phosphine oxide group could change its conformation to set the coordination to a cationic gold center.



Scheme 5. Synthesis of the cationic catalyst ±-17 and XRD structure

DFT calculations were also performed to get insight into the mechanism of the cyclisation of a γ -allenol in order to figure out the crucial role of the phosphine oxide moiety and of the tosylate anion in the cycloisomerization process. Indeed, it appeared that the enantioselective induction was rather effective only in the case where a phosphine oxide was present on the carbene ligand in combination with a tosylate as counterion pointing out a possible participation of these two polar functional groups (see Table 2).

Although an *inner*-sphere mechanism was initially proposed by Telles in 1998 to describe the gold(I)-catalyzed addition of methanol onto alkynes,^[60] the *outer*-sphere mechanism for the addition of an O- or N-nucleophile to a gold-coordinated C–C multiple bond complex has received much more experimental and theoretical supports.^[61] Nevertheless, gold complexes with ligands bearing a judiciously positioned pendant donor function, such as a phosphine oxide^[62] or an amine,^[63] may favour a *syn*addition through an *inner*-sphere mechanism. The polar function might also play an active role in the proton transfer steps thanks to its hydrogen-bond acceptor capability.^[62] Concerning transformations involving allenes, an *inner*-sphere mechanism supported by calculations has already been put forward for the intramolecular cyclisation of γ -amino allenes when a bis-gold phosphine complex was used.^[64]

In the following DFT study, we considered the *inner*- and *outer*-sphere mechanisms for the cycloisomerization of a γ -allenol yielding to a respective *syn*- and *anti*-addition of the hydroxy group onto the gold-activated allene.^[65] As mentioned earlier, the presence of the tosylate anion is crucial and therefore its involvement has been taken into consideration. Calculations were performed with the complex (*R*)-**2a.OTs**.

Hexa-4,5-dien-1-ol 18 was used as a model to compute the formation of the (R)-vinyltetrahydrofuran product (Figure 3a). As non-polar solvents gave the best results in terms of reactivity and enantioselectivity, it can be assumed that the tosylate anion stays rather close to the gold complex during the whole substrate transformation. Thereby, we considered two starting gold-allene η^2 complexes I and I' with the tosylate anion interacting either with the metal or with the heterocyclic ring, respectively. In the latter complex I', which is slightly less stable than I by 1.5 kcal.mol⁻¹, the electron-rich oxygen of the phosphine oxide is pointing towards the electrophilic metal (d Au-O = 3.62 Å) and therefore is located in a suitable position to adopt an appropriate conformation for interactions with the coordinated substrate. A transition state for the inner-sphere mechanism involving unambiguous hydrogen bonding between the phosphine oxide and the hydroxyl group could be computed (blue pathway, Figure 3a). The optimised structure (S)-TS_{SYN}, displays an activation barrier of 14.0 kcal.mol⁻¹ as a consequence of a beneficial template effect. The nucleophilicity

of the hydroxyl group is enhanced by the proximity of the hydrogen-bond acceptor which also accounts for the lower energy barrier observed. The cyclization is endergonic by 9.8 kcal.mol⁻¹ from I' and lead to the intermediate (R)-II' where the protonated tetrahydrofuran is intramolecularly stabilised by the phosphine oxide. Reversibility in the cyclization of y-allenols due to ring constraints has already been evidenced experimentally by the groups of Gagné and Widenhoefer.[66] The next elementary step to form the desired product is the protodeauration. This step usually requires the assistance of an exogenous molecule playing the role of a proton shuttle.^{[61b],[67]} Once again, the phosphine oxide group is ideally placed to act this way and operates similarly to a molecular crane by first picking up the proton to deliver it to the reactive C-Au bond. This promotes the protodeauration and furnishes the gold(I)complexed product (R)-III'.



Figure 3. a) Gibbs energy profile for the enantioselective cycloisomerization of γ -allenol 18 in the presence of (*R*)-2a.OTs at the M06/def2-QZVP(Au)-6-311+G(2d,p)//B3LYP-D3/SDD(Au)-6-31G(d,p) level of theory (CPCM solvent=toluene). (*R*)-TS_{ANTI} is obtained from an isoenergetic isomer of I with the metal complexed to the other face of the double bond. b) Visualization of the transition states for the cylization and protodeauration steps.

The transition state (R)-TS_{II'-III'} is located 18.3 kcal.mol⁻¹ above the starting intermediate **I**'. The overall cycloisomerization process is highly exergonic with an associated energy gain of about 20 kcal.mol⁻¹ that totally drives the reaction despite the first endergonic step. In such a case, the tosylate does not play an active role in the transformation. This pathway is very similar to the one calculated with the cationic complex of (R)-2a (not taking the anion into consideration, see SI). In such a case, the rate determining step of the sequence is the protodeauration. Even though the stereogenic center is installed during the cyclisation, this internal addition is reversible and the small energy barrier to go backwards (about 4 kcal.mol⁻¹) implies a fast equilibrium. Therefore, the mechanism would follow a

Curtin-Hammett scenario in which the final protodeauration step is the actual stereochemically determining event.^[69]

However, the tosylate anion is not a weakly coordinating anion and therefore can be involved in the cyclization and the protodeauration steps (black pathway, Figure 3a). Indeed, the activation barrier to reach (*R*)-TS_{SYN} from the starting intermediate I is only 9.1 kcal.mol⁻¹. The tosylate, which is coordinated to the gold center, stabilizes the cyclization transition state through hydrogen bonding. The resulting vinylgold intermediate (*R*)-II is more stable than (*R*)-II' by 14.5 kcal.mol⁻¹ and the cyclization becomes exergonic. The *anti*addition without the assistance of the tosylate anion was also computed with a corresponding transition state (*R*)-TS_{ANTI} lying about 9 kcal.mol⁻¹ above the *syn*-addition. The protodeauration step can also be assisted by the tosylate. The transient ptoluenesulfonic acid delivers the proton to the C-Au bond through (R)-TS_{II-III} with an activation barrier of only 8.1 kcal.mol⁻¹ from (R)-II to lead to the final intermediate (R)-III. Interestingly, in this pathway with the tosylate coordinated to the gold center, the oxygen of the phosphine oxide prefers to point towards the electron-deficient N-containing heterocyclic ring (d O-N = 2.96 Å in structure I). The favored interaction brings some rigidity to the system and forces the phosphine oxide moiety to display its two aromatic substituents nearby the chiral active catalytic site, which rationalizes the influence of the aromatic rings substitution on the enantioinduction. In this pathway, the cyclization is the stereochemically-determining step. Gratifyingly, the formation of the (S)-vinyltetrahydrofuran product could be computed as well and the transition state corresponding to the cyclization step is higher than the one calculated for the (R)-product by 1.3 kcal.mol⁻¹ (see SI). Although this DFT study has been carried out with a simpler model, the results are consistent with the finding that allenol 13a is converted into (R)-14a using (R)-2a and analogues as precatalysts.

Conclusion

In summary, we have designed a new family of gold(I) complexes based on indolizine-ylidene (Indolizy) ligands. These complexes are readily available from the gold(I)-promoted cyclization of allenylpyridines. The evaluation of the electronic features of the Indolizy ligand has shown highly σ -donor and π acceptor properties. For that we have relied on a diagnostic reaction and on various NMR analyses. Among others, we used the Huynh's HEP method which had never been used for any carbene other than NHCs. All findings were supported by DFT calculations. Chiral gold(I) complexes could be separated by chiral HPLC and evaluated in asymmetric catalysis. Chiral bifunctional ligands presenting a pending phosphine oxide moiety on the stereogenic center afforded highly efficient and enantioselective cycloisomerizations of y-functionalized allene derivatives. Increasing the steric demand on the phosphine oxides promoted the best enantiomeric ratios. DFT calculations suggested several possible scenarios highlighting the crucial roles of the phosphine oxide moiety and the tosylate counterion.

Acknowledgements

The authors thank Sorbonne Université (PhD grant to TM), the Franco Thai Scholarship Program administered by CAMPUS FRANCE, IUF and CNRS for financial support. We are also grateful to Lise-Marie Chamoreau (IPCM) for the XRD data and Manon Louis for the studies related to **2h**.

Conflic of interest

The authors declare no conflict of interest.

Keywords: carbene • gold • phosphine oxide • asymmetric catalysis • DFT

- (a) Homogenous Gold Catalysis, V. Michelet, F. D. Toste, Eds., Imperial College Press: London, 2014; (b) Modern Gold Catalysis Synthesis, A. S. K. Hashmi, F. D.Toste, Eds.; Wiley, 2012.
- [2] For recent reviews, see: a) L. Fensterbank, M. Malacria, Acc. Chem.Res. 2014, 47, 953–965; b) R. Dorel, A. M. Echavarren, Chem. Rev. 2015, 115, 9028–9072; c) D. Pflästerer, A. S. K. Hashmi, Chem. Soc. Rev. 2016, 45, 1331-1367; d) R. Herrera, M. C. Gimeno, Chem. Rev. 2021, DOI: 10.1021/acs.chemrev.0c00930; e) C.C. Chintawar, A. K. Yadav, A. Kumar, S. P. Sancheti, N. T. Patil, Chem. Rev. 2021, DOI: 10.1021/acs.chemrev.0c00903; f) D. Campeau, D. F. Leon Rayo, A. Mansour, K. Muratov, F. Gagosz, Chem. Rev. 2020, DOI: 10.1021/acs.chemrev.0c00788; g) M. Mato, A. Franchina, C. Garcia-Morales, A. M. Echevarren, Chem. Rev. 2020, DOI: 10.1021/acs.chemrev.0c00697; h) R. L. Reyes, T. Iwai, M. Sawamura, Chem. Rev. 2020, DOI: 10.1021/acs.chemrev.0c00793.
- [3] Z. Lu, G. B. Hammond, B. Xu, Acc. Chem. Res. 2019, 52, 1275-1288.
- [4] W. Zi, F. D. Toste, *Chem. Soc Rev.* **2016**, *45*, 4567-4589.
- [5] a) W. A. Herrmann, Angew. Chem. Int. Ed., 2002, 41, 1290-1309, Angew. Chem. Int. Ed., 2002, 114, 1342-1363; b) S. Díez-González, N. Marion, S. P. Nolan, Chem. Rev. 2009, 109, 3612-3676; c) F. Glorius *N-Heterocyclic Carbenes in Transition Metal Catalysis, Vol. 21* (Ed.: F. Glorius), Springer-Verlag, Berlin/Heidelberg, 2007. d) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, *Nature* 2014, *510*, 485-496.
- [6] a) H. Jacobsen, A. Correa, A. Poater, C. Costabile, L. Cavallo, *Coord. Chem. Rev.* 2009, *253*, 687-703; b) D. J. Nelson, S. P. Nolan, *Chem. Soc. Rev.* 2013, *42*, 6723-6753; b) T. Dröge, F. Glorius, *Angew. Chem.* 2010, *122*, 7094-7107; *Angew. Chem. Int. Ed.* 2010, *49*, 6940-6952; c) D. Gatineau, J.-P. Goddard, V. Mansuy, L. Fensterbank, *Isr. J. Chem.* 2013, *53*, 892-900.
- [7] V. Lavallo, Y. Canac, C. Präsang, B. Donnadieu, G. Bertrand, Angew. Chem. Int. Ed. 2005, 44, 5705-5709; Angew. Chem. 2005, 117, 5851– 5855; b) M. Soleilhavoup, G. Bertrand, Acc. Chem. Res. 2015, 48, 256-266.
- [8] M. Melaimi, R. Jazzar, M. Soleihavoup, G. Bertrand, Angew. Chem. Int. Ed. 2017, 56, 10046-10068; Angew. Chem. 2017, 129,10180-10203.
- [9] R. Jazzar, M. Soleilhavoup, G. Bertrand, Chem. Rev. 2020, 120, 4141-4168.
- a) V. Lavallo, G. D. Frey, B. Donnadieu, M. Soleilhavoup, G. Bertrand, *Angew. Chem. Int. Ed.* 2008, 47, 5224–5228; *Angew. Chem.* 2008, 120, 5302-5306; b) R. Kinjo, B. Donnadieu, G. Bertrand, *Ang. Chem. Int. Ed.* 2011, 50, 5560–5563; *Ang. Chem.* 2011, 123, 5674-5677.
- [11] B. Rao, H. Tang, X. Zeng, L. Liu, M. Melaimi, G. Bertrand, Angew. Chem. Int. Ed. 2015, 54, 14915-14919; Angew. Chem. 2015, 127, 15128-15132.
- [12] D. Janssen-Müller, C. Schlepphorst, F. Glorius, Chem. Soc. Rev. 2017, 46, 4845-4854.
- [13] For review on complexes of coinage metals with chiral NHCs, see: S. Budagumpi, R. S. Keri, G. Achar, K. N. Brinda, *Adv. Synth. Catal.* 2020, 362, 970-997.
- [14] Y. Matsumoto, K. S. Selim, H. Nakanishi, K.-I. Yamada, Y. Yamamoto, K. Tomioka, *Tetrahedron Lett.* **2010**, *51*, 404-406.
- [15] D. Banerjee, A. K. Buzas, C. Besnard, E. P. Kündig, *Organometallics* 2012, *31*, 8348-8354.
- [16] B. W. Gung, M. R. Holmes, C. A. Jones, R. Ruoyu Ma, C. L. Barnes, *Tet. Lett.* **2016**, *57*, 3912-3915.
- [17] a) For a recent review on chiral NHCs see: M. Michalak, W. Kośnik, *Catalysts* **2019**, *9*, 890, doi:10.3390/catal9110890; b) for seminal work on acyclic chiral carbenes, see: S. Handa, L. M. Slaughter, *Angew. Chem. Int. Ed.* **2012**, *51*, 2912-2915; *Angew. Chem. Int. Ed.* **2012**, *124*, 2966-2969.
- [18] M. Alcarazo, S. J. Roseblade, A. R. Cowley, R. Fernández, J. M. Brown, J. M. Lassaletta, *J. Am. Chem. Soc.* **2005**, *127*, 3290-3291.
- [19] C. Burstein, C. W. Lehmann, F. Glorius, *Tetrahedron* 2005, *61*, 6207-6217.

- [20] For robust gold(I) complexes bearing a barbituric heterocycle at the C5 position, see: Y. Tang, I. Benaissa, M. Huynh, L. Vendier, N. Lugan, S. Bastin, P. Belmont, V. César, V. Michelet, *Angew. Chem.* 2019, 131,8061-8065; *Angew. Chem. Int. Ed.* 2019, 58, 7977-7981.
- [21] M. Alcarazo, S. J. Roseblade, A. R. Cowley, R. Fernández, J. M. Brown, J. M. Lassaletta, *J. Am. Chem. Soc.* **2005**, *127*, 3290-3291.
- [22] J. Francos, F. Grande-Carmona, H. Faustino, J. Iglesias-Sigüenza, E. Díez, I. Alonso, R. Fernández, J. M. Lassaletta, F. López, J. L. Mascareñas, J. Am. Chem. Soc. 2012, 134, 14322–14325.
- [23] J.-Q. Zhang, Y. Liu, X.-W. Wang, L. Zhang, Organometallics 2019, 38, 3931-3938.
- [24] D. Pichon, M. Soleilhavoup, J. Morvan, G. P. Junor, T. Vives, C. Crévisy, V. Lavallo, J.-M. Campagne, M. Mauduit, R. Jazzar, G. Bertrand, *Chem. Sci.* 2019, *10*, 7807-7811.
- [25] For Ru complexes bearing C1-symmetric cyclic CAAC ligands giving high enantioselectivities in asymmetric olefin metathesis, see: J. Morvan, F. Vermersch, Z. Zhang, L. Falivene, T. Vives, V. Dorcet, T. Roisnel, C. Crévisy, L. Cavallo, N. Vanthuyne, G. Bertrand, R. Jazzar, M. Mauduit, *J. Am. Chem. Soc.* **2020**, *142*, 19895.
- [26] F. Kolundžić, A. Murali, P. Pérez-Gálan, J. O. Bauer, C. Strohmann, K. Kumar, H. Waldmann, *Angew. Chem.* 2014,126, 8260-8264; *Angew. Chem. Int. Ed.*, 2014, 53, 8122-8126.
- [27] F. Schröder, C. Tugny, E. Salanouve, H. Clavier, L. Giordano, D. Moraleda, Y. Gimbert, V. Mouriès-Mansuy, J.-P. Goddard, L. Louis Fensterbank, Organometallics, 2014, 33, 4051-4056.
- [28] A. Vanitcha, C. Damelincourt, G. Gontard, N. Vanthuyne, V. Mouriès-Mansuy, L. Fensterbank, *Chem. Commun.* **2016**, *52*, 6785-6788.
- [29] C. Tugny, N. del Rio, M. Koohgard, N. Vanthuyne, D. Lesage, K. Bijouard, P. Zhang, J. Meijide Suárez, S. Roland, E. Derat, O. Bistri-Aslanoff, M. Sollogoub, L. Fensterbank, V. Mouriès-Mansuy, ACS Catal. 2020, 10, 5964–5972.
- [30] For a recent example of bifunctional NHC-AuCl complex, see: O. Seppänen, S. Aikonen, M. Muuronen, C. Alamillo-Ferrer, J. Burès, J. Helaja, *Chem. Commun.* **2020**, *56*, 14697-14700.
- [31] A. Vanitcha, G. Gontard, N. Vanthuyne, E. Derat, V. Mouriès-Mansuy, L. Fensterbank, Adv. Synth. Catal. 2015, 357, 2213-2218.
- [32] For recent similar approaches, see: a) H. K. Wagner, H. Wadepohl, J. Ballmann, *Chem. Science* 2021,12, 3693-3701; b) S. Arndt, M. M. Hansmann, P. Motloch, M. Rudolph, F. Rominger, A. S. K. Hashmi *Chem. Eur. J.* 2017, 23, 2542-2537; c) A. S. K. Hashmi, D. Riedel, M. Rudolph, F. Rominger, T. Oeser, Chem. Eur. J. 2012, *18*, 3827-3830.
- [33] CCDC deposition numbers: 2a, 1055996; 2c, 2064282; 2d, 2064283;
 2e, 2064284; 2f, 2064285.
- [34] a) For a review, see: Y. Wang, M. E. Muratore, A. M. Echavarren, *Chem. Eur. J.* 2015, *21*, 7332-7339; b) See also: M. Joost, L. Estevez, S. Mallet-Ladeira, K. Miqueu, A. Amgoune, D. Bourissou, *Angew. Chem. Int. Ed.* 2014, 53, 14512-14516; *Angew. Chem.* 2014, *126*, 14740-14744.
- [35] H. K. Maliszewska, D. L. Hughes, M. P. Muñoz, *Dalton Trans.* 2020, 49, 4034–4038.
- [36] CCDC deposition number for (-)-(S)-**2a**: 2081677.
- [37] See supporting information and also ref. 28.
- [38] J. K. E. T. Berton, H. Salemi, J.-L. Pirat, D. Virieux, C. V. Stevens, J. Org. Chem. 2017, 82, 12439-12446.
- [39] CCDC deposition number for **2k**: 2064286.
- [40] H. V. Huynh, Chem. Rev. 2018, 118, 9457-9492.
- [41] A. A. Tukov, A. T. Normand, M. S. Nechaev, *Dalton Trans.* 2009, 7015– 7028.
- [42] H. Song, H. Kim, E. Lee, Angew. Chem. Int. Ed. 2018, 57, 8603-8607; Angew. Chem. 2018, 130, 8739-8743.
- [43] a) D. Munz, Organometallics 2018, 37, 275-289; b) D. M. Andrada, N. Holzmann, T. Hamadi, G. Frenking, *Beilstein J. Org. Chem.* 2015, *11*, 2727-2736.
- [44] E. A. Carter, W. A. Goddard III, J. Phys. Chem. 1986, 90, 998-1001.
- [45] S. Guo, H. Sivaram, D. Dan Yuan, H. V. Huynh, Organometallics 2013, 32, 3685-3696.
- [46] Q. Teng, H. V. Huynh, *Dalton Trans.* **2017**, *46*, 614-627.
- [47] CCDC deposition number for 6: 2064288.

- [48] A. Liske, K. Verlinden, H. Buhl, K. Schaper, C. Ganter, Organometallics 2013, 32, 5269-5272.
- [49] S. V. C. Vummaleti, D. J. Nelson, A. Poater, A. Gomez-Suarez, D. B. Cordes, A.M. Z. Slawin, S. P. Nolan, L. Cavallo, *Chem. Sci.* 2015, *6*, 1895-1904.
- [50] For the reaction of related allenyl pyridines with I₂, see: T. Martinez, I. Alahyen, G. Lemière, V. Mouriès-Mansuy, L. Fensterbank *Org. Process. Res. Dev.* 2020, 24, 817-821.
- [51] CCDC deposition number for 6: 2064288.
- [52] a) B. Trillo, F. Lopez, S. Montserrat, G. Ujaque, L. Castedo, A. Lledos, J. L. Mascarenas, *Chem. Eur. J.* 2009, 15, 3336–3339; b) I. Alonso, B. Trillo, F. Lopez, S. Montserrat, G. Ujaque, L. Castedo, A. Lledos, J. L. Mascarenas, *J. Am. Chem. Soc.* 2009, 131, 13020–13030; c) P. Mauleon, R. M. Zeldin, A. Z. Gonzalez, F. D. Toste, *J. Am. Chem. Soc.* 2009, 131, 6348–6348; d) J. Petuskova, H. Bruns, M. Alcarazo, *Angew. Chem. Int. Ed.* 2011, *50*, 3799–3802; *Angew. Chem. Int. Ed.* 2011, *123*, 3883-3886; e) A. H. Christian, Z. L. Niemeyer, M. S. Sigman, F. D. Toste, *ACS Catal.* 2017, *7*, 3973-3978.
- [53] It should be noted that thiophosphine complex 2h is not stable in the catalytic reaction conditions.
- [54] Z. Zhang, R. A. Widenhoefer, Angew. Chem. Int. Ed. 2007, 46, 283-285; Angew. Chem. 2007, 119, 287-289.
- [55] a) Z. Wang, C. Nicolini, C. Hervieu, Y.-F. Wong, G. Zanoni, L. Zhang J. *Am. Chem. Soc.* 2017, 139, 16064-16067; b) J. L. Arbour, H. S. Rzepa, J. Contreras-García, L. A. Adrio, E. M. Barreiro, K. K. (Mimi) Hii *Chem. Eur. J.* 2012, *18*, 11317-11324.
- [56] C. Bartomolé, D. Garcia-Cuadrado, Z. Ramiro, P. Espinet *Inorg. Chem.* 2010, 49, 9758–9764.
- [57] N. Mézailles, L. Ricard, F. Gagosz, Org. Lett. 2005, 7, 4133-4136.
- [58] CCDC deposition number for **17**, 2064289.
- [59] H. Schmidbaur, A. Schier, Chem. Soc. Rev. 2008, 37, 1931-1951.
- [60] J. H. Teles, S. Brode, M. Chabanas, Angew. Chem., Int. Ed. 1998, 37, 1415-1418; Angew. Chem. 1998,
- [61] a) Z. Zhang, R. A. Widenhoefer, Angew. Chem. 2007, 119, 287-292; Angew. Chem. Int. Ed. 2007, 46, 283 -285; b) R. S. Paton, F. Maseras, Org. Lett. 2009, 11, 2237-2240; c) Z. J. Wang, D. Benitez, E. Tkatchouk, W. A. Goddard III, F. D. Toste, J. Am. Chem. Soc. 2010, 132, 13064-13071; d) T. J. Brown, D. Weber, M. R. Gagné, R. A. Widenhoefer, J. Am. Chem. Soc. 2012, 134, 9134-9137; e) R. J. Harris, R. G. Carden, A. N. Duncan, R. A. Widenhoefer, ACS Catal. 2018, 8, 8941-8952.
- [62] C. Hahn, L. Cruz, A. Villalobos, L. Garzaa, S. Adeosun, *Dalton Trans.* 2014, 43, 16300-16309.
- [63] K. D. Hesp, M. Stradiotto, J. Am. Chem. Soc. 2010, 132, 18026-18029.
- [64] J. H. Kim, S.-W. Park, S. R. Park, S. Lee, E. J. Kang, Chem. Asian J. 2011, 6, 1982-1986.
- [65] a) E. Soriano, I. Fernández, *Chem. Soc. Rev.*, **2014**, *43*, 3041-3105; b)
 M. Malacria, L.Fensterbank, V. Gandon, *Top. Curr. Chem.* **2011**, *302*, 157-182.
- [66] T. J. Brown, D. Weber, M. R. Gagné, R. A. Widenhoefer, J. Am. Chem. Soc. 2012, 134, 9134–9137.
- [67] a) C. M. Krauter, A. S. K. Hashmi, M. Pernpointner, *ChemCatChem* **2010**, *2*, 1226-1230. b) Ciancaleoni, G.; Belpassi, L.; Zuccaccia, D.; Tarantelli, F.; Belanzoni, P. ACS Catal. **2015**, *5*, 803–814.
- [68] W. Wang, G. B. Hammond, B. Xu, J. Am. Chem. Soc. 2012, 134, 5697-5705.
- [69] R. E. M. Brooner, T. J. Brown, M. A. Chee, R. A. Widenhoefer, Organometallics 2016, 35, 2014–2021.

Entry for the Table of Contents



Au(I) complexes coordinated to the Indolizy ligand, a new carbene ligand that results from the gold-promoted cyclization of a pyridylallene were investigated. The electronic properties of the Indolizy ligand were gauged thanks to physical analyses and DFT calculations. The introduction of stereogenic center in α -position to the carbene center coupled to the introduction of a pending phosphine oxide group affords highly efficient and enantioselective Au(I) catalysts.

Institute and/or researcher Twitter usernames: @IPCM_Sorbonne