

**MtExpress, a Comprehensive and Curated RNAseq-based Gene Expression Atlas for the Model Legume *Medicago truncatula***

Running head: MtExpress database

Corresponding author: S. Carrere, LIPME, Université de Toulouse, INRAE, CNRS, Castanet-Tolosan, France , [sebastien.carrere@inrae.fr](mailto:sebastien.carrere@inrae.fr)

Subject areas: regulation of gene expression; genomics

Number of colour figures: 3

Code de champ modifié

**MtExpress, a Comprehensive and Curated RNAseq-based Gene Expression Atlas for the Model Legume  
*Medicago truncatula***

Running head: MtExpress database

Sebastien Carrere<sup>\*1</sup>, Jerome Verdier<sup>2</sup> and Pascal Gamas<sup>1</sup>

<sup>1</sup> LIPME, Université de Toulouse, INRAE, CNRS, Castanet-Tolosan, France

<sup>2</sup> Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, F-49000 Angers, France

\*Corresponding author E-mail: [Sebastien.Carrere@inrae.fr](mailto:Sebastien.Carrere@inrae.fr)

## Abstract

Although RNA sequencing has been becoming the main transcriptomic approach in the model legume *Medicago truncatula*, there is currently no genome-wide gene expression atlas covering the whole set of RNAseq data published for this species. Nowadays, such tool is highly valuable to provide a global view of gene expression in a wide range of conditions and tissues/organs.

Here, we present MtExpress, a gene expression atlas that compiles an exhaustive set of published *M. truncatula* RNAseq data (<https://medicago.toulouse.inrae.fr/MtExpress>). MtExpress makes use of recent releases of *M. truncatula* genome sequence and annotation, as well as up-to-date tools to perform mapping, quality control, statistical analysis and normalization of RNAseq data. MtExpress combines semi-automated pipelines with manual re-labelling and organization of samples, to produce an attractive and user-friendly interface, fully integrated with other available Medicago genomic resources. Importantly, MtExpress is highly flexible, in terms of both queries, *e.g.* allowing searches with gene names and orthologous gene IDs from Arabidopsis and other legume species, and outputs, to customize visualization and redirect gene study to relevant Medicago webservers.

Thanks to its semi-automated pipeline, MtExpress will be frequently updated to follow the rapid pace of *M. truncatula* RNAseq data publications, as well as the constant improvement of genome annotation. MtExpress also hosts legacy GeneChip expression data originally stored in the Medicago Gene Expression Atlas (MtGEA), as a very valuable and complementary resource.

## Keywords

RNAseq, SRA, TMM, Medicago, legumes, atlas, GeneChip

Code de champ modifié

## Introduction

Legumes represent a plant family of major importance for food, feed and the agro-ecosystem, thanks to their nutritional qualities and their ability to grow efficiently in the absence of nitrogen fertilizers due to their endosymbioses with soil microorganisms, notably with nitrogen-fixing soil bacteria (rhizobia). Indeed, legumes are becoming a keystone in tomorrow's agriculture to ensure food security, sustainable practices and mitigation of greenhouse effects. *Medicago truncatula* was proposed as a model legume about thirty years ago, and since then it has been widely used to study a wide range of biological issues relevant for plant biology and legume crop improvement. Its model legume status has led to the development of important resources for genetics, reverse genetics, structural and functional genomics (Kang et al. 2016). This notably included a gene expression atlas (MtGEA; <https://mtgea.noble.org/v3>) (Benedito et al. 2008) based upon GeneChip technology developed by Affymetrix. This atlas integrated 270 different samples (not considering biological replicates) in its last release and has been extensively used by the Medicago community. Medicago GeneChip data are also displayed with other visualization tools available from the Medicago ePlant BAR website (Waese et al. 2017) ([http://bar.utoronto.ca/eplant\\_medicago/](http://bar.utoronto.ca/eplant_medicago/)). Unfortunately, the MtGEA website does not provide expression profiles anymore, due to technical issues. Moreover, this resource was built upon first genome versions and only represents 45% of the protein-coding and long non-coding RNA genes of Mt5.1.7 genome release/annotation.

In the past years, RNA sequencing has become increasingly popular to study gene expression at the plant, organ and tissue levels. The advantages of this approach are its sensitivity and its flexibility to exploit new genome releases and annotations by the possibility of (re)mapping sequencing reads, as well as to discover transcript isoforms or novel transcripts such as long non-coding RNAs. Two databases (db) give access to some *M. truncatula* RNAseq data, the first one as part of a generic db developed by EBI on many species (<https://www.ebi.ac.uk/gxa/plant/experiments>), and the second one as part of a specific study dedicated to nutrient- and nodulation- responsive peptides (<http://mtsspdb.noble.org/atlas-internal/3880/transcript/profile/0>) (de Bang et al. 2017). However, these db only show a small selection of publicly available RNAseq data. Here we present MtExpress, a gene expression atlas based upon *M. truncatula* RNAseq data supported by peer-reviewed publications (<https://medicago.toulouse.inrae.fr/MtExpress>). We ensured that MtExpress was a comprehensive, reliable, user-friendly, interconnected and upgradeable database, by combination of automated analyses and manual curation.

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

## Results

### **Retrieval, quality checking and organization of RNAseq data**

To build MtExpress, RNAseq data (*i.e.* sequenced reads, fastq) generated from Medicago were downloaded from the Sequence Read Archive (SRA) and the European Nucleotide Archive (ENA) (Fig. 1), along with associated metadata. In order to ensure that MtExpress data are reliable and well documented, only sequencing data associated with peer-reviewed publications were conserved, representing 431 different samples (1374 if considering biological replicates) (supplemental data set 1). The nf-core/rnaseq bioinformatic pipeline version 3.0 (Patel et al. 2020) was used to automatically analyze the RNAseq libraries. First, FastQC/MultiQC were used to perform quality checking and reporting of sequencing data. Sequencing reads were, then, mapped to the Medicago genome version 5.1.7 (Pecrix et al. 2018) (52,723 genes) and reference genomes of Medicago symbionts or pathogens used in the selected SRA projects (listed in supplemental table 1), using STAR mapper (Dobin et al. 2013). Finally, SALMON (Patro et al. 2017) was used to quantify read counts of transcripts. To enable cross-comparisons between different experiments, we normalized the complete dataset by the trimmed mean of M-values method (TMM) (Robinson and Oshlack 2010), using the EdgeR package. All samples were then manually labelled using a coherent nomenclature and organized into different categories based on organs or physiological conditions (18 categories in this first release; *e.g.* leaf, seed, abiotic factors, biotic stress, root nodule symbiosis...; supplemental data set 1), with keywords attributed to each experiment, thereby enabling rapid search/selection.

### **A flexible and user-friendly interface**

Thanks to synonymy links previously computed in the LeGOO knowledge base (Carrère et al. 2019), MtExpress can be queried by any gene locus identifier from the different *M. truncatula* genome releases (from Mt3.0 to Mt5.0), and identifiers of Mt16kOLI1 microarrays (Kuster et al. 2004) or Affymetrix (Benedito et al. 2008) and Nimblegen (Verdier et al. 2013) gene chips. Moreover, we extended this query flexibility to gene acronyms, based on a list manually extracted from publications, available on the Mt5.0 genome browser <https://medicago.toulouse.inrae.fr/MtrunA17r5.0-ANR/>. Furthermore, gene IDs from *Arabidopsis thaliana* (*A. thaliana* Col-0 [Araport11]) (Cheng et al. 2017) and four major legumes, namely soybean (*Glycine max* Williams 82 [Phytozome.12]) (Schmutz et al. 2010), *Lotus japonicus* (*L. japonicus* Miyakojima MG-20 [3.0] (Li et al. 2020)), common bean (*Phaseolus vulgaris* BAT93 [EnsemblPlants.38] (Schmutz et al. 2014)) and pea (*Pisum sativum* cv.

Code de champ modifié

'Caméor' v1a (Kreplak et al. 2019), can be used to query the expression pattern of putative orthologues in *M. truncatula* (Carrère et al. 2019).

The query output (Fig. 2) was set up to display by default the expression profile of the queried gene using a 'reference dataset', containing 28 samples which provide an overview of gene expression in various plant organs as well as symbiotic interactions. Then, the user can switch to one of the 18 pre-arranged categories, each with a more extensive or specialized set of samples: e.g. 'abiotic factors' and 'biotic stress' categories with all RNAseq samples related to different stresses, or 'seed' and 'nodule' categories, dedicated to samples specific to these organs. If needed, the complete dataset containing the 431 samples can also be selected for visualization. Finally, custom-made selections can also be defined by the user by selecting/unselecting specific experiments to visualize and export the desired graphical outputs. All of the information associated to each experiment is provided, including the MultiQC (Ewels et al. 2016) report, and the corresponding publication, metadata and raw counts. Gene expression levels (with a choice between log<sub>2</sub> TMM or raw TMM values) are displayed as box plots, with various informative elements (notably project name from SRA and descriptive statistics from biological replicates such as mean, median, min, max, Q1, Q3), each project being distinguished by a different color. Information about the whole dataset is also available via the 'metadata' button, including global statistics, used reference genomes and tools. Additional information on queried genes is easily accessible via the 'synonymous' and 'annotation' buttons, as well as links to the Mt5.0 genome browser and the LeGOO knowledge database. Conversely, a link to MtExpress is provided for all annotated genes in the Mt5.0 genome browser.

As case studies, we examined the expression profile of various gene markers described to be specifically expressed in certain organs, tissues, or conditions. We found that their expression patterns were as expected from the literature, thereby supporting our automated processes. Three of them are shown in Fig. 3, using the reference sample set: *MtPT4* (*PHOSPHATE TRANSPORTER 4*) (Harrison et al. 2002) (arbuscular mycorrhizal symbiosis: arbuscule-containing cells), *MtNF-YA1* (*NUCLEAR FACTOR YA1*) (Combier et al. 2006) (root nodule symbiosis: response to bacterial Nod factors, rhizobium infection, nodule inception and apical nodule zones), *MtDME* (*DEMETER*) (Satgé et al. 2016) (endosperm and nodule differentiation zone).

### **Retrieval and presentation of MtGEA GeneChip data**

As an additional tool to MtExpress, we downloaded publicly available Medicago Affymetrix GeneChip data (supplemental data set 2), previously stored in the Medicago Gene Expression Atlas (MtGEA) site hosted by the Noble Research Institute at <https://mtgea.noble.org/v3>. We implemented these data into the MtExpress webserver to provide an alternative solution to the inoperative MtGEA webserver. Moreover, we added features of MtExpress, such as the flexible input queries and pre-defined or custom-made categories. This alternative resource is also available at <https://medicago.toulouse.inrae.fr/MtExpress>.

Code de champ modifié

### **Discussion**

MtExpress benefits from several important assets: (i) the use of recent releases of genome sequences/annotations to provide an up-to-date tool; (ii) semi-automated analysis pipelines allowing frequent updates (at least twice a year, depending on the number of new RNAseq data publications and possible new annotation release); (iii) manual re-labelling and organization of samples to conserve an unified structure through the updates; (iv) an attractive and flexible interface; (v) a transcriptomic platform fully integrated with other Medicago tools such as the Medicago Mt5.0 genome browser (Pecrix et al. 2018) and the LeGOO knowledge base (Carrère et al. 2019). The combination of manual curation with automated quality control and statistical analyses is key to ensure the reliability of MtExpress. Furthermore, the immediate access to accompanying metadata and publications represent an obvious gain of time and added value for users to better evaluate the data quality.

We therefore believe that this user-friendly, exhaustive and integrated resource will be of great use to the Medicago and, widely, to the legume community to quickly obtain the expression profiles of genes of interest in a variety of organs, tissues and conditions. This is, for instance, essential to define gene regulation patterns and determine gene expression specificity of developmental stage(s) or plant responses to biotic or abiotic environments. Considering the current pace of *M. truncatula* RNAseq data production and publication, we anticipate that the number and diversity of sequenced RNA samples will continue to grow rapidly. MtExpress will be updated accordingly, with creation of additional categories if needed. In the near future, we also plan to implement analytical tools such as gene co-expression analysis and pair-wise comparisons between experiments.

## Material and Methods

### Collecting Data

*RNAseq dataset.* We collected all data available as of April 1st 2021 in the Sequence Read Archive (SRA), using the Run Selector with the query "Medicago truncatula [ORG]" and the filter "AssayType = RNA-seq". We kept all studies associated with a peer-reviewed publication, representing 51 projects altogether (supplemental data set 1). All metadata coming from SRA, ENA and PubMed were associated to each selected project and stored for downstream treatment under a GIT repository [<https://lipm-gitlab.toulouse.inra.fr/EXPRESSION-ATLAS/ExpressionAtlasData>]. Fastq files were directly downloaded from EBI SRA FTP mirror (<ftp.sra.ebi.ac.uk>) and listed in csv files with curated sample names, replicates and library strandedness to be properly processed.

*Affymetrix GeneChip MtGEA dataset.* We exported experiment replicates for all probesets (including Mtr, Msa, Sme and control probesets) from <https://mtgea.noble.org/v3/experiments.php> downloading webpage (Benedito et al. 2008). We excluded eight duplicated libraries as well as seven libraries for which biological replicate data were not available, and added one missing set of published data (supplemental data set 2). All samples were manually ordered, classified into 14 categories (supplemental data set 2) and aliases were added to homogenize labelling. Metadata such as publication, organ, experimental factors were manually extracted from the MtGEA website.

### Expression Measure of RNAseq dataset

nf-core/rnaseq version 3.0 (Patel et al. 2020) was used with default parameters to run expression measure separately on each SRA project (Fig. 1). We chose STAR (Dobin et al. 2013) as a read mapper with 20 multiple alignments allowed for one read, and SALMON as a read quantifier (Patro et al. 2017) (see supplemental data set 3 for software versions). Of note, SALMON can deal with multi-mapped reads, that can occur with *M. truncatula* gene duplications, using an expectation maximisation (EM) algorithm to attribute read counts to the most likely transcripts (Deschamps-Francoeur, 2020). We used the version 5.1.7 of *M. truncatula* A17 genome annotation (Pecrix et al. 2018) to map RNAseq data from different accessions of *M. truncatula*, as well as reference genomes of Medicago symbionts or pathogens used in the selected SRA Studies (*Sinorhizobium meliloti* 2011, *Sinorhizobium medicae* WSM419, *Fusarium oxysporum* f. sp. medicaginis, *Rhizophagus irregularis*, *Ralstonia pseudosolanacearum* GMI1000, *Verticillium albo-atrum* V31-2, *Gigaspora rosea* and *Rhizoctonia solani* AG-3)



(supplemental table 1). Using quality control reports and statistical analyses provided by MultiQC (Ewels et al. 2016), we defined strandedness for each library and verified the consistency of biological replicates.

#### *Normalization*

*RNAseq dataset.* The complete dataset (51 SRA projects representing 431 samples) was normalized together in order to provide consistent values across gene expression comparisons in sub-categories, using TMM normalization from EdgeR package (Robinson and Oshlack 2010).

*Affymetrix GeneChip MtGEA dataset.* The expression data downloaded from the MtGEA site website were already normalized using a robust multichip average method (RMA normalization).

#### *Consolidation*

For RNAseq and GeneChip datasets, we added functional annotations provided by an automatic pipeline (Pecrix et al. 2018) as well as synonymy links to identifiers of different generations of *M. truncatula* microarrays, GeneChips and genome sequences/annotations (Carrère et al. 2019). We also used a manually curated list of 3,760 gene names/acronyms and products (Gene acronyms and IDs, release 20210409; <https://medicago.toulouse.inra.fr/MtrunA17r5.0-ANR/>). Normalized expression values (TMM or RMA) from biological replicates were summarized using descriptive statistics (mean, median, min, max, Q1, Q3) using ad-hoc Scala/Spark scripts to ensure future scalability. All these information were merged with dataset metadata to build a consistent database.

#### *Database*

The database consists of a repository of JSON and PARQUET files. Each dataset is documented through a JSON definition file along with run definition files including information related to reference genome and tools versions used for mapping. Database is accessed using Drill software (<https://drill.apache.org/>) version 1.16.0. Database structure and scheme are available at <https://lipm-gitlab.toulouse.inra.fr/EXPRESSION-ATLAS>

Code de champ modifié

#### *API*

All normalized data are accessible through a REST API developed using the Scalatra framework and exposed with Jetty servlet container (supplemental data set 3). The API is fully documented using Swagger 2.0. Documentation can be found at: <https://lipm-browsers.toulouse.inra.fr/expression-atlas-api>

Code de champ modifié

### *User Interface*

The user interface has been developed using the Vue.js library (v. 2.6.12) with a set of components, and plugins (see supplemental data set 3), notably Chart.js for boxplot display. It aims at providing simple functionalities such as gene searching through different accession identifiers (based on synonymous computed across different generations of genome assemblies/annotations and arrays), gene names or functions. Expression patterns are displayed using box-plots to show the level of variability between biological replicates. Users can easily select/unselect SRA projects to be plotted, navigate through different subsets of projects or locus and display raw or log2 normalized expression values, access to gene annotations, synonymous and links to relevant external resources (genome browser, knowledge base, original data repository). For each SRA project, links to MultiQC (Ewels et al. 2016) report, raw count and metadata files are provided.

### *Availability*

Project name: MtExpress

Project home page: <https://medicago.toulouse.inrae.fr/MtExpress>

Source Code and Data: <https://lipm-gitlab.toulouse.inra.fr/EXPRESSION-ATLAS>

Operating system(s): Platform independent

Programming languages: Scala, R, Spark, Javascript, Perl

Other requirements: Drill 1.16, Apache web server

License: Apache License 2.0

### **Data Availability**

The data underlying this article are publicly available in SRA (<https://trace.ncbi.nlm.nih.gov/Traces/sra/>) and MtGEA (<https://mtgea.noble.org/v3/>) databases.

### **Funding**

This work was supported by the Agence Nationale de la Recherche [EPISYM ANR-15-CE20-0002 and DESWITCH ANR-19-CE20-0027], Laboratoire d'Excellence TULIP ANR-10-LABX-41]; and Plant2Pro<sup>®</sup> Carnot Institute [agreement #18-CARN-024-01 – 2018].

Code de champ modifié

Code de champ modifié

## Disclosures

Conflicts of interest: No conflicts of interest declared

## Acknowledgments

We thank Jérôme Gouzy and Marie-Françoise Jardinaud (LIPME, Castanet-Tolosan) for valuable discussions on the MtExpress set up and Nathalie Vialaneix (MIAT, Castanet-Tolosan) for her advice on statistical analyses. We also thank Erika Sallet, Ludovic Cottret, Ludovic Legrand and Léo Géré (LIPME) for technical advice. Finally, we are grateful to the Genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, <https://doi.org/10.15454/1.5572369328961167E12>) for providing computing resources.

## References

- Benedito, V.A., Torres-Jerez, I., Murray, J.D., Andriankaja, A., Allen, S., Kakar, K., et al. (2008) A gene expression atlas of the model legume *Medicago truncatula*. *Plant J* 55: 504-513.
- Carrère, S., Verdenaud, M., Gough, C., Gouzy, J. and Gamas, P. (2019) LeGOO: An Expertized Knowledge Database for the Model Legume *Medicago truncatula*. *Plant Cell Physiol.* 61: 203–211.
- Cheng, C.Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S. and Town, C.D. (2017) Araport11: a complete reannotation of the *Arabidopsis thaliana* reference genome. *Plant J* 89: 789-804.
- Combiér, J.P., Frugier, F., de Billy, F., Boualem, A., El-Yahyaoui, F., Moreau, S., et al. (2006) MthAP2-1 is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula*. *Genes Dev* 20: 3084-3088.
- de Bang, T.C., Lundquist, P.K., Dai, X., Boschiero, C., Zhuang, Z., Pant, P., et al. (2017) Genome-Wide Identification of *Medicago* Peptides Involved in Macronutrient Responses and Nodulation. *Plant Physiol* 175: 1669-1689.
- Deschamps-Francoeur, G., Simoneau, J. and Scott, M.S. (2020) Handling multi-mapped reads in RNA-seq. *Comput Struct Biotechnol J* 18: 1569-1576.
- Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21.

Ewels, P., Magnusson, M., Lundin, S. and Kaller, M. (2016) MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 32: 3047-3048.

Harrison, M.J., Dewbre, G.R. and Liu, J. (2002) A phosphate transporter from *Medicago truncatula* involved in the acquisition of phosphate released by arbuscular mycorrhizal fungi. *Plant Cell* 14: 2413-2429.

Kang, Y., Li, M., Sinharoy, S. and Verdier, J. (2016) A Snapshot of Functional Genetic Studies in *Medicago truncatula*. *Front Plant Sci* 7: 1175.

Kreplak, J., Madoui, M.A., Capal, P., Novak, P., Labadie, K., Aubert, G., et al. (2019) A reference genome for pea provides insight into legume genome evolution. *Nat Genet* 51: 1411-1422.

Kuster, H., Hohnjec, N., Krajinski, F., El Yahyaoui, F., Manthey, K., Gouzy, J., et al. (2004) Construction and validation of cDNA-based Mt6k-RIT macro- and microarrays to explore root endosymbioses in the model legume *Medicago truncatula*. *Journal of Biotechnology* 108: 95-113.

Li, H., Jiang, F., Wu, P., Wang, K. and Cao, Y. (2020) A High-Quality Genome Sequence of Model Legume *Lotus japonicus* (MG-20) Provides Insights into the Evolution of Root Nodule Symbiosis. *Genes (Basel)* 11.

Patel, H., Ewels, P., Peltzer, A., Hammarén, R., Botvinnik, O. and Sturm, G. (2020) nf-core/rnaseq: nf-core/rnaseq v3.0 - Silver Shark. Zenodo.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. and Kingsford, C. (2017) Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* 14: 417-419.

Pecrix, Y., Staton, S.E., Sallet, E., Lelandais-Briere, C., Moreau, S., Carrere, S., et al. (2018) Whole-genome landscape of *Medicago truncatula* symbiotic genes. *Nat Plants* 4: 1017-1025.

Robinson, M.D. and Oshlack, A. (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* 11: R25.

Satgé, C., Moreau, S., Sallet, E., Lefort, G., Auriac, M.C., Remblière, C., et al. (2016) Reprogramming of DNA methylation is critical for nodule development in *Medicago truncatula*. *Nat Plants* 2: 16166.

Schmutz, J., Cannon, S.B., Schlueter, J., Ma, J., Mitros, T., Nelson, W., et al. (2010) Genome sequence of the palaeopolyploid soybean. *Nature* 463: 178-183.

Schmutz, J., McClean, P.E., Mamidi, S., Wu, G.A., Cannon, S.B., Grimwood, J., et al. (2014) A reference genome for common bean and genome-wide analysis of dual domestications. *Nat Genet* 46: 707-713.

Verdier, J., Lalanne, D., Pelletier, S., Torres-Jerez, I., Righetti, K., Bandyopadhyay, K., et al. (2013) A regulatory network-based approach dissects late maturation processes related to the acquisition of desiccation tolerance and longevity of *Medicago truncatula* seeds. *Plant Physiol* 163: 757-774.

Waese, J., Fan, J., Pasha, A., Yu, H., Fucile, G., Shi, R., et al. (2017) ePlant: Visualizing and Exploring Multiple Levels of Data for Hypothesis Generation in Plant Biology. *Plant Cell* 29: 1806-1821.

### Figures and Tables

Figure 1. RNAseq analysis pipeline used to build MtExpress. The first step consists in collecting and curating metadata (sample and project descriptions, experiment design) and raw data (fastq files) from public databases. Then these data are processed for quality controls and expression measure using the nf-core/rnaseq pipeline. Once all datasets are processed, count tables are merged and values are normalized with the TMM method from EdgeR package. Normalized values, gene annotations and datasets metadata are consolidated into a database made of parquet files accessible through an Application Programming Interface (API) to feed the web interface.

Figure 2. Interface of MtExpress, a user-friendly gene expression atlas. (A) drop-down menu for selecting dataset or sample category. (B) query boxes, using either Mt5.0 locus IDs, keywords, gene IDs from other genome releases or gene acronyms. (C) example of a graphical output using the reference dataset, with color-specific box plots per SRA project showing log<sub>2</sub>(TMM) values; buttons to switch between Raw and log<sub>2</sub> TMM values and to display metadata, synonymous IDs and annotation information are indicated next to links to the Mt5.0 genome portal and the Legoo knowledge base. (D) list of plotted SRA projects with associated metadata and checkboxes to select/ unselect projects/samples.

Figure 3. Expression profile of three genes with specific expression patterns. (A) *MtPT4* (*PHOSPHATE TRANSPORTER 4*), expressed in arbuscule containing cells of mycorrhizal roots. (B) *MtNF-YA1* (*NUCLEAR FACTOR YA1*), expressed in bacterial Nod factor-treated epidermal root cells, in Rhizobium-inoculated root cells, and in nodule apical zones (zone 1, distal zone 2 and proximal zone 2). (C) *MtDME* (*DEMETER*), expressed in seed endosperm and in the nodule differentiation zones (proximal zone and interzone).

Supplemental table 1. Reference genomes used for mapping

Supplemental data set 1. RNAseq data

Supplemental data set 2. GeneChip data

Supplemental data set 3. Software, API, User interface