



HAL
open science

Ram pressure candidates in UNIONS

Ian Roberts, Laura Parker, Stephen Gwyn, Michael Hudson, Raymond Carlberg, Alan Mcconnachie, Jean-Charles Cuillandre, Kenneth Chambers, Pierre-Alain Duc, Hisanori Furusawa, et al.

► **To cite this version:**

Ian Roberts, Laura Parker, Stephen Gwyn, Michael Hudson, Raymond Carlberg, et al.. Ram pressure candidates in UNIONS. Monthly Notices of the Royal Astronomical Society, 2021, 10.1093/mnras/stab3101 . hal-03421079

HAL Id: hal-03421079

<https://hal.science/hal-03421079>

Submitted on 6 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ram pressure candidates in UNIONS

Ian D. Roberts¹,^{*} Laura C. Parker², Stephen Gwyn³, Michael J. Hudson^{4,5,6}, Raymond Carlberg,⁷ Alan McConnachie,³ Jean-Charles Cuillandre,⁸ Kenneth C. Chambers,⁹ Pierre-Alain Duc,¹⁰ Hisanori Furusawa¹¹, Raphael Gavazzi,¹² Vanessa Hill,¹³ Mark E. Huber⁹, Rodrigo Ibata,¹⁰ Martin Kilbinger¹⁴, Simona Mei,¹⁴ Yannick Mellier,¹⁵ Satoshi Miyazaki,¹¹ Masamune Oguri^{16,17,18} and Richard J. Wainscoat⁹

¹Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands

²Department of Physics and Astronomy, McMaster University, Hamilton, ON L8S 4M1, Canada

³NRC Herzberg Astronomy and Astrophysics, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

⁴Department of Physics & Astronomy, University of Waterloo, Waterloo, ON N2L 3G1, Canada

⁵Waterloo Centre for Astrophysics, University of Waterloo, 200 University Ave W, Waterloo, ON N2L 3G1, Canada

⁶Perimeter Institute for Theoretical Physics, 31 Caroline St North, Waterloo, ON N2L 2Y5, Canada

⁷Department of Astronomy and Astrophysics, University of Toronto, Toronto, ON M5S 3H4, Canada

⁸AIM, CEA, CNRS, Université Paris-Saclay, Université de Paris, F-91191 Gif-sur-Yvette, France

⁹Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

¹⁰Université de Strasbourg, CNRS, Observatoire astronomique de Strasbourg, UMR 7550, F-67000 Strasbourg, France

¹¹National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan

¹²Sorbonne Université, CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98bis Bd Arago, F-75014 Paris, France

¹³Université Côte d'Azur, OCA, CNRS, Observatoire de la Côte d'Azur, CNRS, Bd de l'Observatoire, CS 34229, 06304 Nice cedex 4, France

¹⁴Université de Paris, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France

¹⁵Institut d'Astrophysique de Paris, UMR 7095 CNRS, Université Pierre et Marie Curie, 98bis boulevard Arago, F-75014 Paris, France

¹⁶Kavli Institute for the Physics and Mathematics of the Universe (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

¹⁷Department of Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

¹⁸Research Center for the Early Universe, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Accepted 2021 October 22. Received 2021 October 22; in original form 2021 March 26

ABSTRACT

We present a search for disturbed, candidate ram pressure stripping galaxies across more than 50 spectroscopically selected SDSS groups and clusters. 48 ram pressure candidates are visually identified in these systems using high-quality UNIONS imaging from the Canada-France-Hawaii Telescope, covering $\sim 6200 \text{ deg}^2$ and $\sim 2800 \text{ deg}^2$ in the u - and r -bands, respectively. Ram pressure candidates are found in groups and clusters spanning a wide range in halo mass and include ~ 30 ram pressure candidates in the group regime ($M_h < 10^{14}$). The observed frequency of ram pressure candidates shows substantial scatter with group/cluster mass, but on average is larger in clusters ($M_h \geq 10^{14} M_\odot$) than groups ($M_h < 10^{14} M_\odot$) by a factor of ~ 2 . We find that ram pressure candidates are most commonly low-mass galaxies and have enhanced star formation rates relative to star-forming field galaxies. The enhancement in star formation is largely independent of galaxy mass and strongest for galaxies in clusters. As a result of the large survey footprint and excellent image quality from UNIONS, we are able to identify disturbed galaxies, potentially affected by ram pressure stripping, across a wide range of host environment.

Key words: galaxies: clusters: general – galaxies: evolution – galaxies: irregular.

1 INTRODUCTION

Galaxy clusters host populations of galaxies with properties that clearly differ from galaxies in low-density environments. Relative to galaxies in low-mass groups, or isolated in the field, galaxy populations in clusters are systematically redder, more gas poor, and have enhanced quiescent fractions. This prevalence of ‘red and dead’ galaxies in dense environments was noted by early, formative papers (e.g. Dressler 1980) and has since been confirmed by large,

modern galaxy surveys (e.g. Wetzel, Tinker & Conroy 2012; Haines et al. 2015; Brown et al. 2017). As these observed environmental trends have become more clear, the focus has shifted towards a physical understanding of the mechanisms causing these galaxy transformations. In particular, substantial effort has been devoted to understanding which physical processes are responsible for shutting off star formation (‘quenching’) in cluster galaxies.

The cluster environment subjects galaxies to gravitational/dynamical interactions with other member galaxies, as well as hydrodynamic interactions with the hot ($T \sim 1\text{--}10 \text{ keV}$; e.g. Mohr, Mathiesen & Evrard 1999) intracluster medium (ICM). These interactions are unique to dense environments and likely drive the

* E-mail: iroberts@strw.leidenuniv.nl

strong environmental dependence of galaxy properties which is observed. Specific physical processes capable of quenching star formation in clusters have been proposed, all of which involve gravitational and/or hydrodynamic interactions. Examples of these quenching mechanisms include: directly stripping gas from galaxy discs (ram pressure, RP, stripping, tidal stripping, viscous stripping; e.g. Gunn & Gott 1972; Nulsen 1982; Quilis, Moore & Bower 2000), preventing gas cooling within galaxies (starvation/strangulation; e.g. Larson, Tinsley & Caldwell 1980; Peng, Maiolino & Cochrane 2015), or inducing strong starbursts, typically as gas is funneled towards galaxy centres, which quickly consume cold-gas reserves (mergers, galaxy harassment; e.g. Mihos & Hernquist 1994a, b; Moore et al. 1996).

The build-up of massive clusters is a hierarchical process involving mergers with other galaxy clusters as well as lower mass groups. Therefore, some passive cluster galaxies may have actually been quenched in a lower mass group prior to accretion on to the more massive host cluster. This scenario is referred to as pre-processing and numerous works have attempted to constrain the relative contribution of pre-processing to the cluster red sequence (e.g. McGee et al. 2009; von der Linden et al. 2010; Haines et al. 2015; Roberts & Parker 2017; Pallerio et al. 2020). It is also likely that the efficiency of different quenching mechanisms differs between massive clusters and lower mass groups. For example, galaxy mergers are more common in groups where relative velocities are smaller, whereas RP stripping should be stronger in massive clusters with dense ICMs and large velocity dispersions (e.g. Hickson 1997; Hester 2006; Rasmussen, Ponman & Mulchaey 2006; Darg et al. 2010). Large redshift surveys at low- z , which contain thousands of groups and clusters across a wide range in mass ($\sim 10^{13}$ – $10^{15} M_{\odot}$), provide an excellent avenue for constraining this pre-processing.

Of key importance is identifying signatures of specific quenching mechanisms that can be observed and used to disentangle the relative contributions of various processes. For galaxies undergoing strong RP stripping, they are expected to leave a wake of stripped gas opposite to the direction of motion with respect to the ICM. Furthermore, bow shocks and enhanced star formation are expected on the galaxy leading edge (e.g. Clemens, Alexander & Green 2000; Yun et al. 2019). Typically, these stripped features are observed with the 21 cm line from neutral atomic hydrogen (e.g. Kenney, van Gorkom & Vollmer 2004; Chung et al. 2007, 2009; Kenney, Abramson & Bravo-Alfaro 2015) or the $H\alpha$ line tracing (partially ionized gas (e.g. Poggianti et al. 2017; Boselli et al. 2018)). That said, stripped features have also been observed in molecular hydrogen (e.g. Vollmer et al. 2012; Lee et al. 2017; Lee & Chung 2018; Jáchym et al. 2019), dust (e.g. Crowl et al. 2005), far-ultraviolet (e.g. Smith et al. 2010; Boissier et al. 2012; George et al. 2018), X-rays (e.g. Sun et al. 2010; Poggianti et al. 2019; Sun et al. 2021), and the radio continuum (e.g. Gavazzi & Jaffe 1987; Vollmer et al. 2009; Chen et al. 2020; Roberts et al. 2021a). While these methods provide an accurate means for identifying galaxies experiencing stripping, the required observations are often expensive and therefore target galaxies are typically known *a priori* to be morphologically disturbed. Alternatively, it is possible to identify galaxies that are likely undergoing RP stripping with rest-frame optical imaging (McPartland et al. 2016; Poggianti et al. 2016; Roberts & Parker 2020; Durret et al. 2021). In particular, blue filters (such as u -band) can be used to identify disturbed morphological features associated with RP stripping (Smith et al. 2010). These features are typically identified by-eye, but these visual classifications have been shown to agree well with quantitative morphological measures such as Gini- M_{20} and concentration-asymmetry (Conselice 2003; Lotz, Primack &

Madau 2004; McPartland et al. 2016; Roberts & Parker 2020). Since broad-band imaging is primarily tracing stellar-light, and the galaxy stellar components are less perturbed by RP stripping than neutral and ionized gas, RP classifications from optical imaging will be less accurate and/or less complete than corresponding $H I$ or $H\alpha$ surveys. That said, the advantage of broad-band imaging is the large areas which can be surveyed compared to spectroscopic observations, allowing for an unbiased, blind search for RP stripped galaxies across many groups and clusters.

In this work, we take advantage of the extremely large survey footprint (\sim thousands of square degrees of the northern sky) and high-resolution imaging (sub-arcsecond image quality, IQ) from the Ultraviolet Near-Infrared Optical Northern Survey (UNIONS). This multiband imaging survey includes the deep u - and r -band photometry from the Canada France Imaging Survey (CFIS) that is used in this work. Photometry in blue and/or near-UV filters is valuable for visually identifying candidate RP galaxies (Roberts & Parker 2020), as it traces a combination of young stellar populations and ionized gas. By matching the UNIONS footprint to SDSS spectroscopy, we perform an unbiased search for RP stripping across more than 50 groups and clusters with ~ 1000 star-forming, spectroscopic members. The size of this sample allows the investigation of the frequency of RP stripping as a function of both galaxy stellar mass and group/cluster mass.

The layout of this paper is the following. In Section 2, we describe the UNIONS imaging as well as the SDSS group/cluster and field galaxy samples used in this work. In Section 3, we describe the procedure that we employ to identify candidate RP galaxies. In Section 4, we present the primary results of this work including the frequency of RP candidates as a function of galaxy stellar mass and group/cluster mass (4.1), the position of RP candidates in projected phase space and the orientation of observed asymmetries (4.2), and the star formation properties of RP candidates relative to ‘normal’ star-forming cluster and field galaxies (4.3). We discuss these results in Section 5 and give our main conclusions in Section 6.

Throughout the paper, we assume a flat, Λ cold dark matter cosmology with $\Omega_M = 0.3$, $\Omega_{\Lambda} = 0.7$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 DATA

2.1 UNIONS imaging

UNIONS is a new consortium of wide field imaging surveys of the Northern hemisphere. UNIONS consists of CFIS, members of the Pan-STARRS team, and the Wide Imaging with Subaru HyperSuprimeCam of the Euclid Sky (WISHES) team. Each team is currently collecting imaging at their respective telescopes: Canada-France-Hawaii Telescope (CFHT)/CFIS is targeting deep u - and r -band photometry, Pan-STARRS is obtaining deep i - and moderate-deep z -bands, and Subaru/WISHES is acquiring deep z . These independent efforts are directed, in part, to securing optical imaging to complement the *Euclid* space mission, although UNIONS is a separate consortium aimed at maximizing the science return of these large and deep ground-based surveys of the northern skies. In the current analysis, we use the UNIONS/CFIS u - and r -band data only.

CFIS (Ibata et al. 2017) is a CFHT Large Program that will image the northern sky in u - and r -band over $\sim 10\,000 \text{ deg}^2$ and 5000 deg^2 , respectively. r -band imaging will be obtained above a declination of 30° and the u -band imaging will cover declinations above 0° . At the time of writing, $\sim 6200 \text{ deg}^2$ of u -band imaging and $\sim 2800 \text{ deg}^2$ of r -band imaging has been collected and reduced. CFIS provides sub-arcsecond IQ and reaches depths typically 3 mag deeper than SDSS

photometry. The u - and r -band imaging in this work is taken with the MegaCam wide-field imaging instrument at the CFHT, which covers a full $1 \times 1 \text{ deg}^2$ field-of-view with a pixel scale of $0.187''/\text{px}$. Raw images are flat-fielded and bias subtracted, astrometry is calibrated with Gaia DR2 (Gaia Collaboration 2016, 2018), and photometric calibration is done with Pan-STARRs (Chambers et al. 2016). Individual images are stacked on to $0.5 \times 0.5 \text{ deg}$ tiles, and images used in this work are extracted from these tiles. The median IQ across all CFIS imaging is 0.92 arcsec and 0.68 arcsec for the u - and r -bands, and the median depths (5σ point source) are 24.5 and 25.3 for u and r . This excellent IQ, which is relatively uniform across the sky, is an asset that makes it possible to identify RP candidates (see Section 3) across a wide range in galaxy mass and environment.

2.2 Cross-matching UNIONS with SDSS groups and clusters

To complement the imaging provided by UNIONS, we cross-match the CFIS footprint to galaxies with SDSS spectroscopic redshifts, focusing on galaxies in the vicinity of SDSS groups and clusters. We use the Lim et al. (2017) SDSS group catalogue, which follows the group finding algorithm from the Yang et al. (2005, 2007), with an improved method for halo mass assignment. We select all groups and clusters with seven or more spectroscopic members, halo masses greater than $10^{13} h^{-1} M_{\odot}$, and u - and r -band CFIS imaging. Seven galaxies correspond to the typical galaxy membership for groups in the Lim et al. catalogue at $z < 0.04$ with $M_h \sim 10^{13} M_{\odot}$. This also still gives enough spectroscopic members to derive reasonably accurate luminosity-weighted centres. We only include groups/clusters at $z < 0.04$ to ensure sub-kpc IQ for our entire galaxy sample. In Fig. 1 (top), we show the distributions of median IQ for each group/cluster in both u - and r -band. The IQ is best for the r -band observations, but for both u - and r -band the median IQ is sub-kpc for all groups/clusters. In Fig. 1 (bottom), we show the median 5σ magnitude limits for each group/cluster. The u -band limiting magnitudes range between 24th and 25th magnitude and the r -band limiting magnitudes are almost all fainter than 25th magnitude.

To select member galaxies for each group/cluster we do not use the galaxy memberships given in Lim et al. catalogue, and instead include all galaxies which are within $1 \times R_{180}$ and $3 \times \sigma_v$ of the group/cluster centre and redshift given in Lim et al. (2017). This is a loose membership criterion, which likely includes some galaxies not strictly bound to their host system. We opt for this approach to ensure that we do not exclude galaxies just starting their infall, and note that the qualitative results of this work are unchanged with the stricter galaxy memberships given by Lim et al. (2017). For each group/cluster the virial radius, R_{180} , and velocity dispersion, σ_v , are calculated following Lim et al. (2017) as

$$R_{180} = 1.33 h^{-1} \text{Mpc} \left(\frac{M_h}{10^{14} h^{-1} M_{\odot}} \right)^{1/3} (1 + z_{\text{cluster}})^{-1} \quad (1)$$

$$\sigma_v = 418 \text{ km s}^{-1} \left(\frac{M_h}{10^{14} h^{-1} M_{\odot}} \right)^{0.3367}, \quad (2)$$

where M_h is the halo mass and z_{cluster} is the cluster redshift. The criteria outlined above select 69 groups and clusters with halo masses ranging from $10^{13.1} M_{\odot}$ to $10^{14.4} M_{\odot}$. With our loose membership criteria, we identify a total of 2059 member galaxies with SDSS redshifts across these systems. For all galaxies, we obtain stellar masses (M_{star}) and star formation rates (SFRs) from the GSWLC-2 catalogue (Salim et al. 2016; Salim, Boquien & Lee 2018), which derives stellar masses and SFRs for SDSS galaxies via SED fitting




Figure 1. *Top:* Median u - and r -band IQ, in kpc, for each of the CFIS groups/clusters in the sample. *Bottom:* Median 5σ u - and r -band magnitude limits for each of the CFIS groups/clusters in the sample.

from the UV to the mid-IR. To ensure uniform coverage for the galaxies in our sample, we use the shallow, all-sky catalogue. For the remainder of the paper, we define star-forming galaxies to be all galaxies with specific SFRs above 10^{-11} yr^{-1} ($\text{sSFR} = \text{SFR}/M_{\text{star}}$). Of the member galaxies, 989 are star forming by this criteria.

2.3 Field sample

We use the SDSS isolated field galaxy sample from Roberts & Parker (2017). In brief, this field sample is compiled by selecting all galaxies in single-member groups from the Yang et al. (2005, 2007) catalogue, which are isolated from the nearest ‘bright’ galaxy by at least 1 Mpc and 1000 km s^{-1} . Bright galaxies are defined to be any galaxy with an r -band absolute magnitude brighter than the SDSS magnitude limit at $z = 0.04$ (the redshift upper limit of this sample). We further restrict the field sample by only including galaxies which overlap with both u - and r -band imaging tiles from the CFIS survey. This gives a field sample of 1352 SDSS galaxies with multiband CFIS imaging. Stellar masses and SFRs for the field sample are also obtained from the GSWLC-2 catalogue (Salim et al. 2016, 2018). The stellar mass and redshift distributions for the field sample are well matched to the sample of group galaxies described in the previous section.

3 IDENTIFYING RAM PRESSURE CANDIDATES

We identify RP candidates from pseudo three-colour images. For each star-forming galaxy an rgb thumbnail is made with the r -band in the r -channel, the u -band in the b -channel, and the average of the u - and r -bands in the g -channel. 50×50 kpc fits cutouts are made for each channel, and the three channels are combined to make rgb thumbnails with STIFF¹ (Bertin 2012).

To visually identify RP candidates we follow the criteria given in Roberts & Parker (2020) for galaxies in the Coma Cluster. These criteria were designed to identify galaxies which may be experiencing RP stripping; however, more generally they will flag strongly disturbed galaxies with asymmetric star formation. In Roberts & Parker (2020), we show that, on average, galaxies matching these criteria are consistent with RP stripping. That said, for some galaxies, other cluster processes such as tidal effects or impulsive interactions (i.e. galaxy harassment) may certainly play a role. We note that 13/17 of the Coma RP candidates identified by Roberts & Parker (2020) with stellar masses $> 10^{9.5} M_{\odot}$ (9/11 with stellar mass $> 10^{10} M_{\odot}$) have been subsequently observed to have clear RP tails visible in the radio continuum with LOFAR at 144 MHz (Roberts et al. 2021a). The LOFAR Two-metre Sky Survey (LoTSS) suffers from substantial incompleteness for star-forming galaxies with $M_{\text{star}} \lesssim 10^{9.5-10} M_{\odot}$, therefore sensitivity limitations likely also play a role for those RP candidates from Roberts & Parker (2020) without detected radio continuum tails. The high recovery rate of radio continuum tails for Coma Cluster RP candidates reinforces that these visual criteria are relatively effective at identifying RP stripping. RP candidates are identified according to one (or more) of the following visual criteria:

- (i) *The presence of asymmetric tails.* Observed on one side of the galaxy either in u -band (blue emission) or in dust (dark red extinction).
- (ii) *Asymmetric star formation.* u -band (blue) emission that is knotty and clearly asymmetric about the galaxy centre.
- (iii) *The presence of a bow shock morphology.* u -band (blue emission) or dust (dark red extinction) concentrated along one edge of the galaxy, potentially tracing enhanced star formation along the leading edge at the galaxy-ICM interface.

Roberts & Parker (2020) show that these visual criteria identify galaxies that are clearly disturbed according to quantitative morphology measures such as Asymmetry and M_{20} . Here, Asymmetry is a measure of the rotational asymmetry and corresponds to the fractional galaxy flux that is contained in asymmetric components (e.g. Abraham et al. 1996; Conselice 2003), and M_{20} is a statistic derived from the second-order moment brightest 20 percent of a galaxy’s flux that is particularly sensitive to bright features that are offset from galaxy centres (e.g. Lotz et al. 2004).

We note that the tails referenced above should not be thought of as large-scale extensions of material trailing the galaxy (for instance $H\alpha$ or $H\text{I}$ RP tails; e.g. Chung et al. 2009; Poggianti et al. 2017; Boselli et al. 2018), but instead we use this ‘tail’ notation to refer to filamentary or spiral arm like extensions of the galaxy that are asymmetric about the bright central component of the galaxy (i.e. these features are more extended towards one side of the galaxy than the other). This is in contrast to the ‘asymmetric star formation’ condition that is tracing asymmetric u -band emission (often knotty and compact) that could be embedded in an otherwise symmetric galaxy disc. Shocks from RP can induce enhanced gas

densities in interstellar medium (ISM) of galaxies, in particular in a bow-like morphology along the leading edge of the galaxy (e.g. Rasmussen et al. 2006; Ramos-Martínez, Gómez & Pérez-Villegas 2018; Troncoso-Iribarren et al. 2020). These enhanced densities in turn can lead to strong star formation along the leading edge (e.g. Gavazzi et al. 2001; Tomičić et al. 2018; Boselli et al. 2021). It is the result of this ICM-ISM interaction that we are looking to identify when we consider ‘bow shock morphologies’ in the u -band or dust.

Visual inspections were completed by a single classifier (IDR). We note that the exact criteria from Roberts & Parker (2020) were used, and in Roberts & Parker (2020) there was good agreement shown between multiple classifiers (e.g. 83 per cent of RP candidates were identified by at least 4/5 classifiers). Mergers are also flagged in this visual classification process, and galaxies identified as mergers are removed from the sample. As in Roberts & Parker (2020), mergers are flagged on the basis of two clearly interacting galaxies, or the presence of multiple, bright galaxy nuclei for the case of more evolved mergers. This does not amount to an exhaustive merger identification, but it serves the purpose of ensuring that obvious mergers are not contaminating the RP candidate sample. Less than 2 per cent of classified galaxies were identified as mergers in this way. Galaxies are classified in a randomized order and the coordinates of the galaxy and the ID of the host cluster are not known in the classification process. 48 RP candidates are identified by this process, which span the entire stellar mass range of the sample from $\sim 10^9 M_{\odot}$ to $\sim 10^{11} M_{\odot}$. 29/48 RP candidates are hosted by group mass haloes ($< 10^{14} M_{\odot}$) and 19/48 RP candidates are hosted by cluster mass haloes ($\geq 10^{14} M_{\odot}$). In Fig. 2, an example of the u -band, r -band, and rgb imaging from CFIS is shown. For comparison, we also show the u , r , and rgb SDSS images (with the same filters as the CFIS images). The differences in resolution and depth are clear for both the u and the r band, highlighting the superior IQ of CFIS. In Fig. 2, faint blue tails are barely visible in the SDSS image but clear in CFIS. Furthermore, \sim arcsecond scale sources are resolved by the CFIS imaging, many of which are coincident with these tails, that are not seen at all in the SDSS imaging. In Appendix A, we show u , r , and rgb CFIS images for all of the RP candidates (Figs A1–A3) and we also include a table with the coordinates of all of the RP candidates as well as some brief notes on the morphological features used to classify each galaxy.

As a test of our classification procedure, when visually classifying group galaxies we also randomly inject 500 galaxies from the field sample. In Fig. 2, we also show an example of the UNIONS and SDSS imaging for a randomly selected galaxy from the field sample. These field galaxies are randomly injected and when classifying a galaxy the classifier does not know whether the galaxy is from the cluster or the field sample. This is an important test of the process, as we are trying to flag galaxies according to cluster-specific processes (i.e. RP stripping), field galaxies should not pass the classification criteria. After all galaxies have been classified (group galaxies + 500 random field galaxies), we go back and check the false-positive rate of our classifications – i.e. what fraction of galaxies from the field sample were classified as RP candidates. We find that this false positive rate is 1.2 per cent, with 6/500 field galaxies passing the visual criteria for RP candidates. All of the field galaxies flagged as RP candidates were selected on the basis of asymmetric star formation within the galaxy, no galaxies in the field sample were flagged as showing asymmetric tails of bow shock morphological features. We note that this false positive rate is clearly below the fraction of RP candidates that we find among group galaxies (see Fig. 3), therefore our method appears reliable at morphologically identifying galaxies according to cluster-specific processes.

¹<https://www.astromatic.net/software/stiff>




Figure 2. Example u -band, r -band, and rgb images of an RP candidate and a randomly selected field galaxy. For each, in the top row we show the UNIONS imaging used for classification in this work, and for comparison in the bottom row we show the analogous images from the SDSS, highlighting the excellent IQ in UNIONS. The scale bar in each panel corresponds to a physical size of 10 kpc.




Figure 3. RP candidate fraction (see equation 3) as function of stellar mass. Numbers along the top of the panel correspond to the number of RP candidates in each stellar mass bin (purple) and the total number of galaxies in each stellar mass bin (black). The dashed black line shows the ‘false-positive rate’ from our classifications, measured by randomly injecting field galaxies into the group galaxy classifications (see Section 3). Error bars correspond to 68 per cent binomial confidence intervals (Cameron 2011).

As is shown in Fig. 1, CFIS provides excellent IQ and relatively uniform depth across the diverse set of groups and clusters in our sample. The methods used in this work are only possible thanks to this uniform imaging. If there were large variations in IQ or depth across the groups and clusters in the sample, this could bias the selection RP candidates towards the systems with the highest quality imaging. We note that we do not find any variations in the fraction of RP candidates identified as a function of IQ or depth, over the narrow range covered by the CFIS imaging. Finally, we reiterate the gains provided by CFIS over other wide-area imaging surveys such as the SDSS. Without the depth and resolution provided by CFIS it becomes very difficult to identify these morphological features with confidence (Fig. 2). This is particularly true for faint, low-mass galaxies which are most strongly influenced by environment (e.g. Haines et al. 2006; Bamford et al. 2009; Peng et al. 2010).

3.1 A note on nomenclature

Throughout this paper we make comparisons between three main galaxy samples, and will use the following nomenclature to differentiate them. ‘Field galaxies’ refers to star-forming galaxies from the field sample described in Section 2.3. ‘Group galaxies’ will refer to star-forming galaxies that are group/cluster members (as described in Section 2.2), but are not classified as RP candidates. We refer to these galaxies as group galaxies as the majority of the systems in this sample have $M_h < 10^{14} M_{\odot}$, that said the group galaxy sample does include some galaxies in haloes with $M_h \geq 10^{14} M_{\odot}$ which are typically considered clusters. ‘RP candidates’ will refer to star-forming galaxies that were identified as RP candidates according to the criteria described above. Galaxies undergoing mergers have been visually identified and removed from all samples. All galaxies considered in this paper are star-forming according to the criteria, $s\text{SFR} > 10^{-11} \text{ yr}^{-1}$.

Additionally, this work considers a sample of SDSS groups and clusters spanning two orders of magnitude in halo mass. Throughout the paper we will generally refer to groups as systems with $M_h < 10^{14} M_{\odot}$ and clusters as systems with $M_h \geq 10^{14} M_{\odot}$.

4 RESULTS

4.1 Ram-pressure candidate demographics

4.1.1 Stellar mass

We first consider the frequency of RP candidates as a function of galaxy stellar mass. We measure the frequency of RP candidates, which we define relative to all star-forming galaxies as

$$f_{\text{RP}} = \frac{N_{\text{RP}}}{N_{\text{SF}} + N_{\text{RP}}}, \quad (3)$$

where N_{RP} is the number of RP candidate galaxies, and N_{SF} is the number of normal star-forming galaxies (non-stripping). f_{RP} gives the fraction of all star-forming galaxies in groups and clusters that have been identified as RP candidates.

In Fig. 3, we plot f_{RP} versus stellar mass for the sample of RP candidates, showing that f_{RP} increases from intermediate stellar masses to low stellar masses. This is consistent with previous studies that argue low-mass galaxies are preferentially influenced by RP stripping (e.g. Fillingham et al. 2015; Roberts et al. 2019; Yun et al. 2019), likely due to their shallow potential wells. We note that the stellar mass distributions for galaxies in low-mass groups ($M_h < 10^{13.5} M_{\odot}$), high-mass groups ($10^{13.5} \leq M_h < 10^{14} M_{\odot}$), and clusters ($M_h \geq 10^{14} M_{\odot}$) are virtually identical. Therefore, the trend with stellar mass in Fig. 3 is not due to different host halo masses for low- versus high-mass galaxies.

4.1.2 Group/cluster mass

In Fig. 4, we plot the frequency of RP candidates as a function of group/cluster mass, ranging from low-mass groups ($\sim 10^{13} M_{\odot}$) to high-mass clusters ($\sim 10^{14.5} M_{\odot}$). The data points mark f_{RP} for individual groups and clusters, and solid bands show the combined f_{RP} separately for groups ($M_h < 10^{14} M_{\odot}$) and clusters ($M_h \geq 10^{14} M_{\odot}$). Open markers and the dashed line correspond to galaxy groups and filled markers and the solid line correspond to galaxy clusters. For reference, we also plot the frequency of RP candidates in the Coma Cluster presented in Roberts & Parker (2020).

On a system-by-system basis there is significant scatter in Fig. 4, with the bulk of the scatter occurring at the low-halo mass end. The reason for the large scatter at low group masses is simply low-number statistics, as most groups in this sample host < 10 star-forming members. The observed frequency of RP candidates in individual groups/clusters ranges from 0 to ~ 30 per cent. When considering the combined value of f_{RP} for groups and clusters separately, we find a small enhancement in the frequency of RP candidates for clusters compared to groups. f_{RP} for clusters is larger than f_{RP} for groups by roughly a factor of 2, and this difference is moderately significant at the $\sim 2\sigma$ level. Notably, all of systems with zero RP candidates fall in the group regime with $M_h < 10^{14} M_{\odot}$, whereas all systems with $M_h \geq 10^{14} M_{\odot}$ have detected RP candidates. This may be a physical effect, in that the less dense ICM and smaller velocity dispersions in the group regime are less conducive to RP stripping. The high number of groups with no RP candidates could also be a result of the smaller galaxy memberships in lower mass haloes. Even if the intrinsic frequency of RP candidates does not




Figure 4. RP candidate fraction as a function of group/cluster halo mass. Open stars correspond to individual galaxy groups ($M_H < 10^{14} M_\odot$) and filled stars correspond to individual galaxy clusters ($M_H < 10^{14} M_\odot$). The horizontal dashed line shows the total fraction for all galaxy groups and the solid line shows the total fraction for all clusters. Error bars are 68 per cent binomial confidence intervals and shaded regions correspond to 68 and 95 per cent confidence regions for the binned fractions (Cameron 2011). We also show the RP candidate frequency for galaxies in the Coma Cluster from Roberts & Parker (2020).

depend on halo mass, there would naturally be more low-mass groups with zero RP candidates simply because they host fewer galaxies.

Of the 61 groups in our sample with $M_h < 10^{14} M_\odot$, we find that 41 (67 per cent) have zero star-forming galaxies classified as RP candidates. We can now gauge whether this high fraction of systems with no RP candidates can be explained solely by the smaller galaxy memberships (as described above). In other words, if we assume that galaxy groups ($M_h < 10^{14} M_\odot$) and galaxy clusters ($M_h \geq 10^{14} M_\odot$) have the same intrinsic frequency of RP candidates, and given the fact that they have fewer member galaxies, what fraction of groups would we expect to observe with zero RP candidates due to random chance alone? To accomplish this we take the following procedure:

(i) We take the observed frequency of RP candidates for clusters ($M_h \geq 10^{14} M_\odot$) in our sample, which is 8 per cent.

(ii) We take the observed number of star-forming galaxies in each group ($M_h < 10^{14} M_\odot$), which ranges from 4 to 42 galaxies for our sample.

(iii) We randomly draw a binary array (0 \rightarrow normal star-forming galaxy, 1 \rightarrow RP candidate) for each group with length equal to the number of star-forming galaxies in that system. The probability of drawing a 0 or 1 is weighted by 0.92 and 0.08, respectively, to reflect the frequency of RP candidates in clusters.

(iv) For each group, we count the number of non-zero elements in the random array, corresponding to the number of RP candidates expected in that system assuming the same frequency as clusters (8 per cent). We then determine the fraction of groups expected to have zero RP candidates according to this random draw.

(v) We repeat this process for a total of 5000 iterations to generate a probability distribution function (PDF) for the fraction of groups that are expected to be observed with zero RP candidates.




Figure 5. PDF corresponding to the fraction of groups ($M_h < 10^{14} M_\odot$) expected to host zero RP candidates, assuming the same RP frequency as clusters (8 per cent). The observed fraction of groups with zero RP candidates (68 per cent) is also shown.

The resulting PDF is shown in Fig. 5 along with the best-fitting normal distribution, as well as the observed fraction of groups with zero RP candidates, of 68 per cent. The observed fraction is clearly larger than expected given the assumption that the frequency of RP candidates does not depend on halo mass. The significance of this discrepancy is 4σ . This implies that the large fraction of groups with zero RP candidates cannot be explained by galaxy membership alone, and instead the intrinsic frequency of RP candidates must be lower in groups than clusters. In order to bring the observed fraction in Fig. 5 within the 1σ scatter of the PDF, one must assume that the frequency of RP candidates is smaller by a factor of 2 in groups compared to clusters.

4.2 Projected phase space and u -band asymmetry orientations

In this section, we explore the distribution of RP candidates in projected phase space (velocity offset versus projected radius) as well as estimate the direction of RP based off on the orientation of the observed asymmetries in the u -band.

4.2.1 Phase space

Previous works have shown that galaxies undergoing RP stripping in clusters are systematically found at large velocity offsets in phase space (e.g. Yoon et al. 2017; Jaffé et al. 2018; Roberts & Parker 2020; Roberts et al. 2021a). This corresponds to regions of phase space where RP should be strong (since $P_{\text{ram}} \sim \rho_{\text{ICM}} v^2$), and also suggests that galaxies in clusters are primarily being stripped on their first infall (Jaffé et al. 2018). The majority ($\gtrsim 60$ per cent) of RP candidates in this work are hosted by galaxy groups with $M_h < 10^{14} M_\odot$, and the rest are primarily hosted by low-mass clusters ($M_h \sim 10^{14} M_\odot$). Therefore, we can test whether the phase space trends reported for RP stripping galaxies in clusters also hold in lower mass haloes. Unfortunately, we do not have sufficient statistics to do a group/cluster split from our sample alone.

In Fig. 6, we show the distribution of group galaxies (grey-scale 2D histogram) and RP candidates (purple stars) in projected phase space.




Figure 6. Projected phase space diagram. The phase space distribution for group galaxies is shown with 2D histogram and RP candidates are marked by purple stars. The dashed line shows the virialized region (dominated by ‘ancient infallers’) of phase space as given in Rhee et al. (2017). We also show the marginal kernel density distributions for projected radius and velocity offset above the corresponding axes.

We also show the marginal distributions for normalized radius and velocity offsets above the corresponding axes. Contrary to previous results for rich galaxy clusters, we find no difference between the phase space distribution for RP candidates and normal star-forming galaxies from this work. There is no evidence in Fig. 6 that RP candidates are shifted to large velocity offsets or small group/cluster-centric radii. This can be seen from the similarity of the marginal distributions for RP candidates (purple) and field galaxies (black) in Fig. 6. It is confirmed quantitatively with the two-sample Anderson–Darling test (Scholz & Stephens 1987). This could be a product of the weaker RP, on average, in the lower mass haloes that our sample probes, meaning that galaxies in groups may not show signs of stripping shortly after infall as is the case in clusters. If galaxies in groups have RP stripping features that persist beyond first infall, then that could explain the RP candidate distribution in Fig. 6 which appears well mixed and indistinguishable from the group galaxies. This picture is supported by the work of Oman et al. (2021) who model the time-scales of gas stripping as a function of halo mass. Oman et al. (2021) find that galaxy groups ($M \sim 10^{13.5} M_{\odot}$) strip their galaxies on longer time-scales than clusters ($M \sim 10^{14.5} M_{\odot}$), with a typical difference of ~ 3 Gyr. The distribution of RP candidates in Fig. 6 does not show strong evidence for galaxies (primarily in groups) being stripped primarily on first infall, as has been suggested for galaxies in clusters (e.g. Yoon et al. 2017; Jaffé et al. 2018; Roberts & Parker 2020).

4.2.2 *u*-band asymmetry orientations

We now attempt to constrain the direction of RP based on the observed asymmetries in RP candidates. The distribution of these orientations contains information on orbital and infall history for galaxies undergoing RP stripping; namely, galaxies stripped on first infall should have asymmetries oriented away from the group/cluster centre and galaxies being stripped after orbiting past pericentre should have asymmetries oriented towards the group/cluster centre.




Figure 7. Orientation of *u*-band asymmetries for identified RP candidates. An orientation of 0° corresponds to an asymmetry directed towards the centre of the host group/cluster and an orientation of 180° corresponds to an asymmetry oriented directly away from the centre of the host group/cluster.

While useful to gather information on galaxy orbits, this technique is also complicated by observational uncertainties. Projection effects are a significant concern as observed orientations that are projected on the sky will not always be representative of the true 3D directions. For this work specifically, there is added uncertainty given that we are observing RP features in the rest-frame optical and therefore primarily tracing stellar distributions. RP features are best traced by lower density components of the ISM (e.g. H I, H α , radio continuum, etc.) that are more easily stripped by RP. Perturbations to the optical morphologies are more subtle, and therefore more ambiguous in terms of connecting to RP. That said, for most jellyfish galaxies in the Coma Cluster, observed radio continuum tails are roughly consistent with stripping directions estimated from optical imaging (see the appendices of Roberts & Parker 2020 and Roberts et al. 2021a).

To estimate tail directions we use visual inspection of the colour images to assign the direction of the observed asymmetry for each galaxy an angle between 0° and 360° , where 0° = west and 90° = north. This is the same methodology used by Roberts & Parker (2020) for optically selected RP candidates in Coma and by Roberts et al. (2021a, b) for jellyfish galaxies in groups and clusters with radio continuum tails. These directions are meant to be representative of the direction of RP stripping; namely, if prominent, extended tails were present these are the directions along which we would expect to see them. All of the estimated asymmetry directions are indicated on the panel images in Appendix A. From these asymmetry directions we then compute the angle with respect to the centre of the corresponding host group or cluster such that an orientation of 0° corresponds to an asymmetry pointed directly at the group/cluster centre and an orientation of 180° corresponds to an asymmetry pointed directly away from the group/cluster centre. In Fig. 7, we show the distribution of asymmetry orientations for the RP candidates in this work. Fig. 7 shows a relatively uniform distribution of angles, with the exception for the $[150^{\circ}, 180^{\circ}]$ bin that shows evidence for an excess of RP candidates with observed asymmetries that are oriented away from the centre of the host group/cluster. Given the uncertainties with regards to estimating orientations from rest-frame optical imaging (see previous paragraph), projection effects, and the relatively small




Figure 8. SFR versus stellar mass for field galaxies (grey-scale), group galaxies (contours), and RP candidates (purple stars). The solid line shows the best-fitting SFMS relation for isolated field galaxies.

number of RP candidates in our sample, it is difficult to make strong conclusions from Fig. 7. That said, the broad distribution of angles in Fig. 7 is consistent with a fairly mixed set of orbits among the RP candidates in this work, similar to the phase space results above. This is in contrast to asymmetry directions from optical imaging for RP candidates in the Coma Cluster that are almost exclusively above 120° (Roberts & Parker 2020). In Fig. 7, there is a small excess of orientations above 150° which may indicate a slight overabundance of RP candidates on their first infall towards the group/cluster centre.

4.3 Star formation properties

To probe cluster specific effects on galaxy star formation, we compare group galaxies to the isolated field sample. We use the star-forming galaxies in the field sample to fit a field star-forming main sequence (SFMS) and compare cluster SFRs relative to this standard. We consider all galaxies in the field sample with $sSFR > 10^{-11} \text{ yr}^{-1}$ and fit a power law to the SFR– M_{star} relation. The fit is done with the LINMIX² (Kelly 2007) package and accounts for uncertainties on both SFR and M_{star} for each field galaxy. The best-fitting SFMS relation for the field sample is $\log \text{SFR} = 0.55 \times \log M_{\text{star}} - 5.7$.

In Fig. 8, we plot SFR versus stellar mass. Star-forming field galaxies are shown by the grey-scale 2D histogram, the distribution of star-forming group galaxies are shown by the contours, and RP candidates are shown by the purple stars. The solid black line marks the best-fitting SFMS for the field sample. It is clear from Fig. 8 that RP candidates are systematically found above the field SFMS, quantitatively, 77 per cent of RP candidates lie above the field SFMS. This enhanced star formation in RP candidates has been reported by Roberts & Parker (2020) for galaxies in the Coma Cluster, and here is confirmed for RP candidates across a wide range in cluster mass.

For all RP candidates, we now compute the offset in SFR relative to the field SFMS, Δ_{MS} . This allows us to quantify the enhancement of star formation for RP candidates as a function of stellar and halo mass. In Fig. 9, we plot the offset from the SFMS as a function of

stellar mass (left) and halo mass (right). Data points show the offsets for individual RP candidates. An enhancement in star formation ($\Delta_{\text{MS}} > 0$) is present across the entire range in stellar mass, except, perhaps, the high-mass end where the number of RP candidates is low. The typical value for Δ_{MS} is ~ 0.2 dex and there is no clear trend between Δ_{MS} and stellar mass.

In Fig. 9 (right), we show Δ_{MS} as a function of halo mass, with RP candidates in groups marked by the open symbols and RP candidates in clusters marked by the filled symbols. We also show Δ_{MS} for Coma RP candidates from Roberts & Parker (2020) at the high halo mass end. The median Δ_{MS} for RP candidates in groups, clusters, and Coma are shown by the horizontal lines. A mild difference is seen in Fig. 9 with a larger median Δ_{MS} in more massive haloes. RP candidates hosted by clusters in this sample as well as the Coma Cluster (Roberts & Parker 2020) have larger offsets from the SFMS than RP candidates hosted by groups. Furthermore, RP candidates in clusters scatter to much larger Δ_{MS} values than is seen for group RP candidates. The RP candidates found far above the SFMS in Fig. 8 are largely hosted by relatively massive clusters, which may be connected to stronger RP in these systems. We reiterate that all offsets from the SFMS are measured relative to the field SFMS. When compared to normal star-forming galaxies in clusters, RP candidates show enhanced star formation at all stellar masses and within clusters of all halo masses.

5 DISCUSSION

Roberts & Parker (2020) show that the visual criteria listed in Section 3 select a population of galaxies that are consistent with RP stripping as the primary driver behind the observed disturbed morphologies. Furthermore, many of the Coma RP candidates identified by Roberts & Parker (2020) have been subsequently observed to have clear RP tails visible in the radio continuum with LOFAR at 144 MHz (Roberts et al. 2021a), reinforcing that these visual criteria are effective at identifying RP stripping. In this work, the stellar-mass dependence of RP candidates (i.e. preferentially low-mass galaxies) and the group/cluster mass dependence of RP candidates (i.e. preferentially found in higher mass clusters) are measured and are consistent with simple predictions from models of RP. That said, it is possible that some galaxies which we identify as RP candidates may owe their disturbed morphologies (at least in part) to other physical processes, for example tidal effects or galaxy harassment. In the future more multiwavelength data on these galaxies, such as H α or H I, will be an asset to more definitively constrain the origins their disturbed morphologies (i.e. ram pressure or a different perturbing mechanism?). The LoTSS (Shimwell et al. 2017, 2019) will finish imaging the entire northern extragalactic sky at 144 MHz within the next few years, therefore we will be able to measure how many of these RP candidates show extended radio continuum tails. Though the LoTSS sensitivity may not be high enough to detect the lowest mass RP candidates at 144 MHz (Roberts et al. 2021a).

Given that the criteria in Section 3 were designed to identify RP stripping galaxies, and that the results from this work and Roberts & Parker (2020) are consistent with expectations from RP stripping, for the remainder of the discussion we will assume that the morphologies of RP candidates in this work are primarily being driven by RP.

5.1 Stellar mass dependence and star formation

In Fig. 3, we find a clear mass dependence for the frequency of RP candidates. The highest values for f_{RP} correspond to the lowest mass galaxies in the sample. This increase in f_{RP} towards the lowest masses

²<https://linmix.readthedocs.io/en/latest/index.html>




Figure 9. *Left:* Offset from the field SFMS as a function of stellar mass for RP candidates. The dashed line shows the median offset from the SFMS and the shaded region shows the standard error on the median. *Right:* Offset from the SFMS as a function of halo mass. Open stars correspond to RP candidates in groups, filled stars correspond to RP candidates in clusters, and crosses are Coma Cluster RP candidates from Roberts & Parker (2020). Horizontal bars show median offsets for galaxies in groups (dashed purple), clusters (solid purple), and Coma (solid orange).

is likely to be physical in nature. It has been previously shown that environmental quenching is most important for low-mass galaxies ($M_{\text{star}} \lesssim \text{few} \times 10^{10} M_{\odot}$; Haines et al. 2006; Bamford et al. 2009), therefore there is some expectation that low-mass galaxies should be strongly affected by environmental processes. Given the relatively shallow potential wells of low-mass galaxies, they may be more susceptible to environmental processes such as RP stripping. This is consistent with the observed trend in Fig. 3. Yun et al. (2019) visually identify jellyfish galaxies (according to the gas column density maps) in the Illustris TNG simulation, finding that low-mass galaxies are most frequently jellyfish, and the fraction of jellyfish galaxies (relative to all gas-rich galaxies) decreases monotonically with increasing galaxy stellar mass. Other observational studies have also argued that low-mass galaxies are preferentially affected by RP stripping relative to higher mass counterparts (Fillingham et al. 2015; Wetzel, Tollerud & Weisz 2015; Roberts et al. 2019; Rodriguez Wimberly et al. 2019; Durret et al. 2021).

Across the stellar mass range, we see evidence for enhanced star formation (Figs 8 and 9) for RP candidates. The star formation enhancement is clear through the SED SFRs (Salim et al. 2016, 2018) used in the text. We also confirm that SDSS fibre $H\alpha$ fluxes are strongly enhanced in RP candidates, which trace only the central 3 arcsec of each galaxy. In an RP stripping context, the enhanced SFRs may be a result of compression and increased gas densities in the disc due to the galaxy-ICM interaction. When comparing RP candidates to the field SFMS, we find enhancements of roughly 0.2 dex, with no strong dependence on stellar mass. The enhanced SFRs in RP stripping galaxies are consistent with previous work from observations (Dressler & Gunn 1983; Ebeling, Stephenson & Edge 2014; Poggianti et al. 2016; Vulcani et al. 2018b; Roberts & Parker 2020; Wang et al. 2020; Durret et al. 2021) and simulations (Steinhauser et al. 2012; Bekki 2014; Troncoso Iribarren et al. 2016). In Fig. 9 (right), we see evidence for a weak dependence of SFMS offset on host halo mass. The observed offsets are slightly larger for RP candidates hosted by clusters ($M_h \geq 10^{14} M_{\odot}$) compared to groups ($M_h < 10^{14} M_{\odot}$), and there is significant scatter towards large main-sequence offsets for group galaxies. If robust, this trend

could be related to stronger RP in clusters, leading to more extreme gas compression which catalyses stronger bursts of star formation. This can potentially be tested observationally with recent and future surveys of molecular gas in group galaxies (e.g. VERTICO,³ the ALMA Fornax Cluster Survey, GASP; Zabel et al. 2019; Moretti et al. 2020; Brown et al. 2021). It is also important to note that we identify a relatively small number of RP candidates in the group regime, therefore it is difficult to make strong conclusions regarding offsets from the SFMS in groups without a larger sample.

5.2 Low-mass groups to massive clusters

Galaxies in our RP candidate sample span host halo masses ranging from $\sim 10^{13}$ – $10^{14.5} M_{\odot}$, and even up to $\sim 10^{15} M_{\odot}$ when including Coma Cluster RP candidates from Roberts & Parker (2020). The fact that we observe RP candidates in the group regime is consistent with previous works that have shown evidence for RP stripping in galaxy groups (Rasmussen et al. 2006; Vulcani et al. 2018a). RP scales as $\rho_{\text{ICM}} v^2$ (e.g. Gunn & Gott 1972), and both the density of the ICM and the relative velocities of satellite galaxies increase with cluster mass, at least on average. Therefore, galaxies should experience stronger RP in high-mass clusters than low-mass groups. We do find evidence for more RP candidates in clusters versus groups by roughly a factor of 2 (see Figs 4 and 5). This is consistent with simulation results from Yun et al. (2019) that show a factor of ~ 2 increase in the jellyfish fraction from low-mass groups ($M_h \sim 10^{13} M_{\odot}$) to galaxy clusters ($M_h > 10^{14} M_{\odot}$). It is difficult to directly compare the jellyfish fraction in Yun et al. (2019) to the RP candidate frequencies in this work, due to the very different methodologies (i.e. jellyfish galaxies identified from simulated gas column densities versus RP candidates from rest-frame optical imaging). That said the general trends with stellar mass and halo mass from Yun et al. are consistent

³<https://sites.google.com/view/verticosurvey/home>

with the observed trends in this work. Most of the groups in our sample have a relatively small number of spectroscopic members, with a median number of ~ 10 star-forming members. Therefore, on an individual group-by-group basis the frequency of RP candidates is highly susceptible to low-number statistics, as is apparent from the scatter in Fig. 4.

Given that RP is expected to be weaker in the group regime, galaxies in groups retain a larger fraction of their gas over the course of an orbit, whereas galaxies in clusters should be stripped more quickly. Galaxies in groups may strip more slowly relative to galaxies in clusters. This picture is also consistent with the quenching models in Oman et al. (2021). If RP stripping features survive for longer in galaxy groups this would increase the number of RP candidates identified in groups relative to clusters, contributing to the mild dependence of RP candidate frequency on halo mass seen in this work. The phase space distribution in Fig. 6 suggests that RP candidates in groups persist beyond first infall, consistent with this picture.

6 CONCLUSIONS

In this work, we present a sample of RP candidate galaxies, visually identified from the Canada-France Imaging Survey, which covers >50 SDSS groups and clusters ranging in mass from $\sim 10^{13}$ – $10^{14.5} M_{\odot}$. This large sample allows us to study the properties and demographics of galaxies likely being influenced by RP in nearby groups and clusters. The primary findings of this work are as follows:

(i) The frequency of RP candidates is highest for low-mass galaxies.

(ii) The average frequency of RP candidates is larger in clusters ($M_h \geq 10^{14} M_{\odot}$) than groups ($M_h < 10^{14} M_{\odot}$) by roughly a factor of 2. On an individual group/cluster basis, the frequency of RP candidates shows significant scatter with halo mass.

(iii) RP candidates have projected phase space distributions that are consistent with normal star-forming group galaxies. Since the majority of RP candidates in this work are hosted by galaxy groups, this suggests that galaxies in groups may show signature of RP stripping well after first infall.

(iv) RP candidates show a broad distribution of observed u -band asymmetry orientations; however, there is a small excess of RP candidates with asymmetric features oriented directly away from the group/cluster centre.

(v) RP candidates have enhanced SFRs, relative to both isolated field galaxies and normal star-forming group galaxies. Median offsets from the SFMS for RP candidates do not depend strongly on stellar mass, but are slightly larger for RP candidates in clusters compared to groups.

In this paper, we have presented an introductory look at this sample of CFIS RP candidates and the properties of these disturbed galaxies. In the next paper in this series we will move beyond visual classifications and explore the quantitative morphological properties (e.g. Gini- M_{20} , concentration-asymmetry) of the RP candidates identified in this work. Ultimately looking to determine whether an automated selection based on quantitative morphology measures can reproduce the sample of visually identified RP candidates from this work. This will be especially important in the context of next-generation photometric galaxy surveys (e.g. EUCLID, the Vera C. Rubin Observatory, the Roman Space Telescope), where the number of galaxies with high-quality imaging will surpass what is practical for by-eye classifications. The large number of RP candidates, across a range in redshifts, that could be identified with these

upcoming wide-field imaging projects will significantly advance our understanding of the prevalence and importance of RP stripping in dense environments.

ACKNOWLEDGEMENTS

IDR acknowledges support from the ERC Starting Grant Cluster Web 804208. LCP and MJH acknowledge funding from the Natural Sciences and Engineering Research Council of Canada.

This work is based on data obtained as part of the Canada-France Imaging Survey, a Canada-France-Hawaii Telescope (CFHT) large program of the National Research Council (NRC) of Canada and the French Centre National de la Recherche Scientifique. Based on observations obtained with MegaPrime/MegaCam, a joint project of CFHT and CEA Saclay, at the CFHT which is operated by the NRC of Canada, the Institut National des Science de l'Univers (INSU) of the Centre National de la Recherche Scientifique (CNRS) of France, and the University of Hawaii.

DATA AVAILABILITY

The SDSS catalogues used in this work are available at <https://salims.pages.iu.edu/gswlc/> and <https://gax.sjtu.edu.cn/data/Group.html>. A subset of the raw imaging data underlying this article are publicly available via the Canadian Astronomical Data Center at <http://www.cadc-ccda.hia-ihp.nrc-cnrc.gc.ca/en/megapipe/>. The remaining raw data and all processed data are available to members of the Canadian and French communities via reasonable requests to the principal investigators of the Canada-France Imaging Survey, Alan McConnachie and Jean-Charles Cuillandre. All data will be publicly available to the international community at the end of the proprietary period, scheduled for 2023.

REFERENCES

- Abraham R. G., Tanvir N. R., Santiago B. X., Ellis R. S., Glazebrook K., van den Bergh S., 1996, *MNRAS*, 279, L47
- Bamford S. P. et al., 2009, *MNRAS*, 393, 1324
- Bekki K., 2014, *MNRAS*, 438, 444
- Bertin E., 2012, in Ballester P., Egret D., Lorente N., eds, *Astronomical Data Analysis Software and Systems XXI*, Vol. 461, p. 263
- Boissier S. et al., 2012, *A&A*, 545, A142
- Boselli A. et al., 2018, *A&A*, 614, A56
- Boselli A. et al., 2021, *A&A*, 646, A139
- Brown T. et al., 2021, preprint ([arXiv:2111.00937](https://arxiv.org/abs/2111.00937))
- Brown T. et al., 2017, *MNRAS*, 466, 1275
- Cameron E., 2011, *Publ. Astron. Soc. Aust.*, 28, 128
- Chambers K. C. et al., 2016, preprint ([arXiv:1612.05560](https://arxiv.org/abs/1612.05560))
- Chen H. et al., 2020, *MNRAS*, 496, 4654
- Chung A., van Gorkom J. H., Kenney J. D. P., Crowl H., Vollmer B., 2009, *AJ*, 138, 1741
- Chung A., van Gorkom J. H., Kenney J. D. P., Vollmer B., 2007, *ApJ*, 659, L115
- Clemens M. S., Alexander P., Green D. A., 2000, *MNRAS*, 312, 236
- Conselice C. J., 2003, *ApJS*, 147, 1
- Crowl H. H., Kenney J. D. P., van Gorkom J. H., Vollmer B., 2005, *AJ*, 130, 65
- Darg D. W. et al., 2010, *MNRAS*, 401, 1552
- Dressler A., 1980, *ApJ*, 236, 351
- Dressler A., Gunn J. E., 1983, *ApJ*, 270, 7
- Durret F., Chiche S., Lobo C., Jauzac M., 2021, *A&A*, 648, A63
- Ebeling H., Stephenson L. N., Edge A. C., 2014, *ApJ*, 781, L40
- Fillingham S. P., Cooper M. C., Wheeler C., Garrison-Kimmel S., Boylan-Kolchin M., Bullock J. S., 2015, *MNRAS*, 454, 2039

- Gaia Collaboration, 2016, *A&A*, 595, A1
 Gaia Collaboration, 2018, *A&A*, 616, A1
 Gavazzi G., Boselli A., Mayer L., Iglesias-Paramo J., Vílchez J. M., Carrasco L., 2001, *ApJ*, 563, L23
 Gavazzi G., Jaffe W., 1987, *A&A*, 186, L1
 George K. et al., 2018, *MNRAS*, 479, 4126
 Gunn J. E., Gott J. R. III, 1972, *ApJ*, 176, 1
 Haines C. P. et al., 2015, *ApJ*, 806, 101
 Haines C. P., La Barbera F., Mercurio A., Merluzzi P., Busarello G., 2006, *ApJ*, 647, L21
 Hester J. A., 2006, *ApJ*, 647, 910
 Hickson P., 1997, *ARA&A*, 35, 357
 Ibata R. A. et al., 2017, *ApJ*, 848, 128
 Jáchym P. et al., 2019, *ApJ*, 883, 145
 Jaffé Y. L. et al., 2018, *MNRAS*, 476, 4753
 Kelly B. C., 2007, *ApJ*, 665, 1489
 Kenney J. D. P., Abramson A., Bravo-Alfaro H., 2015, *AJ*, 150, 59
 Kenney J. D. P., van Gorkom J. H., Vollmer B., 2004, *AJ*, 127, 3361
 Larson R. B., Tinsley B. M., Caldwell C. N., 1980, *ApJ*, 237, 692
 Lee B. et al., 2017, *MNRAS*, 466, 1382
 Lee B., Chung A., 2018, *ApJ*, 866, L10
 Lim S. H., Mo H. J., Lu Y., Wang H., Yang X., 2017, *MNRAS*, 470, 2982
 Lotz J. M., Primack J., Madau P., 2004, *AJ*, 128, 163
 McGee S. L., Balogh M. L., Bower R. G., Font A. S., McCarthy I. G., 2009, *MNRAS*, 400, 937
 McPartland C., Ebeling H., Roediger E., Blumenthal K., 2016, *MNRAS*, 455, 2994
 Mihos J. C., Hernquist L., 1994a, *ApJ*, 425, L13
 Mihos J. C., Hernquist L., 1994b, *ApJ*, 431, L9
 Mohr J. J., Mathiesen B., Evrard A. E., 1999, *ApJ*, 517, 627
 Moore B., Katz N., Lake G., Dressler A., Oemler A., 1996, *Nature*, 379, 613
 Moretti A. et al., 2020, *ApJ*, 889, 9
 Nulsen P. E. J., 1982, *MNRAS*, 198, 1007
 Oman K. A., Bahé Y. M., Healy J., Hess K. M., Hudson M. J., Verheijen M. A. W., 2021, *MNRAS*, 501, 5073
 Pallero D., Gómez F. A., Padilla N. D., Bahé Y. M., Vega-Martínez C. A., Torres-Flores S., 2020, preprint ([arXiv:2012.08593](https://arxiv.org/abs/2012.08593))
 Peng Y.-j. et al., 2010, *ApJ*, 721, 193
 Peng Y., Maiolino R., Cochrane R., 2015, *Nature*, 521, 192
 Poggianti B. M. et al., 2016, *AJ*, 151, 78
 Poggianti B. M. et al., 2017, *ApJ*, 844, 48
 Poggianti B. M. et al., 2019, *ApJ*, 887, 155
 Quilis V., Moore B., Bower R., 2000, *Science*, 288, 1617
 Ramos-Martínez M., Gómez G. C., Pérez-Villegas Á., 2018, *MNRAS*, 476, 3781
 Rasmussen J., Ponman T. J., Mulchaey J. S., 2006, *MNRAS*, 370, 453
 Rhee J., Smith R., Choi H., Yi S. K., Jaffé Y., Candlish G., Sánchez-Jánsen R., 2017, *ApJ*, 843, 128
 Roberts I. D. et al., 2021a, *A&A*, 650, A111
 Roberts I. D., Parker L. C., 2017, *MNRAS*, 467, 3268
 Roberts I. D., Parker L. C., 2020, *MNRAS*, 495, 554
 Roberts I. D., Parker L. C., Brown T., Joshi G. D., Hlavacek-Larrondo J., Wadsley J., 2019, *ApJ*, 873, 42
 Roberts I. D., van Weeren R. J., McGee S. L., Botteon A., Ignesti A., Rottgering H. J. A., 2021b, *A&A*, 652, A153
 Rodríguez Wimberly M. K., Cooper M. C., Fillingham S. P., Boylan-Kolchin M., Bullock J. S., Garrison-Kimmel S., 2019, *MNRAS*, 483, 4031
 Salim S. et al., 2016, *ApJS*, 227, 2
 Salim S., Boquien M., Lee J. C., 2018, *ApJ*, 859, 11
 Scholz F. W., Stephens M. A., 1987, *J. Am. Stat. Assoc.*, 82, 918
 Shimwell T. W. et al., 2017, *A&A*, 598, A104
 Shimwell T. W. et al., 2019, *A&A*, 622, A1
 Smith G. P. et al., 2010, *MNRAS*, 409, 169
 Steinhauser D., Haider M., Kapferer W., Schindler S., 2012, *A&A*, 544, A54
 Sun M. et al., 2021, preprint ([arXiv:2103.09205](https://arxiv.org/abs/2103.09205))
 Sun M., Donahue M., Roediger E., Nulsen P. E. J., Voit G. M., Sarazin C., Forman W., Jones C., 2010, *ApJ*, 708, 946
 Tomičić N. et al., 2018, *ApJ*, 869, L38
 Troncoso Iribarren P., Padilla N., Contreras S., Rodríguez S., García-Lambas D., Lagos C., 2016, *Galaxies*, 4, 77
 Troncoso-Iribarren P., Padilla N., Santander C., Lagos C. D. P., García-Lambas D., Rodríguez S., Contreras S., 2020, *MNRAS*, 497, 4145
 Vollmer B. et al., 2012, *A&A*, 537, A143
 Vollmer B., Soida M., Chung A., Chemin L., Braine J., Boselli A., Beck R., 2009, *A&A*, 496, 669
 von der Linden A., Wild V., Kauffmann G., White S. D. M., Weinmann S., 2010, *MNRAS*, 404, 1231
 Vulcani B. et al., 2018a, *MNRAS*, 480, 3152
 Vulcani B. et al., 2018b, *ApJ*, 866, L25
 Wang J., Xu W., Lee B., Du M., Overzier R., Shao L., 2020, *ApJ*, 903, 103
 Wetzel A. R., Tinker J. L., Conroy C., 2012, *MNRAS*, 424, 232
 Wetzel A. R., Tollerud E. J., Weisz D. R., 2015, *ApJ*, 808, L27
 Yang X., Mo H. J., van den Bosch F. C., Jing Y. P., 2005, *MNRAS*, 356, 1293
 Yang X., Mo H. J., van den Bosch F. C., Pasquali A., Li C., Barden M., 2007, *ApJ*, 671, 153
 Yoon H., Chung A., Smith R., Jaffé Y. L., 2017, *ApJ*, 838, 81
 Yun K. et al., 2019, *MNRAS*, 483, 1042
 Zabel N. et al., 2019, *MNRAS*, 483, 2251

APPENDIX A: RAM PRESSURE CANDIDATE IMAGES

Table A1. Ram pressure candidates.

Galaxy Number	RA [deg]	Dec. [deg]	z	Classification Notes
1	123.6493	58.3661	0.0293	Asymmetric u to the west
2	124.5442	57.7588	0.0268	Tails to the southeast, bright u to the north-west
3	125.0641	56.3767	0.0300	Tails to the south
4	125.1572	58.0453	0.0265	Tail to the south
5	128.3571	41.0544	0.0250	Tail to the west
6	130.0765	40.7965	0.0307	Tails and bright u to the south
7	136.8169	37.2154	0.0238	Asymmetric u to the north
8	137.3513	37.6858	0.0236	Tails to the southeast, u sources to the north-west
9	138.5934	30.4330	0.0215	Tail to the east, bright u on western edge
10	139.8783	38.9660	0.0278	Tails to the southeast, bow feature on north-west edge?
11	140.0890	34.2384	0.0246	Tail to the north, bright u source on southern edge
12	143.6829	33.9727	0.0289	Tail to the northeast
13	144.1025	31.8513	0.0222	Asymmetric u to northeast, tail to the south?
14	147.3384	34.6568	0.0400	Tail to the north
15	147.7212	34.6549	0.0399	Tail to the north
16	154.7025	38.4697	0.0223	Tail to the south, asymmetric u on northern edge (tidal?)
17	160.0280	39.1089	0.0342	Tails to the southwest, asymmetric u in the north
18	160.0464	38.5785	0.0360	Tail to the southwest
19	160.2714	37.3860	0.0234	Tail to the north
20	160.7922	39.0387	0.0339	Asymmetric u , bow feature on western edge
21	161.0554	39.2021	0.0356	Asymmetric u to the east
22	161.4409	39.0194	0.0359	Tail to the south, faint tail to the north? (tidal?)
23	161.6613	38.3612	0.0351	Tail to the south
24	161.8615	38.9364	0.0350	Tail to the east, bright u on western edge
25	165.2576	46.5653	0.0222	Asymmetric u to the northeast
26	165.2820	45.6527	0.0248	Tail to the southwest
27	168.7345	30.8958	0.0274	Asymmetric u to the east
28	172.0338	35.3783	0.0377	Asymmetric u to the southeast
29	173.0356	35.5109	0.0385	Asymmetric u to the west
30	173.0633	35.5125	0.0377	Tails to the south
31	175.7723	33.8349	0.0315	Tail to the south
32	175.8477	32.8801	0.0343	Bow feature on eastern edge, faint tail to the north-west?
33	175.9726	33.3490	0.0322	Tail to the northeast, bright u on southwestern edge
34	176.4507	33.1483	0.0321	Tail to the south, bright u on northern edge
35	176.5412	33.2083	0.0329	Tail to the west
36	197.5338	34.9055	0.0356	Asymmetric u to the south
37	197.6735	34.8172	0.0367	Tail/asymmetric u to the south
38	200.0656	33.6637	0.0391	Tail to the north
39	200.7128	31.8259	0.0178	Asymmetric u to the north-west
40	200.8339	32.0635	0.0167	Tails to the southeast
41	201.2553	36.4372	0.0195	Asymmetric u to the north
42	201.4014	36.3812	0.0188	Tails to the north/northeast (tidal?)
43	231.9340	43.0689	0.0175	Tail to the southwest
44	244.8756	37.7874	0.0327	Asymmetric u to the west
45	244.9793	38.1306	0.0304	Asymmetric u to the west
46	245.5684	38.4499	0.0292	Asymmetric u to the southeast
47	245.8868	38.0366	0.0338	Tail to the southeast
48	248.7128	36.7517	0.0337	Tail to the east




Figure A1. u -band (left), r -band (center), and rgb (right) for the RP candidates identified in this work. White line indicates the estimated 'tail' direction for each RP candidate (see Section 4.2.2).




Figure A2. u -band (left), r -band (centre), and rgb (right) for the RP candidates identified in this work. White line indicates the estimated 'tail' direction for each RP candidate (see Section 4.2.2).




Figure A3. u -band (left), r -band (centre), and rgb (right) for the RP candidates identified in this work. White line indicates the estimated ‘tail’ direction for each RP candidate (see Section 4.2.2).

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.