



HAL
open science

Existence and decay of traveling waves for the nonlocal Gross-Pitaevskii equation

André de Laire, Salvador López-Martínez

► **To cite this version:**

André de Laire, Salvador López-Martínez. Existence and decay of traveling waves for the nonlocal Gross-Pitaevskii equation. 2021. hal-03422447v1

HAL Id: hal-03422447

<https://hal.science/hal-03422447v1>

Preprint submitted on 9 Nov 2021 (v1), last revised 20 Apr 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence and decay of traveling waves for the nonlocal Gross–Pitaevskii equation

André de Laire¹ and Salvador López-Martínez²

Abstract

We consider the nonlocal Gross–Pitaevskii equation that models a Bose gas with general nonlocal interactions between particles in one spatial dimension, with constant density far away. We address the problem of the existence of traveling waves with nonvanishing conditions at infinity, i.e. dark solitons. Under general conditions on the interactions, we prove existence of dark solitons for almost every subsonic speed. Moreover, we show existence in the whole subsonic regime for a family of potentials. The proofs are based on a Mountain Pass argument combined with the so-called “monotonicity trick”, as well as on a priori estimates for the Palais–Smale sequences. Finally, we establish properties of the solitons such as exponential decay at infinity and analyticity.

Keywords: Nonlocal Schrödinger equation, Gross–Pitaevskii equation, traveling waves, dark solitons, nonzero conditions at infinity, decay, analyticity.

2010 *Mathematics Subject Classification:* 35Q55; 35J20; 35C07; 37K05; 35C08; 35A01, 37K40

1 Introduction

1.1 The problem

In order to describe the dynamics of a weakly interacting Bose gas, Gross [30] and Pitaevskii [51] found that the wavefunction Ψ governing the condensate satisfies a Schrödinger equation, that in dimension one and in its dimensionless form, is given by

$$i\partial_t\Psi = -\partial_{xx}\Psi + \Psi \int_{\mathbb{R}} |\Psi(y,t)|^2 V(x-y) dy, \quad \text{in } \mathbb{R} \times \mathbb{R}. \quad (1.1)$$

Here, $\Psi : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C}$ and V describes the interaction between bosons. In their works, they are interested in a function Ψ satisfying the nonzero condition at infinity:

$$\lim_{|x| \rightarrow \infty} |\Psi(x, \cdot)| = 1, \quad (1.2)$$

representing the fact that the density is constant far away.

Equation (1.1) also appears as the model for the evolution of a one-dimensional optical beam of intensity $|\Psi|^2$ in a self-defocusing nonlocal Kerr-like medium, where V characterizes the nonlocal response of the medium [41, 49]. In this case, the condition (1.2) is natural when studying dark optical solitons. In all of these physical situations, V is assumed to be real-valued and symmetric. Moreover, in the most typical first approximation, V is considered as a

¹Univ. Lille, CNRS, Inria, UMR 8524 - Laboratoire Paul Painlevé, Inria, F-59000 Lille, France.
E-mail: andre.de-laure@univ-lille.fr

²Univ. Lille, Inria, CNRS, UMR 8524 - Laboratoire Paul Painlevé, F-59000 Lille, France.
E-mail: salvador.lopez-martinez@inria.fr

Dirac delta function, which leads to the standard Gross–Pitaevskii equation with nonvanishing condition at infinity, that has been intensively investigated (see e.g. [17, 34, 35, 39]).

To provide a clear mathematical context to the problem, it is useful to perform the change of variables $\Psi \rightarrow e^{-it}\bar{\Psi}$, which leads to the equation

$$i\partial_t\Psi = \partial_{xx}\Psi + \Psi(\mathcal{W} * (1 - |\Psi|^2)) \quad \text{in } \mathbb{R} \times \mathbb{R}, \quad (1.3)$$

where we assumed that $V * 1 = 1$ and denoted by \mathcal{W} the potential to make our following assumptions more clear. Here $*$ denotes the convolution in \mathbb{R} . We assume from now on that \mathcal{W} is a real-valued even tempered distribution. In this manner, (1.3) is Hamiltonian and its energy

$$E(\Psi(t)) = \frac{1}{2} \int_{\mathbb{R}} |\partial_x \Psi(t)|^2 dx + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (1 - |\Psi(t)|^2))(1 - |\Psi(t)|^2) dx,$$

is formally conserved. The (renormalized) momentum

$$p(\Psi(t)) = \int_{\mathbb{R}} \langle i\partial_x \Psi(t), \Psi(t) \rangle_{\mathbb{C}} \left(1 - \frac{1}{|\Psi(t)|^2} \right) dx,$$

is formally conserved too whenever $\inf_{x \in \mathbb{R}} |\Psi(x, t)| > 0$, where $\langle z_1, z_2 \rangle_{\mathbb{C}} = \text{Re}(z_1 \bar{z}_2)$, for $z_1, z_2 \in \mathbb{C}$ (see [19]).

We will be interested in special solutions to (1.3) with boundary condition (1.2), the so-called dark solitons. Roughly speaking, these are localized density notches that propagate without spreading [38]. They have been observed for example in Bose–Einstein condensates [5, 23]. More precisely, dark solitons in our context will be nontrivial finite energy solutions to (1.3) of the form

$$\Psi_c(x, t) = u(x - ct),$$

which represents a traveling wave with profile $u : \mathbb{R} \rightarrow \mathbb{C}$ propagating at speed $c \in \mathbb{R}$. Hence, the soliton u satisfies

$$icu' + u'' + u(\mathcal{W} * (1 - |u|^2)) = 0 \quad \text{in } \mathbb{R}. \quad (S(\mathcal{W}, c))$$

Notice that taking the complex conjugate of u in equation $(S(\mathcal{W}, c))$, we are reduced to the case $c \geq 0$.

By finite energy solution to $(S(\mathcal{W}, c))$ we mean a solution belonging to the energy space

$$\mathcal{E}(\mathbb{R}) = \{v \in H_{\text{loc}}^1(\mathbb{R}) : 1 - |v|^2 \in L^2(\mathbb{R}), v' \in L^2(\mathbb{R})\}.$$

This is justified by assuming that the Fourier transform of \mathcal{W} is bounded, i.e. that $\widehat{\mathcal{W}} \in L^\infty(\mathbb{R})$. Indeed, by Plancherel’s identity,

$$E(u) \leq \frac{1}{2} \|u'\|_{L^2(\mathbb{R})}^2 + \frac{1}{4} \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})} \|1 - |u|^2\|_{L^2(\mathbb{R})}^2.$$

We point out that any function in the energy space satisfies (1.2) (see Theorem 1.8 in [26]).

The simplest case for $(S(\mathcal{W}, c))$ that one may consider corresponds to the *contact* interaction $\mathcal{W} = \delta_0$. In this way, $(S(\delta_0, c))$ becomes the classical Gross–Pitaevskii equation, which is a local equation. In our one-dimensional case, $(S(\delta_0, c))$ can be solved explicitly. More precisely, as explained in [9], if $c \geq \sqrt{2}$ the only solutions in $\mathcal{E}(\mathbb{R})$ are the trivial ones (i.e. the constant functions of modulus one). On the contrary, if $0 \leq c < \sqrt{2}$, the nontrivial solutions in $\mathcal{E}(\mathbb{R})$ are given, up to invariances (translations and multiplication by constants of modulus one), by

$$u_c(x) = \sqrt{\frac{2 - c^2}{2}} \tanh\left(\frac{\sqrt{2 - c^2}}{2}x\right) - i\frac{c}{\sqrt{2}}. \quad (1.4)$$

Thus there is a family of dark solitons belonging to the nonvanishing energy space

$$\mathcal{NE}(\mathbb{R}) = \{v \in \mathcal{E}(\mathbb{R}) : \inf_{\mathbb{R}} |v| > 0\},$$

for $c \in (0, \sqrt{2})$. We refer to them as the *vortexless* solutions, as usual in nonlinear optics. There is also one stationary soliton that vanishes at exactly one point, associated with the speed $c = 0$, that is called the black soliton. Notice also that the values of $u_c(\infty)$ and $u_c(-\infty)$ are different if $c \neq 0$, and thus we cannot relax the condition (1.2) to $\lim_{|x| \rightarrow \infty} \Psi = 1$.

In the case of spatial dimension equal to two or three, the study of traveling waves for the contact interaction $\mathcal{W} = \delta_0$ started with numerical simulations in the Jones–Roberts program [34, 35]. There, it was observed numerically that finite energy traveling waves should exist for every $c \in [0, \sqrt{2})$, and should not otherwise. Rigorous proofs of these conjectures have been established by Béthuel and Saut [11], Béthuel, Gravejat and Saut [10], Mariş [47], Ruiz and Bellazzini [7], among others.

Despite the physical interest of the most realistic case where \mathcal{W} is a more general distribution, there are very few mathematical results concerning nonlocal interactions with nonzero conditions at infinity. To our knowledge, most of the mathematical results concerning the existence of traveling waves deal with functions vanishing at infinity (see e.g. [3, 6, 16, 42, 46]) and the techniques used in these works cannot be adapted to include solutions satisfying (1.2). Recently, de Laire and Mennuni [22] proved the existence of a branch of solutions to $(S(\mathcal{W}, c))$, by using a minimization approach. For $\mathfrak{q} \geq 0$, they consider the (nondecreasing) minimization curve

$$E_{\min}(\mathfrak{q}) := \inf\{E(v) : v \in \mathcal{E}(\mathbb{R}), p(v) = \mathfrak{q}\}, \quad (1.5)$$

and set

$$\mathfrak{q}_* = \sup\{\mathfrak{q} > 0 \mid \forall v \in \mathcal{E}(\mathbb{R}), E(v) \leq E_{\min}(\mathfrak{q}) \Rightarrow \inf_{\mathbb{R}} |v| > 0\}. \quad (1.6)$$

Under certain technical conditions on \mathcal{W} , they show that $\mathfrak{q}_* > 0.027$ and that for any $\mathfrak{q} \in (0, \mathfrak{q}_*)$, the minimum associated with $E_{\min}(\mathfrak{q})$ is attained and the corresponding Euler–Lagrange equation satisfied by the minimizers is exactly $(S(\mathcal{W}, c))$, where $c \in (0, \sqrt{2})$ appears as a Lagrange multiplier. In addition, their solutions are orbitally stable. Therefore, their result establishes the existence of a family of solutions to $(S(\mathcal{W}, c))$ parametrized by the momentum. This theorem applies for instance to the potential $\mathcal{W}_{\alpha, \beta} = \frac{\beta}{\beta - 2\alpha}(\delta_0 - \alpha e^{-\beta|x|})$, for $\beta > 2\alpha > 0$, which describes a strong repulsion force when two particles are in the same place, but an attractive force otherwise. However, the results in [22] do not apply to an interaction of the form $\mathcal{W}(x) = \exp(-\alpha x^2)$, $\alpha > 0$, since its Fourier transform grows exponentially in the complex plane. The first goal of this paper is to provide simple conditions on \mathcal{W} that guarantee the existence of nontrivial finite energy solutions to $(S(\mathcal{W}, c))$, covering a large variety of relevant nonlocal interactions, such as the Gaussian potential. The second one is to determine an optimal range for c (depending on \mathcal{W}) for which there exist finite energy traveling waves. Finally, a third goal is to establish the regularity and the decay of these solutions, and their nonexistence at critical speed.

1.2 Main results

From now on we assume that $\widehat{\mathcal{W}}$ satisfies the following minimal regularity assumption:

(H0) \mathcal{W} is an even tempered distribution such that $\widehat{\mathcal{W}} \in L^\infty(\mathbb{R})$.

Let us remark that the condition $\widehat{\mathcal{W}} \in L^\infty(\mathbb{R})$ is equivalent to the continuity of the application $\eta \in L^2(\mathbb{R}) \mapsto \mathcal{W} * \eta \in L^2(\mathbb{R})$ (see e.g. [27]). The parity assumption is necessary to have a variational formulation (see Lemma 3.1).

As explained in [22], the Bogoliubov dispersion relation [12] is given by

$$w(\xi) = \sqrt{\xi^4 + 2\widehat{\mathcal{W}}(\xi)\xi^2}. \quad (1.7)$$

We formally get $w(\xi) \approx (2\widehat{\mathcal{W}}(0))^{1/2}|\xi|$, for $\xi \approx 0$. The critical speed $c_*(\mathcal{W}) = (2\widehat{\mathcal{W}}(0))^{1/2}$ corresponds to the so-called speed of sound. It is conjectured that there is no nontrivial solution to $(S(\mathcal{W}, c))$ with finite energy when $c(\mathcal{W}) \geq c_*(\mathcal{W})$. Observe also that if $\widehat{\mathcal{W}}$ is continuous at the origin, then we can assume without loss of generality that \mathcal{W} fulfills the normalization condition (see [22]) $\widehat{\mathcal{W}}(0) = 1$, so that the speed of sound is well-defined and equal to $c_*(\mathcal{W}) = \sqrt{2}$.

Here and in what follows we use the convention that the Fourier transform of an integrable function is

$$\widehat{f}(\xi) = \int_{\mathbb{R}} e^{-ix\xi} f(x) dx. \quad (1.8)$$

In particular, the Fourier transform of the Dirac delta is $\widehat{\delta}_0 = 1$, so that $c_*(\delta_0) = \sqrt{2}$, and the nonexistence conjecture holds when $\mathcal{W} = \delta_0$, as explained before.

Our first result establishes the existence of dark solitons under the following assumption:

(H1) There exist $\sigma \in (0, 1]$ and $\kappa \in [0, 1/2)$ such that $\widehat{\mathcal{W}}(\xi) \geq \sigma - \kappa\xi^2$ a.e. on \mathbb{R} .

Theorem 1.1. *Assume that \mathcal{W} satisfies (H0) and (H1). Then there exists a nontrivial solution to $(S(\mathcal{W}, c))$ in $\mathcal{E}(\mathbb{R})$, for almost every $c \in (0, \sqrt{2\sigma})$.*

As an easy consequence of Theorem 1.1, we prove the following existence result for nonnegative potentials satisfying (H1) in the critical case $\kappa = 1/2$. This critical lower bound was already considered in [22].

Corollary 1.2. *Assume that \mathcal{W} satisfies (H0), with $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} , and that there is $\sigma \in (0, 1]$ such that $\widehat{\mathcal{W}}(\xi) \geq \sigma - \xi^2/2$, for a.e. $|\xi| \leq \sqrt{2\sigma}$. Then there exists a nontrivial solution to $(S(\mathcal{W}, c))$ in $\mathcal{E}(\mathbb{R})$, for almost every $c \in (0, \sqrt{2\sigma})$.*

Notice that there is no assumption on the continuity of \mathcal{W} . For instance, Theorem 1.1 applies to the potential

$$\mathcal{W} = 3\delta_0 - \frac{J_1(2|\cdot|^{1/2})}{2|\cdot|^{1/2}}, \quad \text{so that} \quad \widehat{\mathcal{W}}(\xi) = 2 + \cos(1/|\xi|), \quad \text{for all } \xi \in \mathbb{R} \setminus \{0\},$$

where J_1 is the Bessel function of first kind, with $\sigma = 1$ and $\kappa = 0$. This gives us the existence of nontrivial finite energy solutions to $(S(\mathcal{W}, c))$ for a.e. $c \in (0, \sqrt{2})$.

We can also apply Theorem 1.1 to the potential

$$\widehat{\mathcal{W}}_{a,b,\lambda}(\xi) = (1 + a\xi^2 + b\xi^4)e^{-\lambda\xi^2}, \quad (1.9)$$

that has been proposed in [8, 53] to describe a quantum fluid exhibiting a roton-maxon spectrum such as Helium 4. Indeed, as predicted by the Landau theory, in such a fluid, the dispersion curve (1.7) cannot be monotone, and it should have a local maximum and a local minimum, the so-called maxon and roton, respectively. In [22], some numerical simulations were done for $a = -36$, $b = 2687$, $\lambda = 30$, and a branch of solitons was found with speeds in $(0, \sqrt{2})$. These values are relevant because they provide the existence of a maxon and a roton. However, the existence theorem in [22] does not apply to this potential. On the other hand, it can be checked that, for these values of a , b and λ , condition (H1) is fulfilled with $\sigma = 0.175$ and any $\kappa \in (0, 1/2)$. Consequently, Theorem 1.1 provides the existence of nontrivial finite energy solutions to $(S(\mathcal{W}_{a,b,\lambda}, c))$ for a.e. $c \in (0, \sqrt{0.35})$.

The first step to prove Theorem 1.1 is to show that, if $c > 0$, then any solution $u \in \mathcal{E}(\mathbb{R})$ must belong to $\mathcal{N}\mathcal{E}(\mathbb{R})$, which allows us to lift the function as $u = \rho e^{i\theta}$, for some real-valued functions

ρ and θ . Then, we prove in Section 2 the crucial fact that $(S(\mathcal{W}, c))$ is actually equivalent to the nonlocal singular equation

$$-\rho'' + \frac{c^2}{4} \frac{1 - \rho^4}{\rho^3} = \rho \left(\mathcal{W} * (1 - \rho^2) \right) \quad \text{in } \mathbb{R}. \quad (1.10)$$

Therefore, we only need to look for a *real* solution $\rho \in \mathcal{NE}(\mathbb{R})$. Moreover, we know now that $\rho(x) = |u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$, so that we can suppose that ρ recast as $\rho = 1 + v$, for some $v \in H^1(\mathbb{R})$. The drawback is that we have introduced a singularity in the equation, we need thus to take care of the possible vanishing of functions on the variational approximation. Indeed, in Section 3 we will show that the solutions to (1.10) correspond to critical points of the functional $J_c : H^1(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ given by

$$J_c(1 - \rho) = \mathcal{A}(1 - \rho) - c^2 \mathcal{B}(1 - \rho), \quad \text{for } \rho \in 1 + H^1(\mathbb{R}),$$

where

$$\mathcal{A}(1 - \rho) = \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (1 - \rho^2))(1 - \rho^2) \quad \text{and} \quad \mathcal{B}(1 - \rho) = \frac{1}{8} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2}.$$

More precisely, it will be obtained as a mountain pass point. However, we cannot directly apply the classical version of the mountain pass theorem for several reasons. First, to handle the singularity of J_c , we do not work in H^1 , but in the nonvanishing open set

$$\mathcal{NV}(\mathbb{R}) = \{v \in H^1(\mathbb{R}) : 1 - v > 0 \text{ in } \mathbb{R}\},$$

i.e. we consider $\rho = 1 - v$ with $v \in \mathcal{NV}(\mathbb{R})$. Hence we need to verify that we can adapt the deformation lemma in this setting. Second, our formulation does not give the boundedness of Palais–Smale sequences. Nevertheless, we can apply the monotonicity trick introduced by Struwe [54] and generalized by Jeanjean [33], that, roughly speaking, will provide bounded Palais–Smale sequences for almost every $c \in (0, \sqrt{2}\sigma)$.

In order to prove existence in the whole subsonic regime, we have to restrict the potentials we work with; in particular, they will satisfy (H1) with $\sigma = 1$, so the subsonic interval becomes $(0, \sqrt{2})$. One of the new assumptions on the potentials implies some a priori estimates on the solutions. To see how this hypothesis is used, observe that, for any $c \in (0, \sqrt{2})$, one can apply Theorem 1.1 to get the existence of a sequence of speeds $\{c_n\}$ and a sequence of associated solutions $\{u_n\} \subset \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, c_n))$, such that $c_n \rightarrow c$. At this point, some a priori estimates allow us to conclude that $u_n \rightarrow u \in C_{\text{loc}}^k(\mathbb{R})$, so that u is a solution to $(S(\mathcal{W}, c))$. However, from our estimates, it is not clear that u has finite energy. Indeed, there are solutions with infinite energy to $(S(\mathcal{W}, c))$ that we need to avoid, for instance

$$u_r^\pm(x) = r \exp\left(ix \left(\frac{-c \pm \sqrt{c^2 + 4(1 - r^2)}}{2} \right)\right), \quad \text{for all } r \in (0, 1].$$

By imposing more restrictive conditions on the potential \mathcal{W} , we by-pass this difficulty in Section 4 by performing a more refined study of the Palais–Smale sequence and using the profile decomposition theorem for bounded functions in $H^1(\mathbb{R})$.

For the sake of simplicity, we only state in this introduction the existence result in the whole interval $(0, \sqrt{2})$ for potentials of the form

$$\mathcal{W}_f = A_f(\delta_0 + f), \quad \text{with } f \text{ even, } f \in L^1(\mathbb{R}; \mathbb{R}), \quad \|f^-\|_{L^1(\mathbb{R})} < 1 \text{ and } A_f = (1 + \widehat{f}(0))^{-1}. \quad (1.11)$$

Here we use the notation $f^+ = \max\{f, 0\}$ and $f^- = \min\{f, 0\}$. Notice that the normalization constant A_f^{-1} can be recast as $1 + \|f^+\|_{L^1(\mathbb{R})} - \|f^-\|_{L^1(\mathbb{R})}$, so that $A_f > 0$.

Theorem 1.3. *Let \mathcal{W}_f be as in (1.11). Assume that \widehat{f} is nondecreasing a.e. on \mathbb{R}^+ and that the map $\xi \mapsto \xi(\widehat{f})'(\xi)$ is bounded and continuous a.e. on \mathbb{R} . Then there exists a nontrivial solution to $(S(\mathcal{W}, c))$ in $\mathcal{NE}(\mathbb{R})$ for all $c \in (0, \sqrt{2})$.*

In Section 5 we study further properties of the solutions by considering the variable $\eta = 1 - |u|^2$, that satisfies the equation

$$-\eta'' + 2\mathcal{W} * \eta - c^2\eta = 2|u'|^2 + 2\eta(\mathcal{W} * \eta) := F. \quad (1.12)$$

From (1.12) we can deduce that every finite energy solution is smooth and that, if $c > 0$, then $|u|$ does not vanish, i.e. $\eta < 1$ on \mathbb{R} (see Proposition 2.2). On the other hand, equation (1.12) can be written as

$$M_c(\xi)\widehat{\eta}(\xi) = \widehat{F}(\xi), \quad \text{with } M_c(\xi) = \xi^2 + 2\widehat{\mathcal{W}}(\xi) - c^2. \quad (1.13)$$

If $c \in [0, \sqrt{2})$ and \mathcal{W} satisfies (H1), then M_c is strictly positive, so (1.12) is an elliptic equation. In this case, (1.12) can be written as the convolution equation:

$$\eta = \mathcal{L}_c * F, \quad \text{where } \widehat{\mathcal{L}}_c = M_c^{-1}.$$

In this way, $\widehat{\mathcal{L}}_c$ appears as a Fourier multiplier.

Let us remark that this kind of convolution formulation has been used in several contexts to get further properties of the solutions, see e.g. [18, 20, 29]. In our case, we can adapt and extend the classical theory of Bona–Li [13, 14] to handle the nonlocal function F , and to deduce the algebraic or exponential decay, and analyticity of the solutions depending on the properties of \mathcal{W} . For instance, our main result concerning the exponential decay reads as follows.

Theorem 1.4. *Assume that \mathcal{W} satisfies (H0). Let $c \geq 0$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Suppose that*

$$e^{m|\cdot|}\mathcal{L}_c \in L^p(\mathbb{R}) \quad \text{for some } m > 0, p \in (1, \infty]. \quad (1.14)$$

Then, for all $\ell \in [0, m)$, the function $\eta = 1 - |u|^2$ has the exponential decay:

$$\lim_{|x| \rightarrow \infty} e^{\ell|x|} D^k \eta(x) = 0, \quad \text{for all } k \in \mathbb{N}.$$

We refer to Section 5 for the precise statements for algebraic decay and the real analyticity of u , in the sense that $\text{Re}(u)$ and $\text{Im}(u)$ are real analytic functions.

We now discuss the nonexistence conjecture of nontrivial finite energy solution to $(S(\mathcal{W}, c))$ for $c \geq \sqrt{2}$. In the case $\mathcal{W} = \delta_0$, the proof in [9] uses the Cauchy–Lipschitz theorem for ODEs. Thus, this argument seems difficult to apply to the nonlocal equations $(S(\mathcal{W}, c))$ or (1.10). In the limit case $c = \sqrt{2}$, we can use (1.13) again to get the following nonexistence result.

Theorem 1.5. *Assume that \mathcal{W} satisfies (H0) and that $\widehat{\mathcal{W}}$ is of class \mathcal{C}^2 in a neighborhood of the origin, with $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} and $\widehat{\mathcal{W}}(0) = 1$. Suppose that either $(\widehat{\mathcal{W}})''(0) \neq -1$, or $(\widehat{\mathcal{W}})'' = -1$ on a neighborhood of the origin. Then $(S(\mathcal{W}, \sqrt{2}))$ admits no nontrivial solution in $\mathcal{E}(\mathbb{R})$.*

As a consequence of the real analyticity of the solutions to $(S(\mathcal{W}, c))$, we can deduce that the solutions obtained by minimization by de Laire and Mennuni in [22] are symmetric. By combining the analyticity with a reflection argument, we get the following result.

Corollary 1.6. *Let \mathcal{W} be a potential satisfying the hypotheses in Theorem 1 in [22]. Let $\mathbf{q} \in (0, \mathbf{q}_*)$ and let $u = \rho e^{i\theta} \in \mathcal{NE}(\mathbb{R})$ be the nontrivial solution to $(S(\mathcal{W}, c))$, for some $c \in (0, \sqrt{2})$, satisfying $p(u) = \mathbf{q}$, given by Theorem 1 in [22]. Then, up to translations, ρ is an even function and, up to multiplying u by a constant of modulus one, θ is an odd function.*

The uniqueness of solutions to $(S(\mathcal{W}, c))$ is a difficult problem due to the nonlocal potential. We do not know if the solutions to $(S(\mathcal{W}, c))$ are unique (up to invariances). Actually, the uniqueness for nonlocal equations such as $(S(\mathcal{W}, c))$ can be hard to establish (see e.g. [2, 43]). We cannot prove that our solutions are equal to the solutions in [22], except in the case $\mathcal{W} = \delta_0$ (where the solutions are explicitly given in (1.4)). However, we think that the uniqueness holds, at least for the potentials in the examples in the next subsection.

1.3 Examples

For $\beta > 2\alpha > 0$, we consider the potential

$$\mathcal{W}_{\alpha,\beta} = \frac{\beta}{\beta - 2\alpha}(\delta_0 - \alpha e^{-\beta|\cdot|}), \quad \text{so that} \quad \widehat{\mathcal{W}}_{\alpha,\beta}(\xi) = \frac{\beta}{\beta - 2\alpha} \left(1 - \frac{2\alpha\beta}{\xi^2 + \beta^2} \right), \quad \xi \in \mathbb{R}. \quad (1.15)$$

This kind of potential has been used in [49] for the study of dark solitons in a self-defocusing nonlocal Kerr-like medium. The kernel $\mathcal{W}_{\alpha,\beta}$ represents a strong repulsive interaction between particles that coincide in space, while the interaction becomes attractive otherwise, being this attraction significant at short distances.

From a mathematical point of view, this potential satisfies all the conditions to apply Theorems 1.3, 1.4, 1.5 and 5.12. The result reads as follows.

Theorem 1.7. *Let $\mathcal{W}_{\alpha,\beta}$ be given by (1.15) with $\beta > 2\alpha > 0$. Then for every $c \in (0, \sqrt{2})$, there exists a nontrivial solution $u_c \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}_{\alpha,\beta}, c))$. In addition, u_c is real-analytic, the limits $u_c(\pm\infty)$ exist, and there exists $\ell > 0$, depending only on c , α and β , such that $\eta_c = 1 - |u_c|^2$ satisfies*

$$\lim_{|x| \rightarrow \infty} e^{\ell|x|} D^k \eta_c(x) = 0, \quad \text{for all } k \in \mathbb{N}. \quad (1.16)$$

Furthermore, there is no nontrivial solution to $(S(\mathcal{W}_{\alpha,\beta}, \sqrt{2}))$ in $\mathcal{E}(\mathbb{R})$.

The following three examples will provide similar mathematical results, so we will gather them in a single theorem after some comments. The first one was proposed in [58] as simple model for interactions in a Bose–Einstein condensate. For $\lambda > 0$, it is given by a contact interaction δ_0 and two Dirac delta functions centered at $\pm\lambda$, as

$$\mathcal{W}_\lambda = 2\delta_0 - \frac{1}{2}(\delta_{-\lambda} + \delta_\lambda), \quad \text{so that} \quad \widehat{\mathcal{W}}_\lambda(\xi) = 2 - \cos(\lambda\xi), \quad \xi \in \mathbb{R}. \quad (1.17)$$

Notice that, as well as (1.15), (1.17) models a competition between repulsive and attractive interactions.

Another interesting example proposed in [45] is the Gaussian function,

$$\mathcal{W}_\lambda(x) = \frac{1}{2\lambda\sqrt{\pi}} e^{-\frac{x^2}{4\lambda^2}}, \quad \text{so that} \quad \widehat{\mathcal{W}}_\lambda(\xi) = e^{-\lambda\xi^2}, \quad \xi \in \mathbb{R}. \quad (1.18)$$

where $\lambda > 0$. In fact, for $\lambda > 0$ small, this potential represents a smooth approximation of the Dirac delta.

Finally, we introduce the so-called soft core potential, which was used in [1, 36] to study supersolids. It can be seen as a nonsmooth approximation of the Dirac delta when $\lambda > 0$ is small,

$$\mathcal{W}_\lambda(x) = \begin{cases} \frac{1}{2\lambda}, & \text{if } |x| < \lambda, \\ 0, & \text{otherwise,} \end{cases} \quad \text{so that} \quad \widehat{\mathcal{W}}_\lambda(\xi) = \frac{\sin(\lambda\xi)}{\lambda\xi}, \quad \xi \in \mathbb{R}. \quad (1.19)$$

Unlike Theorem 1.7, for this three potentials we prove existence of nontrivial finite energy traveling waves for *almost every* $c \in (0, \sqrt{2})$. We summarize our main results for (1.17), (1.18) and (1.19) as follows.

Theorem 1.8. *Assume that one of the following cases holds.*

- (i) \mathcal{W}_λ is given by (1.17) with $0 < \lambda$.
- (ii) \mathcal{W}_λ is given by (1.18) with $0 < \lambda < 1/2$.
- (iii) \mathcal{W}_λ is given by (1.19) with $0 < \lambda < \sqrt{3}$.

Then, for almost every $c \in (0, \sqrt{2})$, there exists a nontrivial solution $u_c \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}_\lambda, c))$. In addition, u_c is real-analytic, the limits $u_c(\pm\infty)$ exist, and there exists $\ell > 0$, depending only on c and λ , such that the function $\eta_c = 1 - |u_c|^2$ satisfies

$$\lim_{|x| \rightarrow \infty} e^{\ell|x|} D^k \eta_c(x) = 0, \quad \text{for all } k \in \mathbb{N}. \quad (1.20)$$

Moreover, in the cases (i) and (ii), there is no nontrivial solution to $(S(\mathcal{W}_\lambda, \sqrt{2}))$ in $\mathcal{E}(\mathbb{R})$.

Some comments on this theorem are in order.

- If the Fourier transform of the potential is nonnegative, we can also apply Corollary 1.2 to study other ranges of λ . For instance, if \mathcal{W}_λ is given by (1.18), a simple computation shows that for $\lambda \geq 1/2$, we have the estimate $\widehat{\mathcal{W}}_\lambda(\xi) \geq \sigma_\lambda - \xi^2/2$, for all $\xi \in \mathbb{R}$, where $\sigma_\lambda = \frac{1+\ln(2\lambda)}{2\lambda}$. Therefore, we can deduce the existence of nontrivial solitons for a.e. $c \in (0, \sqrt{2\sigma_\lambda})$.
- If \mathcal{W}_λ is given by (1.19), then $\widehat{\mathcal{W}}_\lambda$ changes sign. For this reason we cannot guarantee nonexistence of finite energy traveling waves for the critical speed $c = \sqrt{2}$ in Theorem 1.8.
- Filling the existence in the complete interval $(0, \sqrt{2})$ in Theorem 1.8 would require to prove several estimates, a task far from being trivial without our assumptions (H3) and (H5) (see Section 4 for details).
- The arguments in the proof of Theorem 1.8 also apply to the potential (1.9), with the values $a = -36$, $b = 2687$, $\lambda = 30$. Therefore, for a.e. $c \in (0, \sqrt{0.35})$, the solution u_c is real-analytic and decays exponentially at infinity. Also, there is no nontrivial solution with critical speed $c = \sqrt{2}$ in $\mathcal{E}(\mathbb{R})$.

The last example that we consider is given, for $\kappa \in (0, 1/2]$, by

$$\mathcal{W}_\kappa(x) = \frac{2\kappa}{\pi x^2} \left(\frac{\sin(x/\sqrt{\kappa})}{x} - \frac{1}{\sqrt{\kappa}} \cos(x/\sqrt{\kappa}) \right), \quad \text{so that } \widehat{\mathcal{W}}_\kappa(\xi) = (1 - \kappa\xi^2)^+, \quad \xi \in \mathbb{R}. \quad (1.21)$$

It is simple to check that \mathcal{W}_κ is bounded continuous, with $\mathcal{W}_\kappa \in L^1(\mathbb{R})$. From a mathematical point of view, this is an interesting example since it represents the limiting case among the normalized potentials (i.e. $\widehat{\mathcal{W}}(0) = 1$) satisfying (H1) with nonnegative Fourier transform. This kernel also appears in the Bochner–Riesz means, and it is important in the Fourier multipliers theory.

Due to the lack of regularity of $\widehat{\mathcal{W}}_\kappa$, we do not expect an exponential decay of the solution. Nevertheless, we will show that, in this case, \mathcal{L}_c decays as $1/x^2$, which will lead us to the following result.

Theorem 1.9. *Let \mathcal{W}_κ be given by (1.21), with $\kappa \in (0, 1/2]$. Then for almost every $c \in (0, \sqrt{2})$, there exists a nontrivial solution $u_c \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}_\kappa, c))$. In addition, u_c is real-analytic, the limits $u_c(\pm\infty)$ exist, and the function $\eta_c = 1 - |u_c|^2$ satisfies the following algebraic decay*

$$|\cdot|^\ell D^k \eta_c \in L^1(\mathbb{R}), \quad \lim_{|x| \rightarrow \infty} |x|^\ell D^k \eta_c(x) = 0, \quad \text{for all } \ell \in [0, 1), \quad \text{for all } k \in \mathbb{N}. \quad (1.22)$$

Moreover, there is no nontrivial solution to $(S(\mathcal{W}_\kappa, \sqrt{2}))$ in $\mathcal{E}(\mathbb{R})$.

We will study numerically the solitons given in this subsection in the forthcoming paper [21].

1.4 Notation

The usual real-valued Sobolev and Lebesgue spaces will be denoted, respectively, by $W^{k,p}(\mathbb{R})$ and $L^p(\mathbb{R})$ for $p \in [1, \infty]$ and $k \in \mathbb{N}$. Moreover, $W^{k,2}(\mathbb{R}) = H^k(\mathbb{R})$. The notation for the Lebesgue spaces of complex-valued functions will be $L^p(\mathbb{R}; \mathbb{C})$, and analogously for the Sobolev spaces of complex-valued functions, or simply $L^p(\mathbb{R})$ if there is no ambiguity. For a real-valued function f , we write $f^+ = \max\{f, 0\}$ and $f^- = \min\{f, 0\}$, so that $f = f^+ + f^-$.

In this paper we *always* assume that \mathcal{W} satisfies (H0). In particular, this implies that

$$\|\mathcal{W} * f\|_{L^2(\mathbb{R})} \leq \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})} \|f\|_{L^2(\mathbb{R})}, \quad \text{for all } f \in L^2(\mathbb{R}; \mathbb{C}), \quad (1.23)$$

and that Plancherel's identity reads, with our convention for the Fourier transform,

$$\int_{\mathbb{R}} (\mathcal{W} * f)g = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{\mathcal{W}}(\xi) \widehat{f}(\xi) \overline{\widehat{g}}(\xi), \quad \text{for all } f, g \in L^2(\mathbb{R}; \mathbb{C}). \quad (1.24)$$

2 Some identities

We start recalling that any finite energy solution to $(S(\mathcal{W}, c))$ is smooth and admits a lifting at infinity (without restriction on c). This result corresponds to Corollary 2.4 in [18], where it was proved in dimension greater or equal than two, but the same proof applies in our one-dimensional setting.

Lemma 2.1. *Let $c \geq 0$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Then u is bounded and of class $\mathcal{C}^\infty(\mathbb{R})$. Moreover, $\eta := 1 - |u|^2$ and u' belong to $W^{k,p}(\mathbb{R})$, for all $k \in \mathbb{N}$ and $p \in [2, \infty]$. Furthermore, there exists a smooth lifting of u . More precisely, there exist $R > 0$, $\delta \in (0, 1)$ and $\theta \in \mathcal{C}^\infty((-R, R)^c)$ such that $u = \rho e^{i\theta}$ on $(-R, R)^c$, with $\rho \geq \delta$ on $(-R, R)^c$. In particular, $\theta', 1 - \rho \in W^{k,p}((-R, R)^c)$ for all $k \in \mathbb{N}$, $p \in [2, \infty]$, and*

$$\rho(\pm\infty) = 1, \quad D^j u(\pm\infty) = D^j \rho(\pm\infty) = D^j \theta(\pm\infty) = D^j \eta(\pm\infty) = 0, \quad \text{for all } j \geq 1. \quad (2.1)$$

Finally, if $u \in \mathcal{NE}(\mathbb{R})$, then the above conclusions still hold true in \mathbb{R} , i.e. for $R = 0$.

Proof. The proof is contained in [18] except for the regularity of $1 - \rho$. Notice that

$$1 - \rho = \frac{1 - \rho^2}{1 + \rho} = \frac{\eta}{1 + \rho} \quad \text{on } \mathbb{R}. \quad (2.2)$$

Therefore, $|1 - \rho| \leq |\eta|$, so $1 - \rho \in L^p(\mathbb{R})$ for every $p \in [2, \infty]$. Moreover, using that $|u| \geq \delta$ on $(-R, R)^c$ and also Cauchy-Schwarz inequality, we deduce that

$$|\rho'| = \frac{|\langle u, u' \rangle|}{|u|} \leq |u'| \quad \text{on } (-R, R)^c.$$

Thus $1 - \rho \in W^{1,p}((-R, R)^c)$ for all $p \in [2, \infty]$. Similarly, using that

$$\rho'' = \frac{|u'|^2 + \langle u, u'' \rangle}{|u|} - \frac{|\langle u, u' \rangle|^2}{|u|^3} \quad \text{on } (-R, R)^c,$$

and that $u, u' \in L^\infty(\mathbb{R})$, we conclude that $\rho'' \in W^{2,p}((-R, R)^c)$ for all $p \in [2, \infty]$. Repeating the previous arguments inductively, we arrive up to $1 - \rho \in W^{k,p}((-R, R)^c)$ for all $k \in \mathbb{N}$ and $p \in [2, \infty]$. Finally, if $u \in \mathcal{NE}(\mathbb{R})$, then $\rho \geq \delta$ on \mathbb{R} for some $\delta \in (0, 1)$, so that the previous arguments are valid on \mathbb{R} . \square

We now establish some key identities in terms of $\eta = 1 - |u|^2$. In particular, we derive equation (1.13) and we deduce that, if $c > 0$, the finite energy solutions to $(S(\mathcal{W}, c))$ do not vanish. Notice that $\eta \leq 1$ on \mathbb{R} , but η could be negative.

Proposition 2.2. *Let $c \geq 0$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Setting $K = |u'|^2$ and $\eta = 1 - |u|^2$, the following identities are satisfied on \mathbb{R} :*

$$\frac{c}{2}\eta = -\langle iu', u \rangle, \quad (2.3)$$

$$-\eta'' + 2\mathcal{W} * \eta - c^2\eta = 2K + 2\eta(\mathcal{W} * \eta), \quad (2.4)$$

$$K' = \eta'(\mathcal{W} * \eta), \quad (2.5)$$

$$c^2\eta^2 + (\eta')^2 = 4K(1 - \eta). \quad (2.6)$$

As a consequence, if $c > 0$, then $\eta < 1$ on \mathbb{R} , $u \in \mathcal{NE}(\mathbb{R})$ and

$$2K = \frac{c^2\eta^2}{2(1 - \eta)} + \frac{(\eta')^2}{2(1 - \eta)} \quad \text{on } \mathbb{R}. \quad (2.7)$$

Notice that (2.4) corresponds to equation (1.13), that we will use in Section 5 to establish the decay and analyticity of solutions, as well as the nonexistence for the critical speed.

Proof of Proposition 2.2. Let $u = u_1 + iu_2$. Taking real and imaginary parts in $(S(\mathcal{W}, c))$, we obtain

$$u_1'' - cu_2' + u_1(\mathcal{W} * (1 - u_1^2 - u_2^2)) = 0, \quad (2.8)$$

$$u_2'' + cu_1' + u_2(\mathcal{W} * (1 - u_1^2 - u_2^2)) = 0. \quad (2.9)$$

Multiplying (2.8) by u_2 and (2.9) by u_1 , we get

$$\frac{c}{2}\eta' = (u_1u_2' - u_2u_1')'.$$

Integrating over \mathbb{R} and taking into account (2.1) yields

$$\frac{c}{2}\eta = u_1u_2' - u_2u_1',$$

which is exactly (2.3).

On the other hand, multiplying (2.8) by u_1 and (2.9) by u_2 , and using (2.3), we deduce that

$$u_1''u_1 + u_2''u_2 = \frac{c^2}{2}\eta - (1 - \eta)(\mathcal{W} * \eta). \quad (2.10)$$

Hence, by differentiating,

$$\eta'' = -2|u'|^2 - 2\langle u, u'' \rangle = -2K - 2(u_1''u_1 + u_2''u_2), \quad (2.11)$$

which allows us to obtain (2.4) by combining (2.10) and (2.11).

Let us now multiply (2.8) by u_1' and (2.9) by u_2' . This gives

$$u_1'u_1'' + u_2'u_2'' + (u_1u_1' + u_2u_2')(\mathcal{W} * \eta) = 0.$$

Therefore, (2.5) follows directly by the definitions of η and K .

In order to show (2.6), let us multiply both sides of (2.4) by $2\eta'$. Taking (2.5) into account, it is easy to see that

$$-((\eta')^2)' + 4K' - c^2(\eta^2)' = 4(K\eta)'$$

Hence, (2.6) follows by integrating this equality.

Let us now show that if $c > 0$, then $\eta < 1$ on \mathbb{R} . Since $\eta \leq 1$ on \mathbb{R} , we assume by contradiction that there exists $x_0 \in \mathbb{R}$ such that $\eta(x_0) = 1$. Then, x_0 is a maximum of η , so that $\eta'(x_0) = 0$, and substituting into (2.6) we get $c^2 = 0$, a contradiction.

Finally, equation (2.7) is an immediate consequence of (2.6). \square

Remark 2.3. Recall that the energy functional is defined by

$$E(u) = \frac{1}{2} \int_{\mathbb{R}} |u'|^2 + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * \eta) \eta \quad \text{for all } u \in \mathcal{E}(\mathbb{R}),$$

where $\eta = 1 - |u|^2$. Moreover, the momentum reads

$$p(u) = -\frac{1}{2} \int \langle iu', u \rangle \frac{\eta}{1 - \eta}.$$

Taking (2.3) and (2.7) into account, we see that the energy and the momentum of any *solution* $u \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c))$, with $c > 0$, can be written only in terms of η and c as

$$E(u) = \frac{c^2}{8} \int_{\mathbb{R}} \frac{\eta^2}{1 - \eta} + \frac{1}{8} \int_{\mathbb{R}} \frac{(\eta')^2}{1 - \eta} + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * \eta) \eta \quad \text{and} \quad p(u) = \frac{c}{4} \int_{\mathbb{R}} \frac{\eta^2}{1 - \eta}.$$

Furthermore, since $\eta = 1 - \rho^2$ and $\eta' = -2\rho\rho'$, we also have expressions for the energy and the momentum of a solution $u \in \mathcal{NE}(\mathbb{R})$ in terms of ρ ,

$$E(u) = \frac{c^2}{8} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2} + \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (1 - \rho^2))(1 - \rho^2) \quad \text{and} \quad p(u) = \frac{c}{4} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2}.$$

That is the key observation in order to establish the variational framework in Section 3.

The next result gives an essential reformulation of the complex-valued $(S(\mathcal{W}, c))$ for vortexless solutions. In this case, we can reduce the problem to a single real-valued equation.

Proposition 2.4. *Let $c \geq 0$. If $u = \rho e^{i\theta} \in \mathcal{NE}(\mathbb{R})$ is a solution to $(S(\mathcal{W}, c))$, then*

$$\theta' = \frac{c}{2} \left(\frac{1}{\rho^2} - 1 \right) \quad \text{on } \mathbb{R}, \tag{2.12}$$

$$-\rho'' + \frac{c^2}{4} \frac{1 - \rho^4}{\rho^3} = \rho (\mathcal{W} * (1 - \rho^2)) \quad \text{on } \mathbb{R}. \tag{2.13}$$

Reciprocally, let $\rho \in \mathcal{C}^2(\mathbb{R})$ be such that $\rho > 0$ on \mathbb{R} and assume that it satisfies (2.13). For any $a \in \mathbb{R}$, let us define

$$\theta(x) = \frac{c}{2} \int_a^x \left(\frac{1}{\rho(y)^2} - 1 \right) dy \quad \text{for all } x \in \mathbb{R}. \tag{2.14}$$

Then, the function $u = \rho e^{i\theta}$ belongs to $\mathcal{C}^2(\mathbb{R}; \mathbb{C})$ and is a solution to $(S(\mathcal{W}, c))$. If in addition $1 - \rho \in H^1(\mathbb{R})$, then $u \in \mathcal{NE}(\mathbb{R})$.

Proof. Let $u = \rho e^{i\theta}$, with $|u| > 0$ on \mathbb{R} and $\theta \in \mathcal{C}^2(\mathbb{R})$. By computing the derivatives of u and taking real and imaginary parts, we check that u satisfies $(S(\mathcal{W}, c))$ if and only if the couple (ρ, θ) satisfies

$$\begin{cases} -c\theta'\rho + \rho'' - \rho(\theta')^2 + \rho(\mathcal{W} * (1 - \rho^2)) = 0 & \text{on } \mathbb{R}, \\ c\rho' + 2\theta'\rho' + \theta''\rho = 0 & \text{on } \mathbb{R}. \end{cases} \tag{2.15}$$

Let $u = \rho e^{i\theta} \in \mathcal{NE}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. By Lemma 2.1, we have $\rho, \theta \in C^\infty(\mathbb{R})$, so that (ρ, θ) satisfies (2.15). By multiplying the second equation in (2.15) by ρ , we obtain

$$(c\rho^2 + 2\theta'\rho^2)' = 0. \quad (2.16)$$

Bearing in mind (2.1), we can integrate (2.16) and obtain (2.12). Plugging (2.12) into the first equation of (2.15), we get (2.13).

We turn now to the second part of the result. Indeed, let $\rho \in C^2(\mathbb{R})$ be such that $\rho > 0$ on \mathbb{R} . Assume that ρ satisfies (2.13) and consider θ defined by (2.14). Then, one may immediately check that the equations (2.15) are satisfied. Hence, $u = \rho e^{i\theta}$ is a solution to $(S(\mathcal{W}, c))$. It only remains to verify that $u \in \mathcal{NE}(\mathbb{R})$ if $1 - \rho \in H^1(\mathbb{R})$. Indeed, by the Sobolev embedding theorem, we get $\rho \in L^\infty(\mathbb{R})$, with $\rho(\pm\infty) = 1$ and $\rho \geq \delta$ on \mathbb{R} for some $\delta \in (0, 1)$. Thus

$$1 - |u|^2 = 1 - \rho^2 = (1 - \rho)(1 + \rho) \in L^2(\mathbb{R}).$$

Moreover, by definition of θ ,

$$|u'|^2 = (\rho')^2 + \rho^2(\theta')^2 = (\rho')^2 + \frac{c^2}{4\rho^2}(1 - \rho^2)^2,$$

which also belongs to $L^2(\mathbb{R})$, since $(1 - \rho^2)^2 = (1 - \rho)^2(1 + \rho)^2$ and $\rho \in L^\infty(\mathbb{R})$. \square

In view of Proposition 2.4, the problem of existence of vortexless finite energy solution to $(S(\mathcal{W}, c))$ is reduced to the existence of positive solution to (2.13) with $1 - \rho \in H^1(\mathbb{R})$. Abusing of the concept of energy, we will say that a solution to (2.13) has finite if $\rho \in 1 + H^1(\mathbb{R})$.

3 The variational formulation

In this section we introduce a variational formulation that will lead to the proof of Theorem 1.1. Formally speaking, it is showed in [22] that critical points of the functional $E(u) - cp(u)$ are (complex-valued) solutions to $(S(\mathcal{W}, c))$. Thanks to Proposition 2.4, we may simplify the setting and work in a space of *real-valued* functions. More precisely, we will find solutions to (2.13) as critical points of the functional $J_c : H^1(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ formally defined by

$$J_c(1 - \rho) = \mathcal{A}(1 - \rho) - c^2\mathcal{B}(1 - \rho), \quad \text{for } \rho \in 1 + H^1(\mathbb{R}),$$

where

$$\mathcal{A}(1 - \rho) = \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 + \frac{1}{4} (\mathcal{W} * (1 - \rho^2))(1 - \rho^2) \quad \text{and} \quad \mathcal{B}(1 - \rho) = \frac{1}{8} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2}.$$

It is easy to see, thanks to Remark 2.3, that for every solution $u \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c))$, the equality $J_c(1 - \rho) = E(u) - cp(u)$ holds, where $\rho = |u|$.

Notice that if $\rho \in 1 + H^1(\mathbb{R})$ with $\rho \geq 0$, then $1 - \rho \in L^2(\mathbb{R})$ iff $1 - \rho^2 \in L^2(\mathbb{R})$ by (2.2). To avoid ambiguities in the definition, we will restrict J_c to the nonvanishing set

$$\mathcal{NV}(\mathbb{R}) = \{v \in H^1(\mathbb{R}) : 1 - v > 0 \text{ in } \mathbb{R}\},$$

which is an open in $H^1(\mathbb{R})$ due to the continuous embedding $H^1(\mathbb{R}) \subset L^\infty(\mathbb{R})$. Thus J_c is defined in the variable $v = 1 - \rho \in \mathcal{NV}(\mathbb{R})$ by

$$J_c(v) = \frac{1}{2} \int_{\mathbb{R}} (v')^2 + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (v(2 - v)))v(2 - v) - \frac{c^2}{8} \int_{\mathbb{R}} \frac{v^2(2 - v)^2}{(1 - v)^2}.$$

It is not difficult to show that the functional satisfies a mountain pass geometry (see Lemmas 3.5 and 3.6). However, it is not clear at all that the Palais–Smale sequences are bounded. In order to overcome this issue, we take advantage of the “monotonicity trick” of Struwe [54]. More precisely, we are deeply inspired by the work of Jeanjean [33]. We adapt some of his results, since several nontrivial modifications are needed due to the singular behavior of J_c . This way, we are able to obtain *bounded* Palais–Smale sequences for *almost every* speed $c \in (0, \sqrt{2})$.

We start by showing that J_c is smooth on $\mathcal{NV}(\mathbb{R})$ and that the critical points provide solutions to $(S(\mathcal{W}, c))$.

Lemma 3.1. *Let $c > 0$. The functional J_c is of class $\mathcal{C}^2(\mathcal{NV}(\mathbb{R}))$. Moreover, for any $v \in \mathcal{NV}(\mathbb{R})$, its Fréchet derivatives are given by*

$$J'_c(v)(\phi) = \int_{\mathbb{R}} v' \phi' + \int_{\mathbb{R}} (\mathcal{W} * f(v))(1-v)\phi - c^2 \int_{\mathbb{R}} h(v)\phi, \quad (3.1)$$

$$J''_c(v)(\phi, \psi) = \int_{\mathbb{R}} \phi' \psi' - c^2 \int_{\mathbb{R}} h'(v)\phi\psi + \int_{\mathbb{R}} (\mathcal{W} * (f'(v)\psi))(1-v)\phi - \int_{\mathbb{R}} (\mathcal{W} * f(v))\phi\psi, \quad (3.2)$$

for all $\phi, \psi \in H^1(\mathbb{R})$, where $f(s) = s(2-s)$ and $h(s) = \frac{s(2-s)(s^2-2s+2)}{4(1-s)^3}$ for all $s < 1$.

Remark 3.2. Observe that if $v \in \mathcal{NV}(\mathbb{R}) \setminus \{0\}$ satisfies $J'_c(v) = 0$, then

$$-v'' + (\mathcal{W} * f(v))(1-v) - c^2 h(v) = 0 \quad \text{on } \mathbb{R}. \quad (3.3)$$

Hence, setting $\rho = 1-v$ and noticing that $h(1-\rho) = (1-\rho^4)/(4\rho^3)$ and that $f(1-\rho) = 1-\rho^2$, we conclude that ρ is a nontrivial solution to (2.13). Therefore, by Proposition 2.4, this provides a nontrivial finite energy solution u to $(S(\mathcal{W}, c))$.

Proof of Lemma 3.1. First, we recall that since \mathcal{W} is even, we have

$$\int_{\mathbb{R}} (\mathcal{W} * g_1)g_2 = \int_{\mathbb{R}} (\mathcal{W} * g_2)g_1, \quad \text{for all } g_1, g_2 \in L^2(\mathbb{R}). \quad (3.4)$$

In order to differentiate the nonlocal term of J_c , by using the dominated convergence theorem, we conclude that the functional $v \in H^1(\mathbb{R}) \mapsto \int_{\mathbb{R}} (\mathcal{W} * f(v))f(v) \in \mathbb{R}$ admits a Fréchet derivative, given by

$$2 \int_{\mathbb{R}} (\mathcal{W} * f(v))f'(v)\phi, \quad \text{for all } \phi \in H^1(\mathbb{R}).$$

Therefore, we easily deduce (3.1). Computing (3.2) is also straightforward.

In order to prove the continuity of J'_c in $\mathcal{NV}(\mathbb{R})$, let $v \in \mathcal{NV}(\mathbb{R})$ and consider a sequence $\{v_n\} \subset H^1(\mathbb{R})$ such that $v_n \rightarrow v$ in $H^1(\mathbb{R})$. Then, $v_n \rightarrow v$ in $L^\infty(\mathbb{R})$. In particular, since there exists $\delta \in (0, 1)$ such that $v \leq 1-\delta$ on \mathbb{R} , it follows that $v_n \leq 1-\delta/2$ on \mathbb{R} , for n large enough. Thus, $h'(v_n) \rightarrow h'(v)$ in $L^2(\mathbb{R})$ by using the dominated convergence theorem. The continuity of the other terms is standard. \square

To apply a mountain pass argument, we need to invoke a deformation lemma. Although there are many versions of this classical lemma, we did not find one that fits in our framework since our functional is well-defined only on the open set $\mathcal{NV}(\mathbb{R})$, and not in the whole space. For this reason, we give here a modification of Lemma 2.3 in [59] that can be applied to our purposes. Furthermore, such a version does not require any Palais–Smale condition. For the sake of completeness, we also include its proof in the appendix.

Lemma 3.3. *Let $c > 0$. For some $R > 0$ and for every $\delta \in (0, 1)$, let us consider the set*

$$Z_\delta = \{v \in \mathcal{NV}(\mathbb{R}) : \|v\|_{H^1(\mathbb{R})} \leq R + 1 - \delta, v \leq 1 - \delta \text{ in } \mathbb{R}\}. \quad (3.5)$$

Assume that there exist constants $0 < \delta_1 < \delta_2 < \delta_3 < 1$, $\varepsilon > 0$ and $\gamma \in \mathbb{R}$ such that

$$\|J'_c(v)\|_{H^{-1}(\mathbb{R})} \geq \frac{2\varepsilon}{\delta_3 - \delta_2}, \quad \text{for all } v \in J_c^{-1}([\gamma - 2\varepsilon, \gamma + 2\varepsilon]) \cap Z_{\delta_1}.$$

Then there exists a continuous function $h : [0, 1] \times \mathcal{NV}(\mathbb{R}) \rightarrow \mathcal{NV}(\mathbb{R})$ such that

- (i) $h(0, v) = v$, for all $v \in \mathcal{NV}(\mathbb{R})$,
- (ii) $h(t, v) = v$, for all $v \in \mathcal{NV}(\mathbb{R}) \setminus (J_c^{-1}([\gamma - 2\varepsilon, \gamma + 2\varepsilon]) \cap Z_{\delta_1})$, for all $t \in [0, 1]$,
- (iii) $h(t, Z_{\delta_3}) \subset Z_{\delta_2}$, for all $t \in [0, 1]$,
- (iv) $J_c(h(t, v)) \leq J_c(v)$, for all $v \in \mathcal{NV}(\mathbb{R})$, for all $t \in [0, 1]$,
- (v) $h(1, J_c^{\gamma+\varepsilon} \cap Z_{\delta_3}) \subset J_c^{\gamma-\varepsilon} \cap Z_{\delta_2}$,

where $J_c^d = J_c^{-1}((-\infty, d])$ for every $d \in \mathbb{R}$.

Remark 3.4. To simplify the statement and the proof of Lemma 3.3, we used the sharp constant in the Sobolev embedding (see [15, p. 138, Theorem 4]), so that

$$\|v\|_{L^\infty(\mathbb{R})} \leq \frac{1}{2}(\|v\|_{L^2(\mathbb{R})} + \|v'\|_{L^2(\mathbb{R})}) \leq \sqrt{\|v\|_{L^2(\mathbb{R})}^2 + \|v'\|_{L^2(\mathbb{R})}^2} = \|v\|_{H^1(\mathbb{R})}, \quad (3.6)$$

for all $v \in H^1(\mathbb{R})$.

The following two lemmata provide the mountain pass geometry of J_c .

Lemma 3.5. Assume that \mathcal{W} satisfies (H1), and let $c \in (0, \sqrt{2\sigma})$. Then, there is a constant $r_c > 0$ such that, for every $r \in (0, r_c]$, there exists $l_r > 0$, depending on κ , σ , c and r , such that $J_c(1 - \rho) \geq l_r$ for every $\rho \in 1 + H^1(\mathbb{R})$ with $\|1 - \rho\|_{H^1(\mathbb{R})} = r$.

Proof. Let $\rho \in 1 + H^1(\mathbb{R})$ be such that $\|1 - \rho\|_{H^1(\mathbb{R})} \leq r_c$ for some $r_c \in (0, 1)$ to be chosen later, and let $\eta = 1 - \rho^2$. By applying Plancherel's identity and (H1), we deduce that

$$J_c(1 - \rho) \geq \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 + \frac{1}{8\pi} \int_{\mathbb{R}} (\sigma - \kappa\xi^2) |\widehat{\eta}(\xi)|^2 d\xi - \frac{c^2}{8} \int_{\mathbb{R}} \frac{\eta^2}{\rho^2}.$$

Using that

$$\frac{1}{2\pi} \int_{\mathbb{R}} (\sigma - \kappa\xi^2) |\widehat{\eta}(\xi)|^2 d\xi = \sigma \int_{\mathbb{R}} \eta^2 - \kappa \int_{\mathbb{R}} (\eta')^2,$$

and that $\eta' = -2\rho\rho'$, we get the lower bound

$$J_c(1 - \rho) \geq \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 (1 - 2\kappa\rho^2) + \frac{1}{4} \int_{\mathbb{R}} \left(\sigma - \frac{c^2}{2\rho^2} \right) \eta^2.$$

By invoking (3.6), we see that $1 - r_c \leq \rho \leq 1 + r_c$ on \mathbb{R} . Thus, recalling that $2\kappa < 1$ and $c^2 < \sigma$, we choose $r_c > 0$ small enough so that $1 - 2\kappa(1 + r_c)^2 > 0$ and $\sigma - \frac{c^2}{2(1-r_c)^2} > 0$. Consequently, using that $\eta^2 = (1 - \rho)^2(1 + \rho)^2 \geq (1 - \rho)^2$, we finally obtain

$$J_c(1 - \rho) \geq \ell_r \|1 - \rho\|_{H^1(\mathbb{R})}^2, \quad \text{with } \ell_r = \min \left\{ \frac{1 - 2\kappa(1 + r_c)^2}{2}, \frac{1}{4} \left(\sigma - \frac{c^2}{2(1 - r_c)^2} \right) \right\}.$$

The result follows by choosing $l_r = \ell_r r^2$. □

Lemma 3.6. *For every $\mathfrak{c} > 0$, there exists $\phi_{\mathfrak{c}} \in 1 + H^1(\mathbb{R})$ such that $J_c(1 - \phi_{\mathfrak{c}}) \in (-\infty, 0)$ for every $c \geq \mathfrak{c}$.*

Proof. For any $\delta \in (0, 1)$ and $r > 0$ to be chosen later, let us consider a nonnegative even function ϕ , with $0 \leq 1 - \phi^2 \leq 1 - \delta$ in \mathbb{R} , satisfying the following properties:

$$\phi^2 = \delta \text{ in } [0, r], \quad \phi = 1 \text{ in } [r + 1, \infty), \quad \phi(x + r) = \psi(x), \text{ for all } x \in [0, 1],$$

where $\psi : \mathbb{R} \rightarrow \mathbb{R}$ is a function independent of r that we choose such that $\phi \in C^\infty(\mathbb{R})$. In particular, $1 - \phi \in H^1(\mathbb{R})$. Thus, by Plancherel's identity, we get for all $c \geq \mathfrak{c}$,

$$\begin{aligned} J_c(1 - \phi) &\leq \frac{1}{2} \int_{\mathbb{R}} (\phi')^2 + \frac{\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}}{4} \int_{\mathbb{R}} (1 - \phi^2)^2 - \frac{\mathfrak{c}^2}{8} \int_{\mathbb{R}} \frac{(1 - \phi^2)^2}{\phi^2} \\ &= \frac{(1 - \delta)^2}{2} \left(\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})} - \frac{\mathfrak{c}^2}{2\delta} \right) r + \int_r^{r+1} \left[(\phi')^2 + \left(\frac{\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}}{2} - \frac{\mathfrak{c}^2}{4\phi^2} \right) (1 - \phi^2) \right]. \end{aligned}$$

Let us choose $\delta \in (0, 1)$ so that $\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})} - \mathfrak{c}^2/2\delta < 0$. Notice that the last integral depends on δ and \mathfrak{c} , but not on r , since $\phi(x + r) = \psi(x)$ for all $x \in [0, 1]$. Therefore we may take $r > 0$ large enough so that $J_c(1 - \phi) < 0$. In this way, δ and r depend on only \mathfrak{c} and $\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}$. The proof concludes by taking $\phi_{\mathfrak{c}} = \phi$. \square

In the rest of this section we assume that \mathcal{W} satisfies (H1), we fix $\mathfrak{c} \in (0, \sqrt{2\sigma})$ and we focus on speeds c on the interval $(\mathfrak{c}, \sqrt{2\sigma})$. We consider the paths connecting the origin with $1 - \phi_{\mathfrak{c}}$ given by Lemma 3.6, as follows

$$\Gamma(\mathfrak{c}) = \{g \in \mathcal{C}([0, 1], \mathcal{NV}(\mathbb{R})) : g(0) = 0, g(1) = 1 - \phi_{\mathfrak{c}}\}.$$

Thanks to the mountain pass geometry (Lemma 3.5 and Lemma 3.6) and to the continuity of J_c in $\mathcal{NV}(\mathbb{R})$, the so-called mountain pass level is well-defined and is positive:

$$\gamma_{\mathfrak{c}}(c) := \inf_{g \in \Gamma(\mathfrak{c})} \max_{t \in [0, 1]} J_c(g(t)) > 0, \quad \text{for all } c \in [\mathfrak{c}, \sqrt{2\sigma}).$$

Notice that the $\Gamma(\mathfrak{c})$ and $\gamma_{\mathfrak{c}}(c)$ are not standard since the paths take values on the set $\mathcal{NV}(\mathbb{R})$ (and not on a vector space). We recall that the advantage of working only on $\mathcal{NV}(\mathbb{R})$ is that J_c is smooth. However, the drawback of this setting is that one needs to control J_c near $\partial\mathcal{NV}(\mathbb{R})$ in some sense. In fact, in principle there might exist a sequence $\{\rho_n\} \subset 1 - \mathcal{NV}(\mathbb{R})$ such that $\inf_{\mathbb{R}} \rho_n$ tends to zero but $J_c(1 - \rho_n)$ remains finite for all n . The next result provides some properties of the functional \mathcal{B} that prevent undesirable phenomena to happen, so we can deal with the singular behavior near $\partial\mathcal{NV}(\mathbb{R})$.

Lemma 3.7. *Given $\rho \in 1 + H^1(\mathbb{R})$, it holds that*

$$\mathcal{B}(1 - \rho) < +\infty \iff 1 - \rho \in \mathcal{NV}(\mathbb{R}).$$

More precisely, if $\rho \in 1 + H^1(\mathbb{R})$ satisfies

$$\|1 - \rho\|_{H^1(\mathbb{R})} + \mathcal{B}(1 - \rho) \leq R. \tag{3.7}$$

for some $R > 0$, then there exists $\delta \in (0, 1)$ such that $\rho \geq \delta$ on \mathbb{R} .

Proof. We first prove the equivalence. Let $1 - \rho \in \mathcal{NV}(\mathbb{R})$, so that ρ is continuous, $\rho > 0$ on \mathbb{R} and $\rho(\pm\infty) = 1$. Thus, taking $x_0 \in \mathbb{R}$ such that $\rho(x_0) = \min_{\mathbb{R}} \rho > 0$, we have

$$8\mathcal{B}(1 - \rho) = \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2} = \int_{\mathbb{R}} \frac{(1 - \rho)^2(1 + \rho)^2}{\rho^2} \leq \frac{\|1 - \rho\|_{L^2(\mathbb{R})}^2 \|1 + \rho\|_{L^\infty(\mathbb{R})}^2}{\rho(x_0)^2} < +\infty.$$

For the converse implication, we consider $\rho \in 1 + H^1(\mathbb{R})$ such that $\min_{\mathbb{R}} \rho \leq 0$. Since $\rho(\pm\infty) = 1$, we deduce that there is $x_0 \in \mathbb{R}$ such that $\rho(x_0) = 0$ and $\rho(x) > 0$ for all $x > x_0$. Thus, the continuous embedding $H^1(\mathbb{R}) \subset C^{0,1/2}(\mathbb{R})$ implies that

$$\rho(x)^2 = (\rho(x) - \rho(x_0))^2 \leq \|1 - \rho\|_{C^{0,1/2}(\mathbb{R})}^2 (x - x_0), \quad \text{for all } x > x_0,$$

so that $\int_{x_0}^{x_0+\delta} 1/\rho^2 = +\infty$, for every $\delta > 0$. Now we choose $\delta > 0$ such that $\min_{[x_0, x_0+\delta]} (1 - \rho^2)^2 > 0$. Hence,

$$8\mathcal{B}(1 - \rho) \geq \int_{x_0}^{x_0+\delta} \frac{(1 - \rho^2)^2}{\rho^2} \geq \min_{[x_0, x_0+\delta]} (1 - \rho^2)^2 \int_{x_0}^{x_0+\delta} \frac{1}{\rho^2} = +\infty.$$

Thus the equivalence $\mathcal{B}(1 - \rho) < +\infty \iff 1 - \rho \in \mathcal{NV}(\mathbb{R})$ holds true.

We turn now to the proof of the fact that $\rho \geq \delta$ provided that (3.7) holds. First, we have already proved that any $\rho \in 1 + H^1(\mathbb{R})$ satisfying (3.7) belongs to $1 - \mathcal{NV}(\mathbb{R})$. We argue now by contradiction and assume that there exist $R > 0$ and a sequence $\{\rho_n\} \subset 1 - \mathcal{NV}(\mathbb{R})$ such that ρ_n satisfies (3.7) for all n but $\min_{\mathbb{R}} \rho_n \rightarrow 0$ as $n \rightarrow \infty$. Since $\{\rho_n(\cdot + x_n)\} \subset 1 + H^1(\mathbb{R})$ still satisfies (3.7) for any sequence $\{x_n\} \subset \mathbb{R}$, then we may assume without loss of generality that $\rho_n(0) = \min_{\mathbb{R}} \rho_n \rightarrow 0$ as $n \rightarrow \infty$. By the embedding $H^1(\mathbb{R}) \subset C^{0,1/2}(\mathbb{R})$, we deduce that there is a constant $C > 0$ such that $\|1 - \rho_n\|_{C^{0,1/2}(\mathbb{R})} \leq C\|1 - \rho_n\|_{H^1(\mathbb{R})} \leq CR$ for all n , so that

$$\rho_n(x) \leq \rho_n(0) + CR\sqrt{x}, \quad \text{for all } x > 0, \text{ for all } n.$$

We conclude as before that $\lim_{n \rightarrow \infty} \int_0^\delta 1/\rho_n^2 = +\infty$, for every $\delta > 0$. If there exists $\delta > 0$ such that the sequence $\{\min_{[0, \delta]} (1 - \rho_n^2)^2\}$ is bounded away from zero, then we may argue as above to conclude that $\lim_{n \rightarrow \infty} \mathcal{B}(1 - \rho_n) = +\infty$, a contradiction. Otherwise, for every $\delta > 0$, up a subsequence (that depends on δ), $\lim_{n \rightarrow \infty} \min_{[0, \delta]} (1 - \rho_n^2)^2 = 0$. For some fixed $\delta > 0$ to be chosen below, we consider the mentioned subsequence, which we do not relabel, and we take $\{x_n\} \subset [0, \delta]$ such that $\rho_n(x_n) = \max_{[0, \delta]} \rho_n$. Observe that $\rho_n(x_n) \rightarrow 1$ as $n \rightarrow \infty$ and

$$\rho_n(x_n) - \rho_n(0) \leq CR\sqrt{x_n} \leq CR\sqrt{\delta}, \quad \text{for all } n.$$

As the left hand side of the previous inequality tends to one as $n \rightarrow \infty$, choosing $0 < \delta < 1/(CR)^2$ leads one more time to a contradiction. The proof is now concluded. \square

Following the ideas of [33], we observe that $\gamma_c(c)$ is a nonincreasing function of c . Therefore, its derivative $\gamma'_c(c)$ exists for almost every $c \in [c, \sqrt{2\sigma}]$. The points of differentiability of γ_c will be crucial in our arguments. For this reason we introduce the set

$$\mathcal{D}_c = \{c \in (c, \sqrt{2\sigma}) : \exists \gamma'_c(c)\}.$$

As we have pointed out,

$$|\mathcal{D}_c| = |(c, \sqrt{2\sigma})| = \sqrt{2\sigma} - c. \quad (3.8)$$

Now we can state the following result due to Jeanjean [33] adapted to our setting.

Lemma 3.8. *Assume that \mathcal{W} satisfies (H1). Let $c \in \mathcal{D}_c$ and let $\{c_n\}$ be an increasing sequence such that $c_n \rightarrow c$. Then there exist a sequence $\{g_n\} \subset \Gamma(c)$ and a constant $R = R(\gamma'_c(c)) > 0$ such that the following holds:*

- (i) *For any $t \in [0, 1]$ such that $J_c(g_n(t)) \geq \gamma_c(c) - (c - c_n)$, we have the estimate*

$$\|g_n(t)\|_{H^1(\mathbb{R})} + \mathcal{B}(g_n(t)) \leq R.$$

- (ii) $\max_{t \in [0, 1]} J_c(g_n(t)) \leq \gamma_c(c) + (-\gamma'_c(c) + 2)(c - c_n)$.

Proof. The proof is exactly as the one of [33, Proposition 2.1]. We only point out that, in our case, we conclude using the coerciveness of \mathcal{A} . We also stress that the estimate $\mathcal{B}(g_n(t)) \leq R$ follows directly from the proof of [33, Proposition 2.1], as it is also observed in [7, Lemma 4.5]. \square

The next result is also mainly due to Jeanjean [33], but some crucial modifications are needed since J_c is not of class \mathcal{C}^2 in the whole space $H^1(\mathbb{R})$. Thus we adapt the proof thanks to Lemmas 3.3 and 3.7.

Proposition 3.9. *Assume that \mathcal{W} satisfies (H1). Let $c \in \mathcal{D}_c$, let $R_c := R(\gamma'_c(c)) > 0$ be given by Lemma 3.8 and let $\delta_c := \delta(R_c) \in (0, 1)$ be given by Lemma 3.7. For any $\alpha > 0$ and $\delta \in (0, 1)$, let us consider the set*

$$Y_{\alpha, \delta} = J_c^{-1}([\gamma_c(c) - \alpha, \gamma_c(c) + \alpha]) \cap Z_\delta,$$

where Z_δ is defined by (3.5). Then, Y_{α, δ_c} is nonempty for every $\alpha > 0$. Moreover,

$$\inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in Y_{\alpha, \delta_c}\} = 0, \quad \text{for all } \alpha > 0. \quad (3.9)$$

Proof. We first show that $Y_{\alpha, \delta_c} \neq \emptyset$ for any $\alpha > 0$. Indeed, let $\{c_n\} \subset \mathbb{R}$ be an increasing sequence such that $c_n \rightarrow c$, and let $\{g_n\} \subset \Gamma(\mathbf{c})$ be the sequence given by Lemma 3.8. For some n to be chosen later, let $t_n \in [0, 1]$ be such that

$$J_c(g_n(t_n)) = \max_{t \in [0, 1]} J_c(g_n(t)).$$

We check now that $g_n(t_n) \in Y_{\alpha, \delta_c}$. On the one hand, by definition of $\gamma_c(c)$, it follows that

$$J_c(g_n(t_n)) \geq \gamma_c(c) - (c - c_n). \quad (3.10)$$

In consequence, Lemmas 3.8 and 3.7 imply that $g_n(t_n) \in Z_{\delta_c}$. On the other hand, from Lemma 3.8 we deduce that

$$J_c(g_n(t_n)) \leq \gamma_c(c) + (-\gamma'_c(c) + 2)(c - c_n). \quad (3.11)$$

Hence, bearing (3.10) and (3.11) in mind, n can be chosen large enough so that $g_n(t_n) \in Y_{\alpha, \delta_c}$.

We turn now to the proof of (3.9). Arguing by contradiction, let us assume that $\inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in Y_{2\alpha, \delta_c}\} > 0$ for some $\alpha > 0$. Let us denote

$$X_{2\alpha, \delta_c} = J_c^{-1}((\gamma_c(c) - 2\alpha, \gamma_c(c) + 2\alpha)) \cap Z_{\delta_c}.$$

Of course, $\inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in X_{2\alpha, \delta_c}\} > 0$ too. By Lemma 6.3 in the appendix, there exists $\tilde{\delta} \in (0, \delta_c/2)$ such that

$$\inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in X_{2\alpha, \delta_c - 2\tilde{\delta}}\} > 0.$$

Therefore, we immediately have that

$$\inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in Y_{\alpha, \delta_c - 2\tilde{\delta}}\} > 0.$$

Let us choose $\varepsilon > 0$ small enough so that

$$2\varepsilon < \alpha, \quad \|J'_c(v)\|_{H^{-1}(\mathbb{R})} \geq \frac{2\varepsilon}{\tilde{\delta}}, \quad \text{for all } v \in Y_{\alpha, \delta_c - 2\tilde{\delta}}.$$

Thus, we may apply Lemma 3.3 with $\delta_1 = \delta_c - 2\tilde{\delta}$, $\delta_2 = \delta_c - \tilde{\delta}$ and $\delta_3 = \delta_c$, so that there exists a continuous function $h : [0, 1] \times \mathcal{NV}(\mathbb{R}) \rightarrow \mathcal{NV}(\mathbb{R})$ satisfying items (i)-(v). In the rest of the proof we will abuse of the notation and write $h(v) = h(1, v)$ for simplicity.

Recall that $\{c_n\}$ is an increasing sequence such that $c_n \rightarrow c$ and $\{g_n\} \subset \Gamma(\mathbf{c})$ is given by Lemma 3.8. We show now that $h \circ g_n \in \Gamma(\mathbf{c})$ for all n . Notice that, since $\gamma_c(c) > 0$, for every $v \in \mathcal{NV}(\mathbb{R})$ with $J_c(v) \leq 0$ one may choose $\varepsilon > 0$ small enough so that $J_c(v) \notin [\gamma_c(c) - 2\varepsilon, \gamma_c(c) + 2\varepsilon]$. In consequence, recalling that $J_c(0) = 0$ and $J_c(1 - \phi_c) < 0$, we deduce that

$$0, 1 - \phi_c \in \mathcal{NV}(\mathbb{R}) \setminus J_c^{-1}([\gamma_c(c) - 2\varepsilon, \gamma_c(c) + 2\varepsilon]).$$

Therefore, item (ii) of Lemma 3.3 implies that $h(g_n(0)) = h(0) = 0$ and $h(g_n(1)) = h(1 - \phi_c) = 1 - \phi_c$ for all n . In sum, $h \circ g_n \in \Gamma(\mathbf{c})$ for all n .

We now claim that

$$\max_{t \in [0,1]} J_c(h(g_n(t))) \leq \gamma_c(c) - (c - c_n) \quad (3.12)$$

for some n large enough. In order to prove the claim, let us fix $t \in [0, 1]$. On the one hand, if $J_c(g_n(t)) \leq \gamma_c(c) - (c - c_n)$, then item (iv) of Lemma 3.3 implies that $J_c(h(g_n(t))) \leq \gamma_c(c) - (c - c_n)$.

On the other hand, if $J_c(g_n(t)) > \gamma_c(c) - (c - c_n)$, then Lemma 3.8 implies that $\|g_n(t)\|_{H^1(\mathbb{R})} + \mathcal{B}(g_n(t)) \leq R_c$. In turn, Lemma 3.7 yields that $1 - g_n(t) \geq \delta_c$ on \mathbb{R} . In particular, $g_n(t) \in Z_{\delta_c}$. Moreover, by Lemma 3.8, we have $J_c(g_n(t)) \leq \gamma_c(c) + (-\gamma'_c(c) + 2)(c - c_n)$. Thus, by taking n large enough (independent of t) so that $(-\gamma'_c(c) + 2)(c - c_n) \leq \varepsilon$, we derive that $g_n(t) \in J_c^{\gamma_c(c) + \varepsilon}$. Therefore, item (v) of Lemma 3.3 implies that $J_c(h(g_n(t))) \leq \gamma_c(c) - \varepsilon \leq \gamma_c(c) - (c - c_n)$.

In any case, since t was arbitrary, we have shown that (3.12) holds for some n large enough. This contradicts the definition of $\gamma_c(c)$. \square

Next result shows the existence of a bounded Palais–Smale sequence of J_c in $\mathcal{NV}(\mathbb{R})$, for $c \in \mathcal{D}_c$.

Proposition 3.10. *Assume that \mathcal{W} satisfies (H1). For any $c \in \mathcal{D}_c$, there exist $R_c > 0$, $\delta_c \in (0, 1)$ and a sequence $\{v_n\} \subset \mathcal{NV}(\mathbb{R})$ such that*

$$\{v_n\} \subset Z_{\delta_c}, \quad J_c(v_n) \rightarrow \gamma_c(c), \quad \text{and} \quad \|J'_c(v_n)\|_{H^{-1}(\mathbb{R})} \rightarrow 0, \quad (3.13)$$

where Z_{δ_c} is defined by (3.5). Moreover, up a subsequence, $v_n \rightharpoonup v$ weakly in $H^1(\mathbb{R})$, for some $v \in Z_{\delta_c}$ with $v \not\equiv 0$.

Before proving this proposition, let us recall a well-known result (see e.g. Lemma 3.3 in [56]).

Lemma 3.11. *Let $\{v_n\} \subset H^1(\mathbb{R})$ be a bounded sequence and let $q \in (2, \infty)$. Then $v_n \rightarrow 0$ in $L^q(\mathbb{R})$ if and only if $v_n(\cdot + y_n) \rightarrow 0$, for every sequence $\{y_n\} \subset \mathbb{R}$.*

We will also need the following key lemma.

Lemma 3.12. *Let $v \in \mathcal{NE}(\mathbb{R})$ and $c > 0$. Then the following identity holds*

$$2J_c(v) - J'_c(v)(v) = \frac{1}{2} \int_{\mathbb{R}} (\mathcal{W} * (v(2-v)))v^2 + \frac{c^2}{4} \int_{\mathbb{R}} \frac{v(2-v)v^2}{(1-v)^3}. \quad (3.14)$$

In particular, if $v \leq 1 - \delta$ on \mathbb{R} , then

$$|2J_c(v) - J'_c(v)(v)| \leq \max \left\{ \frac{\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}}{2}, \frac{c^2}{4\delta^3} \right\} (2 + \|v\|_{L^\infty(\mathbb{R})}) \|v\|_{L^2(\mathbb{R})} \|v\|_{L^4(\mathbb{R})}^2. \quad (3.15)$$

Proof. Let $\rho = 1 - v$ and $\eta = 1 - \rho^2$, so that $\eta = v(2 - v)$. From (3.1), we have

$$\begin{aligned} 2J_c(v) - J'_c(v)(v) &= \frac{1}{2} \int_{\mathbb{R}} (\mathcal{W} * \eta)(\eta - 2(1-v)v) - \frac{c^2}{4} \int_{\mathbb{R}} \left(\frac{\eta^2}{\rho^2} - \frac{(1-\rho^4)(1-\rho)}{\rho^3} \right) \\ &= \frac{1}{2} \int_{\mathbb{R}} (\mathcal{W} * \eta)v^2 + \frac{c^2}{4} \int_{\mathbb{R}} \frac{\eta v^2}{\rho^3}, \end{aligned}$$

which is (3.14). Finally, the right-hand side can be bounded from above by

$$\max \left\{ \frac{\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}}{2}, \frac{c^2}{4\delta^3} \right\} \|\eta\|_{L^2(\mathbb{R})} \|v\|_{L^4(\mathbb{R})}^2,$$

which, using that $\|\eta\|_{L^2(\mathbb{R})} \leq \|2 - v\|_{L^\infty(\mathbb{R})} \|v\|_{L^2(\mathbb{R})}$, yields (3.15). \square

Proof of Proposition 3.10. Applying Proposition 3.9 with $\alpha = 1/n$, we deduce the existence of R_c , δ_c and a sequence $\{w_n\} \subset \mathcal{NV}(\mathbb{R})$ satisfying (3.13). Since $\{w_n\}$ is bounded in $H^1(\mathbb{R})$ and $\gamma_c(c) \neq 0$, we infer from (3.15) that $w_n \not\rightarrow 0$ strongly in $L^4(\mathbb{R})$. Then we deduce from Lemma 3.11 the existence of a sequence $\{y_n\} \subset \mathbb{R}$ such that $v_n := w_n(\cdot + y_n)$ does not converge weakly to 0 in $H^1(\mathbb{R})$. Of course, $\{v_n\} \subset Z_{\delta_c}$ still satisfies (3.13). Thus, there exists a subsequence (that we do not relabel) such that $v_n \rightharpoonup v$ in $H^1(\mathbb{R})$ for some $v \neq 0$. In addition, since the $\|\cdot\|_{H^1(\mathbb{R})}$ -norm is weakly lower semicontinuous, then $\|v\|_{H^1(\mathbb{R})} \leq R_c + 1 - \delta_c$. Moreover, since $v_n \rightarrow v$ pointwise on \mathbb{R} , it follows that $v \leq 1 - \delta_c$ on \mathbb{R} , so that $v \in Z_{\delta_c}$. \square

Finally, we proceed to prove Theorem 1.1, that establishes the existence of solitons for a.e. $c \in (0, \sqrt{2\sigma})$.

Proof of Theorem 1.1. Let us fix $c \in \mathcal{D}_c$ and let $\{v_n\} \subset H^1(\mathbb{R})$ be the sequence given by Proposition 3.10. We set as usual $\rho_n = 1 - v_n$ and $\eta_n = 1 - \rho_n^2$. Then, by Remark 3.2,

$$J'_c(v_n)(\phi) = \int_{\mathbb{R}} v'_n \phi' + \int_{\mathbb{R}} (\mathcal{W} * \eta_n) \rho_n \phi - \frac{c^2}{4} \int_{\mathbb{R}} \frac{1 - \rho_n^4}{\rho_n^3} \phi \rightarrow 0,$$

for all $\phi \in \mathcal{C}_0^\infty(\mathbb{R})$, and $v_n \rightharpoonup v$ weakly in $H^1(\mathbb{R})$ for some $v \in Z_\delta$ with $v \neq 0$. Observe that, by the Sobolev embedding (3.6), η_n and $1 - \rho_n^4$ are also bounded in $H^1(\mathbb{R})$. Thus, by setting $\rho = 1 - v$ and $\eta = 1 - v^2$, we deduce by the uniqueness of the limit that, up to subsequences,

$$\eta_n \rightharpoonup \eta \quad \text{and} \quad 1 - \rho_n^4 \rightharpoonup 1 - \rho^4 \quad \text{in } H^1(\mathbb{R}), \quad (3.16)$$

$$v_n \rightarrow v \quad \text{and} \quad \rho_n \rightarrow \rho, \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}). \quad (3.17)$$

We will show now that we can pass to the limit the three integral terms. The first one is trivial. For the second term, observe that by (1.23), the convolution is continuous in $L^2(\mathbb{R})$, which combined with (3.16) and the Rellich theorem implies that,

$$\mathcal{W} * \eta_n \rightarrow \mathcal{W} * \eta \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}).$$

Since $\phi \in \mathcal{C}_0^\infty(\mathbb{R})$, using also (3.17), we conclude that $(\mathcal{W} * \eta_n) \rho_n \phi \rightarrow (\mathcal{W} * \eta) \rho \phi$ in $L^1(\mathbb{R})$.

For the last term, using that $(1 - v_n)^3 \geq \delta^3$ on \mathbb{R} for all n , we similarly deduce that

$$\int_{\mathbb{R}} \frac{1 - \rho_n^4}{\rho_n^3} \phi \rightarrow \int_{\mathbb{R}} \frac{1 - \rho^4}{\rho^3} \phi.$$

Gathering all together, we have proved that $J'(v) = 0$, so that $\rho = 1 - v \in 1 + H^1(\mathbb{R})$ is a nontrivial positive solution to (2.13). Moreover, by elliptic regularity, we infer that $\rho \in \mathcal{C}^\infty(\mathbb{R})$. Therefore, by virtue of Proposition 2.4, it follows that $u = \rho e^{i\theta}$, with θ defined by (2.14), belongs to $\mathcal{NE}(\mathbb{R})$ and is a nontrivial solution to $(S(\mathcal{W}, c))$.

In conclusion, we have shown that there exists a solution $u \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c))$ for any $c \in \mathcal{D}_c$. Thus, the same holds for every $c \in \mathcal{D}$, where

$$\mathcal{D} := \bigcup_{c \in (0, \sqrt{2\sigma})} \mathcal{D}_c \subset (0, \sqrt{2\sigma}).$$

By (3.8) we have

$$\sqrt{2\sigma} - c = |\mathcal{D}_c| \leq |\mathcal{D}| \leq \sqrt{2\sigma}, \quad \text{for all } c \in (0, \sqrt{2\sigma}).$$

Taking limits as $c \rightarrow 0$, we conclude that $|\mathcal{D}| = \sqrt{2\sigma}$, which proves the theorem. \square

Proof of Corollary 1.2. Since $\widehat{\mathcal{W}} \geq 0$ a.e. and $\widehat{\mathcal{W}}(\xi) \geq \sigma - \xi^2/2$, for a.e. $|\xi| \leq \sqrt{2\sigma}$, it is immediate to check that for every $\tilde{\sigma} \in (0, \sigma)$, we have $\widehat{\mathcal{W}}(\xi) \geq \tilde{\sigma} - \kappa_\sigma \xi^2/2$, for a.e. $\xi \in \mathbb{R}$, where $\kappa_\sigma = \tilde{\sigma}/(2\sigma)$. By Theorem 1.1, we conclude the existence for a.e. speed in every the interval $]0, \sqrt{2\tilde{\sigma}}[$, for any $\tilde{\sigma} \in (0, \sigma)$, which yields the existence for a.e. $c \in (0, \sqrt{2\sigma})$. \square

4 Existence in the whole subsonic regime

Our next goal is to provide conditions on \mathcal{W} in order to extend Theorem 1.1 and conclude the existence of solution for every subsonic speed. For this reason, we introduce the following assumptions.

- (H2) $\widehat{\mathcal{W}}$ is differentiable a.e. and the map $\xi \mapsto \xi(\widehat{\mathcal{W}})'(\xi)$ is bounded and continuous a.e. on \mathbb{R} .
- (H3) $\widehat{\mathcal{W}}$ is nondecreasing on \mathbb{R}^+ , with $\widehat{\mathcal{W}}(0) = 1$.
- (H4) \mathcal{W} is given by a finite Borel measure. In particular, there is a constant $\|\mathcal{W}\|_p$ such that $\|\mathcal{W} * f\|_{L^p(\mathbb{R})} \leq \|\mathcal{W}\|_p \|f\|_{L^p(\mathbb{R})}$, for all $f \in L^p(\mathbb{R})$, $p \in [1, \infty]$.
- (H5) There exists a continuous function $V_0 : (0, \sqrt{2}) \rightarrow (0, \infty)$ such that for any $u \in \mathcal{NE}(R)$ solution to $(S(\mathcal{W}, c))$, with $c \in (0, \sqrt{2})$, we have $\|u\|_{L^\infty(\mathbb{R})} \leq V_0(c)$.

Let us remark that the condition $\widehat{\mathcal{W}}(0) = 1$ is only a normalization of the potential, so that the speed of sound is from now on fixed and equal to $c_*(\mathcal{W}) = \sqrt{2}$. Indeed, if $\widehat{\mathcal{W}}$ is increasing on \mathbb{R}^+ , then the limit $\lim_{\xi \rightarrow 0^+} \widehat{\mathcal{W}}(\xi)$ exists. Since we always assume that $\widehat{\mathcal{W}}$ is even, this yields that we can suppose that $\widehat{\mathcal{W}}$ is continuous at the origin. Therefore, if $\widehat{\mathcal{W}}(0) \neq 0$, making a change of variable we can replace $\widehat{\mathcal{W}}(\xi)$ with $\widehat{\mathcal{W}}(\xi)/\widehat{\mathcal{W}}(0)$ as in [22], which gives the normalization. Also, observe that (H3) implies that

$$\min_{\xi \in \mathbb{R}} \widehat{\mathcal{W}}(\xi) = 1, \quad (4.1)$$

so that (H1) is satisfied with $\sigma = 1$ and $\kappa = 0$. In particular, if \mathcal{W} satisfies (H0) and (H3), Theorem 1.1 gives the existence of solitons for a.e. $c \in (0, \sqrt{2})$.

Concerning (H4), invoking the results in [28, Section 2] and [25], we see that if $f \mapsto \mathcal{W} * f$ is a bounded linear operator from $L^1(\mathbb{R})$ to itself, then \mathcal{W} is given by a (signed) finite Borel measure μ , i.e.

$$(\mathcal{W} * f)(x) = \int_{\mathbb{R}} f(x - y) d\mu(y), \quad \text{for all } f \in L^p(\mathbb{R}), \quad p \in [1, \infty]. \quad (4.2)$$

Thus $\|\mathcal{W}\|_p \leq \int_{\mathbb{R}} d|\mu(y)|$ and (1.24) holds for $f \in L^p(\mathbb{R})$, $g \in L^{p'}(\mathbb{R})$. In addition, $\widehat{\mu}$ is continuous, with $\widehat{\mu}(\xi) = \int_{\mathbb{R}} e^{-ix\xi} d\mu(x)$. Thus, if (H3) also holds, then

$$\widehat{\mathcal{W}}(0) = \widehat{\mu}(0) = \mathcal{W} * 1 = 1. \quad (4.3)$$

Now we can state our main theorem concerning the existence of solitons for every subsonic speed.

Theorem 4.1. *Let $c \in (0, \sqrt{2})$ and assume that \mathcal{W} satisfies (H2), (H3), (H4) and (H5). Then there exists a nontrivial solution $u \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c))$.*

Notice that assumption (H5) can be seen as an alternative way of imposing that the equation $(S(\mathcal{W}, c))$ satisfies some type of maximum principle. Clearly, given a potential \mathcal{W} , this is the only hypothesis difficult to verify. Remark that if one we can show the existence of a constant

$C > 0$ such that any solution $u \in \mathcal{NE}(R)$ to $(S(\mathcal{W}, c))$, with $c \in (0, \sqrt{2})$, satisfies $\|u\|_{L^\infty(\mathbb{R})} \leq C$, then (H5) holds true.

In Proposition 4.5, we will prove that (H5) holds for a potential of the form (1.11). Moreover, in this case $(\widehat{\mathcal{W}}_f)' = A_f(\widehat{f})'$, so that Theorem 1.3 follows immediately from Theorem 4.1.

As explained in the introduction, for the proof of Theorem 4.1, we can take $c \in (0, \sqrt{2})$ and apply Theorem 1.1 to get the existence of a sequence of speeds $\{c_n\}$ and a sequence of associated solutions $\{v_n\} \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c_n))$ such that $c_n \rightarrow c$. To conclude that $\{v_n\}$ converges to a *finite energy* solution v to $(S(\mathcal{W}, c))$, we need to obtain uniform estimates for $\{v_n\}$, and to get a more precise information of v , using the fact that each v_n is the limit of a Palais–Smale sequence for J_c . We deal with these problems in the following subsections.

4.1 Uniform estimates

We start by recalling a Pohozaev identity that was proved by the first author in [18] in a more general framework.

Proposition 4.2 (Propositions 5.1 and 5.3 in [18]). *Let $c \geq 0$ and assume that \mathcal{W} satisfies (H0) and (H2). Let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $S(\mathcal{W}, c)$. Then*

$$\int_{\mathbb{R}} |u'|^2 = \frac{1}{4\pi} \int_{\mathbb{R}} \left(\widehat{\mathcal{W}}(\xi) - \xi(\widehat{\mathcal{W}})'(\xi) \right) |\widehat{\eta}(\xi)|^2 d\xi, \quad (4.4)$$

where $\eta = 1 - |u|^2$.

Notice that Proposition 4.2 can be applied to the focusing case $\mathcal{W} = -\delta_0$, and it is immediate to deduce that then the only finite energy solution to $S(\mathcal{W}, c)$ are the constants, since $\widehat{\mathcal{W}} = -1$.

More general examples can be constructed by defining an even function such that

$$\widehat{\mathcal{W}}(\xi; g) = -\xi \int_{\xi}^{\infty} \frac{g(y)}{y^2} dy, \quad \text{for } \xi > 0,$$

with g a bounded nonnegative function, since $\widehat{\mathcal{W}}(\xi; g) - \xi(\widehat{\mathcal{W}})'(\xi; g) \leq 0$ a.e. on \mathbb{R} . For instance, if $g = 1$, then we obtain $\widehat{\mathcal{W}}(\xi; g) = -1$. Consequently, we can construct potentials with arbitrary L^1 -norm, such that the only finite energy solution to $(S(\mathcal{W}, c))$ are the trivial ones.

Corollary 4.3. *For any $\alpha > 0$, there exists a function $\mathcal{W}_\alpha \in C^\infty(\mathbb{R})$ satisfying (H0) and (H2), with $\|\mathcal{W}_\alpha\|_{L^1(\mathbb{R})} = \alpha$ and $\widehat{\mathcal{W}}_\alpha \leq 0$ on \mathbb{R} , such that if $u \in \mathcal{E}(\mathbb{R})$ is a solution to $(S(\mathcal{W}_\alpha, c))$ for some $c \geq 0$, then u is constant.*

Proof. Let us take $g(y) = y^3 \exp(-y^2)$, which gives $\widehat{\mathcal{W}}(\xi) = -|\xi|e^{-\xi^2}$, for $\xi \in \mathbb{R}$. In this manner, \mathcal{W} is a smooth function with exponential decay, so that it suffices to consider $\mathcal{W}_\alpha = \alpha\mathcal{W}/\|\mathcal{W}\|_{L^1(\mathbb{R})}$. \square

Identity (4.4), together with (2.4) and (2.6), allow us to prove the following nonvanishing property of nontrivial finite energy solutions to $(S(\mathcal{W}, c))$.

Proposition 4.4. *Let $c \geq 0$. Assume that \mathcal{W} satisfies (H2) and that*

$$3\xi^2 + 2\widehat{\mathcal{W}}(\xi) + 2\xi(\widehat{\mathcal{W}})'(\xi) \geq 2 \quad \text{a.e. on } \mathbb{R}. \quad (4.5)$$

Then every nontrivial solution $u \in \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, c))$ satisfies the estimate

$$\|\mathcal{W} * \eta\|_{L^\infty(\mathbb{R})} \geq \frac{2 - c^2}{4}. \quad (4.6)$$

Proof. Multiplying (2.4) by $2\eta'$ and integrating by parts leads to

$$2 \int_{\mathbb{R}} (\eta')^2 + 4 \int_{\mathbb{R}} \eta(\mathcal{W} * \eta) - 2c^2 \int_{\mathbb{R}} \eta^2 = 4 \int_{\mathbb{R}} |u'|^2 \eta + 4 \int_{\mathbb{R}} \eta^2(\mathcal{W} * \eta).$$

Using now (2.6), we deduce that

$$3 \int_{\mathbb{R}} (\eta')^2 + 4 \int_{\mathbb{R}} \eta(\mathcal{W} * \eta) - c^2 \int_{\mathbb{R}} \eta^2 = 4 \int_{\mathbb{R}} |u'|^2 + 4 \int_{\mathbb{R}} \eta^2(\mathcal{W} * \eta).$$

Combining with (4.4) and applying Plancherel's identity, we derive

$$3 \int_{\mathbb{R}} (\eta')^2 + 2 \int_{\mathbb{R}} \eta(\mathcal{W} * \eta) - c^2 \int_{\mathbb{R}} \eta^2 + \frac{1}{\pi} \int_{\mathbb{R}} \xi(\widehat{\mathcal{W}})'(\xi) |\widehat{\eta}(\xi)|^2 d\xi = 4 \int_{\mathbb{R}} \eta^2(\mathcal{W} * \eta).$$

Again by Plancherel's identity, this equality can be recast as

$$\frac{1}{2\pi} \int_{\mathbb{R}} (3\xi^2 + 2\widehat{\mathcal{W}}(\xi) + 2\xi(\widehat{\mathcal{W}})'(\xi)) |\widehat{\eta}(\xi)|^2 d\xi - c^2 \int_{\mathbb{R}} \eta^2 = 4 \int_{\mathbb{R}} \eta^2(\mathcal{W} * \eta).$$

Therefore, inequality (4.5) implies that

$$(2 - c^2) \int_{\mathbb{R}} \eta^2 \leq 4 \int_{\mathbb{R}} \eta^2(\mathcal{W} * \eta) \leq 4 \|\mathcal{W} * \eta\|_{L^\infty(\mathbb{R})} \int_{\mathbb{R}} \eta^2.$$

The result follows by taking into account that η is nontrivial, i.e. $\|\eta\|_{L^2(\mathbb{R})} > 0$. \square

The next proposition shows that the potential in (1.11) satisfies (H5).

Proposition 4.5. *Let $c > 0$ and assume that $\mathcal{W}_f = A_f(\delta_0 + f)$ is as in (1.11). Then, for every solution $u \in \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, c))$, the following estimates hold:*

$$\|u\|_{L^\infty(\mathbb{R})}^2 \leq B_0(f) \left(1 + \frac{c^2}{4}\right), \quad (4.7)$$

$$\|u'\|_{L^\infty(\mathbb{R})} \leq B_1(f) \left(1 + \frac{c^2}{4}\right)^2. \quad (4.8)$$

where $B_0(f) = 1 + \frac{\|f^+\|_{L^1(\mathbb{R})}}{1 - \|f^-\|_{L^1(\mathbb{R})}}$ and $B_1(f)$ is a constant depending only on $\|f^+\|_{L^1(\mathbb{R})}$ and $\|f^-\|_{L^1(\mathbb{R})}$. Moreover, for any $k \geq 2$ there is a constant $C_k(c) > 0$, depending only on c and k , and $B_k(f) > 0$, depending only on $\|f^+\|_{L^1(\mathbb{R})}$, $\|f^-\|_{L^1(\mathbb{R})}$ and k , such that

$$\|D^k u\|_{L^\infty(\mathbb{R})} \leq B_k(f) C_k(c). \quad (4.9)$$

Proof. Since $c > 0$, by Propositions 2.2 and 2.4, the function $\rho = |u|$ satisfies the equation (2.13). By using Young's inequality and the fact that $f * 1 = \int_{\mathbb{R}} f(x) dx = \widehat{f}(0)$, we estimate the term on the right-hand side of the equation as follows, where we drop the subscript f for simplicity,

$$\begin{aligned} \mathcal{W} * (1 - \rho^2) &= A(1 - \rho^2) + Af * (1 - \rho^2) = A(1 - \rho^2) + A\widehat{f}(0) - Af * (\rho^2) \\ &= 1 - A\rho^2 - Af * (\rho^2), \end{aligned}$$

where we used that $A(1 + \widehat{f}(0)) = 1$. Therefore

$$\mathcal{W} * (1 - \rho^2) \leq 1 - A\rho^2 - A(f^-) * (\rho^2) \leq 1 - A\rho^2 + A\|f^-\|_{L^1(\mathbb{R})} \|\rho\|_{L^\infty(\mathbb{R})}^2, \quad (4.10)$$

and

$$\|\mathcal{W} * (1 - \rho^2)\|_{L^\infty(\mathbb{R})} \leq 1 + A\|\rho\|_{L^\infty(\mathbb{R})}^2 + A\|f^-\|_{L^1(\mathbb{R})} \|\rho\|_{L^\infty(\mathbb{R})}^2. \quad (4.11)$$

Plugging (4.10) into (2.13) leads to

$$-\rho'' + \rho \left(A\rho^2 - 1 - \frac{c^2}{4} - A\|f^-\|_{L^1(\mathbb{R})}\|\rho\|_{L^\infty(\mathbb{R})}^2 \right) \leq 0 \quad \text{on } \mathbb{R}.$$

By applying the maximum principle or Proposition 2.1 in [24], we conclude that

$$\rho(x)^2 \leq \frac{1}{A} \left(1 + \frac{c^2}{4} \right) + \|f^-\|_{L^1(\mathbb{R})}\|\rho\|_{L^\infty(\mathbb{R})}^2, \quad \text{for all } x \in \mathbb{R}. \quad (4.12)$$

Let us assume that there exists some $\bar{x} \in \mathbb{R}$ such that $\rho(\bar{x}) > 1$; otherwise, the result is trivial. Since $\rho(\pm\infty) = 1$, there exists $\tilde{x} \in \mathbb{R}$ such that $\rho(\tilde{x}) = \|\rho\|_{L^\infty(\mathbb{R})}$. Thus, using (4.12) in \tilde{x} , we get

$$(1 - \|f^-\|_{L^1(\mathbb{R})})\|\rho\|_{L^\infty(\mathbb{R})}^2 \leq \frac{1}{A} \left(1 + \frac{c^2}{4} \right),$$

which proves (4.7).

In order to establish (4.8), we follow [10, 24] and define $v(x) = u(x)e^{\frac{ic}{2}x}$, for $x \in \mathbb{R}$. It is immediate to verify that $v \in \mathcal{E}(\mathbb{R})$ and that it solves the equation

$$-v'' = \left(\frac{c^2}{4} + \mathcal{W} * (1 - |v|^2) \right) v \quad \text{on } \mathbb{R}. \quad (4.13)$$

From (4.7) and (4.11), it follows that

$$\begin{aligned} \|v''\|_{L^\infty(\mathbb{R})} &\leq \|v\|_{L^\infty(\mathbb{R})} \left(\frac{c^2}{4} + 1 + A\|\rho\|_{L^\infty(\mathbb{R})}^2 + A\|f^-\|_{L^1(\mathbb{R})}\|\rho\|_{L^\infty(\mathbb{R})}^2 \right) \\ &\leq \left(1 + \frac{c^2}{4} \right)^{\frac{3}{2}} B_0^{1/2} (1 + AB_0 + AB_0\|f^-\|_{L^1(\mathbb{R})}). \end{aligned}$$

Recalling that $A(1 + \|f^+\|_{L^1(\mathbb{R})} - \|f^-\|_{L^1(\mathbb{R})}) = 1$, it is clear that

$$\|v''\|_{L^\infty(\mathbb{R})} \leq 2 \left(1 + \frac{c^2}{4} \right)^{\frac{3}{2}} B_0^{1/2}.$$

Thus, using the Landau–Kolmogorov interpolation inequality (see e.g. p.133 in [15])

$$\|v'\|_{L^\infty(\mathbb{R})} \leq \sqrt{2} \|v\|_{L^\infty(\mathbb{R})} \|v''\|_{L^\infty(\mathbb{R})},$$

we infer that

$$\|v'\|_{L^\infty(\mathbb{R})} \leq 2\sqrt{2} B_0 \left(1 + \frac{c^2}{4} \right)^2.$$

Therefore, by definition of v and using that $c/2 \leq 1 + c^2/4$, we deduce that

$$\|u'\|_{L^\infty(\mathbb{R})} \leq \frac{c}{2} \|u\|_{L^\infty(\mathbb{R})} + \|v'\|_{L^\infty(\mathbb{R})} \leq \left(1 + \frac{c^2}{4} \right)^2 B_0^{1/2} (1 + 2\sqrt{2} B_0^{1/2}).$$

Hence, taking $B_1(f) := B_0^{1/2} (1 + 2\sqrt{2} B_0^{1/2})$, we have (4.8). Differentiating (4.13) and using the higher order Landau–Kolmogorov inequalities, we finally conclude the proof of (4.9). \square

Next two propositions show that, for general potentials satisfying the continuity property (H4), an L^∞ estimate for the solutions (i.e. condition (H5)) implies a priori estimates also for the derivatives as well as a uniform lower bound.

Proposition 4.6. *Assume that \mathcal{W} satisfies (H4) and (H5). Then, for every $k \in \mathbb{N}$, there exist continuous functions $V_k : (0, \sqrt{2}) \rightarrow (0, \infty)$ such that for any $u \in \mathcal{NE}(\mathbb{R})$ solution to $(S(\mathcal{W}, c))$, with $c \in (0, \sqrt{2})$, we have $\|D^k u\|_{L^\infty(\mathbb{R})} \leq V_k(c)$. In particular, if $\mathcal{W} = \mathcal{W}_f$ is given by (1.11), then $V_1(c) = B_1(f)(1 + c^2/4)^2$, where $B_1(f)$ is the constant in Proposition 4.5.*

Proof. By using (4.13) and (H4), the proof follows the same line as Proposition 4.5. \square

Proposition 4.7. *Assume that \mathcal{W} satisfies (H4) and (H5). Let $c \in (0, \sqrt{2})$ and let $u \in \mathcal{NE}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Then*

$$|u(x)| \geq \frac{\sqrt{1 + 4c^2/V_1(c)} - 1}{\sqrt{1 + 4c^2/V_1(c)} + 1}, \quad \text{for all } x \in \mathbb{R}, \quad (4.14)$$

where V_1 is the function given by Proposition 4.6.

Proof. Since $u \in \mathcal{NE}(\mathbb{R})$, we have that $\min_{\mathbb{R}} |u| > 0$. Let $x_0 \in \mathbb{R}$ be such that $u(x_0) = \min_{\mathbb{R}} |u|$. From the identity (2.6), we deduce that the function $\eta = 1 - |u|^2$ satisfies

$$c^2 \eta(x_0)^2 \leq \|u'\|_{L^\infty(\mathbb{R})} (1 - \eta(x_0)). \quad (4.15)$$

By using the estimate in Proposition 4.6, we get

$$c^2 \eta(x_0)^2 + V_1(c) \eta(x_0) - V_1(c) \leq 0,$$

which implies that

$$\eta(x_0) \leq \frac{-V_1(c) + \sqrt{V_1(c)^2 + 4V_1(c)c^2}}{2c^2}.$$

In terms of $|u(x_0)|$ we get

$$|u(x_0)|^2 \geq 1 + \frac{V_1(c) - \sqrt{V_1(c)^2 + 4V_1(c)c^2}}{2c^2} = \frac{\sqrt{V_1(c)^2 + 4V_1(c)c^2} - V_1(c)}{\sqrt{V_1(c)^2 + 4V_1(c)c^2} + V_1(c)},$$

which completes the proof. \square

The following nonvanishing property of the functional \mathcal{A} will be useful.

Lemma 4.8. *Assume that \mathcal{W} satisfies (H2), (H4), (H5) and (4.5). Then there exists $C > 0$ such that for any nonzero solution $v \in H^1(\mathbb{R})$ to (3.3), we have*

$$\mathcal{A}(v) \geq \frac{C(2 - c^2)^2}{16}. \quad (4.16)$$

Proof. Let $\eta = 1 - |v|^2$. Then

$$\begin{aligned} \mathcal{A}(v) &= \frac{1}{2} \int_{\mathbb{R}} (v')^2 + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * \eta) \eta = \frac{1}{8} \int_{\mathbb{R}} \frac{(\eta')^2}{1 - \eta} + \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * \eta) \eta \\ &\geq \frac{1}{8 \|1 - \eta\|_{L^\infty(\mathbb{R})} \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}^2} \int_{\mathbb{R}} (\mathcal{W} * \eta')^2 + \frac{1}{4 \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}} \int_{\mathbb{R}} (\mathcal{W} * \eta)^2. \end{aligned}$$

By using the Sobolev's embedding, (4.6) and (4.14), we conclude that there exists $C > 0$ such that

$$\mathcal{A}(v) \geq C \|\mathcal{W} * \eta\|_{H^1(\mathbb{R})}^2 \geq C \|\mathcal{W} * \eta\|_{L^\infty(\mathbb{R})}^2 \geq \frac{C(2 - c^2)^2}{16}.$$

\square

4.2 Refined study of Palais–Smale sequences

We start by recalling a classical result of profile decomposition of a bounded sequence, that is a refinement of the Banach–Alaoglu theorem. We use here the version given in Theorem 4.6.5 in [55] (see also [4, 31]).

Theorem 4.9. *Let $\{v_n\} \subset H^1(\mathbb{R})$ be a bounded sequence. Then there exist a family of concentration profiles $\{w_j\} \subset H^1(\mathbb{R})$ and points $\{y_{n,j}\} \subset \mathbb{R}$ such that, on a renumbered subsequence,*

$$y_{n,1} = 0, \quad \lim_{n \rightarrow \infty} |y_{n,i} - y_{n,j}| \rightarrow \infty, \quad \text{if } i \neq j,$$

$v_n(\cdot + y_{n,j}) \rightharpoonup w_j$ in $H^1(\mathbb{R})$ and $v_n(\cdot + y_{n,j}) \rightarrow w_j$ in $L^\infty_{\text{loc}}(\mathbb{R})$,

$$v_n - S_n \rightarrow 0 \text{ in } L^q(\mathbb{R}), \quad \text{where } S_n = \sum_{j=1}^{\infty} w_j(\cdot - y_{n,j}), \quad (4.17)$$

for all $q \in (2, \infty)$. Moreover, the series S_n converges in $H^1(\mathbb{R})$ unconditionally and uniformly in n , and for all $\varphi \in L^2(\mathbb{R})$, $\{\alpha_n\} \subset \mathbb{R}$ and $q \in (2, \infty)$, we have

$$v_n = \sum_{j=1}^k w_{n,j} + r_{n,k}, \quad \text{with } \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}} r_{n,k}(\cdot - \alpha_n) \varphi \right| = \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \|r_{n,k}\|_{L^q(\mathbb{R})} = 0, \quad (4.18)$$

where $w_{n,j} = w_j(\cdot - y_{n,j})$. In addition,

$$\|D^m v_n\|_{L^2(\mathbb{R})}^2 = \sum_{j=1}^k \|D^m w_j\|_{L^2(\mathbb{R})}^2 + \|D^m r_{n,k}\|_{L^2(\mathbb{R})}^2 + o_n(1), \quad \text{for } m \in \{0, 1\}. \quad (4.19)$$

In Theorem 4.9 and for the rest of the article, the notation $o_n(1)$ stands for a sequence in \mathbb{R} such that $o_n(1) \rightarrow 0$, as $n \rightarrow \infty$. Besides, from now on $o_n(1; L^1)$ will denote a function such that $\|o_n(1; L^1)\|_{L^1(\mathbb{R})} \rightarrow 0$, as $n \rightarrow \infty$.

Notice also that we added to the statement in [55] that $v_n(\cdot + y_{n,j})$ converges to w_j in $L^\infty_{\text{loc}}(\mathbb{R})$, by invoking the Rellich theorem.

We recall now a version of the Brezis–Lieb lemma given in [57].

Lemma 4.10. *Assume that $G \in C^1(\mathbb{R}; \mathbb{R})$, with $G(0) = 0$, and that there exist $a > 0$ and $q > 1$ such that*

$$|G'(t)| \leq a(|t| + |t|^q), \quad \text{for all } t \in \mathbb{R}. \quad (4.20)$$

If the sequence $\{v_n\}$ is bounded in $H^1(\mathbb{R})$ and converges a.e. to v , then

$$G(v_n) = G(v_n - v) + G(v) + o_n(1; L^1). \quad (4.21)$$

Moreover, using the notations in Theorem 4.9, if the profile decomposition is finite, i.e. there exists $k \geq 1$ such that

$$v_n = \sum_{j=1}^k w_{n,j} + r_n, \quad \text{with } r_n \rightarrow 0 \text{ in } H^1(\mathbb{R}) \text{ and } r_n \rightarrow 0 \text{ in } L^q(\mathbb{R}) \text{ for all } q \in (2, \infty), \quad (4.22)$$

then

$$G(v_n) = \sum_{j=1}^k G(w_{n,j}) + G(r_n) + o_n(1; L^1). \quad (4.23)$$

Proof. The decomposition in (4.21) corresponds to Lemma 2.3 in [57]. To show (4.23), let us denote by $\tau_{n,j}$ the translation by $y_{n,j}$, i.e. $\tau_{n,j}v = v(\cdot + y_{n,j})$. Using that $y_{n,1} = 0$, we have by Theorem 4.9 that $v_n \rightharpoonup w_1$ in $H^1(\mathbb{R})$. Then, by (4.21) we obtain

$$G(v_n) = G(w_1) + G(v_n - w_1) + o_n(1; L^1). \quad (4.24)$$

Now, again by Theorem 4.9 and using that $|y_{n,2}| \rightarrow \infty$ as $n \rightarrow \infty$, we derive that $\tau_{n,2}v_n - \tau_{n,2}w_1 \rightarrow w_2$ a.e. on \mathbb{R} . Thus (4.21) implies that

$$G(\tau_{n,2}v_n - \tau_{n,2}w_1) - G(w_2) - G(\tau_{n,2}v_n - \tau_{n,2}w_1 - w_2) = o_n(1; L^1).$$

Therefore, by a change of variables,

$$G(v_n - w_1) - G(w_{n,2}) - G(v_n - w_1 - w_{n,2}) = o_n(1; L^1).$$

Combining with (4.24), we conclude that

$$G(v_n) = G(w_1) + G(w_{n,2}) + G(v_n - w_1 - w_{n,2}) + o_n(1; L^1).$$

By repeating the same argument k times, we get (4.23). \square

In the following lemma we deal with the splitting of the singular term \mathcal{B} .

Lemma 4.11. *Let $\{v_n\} \subset H^1(\mathbb{R})$ be a bounded sequence such that $v_n \rightharpoonup v$ in $H^1(\mathbb{R})$ for some $v \in H^1(\mathbb{R})$. Assume that there exists $\delta \in (0, 1)$ such that $v_n \leq 1 - \delta$ on \mathbb{R} . Then, there is $N \in \mathbb{N}$ such that,*

$$v \leq 1 - \delta, \quad v_n - v \leq 1 - \delta/2 \quad \text{on } \mathbb{R}, \quad \text{for all } n \geq N, \quad (4.25)$$

and

$$\mathcal{B}(v_n) = \mathcal{B}(v_n - v) + \mathcal{B}(v) + o_n(1). \quad (4.26)$$

Moreover, if the profile decomposition is finite as in (4.22), then

$$w_j \leq 1 - \delta, \quad \text{on } \mathbb{R}, \quad \text{for all } j = 1, \dots, k, \quad (4.27)$$

$$r_n \leq 1 - \frac{\delta}{2}, \quad \text{on } \mathbb{R}, \quad \text{for all } n \geq N, \quad (4.28)$$

and

$$\mathcal{B}(v_n) = \sum_{j=1}^k \mathcal{B}(w_j) + \mathcal{B}(r_n) + o_n(1). \quad (4.29)$$

Proof. We first prove (4.25). Since $v_n \rightarrow v$ a.e. on \mathbb{R} and $v_n \leq 1 - \delta$, it follows that $v \leq 1 - \delta$. Now, since $v \in H^1(\mathbb{R})$, we can fix $R > 0$ such that $|v| \leq \delta/2$ a.e. on $\mathbb{R} \setminus B_R(0)$. Then, for all n ,

$$v_n - v \leq 1 - \delta + \delta/2 = 1 - \delta/2 \quad \text{on } \mathbb{R} \setminus B_R(0).$$

Moreover, since $v_n \rightarrow v$ in $L^\infty(B_R(0))$, then, for any n large enough,

$$v_n - v \leq \|v_n - v\|_{L^\infty(B_R(0))} \leq 1 - \delta/2 \quad \text{on } B_R(0).$$

In any case, (4.25) holds.

We turn now to proving (4.26). Using the notation in Lemma 3.1, we see that

$$\mathcal{B}(v) = \int_{\mathbb{R}} H(v(x)) dx, \quad \text{where } H(t) = \int_0^t h(s) ds, \quad h(s) = \frac{s(2-s)(s^2 - 2s + 2)}{4(1-s)^3}.$$

We remark that we can easily construct a bounded function $\chi_\delta \in \mathcal{C}^1(\mathbb{R})$ such that

$$\chi_\delta(s) = \frac{1}{4(1-s)^3} \quad \text{for all } s \leq 1 - \frac{\delta}{2}, \quad \|\chi_\delta\|_{L^\infty(\mathbb{R})} \leq B_\delta, \quad (4.30)$$

for some constant $B_\delta > 0$ depending only on δ . Then the function $\tilde{h}(s) = s(2-s)(s^2-2s+2)\chi_\delta(s)$ clearly satisfies

$$|\tilde{h}(s)| \leq C_\delta(|s| + s^4), \quad \text{for all } s \in \mathbb{R},$$

for some $C_\delta > 0$ depending only on δ . Therefore, condition (4.20) holds for $G = \tilde{H}$, being $\tilde{H}(t) = \int_0^t \tilde{h}(s)ds$. Thus we obtain

$$\tilde{H}(v_n) = \tilde{H}(v_n - v) + \tilde{H}(v) + o_n(1; L^1).$$

Using now (4.25), we conclude that

$$H(v_n) = H(v_n - v) + H(v) + o_n(1; L^1), \quad (4.31)$$

which gives (4.26).

Next, we prove (4.27) and (4.28). In order to do so, let us fix $\varepsilon > 0$ to be chosen later. The density of $\mathcal{C}_0^\infty(\mathbb{R})$ in $H^1(\mathbb{R})$ implies that, for every $j = 1, \dots, k$, there exist $g_j \in \mathcal{C}_0^\infty(\mathbb{R})$ and $\varphi_j \in H^1(\mathbb{R})$ such that

$$w_j = g_j + \varphi_j, \quad \|\varphi_j\|_{L^\infty(\mathbb{R})} < \varepsilon/k. \quad (4.32)$$

Hence, we can take $R > 0$ such that $\cup_{j=1}^k \text{supp}(g_j) \subset B_R(0)$. Let us denote $g_{n,j} = g_j(\cdot - y_{n,j})$. It is clear that

$$\text{supp}(g_{n,j}) \subset B_R(y_{n,j}), \quad \text{for all } j = 1, \dots, k.$$

In particular, since $|y_{n,i} - y_{n,j}| \rightarrow \infty$ for all $i \neq j$, there is $N \in \mathbb{N}$ such that, for all $n \geq N$,

$$\text{supp}(g_{n,i}) \cap \text{supp}(g_{n,j}) = \emptyset \quad \text{for all } i \neq j. \quad (4.33)$$

On the other hand, by Theorem 4.9, $v_n(\cdot + y_{n,j}) \rightarrow w_j$ a.e. on \mathbb{R} , so (4.27) follows directly from the fact that $v_n \leq 1 - \delta$. Moreover, $v_n(\cdot + y_{n,j}) \rightarrow w_j$ in $L^\infty(B_R(0))$, for all $j = 1, \dots, k$. Thus, we may take N larger if necessary in order to get, for all $n \geq N$ and for all $j = 1, \dots, k$,

$$\|v_n(\cdot + y_{n,j}) - w_j\|_{L^\infty(B_R(0))} < \varepsilon/k. \quad (4.34)$$

To show (4.28), fix $x \in \mathbb{R}$ and $n \geq N$. Observe that

$$r_n(x) = v_n(x) - \sum_{j=1}^k g_{n,j}(x) - \sum_{j=1}^k \varphi_j(x - y_{n,j}).$$

Now we have two possibilities. On the one hand, if $x \notin \text{supp}(g_{n,j})$ for any $j = 1, \dots, k$, then, using (4.32), we obtain

$$r_n(x) = v_n(x) - \sum_{j=1}^k \varphi_j(x - y_{n,j}) < 1 - \delta + \varepsilon.$$

On the other hand, if $x \in \text{supp}(g_{n,i})$ for some $i = 1, \dots, k$, then i is unique by virtue of (4.33). We may assume without loss of generality that $i = 1$. Moreover, $x \in B_R(y_{n,1})$, so $z_n := x - y_{n,1} \in B_R(0)$. Therefore, using (4.32) and (4.34), we deduce that

$$\begin{aligned} r_n(x) &= v_n(x) - g_{n,1}(x) - \sum_{j=1}^k \varphi_j(x - y_{n,j}) = v_n(z_n + y_{n,1}) - w_1(z_n) - \sum_{j=2}^k \varphi_j(x - y_{n,j}) \\ &< \|v_n(\cdot + y_{n,1}) - w_1\|_{L^\infty(B_R(0))} + \frac{(k-1)\varepsilon}{k} < \varepsilon. \end{aligned}$$

In any case, we can choose $\varepsilon = \min\{\delta/2, 1 - \delta/2\} = \delta/2$ so that (4.28) holds.

Once (4.27) and (4.28) are proved, (4.29) follows from Lemma 4.10 by applying the same procedure by truncation described above. \square

In order to deal with the splitting of the nonlocal term, we introduce the notation

$$\langle u, v \rangle = \int_{\mathbb{R}} (\mathcal{W} * u)v, \quad \|u\|^2 = \langle u, u \rangle, \quad \mathcal{Q}(u) = \langle u(2-u), u(2-u) \rangle, \quad \text{for all } u, v \in H^1(\mathbb{R}).$$

Notice that $\langle \cdot, \cdot \rangle$ is symmetric and bilinear, so that $\|\cdot\|$ defines a norm provided that $\widehat{\mathcal{W}} \geq 0$.

Lemma 4.12. *Let $\{v_n\} \subset H^1(\mathbb{R})$ be a bounded sequence. Using the notation in Theorem 4.9, we have, up to a subsequence,*

$$\mathcal{Q}(v_n) = \sum_{j=1}^k \mathcal{Q}(w_j) + \mathcal{Q}(r_{n,k}) + \varepsilon_{n,k}, \quad (4.35)$$

where $\{\varepsilon_{n,k}\} \subset \mathbb{R}$ satisfies

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} |\varepsilon_{n,k}| = 0. \quad (4.36)$$

Remark 4.13. It is clear from (4.19) and (4.35) that

$$\mathcal{A}(v_n) = \sum_{j=1}^k \mathcal{A}(w_j) + \mathcal{A}(r_{n,k}) + \varepsilon_{n,k},$$

for some $\{\varepsilon_{n,k}\} \subset \mathbb{R}$ satisfying (4.36). Moreover, if $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} , then $\mathcal{A}(r_{n,k}) \geq 0$, so that

$$\sum_{j=1}^k \mathcal{A}(w_j) \leq \limsup_{n \rightarrow \infty} (\mathcal{A}(v_n) + |\varepsilon_{n,k}|) \leq \limsup_{n \rightarrow \infty} \mathcal{A}(v_n) + \limsup_{n \rightarrow \infty} |\varepsilon_{n,k}|.$$

Then, (4.36) implies that

$$\sum_{k=1}^{\infty} \mathcal{A}(w_k) \leq \limsup_{n \rightarrow \infty} \mathcal{A}(v_n). \quad (4.37)$$

The inequality (4.37) will be used below in order to show that, if $\{v_n\}$ is a Palais–Smale sequence of J_c at level $\gamma_c(c) \neq 0$, then v_n is decomposed only in a finite number of profiles w_1, \dots, w_k .

Proof of Lemma 4.12. To prove (4.35), we first remark that

$$\mathcal{Q}(u) = 4\|u\|^2 + \|u^2\|^2 - 4\langle u, u^2 \rangle.$$

Observe also that, for any $\mathbf{f} = (f_1, f_2, \dots, f_m) \in H^1(\mathbb{R})^m$, we have

$$\left\| \sum_{i=1}^m f_i \right\|^2 = \sum_{i=1}^m \|f_i\|^2 + T_1(\mathbf{f}), \quad T_1(\mathbf{f}) = \sum_{i \neq j}^m \langle f_i, f_j \rangle, \quad (4.38)$$

$$\left\| \left(\sum_{i=1}^m f_i \right)^2 \right\|^2 = \sum_{i=1}^m \|f_i^2\|^2 + T_2(\mathbf{f}), \quad T_2(\mathbf{f}) = \sum_{i \neq j}^m \langle f_i^2, f_j^2 \rangle + 2 \sum_{k; i \neq j}^m \langle f_k^2, f_i f_j \rangle + \sum_{i \neq j; k \neq \ell}^m \langle f_i f_j, f_k f_\ell \rangle, \quad (4.39)$$

$$\left\langle \sum_{i=1}^m f_i, \left(\sum_{i=1}^m f_i \right)^2 \right\rangle = \sum_{i=1}^m \langle f_i, f_i^2 \rangle + T_3(\mathbf{f}), \quad T_3(\mathbf{f}) = \sum_{i \neq j}^m \langle f_i, f_j^2 \rangle + \sum_{k; i \neq j}^m \langle f_k, f_i f_j \rangle. \quad (4.40)$$

In sum,

$$\mathcal{Q}\left(\sum_{i=1}^m f_i\right) = \sum_{i=1}^m \mathcal{Q}(f_i) + T(\mathbf{f}), \quad \text{with } T = 4T_1 + T_2 - 4T_3. \quad (4.41)$$

From now on, the notation $X \lesssim Y$ means that there exists a constant C independent of n and k such that $X \leq CY$. Since we are assuming that $\{v_n\}$ is bounded in $H^1(\mathbb{R})$, we can write $\|v_n\|_{H^1(\mathbb{R})} \lesssim 1$. Thus, it follows from (4.19) and the Sobolev's embedding that

$$\sum_{i=1}^k \|w_i\|_{L^p(\mathbb{R})} \lesssim 1, \quad \|r_{n,k}\|_{L^p(\mathbb{R})} \lesssim 1, \quad \text{for all } p \in [2, \infty]. \quad (4.42)$$

Now we apply (4.41) with $m = k + 1$, $f_i = w_{n,i}$, for $1 \leq i \leq k$, and $f_m = r_{n,k}$. Hence, one obtains (4.35) with

$$\varepsilon_{n,k} = T(w_{n,1}, \dots, w_{n,k}, r_{n,k}).$$

We aim to show that $\varepsilon_{n,k}$ satisfies (4.36).

Let us start with $T_1(w_{n,1}, \dots, w_{n,k}, r_{n,k})$, where there are two types of terms. The first type is of the form $\langle w_{n,i}, w_{n,j} \rangle$ with $i \neq j$. This case is simple to handle by using that $|y_{n,j} - y_{n,i}| \rightarrow \infty$, which leads to

$$\langle w_{n,i}, w_{n,j} \rangle = \langle w_i, w_j(\cdot - y_{n,j} + y_{n,i}) \rangle \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

The other terms in the summation $T_1(w_{n,1}, \dots, w_{n,k}, r_{n,k})$ are of the form $\langle w_{n,i}, r_{n,k} \rangle$. In this case we apply (4.18) with $\varphi = \mathcal{W} * w_i$ and $\alpha_n = -y_{n,i}$, so we get

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} |\langle w_{n,i}, r_{n,k} \rangle| = \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} |\langle w_i, r_{n,k}(\cdot + y_{n,i}) \rangle| = 0.$$

Let us study now $T_2(w_{n,1}, \dots, w_{n,k}, r_{n,k})$ and $T_3(w_{n,1}, \dots, w_{n,k}, r_{n,k})$. Here we find several types of terms. We first remark that the terms of the form $\langle w_{n,i}, w_{n,j}^2 \rangle$ and $\langle w_{n,i}^2, w_{n,j}^2 \rangle$, with $i \neq j$, and $\langle r_{n,k}, w_{n,i}^2 \rangle$ can be dealt with as we did above for the terms in $T_1(w_{n,1}, \dots, w_{n,k}, r_{n,k})$. Next we show how to treat the rest of the terms.

First we consider the terms of the form, for $i \neq j$,

$$F_n = \langle g_n, w_{n,i} w_{n,j} \rangle, \quad (4.43)$$

for some g_n with $\|g_n\|_{L^p(\mathbb{R})} \lesssim 1$ for every $p \in [2, \infty]$. In view of (4.42), this is the case when considering, for $1 \leq \ell \leq m \leq k$,

$$g_n \in \{r_{n,k}, r_{n,k}^2, r_{n,k} w_{n,\ell}, w_{n,\ell} w_{n,m}, w_{n,\ell}\}. \quad (4.44)$$

In order to deal with (4.43), by the density of $\mathcal{C}_0^\infty(\mathbb{R})$ in $H^1(\mathbb{R})$, we may consider two sequences $\{a_m\}, \{b_m\} \subset \mathcal{C}_0^\infty(\mathbb{R})$ such that $a_m \rightarrow w_i$ and $b_m \rightarrow w_j$, in $H^1(\mathbb{R})$. Of course, $\{a_m\}, \{b_m\}$ depend on i, j respectively, we do not denote explicitly this dependence for clarity. Notice that

$$F_n = A_{n,k,m} + B_{n,k,m}, \quad (4.45)$$

where

$$A_{n,k,m} = \langle g_n, w_{n,i} [w_{n,j} - b_m(\cdot - y_{n,j})] \rangle + \langle g_n, b_m(\cdot - y_{n,j}) [w_{n,i} - a_m(\cdot - y_{n,i})] \rangle$$

and

$$B_{n,k,m} = \langle g_n, b_m(\cdot - y_{n,j}) a_m(\cdot - y_{n,i}) \rangle.$$

On the one hand, we have by (1.23) and Hölder's inequality,

$$|A_{n,k,m}| \lesssim \|w_j - b_m\|_{L^2(\mathbb{R})} + \|b_m\|_{L^\infty(\mathbb{R})} \|w_i - a_m\|_{L^2(\mathbb{R})}.$$

Thus, given $\varepsilon > 0$, we may fix m , independent of n , such that $|A_{n,k,m}| < \varepsilon$ for every n . On the other hand, since a_m and b_m have compact support and $|y_{n,i} - y_{n,j}| \rightarrow \infty$ as $n \rightarrow \infty$, it follows that $B_{n,k,m} = 0$ for every n large enough. In sum,

$$\lim_{n \rightarrow \infty} F_n = 0.$$

We focus now on the terms of the form $G_n = \langle g_n, w_{n,i} r_{n,k} \rangle$, with g_n satisfying (4.44). Again by (1.23), (4.42) and Hölder's inequality, we deduce the estimate

$$|G_n| \lesssim \|w_{n,i} r_{n,k}\|_{L^2(\mathbb{R})} \lesssim \|r_{n,k}\|_{L^4(\mathbb{R})}.$$

Using (4.18) with $q = 4$, we conclude that

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} |G_n| = 0.$$

Finally, it remains to consider the terms of the form $\langle w_{n,i}, r_{n,k}^2 \rangle$ and $\langle w_{n,i}^2, r_{n,k}^2 \rangle$, which can be handled as G_n . Consequently, the proof of is complete. \square

Applying the splitting properties that we have proved to bounded Palais–Smale sequences, we obtain the following general theorem.

Theorem 4.14. *Assume that \mathcal{W} satisfies (H2), (H4), (H5), (4.5) and $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} . Let $c > 0$ and let $\{v_n\} \subset \mathcal{NV}(\mathbb{R})$ be a Palais–Smale sequence of J_c at level γ , for some $\gamma \neq 0$, i.e.*

$$J_c(v_n) \rightarrow \gamma, \quad \|J'_c(v_n)\|_{H^{-1}(\mathbb{R})} \rightarrow 0. \quad (4.46)$$

Assume in addition that there exist $R > 0$ and $\delta \in (0, 1)$ such that, for all n ,

$$\|v_n\|_{H^1(\mathbb{R})} \leq R \quad \text{and} \quad v_n \leq 1 - \delta \quad \text{on } \mathbb{R}. \quad (4.47)$$

Then there exist $k \in \mathbb{N}$ and $w_1, w_2, \dots, w_k \in \mathcal{NV}(\mathbb{R})$ such that

$$\sum_{j=1}^k J_c(w_j) = \gamma. \quad (4.48)$$

In addition, for any $1 \leq j \leq k$, the function $\rho_j = 1 - w_j$ is a nontrivial finite energy solution to (2.13) and $\rho_j \geq \delta$ on \mathbb{R} .

Proof. Since $\{v_n\}$ is bounded in $H^1(\mathbb{R})$, by Theorem 4.9, there are profiles $\{w_j\}_{j \geq 1} \subset H^1(\mathbb{R})$ and points $\{y_{n,j}\} \subset \mathbb{R}$ such that (4.17), (4.18) and (4.19) hold. In addition, as in Lemma 4.11, we infer that $w_j \leq 1 - \delta$ on \mathbb{R} , for all $j \geq 1$. Moreover, as in the proof of Theorem 1.1, we conclude that $J'_c(w_j) = 0$, so that w_j is a solution to (3.3) and $\rho_j = 1 - w_j$ is solution to (2.13). Furthermore, we see that $\{\mathcal{A}(v_n)\}$ is bounded. Since $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} , we deduce from (4.37) in Remark 4.13 that

$$\sum_{j=1}^{\infty} \mathcal{A}(w_j) \leq C, \quad (4.49)$$

for some constant $C > 0$ depending only on $\sup_n \|v_n\|_{H^1(\mathbb{R})}$ and $\|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}$.

Let us show that there is $j_0 \geq 1$ such that $w_{j_0} \neq 0$. Indeed, assuming otherwise, i.e. $w_j = 0$, for all $j \geq 1$, we deduce from (4.17) that $S_n = 0$, so that $v_n \rightarrow 0$ in $L^4(\mathbb{R})$. On the other hand, (4.46) implies that $2J_c(v_n) - J'_c(v_n)(v_n) \rightarrow 2\gamma$. This leads to a contradiction with the estimate in (3.15), since $\gamma \neq 0$.

In addition, there can only be a finite number of nonzero profiles. Indeed, if w_j is nonzero, then Lemma 4.8 provides a positive lower bound for $\mathcal{A}(w_j)$, which is independent of j . Therefore, (4.49) implies that the number of nonzero profiles is finite. Consequently, without loss of generality, we can assume that there is $k \geq 1$ such that $w_j \not\equiv 0$, for all $j \leq k$, and $w_j \equiv 0$, for all $j > k$. In this manner, the profile decomposition is finite, and we can write

$$v_n = \sum_{j=1}^k w_{n,j} + r_n, \quad (4.50)$$

with $r_n \leq 1 - \delta/2$, for n large enough, by Lemma 4.11. Also, by (4.19), (4.29) and (4.35),

$$J_c(v_n) = \sum_{j=1}^k J_c(w_j) + J_c(r_n) + o_n(1). \quad (4.51)$$

Therefore, to prove (4.48), it is enough to show that

$$J'_c(r_n)(r_n) \rightarrow 0. \quad (4.52)$$

Indeed, assuming this claim and using that $\|r_n\|_{L^4(\mathbb{R})} \rightarrow 0$, we can invoke the estimate in (3.15) to conclude that $J_c(r_n)$ converges to 0. Thus, taking the limit in (4.51), we obtain (4.48), which concludes the proof of the theorem.

To establish (4.52), recall that by (3.1),

$$J'_c(r_n)(r_n) = \int_{\mathbb{R}} r_n'^2 + \langle f(r_n), (1 - r_n)r_n \rangle - c^2 \int_{\mathbb{R}} h(r_n)r_n.$$

Remark that, if $\{z_n\}$ is bounded in $H^1(\mathbb{R})$ and $\{a_n\}$ is bounded in $L^4(\mathbb{R})$, then

$$\langle f(z_n), a_n r_n \rangle = o_n(1). \quad (4.53)$$

Indeed, this follows from the fact that $\|r_n\|_{L^4(\mathbb{R})} \rightarrow 0$ and the estimate

$$|\langle f(z_n), a_n r_n \rangle| \leq \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})} \|f(z_n)\|_{L^2(\mathbb{R})} \|a_n\|_{L^4(\mathbb{R})} \|r_n\|_{L^4(\mathbb{R})}.$$

Therefore, using also (4.50), we obtain

$$J'_c(r_n)(r_n) = \int_{\mathbb{R}} v_n' r_n' - \sum_{j=1}^k \int_{\mathbb{R}} w_{n,j}' r_n' + \langle f(r_n), r_n \rangle - c^2 \int_{\mathbb{R}} h(r_n)r_n + o_n(1). \quad (4.54)$$

On the other hand, since $J'_c(w_j) = 0$, we deduce that $J'_c(w_{n,j}) = 0$, so that, using also (4.53), we get

$$0 = J'_c(w_{n,j})(r_n) = \int_{\mathbb{R}} w_{n,j}' r_n' + \langle f(w_{n,j}), r_n \rangle - c^2 \int_{\mathbb{R}} h(w_{n,j})r_n + o_n(1). \quad (4.55)$$

Similarly, using that $\|J'_c(v_n)\|_{H^{-1}(\mathbb{R})} \rightarrow 0$ and (4.53), we obtain

$$o_n(1) = J'_c(v_n)(r_n) = \int_{\mathbb{R}} v_n' r_n' + \langle f(v_n), r_n \rangle - c^2 \int_{\mathbb{R}} h(v_n)r_n + o_n(1). \quad (4.56)$$

By putting together (4.54), (4.55) and (4.56), we conclude that

$$J'_c(r_n)(r_n) = \langle r_n, f(r_n) - f(v_n) + \sum_{j=1}^k f(w_{n,j}) \rangle - c^2 \int_{\mathbb{R}} (h(r_n) - h(v_n) + \sum_{j=1}^k h(w_{n,j}))r_n + o_n(1). \quad (4.57)$$

Notice now that, using (4.50) and that $f(s) = 2s - s^2$,

$$f(r_n) - f(v_n) + \sum_{j=1}^k f(w_{n,j}) = -r_n^2 + v_n^2 - \sum_{j=1}^k w_{n,j}^2.$$

Similarly,

$$h(r_n) - h(v_n) + \sum_{j=1}^k h(w_{n,j}) = g(r_n) - g(v_n) + \sum_{j=1}^k g(w_{n,j}), \text{ with } g(s) := h(s) - s = \frac{3s^2 - 8s + 6}{4(1-s)^3} s^2.$$

In sum, (4.57) can be simplified as

$$J'_c(r_n)(r_n) = -\langle r_n, r_n^2 - v_n^2 + \sum_{j=1}^k w_{n,j}^2 \rangle - c^2 \int_{\mathbb{R}} (g(r_n) - g(v_n) + \sum_{j=1}^k g(w_{n,j})) r_n + o_n(1). \quad (4.58)$$

Moreover, since $r_n \leq 1 - \delta/2$, $v_n \leq 1 - \delta$ and $w_{n,j} \leq 1 - \delta$, we can replace g with $\tilde{g}\chi_\delta$, where $\tilde{g}(s) = (3s^2 - 8s + 6)s^2$ and χ_δ is defined in (4.30). Applying Lemma 4.10 to (4.58) with $G(s) = s^2$ and with $G = \tilde{g}\chi_\delta$, and using (H4), we finally conclude that there is a function $o_n(1; L^1)$ such that

$$|J'_c(r_n)(r_n)| \lesssim \|\mathcal{W}\|_\infty \|r_n\|_{L^\infty(\mathbb{R})} \|o_n(1; L^1)\|_{L^1(\mathbb{R})}.$$

This completes the proof of (4.52). \square

The property (4.48) given by Theorem 4.14 can be seen as an a priori estimate for solutions obtained via splitting of Palais–Smale sequences. However, it is not clear how to use this property if J_c changes sign. The following lemma guarantees that, actually, $J_c(1 - \rho)$ is nonnegative if ρ is a finite energy solution.

Lemma 4.15. *Let $c > 0$ and assume that \mathcal{W} satisfies (H2). Then, every solution $u \in \mathcal{NE}(\mathbb{R})$ to $(S(\mathcal{W}, c))$ satisfies that*

$$J_c(1 - \rho) = \int_{\mathbb{R}} (\rho')^2 + \frac{1}{8\pi} \int_{\mathbb{R}} \xi(\widehat{\mathcal{W}})'(\xi) |\widehat{\eta}|^2,$$

where $\rho = |u|$ and $\eta = 1 - |u|^2$. In particular, if \mathcal{W} satisfies (H3), then

$$J_c(1 - \rho) \geq \int_{\mathbb{R}} (\rho')^2 \geq 0.$$

Proof. Combining (4.4) and (2.7) yields

$$\frac{c^2}{4} \int_{\mathbb{R}} \frac{\eta^2}{1 - \eta} + \frac{1}{4} \int_{\mathbb{R}} \frac{(\eta')^2}{1 - \eta} = \frac{1}{4\pi} \int_{\mathbb{R}} (\widehat{\mathcal{W}}(\xi) - \xi(\widehat{\mathcal{W}})'(\xi)) |\widehat{\eta}(\xi)|^2 d\xi.$$

Writing the left-hand side in terms of ρ and multiplying by 1/2, we arrive at

$$\frac{c^2}{8} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2} + \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 = \frac{1}{8\pi} \int_{\mathbb{R}} (\widehat{\mathcal{W}}(\xi) - \xi(\widehat{\mathcal{W}})'(\xi)) |\widehat{\eta}(\xi)|^2 d\xi.$$

Observe now that, by Plancherel's identity,

$$J_c(1 - \rho) = \frac{1}{2} \int_{\mathbb{R}} (\rho')^2 + \frac{1}{8\pi} \int_{\mathbb{R}} \widehat{\mathcal{W}}(\xi) |\widehat{\eta}(\xi)|^2 d\xi - \frac{c^2}{8} \int_{\mathbb{R}} \frac{(1 - \rho^2)^2}{\rho^2}.$$

From both previous identities, we conclude the proof. \square

We are now in position to prove a uniform estimate for solutions obtained from bounded Palais–Smale sequences.

Corollary 4.16. *Assume that \mathcal{W} satisfies (H2), (H3), (H4) and (H5). Then, for any $c \in (0, \sqrt{2})$, there exist sequences $\{c_n\} \subset (c, \sqrt{2})$ and $\{u_n\} \subset \mathcal{NE}(\mathbb{R})$ such that $c_n \rightarrow c$ and u_n is a nontrivial solution to $(S(\mathcal{W}, c_n))$ for all n . In addition, there exist $C > 0$ and $\delta_c \in (0, 1)$, independent of n , such that, denoting $\rho_n = |u_n|$,*

$$\|\rho'_n\|_{L^2(\mathbb{R})}^2 \leq J_{c_n}(1 - \rho_n) \leq \gamma_c(c_n) \leq C, \quad \text{for all } n, \quad (4.59)$$

and

$$\rho_n \geq \delta_c \quad \text{on } \mathbb{R}, \quad \text{for all } n. \quad (4.60)$$

Proof. Notice that \mathcal{W} satisfies (H3), so that (4.5) holds and (H1) is satisfied with $\sigma = 1$ and $\kappa = 0$.

Let $\mathfrak{c} \in (0, c)$. Consider the set $\mathcal{D}_{\mathfrak{c}} = \{s \in (\mathfrak{c}, \sqrt{2}) : \exists \gamma'_{\mathfrak{c}}(s)\}$. Let $\{c_n\} \in \mathcal{D}_{\mathfrak{c}}$ be a nondecreasing sequence such that $c_n \rightarrow c$. Recall that such a sequence exists thanks to (3.8). Proposition 3.10 implies that, for every fixed n , there exists a sequence $\{v_{n,m}\} \subset \mathcal{NV}(\mathbb{R})$ such that

$$\|v_{n,m}\|_{H^1(\mathbb{R})} \leq R_n, \quad v_{n,m} \leq 1 - \delta_n, \quad \lim_{m \rightarrow \infty} J_{c_n}(v_{n,m}) = \gamma_{\mathfrak{c}}(c_n), \quad \lim_{m \rightarrow \infty} \|J'_{c_n}(v_{n,m})\|_{H^{-1}(\mathbb{R})} = 0,$$

where $R_n > 0$ and $\delta_n \in (0, 1)$ do not depend on m . Therefore, Theorem 4.14 yields

$$\sum_{j=1}^{k_n} J_{c_n}(w_{j,n}) = \gamma_{\mathfrak{c}}(c_n),$$

being $1 - w_{j,n}$ nontrivial finite energy solutions to (2.13) for every $j = 1, \dots, k_n$. Let us denote $w_{1,n} = w_n$ and $\rho_n = 1 - w_n$. By Lemma 4.15, we deduce

$$\int_{\mathbb{R}} (\rho'_n)^2 \leq J_{c_n}(w_n) \leq \gamma_{\mathfrak{c}}(c_n).$$

Since $\gamma_{\mathfrak{c}}$ is a nonincreasing function and $\{c_n\}$ is nondecreasing, we conclude

$$\int_{\mathbb{R}} (\rho'_n)^2 \leq J_{c_n}(w_n) \leq \gamma_{\mathfrak{c}}(c_1).$$

Thus (4.59) follows with $C = \gamma_{\mathfrak{c}}(c_1)$.

By Lemma 2.4, there is θ_n such that $u_n = \rho_n e^{i\theta_n}$ is a finite energy solution to $(S(\mathcal{W}, c_n))$. Finally, we deduce from (4.14) in Proposition 4.7, that for all $x \in \mathbb{R}$,

$$|u_n(x)| \geq \delta_c := \inf_{s \in [\mathfrak{c}, c]} \frac{\sqrt{1 + 4s^2/V_1(s)} - 1}{\sqrt{1 + 4c^2/V_1(s)} + 1},$$

which proves (4.60). □

4.3 Passing to the limit

For any $c \in (0, \sqrt{2})$, Corollary 4.16 provides a sequence of nontrivial finite energy solutions u_n to $(S(\mathcal{W}, c_n))$ with $c_n \rightarrow c$. The last step for completing the proof of Theorem 4.1 consists of passing to the limit in $(S(\mathcal{W}, c_n))$, controlling that the limit is nontrivial and has finite energy. To this aim, the estimates proved in the previous subsections will be essential. In this subsection we adapt a technique from [7] in a similar context.

We start with a lemma that provides a sufficient condition for the boundedness of the sequence of energies.

Lemma 4.17. *Assume that the hypotheses in Corollary 4.16 hold, and let $\{\rho_n\}$ be given by Corollary 4.16. Set*

$$S_n^r = \{x \in \mathbb{R} : \rho_n(x) < r\}.$$

If for some $r \in (c/\sqrt{2}, 1)$ the sequence $\{|S_n^r|\}$ is bounded, then $\{E(u_n)\}$ is also bounded.

Proof. Let $\{c_n\}$ and $\delta_c \in (0, 1)$ be given by Corollary 4.16, and let us take some $r \in (0, 1)$. Using (4.59), (4.60), Plancherel's identity and (4.1), we obtain

$$\begin{aligned} C &\geq \gamma_c(c_n) \geq J_{c_n}(1 - \rho_n) \geq \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (1 - \rho_n^2))(1 - \rho_n^2) - \frac{c_n^2}{8} \int_{\mathbb{R}} \frac{(1 - \rho_n^2)^2}{\rho_n^2} \\ &\geq \frac{1}{4} \int_{\mathbb{R}} (\mathcal{W} * (1 - \rho_n^2))(1 - \rho_n^2) - \frac{c_n^2(1 - \delta_c^2)^2}{8\delta_c^2} |S_n^r| - \frac{c_n^2}{8r^2} \int_{\mathbb{R}} (1 - \rho_n^2)^2 \\ &\geq \frac{1}{4} \left(1 - \frac{c_n^2}{2r^2}\right) \int_{\mathbb{R}} (1 - \rho_n^2)^2 - \frac{c_n^2(1 - \delta_c^2)^2}{8\delta_c^2} |S_n^r|. \end{aligned}$$

Since $c_n \rightarrow c$, choosing $r \in (c/\sqrt{2}, 1)$ we end up with

$$\int_{\mathbb{R}} (1 - \rho_n^2)^2 \leq C_1(1 + |S_n^r|), \quad \text{for all } n,$$

for some constant $C_1 > 0$ independent of n . Thus, if $\{|S_n^r|\}$ is bounded, we get

$$\int_{\mathbb{R}} (1 - \rho_n^2)^2 \leq C_2,$$

for some $C_2 > 0$.

Recall that $J_{c_n}(1 - \rho_n) = E(u_n) - c_n p(u_n)$ (see Remark 2.3). Then, using (4.59) and (4.60) again, we conclude

$$E(u_n) = J_{c_n}(1 - \rho_n) + \frac{c_n^2}{4} \int_{\mathbb{R}} \frac{(1 - \rho_n^2)^2}{\rho_n^2} \leq C + \frac{c_n^2}{4\delta_c^2} \int_{\mathbb{R}} (1 - \rho_n^2)^2 \leq C_3,$$

for some $C_3 > 0$. □

Next result is proved as Lemma 6.6 in [7].

Lemma 4.18. *Let $\{f_n\} \subset L^1(\mathbb{R})$ be a bounded sequence, and consider a sequence $\{S_n\}$ of measurable subsets of \mathbb{R} such that $|S_n| \rightarrow \infty$. Then, for every n , there exist $x_n \in S_n$ and $R_n > 0$ with $R_n \rightarrow \infty$ such that*

$$\int_{B(x_n, R_n)} |f_n| \rightarrow 0.$$

So far, we have only used the condition (H4) for $p = 2$ or $p = \infty$. In the next lemma we will use it also for $p = 1$ in order to handle the weak star convergence, denoted by $\overset{*}{\rightharpoonup}$, in L^∞ .

Lemma 4.19. *Assume that \mathcal{W} satisfies (H4) and let $\{\eta_n\}$ be a bounded sequence in $W^{1,\infty}(\mathbb{R})$. Then there exists $\eta \in L^\infty(\mathbb{R})$ such that, up to a subsequence, $\eta_n \rightarrow \eta$ in $L_{\text{loc}}^\infty(\mathbb{R})$ and $\mathcal{W} * \eta_n \overset{*}{\rightharpoonup} \mathcal{W} * \eta$ in $L^\infty(\mathbb{R})$. In addition, for any sequence $\{f_n\} \subset L^\infty(\mathbb{R})$ such that $f_n \rightarrow f$ in $L_{\text{loc}}^\infty(\mathbb{R})$, we have the following convergence in the sense of distributions,*

$$f_n(\mathcal{W} * \eta_n) \rightarrow f(\mathcal{W} * \eta) \quad \text{in } \mathcal{D}'(\mathbb{R}). \quad (4.61)$$

Proof. First, a standard diagonal argument together with Ascoli–Arzela’s theorem imply that there exists $\eta \in L^\infty(\mathbb{R})$ such that, up to a subsequence,

$$\eta_n \rightarrow \eta \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}). \quad (4.62)$$

On the other hand, we also deduce that there is $g \in L^\infty(\mathbb{R})$ such that, up to a subsequence, $\eta_n \xrightarrow{*} g$ in $L^\infty(\mathbb{R})$. By (4.62), we get $g = \eta$. Let us now fix a function $\varphi \in L^1(\mathbb{R})$. By (H4), we have $\mathcal{W} * \varphi \in L^1(\mathbb{R})$ and, using that \mathcal{W} is even, we deduce that

$$\int_{\mathbb{R}} (\mathcal{W} * \eta_n) \varphi = \int_{\mathbb{R}} \eta_n (\mathcal{W} * \varphi) \rightarrow \int_{\mathbb{R}} \eta (\mathcal{W} * \varphi) = \int_{\mathbb{R}} (\mathcal{W} * \eta) \varphi.$$

Therefore,

$$\mathcal{W} * \eta_n \xrightarrow{*} \mathcal{W} * \eta \quad \text{in } L^\infty(\mathbb{R}). \quad (4.63)$$

To prove (4.61), we consider $\phi \in \mathcal{C}_0^\infty(\mathbb{R})$, with $\text{supp } \phi \subset K$, for some compact set K , and notice that

$$\int_{\mathbb{R}} (\mathcal{W} * \eta_n) f_n \phi = \int_{\mathbb{R}} (\mathcal{W} * \eta_n) f \phi + \int_K (\mathcal{W} * \eta_n) (f_n - f) \eta \phi. \quad (4.64)$$

The second term in the right-hand side can be bounded by $\|\mathcal{W}\|_\infty \|\eta_n\|_{L^\infty(\mathbb{R})} \|f_n - f\|_{L^\infty(K)} \|\phi\|_{L^1(K)}$, which goes to zero by hypothesis. Since $f \phi \in L^1(\mathbb{R})$, using (4.63) we can thus pass to the limit in (4.64) and obtain (4.61). \square

We are now in a position to prove Theorem 4.1.

Proof of Theorem 4.1. Let $\{c_n\}$ and $\{u_n\}$ be the sequences given by Corollary 4.16. Thus, $\{c_n\} \subset (0, \sqrt{2})$ with $c_n \rightarrow c$ and $u_n \in \mathcal{NE}(\mathbb{R})$ is a nontrivial solution to $(S(\mathcal{W}, c_n))$ for all n . We start by proving that $\{E(u_n)\}$ remains bounded. Indeed, arguing by contradiction, assume that $\{|S_n^r|\}$ is unbounded for every $r \in (c/\sqrt{2}, 1)$, where S_n^r is defined in Lemma 4.17. Let us thus fix $r \in (c/\sqrt{2}, 1)$ to be chosen later. Observe that, by (4.59), the sequence $\{(\rho'_n)^2\}$ is bounded in $L^1(\mathbb{R})$. Then, applying Lemma 4.18, for every n there exist $x_n \in S_n^r$ and $R_n > 0$ such that $R_n \rightarrow \infty$ and

$$\int_{B(0, R_n)} \rho'_n(x + x_n)^2 dx \rightarrow 0. \quad (4.65)$$

Now we define $\tilde{u}_n = u_n(\cdot + x_n)$ and $\tilde{\rho}_n = |\tilde{u}_n|$. We know from (H5) and Proposition 4.5 that $\{\tilde{u}_n\}$ is bounded in $W^{k, \infty}(\mathbb{R}; \mathbb{C})$ for every $k \in \mathbb{N}$. As a consequence, arguing as in the proof of Lemma 2.1 and taking (4.60) into account, we deduce that $\{\tilde{\rho}_n\}$ is bounded in $W^{k, \infty}(\mathbb{R})$ for every $k \in \mathbb{N}$. In particular, there exists $\rho \in W^{2, \infty}(\mathbb{R})$ such that, up to a subsequence,

$$\tilde{\rho}_n \rightarrow \rho \quad \text{in } W_{\text{loc}}^{2, \infty}(\mathbb{R}).$$

Moreover, thanks to (4.65), we deduce that $\rho' \equiv 0$, so ρ is a constant. Furthermore, the pointwise convergence $\tilde{\rho}_n \rightarrow \rho$ leads to

$$\rho_n(x_n) = \tilde{\rho}_n(0) \rightarrow \rho.$$

Therefore, since $\rho_n(x_n) < r$ for all n , it follows that $\rho \leq r$.

Notice that $\tilde{\rho}_n \in 1 + H^1(\mathbb{R})$ satisfies the equation

$$-\tilde{\rho}_n'' + \frac{c_n^2(1 - \tilde{\rho}_n^4)}{4\tilde{\rho}_n^3} = \tilde{\rho}_n(\mathcal{W} * (1 - \tilde{\rho}_n^2)) \quad \text{on } \mathbb{R}. \quad (4.66)$$

We aim to pass to the limit in (4.66). In order to do so we notice that, since ρ is a constant,

$$\tilde{\rho}_n'' \rightarrow 0 \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}).$$

Besides, by virtue of Lemma 4.19 and (4.3), we have

$$\tilde{\rho}_n(\mathcal{W} * (1 - \tilde{\rho}_n^2)) \rightarrow \rho(\mathcal{W} * (1 - \rho^2)) = \rho(1 - \rho^2)\widehat{\mathcal{W}}(0) \quad \text{in } \mathcal{D}'(\mathbb{R}).$$

Thus, using that $\widehat{\mathcal{W}}(0) = 1$, we can pass to the limit in (4.66) in $\mathcal{D}'(\mathbb{R})$ to obtain

$$c^2(1 - \rho^4) = 4\rho^4(1 - \rho^2).$$

Using that $1 - \rho^4 = (1 - \rho^2)(1 + \rho^2)$, it follows that

$$c^2(1 + \rho^2) = 4\rho^4.$$

From this equation, it is immediate to check that

$$\rho^2 = \frac{c^2 + c\sqrt{c^2 + 16}}{8}.$$

Observe that $\frac{c^2 + c\sqrt{c^2 + 16}}{8} > \frac{c^2}{2}$ since $c < \sqrt{2}$. Therefore, we can choose $\varepsilon > 0$ small enough (independent of r) so that

$$\varepsilon + \frac{c^2}{2} < \frac{c^2 + c\sqrt{c^2 + 16}}{8} = \rho^2 \leq r^2.$$

Finally, if we choose $r = \sqrt{\varepsilon + \frac{c^2}{2}}$ and take $\varepsilon > 0$ possibly smaller so that $r \in (c/\sqrt{2}, 1)$, then we arrive at a contradiction. Therefore, $\{|S_n^r|\}$ must be bounded for some $r \in (c/\sqrt{2}, 1)$. Consequently, Lemma 4.17 implies that $\{E(u_n)\}$ is bounded too.

Arguing as before, there exists $u \in \mathcal{C}^2(\mathbb{R}; \mathbb{C})$ such that, up to a subsequence, $u_n \rightarrow u$ in $W_{\text{loc}}^{2,\infty}(\mathbb{R})$. Moreover, Lemma 4.19 implies that $\mathcal{W} * (1 - |u_n|^2) \xrightarrow{*} \mathcal{W} * (1 - |u|^2)$ in $L^\infty(\mathbb{R})$. Thus, we can pass to the limit in $(S(\mathcal{W}, c_n))$ so that u is a solution to $(S(\mathcal{W}, c))$. Let us now check that $u \in \mathcal{E}(\mathbb{R})$. Indeed, using (4.1), we have

$$\frac{1}{2} \int_{\mathbb{R}} |u_n'|^2 + \frac{1}{4} \int_{\mathbb{R}} (1 - |u_n|^2)^2 \leq \sup_{n \in \mathbb{N}} E(u_n).$$

By virtue of Fatou's lemma,

$$\frac{1}{2} \int_{\mathbb{R}} |u'|^2 + \frac{1}{4} \int_{\mathbb{R}} (1 - |u|^2)^2 \leq \sup_{n \in \mathbb{N}} E(u_n).$$

That is, $u \in \mathcal{E}(\mathbb{R})$. Finally, the estimate (4.60) ensures that $u \in \mathcal{NE}(\mathbb{R})$, while Proposition 4.4 implies that u is nontrivial. The proof is concluded. \square

5 Nonexistence and properties of solitons

This section is devoted to the study of the Fourier transform of equation (2.4), that is

$$M_c(\xi)\widehat{\eta}(\xi) = \widehat{F}(\xi), \quad \text{with } M_c(\xi) = \xi^2 + 2\widehat{\mathcal{W}}(\xi) - c^2, \quad (5.1)$$

where

$$F = 2K + 2\eta(\mathcal{W} * \eta), \quad \eta = 1 - |u|^2 \quad \text{and} \quad K = |u'|^2.$$

We keep this notation for F and η for the rest of the Section, and we assume, as always, that \mathcal{W} satisfies (H0). If $M_c > 0$ a.e., we can recast (5.1) as

$$\widehat{\eta}(\xi) = L_c(\xi)\widehat{F}(\xi), \quad \text{with } L_c(\xi) = \widehat{\mathcal{L}}_c(\xi) = \frac{1}{M_c(\xi)}. \quad (5.2)$$

We will see that the operator \mathcal{L}_c plays an essential role in order to study the regularity and asymptotic behavior at infinity of the solitons given by Theorems 1.1 and 4.1.

We also stress that (H1) is sufficient condition for \mathcal{L}_c to be well defined, for $c \in [0, \sqrt{2\sigma})$. Indeed, we have $M_c(\xi) \geq (1 - 2\kappa)\xi^2 + 2\sigma - c^2 > 0$ for a.e. $\xi \in \mathbb{R}$, so that

$$\int_{\mathbb{R}} |L_c(\xi)| d\xi \leq \int_{\mathbb{R}} \frac{d\xi}{(1 - 2\kappa)\xi^2 + 2\sigma - c^2} < \infty. \quad (5.3)$$

Thus $L_c \in L^1(\mathbb{R})$ and \mathcal{L}_c is a bounded continuous function on \mathbb{R} .

We can now establish our nonexistence result for solitons with critical speed.

Theorem 5.1. *Assume that $\widehat{\mathcal{W}} \geq 0$ a.e. on \mathbb{R} and that there exists $\delta > 0$ such that one of the following holds:*

- (i) $\widehat{\mathcal{W}}(\xi) = 1 - \xi^2/2$, for all $\xi \in (-\delta, \delta)$.
- (ii) $\widehat{\mathcal{W}}$ is differentiable on $(-\delta, \delta)$, $\widehat{\mathcal{W}}(0) = 1$ and $\widehat{\mathcal{W}}(\xi) \neq 1 - \xi^2/2$ for a.e. $\xi \in (-\delta, \delta)$.

Then $(S(\mathcal{W}, \sqrt{2}))$ admits no nontrivial solution in $\mathcal{E}(\mathbb{R})$.

Proof. Arguing by contradiction, assume that there exists a nontrivial solution $u \in \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, \sqrt{2}))$. Then, Proposition 2.2 implies that (5.1) holds, i.e.

$$M(\xi)\widehat{\eta}(\xi) = \widehat{F}(\xi), \quad \text{with } M(\xi) = \xi^2 + 2\widehat{\mathcal{W}}(\xi) - 2, \quad (5.4)$$

On the other hand, by virtue of Lemma 2.1, η and K belong to $W^{k,p}(\mathbb{R})$ for all $k \in \mathbb{N}$ and all $p \in [2, \infty]$.

Let us show that \widehat{F} is continuous and $\widehat{F}(0) > 0$. Indeed, from (2.6) and from the fact that $\eta(\pm\infty) = 0$, we deduce the existence of constants $R, C > 0$ such that

$$|u'(x)|^2 \leq C(\eta(x)^2 + \eta'(x)^2) \quad \text{for all } x \in \mathbb{R} \setminus [-R, R].$$

Hence $K \in L^1(\mathbb{R})$ and, in turn, $F \in L^1(\mathbb{R})$ and \widehat{F} is continuous. Also, it follows from (5.4) that we can assume that $M\widehat{\eta}$ is also continuous. Moreover, since u is not trivial and $\widehat{\mathcal{W}} \geq 0$ a.e., Plancherel's identity yields

$$\widehat{F}(0) = \int_{\mathbb{R}} F(x) dx = 2 \int_{\mathbb{R}} |u'|^2 + \frac{1}{\pi} \int_{\mathbb{R}} \widehat{\mathcal{W}}(\xi) |\widehat{\eta}(\xi)|^2 d\xi > 0. \quad (5.5)$$

If assumption (i) holds, we deduce that $M\widehat{\eta} = 0$ on $(-\delta, \delta)$, so that, by (5.4), $\widehat{F}(0) = 0$, which contradicts (5.5).

If assumption (ii) holds, then M is differentiable on $(-\delta, \delta)$, $M(0) = 0$ and $M(\xi) \neq 0$ for a.e. $\xi \in (-\delta, \delta)$. Therefore,

$$\widehat{\eta}(\xi) = L(\xi)\widehat{F}(\xi) \quad \text{a.e. } \xi \in (-\delta, \delta), \quad (5.6)$$

where $L = 1/M$. Let us show now that $L \notin L^1((-\tilde{\delta}, \tilde{\delta}))$ for every $\tilde{\delta} \in (0, \delta)$ small enough. Expanding M around zero leads to

$$M(\xi) = M'(0)\xi + o(\xi^2), \quad \text{for all } \xi \in (-\tilde{\delta}, \tilde{\delta}),$$

for some $\tilde{\delta} \in (0, \delta)$. Thus, taking $\tilde{\delta}$ even smaller if necessary,

$$|M(\xi)| \leq (|M'(0)| + 1)|\xi|, \quad \text{for all } \xi \in (-\tilde{\delta}, \tilde{\delta}).$$

In consequence,

$$\int_{-\tilde{\delta}}^{\tilde{\delta}} L(\xi) d\xi \geq \frac{1}{|M'(0)| + 1} \int_{-\tilde{\delta}}^{\tilde{\delta}} \frac{d\xi}{|\xi|} = \infty.$$

In particular, $L \notin L^2((-\tilde{\delta}, \tilde{\delta}))$. Hence, taking (5.6) into account, in order $\widehat{\eta}$ to belong to $L^2(\mathbb{R})$, it is necessary that $\widehat{F}(0) = 0$. This is again a contradiction with (5.5). \square

We can prove now the nonexistence result stated in the Introduction.

Proof of Theorem 1.5. Let $\delta > 0$ be such that $\widehat{\mathcal{W}} \in \mathcal{C}^2((-\delta, \delta))$. Recall that $\widehat{\mathcal{W}}(0) = 1$ and $(\widehat{\mathcal{W}})'(0) = 0$. If $(\widehat{\mathcal{W}})''(\xi) = -1$ for all $(-\delta, \delta)$, then $\widehat{\mathcal{W}}(\xi) = 1 - \xi^2/2$, for $\xi \in (-\delta, \delta)$, so that we are in the case (ii) of Theorem 5.1.

Assume now that $(\widehat{\mathcal{W}})''(0) \neq -1$. Then, by decreasing δ if necessary, we deduce by continuity that $(\widehat{\mathcal{W}})'' > -1$ on $(-\delta, \delta)$, or $(\widehat{\mathcal{W}})'' < -1$ on $(-\delta, \delta)$. On the other hand, by Taylor's theorem, and using that $\widehat{\mathcal{W}}$ is even, we deduce that for any $\xi \in (-\delta, \delta)$, there exists $\tilde{\xi} \in (-\delta, \delta)$ such that

$$\widehat{\mathcal{W}}(\xi) = 1 + (\widehat{\mathcal{W}})''(\tilde{\xi}) \frac{\xi^2}{2}.$$

Thus we are in the case (ii) of Theorem 5.1, and the conclusion follows. \square

5.1 Decay at infinity

We assume now that $M_c > 0$ a.e. on \mathbb{R} so that (5.2) holds a.e. Notice also that if $L_c \in \mathcal{S}'(\mathbb{R})$, then

$$D^k \eta = \mathcal{L}_c * D^k F, \quad \text{for all } k \in \mathbb{N}. \quad (5.7)$$

This equation will be the key for analyzing the decay of the solutions $u \in \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, c))$ as $x \rightarrow \pm\infty$. More precisely, it will be provided the decay of $\eta = 1 - |u|^2$ at infinity. We start by showing that we can also recover limits of u at $\pm\infty$. First, we need to establish the integrability of η , which means that u has finite mass.

Lemma 5.2. *Let $c \geq 0$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. If $\mathcal{L}_c \in L^1(\mathbb{R})$, then $\eta \in W^{k,1}(\mathbb{R})$ for every $k \in \mathbb{N}$.*

Proof. In the case $k = 0$, we argue as in the proof of Theorem 5.1 to prove that $F \in L^1(\mathbb{R})$. Then, if $\mathcal{L}_c \in L^1(\mathbb{R})$, Young's inequality applied to (5.7) with $k = 0$ implies that $\eta \in L^1(\mathbb{R})$. The case $k \geq 1$ may be tackled similarly by taking into account that $F' = 4\eta' \mathcal{W} * \eta + 2\eta \mathcal{W} * \eta'$, so the successive derivatives of F have the form $D^k F = \sum_{j=1}^{2k} a_j b_j$, where $a_j, b_j \in L^2(\mathbb{R})$. \square

For any $u \in \mathcal{E}(\mathbb{R})$, the limits $\lim_{x \rightarrow \pm\infty} u(x)$ do not exist in general (see [26]). The following result shows that, if u solves $(S(\mathcal{W}, c))$, then they do exist whenever u presents no vortices and $\eta = 1 - |u|^2$ has finite mass.

Proposition 5.3. *Let $c > 0$ and let $u = \rho e^{i\theta} \in \mathcal{NE}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Assume that $\eta \in L^1(\mathbb{R})$. Then the following limits exist and are finite:*

$$\theta(\pm\infty) = \theta(0) + \frac{c}{2} \int_0^{\pm\infty} \frac{\eta}{1-\eta}. \quad (5.8)$$

In particular,

$$u(+\infty) = e^{i\theta(+\infty)}, \quad u(-\infty) = e^{i\theta(-\infty)}. \quad (5.9)$$

Remark 5.4. On the one hand, we recall that Lemma 5.2 provides sufficient conditions that assure that $\eta \in L^1(\mathbb{R})$. On the other hand, we stress the fact that the limits $u(\pm\infty)$, if they exist, may be different from each other. In fact, from Proposition 5.3 it is easy to see that $u(+\infty) = u(-\infty)$ if and only if $\int_{\mathbb{R}} \frac{\eta}{1-\eta} = 0$.

Proof of Proposition 5.3. By Proposition 2.4, θ satisfies (2.12). By integrating, we have

$$\theta(x) - \theta(0) = \frac{c}{2} \int_0^x \left(\frac{1}{\rho(y)^2} - 1 \right) dy = \frac{c}{2} \int_0^x \frac{\eta(y)}{1-\eta(y)} dy, \quad \text{for all } x \in \mathbb{R}.$$

Therefore,

$$\theta(+\infty) := \lim_{x \rightarrow \infty} \theta(x) = \theta(0) + \int_0^\infty \frac{\eta}{1-\eta}, \quad \theta(-\infty) := \lim_{x \rightarrow -\infty} \theta(x) = \theta(0) - \int_{-\infty}^0 \frac{\eta}{1-\eta}.$$

Since $\inf_{\mathbb{R}}(1-\eta) = \inf_{\mathbb{R}} \rho^2 > 0$ and $\eta \in L^1(\mathbb{R})$, it follows that both limits $\theta(+\infty)$ and $\theta(-\infty)$ are finite. Hence, we deduce directly (5.9) from the fact that $\rho(\pm\infty) = 1$. \square

In the rest of the subsection we will adapt the Bona–Li theory in [13, 14] to our equation. First, we recall the following technical result proved in [50].

Lemma 5.5 ([50]). *For any $0 < \ell < m$ and $\varepsilon > 0$, the following inequality holds,*

$$\int_{\mathbb{R}} \frac{e^{\ell|x|}}{(1+\varepsilon e^{|x|})^m e^{m|x-y|}} dx \leq B \frac{e^{\ell|y|}}{(1+\varepsilon e^{|y|})^m}, \quad \text{for all } y \in \mathbb{R}, \quad (5.10)$$

where $B = (\min\{\ell, m - \ell\})^{-1}$.

Proof of Theorem 1.4. First, we point out that the result holds true for $\ell = 0$. Indeed, (1.14) and Hölder's inequality yield $\mathcal{L}_c \in L^1(\mathbb{R})$. Thus, by virtue of Lemmas 2.1 and 5.2, $D^k \eta \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ for every $k \in \mathbb{N}$. We will then focus only on the case $\ell \in (0, m)$.

From (5.7), Hölder's inequality and (1.14), we deduce the following estimate,

$$|\eta(x)| \leq \int_{\mathbb{R}} |\mathcal{L}_c(x-y)| e^{m|x-y|} \frac{|F(y)|}{e^{m|x-y|}} dy \leq C_1^{\frac{1}{q}} \left(\int_{\mathbb{R}} \frac{|F(y)|^q}{e^{qm|x-y|}} dy \right)^{\frac{1}{q}}, \quad (5.11)$$

where $C_1 = \|e^{m|\cdot|} \mathcal{L}_c\|_{L^p(\mathbb{R})}^q$. We will prove next that $e^{\ell|\cdot|} \eta \in L^q(\mathbb{R})$ and $e^{\ell|\cdot|} \eta' \in L^q(\mathbb{R})$ for all $\ell \in (0, m)$. In order to do so, let $\ell \in (0, m)$ and, for all $\varepsilon \in (0, 1]$, let us consider the functions

$$h_\varepsilon(x) = \frac{e^{\ell|x|}}{(1+\varepsilon e^{|x|})^m} |\eta(x)|, \quad \tilde{h}_\varepsilon(x) = \frac{e^{\ell|x|}}{(1+\varepsilon e^{|x|})^m} |\eta'(x)|.$$

Since $\eta, \eta' \in L^\infty(\mathbb{R})$ and $\ell < m$, it is clear that $h_\varepsilon, \tilde{h}_\varepsilon \in L^q(\mathbb{R})$. Let us take now $r \in (0, q)$ and $R > 1$. Using (5.11) and Hölder's inequality with exponents $\frac{q}{q-r}$ and $\frac{q}{r}$, we deduce that

$$\begin{aligned} \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx &= \int_{\{|x|>R\}} |h_\varepsilon(x)|^{q-r} \frac{e^{\ell r|x|}}{(1+\varepsilon e^{|x|})^{rm}} |\eta(x)|^r dx \\ &\leq C_1^{\frac{r}{q}} \int_{\{|x|>R\}} |h_\varepsilon(x)|^{q-r} \frac{e^{\ell r|x|}}{(1+\varepsilon e^{|x|})^{rm}} \left(\int_{\mathbb{R}} \frac{|F(y)|^q}{e^{qm|x-y|}} dy \right)^{\frac{r}{q}} dx \\ &\leq C_1^{\frac{r}{q}} \left(\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx \right)^{\frac{q-r}{q}} \left(\int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{(1+\varepsilon e^{|x|})^{qm}} \left(\int_{\mathbb{R}} \frac{|F(y)|^q}{e^{qm|x-y|}} dy \right) dx \right)^{\frac{r}{q}}. \end{aligned}$$

From the previous inequality, one gets directly

$$\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx \leq C_1 \int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{(1+\varepsilon e^{|x|})^{qm}} \left(\int_{\mathbb{R}} \frac{|F(y)|^q}{e^{qm|x-y|}} dy \right) dx.$$

Now, by Fubini's theorem and Lemma 5.5, we derive

$$\begin{aligned} \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx &\leq C_1 \int_{\mathbb{R}} |F(y)|^q \left(\int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{(1+\varepsilon e^{|x|})^{qm} e^{qm|x-y|}} dx \right) dy \\ &\leq C_1 \int_{\{|y|>R\}} |F(y)|^q \frac{B e^{\ell q|y|}}{(1+\varepsilon e^{|y|})^{qm}} dy \\ &\quad + C_1 \int_{\{|y|\leq R\}} |F(y)|^q \int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{(1+\varepsilon e^{|x|})^{qm} e^{qm|x-y|}} dx dy. \end{aligned}$$

We will now estimate the last two integrals. On the one hand, using the inequality $\|x\| - \|y\| \leq \|x - y\|$, we obtain

$$\begin{aligned} \int_{\{|y|\leq R\}} |F(y)|^q \int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{(1 + \varepsilon e^{|x|})^{qm} e^{qm|x-y|}} dx dy &\leq \|F\|_{L^\infty(\mathbb{R})}^q \int_{\{|y|\leq R\}} \int_{\{|x|>R\}} \frac{e^{\ell q|x|}}{e^{qm|x-y|}} dx dy \\ &\leq \|F\|_{L^\infty(\mathbb{R})}^q \left(\int_{\{|y|\leq R\}} e^{qm|y|} dy \right) \left(\int_{\{|x|>R\}} e^{-(m-\ell)q|x|} dx \right) := C_2/C_1. \end{aligned}$$

On the other hand, recall that, by Lemma 2.1, $\eta(\pm\infty) = \eta'(\pm\infty) = (\mathcal{W} * \eta)(\pm\infty) = 0$. Hence, equation (2.6) in Proposition 2.2 implies that, for any fixed $\delta > 0$ we may choose $R > 1$ large enough so that

$$|F(y)|^q \leq \delta |\eta(y)|^q + \delta |\eta'(y)|^q, \quad \text{for all } |y| > R. \quad (5.12)$$

Therefore,

$$\int_{\{|y|>R\}} |F(y)|^q \frac{B e^{\ell q|y|}}{(1 + \varepsilon e^{|y|})^{qm}} dy \leq \delta B \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \delta B \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx.$$

In sum,

$$\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx \leq \delta B C_1 \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \delta B C_1 \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx + C_2. \quad (5.13)$$

We will now derive a similar estimate for \tilde{h}_ε . Indeed, from (5.7) with $k = 1$ and using (2.5), it follows that

$$\eta' = \mathcal{L}_c * (4\eta' \mathcal{W} * \eta + 2\eta \mathcal{W} * \eta').$$

Notice that $(\mathcal{W} * \eta')(\pm\infty) = 0$ too. Hence, taking $R > 1$ larger if necessary, we deduce that

$$|4\eta'(y) \mathcal{W} * \eta(y) + 2\eta(y) \mathcal{W} * \eta'(y)|^q \leq \delta |\eta(y)|^q + \delta |\eta'(y)|^q, \quad \text{for all } |y| > R.$$

This estimate allows us to follow the same arguments as we did for h_ε in order to deduce

$$\int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx \leq \delta B C_1 \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx + \delta B C_1 \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + C_3, \quad (5.14)$$

where $C_3 = C_2 \|F'\|_{L^\infty(\mathbb{R})}^q / \|F\|_{L^\infty(\mathbb{R})}^q$. Taking now $\delta \in (0, 1/2BC_1)$, it follows directly from (5.13) and (5.14) that

$$\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx \leq C_4,$$

where $C_4 = (C_2 + C_3)/(1 - 2\delta BC_1)$. By virtue of Fatou's lemma, we take limits as ε tends to zero and obtain

$$\int_{\{|x|>R\}} e^{\ell q|x|} |\eta(x)|^q dx + \int_{\{|x|>R\}} e^{\ell q|x|} |\eta'(x)|^q dx \leq C_4.$$

In conclusion, $e^{\ell \cdot} \eta \in L^q(\mathbb{R})$, $e^{\ell \cdot} \eta' \in L^q(\mathbb{R})$.

Notice that both $e^{\ell x} \eta(x)$ and $e^{-\ell x} \eta(x)$ belong to $W^{1,q}(\mathbb{R})$. Indeed, from what we have already proved, it is clear that

$$e^{\pm \ell x} \eta(x) \in L^q(\mathbb{R}), \quad \left(e^{\pm \ell x} \eta(x) \right)' = (\pm \ell \eta(x) + \eta'(x)) e^{\pm \ell x} \in L^q(\mathbb{R}).$$

Hence, the Sobolev's embedding theorem implies that $e^{\ell \cdot} \eta \in L^\infty(\mathbb{R})$ and $\lim_{x \rightarrow \pm\infty} e^{\ell|x|} \eta(x) = 0$.

We have just proved the result for $k = 0$. Taking $k = 2$ in (5.7), we deduce analogously as before that $\eta'' = \mathcal{L}_c * F''$, where F'' satisfies

$$|F''(y)|^q \leq \delta |\eta(y)|^q + \delta |\eta'(y)|^q + \delta |\eta''(y)|^q, \quad \text{for all } |y| > R.$$

Following the same process as above, and using the estimates we already have for $e^{\ell \cdot} \eta$ and $e^{\ell \cdot} \eta'$, we prove the result for $k = 1$. The complete proof follows easily by induction. \square

Conditions (1.14) in Theorem 1.4 is not easy to check, since the operator \mathcal{L}_c is not simple to compute in general. For this reason, we recall the following Paley–Wiener theorem that provides sufficient conditions on L_c that we will use when applying Theorem 1.4 to our examples in Section 6. We refer to Theorem 5.4.2 in [40] or Theorem IX.13 in [52] for details.

Theorem 5.6. *Let $T \in L^2(\mathbb{R})$. Then $e^{b|x|}T \in L^2(\mathbb{R})$ for all $b < a$, if and only if \widehat{T} has an analytical continuation to the strip $\{z \in \mathbb{C} : |z| < a\}$ with the property that for each $\zeta \in \mathbb{R}$ with $|\zeta| < a$, $\widehat{T}(\cdot + i\zeta) \in L^2(\mathbb{R})$ and for any $b < a$,*

$$\sup_{|\zeta| \leq b} \|\widehat{T}(\cdot + i\zeta)\|_{L^2(\mathbb{R})} < \infty. \quad (5.15)$$

We now tackle the algebraic decay, whose proof will follow similar lines to that of Theorem 1.4. We will employ the following lemma proved in [14].

Lemma 5.7. *For every $m, \ell \in \mathbb{R}$ such that $m > 1$ and $0 < \ell < m - 1$, there exists $B > 0$ such that the following inequality holds:*

$$\int_{\mathbb{R}} \frac{|x|^\ell}{(1 + \varepsilon|x|)^m(1 + |x - y|)^m} dx \leq \frac{B|y|^\ell}{(1 + \varepsilon|y|)^m}, \quad \text{for all } y \in \mathbb{R} \text{ and for all } \varepsilon \in (0, 1]. \quad (5.16)$$

Theorem 5.8. *Let $c \geq 0$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Assume that $L_c \in \mathcal{S}'(\mathbb{R})$ and*

$$(1 + |\cdot|)^s \mathcal{L}_c \in L^p(\mathbb{R}) \text{ for some } p \in (1, \infty], \quad s > 1 - \frac{1}{p}. \quad (5.17)$$

Setting $q = p'$ and $\eta = 1 - |u|^2$, we have

$$|\cdot|^\ell D^k \eta \in L^q(\mathbb{R}) \cap L^\infty(\mathbb{R}), \quad \lim_{x \rightarrow \pm\infty} |x|^\ell D^k \eta(x) = 0, \quad \text{for all } \ell \in \left(0, s - 1 + \frac{1}{p}\right), \quad k \in \mathbb{N}.$$

Proof. First, from (5.7) with $k = 0$, Hölder's inequality and (5.17), we deduce that

$$|\eta(x)| \leq C_1^{\frac{1}{q}} \left(\int_{\mathbb{R}} \frac{|F(y)|^q}{(1 + |x - y|)^{sq}} dy \right)^{\frac{1}{q}},$$

where $C_1 = \|(1 + |\cdot|)^s \mathcal{L}_c\|_{L^p(\mathbb{R})}^q$. Now, for $\ell \in (0, s - 1 + 1/p)$ and $\varepsilon \in (0, 1]$, we consider the functions

$$h_\varepsilon(x) = \frac{|x|^\ell}{(1 + \varepsilon|x|)^s} |\eta(x)|, \quad \tilde{h}_\varepsilon(x) = \frac{|x|^\ell}{(1 + \varepsilon|x|)^s} |\eta'(x)|.$$

Let us take $R > 1$. Arguing as in the proof of Theorem 1.4 and using Lemma 5.7, we obtain the estimate

$$\begin{aligned} \int_{\{|x| > R\}} |h_\varepsilon(x)|^q dx &\leq C_1 \int_{\{|y| > R\}} |F(y)|^q \frac{B|y|^{\ell q}}{(1 + \varepsilon|y|)^{sq}} dy \\ &+ C_1 \int_{\{|y| \leq R\}} |F(y)|^q \int_{\{|x| > R\}} \frac{|x|^{\ell q}}{(1 + \varepsilon|x|)^{sq}(1 + |x - y|)^{sq}} dx dy. \end{aligned} \quad (5.18)$$

On the one hand, since $\ell q < sq - 1$, then the function $x \mapsto \frac{|x|^{\ell q}}{(|x| + 1 - R)^{sq}}$ belongs to $L^1((R, \infty))$. Hence, the inequality $\|x - y\| \leq |x - y|$ leads to

$$\begin{aligned} \int_{\{|y| \leq R\}} |F(y)|^q \int_{\{|x| > R\}} \frac{|x|^{\ell q}}{(1 + \varepsilon|x|)^{sq}(1 + |x - y|)^{sq}} dx dy \\ \leq 2R \|F\|_{L^\infty(\mathbb{R})}^q \int_{\{|x| > R\}} \frac{|x|^{\ell q}}{(|x| + 1 - R)^{sq}} dx := C_2/C_1. \end{aligned}$$

On the other hand, as shown in the proof of Theorem 1.4, for any fixed $\delta > 0$, we may choose $R > 1$ large enough so that (5.12) holds. Therefore,

$$\int_{\{|y|>R\}} |F(y)|^q \frac{B|y|^{\ell q}}{(1 + \varepsilon|y|)^{sq}} dy \leq \delta B \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \delta B \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx.$$

In sum, by (5.18),

$$\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx \leq \delta BC_1 \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \delta BC_1 \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx + C_2.$$

Reasoning as in the proof of Theorem 1.4, we derive the analogous estimate for \tilde{h}_ε :

$$\int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx \leq \delta BC_1 \int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \delta BC_1 \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx + C_3,$$

where $C_3 = C_2 \|F'\|_{L^\infty(\mathbb{R})}^q / \|F\|_{L^\infty(\mathbb{R})}^q$. Combining the last two inequalities and taking $\delta \in (0, 1/2BC_1)$ yields

$$\int_{\{|x|>R\}} |h_\varepsilon(x)|^q dx + \int_{\{|x|>R\}} |\tilde{h}_\varepsilon(x)|^q dx \leq C_4,$$

where $C_4 = (C_2 + C_3)/(1 - 2\delta BC_1)$. By virtue of Fatou's lemma, we take limits as ε tends to zero, and obtain

$$\int_{\{|x|>R\}} |x|^{\ell q} |\eta(x)|^q dx + \int_{\{|x|>R\}} |x|^{\ell q} |\eta'(x)|^q dx \leq C_4.$$

Equivalently, $|\cdot|^\ell \eta \in L^q(\mathbb{R})$ and $|\cdot|^\ell \eta' \in L^q(\mathbb{R})$.

Let us now consider a function $\varphi : \mathbb{R} \rightarrow (0, \infty)$ that is of class \mathcal{C}^1 and satisfies that $\varphi(x) = |x|^\ell$ for every $|x| > 1$. At this point, it is clear that $\varphi \eta \in W^{1,q}(\mathbb{R})$. Hence, the Sobolev's embedding implies that $\varphi \eta \in L^\infty(\mathbb{R})$ and $\lim_{x \rightarrow \pm\infty} \varphi(x) \eta(x) = 0$. This proves the result for $k = 0$. As in the proof of Theorem 1.4, the rest of the proof follows by induction. \square

Conditions (5.17) in Theorem 5.8 can be difficult to verify. The next corollary provides sufficient (and easy to check) conditions on \mathcal{W} that guarantee (5.17) and, in turn, algebraic decay of finite energy traveling waves.

Corollary 5.9. *Assume that \mathcal{W} satisfies (H1) and that weakly differentiable up to order $s \in \mathbb{N} \setminus \{0\}$, with*

$$D^s \widehat{\mathcal{W}} \in L^\infty(\mathbb{R}). \quad (5.19)$$

Let $c \in [0, \sqrt{2\sigma})$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Then,

$$|\cdot|^\ell D^k \eta \in L^2(\mathbb{R}) \cap L^\infty(\mathbb{R}), \quad \lim_{x \rightarrow \pm\infty} |x|^\ell D^k \eta(x) = 0, \quad \text{for all } \ell \in (0, s - 1/2), k \in \mathbb{N}.$$

Proof. By (5.3), since (H1) holds, we see L_c is bounded with $L_c \in L^1(\mathbb{R})$, so that $L_c \in L^2(\mathbb{R})$. Using also (5.19), one may verify that $D^s L_c \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ too. In particular, $D^s L_c \in L^2(\mathbb{R})$. Then, applying Fourier transform, $|\cdot|^s \mathcal{L}_c \in L^2(\mathbb{R})$. Hence, since $\mathcal{L}_c \in L^\infty(\mathbb{R})$, it follows that (5.17) holds for $p = 2$ and we can apply Theorem 5.8. \square

5.2 Analyticity

Let us recall that for $H \in \mathcal{S}'(\mathbb{R})$, the associated multiplier operator \mathcal{H} is defined by

$$\widehat{\mathcal{H}(\varphi)}(\xi) = H(\xi) \widehat{\varphi}(\xi), \quad \text{for all } \varphi \in \mathcal{S}(\mathbb{R}).$$

We say that H is an L^p -multiplier, with $p \in [1, \infty]$, if there exists $\alpha > 0$ such that

$$\|\mathcal{H}(\varphi)\|_{L^p(\mathbb{R})} \leq \alpha \|\varphi\|_{L^p(\mathbb{R})}, \quad \text{for all } \varphi \in L^p(\mathbb{R}).$$

The smallest $\alpha > 0$ for which the previous inequality holds is the norm of the multiplier, and it is denoted by $\|\mathcal{H}\|_p$. For instance, by assumption (H0), \mathcal{W} is an L^2 -multiplier and, by (1.23), $\|\mathcal{W}\|_2 = \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}$.

We recall the so-called Hörmander–Mikhlin multiplier theorem in [32, 48] (see also [44]) adapted to our one-dimensional setting, as follows.

Theorem 5.10 ([32, 48]). *Let $H : \mathbb{R} \rightarrow \mathbb{R}$ be a weakly differentiable function and suppose that there exists $M > 0$ such that*

$$\sup\{|\xi^k D^k H(\xi)| : \xi \in \mathbb{R} \setminus \{0\}, k \in \{0, 1\}\} \leq M. \quad (5.20)$$

Then H is an L^p -multiplier for every $p \in (1, \infty)$. Moreover, there exists a constant $C_p > 0$, depending only on p , such that

$$\|\mathcal{H}\|_p \leq C_p M.$$

Assume that \mathcal{W} satisfies (H1), and including also the limit case $\kappa = 1/2$, and that $\widehat{\mathcal{W}}$ is (weakly) differentiable. We will apply this theorem to the function

$$H_c(\xi) = \frac{-\xi^2}{\xi^2 + 2\widehat{\mathcal{W}}(\xi) - c^2}, \quad \text{for } c \in [0, \sqrt{2\sigma}).$$

Observe that then $H_c \in L^\infty(\mathbb{R})$ and that

$$\xi H'_c(\xi) = \frac{2\xi^3(\widehat{\mathcal{W}})'(\xi) - 4\xi^2\widehat{\mathcal{W}}(\xi) + 2c^2\xi^2}{(\xi^2 + 2\widehat{\mathcal{W}}(\xi) - c^2)^2}.$$

Therefore, $\xi \mapsto \xi H'_c(\xi)$ is a bounded function if

$$|(\widehat{\mathcal{W}})'(\xi)| \leq C(|\xi| + 1) \quad \text{a.e. } \xi \in \mathbb{R}, \quad (5.21)$$

for some $C > 0$. In this case, using Theorem 5.10 we conclude that H_c is an L^p -multiplier for every $p \in (1, \infty)$. More precisely, for every $p \in (1, \infty)$, there exists a constant $\alpha_p > 0$ such that

$$\|\mathcal{H}_c(\varphi)\|_{L^p(\mathbb{R})} \leq \alpha_p \|\varphi\|_{L^p(\mathbb{R})}, \quad \text{for all } \varphi \in L^p(\mathbb{R}), \quad (5.22)$$

where $\widehat{\mathcal{H}}_c = H_c$.

Let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. We will exploit (5.7) in order to prove that η is a real analytic function. First, we prove a technical lemma.

Lemma 5.11. *Assume that there exist $\sigma \in (0, 1]$ and $\kappa \in [0, 1/2]$ such that $\widehat{\mathcal{W}}(\xi) \geq \sigma - \kappa\xi^2$ a.e. on \mathbb{R} . Assume in addition that $\widehat{\mathcal{W}}$ is weakly differentiable and that there exists $C > 0$ such that (5.21) holds. Let $c \in [0, \sqrt{2\sigma})$ and let $u \in \mathcal{E}(\mathbb{R})$ be a solution to $(S(\mathcal{W}, c))$. Let us denote*

$$\mu_k := \max\{\|D^j F'\|_{L^q(\mathbb{R})} : j = 0, \dots, k\}, \quad \text{for all } k \in \mathbb{N}.$$

Then there exist $\beta, \gamma > 0$, depending on η only through $\|D^j \eta\|_{L^2(\mathbb{R})}$ and $\|D^j \eta\|_{L^\infty(\mathbb{R})}$ for $j = 0, 1, 2$, such that

$$\|D^j \eta\|_{L^\infty(\mathbb{R})} \leq \beta \mu_k \quad \text{for all } k \in \mathbb{N}, \quad \text{for all } j = 0, \dots, k + 2, \quad (5.23)$$

$$\|D^j \eta\|_{L^2(\mathbb{R})} \leq \beta \mu_k \quad \text{for all } k \in \mathbb{N}, \quad \text{for all } j = 0, \dots, k + 2, \quad (5.24)$$

$$\mu_k \leq \gamma^k k^{k-1} \quad \text{for all } k \in \mathbb{N} \setminus \{0\}, \quad \mu_0 \leq \frac{\gamma}{24\omega\beta^2} - 1, \quad (5.25)$$

where $\omega = \|\widehat{\mathcal{W}}\|_{L^\infty(\mathbb{R})}$.

Proof. We start by proving (5.23) and (5.24). These estimates hold true for $k = 0$ by simply choosing

$$\beta \geq \frac{\max\{\|D^j \eta\|_{L^\infty(\mathbb{R})}, \|D^j \eta\|_{L^2(\mathbb{R})} : j = 0, 1, 2\}}{\|F'\|_{L^2(\mathbb{R})}}.$$

Let us take $k \geq 1$ and $j \in \{3, \dots, k+2\}$. By (5.7), we have

$$\eta''' = \mathcal{H}_c(F') \quad \text{on } \mathbb{R}, \text{ with } F' = 4\eta'(\mathcal{W} * \eta) + 2\eta(\mathcal{W} * \eta'). \quad (5.26)$$

By using also (5.22) with $p = 2$, it follows that

$$\|D^j \eta\|_{L^2(\mathbb{R})} \leq \alpha_2 \|D^{j-3} F'\|_{L^2(\mathbb{R})} \leq \alpha_2 \mu_k.$$

Moreover, by invoking the Sobolev's embedding (see Remark 3.4), we obtain

$$\begin{aligned} \|D^j \eta\|_{L^\infty(\mathbb{R})} &\leq \frac{1}{2} \left(\|D^j \eta\|_{L^2(\mathbb{R})} + \|D^{j+1} \eta\|_{L^2(\mathbb{R})} \right) \\ &\leq \frac{\alpha_2}{2} \left(\|D^{j-3} F'\|_{L^2(\mathbb{R})} + \|D^{j-2} F'\|_{L^2(\mathbb{R})} \right) \leq \alpha_2 \mu_k. \end{aligned}$$

Therefore, we take $\beta \geq \alpha_2$ so that (5.23) and (5.24) follow.

As far as (5.25) is concerned, we will prove it by induction. Indeed, it holds true for $k = 1$ if one chooses $\gamma \geq \mu_1$. Let us assume as induction hypothesis that there exists $\tilde{k} \in \mathbb{N} \setminus \{0, 1\}$ such that (5.25) holds for every $k \leq \tilde{k}$. Next we compute for $k = \tilde{k}$, taking (5.7) into account,

$$\begin{aligned} \|D^{k+1} F'\|_{L^2(\mathbb{R})} &= \|D^{k+1} (2\eta(\mathcal{W} * \eta') + 4\eta'(\mathcal{W} * \eta))\|_{L^2(\mathbb{R})} \\ &= \|2D^k (\eta'(\mathcal{W} * \eta') + \eta(\mathcal{W} * \eta'')) + 4D^k (\eta'(\mathcal{W} * \eta') + \eta'(\mathcal{W} * \eta''))\|_{L^2(\mathbb{R})} \\ &\leq 2 \sum_{j=0}^k \binom{k}{j} (\|D^{j+1} \eta(\mathcal{W} * D^{k-j+1} \eta)\|_{L^2(\mathbb{R})} + \|D^j \eta(\mathcal{W} * D^{k-j+2} \eta)\|_{L^2(\mathbb{R})}) \\ &\quad + 4 \sum_{j=0}^k \binom{k}{j} (\|(\mathcal{W} * D^{j+1} \eta) D^{k-j+1} \eta\|_{L^2(\mathbb{R})} + \|(\mathcal{W} * D^j \eta) D^{k-j+2} \eta\|_{L^2(\mathbb{R})}). \end{aligned}$$

Using (1.23), (5.23) and (5.24), we deduce that

$$\begin{aligned} \|D^{k+1} F'\|_{L^2(\mathbb{R})} &\leq 2\omega \sum_{j=0}^k \binom{k}{j} (\|D^{j+1} \eta\|_{L^\infty(\mathbb{R})} \|D^{k-j+1} \eta\|_{L^2(\mathbb{R})} + \|D^j \eta\|_{L^\infty(\mathbb{R})} \|D^{k-j+2} \eta\|_{L^2(\mathbb{R})}) \\ &\quad + 4\omega \sum_{j=0}^k \binom{k}{j} (\|D^{j+1} \eta\|_{L^2(\mathbb{R})} \|D^{k-j+1} \eta\|_{L^\infty(\mathbb{R})} + \|D^j \eta\|_{L^2(\mathbb{R})} \|D^{k-j+2} \eta\|_{L^\infty(\mathbb{R})}) \\ &\leq 12\omega \beta^2 \sum_{j=0}^k \binom{k}{j} \mu_j \mu_{k-j}. \end{aligned}$$

The induction hypothesis leads to

$$\begin{aligned} \|D^{k+1} F'\|_{L^2(\mathbb{R})} &\leq 12\omega \beta^2 \left(2\mu_0 \mu_k + \gamma^k \sum_{j=1}^{k-1} \binom{k}{j} j^{j-1} (k-j)^{k-j-1} \right) \\ &= 12\omega \beta^2 \left(2(\mu_0 \mu_k - \gamma^k k^{k-1}) + \gamma^k \sum_{j=0}^k \frac{k!}{j!(k-j)!} j^{(j-1)^+} (k-j)^{(k-j-1)^+} \right), \end{aligned}$$

where we adopt the convention $0^0 = 1$. Now, a combinatorial lemma due to Kahane [37] implies that

$$\|D^{k+1}F'\|_{L^2(\mathbb{R})} \leq 12\omega\beta^2 \left(2(\mu_0\mu_k - \gamma^k k^{k-1}) + 4\gamma^k k^{k-1}\right).$$

Using the induction hypothesis again and choosing $\gamma > 0$ large enough so that $\mu_0 \leq \frac{\gamma}{24\omega\beta^2} - 1$, we deduce that

$$\|D^{k+1}F'\|_{L^2(\mathbb{R})} \leq 24\omega\beta^2(\mu_0 + 1)\gamma^k k^{k-1} \leq \gamma^{k+1} k^{k-1}.$$

In conclusion,

$$\mu_{k+1} = \max\{\mu_k, \|D^{k+1}F'\|_{L^2(\mathbb{R})}\} \leq \max\{\gamma^k k^{k-1}, \gamma^{k+1} k^{k-1}\} = \gamma^{k+1} k^{k-1} \leq \gamma^{k+1} (k+1)^k,$$

which completes the proof. \square

We are thus led to the following analyticity of η as follows.

Theorem 5.12. *Under the hypotheses of Lemma 5.11, for every solution $u \in \mathcal{E}(\mathbb{R})$ to $(S(\mathcal{W}, c))$ with $c \in [0, \sqrt{2\sigma})$, there exists $r > 0$ such that $\eta = 1 - |u|^2$ and u have analytic extensions to the strip $\mathcal{S}_r = \{z \in \mathbb{C} : |\operatorname{Im} z| < r\}$. If $c \in (0, \sqrt{2\sigma})$, then u is real analytic on \mathbb{R} , in the sense that $\operatorname{Re}(u)$ and $\operatorname{Im}(u)$ are real analytic on \mathbb{R} .*

Proof. We need to prove that the Taylor series expansion about any point $x_0 \in \mathbb{R}$ converges with radius of convergence $r > 0$ independent of x_0 . Indeed, let $I_r = [x_0 - r, x_0 + r]$, then, by Taylor's theorem,

$$\eta(x) - \sum_{k=0}^n \frac{D^k \eta(x_0)}{k!} (x - x_0)^k = \frac{D^{n+1} \eta(\zeta)}{(n+1)!} (x - x_0)^{n+1} \quad (5.27)$$

for every $x \in I_r$ and for some $\zeta \in I_r$. Now we deduce from Lemma 5.11 that

$$|D^k \eta(\zeta)| \leq \beta \mu_k \leq \beta \gamma^k k^{k-1}$$

for every $k \in \mathbb{N} \setminus \{0\}$. Since

$$\left(\frac{\gamma^k k^{k-1}}{k!}\right)^{1/k} \rightarrow \gamma e, \quad \text{as } k \rightarrow \infty,$$

we conclude that the left-hand side of (5.27) goes to zero as $n \rightarrow \infty$ and that the radius of convergence satisfies $r \geq (\gamma e)^{-1}$. In addition, η has an analytic extensions to the strip \mathcal{S}_r .

In the case that $c \in (0, \sqrt{2\sigma})$, by Proposition 2.2, we have $\sup_{\mathbb{R}} \eta < 1$, so that $\rho = |u| = \sqrt{1 - u^2}$ is a real analytic function, as a composition of real analytic functions. Also, θ given by (2.14) is real analytic as the integral of a real analytic function. Consequently, $\operatorname{Re}(u) = \rho \cos(\theta)$ and $\operatorname{Im}(u) = \rho \sin(\theta)$ are real analytic functions on \mathbb{R} . \square

Remark 5.13. As pointed out by Corollary 4.1.5 in [14], the fact that η has an analytic extension to the strip \mathcal{S}_r implies the following exponential decay of its Fourier transform:

$$\int_{\mathbb{R}} |\hat{\eta}(\xi)|^2 e^{2\mu|\xi|} d\xi < \infty, \quad \text{for all } \mu \in (0, r).$$

We end this section by proving Corollary 1.6 as a consequence of Theorem 5.12.

Proof of Corollary 1.6. Let $\mathfrak{q} \in (0, \mathfrak{q}_*)$ and let $u = \rho e^{i\theta} \in \mathcal{NE}(\mathbb{R})$ be the nontrivial solution to $(S(\mathcal{W}, c))$, given by Theorem 1 in [22], satisfying

$$E(u) = E_{\min}(\mathfrak{q}). \quad (5.28)$$

Arguing as in the proof of Proposition 3.12 in [22], we see that there exists $a_0 \in \mathbb{R}$ such that

$$\frac{1}{2} \int_{a_0}^{\infty} (1 - \rho^2) \theta' = \frac{\mathfrak{q}}{2},$$

which allows us to define the following function

$$\tilde{u}(x) := \tilde{\rho}(x) e^{i\tilde{\theta}(x)} = \rho(x - a_0) e^{i(\theta(x - a_0) - \theta(-a_0))}.$$

Notice that \tilde{u} is nothing but u multiplied by the constant of modulus one $e^{i\theta(-a_0)}$ and translated in the space variable, so \tilde{u} is still satisfies (5.28), i.e. it is a solution to the minimization problem. Moreover, \tilde{u} satisfies that $\tilde{\theta}(0) = 0$ and

$$\frac{1}{2} \int_0^{\infty} (1 - \tilde{\rho}^2) \tilde{\theta}' = \frac{\mathfrak{q}}{2}. \quad (5.29)$$

Furthermore,

$$\frac{1}{2} \int_{-\infty}^0 (1 - \tilde{\rho}^2) \tilde{\theta}' = p(v) - \frac{1}{2} \int_0^{\infty} (1 - \tilde{\rho}^2) \tilde{\theta}' = \frac{\mathfrak{q}}{2}. \quad (5.30)$$

For notational simplicity, we continue to write u , ρ and θ for \tilde{u} , $\tilde{\rho}$ and $\tilde{\theta}$. By using the reflection operators T^{\pm} and S^{\pm} introduced in the proof of Proposition 3.12 in [22], and the fact that ρ and θ are continuous, it follows that the functions

$$u^{\pm} = (T^{\pm} \rho) e^{iS^{\pm} \theta}$$

belong to $\mathcal{NE}(\mathbb{R})$. Bearing in mind (5.29) and (5.30), we obtain that $p(u^{\pm}) = \mathfrak{q}$, which implies that $E_{\min}(\mathfrak{q}) \leq E(u^{\pm})$.

On the other hand, as in Proposition 3.12 in [22], we get

$$E(u^+) + E(u^-) = 2E_k(u) + E_p(u^+) + E_p(u^-) \quad \text{and} \quad E_p(u^+) + E_p(u^-) \leq 2E_p(u).$$

Since u satisfies (5.28), we deduce that

$$E_{\min}(\mathfrak{q}) \leq \frac{E(u^+) + E(u^-)}{2} \leq E(u) = E_{\min}(\mathfrak{q}).$$

Hence,

$$\frac{E(u^+) + E(u^-)}{2} = E(u) = E_{\min}(\mathfrak{q}).$$

Observe that

$$E_{\min}(\mathfrak{q}) \leq E(u^+) = 2E(u) - E(u^-) \leq E_{\min}(\mathfrak{q}).$$

In consequence, $E(u^+) = E(u^-) = E(u) = E_{\min}(\mathfrak{q})$. This shows that u^{\pm} and u are solutions to the minimization problem (1.5) and therefore, u^{\pm} and u satisfy $(S(\mathcal{W}, c))$, for some c depending on \mathfrak{q} . By virtue of Theorem 5.12, we have that $|u^{\pm}|^2$ and $|u|^2$ are real analytic functions. Thus, since $|u^+| = |u|$ in \mathbb{R}_+ , then $|u^+| = |u|$ in \mathbb{R} . This proves that $\rho = |u|$ is even. On the other hand, from (2.12), from the symmetry of ρ and from the fact that $\theta(0) = 0$, we derive

$$\theta(x) = \frac{c}{2} \int_0^x \left(\frac{1}{\rho(y)^2} - 1 \right) dy = -\frac{c}{2} \int_0^{-x} \left(\frac{1}{\rho(y)^2} - 1 \right) dy = -\theta(-x).$$

This concludes the proof. \square

6 Proofs of the examples

Proof of Theorem 1.7. The existence of a solution u for every $c \in (0, \sqrt{2})$ is an immediate consequence of Theorem 4.1. Also, since $\widehat{\mathcal{W}}_{\alpha, \beta}$ fulfills (5.17), $\eta = 1 - |u|^2$ is real analytic by Theorem 5.12. The nonexistence of finite energy solutions follow from Theorem 1.8 and the fact that $(\widehat{\mathcal{W}}_{\alpha, \beta})''(0) = 4\alpha\beta^{-2}(\beta - 2\alpha)^{-1} \neq -1$.

It remains to prove the exponential decay. By explicit computations, we can find for some $\beta_1, \beta_2 > 0$, depending only on α, β and c such that

$$\mathcal{L}_c(x) = \alpha_1 e^{-\beta_1|x|} + \alpha_2 e^{-\beta_2|x|}, \quad \text{for all } x \in \mathbb{R}, \quad \text{with } \alpha_1 = \frac{\beta^2 - \beta_1^2}{2\beta_1(\beta_2^2 - \beta_1^2)}, \quad \alpha_2 = \frac{\beta_2^2 - \beta^2}{2\beta_2(\beta_2^2 - \beta_1^2)}.$$

Thus, \mathcal{L}_c satisfies the condition (1.14) in Theorem 1.4 with $m = \min\{\beta_1, \beta_2\}$ and $p = \infty$. \square

Proof of Theorem 1.8. It is clear that (H0) holds for the three potentials. Notice that $2 - \cos(\lambda\xi) \geq 1$, for $\xi \in \mathbb{R}$, and using the elementary inequalities $e^x \geq 1 + x$ and $\sin(x)/x \geq 1 - x^2/6$, for $x \in \mathbb{R}$,

$$e^{-\lambda\xi^2} \geq 1 - \lambda\xi^2 \quad \text{and} \quad \frac{\sin(\lambda\xi)}{\lambda\xi} \geq 1 - \frac{\lambda^2\xi^2}{6}, \quad \text{for all } \xi \in \mathbb{R}. \quad (6.1)$$

Hence, (H1) is satisfied, with $(\sigma, \kappa) = (1, 0)$, $(\sigma, \kappa) = (1, \lambda)$ and $(\sigma, \kappa) = (1, \lambda^2/3)$, in case (i), (ii) and (iii), respectively. In particular, in the three cases we have

$$M_c(\xi) = \xi^2 + 2\widehat{\mathcal{W}}_\lambda(\xi) - c^2 \geq 2 - c^2 + \xi^2(1 - 2\kappa) > 0, \quad \text{for all } \xi \in \mathbb{R} \text{ and } c \in (0, \sqrt{2}), \quad (6.2)$$

and the existence of solutions is given by Theorem 1.1. The analyticity of $\eta = 1 - |u|^2$ follows from Theorem 5.12.

The nonexistence of finite energy solutions follows from Theorem 1.5 and the fact that in the case (i) we have $(\widehat{\mathcal{W}}_\lambda)''(0) = \lambda^2$, while in the case (ii), $(\widehat{\mathcal{W}}_\lambda)''(0) = -2\lambda$.

To prove the exponential decay, in view of (6.2), we deduce that in all the cases $\widehat{\mathcal{W}}_\lambda$ can be extended as an analytic function on \mathbb{C} . Hence, we only need to verify that for fixed $c \in (0, \sqrt{2})$ and λ , we can find a constant $\delta = \delta(c, \lambda) > 0$ such that $M_c(z) = z^2 + 2\widehat{\mathcal{W}}_\lambda(z) - c^2$ does not vanish on the strip $\mathcal{S}_\delta := \{z \in \mathbb{C} : |\operatorname{Im} z| < \delta\}$ and that $L_c(z) = (M_c(z))^{-1}$ satisfies the integrability condition in (5.15). This will imply that $e^{\delta|\cdot|}\mathcal{L}_c \in L^2(\mathbb{R})$, so that the decay follows by invoking Theorem 1.4 with $p = 2$.

Let us show that there is $\delta = \delta(c, \lambda) \in (0, 1)$ such that

$$|M_c(\xi + iw)| \geq \delta, \quad \text{for all } |w| \leq \delta, \text{ for all } \xi \in \mathbb{R}. \quad (6.3)$$

Arguing by contradiction, we get the existence of sequences $\delta_n \in (0, 1)$, $|w_n| \leq 1$, $\xi_n \in \mathbb{R}$, with $\delta_n \rightarrow 0$, $w_n \rightarrow 0$, and such that

$$T(\xi_n, w_n) = o_n(1), \quad \text{with } T(\xi, w) := \xi^2 - w^2 + 2\operatorname{Re}(\widehat{\mathcal{W}}_\lambda(\xi + iw)) - c^2, \quad (6.4)$$

$$G(\xi_n, w_n) = o_n(1), \quad \text{with } G(\xi, w) := 2\xi w + 2\operatorname{Im}(\widehat{\mathcal{W}}_\lambda(\xi + iw)). \quad (6.5)$$

By using the explicit expressions for $\widehat{\mathcal{W}}_\lambda$, it is easy to check that in the case (i), we have

$$|\widehat{\mathcal{W}}_\lambda(\xi + iw)| = |2 - \cos(\lambda\xi) \cosh(\lambda w) + i \sin(\lambda\xi) \sinh(\lambda w)| \leq 2 + \cosh(\lambda w) + \sinh(\lambda|w|). \quad (6.6)$$

In the case (ii), we get $|\widehat{\mathcal{W}}_\lambda(\xi + iw)| = |e^{-\lambda(\xi^2 - w^2 + 2i\xi w)}| \leq e^{\lambda w^2}$, while in the case (iii),

$$|\widehat{\mathcal{W}}_\lambda(\xi + iw)| = \frac{|\sin(\lambda\xi) \cosh(\lambda w) + i \cos(\lambda\xi) \sinh(\lambda w)|}{\lambda\sqrt{\xi^2 + w^2}} \leq \cosh(\lambda w) + \frac{\sinh(\lambda|w|)}{\lambda|w|}.$$

Hence, $\widehat{\mathcal{W}}_\lambda$ is bounded on the strip \mathcal{S}_1 , that is, there is $K > 0$ such that $|\widehat{\mathcal{W}}_\lambda(\xi + iw)| \leq K$, for all $\xi + iw \in \mathcal{S}_1$. Therefore, we infer from (6.4) that $\{\xi_n\}$ is bounded, so that there are $\xi_* \in \mathbb{R}$ and subsequence, that we do not relabel, such that $\xi_n \rightarrow \xi_*$. In this manner, passing to the limit in (6.4), we conclude that $T(\xi_*, 0) = 0$, i.e. $M_c(\xi_*) = 0$, which contradicts (6.2). The proof of (6.3) is completed.

By (6.3), the function

$$L_c(\xi + iw) = \frac{1}{M_c(\xi + iw)} = \frac{1}{T(\xi, w) + iG(\xi, w)}$$

defines an analytic function on the strip \mathcal{S}_δ . Also, for all $|w| \leq \delta \leq 1$, we infer the estimate

$$|L_c(\xi + iw)| \leq \begin{cases} \frac{1}{\xi^2 - 3 - 2K}, & \text{if } \xi^2 \geq 4 + 2K, \\ \delta^{-1}, & \text{otherwise.} \end{cases}$$

Consequently, $\sup_{|w| \leq \delta} \|L_c(\cdot + iw)\|_{L^2(\mathbb{R})} < \infty$, which completes the proof. \square

Remark 6.1. A careful study of the functions $T(\xi, w)$ and $G(\xi, w)$ in the proof of Theorem 1.8 should lead to the sharp exponential decay of the solitons.

Remark 6.2. Notice that if \mathcal{W}_λ is given by (1.19), then $(\widehat{\mathcal{W}}_\lambda)''(0) = -\lambda^2/3$, so that $(\widehat{\mathcal{W}}_\lambda)''(0) \neq -1$, for all $\lambda \in (0, \sqrt{3})$. However, we cannot apply Theorem 1.8 due to the change of sign of \mathcal{W}_λ .

Proof of Theorem 1.9. For $\kappa \in (0, 1/2)$, it is obvious that \mathcal{W}_κ satisfies (H0) and (H1), so we can use Theorem 1.1. For $\kappa = 1/2$, we can apply Corollary 1.2. Therefore, for any $\kappa \in (0, 1/2)$, we conclude the existence of nontrivial solutions to $(S(\mathcal{W}_\kappa, c))$ in $\mathcal{NE}(\mathbb{R})$ for almost every $c \in (0, \sqrt{2})$. Moreover, since $(\widehat{\mathcal{W}}_\kappa)'(\xi) = -2\kappa|\xi|$ for $|\xi| < 1/\sqrt{\kappa}$, and $(\widehat{\mathcal{W}}_\kappa)'(\xi) = 0$, for $|\xi| > 1/\sqrt{\kappa}$, we can apply Theorem 5.12 to obtain that $\eta = 1 - |u|^2$ is real analytic. In addition, since \mathcal{W}_κ fulfills condition (i) if $\kappa = 1/2$ in Theorem 5.1, and condition (ii) otherwise, we get the nonexistence for $c = \sqrt{2}$.

It is left to prove the algebraic decay of η . Remark that we can apply Corollary 5.9 with $s = 1$, but we can get a better decay by computing explicitly \mathcal{L}_c . In fact, since $L_c \in L^1(\mathbb{R})$, then

$$\mathcal{L}_c(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{ix\xi} L_c(\xi) d\xi = \frac{1}{2\pi} \int_{-\alpha}^{\alpha} \frac{\cos(x\xi)}{(1 - 2\kappa)\xi^2 + 2 - c^2} d\xi + \frac{1}{2\pi} \int_{\{|\xi| > \alpha\}} \frac{\cos(x\xi)}{\xi^2 - c^2} d\xi,$$

where we have used that L_c is even. Now, after applying integration by parts twice, we get

$$\mathcal{L}_c(x) = \frac{1}{x^2} (A \cos(\alpha x) + g(x)) \quad \text{for all } x \neq 0, \text{ where } A = \frac{4\alpha\kappa}{\pi(\alpha^2 - c^2)^2}$$

and

$$g(x) = -\frac{2}{\pi} \int_{\alpha}^{\infty} \frac{(3\xi^2 + c^2) \cos(x\xi)}{(\xi^2 - c^2)^3} d\xi - \frac{2(1 - 2\kappa)}{\pi} \int_0^{\alpha} \frac{(3(1 - 2\kappa)\xi^2 + 2 - c^2) \cos(x\xi)}{((1 - 2\kappa)\xi^2 + 2 - c^2)^3} d\xi.$$

Observe that $g \in L^\infty(\mathbb{R})$. We are not interested in the value $\mathcal{L}_c(0)$. We simply remark that, since $L_c \in L^1(\mathbb{R})$, it follows that $\mathcal{L}_c \in L^\infty(\mathbb{R})$ and therefore, $(1 + |\cdot|^2)\mathcal{L}_c \in L^\infty(\mathbb{R})$. Consequently, the decay in (1.22) follows by applying Theorem 5.8 with $s = 2$ and $p = \infty$. \square

Acknowledgments. The authors acknowledge support from the Labex CEMPI (ANR-11-LABX-0007-01). A. de Laire was also supported by the ANR project ODA (ANR-18-CE40-0020-01). S. López-Martínez was also supported by PGC2018-096422-B-I00 (MCIU/AEI/FEDER, UE) and Junta de Andalucía FQM-116.

Appendix

We include here the proof of the deformation lemma.

Proof of Lemma 3.3. For $j = 1, 2, 3$, let us denote

$$A_j = J_c^{-1}([\gamma - 2\varepsilon, \gamma + 2\varepsilon]) \cap Z_{\delta_j}.$$

Since these are closed sets in $H^1(\mathbb{R})$, we may define a functional $\psi : H^1(\mathbb{R}) \rightarrow \mathbb{R}$ of class \mathcal{C}^1 such that $0 \leq \psi(v) \leq 1$ for every $v \in H^1(\mathbb{R})$ and

$$\psi(v) = \begin{cases} 1 & \text{for all } v \in A_2, \\ 0 & \text{for all } v \in H^1(\mathbb{R}) \setminus A_1. \end{cases}$$

Let us now consider the vector field $\varphi : H^1(\mathbb{R}) \rightarrow H^1(\mathbb{R})$ given by

$$\varphi(v) = \begin{cases} -\psi(v) \frac{J'_c(v)}{\|J'_c(v)\|_{H^{-1}(\mathbb{R})}} & \text{for all } v \in A_1, \\ 0 & \text{for all } v \in H^1(\mathbb{R}) \setminus A_1. \end{cases}$$

Clearly, $\varphi \in \mathcal{C}^1(H^1(\mathbb{R}))$ (see Lemma 3.1) and $\|\varphi(v)\|_{H^1(\mathbb{R})} \leq 1$ for every $v \in H^1(\mathbb{R})$.

For any $v \in H^1(\mathbb{R})$, we consider the Cauchy problem

$$\begin{cases} w'(t) = \varphi(w(t)), & \text{for all } t \geq 0, \\ w(0) = v. \end{cases}$$

The classical ODE theory, the Cauchy problem has a unique solution $w(\cdot, v) \in H^1(\mathbb{R})$ defined in $[0, +\infty)$. Let us show that $w(t, \mathcal{NV}(\mathbb{R})) \subset \mathcal{NV}(\mathbb{R})$ for every $t \geq 0$. Indeed, let $v \in \mathcal{NV}(\mathbb{R})$. Clearly, $w(0, v) = v \in \mathcal{NV}(\mathbb{R})$. Moreover, since $w(\cdot, v)$ is continuous and $v \in \mathcal{NV}(\mathbb{R})$, there exists $\tilde{t} > 0$ such that $w(t, v) \in \mathcal{NV}(\mathbb{R})$ for every $t \in [0, \tilde{t}]$. Let us assume by contradiction that $s := \sup\{\tilde{t} > 0 : w(t, v) \in \mathcal{NV}(\mathbb{R}) \forall t \in [0, \tilde{t}]\} < +\infty$. Then, $w(s, v) \in \partial\mathcal{NV}(\mathbb{R})$. In particular, $w(s, v) \in H^1(\mathbb{R}) \setminus A_1$, so $\varphi(w(s, v)) = 0$. Actually, since $H^1(\mathbb{R}) \setminus A_1$ is open, then there exists $\tilde{s} \in (0, s)$ such that $w(t, v) \in H^1(\mathbb{R}) \setminus A_1$ for every $t \in [s - \tilde{s}, s]$. Therefore, $\varphi(w(t, v)) = 0$ for every $t \in [s - \tilde{s}, s]$. That is, $w'(t, v) = 0$ for every $t \in [s - \tilde{s}, s]$, so w must be constant in $[s - \tilde{s}, s]$ and, in consequence, $w(s - \tilde{s}, v) = w(s, v)$. But this is a contradiction since, by definition of s , it is necessary that $w(s - \tilde{s}, v) \in \mathcal{NV}(\mathbb{R})$.

On the other hand, for any $v \in H^1(\mathbb{R})$ and $t \geq 0$, we have

$$\|w(t, v) - v\|_{L^\infty(\mathbb{R})} \leq \|w(t, v) - v\|_{H^1(\mathbb{R})} \leq \left\| \int_0^t \varphi(w(s, v)) ds \right\|_{H^1(\mathbb{R})} \leq t.$$

Observe that, in the previous inequality, we have used that the norm of the continuous embedding $H^1(\mathbb{R}) \subset L^\infty(\mathbb{R})$ is equal to one (see Remark 3.4). Hence, we deduce that $w(t, Z_{\delta_3}) \subset Z_{\delta_2}$ for every $t \leq \delta_3 - \delta_2$.

Let us define $h : [0, 1] \times \mathcal{NV}(\mathbb{R}) \rightarrow H^1(\mathbb{R})$ by

$$h(t, v) = w((\delta_3 - \delta_2)t, v), \quad \text{for all } (t, v) \in [0, 1] \times \mathcal{NV}(\mathbb{R}).$$

We have already verified that $h([0, 1] \times \mathcal{NV}(\mathbb{R})) \subset \mathcal{NV}(\mathbb{R})$. Furthermore, with this definition, items (i) and (iii) are obviously satisfied. On the other hand, if $v \in \mathcal{NV}(\mathbb{R}) \setminus A_1$, then $\varphi(v) = 0$, so $w(t) = v$ is the unique solution to the Cauchy problem and item (ii) is satisfied too.

As far as item (iv) is concerned, let $v \in \mathcal{NV}(\mathbb{R})$. Since $w(t, v) \in \mathcal{NV}(\mathbb{R})$ for every $t \geq 0$, then the function $J_c(w(\cdot, v))$ is differentiable and

$$\frac{d}{dt}J_c(w(t, v)) = \langle J'_c(w(t, v)), w'(t, v) \rangle = \langle J'_c(w(t, v)), \varphi(w(t, v)) \rangle \leq 0.$$

Thus, $J_c(w(\cdot, v))$ is nonincreasing and item (iv) holds true.

Lastly, we check item (v). Indeed, let $v \in J_c^{\gamma+\varepsilon} \cap Z_{\delta_3}$. If there exists $t \in [0, \delta_3 - \delta_2)$ such that $J_c(w(t, v)) < \gamma - \varepsilon$, then item (iv) implies that $J_c(w(\delta_3 - \delta_2, v)) < \gamma - \varepsilon$, so $w(\delta_3 - \delta_2, v) \in J_c^{\gamma-\varepsilon} \cap Z_{\delta_2}$. Otherwise, for every $t \in [0, \delta_3 - \delta_2)$, one has

$$\gamma - \varepsilon \leq J_c(w(t, v)) \leq J_c(w(0, v)) = J_c(v) \leq \gamma + \varepsilon.$$

In particular, $w(t, v) \in A_2$ for every $t \in [0, \delta_3 - \delta_2)$. In addition, by the definition of φ , we derive

$$\begin{aligned} J_c(w(\delta_3 - \delta_2, v)) &= J_c(v) + \int_0^{\delta_3 - \delta_2} \frac{d}{dt}J_c(w(t, v))dt \\ &= J_c(v) + \int_0^{\delta_3 - \delta_2} \langle J'_c(w(t, v)), \varphi(w(t, v)) \rangle dt \\ &= J_c(v) - \int_0^{\delta_3 - \delta_2} \|J'_c(w(t, v))\|_{H^{-1}(\mathbb{R})} dt \\ &\leq \gamma + \varepsilon - (\delta_3 - \delta_2) \frac{2\varepsilon}{\delta_3 - \delta_2} = \gamma - \varepsilon. \end{aligned}$$

This concludes the proof. \square

We include also a technical lemma needed in the proof of Proposition 3.9. Even though the proof of the lemma is elementary, it is not straightforward and a sort of uniform continuity of the functional J_c is required.

Lemma 6.3. *Let $c > 0$ and fix $\delta_c \in (0, 1)$, $R_c > 0$ and $\gamma_c > 0$. For every $\alpha > 0$ and $\delta \in (0, 1)$, we consider the set*

$$X_{\alpha, \delta} = J_c^{-1}((\gamma_c - \alpha, \gamma_c + \alpha)) \cap Z_\delta,$$

where Z_δ is defined by (3.5), i.e.

$$Z_\delta = \{v \in \mathcal{NV}(\mathbb{R}) : \|v\|_{H^1(\mathbb{R})} \leq R_c + 1 - \delta, v \leq 1 - \delta \text{ on } \mathbb{R}\}.$$

Let us also denote

$$I_{\alpha, \delta} = \inf\{\|J'_c(v)\|_{H^{-1}(\mathbb{R})} : v \in X_{\alpha, \delta}\}.$$

Assume that there exists $\alpha > 0$ such that $I_{\alpha, \delta_c} > 0$. Then there exists $\bar{\delta} \in (0, \delta_c)$ such that $I_{\alpha, \bar{\delta}} > 0$.

Proof. Let us take $\delta \in (0, \delta_c)$. Observe that $Z_{\delta_c} \subset Z_\delta$. Recall that $J_c \in \mathcal{C}^2(\mathcal{NV}(\mathbb{R}))$, see Lemma 3.1. It is simple to check from (3.2) that $\|J''_c(v)\| \leq C$ for every $v \in Z_\delta$, where $C > 0$ depends only on R_c and δ . Hence, since Z_δ is connected and convex, the Mean Value theorem implies that J'_c is Lipschitz in Z_δ , with Lipschitz constant denoted by $l_\delta > 0$. In consequence, for every $\varepsilon > 0$, if we take $\beta = \varepsilon/l_\delta$, then the following holds for any $v, w \in Z_\delta$ satisfying $\|v - w\|_{H^1(\mathbb{R})} < \beta$,

$$\left| \|J'_c(v)\|_{H^{-1}(\mathbb{R})} - \|J'_c(w)\|_{H^{-1}(\mathbb{R})} \right| \leq \|J'_c(v) - J'_c(w)\|_{H^{-1}(\mathbb{R})} \leq l_\delta \|v - w\|_{H^1(\mathbb{R})} < \varepsilon.$$

In the previous inequality, we take $\varepsilon = I_{\alpha, \delta_c}/2$. Therefore, if $v \in Z_\delta$, $w \in X_{\alpha, \delta_c}$ and $\|v - w\|_{H^1(\mathbb{R})} \leq \beta$, then

$$\|J'_c(v)\|_{H^{-1}(\mathbb{R})} > \|J'_c(w)\|_{H^{-1}(\mathbb{R})} - \frac{I_{\alpha, \delta_c}}{2} \geq \frac{I_{\alpha, \delta_c}}{2} > 0.$$

We have proved that

$$\|J'_c(v)\|_{H^{-1}(R)} \geq \frac{I_{\alpha, \delta_c}}{2} > 0 \quad \text{for all } v \in \mathcal{X}, \quad (6.7)$$

where

$$\mathcal{X} = \{v \in X_{\alpha, \delta} : \text{dist}(v, X_{\alpha, \delta_c}) < \beta\}.$$

Notice that

$$X_{\alpha, \delta_c} \subset \mathcal{X} \subset X_{\alpha, \delta}.$$

Using (6.7), the proof of the lemma will be finished as soon as we show that there exists $\bar{\delta} \in (\delta, \delta_c)$ such that

$$X_{\alpha, \delta_c} \subset X_{\alpha, \bar{\delta}} \subset \mathcal{X} \subset X_{\alpha, \delta}.$$

In order to do so, let $\bar{\delta} \in (\delta, \delta_c)$ to be chosen later, and let $v \in X_{\alpha, \bar{\delta}}$. For some $\lambda > 0$ to be chosen later too, we aim to prove that

$$\lambda v \in X_{\alpha, \delta_c} \quad \text{and} \quad \|v - \lambda v\|_{H^1(\mathbb{R})} < \beta, \quad (6.8)$$

which implies that $v \in \mathcal{X}$.

On the one hand, simple computations show that a sufficient condition for $\lambda v \in Z_{\delta_c}$ and $\|v - \lambda v\|_{H^1(\mathbb{R})} < \beta$ is

$$1 - \frac{\beta}{R_c + 1 - \bar{\delta}} < \lambda < \frac{1 - \delta_c}{1 - \bar{\delta}}. \quad (6.9)$$

Observe that $\lambda > 0$ can be chosen so that (6.9) holds whenever $\bar{\delta}$ is close enough to δ_c .

On the other hand, it is left to prove that $J_c(\lambda v) \in (\gamma_c - \alpha, \gamma_c + \alpha)$. Indeed, since J_c is uniformly continuous in Z_δ and $J_c(v) \in (\gamma_c - \alpha, \gamma_c + \alpha)$, it follows that there exists $\lambda_0 \in (0, 1)$, independent of v , such that $J_c(\lambda v) \in (\gamma_c - \alpha, \gamma_c + \alpha)$ for every $\lambda \in (\lambda_0, 1]$. Now we take $\bar{\delta}$ even closer to δ_c so that $\lambda_0 < \frac{1 - \delta_c}{1 - \bar{\delta}}$. Thus, for every $\lambda \in (\lambda_0, 1)$ satisfying (6.9), we have that (6.8) holds. The proof is finished. \square

References

- [1] A. Aftalion, X. Blanc, and R. Jerrard. Nonclassical rotational inertia of a supersolid. *Phys. Rev. Lett.*, 99(13):135301.1–135301.4, 2007.
- [2] J. Albert. Positivity properties and uniqueness of solitary wave solutions of the intermediate long-wave equation. In *Evolution equations (Baton Rouge, LA, 1992)*, volume 168 of *Lecture Notes in Pure and Appl. Math.*, pages 11–20. Dekker, New York, 1995.
- [3] P. Antonelli and C. Sparber. Existence of solitary waves in dipolar quantum gases. *Phys. D*, 240(4-5):426–431, 2011.
- [4] V. Banica and T. Duyckaerts. Global existence, scattering and blow-up for the focusing NLS on the hyperbolic space. *Dyn. Partial Differ. Equ.*, 12(1):53–96, 2015.
- [5] C. Becker, S. Stellmer, P. Soltan-Panahi, S. Dörscher, M. Baumert, E.-M. Richter, J. Kronjäger, K. Bongs, and K. Sengstock. Oscillations and interactions of dark and dark–bright solitons in Bose-Einstein condensates. *Nature Physics*, 4(6):496, 2008.
- [6] J. Bellazzini and L. Jeanjean. On dipolar quantum gases in the unstable regime. *SIAM J. Math. Anal.*, 48(3):2028–2058, 2016.
- [7] J. Bellazzini and D. Ruiz. Finite energy traveling waves for the Gross-Pitaevskii equation in the subsonic regime, Preprint arXiv:1911.02820.

- [8] N. G. Berloff and P. H. Roberts. Motions in a Bose condensate VI. Vortices in a nonlocal model. *J. Phys. A*, 32(30):5611–5625, 1999.
- [9] F. Béthuel, P. Gravejat, and J.-C. Saut. Existence and properties of travelling waves for the Gross-Pitaevskii equation. In *Stationary and time dependent Gross-Pitaevskii equations*, volume 473 of *Contemp. Math.*, pages 55–103. Amer. Math. Soc., Providence, RI, 2008.
- [10] F. Béthuel, P. Gravejat, and J.-C. Saut. Travelling waves for the Gross-Pitaevskii equation. II. *Comm. Math. Phys.*, 285(2):567–651, 2009.
- [11] F. Béthuel and J.-C. Saut. Travelling waves for the Gross-Pitaevskii equation I. *Ann. Inst. H. Poincaré Phys. Théor.*, 70(2):147–238, 1999.
- [12] N. N. Bogoliubov. On the theory of superfluidity. *J. Phys. USSR*, 11:23–32, 1947. Reprinted in: D. Pines, *The Many-Body Problem* (W. A. Benjamin, New York, 1961), p. 292-301.
- [13] J. Bona and Y. Li. Analyticity of solitary-wave solutions of model equations for long waves. *SIAM J. Math. Anal.*, 27(3):725–737, 1996.
- [14] J. Bona and Y. Li. Decay and analyticity of solitary waves. *J. Math. Pures Appl.*, 76(5):377–430, 1997.
- [15] V. I. Burenkov. *Sobolev spaces on domains*, volume 137 of *Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]*. B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1998.
- [16] R. Carles, P. A. Markowich, and C. Sparber. On the Gross-Pitaevskii equation for trapped dipolar quantum gases. *Nonlinearity*, 21(11):2569–2590, 2008.
- [17] C. Coste. Nonlinear Schrödinger equation and superfluid hydrodynamics. *Eur. Phys. J. B Condens. Matter Phys.*, 1(2):245–253, 1998.
- [18] A. de Laire. Non-existence for travelling waves with small energy for the Gross-Pitaevskii equation in dimension $N \geq 3$. *C. R. Math. Acad. Sci. Paris*, 347(7-8):375–380, 2009.
- [19] A. de Laire. Global well-posedness for a nonlocal Gross-Pitaevskii equation with non-zero condition at infinity. *Comm. Partial Differential Equations*, 35(11):2021–2058, 2010.
- [20] A. de Laire. Minimal energy for the traveling waves of the Landau-Lifshitz equation. *SIAM J. Math. Anal.*, 46(1):96–132, 2014.
- [21] A. de Laire, G. Dujardin, and S. López-Martínez. Numerical computation of some solitons of a nonlocal nonlinear Schrödinger equation. *Preprint*.
- [22] A. de Laire and P. Mennumi. Traveling waves for some nonlocal 1D Gross-Pitaevskii equations with nonzero conditions at infinity. *Discrete Contin. Dyn. Syst.*, 40(1):635–682, 2020.
- [23] J. Denschlag, J. E. Simsarian, D. L. Feder, C. W. Clark, L. A. Collins, J. Cubizolles, L. Deng, E. W. Hagley, K. Helmerson, W. P. Reinhardt, et al. Generating solitons by phase engineering of a Bose-Einstein condensate. *Science*, 287(5450):97–101, 2000.
- [24] A. Farina. From Ginzburg-Landau to Gross-Pitaevskii. *Monatsh. Math.*, 139(4):265–269, 2003.
- [25] G. B. Folland. *Real analysis*. Pure and Applied Mathematics (New York). John Wiley & Sons, Inc., New York, second edition, 1999. Modern techniques and their applications, A Wiley-Interscience Publication.

- [26] P. Gérard. The Gross-Pitaevskii equation in the energy space. In A. Farina and J.-C. Saut, editors, *Stationary and time dependent Gross-Pitaevskii equations. Wolfgang Pauli Institute 2006 thematic program, January–December, 2006, Vienna, Austria*, volume 473 of *Contemporary Mathematics*, pages 129–148. American Mathematical Society.
- [27] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2008.
- [28] L. Grafakos. *Modern Fourier analysis*, volume 250 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [29] P. Gravejat. Decay for travelling waves in the Gross-Pitaevskii equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 21(5):591–637, 2004.
- [30] E. Gross. Hydrodynamics of a superfluid condensate. *J. Math. Phys.*, 4(2):195–207, 1963.
- [31] T. Hmidi and S. Keraani. Blowup theory for the critical nonlinear Schrödinger equations revisited. *Int. Math. Res. Not.*, (46):2815–2828, 2005.
- [32] L. Hörmander. *The analysis of linear partial differential operators I*. Classics in Mathematics. Springer-Verlag, Berlin, 2003.
- [33] L. Jeanjean. On the existence of bounded Palais-Smale sequences and application to a Landesman-Lazer-type problem set on \mathbf{R}^N . *Proc. Roy. Soc. Edinburgh Sect. A*, 129(4):787–809, 1999.
- [34] C. A. Jones, S. J. Putterman, and P. H. Roberts. Motions in a Bose condensate V. Stability of solitary wave solutions of non-linear Schrödinger equations in two and three dimensions. *J. Phys. A, Math. Gen.*, 19(15):2991–3011, 1986.
- [35] C. A. Jones and P. H. Roberts. Motions in a Bose condensate IV. Axisymmetric solitary waves. *J. Phys. A, Math. Gen.*, 15(8):2599–2619, 1982.
- [36] C. Josserand, Y. Pomeau, and S. Rica. Coexistence of ordinary elasticity and superfluidity in a model of a defect-free supersolid. *Phys. Rev. Lett.*, 98(19):195301.1–195301.4, 2007.
- [37] C. Kahane. On the spatial analyticity of solutions of the Navier-Stokes equations. *Arch. Rational Mech. Anal.*, 33:386–405, 1969.
- [38] Y. V. Kartashov and L. Torner. Gray spatial solitons in nonlocal nonlinear media. *Opt. Lett.*, 32(8):946–948, 2007.
- [39] Y. S. Kivshar and B. Luther-Davies. Dark optical solitons: physics and applications. *Phys. Rep.*, 298(2-3):81–197, 1998.
- [40] S. G. Krantz and H. R. Parks. *A primer of real analytic functions*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Boston, Inc., Boston, MA, second edition, 2002.
- [41] W. Królikowski and O. Bang. Solitons in nonlocal nonlinear media: Exact solutions. *Physical Review E*, 63(1):016610, 2000.
- [42] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau. The physics of dipolar bosonic quantum gases. *Reports on Progress in Physics*, 72(12):126401, 2009.
- [43] E. H. Lieb. Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation. *Studies in Appl. Math.*, 57(2):93–105, 1976/77.

- [44] W. Littman. Multipliers in L^p and interpolation. *Bull. Amer. Math. Soc.*, 71:764–766, 1965.
- [45] S. Lopez-Aguayo, A. S. Desyatnikov, Y. S. Kivshar, S. Skupin, W. Krolikowski, and O. Bang. Stable rotating dipole solitons in nonlocal optical media. *Opt. Lett.*, 31(8):1100–1102, 2006.
- [46] Y. Luo and A. Stylianou. Ground states for a nonlocal mixed order cubic-quartic Gross-Pitaevskii equation. *J. Math. Anal. Appl.*, 496(1):124802, 20, 2021.
- [47] M. Mariş. Traveling waves for nonlinear Schrödinger equations with nonzero conditions at infinity. *Ann. of Math. (2)*, 178(1):107–182, 2013.
- [48] S. G. Mikhlin. On the multipliers of Fourier integrals. *Dokl. Akad. Nauk SSSR (N.S.)*, 109:701–703, 1956.
- [49] N. I. Nikolov, D. Neshev, W. Królikowski, O. Bang, J. J. Rasmussen, and P. L. Christiansen. Attraction of nonlocal dark optical solitons. *Optics letters*, 29(3):286–288, 2004.
- [50] L. Pei. Exponential decay and symmetry of solitary waves to Degasperis-Procesi equation. *J. Differential Equations*, 269(10):7730–7749, 2020.
- [51] L. Pitaevskii. Vortex lines in an imperfect Bose gas. *Sov. Phys. JETP*, 13(2):451–454, 1961.
- [52] M. Reed and B. Simon. *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975.
- [53] J. Reneuve, J. Salort, and L. Chevillard. Structure, dynamics, and reconnection of vortices in a nonlocal model of superfluids. *Physical Review Fluids*, 3(11):114602, 2018.
- [54] M. Struwe. The existence of surfaces of constant mean curvature with free boundaries. *Acta Math.*, 160(1-2):19–64, 1988.
- [55] C. Tintarev. *Concentration compactness*, volume 33 of *De Gruyter Series in Nonlinear Analysis and Applications*. De Gruyter, Berlin, 2020. Functional-analytic theory of concentration phenomena.
- [56] K. Tintarev and K.-H. Fieseler. *Concentration compactness*. Imperial College Press, London, 2007. Functional-analytic grounds and applications.
- [57] J. Van Schaftingen and J. Xia. Groundstates for a local nonlinear perturbation of the Choquard equations with lower critical exponent. *J. Math. Anal. Appl.*, 464(2):1184–1202, 2018.
- [58] H. Veksler, S. Fishman, and W. Ketterle. Simple model for interactions and corrections to the Gross-Pitaevskii equation. *Phys. Rev. A*, 90(2):023620, 2014.
- [59] M. Willem. *Minimax theorems*, volume 24 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 1996.