



Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers

Antoine Stier, Bin-Yan Hsu, Coline Marciau, Blandine Doligez, Lars Gustafsson, Pierre Bize, Suvi Ruuskanen

► To cite this version:

Antoine Stier, Bin-Yan Hsu, Coline Marciau, Blandine Doligez, Lars Gustafsson, et al.. Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers. *Biology Letters*, 2020, 16 (11), pp.20200364. 10.1098/rsbl.2020.0364 . hal-03424494

HAL Id: hal-03424494

<https://hal.science/hal-03424494>

Submitted on 29 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers

Journal:	<i>Biology Letters</i>
Manuscript ID	RSBL-2020-0364.R1
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Stier, Antoine; University of Turku, Department of Biology; University of Glasgow, Department of Biology Hsu, Bin-Yan; Turun Yliopisto, Department of Biology Marciau, Coline; Turun Yliopisto, Department of Biology Doligez, Blandine; CNRS, Department of Biometry and Evolutionary Biology Gustafsson, Lars; Uppsala Universitet, Ecology and Genetics/Animal Ecology Bize, Pierre; University of Aberdeen School of Social Science; Ruuskanen, Suvi; University of Turku, Department of Biology
Subject:	Developmental biology < BIOLOGY, Ecology < BIOLOGY
Categories:	Physiology
Keywords:	Ageing, mitochondria, telomere length, bird, fetal programming

Author-supplied statements

Relevant information will appear here if provided.

Ethics

Does your article include research that required ethical approval or permits?:

Yes

Statement (if applicable):

The study was conducted in the long-term monitored population of collared flycatchers on Gotland, Sweden (Jordbruksverkets permit no. ID 872)

Data

It is a condition of publication that data, code and materials supporting your paper are made publicly available. Does your paper present new data?:

Yes

Statement (if applicable):

Data [25] used in this article is publicly available at:
<https://doi.org/10.6084/m9.figshare.11688504.v1>

Conflict of interest

I/We declare we have no competing interests

Statement (if applicable):

CUST_STATE_CONFLICT :No data available.

Authors' contributions

This paper has multiple authors and our individual contributions were as below

Statement (if applicable):

All authors agree to be held accountable for the content therein and approve the final version of the manuscript. AS, BYH & SR designed the study. BYH and SR conducted the fieldwork, AS and CM conducted laboratory work. BD, LG & PB contributed to data collection. AS analyzed the data and wrote the manuscript with input from all authors.

Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers

Antoine Stier^{1,2*}, Bin-Yan Hsu¹, Coline Marciau¹, Blandine Doligez³, Lars Gustafsson⁴, Pierre Bize⁵ and Suvi Ruuskanen¹

¹ Department of Biology, University of Turku, Turku, Finland

² Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow, UK

³ Department of Biometry and Evolutionary Biology, CNRS, Université Lyon 1, France

⁴ Department of Ecology and Genetics / Animal Ecology, University of Uppsala, Uppsala, Sweden

⁵ School of Biological Sciences, University of Aberdeen, Aberdeen, UK

*: corresponding author: antoine.stier@gmail.com / amstie@utu.fi

Antoine Stier: <https://orcid.org/0000-0002-5445-5524>

Bin-Yan Hsu: <https://orcid.org/0000-0002-3799-0509>

Coline Marciau: <https://orcid.org/0000-0001-5559-4289>

Blandine Doligez: <https://orcid.org/0000-0003-3015-5022>

Lars Gustafsson: <https://orcid.org/0000-0001-6566-2863>

Pierre Bize <https://orcid.org/0000-0003-0454-2598>

Suvi Ruuskanen: <https://orcid.org/0000-0001-5582-9455>

Keywords

Ageing, mitochondria, telomere length, bird, fetal programming.

Abstract

The underlying mechanisms of the lifelong consequences of prenatal environmental condition on health and ageing remain little understood. Thyroid hormones (THs) are important regulators of embryogenesis, transferred from the mother to the embryo. Since prenatal THs can accelerate early-life development, we hypothesized that this might occur at the expense of resource allocation in somatic maintenance processes leading to premature ageing. Therefore, we investigated the consequences of prenatal THs supplementation on potential hallmarks of ageing in a free-living avian model in which we previously demonstrated that experimentally elevated prenatal THs exposure accelerates early-life growth. Using a cross-sectional sampling, we first report that mitochondrial DNA (mtDNA) copy number and telomere length significantly decrease from early-life to late adulthood, thus suggesting that these two molecular markers could be hallmarks of ageing in our wild bird model. Elevated prenatal THs had no effect on mtDNA copy number but counter-intuitively increased telomere length both soon after birth and at the end of the growth period (equivalent to offsetting *ca.* 4 years of post-growth telomere shortening). These findings suggest that prenatal THs might have a role in setting the 'biological' age at birth, but raise questions about the nature of the evolutionary costs of prenatal exposure to high THs levels.

Introduction

Prenatal environmental conditions can have lifelong consequences on health and ageing, but much remains to be done to uncover the mechanisms linking the pre- and postnatal stages [1]. Thyroid hormones (THs) are master regulators of development, health and ageing [2-4]. They are transferred from the mother to the embryo [4] and thyroid disorders during pregnancy can induce developmental pathologies in humans [5]. There is natural variation in the amount of TH being transferred from the mother to the embryo, and some of this variation is linked to the environmental conditions experienced by the mother (*e.g.* food availability or temperature [4]). This suggests that maternal THs could be a potential signal transferred from the mother to the embryo to adjust offspring phenotype to current environmental cues in order to maximize fitness [4]. While the potential for prenatal THs to mediate adaptive maternal effects remains mostly untested, some of our recent data in wild collared flycatcher (*Ficedula albicollis*) suggest that increasing prenatal THs exposure increases both hatching success and early-life growth [6]. From an evolutionary perspective, such apparent beneficial effects are likely to come with associated costs if there is no increase in resource acquisition. This could occur for instance through a reduction in resource allocation towards somatic maintenance, which could impair subsequent health and accelerate ageing [7]. Yet, we currently lack experimental data on the effects of prenatal THs on postnatal health and ageing.

Avian models offer a great potential to study this question since prenatal conditions can be directly manipulated through hormonal injection in the egg [4]. Yet, one important challenge remains to be able to assess long-term consequences on ageing under an ecologically-realistic scenario. When direct effects on survival and lifespan cannot be estimated, the use of biomarkers that are mirroring age-related health impairments could be useful. Two promising biomarkers are telomere length and mitochondrial DNA (mtDNA) copy number since both markers are frequently reported to decrease with age and to be associated with increased mortality risks [8-10]. There is evidence that telomere length could reflect phenotypic quality since early-life telomere length has been shown for instance to predict lifespan [11] and fitness [12] in avian species. Interestingly, a large part of the inter-individual

variation in telomere length could already be set at birth, and thus may be partially caused by different exposure to maternal hormones [13,14].

In this study, we aim to evaluate whether mtDNA copy number and telomere length decline with age in a free-living bird population of collared flycatchers, and to test the effects of prenatal THs on those potential hallmarks of ageing. To this aim, we first investigated age-related changes in telomere length and mtDNA copy number using cross-sectional data covering most of the lifespan spectrum for this population. While it is established that telomeres shorten with age in most bird species, there is no information to date regarding age-related variation of mtDNA copy number in avian models [15]. Then, using egg-injection of THs, we investigated the effect of prenatal THs on postnatal mtDNA copy number and telomere length. Based on the known stimulation of mitochondrial biogenesis by THs (among other through the up-regulation of PGC1 α/β [16]), we predicted that increasing prenatal THs might increase early-life mtDNA copy number, which could be a cellular pathway supporting the transient growth-enhancing effect we previously demonstrated in this species [6]. Conversely, since THs have been reported in some studies to increase oxidative stress and in our model species to enhance growth (two pathways accelerating telomere shortening [7,17]), we predicted that increasing prenatal THs levels should shorten telomere length at birth, and/or increase early postnatal telomere shortening.

Materials & Methods

This study was conducted in the long-term monitored population of collared flycatchers on Gotland, Sweden. In 2016, 44 randomly selected adult birds (30 females and 14 males) of known-age (1 to 7 years old, *i.e.* cross-sectional data; maximum lifespan = 9.8 years) were sampled across different woodlands. Data and samples from 30 nests of a previous study [6] were used to investigate the effects of a prenatal THs manipulation on telomeres and mtDNA copy number. We used 14 *Control* (vehicle-injected) and 16 *TH* nests in which all eggs were injected with *ca.* a 2SD increase of THs egg content based on natural range, following the procedure described in detail in [6] and ESM1. The original study

[6] demonstrated that THs-injected eggs had a higher hatching success, that early-life survival was not significantly affected by prenatal THs manipulation, and that chicks from THs-eggs were heavier at day 2 and larger at day 8 but did not significantly differ from controls in mass or size at fledging [6]. Two chicks per nest were randomly selected among those surviving to day 12. These chicks were blood sampled twice: soon after hatching (day 2, < 10µL of blood) and at the end of growth (*i.e.* day 12; < 50µL of blood). Relative mtDNA copy number of blood cells has been measured as described in [18] and ESM2. Both relative telomere length (*rTL* measured using qPCR) and absolute telomere length (*TL*, measured using *in-gel* TRF from day 12 samples only) have been measured as described in [19] and ESM2. Two day 2 samples failed to meet DNA quality control criteria, three samples failed to amplify for the mtDNA copy number analysis, and one day 12 DNA sample was excluded from TRF analysis due to insufficient DNA quantity, explaining the observed differences in sample sizes among analyses. Age-related variation in mtDNA copy number and telomere length were tested using Pearson correlation tests, and age-related regression equations have been calculated to infer the amount of telomere length or mtDNA copy number lost per year post-growth (Fig. 1). The effects of prenatal TH elevation and age on mtDNA copy number and telomere length (*rTL* and *TL*) were tested using linear mixed models, with nest identity and bird identity as random effects (to control for multiple birds per nest and multiple samples per bird), as well as age (2 vs. 12 days), treatment (TH vs. Control) and their interaction as fixed effects. mtDNA copy number and *rTL* were z-transformed prior to statistical analysis [20]. Standardized effect sizes (*ES*) for mixed models have been calculated using the emmeans package in R [21]. Non-significant interactions were removed from final models. The effect of sex was tested but excluded from final analyses since it was never significant.

Results and Discussion

We found strong evidence that both mtDNA copy number ($r = -0.57$, $p < 0.001$, Fig. 1A) and telomere length measured either using a relative qPCR method (*rTL*, $r = -0.30$, $p = 0.022$, Fig. 1B) or an absolute *in-gel* quantification (*TL*, $r = -0.48$, $p < 0.001$, Fig. 1C) significantly decreased from growth

completion (*i.e.* day 12) to late-adulthood using a cross-sectional approach. Restricting our analyses to adult individuals only (*i.e.* age ≥ 1) reveals that mtDNA copy number ($r = -0.42$, $p = 0.004$) and absolute telomere length ($r = -0.43$, $p = 0.003$) still significantly decrease with age, while this relationship is not significant anymore for relative telomere length ($r = -0.23$, $p = 0.14$). These findings in our avian model are in accordance with reports from the human literature [9,10]. Since individuals exhibiting short telomeres (both in human and various avian species [8,10]) or low mtDNA copy number (only tested in human to the best of our knowledge [9]) have been shown to disappear earlier from the population, the age-related slopes from our cross-sectional sampling presented in Fig. 1 could underestimate the real age-related decline in both telomere length and mitochondrial density.

We found no significant impact of prenatal THs on mtDNA copy number (TH: $ES = -0.26$ [-1.03; 0.51], $p = 0.46$, ESM3 Table S1A, Fig. 2A), despite a considerable early-life reduction in mtDNA copy number during the growth period (Age: $p < 0.001$, equivalent to the reduction occurring over 3.5 years post-growth based on the age-related regression equation in Fig. 1A). Contrary to our predictions, increasing prenatal THs led to longer telomeres (measured as rTL) soon after hatching (day 2), and this effect was maintained at the end of the growth (day 12) period (TH: $ES = 1.11$ [0.03;2.19], $p = 0.036$, ESM3 Table S1B, Fig. 2B). This is confirmed by the analysis of absolute telomere length (TL) at day 12, showing longer telomeres in birds hatched from THs-injected eggs (TH: $ES = 1.91$ [0.01;3.81], $p = 0.044$, ESM3 Table S1C, Fig. 2C) compared to control eggs. The effect of increasing prenatal THs on telomere length was substantial (*i.e.* large 'biological' effect size), being equivalent to offsetting *ca.* 4.3 years (rTL) and 3.6 years (TL) of post-growth telomere shortening (based on age-related regression equations in Fig. 1B and 1C). It was unfortunately not possible to evaluate any long-lasting impact of prenatal THs supplementation on adult telomere length. Yet, considering that most telomere dynamics occurs during the growth period and that telomere length is a repeatable trait [13,14], it is likely that the effects of prenatal THs observed here on early-life telomere length would be carried over the adult stage.

The positive effect of prenatal THs on telomere length is unlikely related to oxidative stress prevention, since we previously found no differences in oxidative stress markers in these experimental birds [6], or in a similar experiment in a closely-related species [22]. The selective disappearance of THs-embryos with short telomeres could potentially explain why chicks hatched from THs-eggs have longer telomeres than controls. Yet, this seems unlikely to explain our results since we found a higher hatching success of THs-eggs [6]. One previous study in humans reported that the promoter of *hTERT* (the catalytic subunit of the enzyme telomerase, responsible for elongating telomeres) contains a binding site for THs [23]. Consequently, one hypothesis would be that prenatal THs could elongate telomeres early in life through the activation of the telomerase enzyme. Yet, we are currently lacking both *in vitro* and *in vivo* studies testing such an hypothesis.

While an evolutionary trade-off is generally expected to occur between fast growth and telomere shortening [7], prenatal THs supplementation both enhances early-life growth [6] and leads to longer early-life telomeres in our study system. Egg THs could be a potential signal transferred from the mother to the embryo to adjust offspring phenotype to the expected environmental conditions [4]. Yet, if raising egg THs levels only had benefits, natural selection would have favoured higher maternal transfer of THs into eggs, and/or offspring maximizing growth and telomere length without such a maternal signal. Consequently, it is likely that high egg THs levels have costs that we have not been able to detect here (*e.g.* reduced immunity) or that could only become visible later in life (*e.g.* altered reproductive potential), which clearly deserves further investigation. Importantly, we can expect the potential cost-benefit balance of low or high egg THs levels to depend on environmental conditions [4]. Such potential context-dependent effects will require experimental manipulations of both egg THs and environmental conditions to be revealed (*e.g.* [22]).

While the mechanisms remain to be identified, our study demonstrates that prenatal THs levels have the potential to modulate telomere length in early-life, and thus to influence the 'biological' age at birth. It has previously been shown that prenatal exposure to glucocorticoids could shorten telomeres [24], but our study is the first to show that telomere length at birth could be increased by

modulating the prenatal hormonal environment. Thyroid function is known to influence cardiovascular disease risk and life expectancy in adult humans [3], but no information is currently available regarding the impact of prenatal THs exposure on adult health and lifespan. Epidemiological and long-term experimental studies investigating the impact of prenatal THs on adult phenotype, health and lifespan are now required to establish if the effect observed here on early-life telomere length could be translated into a longevity gain, but also to identify potential costs of elevated prenatal exposure to THs.

Acknowledgements

We are grateful to many students and Szymek Drobnik for their help in the field, to three anonymous reviewers for their constructive feedback and to Pat Monaghan for providing access to TRF facilities.

Funding

The project was funded by a Marie Skłodowska-Curie Postdoctoral Fellowship (#658085) and a 'Turku Collegium for Science and Medicine' Fellowship to AS, and an Academy of Finland grant (# 286278) to SR.

References

1. Preston, J. D., Reynolds, L. J. & Pearson, K. J. 2018 Developmental Origins of Health Span and Life Span: A Mini-Review. *Gerontology* **64**, 237–245. (doi:10.1159/000485506)
2. Boelaert, K. & Franklyn, J. A. 2005 Thyroid hormone in health and disease. *Journal of Endocrinology* **187**, 1–15. (doi:10.1677/joe.1.06089)
3. Bano, A., Dhana, K., Chaker, L., Kavousi, M., Ikram, M. A., Mattace-Raso, F. U. S., Peeters, R. P. & Franco, O. H. 2017 Association of Thyroid Function With Life Expectancy With and Without Cardiovascular Disease. *JAMA Intern Med* **177**, 1650–8. (doi:10.1001/jamainternmed.2017.4836)
4. Ruuskanen, S. & Hsu, B.-Y. 2018 Maternal Thyroid Hormones: An Unexplored Mechanism Underlying Maternal Effects in an Ecological Framework. *Physiol Biochem Zool* **91**, 904–916. (doi:10.1086/697380)
5. Delitala, A. P., Capobianco, G., Cherchi, P. L., Dessole, S. & Delitala, G. 2019 Thyroid function and thyroid disorders during pregnancy: a review and care pathway. *Archives of Gynecology and Obstetrics* **299**, 327–338. (doi:10.1007/s00404-018-5018-8)
6. Hsu, B.-Y., Doligez, B., Gustafsson, L. & Ruuskanen, S. 2019 Transient growth-enhancing effects of elevated maternal thyroid hormones at no apparent oxidative cost during early postnatal period. *Journal of Avian Biology* **50**, 4692–10. (doi:10.1111/jav.01919)
7. Monaghan, P. & Ozanne, S. E. 2018 Somatic growth and telomere dynamics in vertebrates: relationships, mechanisms and consequences. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **373**, 20160446–15. (doi:10.1098/rstb.2016.0446)
8. Wilbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H. & Boonekamp, J. J. 2018 The relationship between telomere length and mortality risk in non-model vertebrate systems: a meta-analysis. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **373**, 20160447–9. (doi:10.1098/rstb.2016.0447)
9. Mengel-From, J., Thinggaard, M., Dalgård, C., Kyvik, K. O., Christensen, K. & Christiansen, L. 2014 Mitochondrial DNA copy number in peripheral blood cells declines with age and is associated with general health among elderly. **133**, 1149–1159. (doi:10.1007/s00439-014-1458-9)
10. Arbee, K. G. et al. 2020 Association of Leukocyte Telomere Length With Mortality Among Adult Participants in 3 Longitudinal Studies. *JAMA Netw Open* **3**, e200023–12. (doi:10.1001/jamanetworkopen.2020.0023)
11. Heidinger, B. J., Blount, J. D., Boner, W., Griffiths, K., Metcalfe, N. B. & Monaghan, P. 2012 Telomere length in early life predicts lifespan. *Proceedings of the National Academy of Sciences* **109**, 1743–1748. (doi:10.1073/pnas.1113306109)
12. Eastwood, J. R., Hall, M. L., Teunissen, N., Kingma, S. A., Hidalgo Aranzamendi, N., Fan, M., Roast, M., Verhulst, S. & Peters, A. 2019 Early-life telomere length predicts lifespan and lifetime reproductive success in a wild bird. *Mol Ecol* **28**, 1127–1137. (doi:10.1111/mec.15002)
13. Entringer, S., de Punder, K., Buss, C. & Wadhwa, P. D. 2018 The fetal programming of telomere biology hypothesis: an update. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **373**, 20170151–15. (doi:10.1098/rstb.2017.0151)
14. Stier, A., Metcalfe, N. B. & Monaghan, P. 2020 Pace and stability of embryonic development affect telomere dynamics: an experimental study in a precocial bird model. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **287**, 20201378–9. (doi:10.1098/rspb.2020.1378)
15. Stier, A., Reichert, S., Criscuolo, F. & Bize, P. 2015 Red blood cells open promising avenues for

- 232 longitudinal studies of ageing in laboratory, non-model and wild animals. *Experimental Gerontology* **71**,
233 118–134. (doi:10.1016/j.exger.2015.09.001)
- 234 16. Weitzel, J. M. & Iwen, K. A. 2011 Coordination of mitochondrial biogenesis by thyroid hormone.
235 *Molecular and Cellular Endocrinology* **342**, 1–7. (doi:10.1016/j.mce.2011.05.009)
- 236 17. Reichert, S. & Stier, A. 2017 Does oxidative stress shorten telomeres in vivo? A review. *Biol Letters* **13**,
237 20170463–7. (doi:10.1098/rsbl.2017.0463)
- 238 18. Stier, A., Bize, P., Hsu, B.-Y. & Ruuskanen, S. 2019 Plastic but repeatable: rapid adjustments of
239 mitochondrial function and density during reproduction in a wild bird species. *Biol Letters* **15**,
240 20190536. (doi:10.1098/rsbl.2019.0536)
- 241 19. Kärkkäinen, T., Teerikorpi, P., Panda, B., Helle, S., Stier, A. & Laaksonen, T. 2019 Impact of continuous
242 predator threat on telomere dynamics in parent and nestling pied flycatchers. *Oecologia* **191**, 757–766.
243 (doi:10.1007/s00442-019-04529-3)
- 244 20. Verhulst, S. 2019 Improving comparability between qPCR-based telomere studies. *Mol Ecol Resour* **20**,
245 11–13. (doi:10.1111/1755-0998.13114)
- 246 21. Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. 2018 Emmeans: Estimated marginal means,
247 aka least-squares means. *R package version 1.3.4* (doi:10.1080/00031305.1980.10483031>)
- 248 22. Hsu, B.-Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A. & Ruuskanen, S. 2020 Testing for
249 context-dependent effects of maternal thyroid hormones on offspring survival and physiology: an
250 experimental approach manipulating temperature in a wild bird species. *Scientific Report* **10**, 14563.
251 (doi:10.1038/s41598-020-71511-y)
- 252 23. Cong, Y. S., Wen, J. & Bacchetti, S. 1999 The human telomerase catalytic subunit hTERT: organization of
253 the gene and characterization of the promoter. *Human Molecular Genetics* **8**, 137–142.
254 (doi:10.1093/hmg/8.1.137)
- 255 24. Haussmann, M. F., Longenecker, A. S., Marchetto, N. M., Juliano, S. A. & Bowden, R. M. 2012 Embryonic
256 exposure to corticosterone modifies the juvenile stress response, oxidative stress and telomere length.
257 *Proceedings of the Royal Society B: Biological Sciences* **279**, 1447–1456. (doi:10.1098/rspb.2011.1913)
- 258 25. Stier A., Hsu B.-Y., Marciau C., Doligez B., Gustafsson L., Bize P. & Ruuskanen S. 2020. Data from: Born
259 to be young? Prenatal thyroid hormones increase telomere length in wild collared flycatchers.
260 *FigShare*. (<https://doi.org/10.6084/m9.figshare.11688504.v1>)

261

Figure captions

Fig. 1: Age-related variation in potential hallmarks of ageing in wild collared flycatchers: (A) decrease in relative mtDNA copy number, (B) decrease in relative telomere length measured with qPCR, and **(C) decrease in absolute telomere length** measured with *in-gel* TRF. Data is cross-sectional, adult birds were of known-age and chicks (*i.e.* age 0) were 12 days old (from control group only, 1 chick per nest). *rTL* and *mtDNA copy number* have been z-transformed, Pearson's coefficients of correlation *r* and associated *p*-values are presented along with the age-related regression equations. Regression lines are plotted \pm 95% C.I., *N* = 58 (44 adults, 14 nestlings). See also the main text for statistical analyses restricted to adult individuals.

Fig. 2: Effects of experimental prenatal thyroid hormone elevation on: (A) early-life dynamics of mtDNA copy number, (B) early-life dynamics of relative telomere length, and (C) absolute telomere length at the end of growth period (day 12). *rTL* and *mtDNA copy number* have been z-transformed. Means are plotted as symbols \pm SE, individual responses are plotted as solid lines, and *p*-values are indicated within each panel. See ESM3 Table S1 for details on statistics.




Fig. 1: Age-related variation in potential hallmarks of ageing in wild collared flycatchers: (A) decrease in relative mtDNA copy number, (B) decrease in relative telomere length measured with qPCR, and (C) decrease in absolute telomere length measured with in-gel TRF. Data is cross-sectional, adult birds were of known-age and chicks (i.e. age 0) were 12 days old (from control group only, 1 chick per nest). rTL and mtDNA copy number have been z-transformed, Pearson's coefficients of correlation r and associated p -values are presented along with the age-related regression equations. Regression lines are plotted \pm 95% C.I., $N = 58$ (44 adults, 14 nestlings). See also the main text for statistical analyses restricted to adult individuals.

240x62mm (300 x 300 DPI)




Fig. 2: Effects of experimental prenatal thyroid hormone elevation on: (A) early-life dynamics of mtDNA copy number, (B) early-life dynamics of relative telomere length, and (C) absolute telomere length at the end of growth period (day 12). rTL and mtDNA copy number have been z-transformed. Means are plotted as symbols \pm SE, individual responses are plotted as solid lines, and p-values are indicated within each panel. See ESM3 Table S1 for details on statistics.

261x65mm (300 x 300 DPI)