



Body size and digestive system shape resource selection by ungulates: A cross-taxa test of the forage maturation hypothesis

Saeideh Esmaeili, Brett Jesmer, Shannon Albeke, Ellen Aikens, Kathryn Schoenecker, Sarah King, Briana Abrahms, Bayarbaatar Buuveibaatar, Jeffrey Beck, Randall Boone, et al.

► To cite this version:

Saeideh Esmaeili, Brett Jesmer, Shannon Albeke, Ellen Aikens, Kathryn Schoenecker, et al.. Body size and digestive system shape resource selection by ungulates: A cross-taxa test of the forage maturation hypothesis. *Ecology Letters*, 2021, 24 (10), pp.2178-2191. 10.1111/ele.13848 . hal-03425900

HAL Id: hal-03425900

<https://hal.science/hal-03425900>

Submitted on 11 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Body size and digestive system shape resource selection by ungulates: a cross-taxa test of the Forage Maturation Hypothesis

Saeideh Esmaeili^{1,*}, Brett R. Jesmer^{2,3,4}, Shannon E. Albeke⁵, Ellen O. Aikens¹, Sarah R. B. King⁶, Kathryn A. Schoenecker^{6,7}, Briana Abrahms⁸, Bayarbaatar Buuveibaatar⁹, Jeffrey L. Beck¹⁰, Randall B. Boone¹¹, Francesca Cagnacci¹², Simon Chamaillé-Jammes^{13,14}, Buyanaa Chimeddorj¹⁵, Paul C. Cross¹⁶, Nandintsetseg Dejid¹⁷, Jagdag Enkhbyar¹⁸, Ilya R. Fischhoff¹⁹, Adam T. Ford²⁰, Kate Jenks²¹, Mahmoud-Reza Hemami²², Jacob D. Hennig¹⁰, Takehiko Y. Ito^{23,24}, Petra Kaczensky^{25,26,27}, Matthew J. Kauffman^{1,28}, John D.C. Linnell^{25,29}, Badamjav Lkhagvasuren³⁰, John F. McEvoy³¹, Joerg Melzheimer³², Jerod A. Merkle¹, Thomas Mueller^{17,33}, Jeff Muntifering^{21,34}, Atle Mysterud³⁵, Kirk A. Olson⁹, Manuela Panzacchi³⁶, John C. Payne^{9,36}, Luca Pedrotti³⁷, Geir R. Rauset³⁶, Daniel I. Rubenstein³⁸, Hall Sawyer³⁹, John. D. Scasta¹⁰, Johannes Signer⁴⁰, Melissa Songer³¹, Jared A. Stabach³¹, Seth Stapleton²¹, Olav Strand³⁶, Siva R. Sundaresan⁴¹, Dorj Usukhjargal¹⁸, Ganbold Uuganbayar¹⁸, John M. Fryxell⁴², and Jacob R. Goheen¹.

¹ Department of Zoology and Physiology and Program In Ecology, University of Wyoming, Laramie, USA.

² Department of Fish and Wildlife Conservation, Virginia Tech, Blacksburg, VA 24061, USA.

³ Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06520, USA.

⁴ Center for Biodiversity and Global Change, Yale University, New Haven, CT 06520, USA.

⁵ Wyoming Geographic Information Science Center, University of Wyoming, Laramie WY, USA.

⁶ Natural Resource Ecology Laboratory, Warner College of Natural Resources, Colorado State University, Fort Collins, CO, USA.

⁷ US Geological Survey, Fort Collins Science Center, Fort Collins, CO, USA.

⁸ Center for Ecosystem Sentinels, Department of Biology, University of Washington, Seattle, WA, USA.

⁹ Wildlife Conservation Society, Mongolia Program, Ulaanbaatar, Mongolia.

¹⁰ Department of Ecosystem Science and Management, University of Wyoming, Laramie WY, USA.

¹¹ Department of Ecosystem Science and Sustainability and the Natural Resources Ecology Laboratory, Colorado State University, Fort Collins, CO, USA.

¹² Department of Biodiversity and Molecular Ecology, Research and Innovation Centre, Fondazione Edmund Mach, 38010 San Michele all'Adige, Italy.

¹³ CEFE, Univ. Montpellier, CNRS, EPHE, IRD, Université Paul Valéry Montpellier 3, Montpellier, France.

¹⁴ Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

¹⁵ Mongolia Program Office, World Wide Fund for Nature, Ulaanbaatar, Mongolia.

¹⁶ U.S. Geological Survey, Northern Rocky Mountain Science Center, Bozeman, MT, USA.

¹⁷ Senckenberg Biodiversity and Climate Research Centre, Frankfurt (Main), Germany.

¹⁸ Hustai National Park Trust, Mongolia.

¹⁹ Cary Institute of Ecosystem Studies, Millbrook, NY, USA.

²⁰ Department of Biology, the University of British Columbia, Okanagan, BC, Canada.

²¹ Minnesota Zoo, 13000 Zoo Boulevard, Apple Valley, MN 55124, USA.

²² Department of Natural Resources, Isfahan University of Technology, Isfahan, Iran.

²³ Arid Land Research Center, Tottori University, Tottori, Japan.

²⁴ International Platform for Dryland Research and Education, Tottori University, Tottori, Japan.

²⁵ Department of Terrestrial Biodiversity, Norwegian Institute for Nature Research, Trondheim, Norway.

²⁶ Research Institute of Wildlife Ecology, University of Veterinary Sciences, Vienna, Austria.

²⁷ Inland Norway University of Applied Sciences, Campus Evenstad, Norway.

²⁸ U.S Geological Survey, Wyoming Cooperative Fish and Wildlife Research Unit, Laramie, WY, USA.

²⁹ Department of Forestry and Wildlife Management, Inland Norway University of Applied Sciences, NO-2480 Koppang, Norway.

³⁰ Institute of General and Experimental Biology, Mongolian Academy of Sciences, Ulaanbaatar, Mongolia, Deceased August 1, 2019.

³¹ Conservation Ecology Center, Smithsonian Conservation Biology Institute, National Zoological Park, Front Royal, VA, USA.

52 32 Department of Evolutionary Ecology, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany.
53 33 Department of Biological Sciences, Goethe University, Frankfurt (Main), Germany.
54 34 Namibia University of Science and Technology, Windhoek, Namibia.
55 35 Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo,
56 Norway.
57 36 Department of Terrestrial ecology, Norwegian Institute for Nature Research, Trondheim, Norway.
58 37 Stelvio-Stilfserjoch National Park, Bormio, SO, Italy.
59 38 Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA.
60 39 Western Ecosystems Technology, Inc., Laramie, WY, USA.
61 40 Wildlife Sciences, Faculty of Forest Sciences, University of Göttingen, Göttingen, Germany.
62 41 Greater Yellowstone Coalition, Bozeman, MT, USA.
63 42 Department of Integrative Biology, University of Guelph, Guelph, ON, Canada.
64 * Corresponding author: saeideh.esmaeili@colostate.edu

65 **Running title:** A cross-taxa test of the Forage Maturation Hypothesis

66 **Keywords:** Equidae, forage biomass, hindgut fermentation, ruminant, step-selection function,
67 water requirements.

68 **Type of article:** Letters

69 **Number of words in the abstract:** 142

70 **Number of words in the main text:** 5,293

71 **Number of references:** 99

72 **Number of figures, tables, and text boxes:** 3 figures

73 **Corresponding author:** Saeideh Esmaeili

74 Department of Zoology and Physiology

75 University of Wyoming, 1000 E. University Ave.

76 Laramie, WY 82071

77 **Email:** saeideh.esmaeili@colostate.edu

78 **Statement of Authorship:** SE and JRG developed the research idea and designed the study. SE
79 compiled, managed, and analyzed the data (and collected GPS locations for onagers in southern
80 Iran), and drafted the manuscript. BRJ, SEA, and EOA provided analytical assistance, and BRJ
81 provided feedback on the methods. JRG and BRJ revised the drafts. All coauthors collected data,
82 contributed to revisions, and gave final approval for publication.

Data accessibility statement: The GPS relocation data associated with the current study have not been deposited in a public repository because they are part of ongoing investigations. Please contact owners of the datasets for requests.

Abstract

The Forage Maturation Hypothesis (FMH) states that energy intake for ungulates is maximized when forage biomass is at intermediate levels. Nevertheless, metabolic allometry and different digestive systems suggest that resource selection should vary across ungulate species. By combining GPS relocations with remotely-sensed data on forage characteristics and surface water, we quantified the effect of body size and digestive system in determining movements of 30 populations of hindgut fermenters (equids) and ruminants across biomes. Selection for intermediate forage biomass was negatively related to body size, regardless of digestive system. Selection for proximity to surface water was stronger for equids relative to ruminants, regardless of body size. To be more generalizable, we suggest that the FMH explicitly incorporate contingencies in body size and digestive system, with small-bodied ruminants selecting more strongly for potential energy intake, and hindgut fermenters selecting more strongly for surface water.

Introduction

Animal movements should be influenced by the need to maintain a positive balance between energetic gains and losses (Senft *et al.* 1987; Nathan *et al.* 2008; Owen-Smith *et al.* 2010; Middleton *et al.* 2013). Foraging behaviors employed by ungulates vary according to body size and digestive system (ruminant versus hindgut fermenters; Bell 1971; Demment & Van Soest 1985; Olff *et al.* 2002; Hopcraft *et al.* 2012). Smaller-bodied ungulates (i.e., ruminants, primarily) should forage in patches with highly digestible forage (high energy and nutrient concentrations) because they possess high mass-specific metabolic rates. In contrast, larger-bodied ungulates (i.e., both larger-bodied ruminants and hindgut fermenters) should forage in high biomass patches because of higher absolute energy demands (Jarman 1974; McNab 1974; Illius & Gordon 1992; Barboza & Bowyer 2000). Consequently, body size creates trade-offs in how ungulates should prioritize forage biomass and forage digestibility (Bailey *et al.* 1996; Wilmshurst *et al.* 2000; Olff *et al.* 2002).

For nearly 30 years, the Forage Maturation Hypothesis (hereafter “FMH”; Fryxell 1991) has provided a lens for understanding resource selection and movements of free-ranging ungulates (Fryxell 1991; Hebblewhite *et al.* 2008; Merkle *et al.* 2016). Digestibility of forage is highest at the beginning of the growing season because plants lack structural fiber needed to support high biomass (Van Soest 1996). At this earliest phenological stage, however, plants are small and biomass is limiting, so energy intake of ungulates may be limited by bite size (Spalinger & Hobbs 1992; Wilmshurst *et al.* 2000; Hebblewhite *et al.* 2008). Digestibility then declines as plants mature, gain biomass, and become more fibrous (Van Soest 1996). Because of this phenological trade-off between biomass and digestibility of plants, and according to the FMH, energy intake is maximized for ungulates at intermediate phenological stages (i.e., when

rates of both forage intake and digestibility are at intermediate levels; Fig 1A). Although the FMH was originally developed to explain resource selection in tropical, bulk-grazing ruminants, its predictions are relevant for temperate ruminants that display a diversity of feeding strategies (e.g., unselective grazing, selective browsing, mixed feeding; Hofmann 1989; Albon & Langvatn 1992; Mueller *et al.* 2008; Middleton *et al.* 2018). As a result, the FMH frequently forms the basis for conceptualizing resource selection and movements of ungulates—including large- and small-bodied species, and hindgut and ruminant fermenters—across the globe (e.g., Drescher *et al.* 2006; Edouard *et al.* 2010; Fleurance *et al.* 2010; St-Louis & Cote 2014).

The central prediction of the FMH—that ungulates should select forage of intermediate biomass and intermediate digestibility to maximize energy intake—should hinge largely on body size. In 1971, Bell first articulated how metabolic rate should interact with resource selection by free-ranging ungulates, stating that relative requirements (i.e., energy and protein per unit weight, per unit time) are higher for smaller ungulates, while absolute requirements (i.e., energy and protein per individual, per unit time) are higher for larger ungulates (Bell 1971). Over two decades ago, Wilmschurst *et al.* (2000) extended the work of Bell (1971) and Fryxell (1991) by testing quantitative predictions for selection of optimal biomass by five co-occurring ruminants in Serengeti National Park. Wilmschurst *et al.* (2000) predicted that optimal biomass of plants selected by large herbivores would vary as a function of body size, at least among ruminants. Specifically, resource selection should occur along a gradient from high digestibility/low biomass to low digestibility/high biomass, dependent on body size. Explicitly considering and testing for this contingency among a diversity of ungulates would refine the predictive capacity of the FMH, thereby advancing its application to ungulates more generally (Fig. 1A, Fig. 1B).

In addition to forage biomass and energy content, reliance on surface water restricts the movement and distribution of ungulates across some landscapes (Rozen-Rechels *et al.* 2019; Veldhuis *et al.* 2019). Ruminants efficiently extract protein from fresh plant tissue because the fermentation site (rumen) occurs anterior to the absorption site (small intestine) for proteins. Ruminants therefore employ an “efficient” urea cycle in which available nitrogen is recycled throughout the digestive tract, thereby conserving water (Janis 1976; Santos *et al.* 2011; Owens & Basalan 2016). In contrast, hindgut fermenters have reduced retention time, but lower rates of protein extraction (Janis 1976; Duncan *et al.* 1990; Van Soest 1996). Consequently, hindgut fermenters excrete more urea, which must be accompanied by water. Larger-bodied ungulates—and larger-bodied hindgut fermenters in particular—consume large amounts of biomass and should therefore be further tied to surface water (i.e., water available for drinking, as opposed to contained within plants) because high-biomass forage tends to be drier (Bell 1971; Redfern *et al.* 2003; Schoenecker *et al.* 2016). Therefore, constraints associated with conserving water in the body should tie hindgut fermenters to water, and potentially result in deviations from the central prediction of the FMH (which is based purely on biomass and digestibility of forage). In sum, natural selection should favor ungulate movements that optimize net energy gain despite constraints imposed by forage biomass, the energy contained within forage, and surface water, with the relative importance of each arising as a function of body size and digestive system (Redfern *et al.* 2003; Fig. 1B, 1C).

We tested the central prediction of the FMH by linking high-resolution movement trajectories (unavailable when the FMH was conceptualized) with selection of forage biomass, potential energy intake, and surface water by free-ranging ungulates. We evaluated the relative influence of body size and digestive system on selection for these resources by ruminants

(foregut fermenters) and equids (Family Equidae, representing hindgut fermenters) using GPS telemetry data from 30 populations of 19 species distributed across the biomes. We predicted that (1) smaller-bodied ungulates (all of which exhibit ruminant fermentation) would select for resource patches that maximize energy intake, thereby conforming to the central prediction of the FMH (Fig. 1A); whereas (2) by virtue of their larger body size, resource selection patterns of both larger-bodied ruminants and equids would deviate from this prediction (Fig. 1B). Specifically, we expected that larger-bodied ruminants would select most strongly for patches of high forage biomass, whereas equids (because of their hindgut fermentation) would select most strongly for patches close to water.

Methods

Data Collection

We compiled a global data set of GPS locations for 11 populations of equids and 19 populations of ruminants totaling 580 individuals; data for all study populations were collected between 2005 and 2019 (Figure 2A; Table S1). The equid data set comprised GPS relocations for seven (out of nine) extant species of wild and feral equids: Asiatic wild ass (khulan, *Equus hemionus hemionus* and onager, *E. h. onager*), feral burro (*E. asinus*), feral horse (*E. caballus*), Grevy's zebra (*E. grevyi*), mountain zebra (*E. zebra*), plains zebra (*E. quagga*), and Przewalski's horse (*E. ferus*). The ruminant data set included GPS relocations for 12 species: African buffalo (*Syncerus caffer*), elk (*Cervus canadensis*), goitered gazelle (*Gazella subgutturosa*), impala (*Aepyceros melampus*), Mongolian gazelle (*Procapra gutturosa*), mule deer (*Odocoileus hemionus*), red deer (*Cervus elaphus*), reindeer (*Rangifer tarandus tarandus*), roe deer (*Capreolus capreolus*), saiga antelope (*Saiga tatarica*), springbok (*Antidorcas marsupialis*), and white-bearded wildebeest

(*Connochaetes taurinus*). Equid species ranged in body size from 180 kg (feral burro) to 430 kg (feral horse; Wilson & Mittermeier 2011); ruminant species ranged in body size from 24 kg (roe deer) to 592 kg (African buffalo; Wilson & Mittermeier 2011). Hereafter, we distinguish between “study areas” (the geographic locales in which resource selection were quantified) and “study populations” (individuals fitted with GPS collars within a study area; Table S1). Five of our study areas contained multiple study populations, and eight species were represented by two or more study populations spread across different study areas (Table S1). Inaccurate GPS locations were removed either by coauthors or by excluding locations with dilution of precision > 10.

We restricted our analyses to habitats with minimal canopy cover, including rangelands, savannas, open forest, and tundra. We intended to include sufficiently open canopies for remotely-sensed vegetation indices to work properly. Therefore, we applied two criteria in selecting study populations. The first criterion required that a study area be comprised of <20% “dense forest”, as classified by the Food and Agriculture Organization (FAO; Friedl & Sulla-Menashe 2015). The second criterion required that percent (%) tree cover across the entire study area was <40% (Hansen *et al.* 2013; Table S1). We calculated percent of the study area covered with dense forest and mean % tree cover within minimum convex polygons (MCP) constructed for each study population using the Terra and Aqua combined Moderate Resolution Imaging Spectroradiometer (MODIS) Land Cover Type, MCD12Q1, Version 6, layer 7, spatial resolution of 500 meters and temporal resolution of a year (Friedl & Sulla-Menashe 2015) and Global 2010 Tree Cover (spatial resolution 30 meters, Hansen *et al.* 2013), respectively.

Forage Biomass and Potential Energy Intake

To delineate study areas, we computed the MCP around each population's GPS locations and buffered the MCP by 5 km to ensure that we captured adjacent habitats that were available to telemetered individuals. Within each study area, we extracted Modified Soil-Adjusted Vegetation Index (MSAVI; MODIS terra satellite imagery Version 6.0 MOD09Q1; spatial resolution 250 x 250 m, temporal resolution eight days) using the MSAVI2 method and equations described in Qi *et al.* (1994) as a proxy of forage biomass (Pettorelli *et al.* 2005; Borowik *et al.* 2013). In arid environments, MSAVI and other soil-adjusted indices of vegetation are more appropriate than the more commonly used Normalized Difference Vegetation Index (NDVI) because they minimize the influence of bare ground on estimates of vegetation (Qi *et al.* 1994). Additionally, NDVI and MSAVI are extracted from similar spectral bands, and usually are strongly correlated in temperate conditions (Jin *et al.* 2014; Liang *et al.* 2016). We removed pixels categorized as snow, cloud, or shadow using quality assessment bands, then smoothed each time-series of MSAVI using a moving three-scene median filter and applied a linear interpolation (Branco *et al.* 2019).

To estimate the instantaneous rate of green-up (IRG), we calculated the rate of change in MSAVI for every three consecutive dates by using a three-scene moving window (Avgar *et al.* 2013; Branco *et al.* 2019). The IRG is a metric that combines both forage biomass and forage digestibility, which collectively equate to potential energy intake (i.e., a proxy of energy that could be effectively available to the feeder, Bischof *et al.* 2012; Avgar *et al.* 2013; Geremia *et al.* 2019). The IRG is positively correlated with the peak in fecal crude protein in ungulates (Hamel *et al.* 2009) and has been used widely as an index of the energy contained in forage across space and time (Merkle *et al.* 2016; Rivrud *et al.* 2016; Branco *et al.* 2019); days from peak IRG is strongly correlated with N:C ratios in grasses (Geremia *et al.* 2019). Hereafter, we refer to

selection for IRG as selection for “potential energy intake”. Similarly, and hereafter, we refer to selection for peak biomass of forage as simply selection for “forage biomass”. Although IRG combines both forage biomass and forage digestibility, it is not redundant with metrics that solely represent forage biomass: while high IRG values represent intermediate plant biomass, low values of IRG may represent either low or high biomass (see Fig. 1A). We therefore used both IRG and MSAVI to disentangle selection for potential energy intake and forage biomass. We normalized both MSAVI and IRG values between 0 and 1 based on the lowest and highest value of each pixel in a year, respectively. Thus, for each pixel, an MSAVI value of 1 represented the highest biomass and a value of 0 represented the lowest biomass in a given year, for a given study population. Similarly, an IRG value of 1 represents forage at a state of intermediate biomass (and the peak rate of green-up), whereas an IRG value of 0 represents forage at a low rate of change. Collectively, these two layers therefore represent metrics of plant phenology across space and time (see also Bischof *et al.* 2012; Merkle *et al.* 2016; Branco *et al.* 2019).

To evaluate how body size and digestive system influenced resource selection, we temporally constrained our analysis times when plants were actively growing (to ensure positive IRG was available to the animals). We determined the duration of growing seasons in each study population by randomly generating 10,000 points within each study area, extracting absolute (non-normalized) MSAVI and IRG values associated with each of the random points, and plotting annual MSAVI and IRG profiles (Fig. S1). For study areas with a single “growing season” per year (25 out of 30 study populations), we defined the beginning of the “growing season” as the Julian day when IRG became positive for three consecutive scenes, and the end of the “growing season” as the Julian day when IRG reached the minimum negative point, followed

by IRG values less than or equal to zero (Fig. S1A; *sensu* Jesmer *et al.* 2018). For study areas with multiple growing seasons per year, we attempted to define each “growing season” consistent with the method described above (Fig. S1B). We then filtered our data set to only those relocations that occurred during growing seasons (Table S1).

Distance to Surface Water

Fine-scale data on surface water is challenging to acquire via remote sensing due to computational restrictions (Redfern *et al.* 2003; Pekel *et al.* 2016; Huang *et al.* 2018). Further, globally-collected precipitation data may underestimate water available to wildlife, given artificial water sources (Ogutu *et al.* 2010). We treated the distribution of surface water as fixed throughout each growing season. We used the following data to identify the occurrence of surface water:

1-Remotely-sensed data on surface water: we used monthly data from the Global Surface Water Explorer (Pekel *et al.* 2016) to estimate occurrence of surface water during the growing season for each study area. With a spatial resolution of 30 x 30 m and temporal resolution of one month, the Global Surface Water Explorer constitutes the most precise data on the distribution of surface water (Pekel *et al.* 2016). For each study area and growing season, we extracted monthly time series of pixels, where each pixel was assigned a 1 or 0, indicating presence versus absence of surface water. We then merged monthly layers into a single layer of surface water.

2-Locally-collected data on surface water: to increase accuracy of our estimates of surface water for each study population, we compiled data on springs, streams, small ponds, and man-made water sources (i.e., surface water sources <30 x 30 m that were undetectable using the Global Water Surface Explorer; Text S1 and Table S2). We merged the locations of locally-collected

surface water with the layer on remotely-sensed data on surface water. We then generated a distance to surface water layer by calculating linear distances between centroids of pixels classified as “surface water” and centroids of non-surface water pixels. To ensure that our distance to surface water layer occurred at a comparable scale with MSAVI and IRG, we normalized values for distance to surface water between 0 and 1 for each study area based on the lowest and highest value of pixels in each growing season, respectively.

Statistical Analyses

We built step-selection functions (Thurfjell *et al.* 2014; Avgar *et al.* 2016) to quantify how spatiotemporal patterns of forage biomass, potential energy intake, and surface water influenced movements by ungulates in each study population during growing seasons. To meet the assumption that the speed of successive steps was uncorrelated (since the step-selection method assumes Brownian motion), we used the R package CTMM (Calabrese *et al.* 2016) to estimate the average time at which 99% of the correlation in speed between successive steps had decayed (i.e., about 3 times of τ_V ; Fleming *et al.* 2014) in each population. We then rarified (i.e., temporally subsampled) the GPS locations of each population, which resulted in uncorrelated, successive steps (Table S1; Fleming *et al.* 2014; Dupke *et al.* 2017). For each observed (used) step, we generated 100 potential (unused) steps by fitting a gamma and von Mises distribution to the step lengths and turning angles, respectively (Signer *et al.* 2019), then compared observed and potential steps using conditional logistic regression (Fortin *et al.* 2005). In addition to step length (standardized to a mean of 0.0 and standard deviation of 1.0), we extracted values of forage biomass, potential energy intake, and distance to surface water at the end points of each observed and potential step. We used conditional logistic regression to estimate selection

coefficients, with each stratum consisting of an observed step and its associated 100 potential steps, and each individual as an independent cluster in fitting a separate model for each study population (Roever *et al.* 2010; Merkle *et al.* 2016; Prima *et al.* 2017). Correlation among the independent variables was minimal (mean r between MSAVI and IRG = -0.09, range = -0.34 to 0.20; mean r between MSAVI and distance to surface water = 0.02, range = -0.17 to 0.55; mean r between IRG and distance to water = 0.008, range = -0.28 to 0.22; Table S3). Since layers for forage biomass and potential energy intake encompassed both spatial and temporal variation across landscapes, but surface water layers encompassed only spatial variation across landscapes, we did not include interactions between forage biomass and water, or between potential energy intake and water in our analyses. Further, we did not find a significant correlation between selection for forage biomass and availability of surface water (i.e., mean distance to surface water; left panel of Figure S3), nor did we find a significant correlation between selection for potential energy intake and mean distance to surface water (right panel of Figure S3). Sample sizes (numbers of individuals telemetered, and the range in numbers of individuals telemetered for a given year) are presented for each study population in Table S1. We interpreted that significant, positive selection for IRG was indicative of movements consistent with the FMH (Merkle *et al.* 2016; Aikens *et al.* 2017). We considered variables significant in the models when 95% confidence intervals around parameter estimates did not encompass zero. We performed these analyses using packages “amt” and “survival” in Program R (Therneau & Lumley 2015; R Core Team 2019; Signer *et al.* 2019).

We next tested the effect of body size and digestive system (i.e., equids or hindgut fermenters vs. ruminants or foregut fermenters) on resource selection across populations using weighted least square regressions and analysis of covariance (ANCOVA). Because resource

selection may intensify when a particular resource is scarce, variable, or both (i.e., exhibit functional responses in habitat selection [Mysterud and Ims 1998; Holbrook *et al.* 2019]), we performed pairwise correlations between population-level selection coefficients for each resource and the (1) mean value of MSAVI, IRG, and distance to surface water; and (2) the coefficient of variation (CV) of MSAVI, IRG, and distance to surface water. We did not find strong support for functional responses in resource selection (i.e., the effect of resource availability and variability on resource selection; Fig. S2), so we attributed variation in selection coefficients to some combination of body size and digestive system. For each of the above tests, we used the parameter estimates derived from the step-selection functions as response variables, and the number of telemetered individuals in each population as a weighting factor. Second, we used weighted ANCOVA to control for the effect of body size (i.e., mean species-specific, sex-unspecific body weight of an adult; Wilson & Mittermeier 2011) in resource selection. When we did not find a statistically significant effect of digestive system on resource selection after controlling for body size, we pooled equids and ruminants into weighted regression models to test how resource selection was influenced by body size. When the effect of digestive system on resource selection was statistically significant after controlling for body size, we used weighted regression models with body size as a predictor for equids and for ruminants separately. We switched the direction of parameter estimates for distance to surface water in all analyses and graphs. Therefore, positive and negative values show selection and avoidance for forage biomass, potential energy intake, and surface water, respectively.

Results

Forage biomass, potential energy intake, distance to surface water, or some combination thereof significantly influenced resource selection in 23 out of 30 equid and ruminant populations (Fig. 2; Table S4). Selection for forage biomass was exhibited by two equids (18%) and five ruminants (26%); avoidance of forage biomass was exhibited by one equid (1%) and five ruminants (26%). Seven ruminants (37%) and no equids selected for potential energy intake (i.e., had movement consistent with the FMH), and one ruminant (0.5%) avoided potential energy intake. Six out of 11 equid populations (55%) selected for surface water, with 10 out of 11 equid populations (91%) having a positive parameter estimates for surface water (Fig. 2). Ruminant populations displayed a diversity of selection behaviors toward surface water. Overall, equids consistently selected for surface water, while resource selection of ruminants was variable (Fig. 2).

After controlling for the effect of body size, we did not detect any difference in selection for forage biomass ($P = 0.18$; Fig. 3A) or potential energy intake ($P = 0.37$; Fig. 3C) between equids and ruminants. Body size did not explain variation in selection for forage biomass (Fig. 3B) but did explained 25% of the variation in selection for potential energy intake (Fig. 3D). Equids selected for surface water more strongly than ruminants after controlling for the effect of body size ($F_{(1, 27)} = 7.09$, $P = 0.013$; Fig. 3E). Body size did not explain selection for surface water in equids and ruminants ($P \geq 0.23$; Fig. 3F and 3G).

Discussion

We quantified how selection for forage biomass, potential energy intake, and surface water shaped the movements of free-ranging ungulates across the globe. In support of the FMH, selection for potential energy intake (i.e., intermediate forage biomass and intermediate forage

digestibility) was most frequent among smaller-bodied ungulates. Selecting forage at early to intermediate states of phenology reflects the higher mass-specific energetic requirements of smaller-bodied ungulates (Illius & Gordon 1987; Hopcraft *et al.* 2012). In contrast, larger-bodied ungulates (comprising both equids and larger-bodied ruminants) foraged in a manner distinct from the central prediction of the FMH. In accordance with our expectations based on hindgut fermentation, equids selected consistently for proximity to surface water. In contrast to our expectations based on ruminant digestion coupled with higher absolute energetic requirements, larger-bodied ruminants did not consistently select for forage biomass. We offer two explanations for equivocal support of the hypothesis that larger-bodied ruminants should prioritize high forage biomass: (1) methodological limitations; and (2) taxonomic and functional diversity.

Combined with high-resolution data from GPS-telemetered individuals, remotely-sensed vegetation indices allow for testing of the FMH across study populations (Wilmers *et al.* 2013). However, methodological limitations are inherent to such indices, and synthetic studies like ours should be viewed as complementary to (rather than as a substitute for) intensive, field-based investigations, in which forage biomass and quality are quantified directly (e.g., Wilmshurst *et al.* 2000, Hebblewhite *et al.* 2008, Geremia *et al.* 2019). This is particularly the case when remotely sensed vegetation indices—MSAVI and its derivative, IRG—are assumed to be correlated with forage biomass and potential energy intake. Although we did not validate these metrics against field-collected data on forage biomass and potential energy intake (i.e., forage quality), several studies have quantified relationships between forage biomass, forage quality, and remotely-sensed metrics directly (e.g., Hebblewhite *et al.* 2008, Wilmers *et al.* 2013, Geremia *et al.* 2019), and found that such metrics are suitable representations of forage biomass

and quality. Therefore, we believe that MSAVI and IRG (i.e., remotely-sensed metrics) are reasonable proxies for forage biomass and potential energy intake, and they are the only way (with which we are familiar) to conduct standardized tests of the FMH across many study populations that range across entire landscapes. A major challenge for future research is to couple field validations of remotely-sensed vegetation indices with the high resolution movement trajectories afforded by GPS telemetry.

Additionally, and at the spatial scale of MODIS, such indices cannot distinguish between sources of “greenness” resulting from different vegetation types, such as woody plants and grasses (Archibald & Scholes 2007; Gaughan *et al.* 2013). So, pixel values could be associated with vegetation that did not necessarily represent forage from an ungulate’s perspective. We attempted to minimize the effect of this potential limitation by restricting our analysis to defined growing seasons, and by restricting our analysis to study areas in which dense forest (per Friedl & Sulla-Menashe 2015) and % tree cover (per Hansen *et al.* 2013) were minimal. Because the green-up profile in leaves of woody plants is usually constant, the dramatic change in MSAVI (which we used to define the growing seasons) is mostly associated with the green-up of grasses rather than green-up of woody plants (Archibald & Scholes 2007; Higgins *et al.* 2011). Therefore, the potential for such phenological confounding should be restricted to ecosystems in which woody plants and grasses co-occur in similar proportions yet exhibit different seasonality (e.g., eastern and southern African savannas and European forests).

Differences in the strength and consistency of resource selection were likely due in part to the relatively high diversity of ruminants in our study, which incorporated 11 genera exhibiting a >20-fold difference in body size (from 24 kg roe deer to 590 kg African buffalo) with additional variation in feeding strategies (e.g., unselective grazing, selective browsing,

mixed feeding). However, all species of ruminants in our study foraged largely or wholly in the understory layer or in open rangelands, for which phenological dynamics were captured by our forage biomass metrics. Our results also are consistent with site-specific studies on ungulate assemblages, where multiple ruminants coexist via resource specialization, and therefore exhibit a diverse array of resource selection (Wilmshurst *et al.* 2000; Kartzinel *et al.* 2015).

In Serengeti National Park, Wilmshurst *et al.* (2000) documented empirical support for theoretical expectations that resource selection of ruminants should be mediated by body size. As phenology progresses, increasing forage biomass goes hand-in-hand with decreasing digestibility, presenting a challenge for smaller-bodied herbivores for which small guts filled with slowly-fermenting vegetation reduces intake rates (Fig. 1; see also Wickstrom *et al.* 1984). As a result, smaller-bodied ruminants were observed on patches of lower biomass than their larger counterparts. Our findings extend those of Wilmshurst *et al.* (2000) across four continents, and by incorporating hindgut fermenters. Hindgut fermenters were represented exclusively by equids in our study which, in contrast to ruminants, are restricted to a single genus (*Equus*), and exhibit limited (~2.5-fold) variation in body size. Consequently, resource selection was relatively consistent across equid populations, with six of 11 populations selecting areas in close proximity to surface water. Equids do not conserve water as efficiently as ruminants, and they excrete proportionately more water (Janis 1976; Ogutu *et al.* 2014); such differences in digestion likely explain the strong selection for surface water by equids across the globe.

Our remotely-sensed imagery of surface water existed at a coarser temporal resolution compared to our remotely-sensed imagery of vegetation indices, and did not comprise exhaustive data on all sources of water available to study populations. For example, ephemeral ponds and streams are not captured by the Global Surface Water Explorer. However, with a resolution of 30

x 30 m, the Global Surface Water Explorer constitutes (by far) the most precise data on the global distribution of surface water (Pekel *et al.* 2016), and we were able to supplement this remotely-sensed imagery with locally-collected locations of surface water. The strength of selection for surface water did not depend on availability of surface water within study areas (but it did depend on variability in the distance to surface water [Fig. S2]) and, per our initial prediction based on digestive system, the strength of selection for surface water was significantly stronger for equids than for ruminants. In sum, we believe that such methodological limitations associated with remotely-sensed imagery of surface water were unlikely to have had undue influence on our results and associated inferences (but see Text S2). More generally, we believe that the Global Surface Water Explorer has strong potential as a tool for wildlife and movement ecologists, and its potential methodological limitations will be overridden by its value in many study systems.

Our study represents a macroecological test of drivers of ungulate resource selection across the globe. By necessity, our synthetic approach sacrifices some area- and population-specific precision in attempt to identify general trends (Levin 1992, Brown 1995). For example, our use of body size as a predictor variable integrates a suite of size-dependent relationships, including retention time in the digestive tract and cropping rate. In particular, variation in cropping rates among species has potential to influence resource selection: for a given body size, a narrower-muzzled species could more efficiently forage on low biomass compared to a wider-muzzled species (Spalinger and Hobbs 1992). As another example, all ruminant populations in our study were sympatric with other wild and domestic ruminants, which could influence resource selection of free-ranging ungulates (e.g., Mishra *et al.* 2004; Kinnaird & O'Brien 2012). Future efforts to synthesize patterns of resource selection for free-ranging ungulates might

incorporate the occurrence and abundance of livestock and population density of the focal species (through, e.g., the Gridded Livestock of the World mapping project by the United Nations). Through resource competition, creating barriers to movement (e.g., fences, roads), and direct interference, humans can suppress the potential for free-ranging ungulates to exploit spatiotemporal variability in forage biomass, potential energy intake, and access to surface water (e.g., Sanderson *et al.* 2002; Ogutu *et al.* 2014; Panzacchi *et al.* 2015; Tucker *et al.* 2018; but see Young *et al.* 2018). By constraining movements of free-ranging ungulates, human activity may dampen any effects of body size and digestive system on resource selection and result in deviations from the central prediction of the FMH. Moreover, individual characteristics such as sex, age, body mass, and reproductive status of individuals strongly affect energy requirements and thus resource selection by large ungulates (e.g., Forsyth *et al.* 2005; Hamel & Côté 2008; Saïd *et al.* 2009; Brivio *et al.* 2014). Additionally, physical constraints inherent to different study areas (e.g., the spatial scale over which variation in forage biomass and potential energy intake arise) likely influence the movement and resource selection of free-ranging ungulates to some degree (Aikens *et al.* 2020). Such area- and population-specific variability almost certainly contributes to differences in resource selection between populations of the same species (e.g., khulan populations in western versus southern Mongolia; Text S2), and could explain variation around the general trends depicted in Figures 2 and 3. Finally, future studies could benefit from separating different movement states (e.g., foraging, resting, travelling) to explicitly investigate resource selection during foraging bouts (Fryxell *et al.* 2008; Gurarie *et al.* 2009). Nevertheless, our analyses point to important generalities—stronger selection of surface water by equids relative to ruminants, and stronger selection for potential energy intake by smaller-bodied ruminants—which conform to expectations based on metabolic allometry and digestive system.

In combination with intensive, longitudinal field observations within each study population, we believe that our comparative cross-taxa study has helped illuminate general rules and contingencies associated with the FMH.

Our synthetic approach provides the first cross-taxa test of the Forage Maturation Hypothesis. By using a combination of remotely-sensed data to quantify forage biomass, potential energy intake, and surface water, we assessed differential selection of resources by free-ranging ungulates across the terrestrial surface. In doing so, we have refined the Forage Maturation Hypothesis relative to its original formulation to explicitly consider variation in ungulate body size and digestive system, thus extending the applicability of this key concept in large herbivore ecology. The forage characteristics that influence population persistence of free-ranging ungulates should differ according to body size and digestive system, such that access to a combination of resource gradients is key to maintaining viable populations of diverse free-ranging ungulates across the globe.

Acknowledgements

The authors are grateful to the following organizations for funding, data collection, and co-ownership: Berry Biodiversity Center at the University of Wyoming, Chicago Zoological Society, Iranian Department of Environment, Isfahan University of Technology, and Saint Louis Zoo (SE, M-RH, JRG); Etosha Ecological Institute (BA); Austrian Science Fund project 18624 (PK); Oyu Tolgoi LLC Core Biodiversity Monitoring Program, Wildlife Conservation Society, Sustainability East Asia LLC (BB, PK, JCP); Bureau of Land Management (JLB, JDH, HS, JDS, SRBK, KAS); University of Wyoming's College of Agriculture and Natural Resources Extension, the Y Cross Ranch Endowment, and the U.S. Department of Agriculture's McIntire

Stennis Project (JLB, JDH, JDS); the U.S. National Science Foundation (RBB, DIR); Zimbabwe Parks and Wildlife Management Authority and the Zone Atelier (LTSER) Hwange program of the CNRS (SC-J); SANParks Scientific Services (PCC); the U.S. Geological Survey (PCC, KAS, SRBK); Smithsonian Conservation Biology Institute, National Zoological Park, Senckenberg Biodiversity and Climate Research Centre, Minnesota Zoo, and Hustai National Park (ND, SS); Robert Bosch Foundation and the German Federal Ministry of Education and Research (TM, ND); Natural Sciences and Engineering Research Council of Canada (ATF, JRG); Japan Ministry of Education, Sports, Science, and Technology (TYI); Federal Ministry of Education and Research (TM); The Nature Conservancy's Africa Programme (JRM); Norwegian Institute for Nature Research and Norwegian Environment Agency (MP, OS, GR); Research Council of Norway (JDCL); the Euromammals/Eurodeer collaborative initiative (www.eurodeer.org) for providing data and expertise; and Princeton University (DIR). We thank B. Maitland, J. Alston, B. Brito, B. Hays, F. Molina, and T. S. M. Stratmann for help with graphics and for valuable discussion. S. Focardi, N. Morellet, N. Owen-Smith, and an anonymous reviewer provided constructive criticism. We thank B. Reineking for assistance with step-selection functions, and J. Evans for assistance with remotely-sensed data layers.

References

- Aikens, E.O., Kauffman, M.J., Merkle, J.A., Dwinell, S.P.H., Fralick, G.L. & Monteith, K.L. (2017). The greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecol. Lett.*, 20, 741–750.
- Aikens, E. O., Myserud, A., Merkle, J. A., Cagnacci, F., Rivrud, I. M., Hebblewhite, M., ... &

526 Kauffman, M. J. (2020). Wave-like patterns of plant phenology determine ungulate
527 movement tactics. *Curr. Biol.*, 30, 3444-3449.

528 Albon, S.D. & Langvatn, R. (1992). Plant phenology and the benefits of migration in a temperate
529 ungulate. *Oikos*, 65, 502-513.

530 Ali, A. H., Ford, A. T., Evans, J. S., Mallon, D. P., Hayes, M. M., et al. (2017). Resource
531 selection and landscape change reveal mechanisms suppressing population recovery for the
532 world's most endangered antelope. *J. Appl. Ecol.*, 54, 1720-1729.

533 Archibald, S. & Scholes, R.J. (2007). Leaf green-up in a semi-arid African savanna –separating
534 tree and grass responses to environmental cues. *J. Veg. Sci.*, 18, 583-594.

535 Arnold, T.W. (2010). Uninformative parameters and model selection using Akaike's Information
536 Criterion. *J. Wildl. Manage.*, 74, 1175–1178.

537 Attias, N., Oliveira-Santos, L. G. R., Fagan, W. F., & Mourão, G. (2018). Effects of air
538 temperature on habitat selection and activity patterns of two tropical imperfect
539 homeotherms. *Anim. Behav.*, 140, 129-140.

540 Avgar, T., Mosser, A., Brown, G.S. & Fryxell, J.M. (2013). Environmental and individual
541 drivers of animal movement patterns across a wide geographical gradient. *J. Anim. Ecol.*,
542 82, 96–106.

543 Avgar, T., Potts, J.R., Lewis, M.A. & Boyce, M.S. (2016). Integrated step selection analysis:
544 Bridging the gap between resource selection and animal movement. *Methods Ecol. Evol.*, 7,
545 619–630.

546 Bailey, D.W., Gross, J.E., Laca, E.A., Rittenhouse, L.R., Coughenour, M.B., Swift, D.M., et al.

547 (1996). Mechanisms that result in large herbivore grazing distribution patterns. *J. Range*
 548 *Manag.*, 49, 386–400.

549 Barboza, P.S. & Bowyer, R.T. (2000). Sexual segregation in demorthic deer: a new gastrocentric
 550 hypothesis. *J. Mammal.*, 81, 473–489.

551 Bell, R.H. V. (1971). A grazing ecosystem in the Serengeti. *Sci. Am.*, 225, 86–93.

552 Bischof, R., Loe, L.E., Meisingset, E.L., Zimmermann, B., van Moorter, B. & Mysterud, A.
 553 (2012). A migratory northern ungulate in the pursuit of spring: Jumping or surfing the green
 554 wave? *Am. Nat.*, 180, 407–424.

555 Boone, R.B., Thirgood, S.J., & Hopcraft, J.G.C. (2006). Serengeti wildebeest migratory patterns
 556 modeled from rainfall and new vegetation growth. *Ecology*, 87, 1987–1994.

557 Borowik, T., N. Pettorelli, L. Soennichsen, and B. Jedrzejewska. (2013). Normalized difference
 558 vegetation index (NDVI) as a predictor of forage availability for ungulates in forest and
 559 field habitats. *Eur. J. Wildl. Res.* 59, 675–682.

560 Branco, P.S., Merkle, J.A., Pringle, R.M., Pansu, J., Potter, A.B., Reynolds, A., *et al.* (2019).
 561 Determinants of elephant foraging behaviour in a coupled human-natural system: Is brown
 562 the new green? *J. Anim. Ecol.*, 88, 780–792.

563 Brivio, F., Grignolio, S., Brambilla, A., & Apollonio, M. (2014). Intra-sexual variability in
 564 feeding behaviour of a mountain ungulate: size matters. *Behav. Ecol. Sociobiol.*, 68, 1649–
 565 1660.

566 Brown, J.H. (1995). *Macroecology*. University of Chicago Press, Chicago, IL.

567 Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and

568 multimodel inference in behavioral ecology: some background, observations, and
569 comparisons. *Behav Ecol Sociobiol*, 65, 23-35.

570 Calabrese, J. M., Fleming, C. H., & Gurarie, E. (2016). ctm: An r package for analyzing animal
571 relocation data as a continuous- time stochastic process. *Methods Ecol. Evol.*, 7, 1124-
572 1132.

573 Demment, M.W. & Van Soest, P.J. (1985). A nutritional explanation for body-size patterns of
574 ruminant and nonruminant herbivores. *Am. Nat.*, 125, 641–672.

575 Drescher, M., Heitkönig, I. M., Van Den Brink, P. J., & Prins, H. H. (2006). Effects of sward
576 structure on herbivore foraging behaviour in a South African savanna: an investigation of
577 the forage maturation hypothesis. *Austral Ecol.*, 31, 76-87.

578 Duncan, P., Foose, T.J., Gordon, I.J., Gakahu, C.G. & Lloyd, M. (1990). Comparative nutrient
579 extraction from forages by grazing bovids and equids: a test of the nutritional model of
580 equid/bovid competition and coexistence. *Oecologia*, 84, 411–418.

581 Dupke, C., Bonenfant, C., Reineking, B., Hable, R., Zeppenfeld, T., Ewald, M., *et al.* (2017).
582 Habitat selection by a large herbivore at multiple spatial and temporal scales is primarily
583 governed by food resources. *Ecography*, 40, 1014–1027.

584 Edouard, N., Duncan, P., Dumont, B., Baumont, R., & Fleurance, G. (2010). Foraging in a
585 heterogeneous environment—An experimental study of the trade-off between intake rate
586 and diet quality. *Appl. Anim. Behav. Sci.*, 126, 27-36.

587 Fleming, C.H., Calabrese, J.M., Mueller, T., Olson, K.A., Leimgruber, P. & Fagan, W.F. (2014).
588 From fine-scale foraging to home ranges: a semivariance approach to identifying movement

589 modes across spatiotemporal scales. *Am. Nat.*, 183, E154–E167.

590 Fleurance, G., Duncan, P., Fritz, H., Gordon, I. J., & Grenier-Loustalot, M. F. (2010). Influence
591 of sward structure on daily intake and foraging behaviour by horses. *Animal*, 4, 480–485.

592 Forsyth, D. M., Duncan, R. P., Tustin, K. G., & Gaillard, J. M. (2005). A substantial energetic
593 cost to male reproduction in a sexually dimorphic ungulate. *Ecology*, 86, 2154–2163.

594 Fortin, D., Beyer, H.L., Boyce, M.S., Smith, D.W., Duchesne, T. & Mao, J.S. (2005). Wolves
595 influence elk movements: behavior shapes a trophic cascade in Yellowstone National Park.
596 *Ecology*, 86, 1320–1330.

597 Friedl, M., Sulla-Menashe, D. (2015). *MCD12C1 MODIS/Terra+Aqua Land Cover Type Yearly*
598 *L3 Global 0.05Deg CMG V006* [Data set]. NASA EOSDIS Land Processes DAAC.
599 Accessed 2021-01-06 from <https://doi.org/10.5067/MODIS/MCD12C1.006>.

600 Fryxell, J.M. (1991). Forage quality and aggregation by large herbivores. *Am. Nat.*, 138, 478–
601 498.

602 Fryxell, J. M., Hazell, M., Börger, L., Dalziel, B. D., Haydon, D. T., Morales, J. M., ... &
603 Rosatte, R. C. (2008). Multiple movement modes by large herbivores at multiple
604 spatiotemporal scales. *Proc. Natl. Acad. Sci. USA*, 105, 19114–19119.

605 Gaughan, A.E., Holdo, R.M. & Anderson, T.M. (2013). Using short-term MODIS time-series to
606 quantify tree cover in a highly heterogeneous African savanna. *Int. J. Remote Sens.*, 34,
607 6865–6882.

608 Geremia, C., Merkle, J.A., Eacker, D.R., Wallen, R.L., White, P.J., Hebblewhite, M., &
609 Kauffman, M.J. (2019). Migrating bison engineer the green wave. *Proc. Natl. Acad. Sci.*
610 *USA*, 116, 25707–25713.

611 Geri, F., Amici, V. & Rocchini, D. (2010). Human activity impact on the heterogeneity of a
612 Mediterranean landscape. *Appl. Geogr.*, 30, 370–379.

613 Gurarie, E., Andrews, R. D., & Laidre, K. L. (2009). A novel method for identifying behavioural
614 changes in animal movement data. *Ecol. Lett.*, 12, 395–408.

615 Hamel, S., & Côté, S. D. (2008). Trade-offs in activity budget in an alpine ungulate: contrasting
616 lactating and nonlactating females. *Anim. Behav.*, 75, 217–227.

617 Hamel, S., Garel, M., Festa-Bianchet, M., Gaillard, J.-M. & Côté, S.D. (2009). Spring
618 Normalized Difference Vegetation Index (NDVI) predicts annual variation in timing of
619 peak faecal crude protein in mountain ungulates. *J. Appl. Ecol.*, 46, 582–589.

620 Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., *et al.*
621 (2013). High-Resolution Global Maps of 21st-Century Forest Cover Change: *Science*, 342,
622 850–853.

623 Hebblewhite, M., Merrill, E. & McDermid, G. (2008). A multi-scale test of the Forage
624 Maturation Hypothesis in a partially migratory ungulate population. *Ecol. Monogr.*, 78,
625 141–166.

626 Higgins, S.I., Delgado-Cartay, M.D., February, E.C. & Combrink, H.J. (2011). Is there a
627 temporal niche separation in the leaf phenology of savanna trees and grasses? *J. Biogeogr.*,
628 38, 2165–2175.

629 Hofmann, R.R. (1989). Evolutionary steps of ecophysiological adaptation and diversification of
630 ruminants: a comparative view of their digestive system. *Oecologia*, 78, 443–457.

631 Holbrook, J. D., Olson, L. E., DeCesare, N. J., Hebblewhite, M., Squires, J. R., & Steenweg, R.

632 (2019). Functional responses in habitat selection: clarifying hypotheses and
 633 interpretations. *Ecol. Appl.*, 29, e01852.

634 Hopcraft, J.G.C., Anderson, T.M., Pérez-Vila, S., Mayemba, E. & Olf, H. (2012). Body size and
 635 the division of niche space: food and predation differentially shape the distribution of
 636 Serengeti grazers. *J. Anim. Ecol.*, 81, 201–213.

637 Hopcraft, J.G.C., Morales, J.M., Beyer, H.L., Borner, M., Mwangomo, E., Sinclair, A.R.E., *et al.*
 638 (2014). Competition, predation, and migration: Individual choice patterns of Serengeti
 639 migrants captured by hierarchical models. *Ecol. Monogr.*, 84, 355–372.

640 Huang, C., Chen, Y., Zhang, S. & Wu, J. (2018). Detecting, extracting, and monitoring surface
 641 water from space using optical sensors: a review. *Rev. Geophys.*, 56, 333–360.

642 Illius, A.W. & Gordon, I.J. (1987). The allometry of food intake in grazing ruminants. *J. Anim.*
 643 *Ecol.*, 56, 989.

644 Illius, A.W. & Gordon, I.J. (1992). Modelling the nutritional ecology of ungulate herbivores:
 645 evolution of body size and competitive interactions. *Oecologia*, 89, 428–434.

646 Janis, C. (1976). The evolutionary strategy of the Equidae and the origins of rumen and cecal
 647 digestion. *Evolution*, 30, 757–774.

648 Jarman, P.J. (1974). The social organisation of antelope in relation to their ecology. *Behaviour*,
 649 48, 215-267.

650 Jesmer, B.R., Merkle, J.A., Goheen, J.R., Aikens, E.O., Beck, J.L., Courtemanch, A.B., *et al.*
 651 (2018). Is ungulate migration culturally transmitted? Evidence of social learning from
 652 translocated animals. *Science*, 361, 1023–1025.

653 Jin, Y., Yang, X., Qiu, J., Li, J., Gao, T., Wu, Q., *et al.* (2014). Remote sensing-based biomass
654 estimation and its spatio-temporal variations in temperate grassland, Northern China.
655 *Remote Sens.*, 6, 1496-1513.

656 Kartzinel, T.R., Chen, P.A., Coverdale, T.C., Erickson, D.L., Kress, W.J., Kuzmina, M.L., *et al.*
657 (2015). DNA metabarcoding illuminates dietary niche partitioning by African large
658 herbivores. *Proc. Natl. Acad. Sci. USA*, 112, 8019–8024.

659 Kinnaird, M.F. & O’Brien, T.G. (2012). Effects of private-land use, livestock management, and
660 human tolerance on diversity, distribution, and abundance of large African mammals.
661 *Conserv. Biol.*, 26, 1026–1039.

662 Levin, S.A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73, 1943-1967.

663 Liang, T., Yang, S., Feng, Q., Liu, B., Zhang, R., Huang, X., *et al.* (2016). Multi-factor modeling
664 of above-ground biomass in alpine grassland: A case study in the Three-River Headwaters
665 Region, China. *Remote Sens. Environ.*, 186, 164-172.

666 McNab, B.K. (1974). The energetics of endotherms. *Ohio J. Sci.*, 74, 370–380.

667 Merkle, J.A., Monteith, K.L., Aikens, E.O., Hayes, M.M., Hersey, K.R., Middleton, A.D., *et al.*
668 (2016). Large herbivores surf waves of green-up during spring. *Proc. R. Soc. B Biol. Sci.*,
669 283, 20160456.

670 Middleton, A.D., Kauffman, M.J., McWhirter, D.E., Cook, J.G., Cook, R.C., Nelson, A.A., *et al.*
671 (2013). Animal migration amid shifting patterns of phenology and predation: lessons from a
672 Yellowstone elk herd. *Ecology*, 94, 1245–1256.

673 Middleton, A.D., Merkle, J.A., McWhirter, D.E., Cook, J.G., Cook, R.C., White, P.J., *et al.*

674 (2018). Green-wave surfing increases fat gain in a migratory ungulate. *Oikos*, 127, 1060–
 675 1068.

676 Mishra, C., Van Wieren, S.E., Ketner, P., Heitkönig, I.M.A. & Prins, H.H.T. (2004).
 677 Competition between domestic livestock and wild bharal *Pseudois nayaur* in the Indian
 678 Trans-Himalaya. *J. Appl. Ecol.*, 41, 344–354.

679 Mueller, T., Olson, K.A., Fuller, T.K., Schaller, G.B., Murray, M.G. & Leimgruber, P. (2008). In
 680 search of forage: predicting dynamic habitats of Mongolian gazelles using satellite-based
 681 estimates of vegetation productivity. *J. Appl. Ecol.*, 45, 649-658.

682 Muntifering, J.R., Ditmer, M.A., Stapleton, S., Naidoo, R. & Harris, T.H. (2019). Hartmann's
 683 mountain zebra resource selection and movement behavior within a large unprotected
 684 landscape in northwest Namibia. *Endanger. Species Res.*, 38, 159-170.

685 Mysterud, A., & Ims, R. A. (1998). Functional responses in habitat use: Availability influences
 686 relative use in trade- off situations. *Ecology*, 79, 1435-1441.

687 Nathan, R., Getz, W.M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. & Smouse, P.E. (2008).
 688 A movement paradigm for unifying organismal movement research. *Proc. Natl. Acad. Sci.*
 689 *USA*, 105, 19052-19059.

690 Ogutu, J.O., Piepho, H.P., Reid, R.S., Rainy, M.E., Kruska, R.L., Worden, J.S., *et al.* (2010).
 691 Large herbivore responses to water and settlements in savannas. *Ecol. Monogr.*, 80, 241–
 692 266.

693 Ogutu, J.O., Reid, R.S., Piepho, H.P., Hobbs, N.T., Rainy, M.E., Kruska, R.L., *et al.* (2014).
 694 Large herbivore responses to surface water and land use in an East African savanna:

695 implications for conservation and human-wildlife conflicts. *Biodivers. Conserv.*, 23, 573–
696 596.

697 Olff, H., Ritchie, M.E. & Prins, H.H. (2002). Global environmental controls of diversity in large
698 herbivores. *Nature*, 415, 901-904.

699 Owen-Smith, N., Fryxell, J.M. & Merrill, E.H. (2010). Foraging theory upscaled: the behavioural
700 ecology of herbivore movement. *Philos. Trans. R. Soc. B Biol. Sci.*, 365, 2267-2278.

701 Owens, F.N. & Basalan, M. (2016). Ruminant fermentation. In: *Rumenology*. pp. 63–102.
702 Springer, Cham.

703 Panzacchi, M., Van Moorter, B., Strand, O., Loe, L. E., & Reimers, E. (2015). Searching for the
704 fundamental niche using individual- based habitat selection modelling across
705 populations. *Ecography*, 38, 659–669.

706 Pekel, J.F., Cottam, A., Gorelick, N. & Belward, A.S. (2016). High-resolution mapping of global
707 surface water and its long-term changes. *Nature*, 540, 418–422.

708 Pettorelli, N., Vik, J.O., Mysterud, A., Gaillard, J.M., Tucker, C.J. & Stenseth, N.C. (2005).
709 Using the satellite-derived NDVI to assess ecological responses to environmental change.
710 *Trends Ecol. Evol.*, 20, 503-510.

711 Prima, M.-C., Duchesne, T. & Fortin, D. (2017). Robust inference from conditional logistic
712 regression applied to movement and habitat selection analysis. *PLoS One*, 12, e0169779.

713 Qi, J., Chehbouni, A., Huete, A.R., Kerr, Y.H. & Sorooshian, S. (1994). A modified soil adjusted
714 vegetation index. *Remote Sens. Environ.*, 48, 119–126.

715 R Core Team. (2019). R: A language and environment for statistical computing. *R Found. Stat.*

716 *Comput. Vienna, Austria. URL <http://www.R-project.org/>.*

717 Redfern, J. V., Grant, R., Biggs, H. & Getz, W.M. (2003). Surface-water constraints on
718 herbivore foraging in the Kruger National Park, South Africa. *Ecology*, 84, 2092–2107.

719 Rivrud, I.M., Heurich, M., Krupczynski, P., Müller, J. & Mysterud, A. (2016). Green wave
720 tracking by large herbivores: an experimental approach. *Ecology*, 97, 3547-3553.

721 Roever, C.L., Boyce, M.S. & Stenhouse, G.B. (2010). Grizzly bear movements relative to roads:
722 Application of step selection functions. *Ecography*, 33, 1113–1122.

723 Rozen-Rechels, D., Dupoué, A., Lourdaïs, O., Chamaillé-Jammes, S., Meylan, S., Clobert, J., *et*
724 *al.* (2019). When water interacts with temperature: Ecological and evolutionary implications
725 of thermo-hydroregulation in terrestrial ectotherms. *Ecol. Evol.*, 9, 10029–10043.

726 Saïd, S., Gaillard, J. M., Widmer, O., Débias, F., Bourgoïn, G., Delorme, D., & Roux, C. (2009).
727 What shapes intra- specific variation in home range size? A case study of female roe
728 deer. *Oikos*, 118, 1299-1306.

729 Sanderson, E.W., Jaiteh, M., Levy, M.A., Redford, K.H., Wannebo, A. V. & Woolmer, G.
730 (2002). The human footprint and the last of the wild. *Bioscience*, 52, 891-904.

731 Santos, A. S., Rodrigues, M. A. M., Bessa, R. J. B., Ferreira, L. M., & Martin-Rosset, W. (2011).
732 Understanding the equine cecum-colon ecosystem: current knowledge and future
733 perspectives. *Animal*, 5, 48-56.

734 Schoenecker, K.A., King, S.R.B., Nordquist, M.K., Nandintsetseg, D. & Cao, Q. (2016). Habitat
735 and diet of equids. In: *Wild Equids: Ecology, Management, and Conservation*. Johns
736 Hopkins University Press, pp. 41–57.

737 Senft, R.L., Coughenour, M.B., Bailey, D.W., Rittenhouse, L.R., Sala, O.E. & Swift, D.M.
738 (1987). Large herbivore foraging and ecological hierarchies. *Bioscience*, 37, 789–799.

739 Signer, J., Fieberg, J. & Avgar, T. (2019). Animal movement tools (amt): R package for
740 managing tracking data and conducting habitat selection analyses. *Ecol. Evol.*, 9, 880–890.

741 Spalinger, D.E., & N.T. Hobbs. (1992). Mechanisms of foraging in mammalian herbivores: new
742 models of functional response. *Am. Nat.*, 140, 325-348.

743 St-Louis, A., & Côté, S. D. (2014). Resource selection in a high-altitude rangeland equid, the
744 kiang (*Equus kiang*): influence of forage abundance and quality at multiple spatial
745 scales. *Can J Zool*, 92, 239-249.

746 van Soest, P.J. (1996). Allometry and ecology of feeding behavior and digestive capacity in
747 herbivores: a review. *Zoo Biol.*, 5, 455-479.

748 Stabach, J.A., Wittemyer, G., Boone, R.B., Reid, R.S. & Worden, J.S. (2016). Variation in
749 habitat selection by white-bearded wildebeest across different degrees of human
750 disturbance. *Ecosphere*, 7, e01428.

751 Therneau, T.M. & T. Lumley. (2015). Package ‘survival’. *R Package*.

752 Thurfjell, H., Ciuti, S. & Boyce, M.S. (2014). Applications of step-selection functions in ecology
753 and conservation. *Mov. Ecol.*, 2/1/4.

754 Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., Fryxell, J.M., Van Moorter, B., Alberts, S.C., *et*
755 *al.* (2018). Moving in the Anthropocene: global reductions in terrestrial mammalian
756 movements. *Science.*, 359, 466–469.

757 Veldhuis, M.P., Kihwele, E.S., Cromsigt, J.P.G.M., Ogutu, J.O., Hopcraft, J.G.C., Owen-Smith,

- N., *et al.* (2019). Large herbivore assemblages in a changing climate: incorporating water dependence and thermoregulation. *Ecol. Lett.*, 22, 1536–1546.
- Wickstrom, M. L., Robbins, C. T., Hanley, T. A., Spalinger, D. E., & Parish, S. M. (1984). Food intake and foraging energetics of elk and mule deer. *J. Wildl. Mgmt.*, 48:1285-1301.
- Wilmers, C., Ram, K., Watson, F.G.R., White, P.J., Smith, D.W. & Levi, T. (2013). Climate and vegetation phenology: predicting the effects of warming temperatures. In: *Yellowstone's wildlife in transition* (eds. White, PJ, Garrott, RA & Plumb, GE). Harvard University Press Cambridge, MA.
- Wilmshurst, J.F., Fryxell, J.M. & Bergman, C.M. (2000). The allometry of patch selection in ruminants. *Proc. R. Soc. B Biol. Sci.*, 267, 345-349.
- Wilson, D.E. & Mittermeier, R.A. (2011). *Handbook of the mammals of the world: v.2: Hoofed mammals*. Barcelona, Spain: Lynx Edicions.
- Young, T.P., Porensky, L.M., Riginos, C., Veblen, K.E., Odadi, W.O., Kimuyu, D.M., *et al.* (2018). Relationships between cattle and biodiversity in a multiuse landscape revealed by the Kenya Long-Term Exclosure Experiment. *Rangel. Ecol. Manag.*, 71, 281–291.

Figures

Figure 1. Illustrated predictions for resource selection by equids and different sizes of ruminants during a hypothetical growing season and in a hypothetical landscape. Forage biomass (dark green line) and potential energy intake (light green line; measured by rate of change in forage biomass) increase at beginning of the growing season. The potential for maximum energy intake occurs at the middle of the growing season, when forage biomass is at intermediate levels.

779 Different sizes and shades of grass represent phenological stages of forage biomass and
 780 digestibility. At early phenological stages (i.e., the lightest shade of green grass), potential
 781 energy intake is low because forage biomass is low. At late phenological and senescent stages
 782 (i.e., the darkest shade of green grass and brown grass, respectively), potential energy intake is
 783 low because forage digestibility is low. (A) The Forage Maturation Hypothesis predicts that
 784 ungulates maximize their energy intake by selecting forage of intermediate biomass at
 785 intermediate phenological stages. (B) However, selection for forage characteristics should also
 786 depend on body size. (C) Smaller-bodied ungulates (which exhibit ruminant digestion) should
 787 select most strongly for maximal energy intake (light green), larger-bodied ruminants should
 788 select for forage biomass (dark green), and equids should select for surface water (blue) to meet
 789 their physiological needs.

790 Figure 2. Locations of the study populations (top panel) and mean \pm SE parameter estimates of
 791 step-selection functions (SSF) for 11 populations of equids (second panel) and 19 populations of
 792 ruminants (third and fourth panels) during growing seasons. Significant variables with 95%
 793 confidence intervals excluding zero are marked by asterisks. Populations are numbered in
 794 increasing order of a focal species' body size: 1-roe deer (n = 23); 2-goitered gazelle (n = 6); 3-
 795 Mongolian gazelle (n = 7); 4-Mongolian gazelle (n = 5); 5-saiga (n = 26); 6-springbok (n = 10);
 796 7-impala (n = 21); 8-mule deer (n = 100); 9-mule deer (n = 78); 10-reindeer (n = 25); 11-red deer
 797 (n = 51); 12-red deer (n = 22); 13-red deer (n = 13); 14-feral burro (n = 10); 15-wildebeest (n =
 798 9); 16-wildebeest (n = 12); 17-wildebeest (n = 13); 18-khulan (n = 7); 19-khulan (n = 9); 20-
 799 onager (n = 9); 21-plains zebra (n = 9); 22-plains zebra (n = 31); 23-Przewalski's horse (n = 14);
 800 24-Przewalski's horse (n = 5); 25-elk (n = 20); 26-elk (n = 7); 27-mountain zebra (n = 5); 28-
 801 Grevy's zebra (n = 7); 29-feral horse (n = 22); and 30-African buffalo (n = 4).

802 Figure 3. Relationship between population-level resource selection coefficients and digestive
803 system (A, C, E; weighted analysis of covariance [ANCOVA]) and body size (B, D, F, G;
804 weighted least squares regression). Equid populations more strongly selected surface water after
805 controlling for the effect of body size (E), but we detected no significant difference in selection
806 for forage biomass (A) and potential energy intake (C) after controlling for the effect of body
807 size differences between equids and ruminants. The effect of body size on selection for potential
808 energy intake was statistically significant for all free-ranging ungulates (D; red dashed line). The
809 effect of body size on selection for surface water was not statistically significant for equids (F)
810 nor for ruminants (G).





