



HAL
open science

Core size of a random partition for the Plancherel measure

Salim Rostam

► **To cite this version:**

| Salim Rostam. Core size of a random partition for the Plancherel measure. 2021. hal-03426306v1

HAL Id: hal-03426306

<https://hal.science/hal-03426306v1>

Preprint submitted on 12 Nov 2021 (v1), last revised 2 Sep 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Core size of a random partition for the Plancherel measure

Salim ROSTAM*

Abstract

We prove that the size of the e -core of a partition taken under the Poissonised Plancherel measure converges in distribution to, as the Poisson parameter goes to $+\infty$ and after a suitable renormalisation, a sum of $e - 1$ mutually independent Gamma distributions with explicit parameters. Such a result already exists for the uniform measure on the set of partitions of n as $n \rightarrow +\infty$, the parameters of the Gamma distributions being all equal. We rely on the fact that the descent set of a partition is a determinantal point process under the Poissonised Plancherel measure and on a central limit theorem for such processes.

1 Introduction

The conjugacy classes of the symmetric group \mathfrak{S}_n on n letters are parametrised by the set \mathcal{P}_n of *partitions* of the integer n . These are the sequences $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$ of positive integers with sum $|\lambda| := n$. In particular, the irreducible complex representations S^λ of \mathfrak{S}_n can be parametrised by $\lambda \in \mathcal{P}_n$. It turns out that one can do this parametrisation such that the dimension of S^λ as a \mathbb{C} -vector space is the number $\#\text{Std}(\lambda)$ of *standard Young tableaux* of shape λ , so that one has the formula

$$n! = \sum_{\lambda \in \mathcal{P}_n} \#\text{Std}(\lambda)^2 \quad (1.1)$$

(see, for instance, [Sa]). Quotienting by $n!$, the above equality exhibits a particular probability measure Pl_n on \mathcal{P}_n , known as the *Plancherel measure*. The equality (1.1) can also be seen as a consequence of the Robinson–Schensted correspondence between permutations in \mathfrak{S}_n and pairs of standard Young tableaux of the same shape with n boxes, for which the Plancherel measure Pl_n is then the image of the uniform measure of \mathfrak{S}_n . This correspondence has many other additional properties, namely: if $\lambda = (\lambda_1 \geq \dots) \in \mathcal{P}_n$ corresponds to $\sigma \in \mathfrak{S}_n$, then λ_1 is exactly the length of a longest increasing subsequence in σ (see, for instance, [Rom, Sa] for more details).

A natural question is the following: what does the probability measure Pl_n on \mathcal{P}_n look like? The question for an arbitrary n is quite complicated, however there is a very nice description as $n \rightarrow +\infty$. Namely, Logan–Shepp [LoSh] and independently Kerov–Vershik [KeVe] proved that there exists a universal limit shape Ω for the partitions taken under Pl_n . More precisely, the rim of a partition $\lambda \in \mathcal{P}_n$ converges uniformly in probability under Pl_n to the limit shape Ω as $n \rightarrow +\infty$. We represent in Figure 1 such a large partition, drawn with the Russian convention for the Young diagram, together with the limit shape¹. Note that there exist limit shapes for other probability measures on \mathcal{P}_n , namely, the uniform measure ([Ve], with many other examples inside), the Gelfand measure [LoSh, Mé10, Mé11] and the family of Schur–Weyl measures [Bi, FéMé], the latter being a generalisation of the Plancherel measure. These limit shape results, which are analogues of the law of large numbers, have analogues of the central limit theorem. The first result of this kind was given by Kerov [Ke] for the Plancherel measure: the difference between the partition and the limit shape converges in distribution to a generalised Gaussian process (see, for instance, [IvOl, FéMé, Mé11, BoSu-a] for further results on this direction).

*Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France

¹Figures 1, 5 and 6 were obtained using SAGEMATH [SM].




Figure 1: Universal limit shape for the partitions under the Plancherel measure (here with $n = 700$).

We now focus on the Plancherel measure \mathcal{P}_n . The derivative of the limit shape Ω represents the proportion of “up” and “down” steps made by the Young diagram at a given position. The “down” steps are the element of the *descent set*

$$\mathcal{D}(\lambda) := \{\lambda_a - a : a \geq 1\} \subseteq \mathbb{Z}.$$

Now if we look at several positions at the same time, for a given (finite) set $X \subseteq \mathbb{Z}$ we are interested in $\text{Pl}_n(X \subseteq \mathcal{D}(\lambda))$. These quantities as functions of X are known as *correlation functions*. It turns out that these correlation functions are well-understood for the *Poissonised* Plancherel measures on $\mathcal{P} = \coprod_{n \geq 0} \mathcal{P}_n$. Given $t > 0$, the Poissonised Plancherel measure pl_t is the Plancherel measure Pl_n where the n is chosen by a Poisson variable of parameter t . In other words, setting $\text{Pl}_n(\lambda) = 0$ if $|\lambda| \neq n$, for any $\lambda \in \mathcal{P}$ we have

$$\text{pl}_t(\lambda) = \exp(-t) \sum_{n=0}^{+\infty} \frac{t^n}{n!} \text{Pl}_n(\lambda).$$

(We denote by \exp the exponential function, to avoid any ambiguity with an integer $e \geq 2$ that we will intensively use in the sequel.) Borodin, Okounkov and Olshanski [BOO] prove that for any fixed $s \geq 1$ and $x_1, \dots, x_s \in \mathbb{Z}$ pairwise distinct, we have

$$\text{pl}_t(x_1, \dots, x_s \in \mathcal{D}(\lambda)) = \det[\mathbf{J}^t(x_a, x_b)]_{1 \leq a, b \leq s}, \quad (1.2)$$

where \mathbf{J}^t is the *discrete Bessel kernel*, built with Bessel functions of the first kind. This means that the discrete point process $\mathcal{D}(\lambda)$ is *determinantal*, with kernel \mathbf{J}^t . Under some conditions on x_1, \dots, x_s , they were able to *de-Poissonise* the result, proving that the limit of $\text{Pl}_n(x_1, \dots, x_s \in \mathcal{D}(\lambda))$ as $n \rightarrow +\infty$ satisfies an equality like (1.2), where the discrete Bessel kernel \mathbf{J}^t is replaced by the *discrete sine kernel*.

Determinantal point processes are well-studied and appear for instance in the theory of random matrices and in mathematical physics (see, for instance, [So00-a, §2]). Given a determinantal point process, for each $t > 0$ let $I_t \subseteq \mathbb{R}$ be an interval and $\#_t$ the number of points of the determinantal point process inside I_t . Then under some conditions we have the following central limit theorem:

$$\frac{\#_t - \mathbb{E}\#_t}{\sqrt{\text{Var}\#_t}} \xrightarrow[t \rightarrow +\infty]{d} \mathcal{N}(0, 1),$$

where $\mathcal{N}(0, 1)$ is the standard Gaussian distribution. This convergence in distribution was first established by Costin–Lebowitz [CoLe] for the sine kernel, and was then generalised for more general kernels by Widom and Soshnikov [So00-b, So00-a]. In the context of the determinantal point process $\mathcal{D}(\lambda)$, this central limit theorem was namely used by Bogachev and Su [BoSu-a, BoSu-b] to prove a pointwise analogue of Kerov’s central limit theorem [Ke].

Now let $e \in \mathbb{Z}_{\geq 0}$. We say that a subset $D \subseteq \mathbb{Z}$ satisfies the property (RH_e) if there exists $b \in D$ such that $b - e \notin D$. If $D \subseteq \mathbb{Z}$ satisfies (RH_e) , we denote by $\text{RH}_e(D)$ the set of diagrams $D' \subseteq \mathbb{Z}$ obtained from D by $D' = (D \setminus \{b\}) \cup \{b - e\}$ where $b \in D$ is as above.

If λ is a partition, saying that its descent set $\mathcal{D}(\lambda)$ satisfies (RH_e) is equivalent to saying that one can remove a certain subset of e adjacent boxes of the rim of the Young diagram of λ (an *e-rim hook*), the resulting diagram being still the Young diagram of a partition. Now $\mathcal{D}(\lambda)$ satisfies $\mathcal{D}(\lambda) \supseteq \mathbb{Z}_{\leq a}$ for some $a \in \mathbb{Z}$. In particular, there exists a (minimal) integer $w_e(\lambda) \in \mathbb{Z}_{\geq 0}$ such that any sequence $D^{(0)}, \dots, D^{(k)}$ with:

- $D^{(0)} = \mathcal{D}(\lambda)$,
- for any $0 \leq a < k$, the set $D^{(a)} \subseteq \mathbb{Z}$ satisfies (RH_e) and $D^{(a+1)} \in \text{RH}_e(D^{(a)})$,

satisfies $k \leq w_e(\lambda)$. The integer $w_e(\lambda)$ is the *e-weight* of λ . A sequence as above with $k = w_e(\lambda)$ is not necessarily unique, however the set $D^{(w_e(\lambda))}$ is uniquely determined from λ . The partition $\bar{\lambda}$ that corresponds to $D^{(w_e(\lambda))}$ is the *e-core* of λ . Saying that two partitions $\lambda, \mu \in \mathcal{P}_n$ have the same *e-core* is equivalent to saying that the Young diagrams of λ and μ have the same number $c_i(\lambda)$ of *i-nodes* for all $i \in \mathbb{Z}/e\mathbb{Z}$ (see [JaKe, §2.7]). It is also equivalent to λ and μ belonging to the same block of a certain Iwahori–Hecke algebra (see, for instance, [Ma, Chapter 5]).

If a limit shape exists for a given probability measure on \mathcal{P}_n for $n \rightarrow +\infty$, it is thus natural to ask what is the behaviour of $\bar{\lambda}$ as $n \rightarrow +\infty$. When \mathcal{P}_n is endowed with the uniform measure, Lulov–Pittel [LuPi] and later Ayyer–Sinha [AySi] proved that $\frac{\pi}{\sqrt{n}}|\bar{\lambda}|$ converges in distribution to the Gamma distribution $\Gamma(\frac{e-1}{2}, \sqrt{6})$ with shape $\frac{e-1}{2}$ and scale $\sqrt{6}$. In this paper, we study the behaviour of $|\bar{\lambda}|$ as $t \rightarrow +\infty$ when λ is taken under the Poissonised Plancherel measure pl_t . As in [LuPi, AySi] we rely on the following identity:

$$|\bar{\lambda}| = \frac{e}{2} \sum_{i \in \mathbb{Z}/e\mathbb{Z}} x_i(\lambda)^2 + \sum_{i=0}^{e-1} i x_i(\lambda),$$

where $x_i(\lambda) := c_i(\lambda) - c_{i+1}(\lambda)$. Now the work of [LuPi, AySi] both namely rely of the asymptotics of Hardy–Ramanujan for the partition function $\#\mathcal{P}_n$. For our means, as in [BoSu-a, BoSu-b] we rely heavily on the fact (1.2) that $\mathcal{D}(\lambda)$ is determinantal [BOO] and we then exploit the first order asymptotics of Bessel functions. Our first result is the following (see Proposition 4.13 and Theorem 4.53), where \mathbb{E}_t , Var_t and Cov_t respectively denote the expectation, variance and covariance with respect to pl_t .

Theorem A. *Under the Poissonised Plancherel measure pl_t , as $t \rightarrow +\infty$ we have:*

$$\begin{aligned} \mathbb{E}_t x_i(\lambda) &= O(1), \\ \text{Cov}_t(x_i(\lambda), x_j(\lambda)) &\sim \frac{2\sqrt{t}}{\pi e^2} \left[\cot(j - i + \frac{1}{2}) \frac{\pi}{e} - \cot(j - i - \frac{1}{2}) \frac{\pi}{e} \right], \end{aligned}$$

for all $i, j \in \mathbb{Z}/e\mathbb{Z}$. In particular, we have $\text{Var}_t x_i(\lambda) \sim \frac{4\sqrt{t}}{\pi e^2} \cot \frac{\pi}{2e}$.

We can already deduce the following asymptotics (Proposition 5.1):

$$\mathbb{E}_t |\bar{\lambda}| \sim \frac{2\sqrt{t}}{\pi} \cot \frac{\pi}{2e},$$

as $t \rightarrow +\infty$. To compare with [LuPi, AySi], when λ is taken under the uniform measure on \mathcal{P}_n then $\mathbb{E}|\bar{\lambda}| \sim \frac{e-1}{2\pi} \sqrt{6n}$ as $n \rightarrow +\infty$. We then show that we can apply the central limit theorem [So00-b] with the variables $x_i(\lambda)$ for $i \in \mathbb{Z}/e\mathbb{Z}$ (Theorem 5.5).

Theorem B. *The random vector*

$$e\sqrt{\frac{\pi}{2}} \left(\frac{x_i(\lambda)}{t^{1/4}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

converges in distribution as $t \rightarrow +\infty$ to the centred normal distribution with covariance matrix $(\cot(j - i + \frac{1}{2})\frac{\pi}{e} - \cot(j - i - \frac{1}{2})\frac{\pi}{e})_{i,j \in \mathbb{Z}/e\mathbb{Z}}$.

In particular, as $t \rightarrow +\infty$ the variables $t^{-1/4}x_i(\lambda)$ and $t^{-1/4}x_j(\lambda)$ for $i \neq j$ are not independent. We show in fact that the variables $\sum_{i=0}^{e-1} \zeta^i t^{-1/4}x_i(\lambda)$ are (Gaussian and) mutually independent as $t \rightarrow +\infty$ for $\zeta \neq 1$ a complex e -th root of unity. We then deduce our main result, Theorem 5.9.

Theorem C. *Under the Poissonised Plancherel measure pl_t , as $t \rightarrow +\infty$ the rescaled size $\frac{\pi}{4\sqrt{t}}|\bar{\lambda}|$ of the e -core of λ converges in distribution to a sum of mutually independent Gamma distributions $\Gamma(\frac{1}{2}, \sin \frac{k\pi}{e})$ for $k \in \{1, \dots, e-1\}$.*

The difference between our main theorem and the result of [LuPi, AySi] is twofold:

- our convergence is for the Poissonised Plancherel measure pl_t on \mathcal{P} as $t \rightarrow +\infty$, and the not for the uniform measure on \mathcal{P}_n as $n \rightarrow +\infty$,
- the Gamma distribution $\Gamma(\frac{e-1}{2}, \sqrt{6}) = \sum_{k=1}^{e-1} \Gamma(\frac{1}{2}, \sqrt{6})$ is for us split into $e-1$ summands with different scale parameters.

Note that the order \sqrt{t} of $|\bar{\lambda}|$ corresponds to the order \sqrt{n} of [LuPi, AySi], at it was already noted for the expectation. As in [BOO, Jo] it should also be possible to state Theorem C for the Plancherel measure Pl_n for $n \rightarrow +\infty$. This process, the so-called *de-Poissonisation*, will not be considered here. Finally, as it is mentioned in [CoLe, So00-b], the convergence in the central limit theorem holds in fact in moments, so that we deduce

$$\text{Var}_t|\bar{\lambda}| \sim \frac{4et}{\pi^2},$$

as $t \rightarrow +\infty$ (Corollary 5.12).

We now give the outline of the paper. In Section 2 we recall the result from Borodin–Okounkov–Olshanski saying that $\mathcal{D}(\lambda)$ is a determinantal point process under pl_t . In Section 3 we recall the notion of e -core of a partition. Namely, in Definition 3.7 we define the variables $x_i(\lambda) := c_i(\lambda) - c_{i+1}(\lambda)$, which we relate in (3.12) to $|\bar{\lambda}|$ by a quadratic polynomial and in Lemma 3.13 to $\mathcal{D}(\lambda)$ in a view to applying the central limit theorem of Costin–Lebowitz and Soshnikov for determinantal point process. Section 4 is devoted to expectation and covariance calculations. In §4.1 we show how we will apply the central limit theorem to $x_i(\lambda)$. In §4.2 we prove that the expectation $\mathbb{E}_t x_i(\lambda)$ of $x_i(\lambda)$ under the Poissonised Plancherel measure pl_t is bounded (Proposition 4.13). Subsection §4.4 is the technical heart of the paper. We compute the first order asymptotics of the covariance $\text{Cov}_t(x_i(\lambda), x_j(\lambda))$ under pl_t for $i \neq j \in \mathbb{Z}/e\mathbb{Z}$ (Theorem 4.53). We deduce in §4.5 the first order asymptotics of the variance $\text{Var}_t x_i(\lambda)$ under pl_t (Corollary 4.55). In Section 5 we apply the preceding results to study the asymptotics of $|\bar{\lambda}|$ when λ is taken under the Poissonised Plancherel measure pl_t as $t \rightarrow +\infty$. We first obtain in Proposition 5.1 the asymptotics of the expectation $\mathbb{E}_t|\bar{\lambda}|$. Using the central limit theorem of Costin–Lebowitz and Soshnikov, we obtain in Theorem 5.5 that the random vector $(x_i(\lambda))_{i \in \mathbb{Z}/e\mathbb{Z}}$ converges in distribution to a certain centred normal vector. Then using a contour integration to compute the eigenvalues of the covariance matrix, we deduce our main result, Theorem 5.9, which says that $|\bar{\lambda}|/\sqrt{t}$ converges in distribution under pl_t as $t \rightarrow +\infty$ to a sum of squares of mutually independent centred Gaussian variables, that is, a sum of mutually independent Gamma variables of shape $\frac{1}{2}$. Since the convergence in the central limit theorem holds in moments, we deduce in Corollary 5.12 the asymptotics of the variance $\text{Var}_t|\bar{\lambda}|$. Finally, in Section 6 we apply the preceding results to compute the first order asymptotics of $c_i(\lambda)$ under the Poissonised Plancherel measure pl_t (Proposition 6.1). It turns out that we can recover this result in the (non-Poissonised) Plancherel setting (Proposition 6.4) using the law of large numbers of Logan–Shepp and Kerov–Vershik.

Acknowledgements The author is thankful to Jean-Christophe Breton, Valentin Féray, Cédric Lecouvey and Pierre-Loïc Méliot for many useful discussions.

2 Setting

Let $n \in \mathbb{Z}_{\geq 0}$.


2.1 Partitions

Partitions A *partition* of n is a non-increasing sequence $\lambda = (\lambda_1 \geq \dots \geq \lambda_h > 0)$ of positive integers with sum $|\lambda| := \sum_{a=1}^h \lambda_a = n$, in which case we write $h =: h(\lambda) \geq 0$. We denote by \mathcal{P}_n the set of partitions of n and $\mathcal{P} := \sqcup_{n \geq 0} \mathcal{P}_n$. The *conjugate* of λ , denoted by λ' , is the partition given by $\lambda'_a = \#\{b \geq 1 : \lambda_b \geq a\}$ for all $a \in \{1, \dots, \lambda_1\}$.

Young diagrams The *Young diagram* of $\lambda \in \mathcal{P}_n$ is the subset $\mathcal{Y}(\lambda)$ of $\mathbb{Z}_{\geq 0}^2$ given by

$$\mathcal{Y}(\lambda) := \{(a, b) \in \mathbb{Z}_{\geq 0}^2 : 0 \leq a < h(\lambda) \text{ and } 0 \leq b < \lambda_a\}.$$

We identify the node $(a, b) \in \mathcal{Y}(\lambda)$ with the (square) box with opposite vertices (a, b) and $(a + 1, b + 1)$.

Example 2.1. The Young diagram associated with the partition $(4, 3, 2, 2)$ is .

The Young diagram of λ' is obtained by flipping $\mathcal{Y}(\lambda)$ with respect to the diagonal, in other words, we have $(a, b) \in \mathcal{Y}(\lambda') \iff (b, a) \in \mathcal{Y}(\lambda)$.

A *standard tableau of shape* λ is a bijection $\mathfrak{t} : \mathcal{Y}(\lambda) \rightarrow \{1, \dots, n\}$ such that \mathfrak{t} increases along the rows and down the columns, in other words for $(a, b) \in \mathcal{Y}(\lambda)$ we have $\mathfrak{t}(a, b) > \mathfrak{t}(a + 1, b)$ if $(a + 1, b) \in \mathcal{Y}(\lambda)$ and $\mathfrak{t}(a, b) > \mathfrak{t}(a, b + 1)$ if $(a, b + 1) \in \mathcal{Y}(\lambda)$. We denote by $\text{Std}(\lambda)$ the set of standard Young tableaux of shape λ . Note that $\text{Std}(\lambda)$ and $\text{Std}(\lambda')$ are naturally in bijection.

Russian convention Rotating the Young diagram of λ' by an angle of $\frac{3\pi}{4}$ and applying a linear homothety of ratio $\sqrt{2}$ gives the *Russian convention* for the Young diagram of λ . We denote by $\omega_\lambda : \mathbb{R} \rightarrow \mathbb{R}$ the upper rim of the resulting diagram, extending ω_λ by $\omega_\lambda(x) := |x|$ outside the diagram. Then ω_λ is a continuous piecewise linear function such that:

- for each $k \in \mathbb{Z}$ we have

$$\omega'_\lambda|_{(k, k+1)} = \pm 1, \tag{2.2}$$

- we have $\omega_\lambda(x) = |x|$ for $|x| \gg 0$ (more precisely, for $x \leq -\lambda'_1$ or $x \geq \lambda_1$),
- we have $\int_{\mathbb{R}} [\omega_\lambda(x) - |x|] dx = 2n$, in particular each box in the Young diagram of λ in the Russian convention has area 2 and semi-diagonal length 1.

An illustration of the construction of ω_λ is given in Figure 2. Note that for any $x \in \mathbb{R}$ we have

$$\omega_{\lambda'}(x) = \omega_\lambda(-x). \tag{2.3}$$

Descent set Yet another way to look at a partition λ is to consider its *descent set*, or *β -set* (in the terminology of [BOO] and [OI], respectively), defined as

$$\mathcal{D}(\lambda) = \{\lambda_a - a : a \geq 1\} \subseteq \mathbb{Z},$$

with $\lambda_a := 0$ if $a > h(\lambda)$. The set $\mathcal{D}(\lambda) \cap \mathbb{Z}_{\geq 0}$ is finite, while $\mathcal{D}(\lambda) \supseteq \mathbb{Z}_{< -h(\lambda)}$. The terminology *descent set* is justified by the following:

$$k \in \mathcal{D}(\lambda) \iff \omega_\lambda(k) > \omega_\lambda(k + 1), \tag{2.4}$$




Figure 2: Russian convention for the Young diagram of $\lambda = (4, 3, 2, 2)$




Figure 3: Descent set $\mathcal{D}(\lambda) = \{3, 1, -1, -2, -5, -6, -7, \dots\}$ for $\lambda = (4, 3, 2, 2)$

for any $k \in \mathbb{Z}$. In particular, using (2.3) we recover the following classical relation:

$$\mathcal{D}(\lambda') = -\mathcal{D}(\lambda)^c - 1. \quad (2.5)$$

An example of descent set is given in Figure 3.

2.2 Plancherel measure

Plancherel measure Recall from the introduction the following identity:

$$n! = \sum_{\lambda \in \mathcal{P}_n} \#\text{Std}(\lambda)^2. \quad (2.6)$$

Definition 2.7. The *Plancherel measure* on the set \mathcal{P}_n of the partitions of n is given by

$$\text{Pl}_n(\lambda) := \frac{\#\text{Std}(\lambda)^2}{n!},$$

for all $\lambda \in \mathcal{P}_n$.

If λ' denotes the conjugate partition of λ , we have $\#\text{Std}(\lambda') = \#\text{Std}(\lambda)$ thus

$$\text{Pl}_n(\lambda) = \text{Pl}_n(\lambda'). \quad (2.8)$$

Definition 2.9. Let $t > 0$. The *Poissonised Plancherel* measure pl_t on the set \mathcal{P} of all partitions is given for $\lambda \in \mathcal{P}$ by

$$\text{pl}_t(\lambda) := \exp(-t)t^{|\lambda|} \left(\frac{\#\text{Std}(\lambda)}{|\lambda|!} \right)^2.$$

In other words, setting $\text{Pl}_n(\lambda) := 0$ if $|\lambda| \neq n$, we have

$$\text{pl}_t = \exp(-t) \sum_{n \geq 0} \frac{t^n}{n!} \text{Pl}_n.$$

We use the notation \exp for the exponential function since the notation e will be used for an integer later (see §3). It follows from (2.8) that if Λ is any set of partitions and $\Lambda' := \{\lambda' : \lambda \in \Lambda\}$ then

$$\text{pl}_t(\Lambda) = \text{pl}_t(\Lambda'). \quad (2.10)$$

Correlation function

Definition 2.11. The *correlation function* ρ^t is defined for any finite subset $X \subseteq \mathbb{Z}$ by

$$\rho^t(X) := \text{pl}_t\{\lambda \in \mathcal{P} : X \subseteq \mathcal{D}(\lambda)\}.$$

In the sequel, when we write $X = \{x_1, \dots, x_s\} \subseteq \mathbb{Z}$ then it will always be understood that $x_a \neq x_b$ if $a \neq b$. In particular, we write

$$\rho^t(x_1, \dots, x_s) := \rho^t(X).$$

Note that (2.10) gives that we also have

$$\rho^t(X) = \text{pl}_t\{\lambda \in \mathcal{P} : X \subseteq \mathcal{D}(\lambda')\}. \quad (2.12)$$

Lemma 2.13. For any $k \in \mathbb{Z}$ we have

$$1 - \rho^t(k) = \rho^t(-k - 1).$$

Proof. By (2.5) we have

$$\begin{aligned} 1 - \rho^t(k) &= 1 - \text{pl}_t(k \in \mathcal{D}(\lambda)) \\ &= \text{pl}_t(k \in \mathcal{D}(\lambda)^c) \\ &= \text{pl}_t(-k - 1 \in -\mathcal{D}(\lambda)^c - 1) \\ &= \text{pl}_t(-k - 1 \in \mathcal{D}(\lambda')), \end{aligned}$$

and we conclude by (2.12). \square

Discrete Bessel kernel For any $x \in \mathbb{R}$, we denote by J_x the Bessel function of order x (see [OLBC, §10]) and we define $J_x^t := J_x(2\sqrt{t})$. We will also write $J_x^{t,2} := (J_x^t)^2$.

Definition 2.14 ([BOO, Jo]). The *discrete Bessel kernel* is defined for any $x, y \in \mathbb{R}$ by

$$\mathbf{J}^t(x, y) := \sqrt{t} \frac{J_x^t J_{y+1}^t - J_{x+1}^t J_y^t}{x - y}.$$

Note that the diagonal values $\mathbf{J}^t(x, x)$ are well-defined since J_x^t is an entire function of x . For instance, we have

$$\mathbf{J}^t(x, x) = \sqrt{t} [L_x^t J_{x+1}^t - J_x^t L_{x+1}^t],$$

where $L_x^t = \frac{\partial}{\partial x} J_x^t$ (see [Jo, (3.27)]). The next result shows that, under pl_t , the point process $\mathcal{D}(\lambda)$ is determinantal.

Theorem 2.15 ([BOO, Theorem 2]). *For any $\{x_1, \dots, x_s\} \subseteq \mathbb{Z}$ we have*

$$\rho^t(x_1, \dots, x_s) = \det[J^t(x_a, x_b)]_{1 \leq a, b \leq s}.$$

Remark 2.16. Under Pl_n , the process $\mathcal{D}(\lambda)$ is a priori not determinantal, however [BOO] proved namely that, under some assumptions,

$$\text{Pl}_n\{\lambda \in \mathcal{P}_n : x_1(n), \dots, x_s(n) \in \mathcal{D}(\lambda)\} \xrightarrow{n \rightarrow +\infty} \det[S(d_{ab}, y)]_{1 \leq a, b \leq s},$$

where S is the *discrete sine kernel*, where y and $(d_{ab})_{ab}$ are determined by $(x_a(n))_{a,n}$.

We now recall some properties of the kernel J^t .

Lemma 2.17. *The kernel J^t is symmetric: for all $x, y \in \mathbb{Z}$ we have $J^t(x, y) = J^t(y, x)$.*

Lemma 2.18 ([BOO, Proposition 2.9], [Jo, (3.31)]). *For all $x, y \in \mathbb{Z}$ we have*

$$J^t(x, y) = \sum_{s \geq 1} J_{x+s}^t J_{y+s}^t.$$

In particular,

$$J^t(x, x) = \sum_{s \geq 1} J_{x+s}^{t,2}.$$

3 Cores

Let $e \in \mathbb{Z}_{\geq 2}$. We will identify $\mathbb{Z}/e\mathbb{Z}$ and $\{0, \dots, e-1\}$ in the natural way. Let $\lambda \in \mathcal{P}_n$ be a partition of n .

Definition 3.1. For any $i \in \mathbb{Z}/e\mathbb{Z}$, we define $c_i(\lambda) \in \mathbb{Z}_{\geq 0}$ to be the number of nodes $(a, b) \in \mathcal{Y}(\lambda)$ of the Young diagram of λ such that $b - a = i \pmod{e}$.

Such a node $(a, b) \in \mathcal{Y}(\lambda)$ with $b - a = i \pmod{e}$ is an *i-node* of $\mathcal{Y}(\lambda)$. Note that

$$\sum_{i \in \mathbb{Z}/e\mathbb{Z}} c_i(\lambda) = |\lambda|. \quad (3.2)$$

The next result is immediate from the definition of ω_λ .

Lemma 3.3. *For any $i \in \mathbb{Z}/e\mathbb{Z}$ we have*

$$c_i(\lambda) = \frac{1}{2} \sum_{\substack{m \in \mathbb{Z} \\ m \equiv i \pmod{e}}} \omega_\lambda(m) - |m| = \frac{1}{2} \sum_{k \in \mathbb{Z}} \omega_\lambda(i + ke) - |i + ke|.$$

Proof. First note that the sum is finite. Recall that, in the Russian convention, each box has diagonal 2. Then for any $m \in \mathbb{Z}$, the number $\frac{1}{2}[\omega_\lambda(m) - |m|]$ (an integer and is) the number of nodes $(a, b) \in \mathcal{Y}(\lambda)$ such that $b - a = m$. \square

If $\lambda, \mu \in \mathcal{P}$ are such that $c_i(\lambda) = c_i(\mu)$ for all $i \in \mathbb{Z}/e\mathbb{Z}$ then we do not necessarily have $\lambda = \mu$ (unless if we are in the, excluded, case $e = 0$). However, we know that λ and μ share the same *e-core*, notion that we now briefly explain (see, for instance, [JaKe, §2.7] for more details). Let $\lambda \in \mathcal{P}$. Given $(a, b) \in \mathcal{Y}(\lambda)$, the *hook* $h_{(a,b)}(\lambda)$ of (a, b) is the set of nodes in $\mathcal{Y}(\lambda)$ that are either directly below or directly to the right of (a, b) , that is:

$$h_{(a,b)}(\lambda) := \{(a', b') \in \mathcal{Y}(\lambda) : a' = a, b' \geq b \text{ or } a' \geq a, b' = b\}.$$

We say that $h_{(a,b)}$ is a *h-hook* if $\#h_{(a,b)}(\lambda) = h$. Removing $h_{(a,b)}(\lambda)$ from $\mathcal{Y}(\lambda)$ and then patching the two disconnected parts together by moving the nodes that were below $h_{(a,b)}(\lambda)$ in north-west direction gives another Young diagram $\lambda_{(a,b)}$. Another way of seeing the construction of $\lambda_{(a,b)}$ is to remove from $\mathcal{Y}(\lambda)$ the *h-rim hook* associated with (a, b) , that is,

the set of nodes of $\mathcal{Y}(\lambda)$ that are both on the rim and between the southmost and eastmost nodes of $h_{(a,b)}(\lambda)$.

The e -core of λ , denoted by $\bar{\lambda}$, is constructed from $\mathcal{Y}(\lambda)$ by successively removing as many e -rim hooks as possible. This construction does not depend on the order we have removed e -rim hooks for the following reason. The number $w_e(\lambda)$ of e -rim hooks that we can remove is the e -weight of λ , in particular, we have the relation

$$|\lambda| = |\bar{\lambda}| + ew_e(\lambda). \quad (3.4)$$

More precisely, since any e -rim hook has exactly one i -node for any $i \in \mathbb{Z}/e\mathbb{Z}$ (see, for instance, [Ros, Remark 2.3]) we deduce that

$$c_i(\lambda) = c_i(\bar{\lambda}) + w_e(\lambda), \quad (3.5)$$

recovering (3.4) by (3.2).

The e -core $\bar{\lambda}$ of λ is the partition whose descent set is obtained from $\mathcal{D}(\lambda)$ by replacing each element $b \in \mathcal{D}(\lambda)$ by $b - e$, provided that $b - e$ is not already in $\mathcal{D}(\lambda)$. We repeat the procedure until the resulting set D satisfies $b - e \in D$ for all $b \in D$ (see, for instance, [Na], [JaKe, Lemma 2.7.13] or [Ma, 5.26 Lemma]).

Example 3.6. Consider $\lambda := (3, 2, 2, 1) \in \mathcal{P}_8$. We have

$$\mathcal{D}(\lambda) = \{2, 0, -1, -3, -5, -6, \dots\}.$$

- To obtain the 3-core of λ , we replace $-1 \in \mathcal{D}(\lambda)$ by $-1 - 3 = -4$, so that we obtain

$$D_1 := \{2, 0, -3, -4, -5, -6, \dots\}.$$

Now we can replace $2 \in D_1$ by $2 - 3 = -1$, so that we obtain

$$D_2 := \{0, -1, -3, -4, -5, -6, \dots\}.$$

Now for each element $b \in D_2$ we have $b - 3 \in D_2$ thus $D_2 = \mathcal{D}(\bar{\lambda})$ and thus $\bar{\lambda} = (1, 1) \in \mathcal{P}_2$.

- To obtain the 4-core of λ , we can replace $2, 0 \in \mathcal{D}(\lambda)$ by $2 - 4 = -2$ and $0 - 4 = -4$ respectively so that we obtain

$$D := \{-1, -2, -3, -4, -5, -6, \dots\}.$$

Now for each $b \in D$ we have $b - 4 \notin D$ thus $D = \mathcal{D}(\bar{\lambda})$ and thus $\bar{\lambda}$ is the empty partition.

- For all $b \in \mathcal{D}(\lambda)$ we have $b - 5 \in \mathcal{D}(\lambda)$ thus $\mathcal{D}(\bar{\lambda}) = \mathcal{D}(\lambda)$ and thus $\bar{\lambda} = \lambda$. In this case we say that λ is an e -core.

Definition 3.7. For any $i \in \mathbb{Z}/e\mathbb{Z}$ we define $x_i(\lambda) := c_i(\lambda) - c_{i+1}(\lambda) \in \mathbb{Z}$.

Note that

$$\sum_{i \in \mathbb{Z}/e\mathbb{Z}} x_i(\lambda) = 0. \quad (3.8)$$

Remark 3.9. The quantities $x_0(\lambda), \dots, x_{e-1}(\lambda)$ can be read on the e -abacus of λ , see, for instance, [Ros, §2.3]. Moreover, we have a surjective map

$$\begin{aligned} \mathcal{P} &\longrightarrow \mathbb{Z}_{\geq 0} \times \{x \in \mathbb{Z}^e : x_0 + \dots + x_{e-1} = 0\} \\ \lambda &\longmapsto (w_e(\lambda), (x_i(\lambda))_{0 \leq i < e}) \end{aligned}$$

The above discussion says that if $\lambda, \mu \in \mathcal{P}_n$ have the same image under this map and if $w_e(\lambda) = w_e(\mu) = 0$ (that is, if both λ and μ are e -cores) then $\lambda = \mu$.

The analogue of (3.2) is the following (non-trivial) equality, where $\|\cdot\|$ denotes the Euclidean norm:

$$c_0(\bar{\lambda}) = \frac{1}{2} \|x(\lambda)\|^2. \quad (3.10)$$

Indeed, by [Fa, Proposition 2.1] (see also [Ros, Proposition 2.14] or also [GKS, Bijection 2]) we have $c_0(\bar{\lambda}) = \frac{1}{2} \|x(\bar{\lambda})\|^2$, but $x(\bar{\lambda}) = x(\lambda)$ by (3.5). Hence, for any $i \in \{1, \dots, e-1\}$ we have

$$c_i(\bar{\lambda}) = \frac{1}{2} \|x(\lambda)\|^2 - x_0(\lambda) - \dots - x_{i-1}(\lambda). \quad (3.11)$$

Note the following consequence of (3.2), (3.8), (3.10) and (3.11):

$$|\bar{\lambda}| = \frac{e}{2} \|x(\lambda)\|^2 + \sum_{i=0}^{e-1} ix_i(\lambda). \quad (3.12)$$

One of the aim of this paper is to study the asymptotic behaviour of $|\bar{\lambda}|$ under the Plancherel measure. By (3.12), for any $i \in \mathbb{Z}/e\mathbb{Z}$ we have

$$x_i(\lambda) = O\left(\sqrt{|\bar{\lambda}|}\right).$$

We will see how to refine this asymptotics when λ is taken under the (Poissonised) Plancherel measure, by using Theorem 2.15 (see Theorem 5.9).

Lemma 3.13. *For any $i \in \{0, \dots, e-1\}$ we have*

$$\begin{aligned} x_i(\lambda) &= \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \#(e\mathbb{Z}_{< 0} + i) \cap \mathcal{D}(\lambda)^c \\ &= \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \#(e\mathbb{Z}_{\geq 1} - i - 1) \cap \mathcal{D}(\lambda') \\ &= \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \#(e\mathbb{Z}_{\geq 0} + i^*) \cap \mathcal{D}(\lambda'), \end{aligned}$$

with $i^* := e - 1 - i \in \{0, \dots, e-1\}$.

Proof. Write $\delta_k := \omega_\lambda(ke + i) - \omega_\lambda(ke + i + 1)$. Recalling (2.2) and (2.4), we have:

$$\delta_k = \begin{cases} 1, & \text{if } ke + i \in \mathcal{D}(\lambda), \\ -1, & \text{otherwise.} \end{cases} \quad (3.14)$$

We have, using Lemma 3.3 and the fact that $i \in \{0, \dots, e-1\}$,

$$\begin{aligned} 2x_i(\lambda) &= 2(c_i(\lambda) - c_{i+1}(\lambda)) \\ &= \sum_{k \in \mathbb{Z}} \left(\omega_\lambda(ke + i) - |ke + i| \right) - \left(\omega_\lambda(ke + i + 1) - |ke + i + 1| \right) \\ &= \sum_{k < 0} \delta_k - |ke + i| + |ke + i + 1| + \sum_{k \geq 0} \delta_k - |ke + i| + |ke + i + 1| \\ &= \sum_{k < 0} \delta_k + (ke + i) - (ke + i + 1) + \sum_{k \geq 0} \delta_k - (ke + i) + (ke + i + 1) \\ &= \sum_{k < 0} (\delta_k - 1) + \sum_{k \geq 0} (\delta_k + 1). \end{aligned}$$

By (3.14), we have

$$\delta_k + 1 = \begin{cases} 2, & \text{if } ke + i \in \mathcal{D}(\lambda), \\ 0, & \text{otherwise,} \end{cases}$$

thus $\frac{1}{2} \sum_{k \geq 0} (\delta_k + 1) = \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda)$. Similarly, we have

$$\delta_k - 1 = \begin{cases} -2, & \text{if } ke + i \in \mathcal{D}(\lambda)^c, \\ 0, & \text{otherwise,} \end{cases}$$

thus $\frac{1}{2} \sum_{k < 0} (\delta_k - 1) = -\#(e\mathbb{Z}_{< 0} + i) \cap \mathcal{D}(\lambda)^c$. We deduce the first equality of the lemma. By (2.5) we have $\#(e\mathbb{Z}_{< 0} + i) \cap \mathcal{D}(\lambda)^c = \#(e\mathbb{Z}_{\geq 1} - i - 1) \cap \mathcal{D}(\lambda')$ thus we deduce the last two announced equalities. \square

4 Covariance calculations

Our aim is to evaluate $x_i(\lambda)$ as $|\lambda| \rightarrow \infty$ under the Poissonised Plancherel measure.

4.1 Central limit theorem

With a view to Lemma 3.13, we will use the following central limit theorem.

Theorem 4.1 (Costin–Lebowitz [CoLe], Soshnikov [So00-b]). *Let $(\mathcal{D}_t)_t$ be a sequence of determinantal point processes on a locally compact Polish space. Assume that each \mathcal{D}_t is associated with a Hermitian non-negative locally trace class operator in L^2 . Let $\{I_t^{(0)}, \dots, I_t^{(e-1)}\}_t$ be a sequence of measurable sets, disjoint for any fixed t . We denote by \mathbb{E}_t , Var_t and Cov_t the expectation, variance and covariance with respect to the probability distribution of the random process \mathcal{D}_t . Let $\#_t^{(i)}$ be the random variable given by the number of points of \mathcal{D}_t inside $I_t^{(i)}$. If $\text{Var}_t \#_t^{(i)} \rightarrow +\infty$ as $t \rightarrow +\infty$ for all i and if for all i, j*

$$\frac{\text{Cov}_t \left(\#_t^{(i)}, \#_t^{(j)} \right)}{\sqrt{\text{Var}_t \#_t^{(i)} \text{Var}_t \#_t^{(j)}}} \xrightarrow{t \rightarrow +\infty} b_{ij},$$

for some $b_{ij} \in \mathbb{R}$ then the vector

$$\left(\frac{\#_t^{(i)} - \mathbb{E}_t \#_t^{(i)}}{\sqrt{\text{Var}_t \#_t^{(i)}}} \right)_{0 \leq i < e}$$

converges in distribution to the k -dimensional centred normal vector with covariance matrix $(b_{ij})_{0 \leq i, j < e}$.

Remark 4.2. Theorem 4.1 in [So00-b] has only been explicitly stated for some particular determinantal processes. However, the proof is valid in the general case (see [So02, Remark 2] and [So00-a, §4]). See also [BKPV].

Remark 4.3. The discrete Bessel kernel J^t satisfies the assumptions of Theorem 4.1, see [BOO, Corollary 2.10]. In particular, we will be able to use Theorem 4.1 in our setting.

Let $i \in \{0, \dots, e-1\}$. Recall from Lemma 3.13 that $x_i(\lambda) = \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \#(e\mathbb{Z}_{< 0} + i) \cap \mathcal{D}(\lambda)^c$. We define

$$\hat{x}_i(\lambda) := \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \#(e\llbracket -t^2, -1 \rrbracket + i) \cap \mathcal{D}(\lambda)^c.$$

We have

$$\hat{x}_i(\lambda) - x_i(\lambda) = \#(e\mathbb{Z}_{< -t^2} + i) \cap \mathcal{D}(\lambda)^c, \quad (4.4)$$

and

$$\begin{aligned} \hat{x}_i(\lambda) &= \#(e\mathbb{Z}_{\geq 0} + i) \cap \mathcal{D}(\lambda) - \left[\#(e\llbracket -t^2, -1 \rrbracket + i) - \#(e\llbracket -t^2, -1 \rrbracket + i) \cap \mathcal{D}(\lambda) \right] \\ &= \#(e\mathbb{Z}_{\geq -t^2} + i) \cap \mathcal{D}(\lambda) - t^2. \end{aligned} \quad (4.5)$$

Lemma 4.6. *For any $0 \leq i < e$ and $r \geq 1$ we have*

$$\mathbb{E}_t |\hat{x}_i(\lambda) - x_i(\lambda)|^r = o(t^{-t}),$$

as $t \rightarrow +\infty$. In particular, the variable $\hat{x}_i(\lambda) - x_i(\lambda)$ converges to 0 in distribution under pl_t as $t \rightarrow +\infty$.

Proof. Recall from (2.5) that $\mathcal{D}(\lambda)^c = -\mathcal{D}(\lambda') - 1$, so that

$$\begin{aligned} \hat{x}_i(\lambda) - x_i(\lambda) &= \#(e\mathbb{Z}_{< -t^2} + i) \cap \mathcal{D}(\lambda)^c \\ &= \#(e\mathbb{Z}_{< -t^2} + i) \cap (-\mathcal{D}(\lambda') - 1) \\ &= \#(e\mathbb{Z}_{> t^2} - i - 1) \cap \mathcal{D}(\lambda') \\ &= \#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda'), \end{aligned}$$

where $i^* = e - 1 - i \in \{0, \dots, e - 1\}$. Hence, by (2.10) it suffices to prove the result for the random variable $\#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda)$.

We first prove that

$$\#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda) \leq \max(T - t^2, 0), \quad (4.7)$$

where $T := |\lambda|$ is (by definition) a Poisson random variable with parameter t . If $\#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda) = n$ with $n \geq 1$ then $\mathcal{D}(\lambda)$ contains an element of the form $ek + i^*$ for $k \geq t^2 + n - 1$. Hence, we have, recalling that $i^* \geq 0$, $n \geq 1$ and $e \geq 2$,

$$\lambda_1 - 1 = \max \mathcal{D}(\lambda) \geq e(t^2 + n - 1) + i^* \geq t^2 + n - 1.$$

Thus, we obtain that

$$T \geq \lambda_1 \geq t^2 + n,$$

and thus $T - t^2 \geq n$ thus $n \leq \max(T - t^2, 0)$ as desired. This inequality is satisfied when $n = 0$ as well, proving (4.7).

We deduce from (4.7) that, using Cauchy–Schwarz inequality,

$$\begin{aligned} \mathbb{E}_t |\#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda)|^r &\leq \mathbb{E}_t [\max(T - t^2, 0)^r] \\ &\leq \mathbb{E}_t [\mathbf{1}_{T \geq t^2} (T - t^2)^r] \\ &\leq \sqrt{\mathbb{E}_t \mathbf{1}_{T \geq t^2}} \sqrt{\mathbb{E}_t (T - t^2)^{2r}} \\ &= \sqrt{\mathbb{P}_t(T \geq t^2)} \sqrt{\mathbb{E}_t (T - t^2)^{2r}}. \end{aligned} \quad (4.8)$$

The second term in the right-hand side is the square root of a polynomial $P_r(t)$ in t since $\mathbb{E}_t T^k$ is a polynomial in t for any $k \geq 0$ (we have $\mathbb{E}_t T^k = \sum_{a=0}^k t^a S_{k,a}$ where $S_{k,a}$ are the Stirling numbers of the second kind, see for instance [Ri]). For the first one, we use the following Chernoff-type bound, valid for $x > t$ (see [MiUp, Theorem 5.4]):

$$\mathbb{P}_t(T \geq x) \leq \frac{\exp(-t) \exp(x) t^x}{x^x} = \exp(x(1 + \ln t) - x \ln x - t).$$

For $t > 1$ and $x = t^2$ we obtain

$$\begin{aligned} \mathbb{P}_t(T \geq t^2) &\leq \exp(t^2(1 + \ln t) - 2t^2 \ln t - t) \\ &\leq \exp(-t^2 \ln t + t^2 - t). \end{aligned}$$

Hence, by (4.8) we deduce that

$$t^t \mathbb{E}_t |\#(e\mathbb{Z}_{\geq t^2} + i^*) \cap \mathcal{D}(\lambda)|^r \leq \sqrt{P_r(t)} \exp\left(\frac{1}{2}(-t^2 \ln t + t^2 - t) + t \ln t\right) \xrightarrow{t \rightarrow +\infty} 0,$$

whence the result. \square

We now want to apply Theorem 4.1 with $\hat{x}_i(\lambda)$. We thus have to compute the asymptotics of the covariances. We will make the calculations for $x_i(\lambda)$ and then see that it gives the desired result.

4.2 Expectation

Lemma 4.9. *We have $\sum_{m=0}^{+\infty} \rho^t(m) \leq t$.*

Proof. Let $m \geq 0$. If $m \in \mathcal{D}(\lambda)$ for $\lambda \in \mathcal{P}$ then $m = \lambda_a - a$ for some $a \geq 1$ thus $|\lambda| \geq \lambda_a = m + a > m$. We deduce that

$$\begin{aligned} \sum_{m=0}^{+\infty} \rho^t(m) &\leq \sum_{m=0}^{+\infty} \text{pl}_t(m \in \mathcal{D}(\lambda)) \\ &\leq \sum_{m=0}^{+\infty} \text{pl}_t(|\lambda| > m) = \mathbb{E}_t |\lambda|, \end{aligned}$$

since $|\lambda| \in \mathbb{Z}_{\geq 0}$. We thus obtain the desired inequality since $|\lambda|$ is a Poisson variable with parameter t under pl_t . \square

Recall that $J_n^t = J_n(2\sqrt{t})$.

Proposition 4.10. *Under pl_t we have,*

$$\mathbb{E}_t x_i(\lambda) = \sum_{k \geq 0} \rho^t(ek + i) - \sum_{k \geq 1} \rho^t(ek - i - 1) = \sum_{s \geq i+1} J_s^{t,2} - \sum_{s=-i}^i \sum_{k=1}^{\infty} J_{ek+s}^{t,2}.$$

Proof. By Lemma 3.13 and (2.12) we have, where $\mathbf{1}_E$ denotes the characteristic function of any set E ,

$$\begin{aligned} \mathbb{E}_t x_i(\lambda) &= \sum_{k \geq 0} \mathbb{E}_t \mathbf{1}_{\mathcal{D}(\lambda)}(ek + i) - \sum_{k \geq 1} \mathbb{E}_t \mathbf{1}_{\mathcal{D}(\lambda')}(ek - i - 1) \\ &= \sum_{k \geq 0} \text{pl}_t(ek + i \in \mathcal{D}(\lambda)) - \sum_{k \geq 1} \text{pl}_t(ek - i - 1 \in \mathcal{D}(\lambda')) \\ &= \sum_{k \geq 0} \rho^t(ek + i) - \sum_{k \geq 1} \rho^t(ek - i - 1). \end{aligned}$$

Note that each sum above is finite by Lemma 4.9. By Theorem 2.15 we obtain

$$\mathbb{E}_t x_i(\lambda) = \sum_{k \geq 0} J^t(ek + i, ek + i) - \sum_{k \geq 1} J^t(ek - i - 1, ek - i - 1).$$

Using Lemma 2.18 we deduce that, noticing that we have a telescopic sum,

$$\begin{aligned} \mathbb{E}_t x_i(\lambda) &= \sum_{k \geq 0} \sum_{s \geq 1} J_{ek+i+s}^{t,2} - \sum_{k \geq 1} \sum_{s \geq 1} J_{ek-i-1+s}^{t,2} \\ &= \sum_{s \geq 1} J_{i+s}^{t,2} + \sum_{k \geq 1} \sum_{s \geq 1} \left(J_{ek+i+s}^{t,2} - J_{ek-i-1+s}^{t,2} \right) \\ &= \sum_{s \geq 1} J_{i+s}^{t,2} - \sum_{k \geq 1} \sum_{s=-i}^i J_{ek+s}^{t,2}. \end{aligned}$$

□

Let us now recall some facts about the Bessel functions that we will use throughout the paper.

Lemma 4.11. *For any $n \in \mathbb{Z}$ and $x \in \mathbb{R}$ we have:*

$$J_n(x) \in \mathbb{R}, \tag{4.12a}$$

$$J_{-n} = (-1)^n J_n, \tag{4.12b}$$

$$|J_n(x)| \leq 1, \tag{4.12c}$$

$$J_n(x)^2 = \frac{2}{\pi} \int_0^{\pi/2} J_{2n}(2x \cos \theta) d\theta, \tag{4.12d}$$

$$J_n(x)^2 = \frac{2}{\pi} \int_0^{\pi/2} J_0(2x \sin \theta) \cos(2n\theta) d\theta, \tag{4.12e}$$

$$\sum_{m \in \mathbb{Z}} J_m(x) z^m = \exp \left[\frac{1}{2} x (z - z^{-1}) \right], \quad z \in \mathbb{C}^*, \tag{4.12f}$$

$$2J'_n = J_{n-1} - J_{n+1}. \tag{4.12g}$$

Moreover, if $n \in \mathbb{Z}_{\geq 0}$ then

$$|J_n(x)| \leq \frac{|x|^n}{2^n n!}, \tag{4.12h}$$

and finally

$$J_0^2 + 2 \sum_{s=1}^{\infty} J_s^2 = 1, \quad (4.12i)$$

$$J_0 + 2 \sum_{s=1}^{+\infty} J_{2s} = 1. \quad (4.12j)$$

Proof. In [OLBC]: for (4.12a) see (10.2.2), for (4.12b) see (10.4.1), for (4.12c) see (10.14.1), for (4.12d) see (10.9.26) and (4.12b), for (4.12e) see (10.22.16) and (4.12b), for (4.12f) see (10.10.1), for (4.12g) see (10.6.1), for (4.12h) see (10.14.4), for (4.12i) see (10.23.3), for (4.12j) see (10.12.4). \square

We will now prove the following result.

Proposition 4.13. *The quantity $\mathbb{E}_t x_i(\lambda)$ is bounded for $t \in \mathbb{R}$.*

By (4.12c) and (4.12i), we already know that the first sum in the expression of $\mathbb{E}_t x_i(\lambda)$ in Proposition 4.10:

$$\sum_{s \geq 1} J_{i+s}^{t,2},$$

is bounded for $t \in \mathbb{R}$. We will now prove that the second sum:

$$\sum_{k \geq 1} \sum_{s=-i}^i J_{ek+s}^{t,2},$$

is bounded, and this will conclude the proof of Proposition 4.13. To that extent, it suffices to prove that for any $s \in \mathbb{Z}$ the sum

$$A_{e,s}(x) := \sum_{k \in \mathbb{Z}} J_{ek+s}^2(x),$$

is bounded for $x \in \mathbb{R}$. By (4.12d), it suffices to prove that

$$B_{e,s}(x) := \sum_{k \in \mathbb{Z}} J_{2ek+2s}(x),$$

is bounded for $x \in \mathbb{R}$. Note that (4.12b) ensures that we can permute the sum and the integral signs. The next result generalises the standard equalities [OLBC, 10.12.3] expressing $\cos(z \cos \theta)$ (resp. $\sin(z \cos \theta)$) as a series involving $J_{2k}(z)$ and $\cos(2k\theta)$ (resp. $J_{2k+1}(z)$ and $\sin((2k+1)\theta)$). To avoid confusion with our $i \in \mathbb{Z}/e\mathbb{Z}$, we denote by \mathbf{i} the complex unit.

Lemma 4.14. *For any $k \in \mathbb{Z}$, $x, t \in \mathbb{R}$, and $e \in \mathbb{Z}_{\geq 1}$ we have, with $\Omega_e := \left\{ \frac{2\ell\pi}{e} : -\left\lfloor \frac{e}{2} \right\rfloor \leq \ell \leq \left\lfloor \frac{e}{2} \right\rfloor \right\}$,*

$$\sum_{m \in e\mathbb{Z}+k} J_m(x) \exp(\mathbf{i}mt) = \frac{1}{e} \sum_{\omega \in \Omega_e} \exp \mathbf{i}[-k\omega + x \sin(\omega + t)].$$

Proof. Let $\mu_e \subseteq \mathbb{C}^*$ be the subgroup of e -th roots of unity. For any $u \in \mathbb{C}^*$ and $\zeta \in \mu_e$, by (4.12f) we have

$$(u\zeta)^{-k} \exp \frac{1}{2} x (u\zeta - (u\zeta)^{-1}) = \sum_{m \in \mathbb{Z}} J_m(x) (u\zeta)^{m-k}.$$

Using the identity

$$\sum_{\zeta \in \mu_e} \zeta^{m-k} = \begin{cases} e, & \text{if } e \mid m-k, \\ 0, & \text{otherwise,} \end{cases} \quad (4.15)$$

we obtain

$$\sum_{\zeta \in \mu_e} (u\zeta)^{-k} \exp \frac{1}{2} x (u\zeta - (u\zeta)^{-1}) = \sum_{m \in e\mathbb{Z}+k} e J_m(x) u^{m-k}$$

and thus

$$\sum_{m \in e\mathbb{Z}+k} J_m(x) u^m = \frac{1}{e} \sum_{\zeta \in \mu_e} \zeta^{-k} \exp \frac{1}{2} x (u\zeta - (u\zeta)^{-1}).$$

We deduce that

$$\begin{aligned} \sum_{m \in e\mathbb{Z}+k} J_m(x) \exp(\mathbf{i}mt) &= \frac{1}{e} \sum_{\omega \in \Omega_e} \exp(-\mathbf{i}k\omega) \exp \frac{1}{2} x \left[\exp \mathbf{i}(t + \omega) - \exp(-\mathbf{i}(t + \omega)) \right] \\ &= \frac{1}{e} \sum_{\omega \in \Omega_e} \exp(-\mathbf{i}k\omega) \exp[\mathbf{i}x \sin(t + \omega)], \end{aligned}$$

which gives the announced formula. \square

It follows immediately from Lemma 4.14 applied with $t = 0$ that $B_{e,s}(x)$ is bounded for $x \in \mathbb{R}$ and thus, by the preceding discussion, this concludes the proof of Proposition 4.13.

4.3 Covariance

We give here an expression for $\text{Cov}_t(x_i(\lambda), x_j(\lambda))$ for $i \neq j$.

Lemma 4.16. *For any $y \neq x$ we have*

$$\text{Cov}_t(\mathbf{1}_{\mathcal{D}(\lambda)}(x), \mathbf{1}_{\mathcal{D}(\lambda)}(y)) = -\mathbf{J}^t(x, y)^2.$$

Proof. Using Theorem 2.15, a direct computation gives

$$\begin{aligned} \text{Cov}_t(\mathbf{1}_{\mathcal{D}(\lambda)}(x), \mathbf{1}_{\mathcal{D}(\lambda)}(y)) &= \mathbb{E}_t \mathbf{1}_{\mathcal{D}(\lambda)}(x, y) - \mathbb{E}_t \mathbf{1}_{\mathcal{D}(\lambda)}(x) \mathbb{E}_t \mathbf{1}_{\mathcal{D}(\lambda)}(y) \\ &= \rho^t(x, y) - \rho^t(x) \rho^t(y) \\ &= (\rho^t(x) \rho^t(y) - \mathbf{J}^t(x, y)^2) - \rho^t(x) \rho^t(y) \\ &= -\mathbf{J}^t(x, y)^2. \end{aligned}$$

\square

Proposition 4.17. *Let $i, j \in \{0, \dots, e-1\}$ with $i \neq j$. We have, under pl_t ,*

$$\text{Cov}_t(x_i(\lambda), x_j(\lambda)) = - \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j} \mathbf{J}^t(m, n)^2.$$

Proof. Recall from Lemma 3.13 that

$$\begin{aligned} x_i(\lambda) &= \sum_{m \in e\mathbb{Z}_{\geq 0}+i} \mathbf{1}_{\mathcal{D}(\lambda)}(m) - \sum_{m \in e\mathbb{Z}_{< 0}+i} (1 - \mathbf{1}_{\mathcal{D}(\lambda)}(m)). \\ &= \sum_{m \in e\mathbb{Z}_{\geq 0}+i} \mathbf{1}_{\mathcal{D}(\lambda)}(m) + \sum_{m \in e\mathbb{Z}_{< 0}+i} (\mathbf{1}_{\mathcal{D}(\lambda)}(m) - 1). \end{aligned}$$

We thus obtain the desired result by Lemma 4.16. \square

4.4 Asymptotics of the covariance

Let $i \neq j \in \mathbb{Z}/e\mathbb{Z}$. Our aim is to compute the asymptotics of $\text{Cov}_t(x_i(\lambda), x_j(\lambda))$. This will require a considerable amount of calculations. For convenience we define

$$\mathcal{J}(m, n)(x) := x \frac{J_m(x) J_{n+1}(x) - J_{m+1}(x) J_n(x)}{m - n}, \quad (4.18)$$

for $x \in \mathbb{R}$ and $m, n \in \mathbb{Z}$ (we will in fact only use the case $m \neq n$). Note that

$$\mathbf{J}^t(m, n) = \frac{1}{2} \mathcal{J}(m, n)(2\sqrt{t}). \quad (4.19)$$

4.4.1 Derivating

Lemma 4.20 ([BOO, Proposition 2.7]). *For any $m, n \in \mathbb{Z}$ we have*

$$\frac{d\mathcal{J}(m, n)}{dx} = J_m J_{n+1} + J_{m+1} J_n.$$

Proof. From Lemma 2.18 and (4.19) we have $\mathcal{J}(m, n) = 2 \sum_{s \geq 1} J_{m+s} J_{n+s}$. By (4.12g) we obtain that

$$\begin{aligned} \frac{d\mathcal{J}(m, n)}{dx} &= 2 \sum_{s \geq 1} (J'_{m+s} J_{n+s} + J_{m+s} J'_{n+s}) \\ &= \sum_{s \geq 1} ([J_{m+s-1} - J_{m+s+1}] J_{n+s} + J_{m+s} [J_{n+s-1} - J_{n+s+1}]) \\ &= \sum_{s \geq 0} J_{m+s} J_{n+s+1} - \sum_{s \geq 1} J_{m+s+1} J_{n+s} + \sum_{s \geq 0} J_{m+s+1} J_{n+s} - \sum_{s \geq 1} J_{m+s} J_{n+s+1} \\ &= J_m J_{n+1} + J_{m+1} J_n, \end{aligned}$$

as desired. Note that we can both derivate inside the sum sign and split the infinite sums thanks to (4.12h). \square

In particular, combining (4.18) and Lemma 4.20 we obtain

$$\frac{d\mathcal{J}(m, n)^2}{dx} = 2x \frac{J_m^2 J_{n+1}^2 - J_{m+1}^2 J_n^2}{m - n}. \quad (4.21)$$

Recalling Proposition 4.17, we are interested in computing the asymptotics of

$$C_{i,j}(x) := \sum_{m \in \mathbb{Z} + i} \sum_{n \in \mathbb{Z} + j} \mathcal{J}(m, n)^2(x), \quad (4.22)$$

as $x \rightarrow +\infty$. We will see that it suffices to compute the asymptotics of $C'_{i,j}(x)$.

Lemma 4.23. *Let $m \neq n \in \mathbb{Z}$. For all $x \in \mathbb{R}$ we have*

$$\left| \frac{d\mathcal{J}(m, n)^2}{dx} \right| \leq \frac{|x|^{2(|m|+|n|+1)+1}}{2^{2(|m|+|n|+1)} (|m|!|n|!)^2} \left(\frac{1}{(|m|+1)^2} + \frac{1}{(|n|+1)^2} \right).$$

Proof. By (4.12b) and (4.21), for all $x \in \mathbb{R}$ we have, since $m \neq n$,

$$\begin{aligned} \left| \frac{d\mathcal{J}(m, n)^2}{dx} \right| &= 2|x| \left| \frac{J_m(x)^2 J_{n+1}(x)^2 - J_{m+1}(x)^2 J_n(x)^2}{m - n} \right| \\ &\leq 2|x| \left(J_m(x)^2 J_{n+1}(x)^2 + J_{m+1}(x)^2 J_n(x)^2 \right) \\ &\leq 2|x| \left(\frac{|x|^{2|m|}}{2^{2|m|} |m|!^2} \frac{|x|^{2(|n|+1)}}{2^{2(|n|+1)} (|n|+1)!^2} + \frac{|x|^{2(|m|+1)}}{2^{2|m|} (|m|+1)!^2} \frac{|x|^{2|n|}}{2^{2|n|} |n|!^2} \right) \quad (4.24) \\ &\leq \frac{|x|^{2(|m|+|n|+1)+1}}{2^{2(|m|+|n|+1)} (|m|!|n|!)^2} \left(\frac{1}{(|n|+1)^2} + \frac{1}{(|m|+1)^2} \right), \end{aligned}$$

as announced. \square

Note that $\mathcal{J}(m, n)(0) = 0$ for all $m \neq n$ by (4.18), thus $C_{i,j}(0) = 0$. Now for any $x > 0$

we have

$$\begin{aligned}
\sum_{m \neq n \in \mathbb{Z}} \frac{x^{2|m|}}{2^{2|m|}|m|!^2} \frac{x^{2(|n|+1)}}{2^{2(|n|+1)}(|n|+1)!^2} &\leq \sum_{m, n \in \mathbb{Z}} \frac{x^{2|m|}}{2^{2|m|}|m|!^2} \frac{x^{2(|n|+1)}}{2^{2(|n|+1)}(|n|+1)!^2} \\
&\leq \left(\sum_{m \in \mathbb{Z}} \frac{x^{2|m|}}{2^{2|m|}|m|!^2} \right) \left(\sum_{n \in \mathbb{Z}} \frac{x^{2(|n|+1)}}{2^{2(|n|+1)}(|n|+1)!^2} \right) \\
&\leq \left(\sum_{m \in \mathbb{Z}} \frac{x^{2|m|}}{2^{2|m|}|m|!} \right) \left(\sum_{n \in \mathbb{Z}} \frac{x^{2(|n|+1)}}{2^{2(|n|+1)}(|n|+1)!} \right) \\
&\leq \left(-1 + 2 \sum_{m \geq 0} \frac{x^{2m}}{2^{2m}m!} \right) \left(-\frac{x^2}{4} + 2 \sum_{n \geq 0} \frac{x^{2(n+1)}}{2^{2(n+1)}(n+1)!} \right) \\
&\leq \left[-1 + 2 \exp\left(\frac{x^2}{4}\right) \right] \left[-\frac{x^2}{4} + 2 \left(\exp\left(\frac{x^2}{4}\right) - 1 \right) \right],
\end{aligned}$$

thus by (4.24) we know that $C_{i,j}(x) < +\infty$ for all $x \in \mathbb{R}$ and $C_{i,j}$ is derivable with

$$C'_{i,j}(x) = \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j} \frac{d\mathcal{J}(m, n)^2}{dx},$$

for all $x \in \mathbb{R}$. By (4.21) we obtain

$$\begin{aligned}
C'_{i,j}(x) &= 2x \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j} \frac{J_m^2 J_{n+1}^2 - J_{m+1}^2 J_n^2}{m-n} \\
&= 2x \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j-i} \frac{J_{m+1}^2 J_{m+n}^2 - J_m^2 J_{m+n+1}^2}{n}.
\end{aligned}$$

Note that by (4.12h) we can distribute the double sum with the difference (we keep the above form for the moment to avoid overloading the equalities). Using (4.12e) we obtain

$$\begin{aligned}
C'_{i,j}(x) &= \frac{4x}{\pi} \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j-i} J_{m+1}^2(x) \int_0^{\pi/2} J_0(2x \sin \theta) \frac{\cos 2(m+n)\theta}{n} d\theta \\
&\quad - J_m^2(x) \int_0^{\pi/2} J_0(2x \sin \theta) \frac{\cos 2(m+n+1)\theta}{n} d\theta.
\end{aligned}$$

4.4.2 Removing the infinite sums

We now define

$$\begin{aligned}
D(x) &:= \frac{4x}{\pi} \sum_{m \in e\mathbb{Z}+i} \sum_{n \in e\mathbb{Z}+j-i} J_{m+1}^2(x) \int_0^{\pi/2} J_0(2x \sin \theta) \frac{\exp 2\mathbf{i}(m+n)\theta}{n} d\theta \\
&\quad - J_m^2(x) \int_0^{\pi/2} J_0(2x \sin \theta) \frac{\exp 2\mathbf{i}(m+n+1)\theta}{n} d\theta, \quad (4.25)
\end{aligned}$$

so that

$$C'_{i,j}(x) = \Re D(x), \quad (4.26)$$

for all $x \in \mathbb{R}$, recalling from (4.12a) that $J_n(y) \in \mathbb{R}$ if $y \in \mathbb{R}$. We now define

$$E_k^{[e]}(x) := \sum_{n \in e\mathbb{Z}+k} \frac{\exp 2\mathbf{i}nx}{n} = 2 \sum_{n \in 2e\mathbb{Z}+2k} \frac{\exp \mathbf{i}nx}{n}, \quad (4.27)$$

for any $k \notin e\mathbb{Z}$ and $x \in \mathbb{R}$. The value of $E_{j-i}^{[e]}$ is given by the Lemma 4.28 below, whose proof is postponed until §A.1. Recall from Lemma 4.14 that $\Omega_{2e} = \left\{ \frac{\ell\pi}{e} : -e < \ell \leq e \right\} \subseteq (-\pi, \pi]$. We define $\omega_\ell := \frac{\ell\pi}{e}$ so that $\Omega_{2e} = \{\omega_\ell\}_{-e < \ell \leq e}$.

Lemma 4.28. For any $k \notin e\mathbb{Z}$ and $x \in (-\pi, \pi) \setminus \Omega_{2e}$ we have

$$E_k^{[e]}(x) = \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \exp(2\mathbf{i}k\omega)(\omega - \epsilon_{x,\omega}\pi),$$

with $\epsilon_{x,\omega} := \operatorname{sgn}(\omega - x)$. In particular, the function $E_k^{[e]}$ is piecewise constant on each interval $(\omega_\ell, \omega_{\ell+1})$ for $-e \leq \ell < e$.

From (4.25) and (4.27) we have

$$D(x) = \frac{4x}{\pi} \int_0^{\pi/2} J_0(2x \sin \theta) E_{j-i}^{[e]}(\theta) \sum_{m \in e\mathbb{Z}+i} [J_{m+1}^2(x) \exp(2\mathbf{i}m\theta) - J_m^2(x) \exp(2\mathbf{i}(m+1)\theta)] d\theta. \quad (4.29)$$

Note that in (4.25) we can permute the integral and the sum in m by (4.12c) and (4.12h). Now to permute the integral and the sum in n , we have to be a bit more careful. By summation by parts, we have, assuming that $k := j - i$ with $0 \leq i < j < e$,

$$\begin{aligned} \sum_{n \in e\mathbb{Z}_{>0}+k} \frac{\exp 2\mathbf{i}n\theta}{n} &= \sum_{n \in e\mathbb{Z}_{>0}+k} \frac{\mathcal{E}_n(\theta) - \mathcal{E}_{n-e}(\theta)}{n} \\ &= -\frac{\mathcal{E}_k(\theta)}{e+k} + e \sum_{n \in e\mathbb{Z}_{>0}+k} \frac{\mathcal{E}_n(\theta)}{n(n+e)}, \end{aligned} \quad (4.30)$$

where

$$\mathcal{E}_n(\theta) := \sum_{\substack{m \in e\mathbb{Z}_{>0}+k \\ m \leq n}} \exp 2\mathbf{i}m\theta$$

(with a similar treatment for the remaining elements $n \in e\mathbb{Z} + k$). Now writing $n = eN + k$ with $N > 0$, by a standard calculation we have, for $\theta \notin \frac{\pi}{e}\mathbb{Z}$,

$$\begin{aligned} \mathcal{E}_n(\theta) &= \sum_{m=1}^N \exp 2\mathbf{i}(em+k)\theta \\ &= \exp(2\mathbf{i}k\theta) \sum_{m=1}^N \exp 2\mathbf{i}em\theta \\ &= \exp(2\mathbf{i}k\theta) \exp(\mathbf{i}e(N+1)\theta) \frac{\sin eN\theta}{\sin e\theta}, \end{aligned}$$

thus since $|\mathcal{E}_n(\theta)| \leq N \leq n$ we deduce that²

$$|\mathcal{E}_n(\theta)| = \sqrt{|\mathcal{E}_n(\theta)|^2} \leq \frac{\sqrt{N}}{\sqrt{|\sin e\theta|}} \leq \frac{\sqrt{n}}{\sqrt{|\sin e\theta|}}.$$

We thus have, for $n \in e\mathbb{Z}_{>0} + k$,

$$\left| \frac{\mathcal{E}_n(\theta)}{n(n+e)} \right| \leq \frac{\sqrt{n}}{n(n+e)} \frac{1}{\sqrt{|\sin e\theta|}} \leq \frac{1}{(n+e)\sqrt{n}} \frac{1}{\sqrt{|\sin e\theta|}},$$

thus since $\frac{1}{\sqrt{|\sin u|}}$ is integrable at 0 we deduce that

$$\int_0^{\pi/2} \sum_{n \in e\mathbb{Z}_{>0}+k} \frac{|\mathcal{E}_n(\theta)|}{n(n+e)} d\theta < +\infty,$$

²The author is indebted to Arnaud Debussche for this trick.

which, together with (4.12c) and recalling (4.30), justifies the permutation of the integral and the sum in n between (4.25) and (4.29) (the reasoning for $n \in e\mathbb{Z}_{\leq 0} + k$ being similar).

We now go back to (4.29), which we repeat here:

$$D(x) = \frac{4x}{\pi} \int_0^{\pi/2} J_0(2x \sin \theta) E_{j-i}^{[e]}(\theta) \sum_{m \in e\mathbb{Z}+i} [J_{m+1}^2(x) \exp(2\mathbf{i}m\theta) - J_m^2(x) \exp 2\mathbf{i}(m+1)\theta] d\theta.$$

By (4.12d) we have

$$D(x) = \frac{8x}{\pi^2} \iint_{[0, \pi/2]^2} J_0(2x \sin \theta) E_{j-i}^{[e]}(\theta) \sum_{m \in e\mathbb{Z}+i} J_{2(m+1)}(2x \cos \eta) \exp(2\mathbf{i}m\theta) - J_{2m}(2x \cos \eta) \exp 2\mathbf{i}(m+1)\theta d\theta d\eta, \quad (4.31)$$

the permutation between the sum and the integral being justified by (4.12h). By Lemma 4.14 we have

$$\sum_{m \in 2e\mathbb{Z}+2k} J_m(x) \exp(\mathbf{i}mu) = \frac{1}{2e} \sum_{\omega \in \Omega_{2e}} \exp \mathbf{i}[-2k\omega + x \sin(\omega + u)],$$

for any $k \in \mathbb{Z}$ and $u \in \mathbb{R}$, thus the sum in (4.31) becomes, where y stands for $2x \cos \eta$,

$$\begin{aligned} \sum_{m \in e\mathbb{Z}+i} J_{2(m+1)}(y) \exp(2\mathbf{i}m\theta) - J_{2m}(y) \exp 2\mathbf{i}(m+1)\theta &= \sum_{m \in e\mathbb{Z}+i+1} J_{2m}(y) \exp 2\mathbf{i}(m-1)\theta - \exp 2\mathbf{i}\theta \sum_{m \in e\mathbb{Z}+i} J_{2m}(y) \exp 2\mathbf{i}m\theta \\ &= \exp(-2\mathbf{i}\theta) \sum_{m \in 2e\mathbb{Z}+2(i+1)} J_m(y) \exp \mathbf{i}m\theta \\ &\quad - \exp 2\mathbf{i}\theta \sum_{m \in 2e\mathbb{Z}+2i} J_m(y) \exp \mathbf{i}m\theta \\ &= \frac{1}{2e} \sum_{\omega \in \Omega_{2e}} \exp(-2\mathbf{i}\theta) \exp \mathbf{i}[-2(i+1)\omega + y \sin(\omega + \theta)] \\ &\quad - \exp(2\mathbf{i}\theta) \exp \mathbf{i}[-2i\omega + y \sin(\omega + \theta)] \\ &= \frac{1}{2e} \sum_{\omega \in \Omega_{2e}} [\exp(-2\mathbf{i}(\theta + \omega)) - \exp(2\mathbf{i}\theta)] \exp \mathbf{i}[-2i\omega + y \sin(\theta + \omega)]. \end{aligned}$$

Defining, for any $u, \theta \in \mathbb{R}$ and $\omega \in \Omega_{2e}$, the quantities

$$F(u) := \frac{2}{\pi} \int_0^{\pi/2} \exp \mathbf{i}[u \cos \eta] d\eta, \quad (4.32)$$

$$f_\omega(\theta) := E_{j-i}^{[e]}(\theta) [\exp(-2\mathbf{i}(\theta + \omega)) - \exp(2\mathbf{i}\theta)] \exp(-2\mathbf{i}i\omega), \quad (4.33)$$

we thus obtain, recalling (4.31),

$$D(x) = \frac{2x}{e\pi} \sum_{\omega \in \Omega_{2e}} \int_0^{\pi/2} J_0(2x \sin \theta) f_\omega(\theta) F(2x \sin(\theta + \omega)) d\theta.$$

We now fix $\omega \in \Omega_{2e}$ and we study the asymptotics as $x \rightarrow +\infty$ of

$$D_\omega(x) = \frac{2x}{e\pi} \int_0^{\pi/2} J_0(2x \sin \theta) f_\omega(\theta) F(2x \sin(\theta + \omega)) d\theta, \quad (4.34)$$

so that

$$D = \sum_{\omega \in \Omega_{2e}} D_\omega. \quad (4.35)$$

4.4.3 Negligible terms

By [OLBC, 10.9.1], for $y \in \mathbb{R}$ we have

$$J_0(y) = \frac{1}{\pi} \int_0^\pi \cos(y \cos(\eta)) d\eta,$$

thus $J_0(y) = \frac{2}{\pi} \int_0^{\pi/2} \cos(y \cos(\eta)) d\eta = \Re F(y)$. By [OLBC, 10.17.3] we have, for $x > 0$,

$$J_0(x) = \sqrt{\frac{2}{\pi x}} \left(\cos\left(x - \frac{\pi}{4}\right) + M(x) \right), \quad M(x) =_{x \rightarrow +\infty} O\left(\frac{1}{x}\right). \quad (4.36)$$

Since $M(x) = \sqrt{\frac{\pi x}{2}} J_0(x) - \cos\left(x - \frac{\pi}{4}\right)$ for all $x \geq 0$, the function M is continuous on \mathbb{R}_+ and thus there exists a constant $C \geq 0$ such that

$$|M(x)| \leq \frac{C}{1+x},$$

for all $x \geq 0$.

Now, $\Im F(y) = \frac{2}{\pi} \int_0^{\pi/2} \sin(y \cos \eta) d\eta$ is the *Struve function of zero order*, thus by [OLBC, 11.2.5, 11.6.1, 10.17.4] and (4.36) we have, for $x > 0$,

$$F(x) = \sqrt{\frac{2}{\pi x}} \left(\exp \mathbf{i} \left(x - \frac{\pi}{4} \right) + N(x) \right), \quad N(x) =_{x \rightarrow +\infty} O\left(\frac{1}{x}\right).$$

Using the equality $F(-x) = \overline{F(x)}$, we deduce that we have, for $x < 0$,

$$F(x) = \sqrt{\frac{2}{-\pi x}} \left(\exp \mathbf{i} \left(x + \frac{\pi}{4} \right) + \overline{N(-x)} \right), \quad \overline{N(-x)} =_{x \rightarrow +\infty} O\left(\frac{1}{x}\right).$$

so that for all $x \neq 0$,

$$F(x) = \sqrt{\frac{2}{\pi|x|}} \left(\exp \mathbf{i} \left(x - \operatorname{sgn}(x) \frac{\pi}{4} \right) + N(x) \right), \quad N(x) =_{|x| \rightarrow +\infty} O\left(\frac{1}{x}\right). \quad (4.37)$$

Similarly to M , we can take C large enough so that for any $x \in \mathbb{R}$ we have

$$|N(x)| \leq \frac{C}{1+|x|}. \quad (4.38)$$

Let us now recall the following version of the Riemann–Lebesgue lemma (we provide a proof in §A.2).

Lemma 4.39 (Riemann–Lebesgue). *Let f be integrable on (a, b) and ϕ be continuously differentiable on $[a, b]$. If ϕ vanishes at a finite number of points then*

$$\int_a^b f(t) \exp \mathbf{i}(x\phi(t)) dt \xrightarrow{x \rightarrow +\infty} 0.$$

Proposition 4.40. *If $\omega \notin \{0, \pi\}$ then*

$$\lim_{x \rightarrow +\infty} D_\omega(x) = 0.$$

Proof. Recalling (4.34), write

$$D_\omega(x) = \frac{1}{e\pi} \int_0^{\pi/2} \sqrt{2x \sin \theta} J_0(2x \sin \theta) \sqrt{2x |\sin(\theta + \omega)|} |F(2x \sin(\theta + \omega))| g_\omega(\theta) d\theta,$$

where $g_\omega(\theta) := \frac{f_\omega(\theta)}{\sqrt{|\sin(\theta)| |\sin(\theta + \omega)|}}$. Note that f_ω is piecewise continuous, moreover by assumption if $\sin(\theta) = 0$ then $\sin(\theta + \omega) \neq 0$ (since $\omega \neq 0 \pmod{\pi}$) thus $\theta \mapsto \frac{1}{\sqrt{|\sin(\theta)| |\sin(\theta + \omega)|}}$ and thus g_ω is integrable on $(0, \pi/2)$.

Using (4.36) and (4.37) we obtain for $D_\omega(x)$ a sum of these following four integrals (with some multiplicative coefficients that do not depend on x):

$$\begin{aligned} & \int_0^{\pi/2} \cos\left(2x \sin \theta - \frac{\pi}{4}\right) \exp \mathbf{i} \left(2x \sin(\theta + \omega) - \operatorname{sgn}(\sin(\theta + \omega)) \frac{\pi}{4}\right) g_\omega(\theta) d\theta, \\ & \int_0^{\pi/2} \cos\left(2x \sin \theta - \frac{\pi}{4}\right) N(2x \sin(\theta + \omega)) g_\omega(\theta) d\theta, \\ & \int_0^{\pi/2} \exp \mathbf{i} \left(2x \sin(\theta + \omega) - \operatorname{sgn}(\sin(\theta + \omega)) \frac{\pi}{4}\right) M(2x \sin \theta) g_\omega(\theta) d\theta, \\ & \int_0^{\pi/2} M(2x \sin \theta) N(2x \sin(\theta + \omega)) g_\omega(\theta) d\theta. \end{aligned}$$

We can easily see that the last three integrals go to zero as x grows to infinity. For instance, we have, using (4.38),

$$\left| \int_0^{\pi/2} \cos\left(2x \sin \theta - \frac{\pi}{4}\right) N(2x \sin(\theta + \omega)) g_\omega(\theta) d\theta \right| \leq C \int_0^{\pi/2} \frac{|g_\omega(\theta)|}{1 + 2x |\sin(\theta + \omega)|} d\theta,$$

and since

$$\frac{|g_\omega(\theta)|}{1 + 2x |\sin(\theta + \omega)|} \leq |g_\omega(\theta)|,$$

is integrable on $(0, \pi/2)$ and

$$\frac{|g_\omega(\theta)|}{1 + 2x |\sin(\theta + \omega)|} \xrightarrow{x \rightarrow +\infty} 0,$$

for almost all $\theta \in (0, \pi/2)$ (all except maybe the only θ for which $\sin(\theta + \omega) = 0$) we obtain the result by dominated convergence.

For the first integral, we have

$$\begin{aligned} & \int_0^{\pi/2} g_\omega(\theta) \cos\left(2x \sin \theta - \frac{\pi}{4}\right) \exp \mathbf{i} \left(2x \sin(\theta + \omega) - \operatorname{sgn}(\sin(\theta + \omega)) \frac{\pi}{4}\right) d\theta \\ &= \frac{1}{2} \sum_{\epsilon \in \{\pm 1\}} \int_0^{\pi/2} g_\omega(\theta) \exp \mathbf{i} \left(2x \sin \theta - \frac{\pi}{4}\right) \exp \mathbf{i} \left(2x \sin(\theta + \omega) - \operatorname{sgn}(\sin(\theta + \omega)) \frac{\pi}{4}\right) d\theta \\ &= \frac{1}{2} \sum_{\epsilon \in \{\pm 1\}} \int_0^{\pi/2} h_{\omega, \epsilon}(\theta) \exp(2ix \phi_\epsilon(t)) d\theta, \end{aligned} \tag{4.41}$$

with

$$\phi_\epsilon(t) = \epsilon \sin \theta + \sin(\theta + \omega),$$

and

$$h_{\omega, \epsilon}(\theta) = g_\omega(\theta) \exp \left[\frac{-i\pi}{4} (\epsilon + \operatorname{sgn}(\sin(\theta + \omega))) \right].$$

The function $h_{\omega, \epsilon}$ is integrable on $(0, \pi/2)$, and we will now see that ϕ'_ϵ vanishes at at most one point. We have $\phi'_\epsilon(\theta) = \epsilon \cos \theta + \cos(\theta + \omega)$. For $\epsilon = 1$ we have

$$\begin{aligned} \phi'_1(\theta) = 0 & \iff \cos(\theta + \omega) = -\cos \theta \\ & \iff \theta + \omega = \pi \pm \theta \pmod{2\pi}, \end{aligned}$$

thus since $\omega \neq \pi$ we obtain

$$\begin{aligned} \phi'_1(\theta) = 0 & \iff \theta + \omega = \pi - \theta \pmod{2\pi} \\ & \iff \theta = \frac{\pi - \omega}{2} \pmod{2\pi}. \end{aligned}$$

Similarly, for $\epsilon = -1$ we have

$$\begin{aligned}\phi'_{-1}(\theta) = 0 &\iff \cos(\theta + \omega) = \cos \theta \\ &\iff \theta + \omega = \pm \theta \pmod{2\pi}\end{aligned}$$

thus since $\omega \neq 0$ we obtain

$$\begin{aligned}\phi'_{-1}(\theta) = 0 &\iff \theta + \omega = -\theta \pmod{2\pi} \\ &\iff \theta = \pi - \frac{\omega}{2}.\end{aligned}$$

We thus conclude, applying the Riemann–Lebesgue lemma (Lemma 4.39), that the quantity in (4.41) goes to zero as $x \rightarrow +\infty$ and this finishes the proof of the proposition. \square

4.4.4 Contributing terms

If $\omega \in \{0, \pi\}$ then $2\omega = 0 \pmod{2\pi}$ and thus, recalling (4.33),

$$f_\omega(\theta) = -2\mathbf{i} \sin(2\theta) E_{j-i}^{[e]}(\theta). \quad (4.42)$$

We deduce from (4.34) and (4.42) that if $\omega \in \{0, \pi\}$ then

$$D_\omega(x) = -\frac{4\mathbf{i}x}{e\pi} \int_0^{\pi/2} E_{j-i}^{[e]}(\theta) J_0(2x \sin \theta) \sin(2\theta) F(2x \sin(\theta + \omega)) d\theta.$$

In particular, noting from (4.32) that $F(-y) = \overline{F(y)}$ we have

$$D_0(x) = -\frac{4\mathbf{i}x}{e\pi} \int_0^{\pi/2} E_{j-i}^{[e]}(\theta) J_0(2x \sin \theta) \sin(2\theta) F(2x \sin \theta) d\theta, \quad (4.43a)$$

$$D_\pi(x) = -\frac{4\mathbf{i}x}{e\pi} \int_0^{\pi/2} E_{j-i}^{[e]}(\theta) J_0(2x \sin \theta) \sin(2\theta) \overline{F(2x \sin \theta)} d\theta. \quad (4.43b)$$

Recall that we have assumed at the beginning of §4.4 that $i \neq j$. Recall also the notation $\omega_\ell = \frac{\ell\pi}{e}$ for $-e \leq \ell \leq e$.

Proposition 4.44. *We have*

$$\lim_{x \rightarrow +\infty} D_0(x) = \lim_{x \rightarrow +\infty} D_\pi(x) = d_{e,j-i},$$

where

$$d_{e,k} := \left(\frac{2}{e\pi}\right)^2 \left(\sum_{\omega \in \Omega_{2e}} \omega \exp 2\mathbf{i}k\omega + 2\pi \sum_{n=1}^{e'-1} (1 - \sin \omega_n) \exp 2\mathbf{i}k\omega_n \right),$$

for $k \notin e\mathbb{Z}$ and $e' := \lceil \frac{e}{2} \rceil$.

Proof. By (4.36) and (4.37), for $t \geq 0$ we have

$$\begin{aligned}tJ_0(t)F(t) &= \frac{2}{\pi} \cos\left(t - \frac{\pi}{4}\right) \exp \mathbf{i}\left(t - \frac{\pi}{4}\right) + R(t) \\ &= \frac{1}{\pi} \left[2 \cos^2\left(t - \frac{\pi}{4}\right) + \mathbf{i} \sin\left(2t - \frac{\pi}{2}\right) \right] + R(t) \\ &= \frac{1}{\pi} \left[\cos\left(2t - \frac{\pi}{2}\right) + 1 + \mathbf{i} \sin\left(2t - \frac{\pi}{2}\right) \right] + R(t) \\ &= \frac{1}{\pi} + \frac{1}{\pi} (\sin(2t) - \mathbf{i} \cos(2t)) + R(t),\end{aligned} \quad (4.45)$$

where R is continuous on \mathbb{R}_+ and satisfies $R(t) = O(\frac{1}{t})$ when $t \rightarrow +\infty$. In particular, there exists $C \geq 0$ such that $|R(t)| \leq \frac{C}{1+t}$ for all $t \geq 0$.

Recall from Lemma 4.28 that $E_{j-i}^{[e]}$ is constant on each interval $(\omega_\ell, \omega_{\ell+1})$ for each $-e \leq \ell < e$. From (4.43a) and (4.45) we have, for any $0 \leq a < b \leq \frac{\pi}{2}$,

$$\begin{aligned} & \frac{4x}{e\pi} \int_a^b \sin(2\theta) J_0(2x \sin \theta) F(2x \sin \theta) d\theta \\ &= \frac{4}{e\pi} \int_a^b 2x \sin(\theta) J_0(2x \sin \theta) F(2x \sin \theta) \cos(\theta) d\theta \\ &= \frac{2}{e\pi} \int_{2 \sin a}^{2 \sin b} xu J_0(xu) F(xu) du \\ &= \frac{2}{e\pi^2} \int_{2 \sin a}^{2 \sin b} 1 + \sin(2xu) - \mathbf{i} \cos(2xu) + \pi R(xu) du, \end{aligned} \quad (4.46)$$

where we did the variable change $u = 2 \sin \theta$. Moreover, as $x \rightarrow +\infty$ we have

$$\int_{2 \sin a}^{2 \sin b} \sin(2xu) - \mathbf{i} \cos(2xu) du = o(1),$$

by the Riemann–Lebesgue lemma, and

$$\begin{aligned} \left| \int_{2 \sin a}^{2 \sin b} R(xu) du \right| &\leq \int_{2 \sin a}^{2 \sin b} \frac{C}{1+xu} du \\ &= \frac{C}{x} [\ln(1+xu)]_{2 \sin a}^{2 \sin b} \\ &= o(1), \end{aligned}$$

so that (4.46) gives

$$\frac{4x}{e\pi} \int_a^b \sin(2\theta) J_0(2x \sin \theta) F(2x \sin \theta) d\theta = \frac{4}{e\pi^2} (\sin a - \sin b) + o(1), \quad (4.47)$$

as $x \rightarrow +\infty$.

Now define $e' := \lceil \frac{e}{2} \rceil$ so that $\omega_{e'} \geq \pi/2 > \omega_{e'-1}$. Define $\hat{\omega}_\ell$ for $\ell \in \{0, \dots, e'\}$ by

$$\hat{\omega}_\ell := \min \left(\omega_\ell, \frac{\pi}{2} \right) = \begin{cases} \omega_\ell, & \text{if } \ell < e', \\ \frac{\pi}{2}, & \text{if } \ell = e'. \end{cases}$$

We deduce from (4.12a), (4.43) and (4.47) that, denoting by \mathbf{e}_ℓ the value of $E_{j-i}^{[e]}$ on the interval $(\omega_\ell, \omega_{\ell+1})$,

$$D_0(x) = \frac{-4\mathbf{i}}{e\pi^2} \sum_{\ell=0}^{e'-1} \mathbf{e}_\ell (\sin \hat{\omega}_{\ell+1} - \sin \hat{\omega}_\ell) + o(1) = D_\pi(x), \quad (4.48)$$

as $x \rightarrow +\infty$. Now we have, noting that $\sin \hat{\omega}_{e'} = 1$ and $\sin \hat{\omega}_0 = 0$,

$$\begin{aligned} \sum_{\ell=0}^{e'-1} \mathbf{e}_\ell (\sin \hat{\omega}_{\ell+1} - \sin \hat{\omega}_\ell) &= \sum_{\ell=1}^{e'} \mathbf{e}_{\ell-1} \sin \hat{\omega}_\ell - \sum_{\ell=0}^{e'-1} \mathbf{e}_\ell \sin \hat{\omega}_\ell \\ &= \mathbf{e}_{e'-1} \sin \hat{\omega}_{e'} - \mathbf{e}_0 \sin \hat{\omega}_0 + \sum_{\ell=1}^{e'-1} (\mathbf{e}_{\ell-1} - \mathbf{e}_\ell) \sin \hat{\omega}_\ell \\ &= \mathbf{e}_{e'-1} + \sum_{\ell=1}^{e'-1} (\mathbf{e}_{\ell-1} - \mathbf{e}_\ell) \sin \omega_\ell. \end{aligned} \quad (4.49)$$

By Lemma 4.28 we have, for any $x \in (\omega_\ell, \omega_{\ell+1})$ and with $k := j - i \pmod{e}$ (thus $k \notin e\mathbb{Z}$),

$$\begin{aligned}
\epsilon_\ell &= \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \exp(2\mathbf{i}k\omega) (\omega - \operatorname{sgn}(\omega - x)\pi) \\
&= \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \omega \exp 2\mathbf{i}k\omega + \frac{\mathbf{i}\pi}{e} \sum_{n=-e+1}^{\ell} \exp 2\mathbf{i}k\omega_n - \frac{\mathbf{i}\pi}{e} \sum_{n=\ell+1}^e \exp 2\mathbf{i}k\omega_n \\
&= \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \omega \exp 2\mathbf{i}k\omega - \frac{2\mathbf{i}\pi}{e} \sum_{n=\ell+1}^e \exp 2\mathbf{i}k\omega_n \\
&= \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \omega \exp 2\mathbf{i}k\omega + \frac{2\mathbf{i}\pi}{e} \sum_{n=1}^{\ell} \exp 2\mathbf{i}k\omega_n
\end{aligned} \tag{4.50}$$

since $\sum_{\omega \in \Omega_{2e}} \exp 2\mathbf{i}k\omega = 0 = \sum_{n=1}^e \exp 2\mathbf{i}k\omega_n$ since $2k \notin 2e\mathbb{Z}$, thus

$$\begin{aligned}
\epsilon_{\ell-1} - \epsilon_\ell &= \frac{2\mathbf{i}\pi}{e} \sum_{n=1}^{\ell-1} \exp 2\mathbf{i}k\omega_n - \frac{2\mathbf{i}\pi}{e} \sum_{n=1}^{\ell} \exp 2\mathbf{i}k\omega_n \\
&= \frac{-2\mathbf{i}\pi}{e} \exp(2\mathbf{i}k\omega_\ell).
\end{aligned} \tag{4.51}$$

Gathering (4.48), (4.49), (4.50) and (4.51) gives the desired result. \square

4.4.5 Expliciting the limit

We are now interested in a closed form for the elements $\Re d_{e,k}$, where

$$d_{e,k} = \left(\frac{2}{e\pi} \right)^2 \left(\sum_{\omega \in \Omega_{2e}} \omega \exp 2\mathbf{i}k\omega + 2\pi \sum_{n=1}^{e'-1} (1 - \sin \omega_n) \exp 2\mathbf{i}k\omega_n \right),$$

are the elements of Proposition 4.44, where $k \notin e\mathbb{Z}$ and $e' = \lceil \frac{e}{2} \rceil$.

Lemma 4.52. *For any $k \notin e\mathbb{Z}$ we have*

$$\Re d_{e,k} = \frac{2}{\pi e^2} \left[\cot\left(k - \frac{1}{2}\right) \frac{\pi}{e} - \cot\left(k + \frac{1}{2}\right) \frac{\pi}{e} \right].$$

In particular, we have $\Re d_{e,k} \neq 0$.

Proof. First, we have

$$\begin{aligned}
\Re \sum_{\omega \in \Omega_{2e}} \omega \exp(2\mathbf{i}k\omega) &= \sum_{\omega \in \Omega_{2e}} \omega \cos 2k\omega \\
&= \pi,
\end{aligned}$$

noticing that if $\omega \in \Omega_{2e} \setminus \{\pi\}$ then $-\omega \in \Omega_{2e}$. We now have to deal with the sum

$$A := \Re \left(2 \sum_{n=1}^{e'-1} (1 - \sin \omega_n) \exp 2\mathbf{i}k\omega_n \right) = 2 \sum_{n=1}^{e'-1} (1 - \sin \omega_n) \cos 2k\omega_n.$$

Recall that $\omega_n = \frac{\pi n}{e}$. For any $n \in \{1, \dots, e-1\}$ we thus have $\omega_{e-n} = \pi - \omega_n$, hence $\sin \omega_{e-n} = \sin \omega_n$ and $\cos 2k\omega_{e-n} = \cos(2k\pi - 2k\omega_n) = \cos 2k\omega_n$. Hence, we have

$$\begin{aligned}
A &= \sum_{n=1}^{e-1} (1 - \sin \omega_n) \cos 2k\omega_n - \begin{cases} (1 - \sin \omega_{e/2}) \cos 2k\omega_{e/2}, & \text{if } n \text{ is even,} \\ 0, & \text{if } n \text{ is odd,} \end{cases} \\
&= \sum_{n=1}^{e-1} (1 - \sin \omega_n) \cos 2k\omega_n,
\end{aligned}$$

since $\sin \omega_{e/2} = \sin \frac{\pi}{2} = 1$ if e is even. Since $\sum_{n=1}^{e-1} \cos 2k\omega_n = \sum_{n=1}^{e-1} \cos\left(\frac{2k\pi}{e}n\right) = -1$ since $2k \notin 2e\mathbb{Z}$, we obtain that, using the identity $\sin(x)\cos(y) = \frac{1}{2}[\sin(x+y) + \sin(x-y)]$,

$$\begin{aligned} A &= -1 - \sum_{n=1}^{e-1} \sin \omega_n \cos 2k\omega_n \\ &= -1 + \frac{1}{2} \sum_{n=1}^{e-1} [\sin(2k-1)\omega_n - \sin(2k+1)\omega_n]. \end{aligned}$$

We now recall the following standard formula (see, for instance, [GrRy, 1.342.1]):

$$\sum_{n=1}^{e-1} \sin nx = \sin \frac{ex}{2} \sin \frac{(e-1)x}{2} \operatorname{cosec} \frac{x}{2},$$

valid for any $x \notin 2\pi\mathbb{Z}$. We obtain, using the identity $\sin\left(a \pm \frac{\pi}{2}\right) = \pm \cos a$,

$$\begin{aligned} \sum_{n=1}^{e-1} \sin(2k \pm 1)\omega_n &= \sum_{n=1}^{e-1} \sin \frac{(2k \pm 1)\pi n}{e} \\ &= \sin\left[\left(k \pm \frac{1}{2}\right)\pi\right] \sin\left[(1-e^{-1})\left(k \pm \frac{1}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= \pm(-1)^k \sin\left[(1-e^{-1})\left(k \pm \frac{1}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= \pm(-1)^k \sin\left[\left((1-e^{-1})k \mp \frac{e^{-1}}{2} \pm \frac{1}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= (-1)^k \cos\left[\left((1-e^{-1})k \mp \frac{e^{-1}}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= \cos\left[\left(-e^{-1}k \mp \frac{e^{-1}}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= \cos\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \operatorname{cosec}\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right] \\ &= \cot\left[e^{-1}\left(k \pm \frac{1}{2}\right)\pi\right]. \end{aligned}$$

and thus

$$A = -1 + \frac{1}{2} \left(\cot\left[e^{-1}\left(k - \frac{1}{2}\right)\pi\right] - \cot\left[e^{-1}\left(k + \frac{1}{2}\right)\pi\right] \right).$$

Finally, we obtain

$$\begin{aligned} \Re d_{e,k} &= \left(\frac{2}{e\pi}\right)^2 (\pi + \pi A) \\ &= \frac{4}{\pi e^2} (1 + A), \end{aligned}$$

as desired. Note that $\Re d_{e,k} \neq 0$ since the cotangent function is bijective on $(0, \pi)$ (taking $1 \leq k \leq e-1$). \square

We conclude this part by putting (4.26), (4.35), Proposition 4.40, Proposition 4.44 and Lemma 4.52 to find that

$$\lim_{x \rightarrow +\infty} C'_{i,j}(x) = 2\Re d_{e,j-i} = \frac{4}{\pi e^2} \left[\cot\left(j-i-\frac{1}{2}\right)\frac{\pi}{e} - \cot\left(j-i+\frac{1}{2}\right)\frac{\pi}{e} \right],$$

in particular, the limit depends only on $j-i$. Since $\Re d_{e,j-i} \neq 0$ we find that

$$C_{i,j}(x) \sim 2\Re d_{e,j-i} x,$$

as $x \rightarrow +\infty$. By (4.19), (4.22) and Proposition 4.17, we have, with $i \neq j$,

$$\operatorname{Cov}_t(x_i(\lambda), x_j(\lambda)) = \frac{-1}{4} C_{i,j}(2\sqrt{t}),$$

thus $\operatorname{Cov}_t(x_i(\lambda), x_j(\lambda)) \sim -\Re d_{e,j-i} \sqrt{t}$. We thus deduce from Lemma 4.52 the final result of this part.

Theorem 4.53. *If $i \neq j$ then as $t \rightarrow +\infty$ we have, under the Poissonised Plancherel measure pl_t ,*

$$\text{Cov}_t(x_i(\lambda), x_j(\lambda)) \sim \frac{2\sqrt{t}}{\pi e^2} \left[\cot\left(j - i + \frac{1}{2}\right) \frac{\pi}{e} - \cot\left(j - i - \frac{1}{2}\right) \frac{\pi}{e} \right].$$

4.5 Asymptotics of the variance

Recall from (3.8) that $\sum_i x_i(\lambda) = 0$. Hence, we directly obtain that

$$\text{Var}_t x_i(\lambda) = \text{Cov}_t(x_i(\lambda), x_i(\lambda)) = - \sum_{j \neq i} \text{Cov}_t(x_i(\lambda), x_j(\lambda)). \quad (4.54)$$

Corollary 4.55. *As $t \rightarrow +\infty$ we have, under pl_t ,*

$$\text{Var}_t x_i(\lambda) \sim \frac{4\sqrt{t}}{\pi e^2} \cot \frac{\pi}{2e}.$$

Proof. By Theorem 4.53 and (4.54), it suffices to prove that $\sum_{k=1}^{e-1} \Re d_{e,k} = \frac{4}{\pi e^2} \cot \frac{\pi}{2e}$. Recalling from Lemma 4.52 that

$$d_{e,k} = \frac{2}{\pi e^2} \left[\cot\left(k - \frac{1}{2}\right) \frac{\pi}{e} - \cot\left(k + \frac{1}{2}\right) \frac{\pi}{e} \right],$$

we obtain, using the fact that $\cot\left(e - \frac{1}{2}\right) \frac{\pi}{e} = \cot\left(\pi - \frac{\pi}{2e}\right) = -\cot \frac{\pi}{2e}$,

$$\begin{aligned} \sum_{k=1}^{e-1} d_{e,k} &= \frac{2}{\pi e^2} \left[\cot \frac{\pi}{2e} - \cot\left(e - \frac{1}{2}\right) \frac{\pi}{e} \right] \\ &= \frac{4}{\pi e} \cot \frac{\pi}{2e}, \end{aligned}$$

as desired. Note that $\cot \frac{\pi}{2e} \neq 0$ indeed since $e \geq 2$. \square

Remark 4.56. In particular, Theorem 4.53 remains valid in the case $i = j$, and the limit of $t^{-1/2} \text{Var}_t x_i(\lambda)$ as $t \rightarrow +\infty$ does not depend on i .

With Proposition 4.13 and Remark 4.56, we thus have proved Theorem A of the introduction.

5 Size of the core of a random partition

We will now use the results of the previous sections to determine the limit law of $|\bar{\lambda}|$ under pl_t as $t \rightarrow +\infty$.

5.1 Limit expectation

Write $v_t := \frac{4\sqrt{t}}{\pi e^2} \cot \frac{\pi}{2e}$. From (3.12) we have

$$\frac{|\bar{\lambda}|}{v_t} = \frac{e}{2} \sum_{i \in \mathbb{Z}/e\mathbb{Z}} \frac{x_i(\lambda)^2}{v_t} + \sum_{i=0}^{e-1} i \frac{x_i(\lambda)}{v_t}.$$

From Proposition 4.13 and Corollary 4.55, we obtain that

$$\lim_{t \rightarrow +\infty} \mathbb{E}_t \frac{|\bar{\lambda}|}{v_t} = \frac{e^2}{2},$$

and thus the following result.

Proposition 5.1. *Under the Poissonised Plancherel measure pl_t , as $t \rightarrow +\infty$ we have*

$$\mathbb{E}_t |\bar{\lambda}| \sim \frac{2\sqrt{t}}{\pi} \cot \frac{\pi}{2e}.$$

Remark 5.2. (See [LuPi, AySi].) Under the uniform measure on partitions of n , the expectation of $|\bar{\lambda}|$ is $\frac{(e-1)\sqrt{6n}}{2\pi}$. The order of the asymptotics is the same but the constant is quite different.

5.2 Limit law for $x_i(\lambda)$

Recall from (4.5) that for $i \in \{0, \dots, e-1\}$ we have

$$\hat{x}_i(\lambda) = \#_t^{(i)} - t^2, \quad (5.3)$$

where $\#_t^{(i)} = \#(e\mathbb{Z}_{\geq -t^2+i}) \cap \mathcal{D}(\lambda)$. In particular, we have $\text{Cov}_t(\hat{x}_i(\lambda), \hat{x}_j(\lambda)) = \text{Cov}_t(\#_t^{(i)}, \#_t^{(j)})$. Now define

$$b_{ij} := \begin{cases} 2 \cot \frac{\pi}{2e}, & \text{if } i = j, \\ \cot(j - i + \frac{1}{2}) \frac{\pi}{e} - \cot(j - i - \frac{1}{2}) \frac{\pi}{e}, & \text{if } i \neq j, \end{cases} \quad (5.4)$$

for $0 \leq i, j < e$ (the formula for $i \neq j$ being in fact valid for $i = j$), set $c_{ij} := \frac{2}{\pi e^2} b_{ij}$ and let N be an e -dimensional centred normal random vector with covariance matrix $B = (b_{ij})$. By Theorem 4.53 and Remark 4.56, we know that $\text{Cov}_t(x_i(\lambda), x_j(\lambda)) \sim c_{ij} \sqrt{t}$ as $t \rightarrow +\infty$ (recalling that $c_{ij} > 0$). We have

$$\begin{aligned} t^{-1/2} \text{Cov}_t(\#_t^{(i)}, \#_t^{(j)}) &= t^{-1/2} \text{Cov}_t(\hat{x}_i(\lambda), \hat{x}_j(\lambda)) \\ &= t^{-1/2} \text{Cov}_t(x_i(\lambda) + (\hat{x}_i(\lambda) - x_i(\lambda)), x_j(\lambda) + (\hat{x}_j(\lambda) - x_j(\lambda))) \\ &= t^{-1/2} \text{Cov}_t(x_i(\lambda), x_j(\lambda)) + t^{-1/2} o(1). \end{aligned}$$

Indeed, by Lemma 4.6 and Cauchy-Schwarz inequality we have that, as $t \rightarrow +\infty$,

$$\text{Cov}_t(\hat{x}_i(\lambda) - x_i(\lambda), \hat{x}_j(\lambda) - x_j(\lambda)) = o(t^{-t})$$

and

$$\text{Cov}_t(x_i(\lambda), \hat{x}_j(\lambda) - x_j(\lambda)) = o(t^{\frac{1}{4} - \frac{t}{2}}),$$

(and the same quantity with i and j permuted) thus both are $o(1)$. We deduce that $\text{Cov}_t(\#_t^{(i)}, \#_t^{(j)}) \sim_t c_{ij} \sqrt{t}$ and thus

$$\frac{\text{Cov}_t(\#_t^{(i)}, \#_t^{(j)})}{\sqrt{\text{Var}_t \#_t^{(i)}} \sqrt{\text{Var}_t \#_t^{(j)}}} \xrightarrow{t \rightarrow +\infty} \frac{b_{ij}}{b_{ii}}.$$

We can thus apply Theorem 4.1 to find that the vector

$$\left(\frac{\#_t^{(i)} - \mathbb{E}_t \#_t^{(i)}}{t^{1/4} \sqrt{c_{ii}}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

converges in distribution to the e -dimensional centred normal vector with covariance matrix $\begin{pmatrix} b_{ij} \\ b_{ii} \end{pmatrix}$, thus the vector

$$e \sqrt{\frac{\pi}{2}} \left(\frac{\#_t^{(i)} - \mathbb{E}_t \#_t^{(i)}}{t^{1/4}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

converges in distribution to N . Now by (5.3) we have $\mathbb{E}_t \hat{x}_i(\lambda) = \mathbb{E}_t \#_t^{(i)} - t^2$ and thus

$$e \sqrt{\frac{\pi}{2}} \left(\frac{\hat{x}_i(\lambda) - \mathbb{E}_t \hat{x}_i(\lambda)}{t^{1/4}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

converges in distribution to N as well. By Lemma 4.6 and Proposition 4.13, we know that $\mathbb{E}_t \hat{x}_i(\lambda)$ is bounded as $t \rightarrow +\infty$ thus the vector

$$e \sqrt{\frac{\pi}{2}} \left(\frac{\hat{x}_i(\lambda)}{t^{1/4}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

again converges in distribution to N . Finally, by Slutsky's theorem and Lemma 4.6, we obtain the following final result (which is Theorem B of the introduction).

Theorem 5.5. *Under the Poissonised Plancherel measure \mathfrak{pl}_t , as $t \rightarrow +\infty$ the random vector*

$$e\sqrt{\frac{\pi}{2}} \left(\frac{x_i(\lambda)}{t^{1/4}} \right)_{i \in \mathbb{Z}/e\mathbb{Z}},$$

converges in distribution to the e -dimensional centred normal vector with covariance matrix $B = (b_{ij})_{i,j \in \mathbb{Z}/e\mathbb{Z}}$ given by $b_{ij} = \cot(j - i + \frac{1}{2})\frac{\pi}{e} - \cot(j - i - \frac{1}{2})\frac{\pi}{e}$.

Remark 5.6. In many cases of determinantal point processes (as in [CoLe, So00-b, BoSu-b]), the variable with which we use the central limit theorem has a variance logarithmic in the expectation. In our case this does not hold, recalling Proposition 4.13 (the first order asymptotics of the expectation vanishes by the formulas in Lemma 3.13).

5.3 Limit law for $|\bar{\lambda}|$

Note that the covariance matrix $B = (b_{ij})$ of Theorem 5.5 is (symmetric, positive semi-definite and) circulant. Recall from (3.12) that

$$|\bar{\lambda}| = \frac{e}{2} \sum_{i=0}^{e-1} x_i(\lambda)^2 + \sum_{i=0}^{e-1} i x_i(\lambda).$$

Hence, by Theorem 5.5 and Slutsky's theorem we know that $t^{-1/2}|\bar{\lambda}|$ has asymptotically the same law as $t^{-1/2} \frac{e}{2} \sum_{i=0}^{e-1} x_i(\lambda)^2$. To explicit this law, it suffices to compute the eigenvalues of B . Note that since each line of B sums to 0 (by (3.8)), we already know that 0 is an eigenvalue of B .

Lemma 5.7. *The eigenvalues of B are*

$$\lambda_k = 2e \sin \frac{k\pi}{e},$$

for $k \in \mathbb{Z}/e\mathbb{Z}$. In particular, the only zero eigenvalue is λ_0 .

Proof. For any $0 \leq j < e$, we write $b_j := b_{0j}$ the entries of the first row of B . Note that since $\cot(\pi + x) = \cot(x)$, we deduce that b_j is defined for $j \in \mathbb{Z}/e\mathbb{Z}$. Since B is circulant, with the permutation matrix

$$P := \begin{pmatrix} 1 & & & \\ & \ddots & & \\ & & \ddots & \\ 1 & & & 1 \end{pmatrix},$$

we have

$$B = \sum_{j=0}^{e-1} b_j P^j.$$

Now P has each element of $\mu_e(\mathbb{C})$ for eigenvalue, an eigenvector associated with $\zeta \in \mu_e(\mathbb{C})$ being $(\zeta^j)_{0 \leq j < e}$. We deduce that the eigenvalues of B are the elements, with $\zeta_k := \exp(2ik\pi/e)$ for $k \in \mathbb{Z}/e\mathbb{Z}$,

$$\lambda_k = \sum_{j=0}^{e-1} b_j \zeta_k^j = \sum_{j=0}^{e-1} b_j \cos \frac{2kj\pi}{e}, \quad (5.8)$$

recalling that the eigenvalues of B are real since B is real symmetric. Recalling that $b_j = \cot(j + \frac{1}{2})\frac{\pi}{e} - \cot(j - \frac{1}{2})\frac{\pi}{e}$, we thus have to compute the sum

$$\sum_{j=0}^{e-1} \left[\cot(j + \frac{1}{2})\frac{\pi}{e} - \cot(j - \frac{1}{2})\frac{\pi}{e} \right] \cos \frac{2kj\pi}{e},$$

for $0 \leq k < e$. First, if $k = 0$ then $\lambda_0 = 0$ since \cot is π -periodic. We thus now assume $k \neq 0$.

We mimic the calculation in [BeYe, §4] of the following trigonometric sum (the case $n = 1$ having been first studied by Eisenstein):

$$\sum_{j=1}^{e-1} \sin \frac{2\pi kj}{e} \cot^n \left(\frac{\pi j}{e} \right).$$

Let us consider the following meromorphic function over \mathbb{C} :

$$g_1 : z \mapsto \frac{\exp 2\mathbf{i}\pi kz}{\exp(2\mathbf{i}\pi ez) - 1} - \frac{\exp(-2\mathbf{i}\pi kz)}{\exp(-2\mathbf{i}\pi ez) - 1}.$$

The function g_1 has simple poles at each $z = \frac{j}{e}$ for $j \in \mathbb{Z}$, with residue

$$\begin{aligned} \frac{\exp \frac{2\mathbf{i}\pi kj}{e}}{2\mathbf{i}\pi e \exp(2\mathbf{i}\pi j)} - \frac{\exp \frac{-2\mathbf{i}\pi kj}{e}}{-2\mathbf{i}\pi e \exp(-2\mathbf{i}\pi j)} &= \frac{\exp \frac{2\mathbf{i}\pi kj}{e}}{2\mathbf{i}\pi e} - \frac{\exp \frac{-2\mathbf{i}\pi kj}{e}}{-2\mathbf{i}\pi e} \\ &= \frac{1}{\mathbf{i}\pi e} \cos \frac{2\pi kj}{e}. \end{aligned}$$

Now the following meromorphic function over \mathbb{C} :

$$g_2 : z \mapsto \cot \pi \left(z + \frac{1}{2e} \right) - \cot \pi \left(z - \frac{1}{2e} \right),$$

has simple poles at each $\pm \frac{1}{2e} + \mathbb{Z}$. Note that since $e \geq 2$ then each pole is simple indeed. Recalling that $z \cot z \sim 1$ as $z \rightarrow 0$, we find that the residue at $\pm \frac{1}{2e} + \mathbb{Z}$ is $\pm \frac{1}{\pi}$. We thus find that, with $g := g_1 g_2$,

$$\begin{aligned} \operatorname{Res}_{\frac{j}{e}} g &= \operatorname{Res}_{\frac{j}{e}} (g_1) g_2 \left(\frac{j}{e} \right) \\ &= \frac{1}{\mathbf{i}\pi e} \cos \frac{2\pi kj}{e} \left[\cot \pi \left(\frac{j}{e} + \frac{1}{2e} \right) - \cot \pi \left(\frac{j}{e} - \frac{1}{2e} \right) \right] \\ &= \frac{b_j}{\mathbf{i}\pi e} \cos \frac{2\pi kj}{e}, \end{aligned}$$

for all $j \in \mathbb{Z}$, and

$$\begin{aligned} \operatorname{Res}_{\pm \frac{1}{2e}} g &= g_1 \left(\frac{\pm 1}{2e} \right) \operatorname{Res}_{\pm \frac{1}{2e}} g_2 \\ &= \pm \frac{1}{\pi} \left(\frac{\exp \frac{\pm \mathbf{i}\pi k}{e}}{-2} - \frac{\exp \frac{\mp \mathbf{i}\pi k}{e}}{-2} \right) \\ &= \mp \frac{\mathbf{i}}{\pi} \sin \frac{\pm \pi k}{e} \\ &= -\frac{\mathbf{i}}{\pi} \sin \frac{\pi k}{e}. \end{aligned}$$

We now consider the integral of $g := g_1 g_2$ on the contour C_R given by the direct rectangle with vertices $\pm \mathbf{i}R$ and $1 \pm \mathbf{i}R$, with the points 0 and 1 avoided from the left by a small semi-circle of radius $< \frac{1}{2e}$ (see Figure 4). Note that the only poles of g inside C_R are the $\frac{j}{e}$ for $0 \leq j < e$, together with $\frac{1}{2e}$ and $1 - \frac{1}{2e}$. Recalling that $\cot(\pi + z) = \cot(z)$, we obtain that g is 1-periodic thus the integral of g on the two vertical sides of C_R vanishes. Now for $x, y \in \mathbb{R}$ we have

$$\frac{\exp \pm 2\mathbf{i}\pi k(x + \mathbf{i}y)}{\exp(\pm 2\mathbf{i}\pi e(x + \mathbf{i}y)) - 1} = \frac{\exp(\pm 2\mathbf{i}\pi kx) \exp(\mp 2\pi ky)}{\exp(\pm 2\mathbf{i}\pi ex) \exp(\mp 2\pi ey) - 1}$$




Figure 4: Contour C_R

that goes to 0 as $|y| \rightarrow +\infty$ (since $1 \leq k < e$) uniformly in x , thus $g_1(x + iy) \xrightarrow{|y| \rightarrow +\infty} 0$ uniformly in $x \in \mathbb{R}$. Moreover, recalling from [OLBC, 4.21.40] that $\cot(a + ib) = \frac{\sin 2a - i \sinh 2b}{\cosh 2b - \cos 2a}$ we have:

$$\cot \pi \left(x + iy \pm \frac{1}{2e} \right) = \frac{\sin 2\pi \left(x \pm \frac{1}{2e} \right) - i \sinh(2\pi y)}{-\cos 2\pi \left(x \pm \frac{1}{2e} \right) + \cosh(2\pi y)}$$

that goes to $\pm i$ as $|y| \rightarrow +\infty$, uniformly in $x \in \mathbb{R}$. We deduce that $g_2(x + iy) \xrightarrow{|y| \rightarrow +\infty} 0$ uniformly in $x \in [0, 1]$ and thus so does $g(x + iy)$. Since the only poles of g are on the real axis, we deduce that $\int_{C_R} g(z) dz = 0$ for any $R > 0$. Now from the residue theorem we obtain that

$$\begin{aligned} 0 &= \operatorname{Res}_{\frac{1}{2e}} g + \operatorname{Res}_{1 - \frac{1}{2e}} g + \sum_{j=0}^{e-1} \operatorname{Res}_{\frac{j}{e}} g \\ &= -\frac{2i}{\pi} \sin \frac{\pi k}{e} + \frac{1}{i\pi e} \sum_{j=0}^{e-1} b_j \cos \frac{2\pi k j}{e}, \end{aligned}$$

thus, recalling (5.8),

$$\lambda_k = 2e \sin \frac{\pi k}{e}.$$

□

Let $N = (N_i)_{0 \leq i < e}$ be a centred normal (column) vector with covariance matrix B and let Q be an orthogonal matrix such that $QBQ^\top = D$ with $D := \operatorname{diag}(\lambda_j)_{0 \leq j < e}$. The random vector $(N'_i)_{0 \leq i < e}$ given by $N' = QN$ is a centred normal vector with covariance matrix D and we have $\sum_{i=0}^{e-1} N_i^2 = \sum_{i=0}^{e-1} N_i'^2$. Now $N'_i \simeq \mathcal{N}(0, \lambda_i)$ is a centred normal distribution with variance λ_i , moreover if $i \neq j$ then N'_i and N'_j are independent. By Lemma 5.7 we have $\lambda_0 = 0 < \lambda_i$ for all $1 \leq i < e$, in particular $N_0 = 0$ almost surely and each $N_i'^2$ for $i \geq 1$ has a Gamma distribution $\Gamma(\frac{1}{2}, 2\lambda_i)$ with shape $\frac{1}{2}$ and scale $2\lambda_i$. We deduce the main result of the paper (Theorem C of the introduction).

Theorem 5.9. *The rescaled size $\frac{\pi}{4\sqrt{t}}|\bar{\lambda}|$ of the e -core of λ has, under the Poissonised Plancherel measure pl_t , asymptotically as $t \rightarrow +\infty$ the distribution of a sum of $e-1$ mutually independent $\Gamma(\frac{1}{2}, \sin \frac{k\pi}{e})$ for $k \in \{1, \dots, e-1\}$.*

Proof. We saw at the beginning of §5.3 that $t^{-1/2}|\bar{\lambda}|$ has asymptotically the same law as $t^{-1/2} \frac{e}{2} \sum_{i=0}^{e-1} x_i(\lambda)^2$. By Theorem 5.5 and the preceding discussion, since the vector




Figure 5: Case $e = 4$. Convergence to $\Gamma(1, \frac{1}{\sqrt{2}}) + \Gamma(\frac{1}{2}, 1)$.




Figure 6: Case $e = 7$. Convergence to $\sum_{k=1}^3 \Gamma(1, \sin \frac{k\pi}{7})$.

$t^{-1/4} e \sqrt{\frac{\pi}{2}} (x_i(\lambda))_i$ converges in distribution to N , we deduce that $t^{-1/2} e^2 \frac{\pi}{2} \sum_{i=0}^{e-1} x_i(\lambda)^2$ converges in distribution to a random variable G that is a sum of independent $\Gamma(\frac{1}{2}, 2\lambda_i)$ for $1 \leq i < e$. We deduce that $t^{-1/2} |\bar{\lambda}|$ converges in distribution to $\frac{1}{e\pi} G$. Recalling from Lemma 5.7 that $\lambda_i = 2e \sin \frac{\pi i}{e}$, we have that $\frac{1}{4e} G$ is a sum of independent $\Gamma(\frac{1}{2}, \sin \frac{\pi i}{e})$. This concludes the proof since $\frac{\pi}{4\sqrt{t}} |\bar{\lambda}|$ converges in distribution to $\frac{1}{4e} G$. \square

Remark 5.10. If the Plancherel measure is replaced by the uniform one, then by [LuPi, AySi] the random variable $\frac{\pi}{\sqrt{n}} |\bar{\lambda}|$ converges in distribution to $\Gamma(\frac{e-1}{2}, \sqrt{6})$, which is a sum of $e-1$ independent $\Gamma(\frac{1}{2}, \sqrt{6})$. This situation thus “corresponds” to $\frac{2\lambda_k}{e} = \sqrt{6}$ for all $1 \leq k < e$ (of course this is not clear at all whether $t^{-1/4} x(\lambda)$ converges in distribution to a normal vector in the uniform case).

Remark 5.11. Noting that $\lambda_k = \lambda_{-k}$ for $k \in \mathbb{Z}/e\mathbb{Z}$, we obtain that the sum of $e-1$ mutually independent Gamma distributions in Theorem 5.9 is in fact a sum of $\lfloor \frac{e}{2} \rfloor$ mutually independent Gamma distributions.

We illustrate with Figures 5 and 6 the pointwise convergence of the cumulative distribution functions in Theorem 5.9 for the (non-Poissonised) Plancherel measures Pl_n for $n = 100, 500, 3000$ respectively. Each (renormalised) histogram is constructed from 7000 trials, the range $[0, 5]$ being divided into 200 bins. These simulations indicate that Theorem 5.9 should still hold in this non-Poissonised setting.

5.4 Limit variance

The proof of Theorem 4.1 in [So00-b] shows that the convergence in Theorem 4.1 holds in fact in moments. In particular, the convergence of Theorem 5.9 holds in moments and we recover Proposition 5.1, recalling that $\mathbb{E}\Gamma(k, \theta) = k\theta$ and $\sum_{k=1}^{e-1} \sin \frac{k\pi}{e} = \cot \frac{\pi}{2e}$ (see, for instance, [GrRy, 1.344.1]). We also obtain the following result.

Corollary 5.12. *Under the Poissonised Plancherel measure pl_t , as $t \rightarrow +\infty$ we have*

$$\text{Var}_t |\bar{\lambda}| \sim \frac{4et}{\pi^2}.$$

Proof. Recall that $\text{Var} \Gamma(k, \theta) = k\theta^2$, so that by Theorem 5.9 and the preceding discussion

we have

$$\lim_{t \rightarrow +\infty} \frac{\pi^2}{16t} \text{Var}_t |\bar{\lambda}| = \frac{1}{2} \sum_{k=1}^{e-1} \sin^2 \frac{k\pi}{e}.$$

Now recalling the formula

$$\sum_{k=1}^{e-1} \sin^2 kx = \frac{e-1}{2} - \frac{\cos ex \sin(e-1)x}{2 \sin x},$$

for $x \notin \pi\mathbb{Z}$ (see, for instance, [GrRy, 1.351.1]), we obtain

$$\begin{aligned} \sum_{k=1}^{e-1} \sin^2 \frac{k\pi}{e} &= \frac{e-1}{2} - \frac{\cos \pi \sin(e-1)\frac{\pi}{e}}{2 \sin \frac{\pi}{e}} \\ &= \frac{e-1}{2} + \frac{1}{2} \\ &= \frac{e}{2}, \end{aligned}$$

and thus the desired result. \square

6 Number of i -nodes

For $i \in \mathbb{Z}/e\mathbb{Z}$, we now study the number $c_i(\lambda)$ of i -nodes of λ .

6.1 Poissonised version

Recall from (3.4) that the e -weight of λ is the integer $w_e(\lambda)$ such that

$$|\lambda| = |\bar{\lambda}| + ew_e(\lambda),$$

more precisely by (3.5) we have

$$c_i(\lambda) = c_i(\bar{\lambda}) + w_e(\lambda),$$

for all $i \in \mathbb{Z}/e\mathbb{Z}$. If λ is taken under pl_t and $t \rightarrow +\infty$, we know that $|\lambda|$ follows a Poisson distribution with parameter t thus by the law of large numbers we have that $\frac{|\lambda|}{t}$ converges in distribution to 1 (recalling that $|\lambda|$ is then a sum of t independent Poisson variables of parameter 1). By Theorem 5.9, we know that $\frac{|\bar{\lambda}|}{t}$ converges in distribution to 0, thus by Slutsky's theorem we deduce that $\frac{w_e(\lambda)}{t}$ converges in distribution to $\frac{1}{e}$.

By (3.2) and Theorem 5.9, we know that $\frac{c_i(\bar{\lambda})}{t}$ converges to 0 in distribution. We have just seen that $\frac{w_e(\lambda)}{t}$ converges to $\frac{1}{e}$ in distribution, thus we have proven the following result.

Proposition 6.1. *Let $i \in \mathbb{Z}/e\mathbb{Z}$. Under the Poissonised Plancherel measure pl_t , the random variable $\frac{c_i(\lambda)}{t}$ converges in distribution (and thus in probability) to $\frac{1}{e}$ as $t \rightarrow +\infty$.*

We will now prove the de-Poissonised version of the above result, not by using a de-Poissonisation technique but instead by using the fact that ω_λ converges to a limit shape.

6.2 Law of large numbers

Let $n \geq 1$, $\lambda \in \mathcal{P}_n$ and $i \in \mathbb{Z}/e\mathbb{Z}$. Recall from §2.1 the definition of the function ω_λ . We denote by $\tilde{\omega}_\lambda : \mathbb{R} \rightarrow \mathbb{R}$ the function defined by $\tilde{\omega}_\lambda(s) := \frac{1}{\sqrt{n}} \omega_\lambda(s\sqrt{n})$. Note that the area between the graphs of $\tilde{\omega}_\lambda$ and $|\cdot|$ is 2. It follows from Lemma 3.3 that

$$c_i(\lambda) = \frac{\sqrt{n}}{2} \sum_{k \in \mathbb{Z}} \tilde{\omega}_\lambda \left(\frac{i+ke}{\sqrt{n}} \right) - \left| \frac{i+ke}{\sqrt{n}} \right|. \quad (6.2)$$

Theorem 6.3 ([LoSh], [KeVe], [Rom, Theorem 1.26]). *Let $\Omega : \mathbb{R} \rightarrow \mathbb{R}$ be defined by*

$$\Omega(s) := \begin{cases} \frac{2}{\pi} \left(s \arcsin\left(\frac{s}{2}\right) + \sqrt{4 - s^2} \right), & \text{if } |s| \leq 2, \\ |s|, & \text{otherwise.} \end{cases}$$

Then, under the Plancherel measure Pl_n , the function $\tilde{\omega}_\lambda$ converges uniformly in probability to Ω as $n \rightarrow +\infty$. In other words, for any $\delta > 0$ we have

$$\text{Pl}_n \left(\sup_{\mathbb{R}} |\tilde{\omega}_\lambda - \Omega| > \delta \right) \xrightarrow{n \rightarrow +\infty} 0.$$

Moreover, we also have convergence of the supports, that is:

$$\inf \{ s \in \mathbb{R} : \tilde{\omega}_\lambda(s) \neq |s| \} \rightarrow -2,$$

and

$$\sup \{ s \in \mathbb{R} : \tilde{\omega}_\lambda(s) \neq |s| \} \rightarrow 2,$$

in probability under Pl_n as $n \rightarrow +\infty$.

We can now directly deduce the de-Poissonised version of Proposition 6.1.

Proposition 6.4. *For any $i \in \mathbb{Z}/e\mathbb{Z}$, the random variable $\frac{c_i(\lambda)}{n}$ converges in probability to $\frac{1}{e}$ under the Plancherel measure Pl_n as $n \rightarrow +\infty$.*

Proof. Take $i \in \{0, \dots, e-1\}$ and write $\frac{c_i(\lambda)}{n} = S_n^i + R_i(\lambda)$, where

$$S_n^i := \frac{1}{2\sqrt{n}} \sum_{k \in \mathbb{Z}} \Omega \left(\frac{i+ke}{\sqrt{n}} \right) - \left| \frac{i+ke}{\sqrt{n}} \right|,$$

$$R_i(\lambda) := \frac{1}{2\sqrt{n}} \sum_{k \in \mathbb{Z}} \tilde{\omega}_\lambda \left(\frac{i+ke}{\sqrt{n}} \right) - \Omega \left(\frac{i+ke}{\sqrt{n}} \right)$$

(note that both sums are in fact finite). We have

$$S_n^i = \frac{1}{2e} \frac{e}{\sqrt{n}} \sum_{k \in \mathbb{Z}} g \left(\frac{i+ke}{\sqrt{n}} \right),$$

where $g := \Omega - |\cdot|$ is continuous on \mathbb{R} . Since g has compact support, we deduce that S_n^i is a mere Riemann sum thus $(S_n^i)_n$ converges as $n \rightarrow +\infty$ and

$$\lim_{n \rightarrow +\infty} S_n^i = \frac{1}{2e} \int_{\mathbb{R}} g(t) dt = \frac{1}{2e} 2 = \frac{1}{e}$$

(to obtain that $\int_{\mathbb{R}} g(t) dt = 2$ we can just make the explicit calculation, or note from Theorem 6.3 that g is a uniform limit of continuous functions of integral 2).

Thus, it now suffices to prove that $R_i(\lambda)$ converges to 0 in probability. Let $\delta > 0$. By Theorem 6.3, the probability for $\lambda \in \mathcal{P}_n$ to satisfy

$$\|\tilde{\omega}_\lambda - \Omega\|_\infty \leq \delta, \tag{6.5a}$$

$$\text{supp}(\tilde{\omega}_\lambda - |\cdot|) \subseteq [-2 - \delta, 2 + \delta], \tag{6.5b}$$

goes to 1 as $n \rightarrow \infty$. Now if $|k| \geq \frac{(2+\delta)\sqrt{n}}{e} + 1$ then

$$\begin{aligned} \left| \frac{i+ke}{\sqrt{n}} \right| &= \frac{e}{\sqrt{n}} \left| k + \frac{i}{e} \right| \\ &\geq \frac{e}{\sqrt{n}} \left(|k| - \left| \frac{i}{e} \right| \right) \\ &\geq \frac{e}{\sqrt{n}} (|k| - 1) \\ &\geq 2 + \delta, \end{aligned}$$

thus for such a $k \in \mathbb{Z}_{\geq 0}$ we have $(\tilde{\omega}_\lambda - |\cdot|)\left(\frac{i+ke}{\sqrt{n}}\right) = 0$. We deduce that for λ satisfying (6.5) we have

$$\begin{aligned} |R_i(\lambda)| &\leq \frac{1}{2\sqrt{n}} \sum_{k \in \mathbb{Z}} \left| \tilde{\omega}_\lambda \left(\frac{i+ke}{\sqrt{n}} \right) - \Omega \left(\frac{i+ke}{\sqrt{n}} \right) \right| \\ &= \frac{1}{2\sqrt{n}} \sum_{|k| < \frac{(2+\delta)\sqrt{n}}{e} + 1} \left| \tilde{\omega}_\lambda \left(\frac{i+ke}{\sqrt{n}} \right) - \Omega \left(\frac{i+ke}{\sqrt{n}} \right) \right| \\ &\leq \frac{1}{2\sqrt{n}} \left(\frac{2(2+\delta)\sqrt{n}}{e} + 3 \right) \delta \\ &\leq \frac{(2+\delta)\delta}{e} + \frac{3\delta}{2\sqrt{n}}. \\ &\leq \frac{(2+\delta)\delta}{e} + \frac{3\delta}{2}. \end{aligned}$$

We thus have proved that $\text{Pl}_n(|R_i(\tilde{\omega}_\lambda)|) > \frac{2(2+\delta)\delta}{e} + \frac{3\delta}{2} \xrightarrow{n \rightarrow \infty} 0$ and this concludes the proof since $\frac{2(2+\delta)\delta}{e} + \frac{3\delta}{2} \xrightarrow{\delta \rightarrow 0} 0$. \square

A Intermediate proofs

We provide here the proofs of Lemma 4.28 and Lemma 4.39.

A.1 A conditionally convergent series

We prove here Lemma 4.28. Let $k \in \mathbb{Z}$ with $k \notin e\mathbb{Z}$. With $E_k^{[e]}(x) := \sum_{n \in e\mathbb{Z}+k} \frac{\exp 2inx}{n} = 2 \sum_{n \in 2e\mathbb{Z}+2k} \frac{\exp inx}{n}$, we want to prove that

$$E_k^{[e]}(x) = \frac{\mathbf{i}}{e} \sum_{\omega \in \Omega_{2e}} \exp(2\mathbf{i}k\omega)(\omega - \epsilon_{x,\omega}\pi),$$

for any $x \in (-\pi, \pi) \setminus \Omega_{2e}$, where $\epsilon_{x,\omega} = \text{sgn}(\omega - x)$ and $\Omega_{2e} = \left\{ \frac{\ell\pi}{e} : -e < \ell \leq e \right\} \subseteq (-\pi, \pi]$. In particular, with $\omega_\ell = \frac{\ell\pi}{e}$ we have $\Omega_{2e} = \{\omega_\ell\}_{-e < \ell \leq e}$ and the above equality shows that $E_k^{[e]}$ is piecewise constant on each interval $(\omega_\ell, \omega_{\ell+1})$ for $-e \leq \ell < e$.

We denote by $\ln : \mathbb{C} \setminus \mathbb{R}_- \rightarrow \{z \in \mathbb{C} : -\pi < \arg z < \pi\}$ the principal value of the logarithm. Recall that for any $z \in \mathbb{C}$ with $|z| \leq 1$ and $z \neq 1$ we have

$$-\ln(1-z) = \sum_{n=1}^{+\infty} \frac{z^n}{n}.$$

As in the proof of Lemma 4.14, we deduce that if moreover $z \notin \mu_{2e}$ then, recalling (4.15),

$$\begin{aligned} - \sum_{\zeta \in \mu_{2e}} \zeta^{-2k} \ln(1-\zeta z) &= \sum_{n=1}^{+\infty} \frac{z^n}{n} \sum_{\zeta \in \mu_{2e}} \zeta^{n-2k} \\ &= 2e \sum_{\substack{n=1 \\ n \in 2e\mathbb{Z}+2k}}^{+\infty} \frac{z^n}{n}. \end{aligned}$$

We deduce that

$$\begin{aligned}
\sum_{n \in 2e\mathbb{Z}+2k} \frac{z^n}{n} &= \sum_{\substack{n=1 \\ n \in 2e\mathbb{Z}+2k}}^{+\infty} \frac{z^n}{n} + \sum_{\substack{n=-\infty \\ n \in 2e\mathbb{Z}+2k}}^{-1} \frac{z^n}{n} \\
&= \sum_{\substack{n=1 \\ n \in 2e\mathbb{Z}+2k}}^{+\infty} \frac{z^n}{n} - \sum_{\substack{n=1 \\ n \in 2e\mathbb{Z}-2k}}^{+\infty} \frac{z^{-n}}{n} \\
&= -\frac{1}{2e} \sum_{\zeta \in \mu_{2e}} \zeta^{-2k} \ln(1 - \zeta z) + \frac{1}{2e} \sum_{\zeta \in \mu_{2e}} \zeta^{2k} \ln(1 - \zeta z^{-1}) \\
&= \frac{1}{2e} \sum_{\zeta \in \mu_{2e}} \zeta^{2k} [\ln(1 - \zeta z^{-1}) - \ln(1 - \zeta^{-1} z)].
\end{aligned}$$

Now if $|z| = 1$, we have $\arg(1 - \zeta z^{-1}), \arg(1 - \zeta^{-1} z) \in (-\pi/2, \pi/2)$ thus

$$-\pi < \arg(1 - \zeta z^{-1}) - \arg(1 - \zeta^{-1} z) < \pi.$$

We deduce that

$$\ln(1 - \zeta z^{-1}) - \ln(1 - \zeta^{-1} z) = \ln \frac{1 - \zeta z^{-1}}{1 - \zeta^{-1} z}.$$

We have $\frac{1 - \zeta z^{-1}}{1 - \zeta^{-1} z} = -\zeta z^{-1}$ thus we obtain

$$\sum_{n \in 2e\mathbb{Z}+2k} \frac{z^n}{n} = \frac{1}{2e} \sum_{\zeta \in \mu_{2e}} \zeta^{2k} \ln(-\zeta z^{-1}),$$

for $|z| = 1$ and $z \notin \mu_{2e}$. With $z = \exp ix$ with $x \in (-\pi, \pi) \setminus \Omega_{2e}$, we obtain that, with $\epsilon_{x,\omega} := \operatorname{sgn}(\omega - x)$,

$$\begin{aligned}
E_k^{[e]}(x) &= 2 \sum_{n \in 2e\mathbb{Z}+2k} \frac{\exp inx}{n} \\
&= \frac{1}{e} \sum_{\omega \in \Omega_{2e}} \exp(2ik\omega) \ln(-\exp i(\omega - x)) \\
&= \frac{1}{e} \sum_{\omega \in \Omega_{2e}} \exp(2ik\omega) \ln[\exp i(\omega - x - \epsilon_{x,\omega}\pi)].
\end{aligned}$$

We have $x \in (-\pi, \pi)$ and $\omega \in (-\pi, \pi]$ thus $\omega - x - \epsilon_{x,\omega}\pi \in (-\pi, \pi)$ thus

$$\ln[\exp i(\omega - x - \epsilon_{x,\omega}\pi)] = i(\omega - x - \epsilon_{x,\omega}\pi).$$

We deduce that

$$\begin{aligned}
E_k^{[e]}(x) &= \frac{i}{e} \sum_{\omega \in \Omega_{2e}} \exp(2ik\omega)(\omega - x - \epsilon_{x,\omega}\pi) \\
&= \frac{i}{e} \sum_{\omega \in \Omega_{2e}} \exp(2ik\omega)(\omega - \epsilon_{x,\omega}\pi) - \frac{ix}{e} \sum_{\omega \in \Omega_{2e}} \exp(2ik\omega),
\end{aligned}$$

which gives the desired result since

$$\sum_{\omega \in \Omega_{2e}} \exp(2ik\omega) = \sum_{\zeta \in \mu_{2e}} \zeta^{2k} = 0,$$

since $e \nmid k$.

Remark A.1. If $k \in e\mathbb{Z}$, then the sum $E^{[e]}(x) := \sum_{n \in e\mathbb{Z} \setminus \{0\}} \frac{\exp 2inx}{n}$ is not piecewise constant. Indeed, we have

$$E^{[e]}(x) + \sum_{k=1}^{e-1} E_k^{[e]}(x) = \sum_{n \neq 0} \frac{\exp 2inx}{n} = 2 \sum_{n \geq 1} \frac{\sin 2nx}{n} = 2S(2x),$$

where $S(y) := \sum_{n \geq 1} \frac{\sin ny}{n}$. By a standard equality we have $S(y) = \frac{\pi-y}{2}$ for all $y \in (0, 2\pi)$ (see, for instance, [GrRy, 1.441.1]), thus by Lemma 4.28 we know that $E^{[e]}$ is not piecewise constant (but piecewise affine). We can also directly compute $E^{[e]}$ since

$$E^{[e]}(x) = \frac{1}{e} \sum_{n \neq 0} \frac{\sin 2enx}{n} = \frac{2}{e} S(2ex).$$

A.2 Riemann–Lebesgue lemma

We prove here Lemma 4.39. Let f be integrable on (a, b) and ϕ be continuously differentiable on $[a, b]$. We want to prove that if ϕ vanishes at a finite number of points then

$$\int_a^b f(t) \exp i(x\phi(t)) dt \xrightarrow{x \rightarrow +\infty} 0.$$

If ϕ' does not vanish in $[a, b]$ the result reduces to the classical Riemann–Lebesgue lemma after the variable change $u = \phi(t)$. Thus, by additivity it suffices to consider the case where ϕ vanishes only at a . Let $0 < \epsilon \ll 1$ and write

$$\int_a^b f(t) \exp i(x\phi(t)) dt = \int_a^{a+\epsilon} f(t) \exp i(x\phi(t)) dt + \int_{a+\epsilon}^b f(t) \exp i(x\phi(t)) dt.$$

The first integral is bounded in module by the integral of $|f|$ over $(a, a + \epsilon)$, and the second one goes to zero as $x \rightarrow +\infty$ by the first case since ϕ' does not vanish in $[a + \epsilon, b]$. We conclude that for any $\epsilon > 0$ we have

$$\limsup_{x \rightarrow +\infty} \left| \int_a^b f(t) \exp i(x\phi(t)) dt \right| \leq \int_a^{a+\epsilon} |f(t)| dt,$$

and now the quantity in the right-hand side goes to zero as $\epsilon \rightarrow 0$ since f is integrable.

References

- [AySi] A. AYYER and S. SINHA, *The size of t -cores and hook lengths of random cells in random partitions*. SLC **84B** #58 (2020).
- [BKPV] J. BEN HOUGH, M. KRISHNAPUR, Y. PERES and B. VIRÁG, *Determinantal processes and independence*. Probab. Surv. **3** 206–229 (2006).
- [BeYe] B. C. BERNDT and B. P. YEAP, *Explicit evaluations and reciprocity theorems for finite trigonometric sums*. Adv. Appl. Math. **29** 358–385 (2002).
- [Bi] P. BIANE, *Approximate factorization and concentration for characters of symmetric groups*. Internat. Math. Res. Notices **4** 179–192 (2001).
- [BoSu-a] L. V. BOGACHEV and Z. SU, *Central Limit Theorem for Random Partitions under the Plancherel Measure*. Doklady Mathematics **75** (3) 381–384 (2007).
- [BoSu-b] L. V. BOGACHEV and Z. SU, *Gaussian fluctuations of Young diagrams under the Plancherel measure*. Proc. R. Soc. A **263** 1069–1080 (2007).
- [BOO] A. BORODIN, A. OKOUNKOV and G. OLSHANSKI, *Asymptotics of Plancherel measures for symmetric groups*. J. Amer. Math. Soc. **13**(3) 481–515 (2000).

- [CoLe] O. COSTIN and J. LEBOWITZ, *Gaussian fluctuations in random matrices*. Phys. Rev. Lett. **75**(1) 69–72 (1995).
- [Fa] M. FAYERS, *Weights of multipartitions and representations of Ariki–Koike algebras*. Adv. Math. **206**(1) 112–144 (2006). An updated version of this paper is available from <http://www.maths.qmul.ac.uk/~mf/>.
- [FéMé] V. FÉRAY and P.-L. MÉLIOT, *Asymptotics of q -Plancherel measures*. Probab. Theory Relat. Fields **152** 589–624 (2012).
- [GKS] F. GARVAN, D. KIM and D. STANTON, *Cranks and t -cores*. Invent. Math. **101**(1) 1–17 (1990).
- [GrRy] I. S. GRADSHTEYN and I. M. RYZHIK, *Table of integrals, series, and products*. Translated from the Russian. Translation edited and with a preface by Victor Moll and Daniel Zwillinger. 8th updated and revised ed. Amsterdam: Elsevier/Academic Press (2015).
- [IvOl] V. IVANOV and G. OLSHANSKI, *Kerov’s central limit theorem for the Plancherel measure on Young diagrams*. In Symmetric Functions 2001: Surveys of Developments and Perspectives **74** NATO Science Series II, Mathematics, Physics and Chemistry, 93–151 (2002).
- [JaKe] G. JAMES and A. KERBER, *The Representation Theory of the Symmetric Group*. Encyclopedia of Mathematics and its Applications **16**, Addison-Wesley (1981).
- [Jo] K. JOHANSSON, *Discrete orthogonal polynomial ensembles and the Plancherel measure*. Ann. Math. **153** 259–296 (2001).
- [Ke] S. V. KEROV, *Gaussian limit for the Plancherel measure of the symmetric group*. Comptes Rendus Acad. Sci. Paris I **316** 303–308 (1993).
- [KeVe] S. V. KEROV and A. M. VERSHIK, *Asymptotics of the Plancherel measure on the symmetric group and the limiting form of Young tableaux*. Sov. Math. Dokl. **18** 527–531 (1977).
- [LoSh] B. F. LOGAN and L. A. SHEPP, *A variational problem for random Young tableaux*. Adv. Math. **26** 206–222 (1977).
- [LuPi] N. LULOV and B. PITTEL, *On the Random Young Diagrams and Their Cores*. J. Comb. Theory Ser. A **86** 245–280 (1999).
- [Ma] A. MATHAS, *Iwahori–Hecke algebras and Schur algebras of the symmetric group*. Amer. Math. Soc. (1999).
- [Mé10] P.-L. MÉLIOT, *Asymptotics of the Gelfand models of the symmetric groups*. arXiv:1009.4047.
- [Mé11] P.-L. MÉLIOT, *Kerov’s central limit theorem for Schur–Weyl and Gelfand measures (extended abstract)*. Proc. 23rd Internat. Conf. on Formal Power Series and Alg. Comb. (Reykjavík, Iceland) 669–680 (2011).
- [MiUp] M. MITZENMACHER and E. UPFAL, *Probability and Computing: Randomized Algorithms and Probabilistic Analysis*. Cambridge University Press, second edition (2017).
- [Na] T. NAKAYAMA, *On some modular properties of irreducible representations of a symmetric group, I*. Japan. J. Math. **17** 165–184 (1941).
- [Ol] J. B. OLSSON, *Combinatorics and representations of finite groups*. Vorlesungen aus dem Fachbereich Mathematik der Universität Essen **20** (1993).
- [OLBC] F. W. J. OLVER (ed.) et al., *NIST handbook of mathematical functions*. Cambridge: Cambridge University Press (2010).
- [Ri] J. RIORDAN, *Moment Recurrence Relations for Binomial, Poisson and Hypergeometric Frequency Distributions*. Ann. Math. Statist. **8**(2) 103–111 (1937).
- [Rom] D. ROMIK, *The Surprising Mathematics of Longest Increasing Subsequences*. Cambridge University Press (2015).

- [Ros] S. ROSTAM, *Stuttering blocks of Ariki–Koike algebras*. Alg. Comb. **2**(1) 75–118 (2019).
- [Sa] B. E. SAGAN, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, Second Edition. Graduate Texts in Mathematics, Springer (2001).
- [SM] SAGEMATH, the Sage Mathematics Software System. The Sage Developers, <https://www.sagemath.org> (2020).
- [So00-a] A. SOSHNIKOV, *Determinantal random point fields*. Russian Math. Surveys **55**(5) 923–975 (2000).
- [So00-b] A. SOSHNIKOV, *Gaussian Fluctuation for the Number of Particles in Airy, Bessel, Sine, and Other Determinantal Random Point Fields*. J. Statist. Phys. **100**(3/4) 491–522 (2000).
- [So02] A. SOSHNIKOV, *Gaussian limit for determinantal random point fields*. Ann. of Probab. **30**(1) 171–187 (2002).
- [Ve] A. M. VERSHIK, *Statistical Mechanics and Combinatorial Partitions, and Their Limit Shapes*. Funct. Anal. Its Appl. **30**(2) 90–105 (1996).