



HAL
open science

The Vlasov-Poisson-Boltzmann/Landau system with polynomial perturbation near Maxwellian

Chuqi Cao, Dingqun Deng, Xingyu Li

► **To cite this version:**

Chuqi Cao, Dingqun Deng, Xingyu Li. The Vlasov-Poisson-Boltzmann/Landau system with polynomial perturbation near Maxwellian. 2024. hal-03427367v3

HAL Id: hal-03427367

<https://hal.science/hal-03427367v3>

Preprint submitted on 28 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE VLASOV-POISSON-BOLTZMANN/LANDAU SYSTEM WITH POLYNOMIAL PERTURBATION NEAR MAXWELLIAN

CHUQI CAO, DINGQUN DENG AND XINGYU LI

ABSTRACT. In this work, we consider the Vlasov-Poisson-Boltzmann system without angular cutoff and the Vlasov-Poisson-Landau system with Coulomb potential near a global Maxwellian μ in torus or union of cubes. We establish the global existence, uniqueness, and large-time behavior for solutions in a polynomial-weighted Sobolev space $H_{x,v}^2((v)^k)$ for some constant $k > 0$. For the domain union of cubes, We will consider the specular-reflection boundary condition and its high-order compatible specular boundary condition. The proof is based on an extra dissipation term generated from an improved semigroup method including the electrostatic field with the help of macroscopic estimates.

CONTENTS

1. Introduction	1
1.1. Models and equations	1
1.2. Reformulation	2
1.3. Domains	3
1.4. Notations	4
1.5. Main results	5
1.6. Strategies and ideas of the proof	6
1.7. Organization of the paper	9
2. Basic estimates	9
2.1. Preliminaries on the Boltzmann/Landau equation	9
2.2. High-order specular boundary conditions	13
2.3. Estimates on the weight	13
2.4. Weighted estimates	14
3. Global regularity	21
3.1. Macroscopic estimate	21
3.2. Estimates on the energy	25
3.3. Recover the energy from semigroup method	30
3.4. Proof of the main theorem	36
References	38

1. INTRODUCTION

1.1. Models and equations. We consider the Vlasov-Poisson-Boltzmann (VPB) and Vlasov-Poisson-Landau (VPL) systems describing the motion of plasma particles of two species in bounded domain $\Omega \subseteq \mathbb{R}^3$ (cf. [41, 44]):

$$\begin{aligned} \partial_t F_+ + v \cdot \nabla_x F_+ - \nabla_x \phi \cdot \nabla_v F_+ &= Q(F_+, F_+) + Q(F_-, F_+), \\ \partial_t F_- + v \cdot \nabla_x F_- + \nabla_x \phi \cdot \nabla_v F_- &= Q(F_+, F_-) + Q(F_-, F_-), \\ F(0) &= F_0, \quad E(0) = E_0, \end{aligned} \tag{1.1}$$

here $F(t, x, v) \geq 0$ is a distribution function of particles at time $t > 0$ with position $x \in \Omega$ and velocity $v \in \mathbb{R}^3$, $E = -\nabla_x \phi$ is the self-consistent electrostatic field satisfying

$$-\Delta_x \phi = \int_{\mathbb{R}^3} (F_+ - F_-) dv, \quad \int_{\Omega} \phi(t, x) dx = 0. \tag{1.2}$$

2020 *Mathematics Subject Classification.* 35A23; 35J05; 35Q20.

Key words and phrases. Vlasov-Poisson-Boltzmann system, Vlasov-Poisson-Landau system, global existence, polynomial weighted space.

For the VPB system, the bilinear collision operator Q acts only on the velocity variable v given by

$$Q(G, F)(v) = \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} B(v - v_*, \sigma) (G'_* F' - G_* F) d\sigma dv_*.$$

Here we use the standard notation $F = F(v)$, $G_* = G(v_*)$, $F' = F(v')$, $G'_* = G(v'_*)$, where v', v'_* are velocities of two particles after collision given by

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma, \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma, \quad \sigma \in \mathbb{S}^2.$$

This representation follows from the physical law of elastic collision:

$$v + v_* = v' + v'_*, \quad |v|^2 + |v_*|^2 = |v'|^2 + |v'_*|^2.$$

The nonnegative function $B(v - v_*, \sigma)$ is called the Boltzmann collision kernel. It depends only on relative velocity $|v - v_*|$ and the deviation angle θ through $\cos \theta \stackrel{\text{def}}{=} \frac{v - v_*}{|v - v_*|} \cdot \sigma$.

In the present work, we consider the non-cutoff kernel B as the following.

(A1). The Boltzmann kernel B takes the form $B(v - v_*, \sigma) = |v - v_*|^\gamma b(\frac{v - v_*}{|v - v_*|} \cdot \sigma)$, where b is a nonnegative angular function.

(A2). $b(\cos \theta)$ is not locally integrable and there exists some constant $\mathcal{K} > 0$, such that

$$\mathcal{K} \theta^{-1-2s} \leq \sin \theta b(\cos \theta) \leq \mathcal{K}^{-1} \theta^{-1-2s}, \quad \text{with } 0 < s < 1.$$

(A3). The parameter γ and s satisfy the condition $-3 < \gamma \leq 1$, $1/2 \leq s \leq 1$, $\gamma + 2s > -1$.

(A4). Without loss of generality, we may assume that $B(v - v_*, \sigma)$ is supported in the set $0 \leq \theta \leq \pi/2$, i.e. $\frac{v - v_*}{|v - v_*|} \cdot \sigma \geq 0$. Otherwise, B can be replaced by its symmetrized form:

$$\bar{B}(v - v_*, \sigma) = |v - v_*|^\gamma \left(b\left(\frac{v - v_*}{|v - v_*|} \cdot \sigma\right) + b\left(\frac{v - v_*}{|v - v_*|} \cdot (-\sigma)\right) \right) 1_{\frac{v - v_*}{|v - v_*|} \cdot \sigma \geq 0},$$

where 1_A is the indicator function of the set A .

Remark 1.1. For inverse repulsive potential, we have $\gamma = \frac{p-5}{p-1}$ and $s = \frac{1}{p-1}$ with $p > 2$. It is worth noting that the condition $\gamma + 2s > -1$ is met across the entire spectrum of the inverse power law model. Generally, the cases $\gamma > 0$, $\gamma = 0$, and $\gamma < 0$ correspond to so-called hard, Maxwellian, and soft potentials respectively.

For the case of the VPL system, Q is the Landau collision operator given by

$$Q(g, f)(v) = \partial_i \int_{\mathbb{R}^3} \phi^{ij}(v - v_*) (g_* \partial_j f - f \partial_j g_*) dv_*, \quad \phi^{ij}(v) = |v|^{\gamma+2} \left(\delta_{ij} - \frac{v_i v_j}{|v|^2} \right), \quad -3 \leq \gamma \leq 1.$$

We use the convention of summation for repeated indices, and the derivatives are in the velocity variable $\partial_i = \partial_{v_i}$. Define the notations $g_* = g(v_*)$, $f = f(v)$, $\partial_j g_* = \partial_{v_{*j}} g(v_*)$, $\partial_j f = \partial_{v_j} f(v)$. Notice that

$$\partial_j \phi^{ij}(v) = -2|v|^\gamma v_i, \quad \partial_{ij} \phi^{ij}(v) = \begin{cases} -2(\gamma + 3)|v|^\gamma, & \text{if } -3 < \gamma \leq 1, \\ -8\pi \delta_0, & \text{if } \gamma = -3, \end{cases}$$

where δ_0 is the Dirac measure. We call it *hard potential* if $\gamma \geq 0$, and *soft potential* if $\gamma \in [-3, 0)$.

1.2. Reformulation. We reformulate the VPB/VPL system near a global Maxwellian. For simplicity, we assume the initial data F_0 is normalized such that the equilibrium associated with (1.1) will be the standard Gaussian function, i.e. $\mu(v) \stackrel{\text{def}}{=} (2\pi)^{-3/2} e^{-|v|^2/2}$, which enjoys the same mass, momentum, and energy as F_0 . For the perturbation framework, we denote $F = [F_+, F_-]$ and let $f = [f_+, f_-]$ satisfies $F_\pm = \mu + f_\pm$. Then system (1.1) and (1.2) become

$$\begin{cases} \partial_t f_\pm + v \cdot \nabla_x f_\pm \mp \nabla_x \phi \cdot \nabla_v f_\pm \pm \nabla_x \phi \cdot v \mu = Q(f_\pm + f_\mp, \mu) + Q(2\mu + f_\pm + f_\mp, f_\pm), \\ -\Delta_x \phi = \int_{\mathbb{R}^3} (f_+ - f_-) dv, \quad \int_{\Omega} \phi(x) dx = 0, \quad f(0) = f_0, \quad \phi(0) = \phi_0. \end{cases} \quad (1.3)$$

Note that $Q(\mu, \mu) = 0$. We also denote linear operator $L = [L_+, L_-]$ and $\mathcal{L} = [\mathcal{L}_+, \mathcal{L}_-]$ by

$$L_\pm f = 2Q(\mu, f_\pm) + Q(f_\pm + f_\mp, \mu), \quad \mathcal{L} f = -v \cdot \nabla_x f \mp \nabla_x \phi \cdot v \mu + Lf, \quad (1.4)$$

where $\phi(x)$ is solved by the second equation of (1.3). The kernel of L on $L_v^2 \times L_v^2$ is the span of $\{[1, 0]\mu, [0, 1]\mu, [1, 1]v\mu, [1, 1]|v|^2\mu\}$ (cf. [41]) and we define the projection of $L_v^2 \times L_v^2$ onto $\ker L$ by

$$\mathbf{P}f = \left(a_+(t, x)[1, 0] + a_-(t, x)[0, 1] + v \cdot b(t, x)[1, 1] + (|v|^2 - 3)c(t, x)[1, 1] \right) \mu, \quad (1.5)$$

or equivalently by

$$\mathbf{P}_\pm f = \left(a_\pm(t, x) + v \cdot b(t, x) + (|v|^2 - 3)c(t, x) \right) \mu,$$

where function a_\pm, b, c are given by

$$a_\pm = \int_{\mathbb{R}^3} f_\pm dv, \quad b_j = \frac{1}{2} \int_{\mathbb{R}^3} v_j (f_+ + f_-) dv, \quad c = \int_{\mathbb{R}^3} \frac{|v|^2 - 3}{12} (f_+ + f_-) dv. \quad (1.6)$$

Taking inner produce of the first equation of (1.3) with 1 over \mathbb{R}_v^3 , we have the continuity equation

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^3} f_\pm(v) dv + \nabla_x \cdot \int_{\mathbb{R}^3} v f_\pm(v) dv = 0. \quad (1.7)$$

1.3. Domains. In this paper, we suppose the bounded domain Ω is either a torus or the union of finite cubes.

Torus. For the case of a torus, we set $\Omega = \mathbb{T}^3 = [-\pi, \pi]^3$. In this case, the solution (F, ϕ) to the VPB/VPL system (1.1) with the initial data F_0 enjoys the conservation of mass, momentum, and the energy, i.e.

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} \int_{\mathbb{R}^3} F_+ dv dx &= \frac{d}{dt} \int_{\Omega} \int_{\mathbb{R}^3} F_- dv dx = 0, & \frac{d}{dt} \int_{\Omega} \int_{\mathbb{R}^3} v (F_+ + F_-) dv dx &= 0, \\ \frac{d}{dt} \int_{\Omega} \int_{\mathbb{R}^3} |v|^2 (F_+ + F_-) dv dx &+ \frac{d}{dt} \int_{\Omega} |\nabla_x \phi(t, x)|^2 dx &= 0. \end{aligned} \quad (1.8)$$

Suppose f_0 have the same mass, momentum and energy as μ , then conservation laws (1.8) yield

$$\begin{aligned} \int_{\Omega} \int_{\mathbb{R}^3} f_+ dv dx &= \int_{\Omega} \int_{\mathbb{R}^3} f_- dv dx = 0, & \int_{\Omega} \int_{\mathbb{R}^3} v (f_+ + f_-) dv dx &= 0, \\ \int_{\Omega} \int_{\mathbb{R}^3} |v|^2 (f_+ + f_-) dv dx &+ \int_{\Omega} |\nabla_x \phi(t, x)|^2 dx &= 0. \end{aligned} \quad (1.9)$$

Union of cubes. The second type of bounded domain we consider is a union of finitely many cubes, denoted as:

$$\Omega = \cup_{i=1}^N \Omega_i, \quad (1.10)$$

where each Ω_i is defined as a rectangular region: $\Omega_i = (a_{i,1}, b_{i,1}) \times (a_{i,2}, b_{i,2}) \times (a_{i,3}, b_{i,3})$, with $a_{i,j}, b_{i,j} \in \mathbb{R}$ satisfying $a_{i,j} < b_{i,j}$. The boundary of this domain, denoted as $\partial\Omega$, is composed of three types of boundaries, Γ_i ($i = 1, 2, 3$), each of which is orthogonal to one of the coordinate axes x_i . These boundaries are further divided into connected sets. It is important to note that we assume that each Γ_i has a non-zero spherical measure, while the boundaries of these sets themselves have zero spherical measure. Therefore, we do not distinguish between Γ_i and their interiors.

On the interior of Γ_i ($i = 1, 2, 3$), we have a unit normal outer vector $n(x)$ defined almost everywhere with respect to spherical measure. This vector takes the form $n(x) = e_i$ or $-e_i$, where e_i is a unit vector with its i th component equal to 1. Additionally, we denote vectors $\tau_1(x)$ and $\tau_2(x)$ on the boundary $\partial\Omega$ such that $(n(x), \tau_1(x), \tau_2(x))$ forms a unit orthonormal basis for \mathbb{R}^3 . Furthermore, for $j = 1, 2$, τ_j takes the values e_k or $-e_k$ for some k .

The boundary of the phase space is denoted as $\gamma := \{(x, v) \in \partial\Omega \times \mathbb{R}^3\}$. We define $n = n(x)$ to represent the outward normal direction at $x \in \partial\Omega$. This allows us to partition γ into three distinct sets:

$$\begin{aligned} \gamma_- &= \{(x, v) \in \partial\Omega \times \mathbb{R}^3 : n(x) \cdot v < 0\}, & (\text{the incoming set}), \\ \gamma_+ &= \{(x, v) \in \partial\Omega \times \mathbb{R}^3 : n(x) \cdot v > 0\}, & (\text{the outgoing set}), \\ \gamma_0 &= \{(x, v) \in \partial\Omega \times \mathbb{R}^3 : n(x) \cdot v = 0\}, & (\text{the grazing set}). \end{aligned}$$

For these sets, we assume that $F(t, x, v)$ satisfies a *specular-reflection* boundary condition on γ_- , where $(x, v) \in \gamma$ is subject to the following reflection operation:

$$R_x v = v - 2n(x)(n(x) \cdot v). \quad (1.11)$$

This specular reflection condition for f can be expressed as:

$$f(t, x, R_x v) = f(t, x, v), \quad \text{on } \gamma_-. \quad (1.12)$$

Regarding the boundary condition for the electric potential ϕ , we further assume that it satisfies a Neumann boundary condition:

$$\partial_n \phi = 0, \quad \text{on } x \in \partial\Omega. \quad (1.13)$$

In particular, the Poisson equation for potential ϕ becomes a pure Neumann boundary problem. To ensure the existence of a solution, we impose a zero-mean condition:

$$\int_{\Omega} \int_{\mathbb{R}^3} (f_+ - f_-) dv dx = 0, \quad \text{for } t \geq 0,$$

which follows from the conservation laws (1.14). Similar to (1.9), it's also well-known that the solution to (1.3) in the bounded domain Ω given by (1.10) satisfies the conservation laws on mass and energy. Specifically, the solution f to (1.3) satisfies the following identities when the initial data f_0 satisfies them:

$$\begin{aligned} \int_{\Omega \times \mathbb{R}^3} f_+(t) dv dx &= \int_{\Omega \times \mathbb{R}^3} f_-(t) dv dx = 0, \\ \int_{\Omega \times \mathbb{R}^3} (f_+(t) + f_-(t)) |v|^2 dv dx + \int_{\Omega} |\nabla_x \phi(t, x)|^2 dx &= 0. \end{aligned} \quad (1.14)$$

1.4. Notations. Let us first introduce the function spaces and notations. We let the multi-indices α and β be $\alpha = [\alpha_1, \alpha_2, \alpha_3]$, $\beta = [\beta_1, \beta_2, \beta_3]$ and define $\partial_{\beta}^{\alpha} := \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \partial_{x_3}^{\alpha_3} \partial_{v_1}^{\beta_1} \partial_{v_2}^{\beta_2} \partial_{v_3}^{\beta_3}$.

If each component of α' is not greater than that of the α 's, we denote by $\alpha' \leq \alpha$. $\alpha' < \alpha$ means $\alpha' \leq \alpha$, and $|\alpha'| < |\alpha|$. We write $a \lesssim b$ ($a \gtrsim b$) to indicate that there is a uniform constant C , which may be different on different lines, such that $a \leq Cb$ ($a \geq Cb$). We use the notation $a \sim b$ if $a \lesssim b$ and $b \lesssim a$. We denote $a \gg b$ if a, b are two constants such that $a > b$ and a is sufficiently large.

We denote C_{a_1, a_2, \dots, a_n} by a constant depending on parameters a_1, a_2, \dots, a_n . Moreover, we use the parameter ε to represent different positive numbers much less than 1 and determined in different cases. We use $(f, g) = (f, g)_{L_v^2}$ to denote the inner product of f, g over velocity variable for short and, $(f, g)_{L_{x,v}^2}$ to represent the inner product over both spatial and velocity variables. Also, we write

$$(f, g)_{L_k^2} = (\langle v \rangle^{2k} f, g)_{L_v^2}, \quad \|f(\theta)\|_{L_{\theta}^1} = \int_{\mathbb{S}^2} f(\theta) d\sigma = 2\pi \int_0^{\pi} f(\theta) \sin \theta d\theta,$$

and for any $p \in [1, \infty]$, $\|f(t)\|_{L_T^p} = \|f(t)\|_{L^p([0, T])}$. Define the **Japanese brackets** by $\langle v \rangle := (1 + |v|^2)^{1/2}$.

For linear operator \mathcal{L} , we write $S_{\mathcal{L}}$ as the semigroup generated by \mathcal{L} . For real numbers m, l , we define the weighted Sobolev norm $\|\cdot\|_{H_l^m}$ by $\|f\|_{H_l^m} = \|\langle v \rangle^l \langle D_v \rangle^m f(v)\|_{L_v^2}$, and $L_l^2 = H_l^0$ for $m = 0$. here $a(D)$ is a pseudo-differential operator with the symbol $a(\xi)$, and is defined by

$$(a(D)f)(v) := \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} e^{i(v-u)\xi} a(\xi) f(u) du d\xi.$$

The mixed norm $\|\cdot\|_{H_x^s H_l^m}$ is defined as $\|f\|_{H_x^s H_l^m} := \left(\int_{\Omega} \|\langle D_x \rangle^s f(x, \cdot)\|_{H_l^m}^2 dx \right)^{1/2}$, and $H_x^0 H_l^m = L_x^2 H_l^m$.

The entropy $L \log L$ space is defined as $L \log L := \left\{ f(v) : \|f\|_{L \log L} = \int_{\mathbb{R}^3} |f| \log(1 + |f|) dv \right\}$.

For any $k \in \mathbb{R}$, $\gamma \in (-3, 1]$, we define

$$\|f\|_{L_{k+\gamma/2, *}}^2 := \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \mu(v_*) |v - v_*|^{\gamma} |f(v)|^2 \langle v \rangle^{2k} dv dv_* \sim \|f\|_{L_{k+\gamma/2}^2}^2.$$

For Boltzmann case, we denote the dissipation norm $L_{D, k}^2$ by

$$\|f\|_{L_{D, k}^2} = \|f\|_{H_{k+\gamma/2}^s}, \quad \|f\|_{L_D^2} = \|f\|_{L_{D, 0}^2}.$$

For Landau case, we denote the anisotropic norm $L_D^2(m)$ by

$$\|f\|_{L_D^2(m)} := \|f\|_{L^2(m\langle v \rangle^{\gamma/2})} + \|\tilde{\nabla}_v(mf)\|_{L^2(\langle v \rangle^{\gamma/2})}$$

and for brevity, we let $L_{D, k}^2 := L_D^2(\langle v \rangle^k)$ and $L_D^2 := L_{D, 0}^2$. Here $\tilde{\nabla}_v$ is the anisotropic gradient given by

$$\tilde{\nabla}_v f := P_v \nabla_v f + \langle v \rangle (I - P_v) \nabla_v f, \quad P_v \xi := \left(\xi \cdot \frac{v}{|v|} \right) \frac{v}{|v|}, \quad \forall \xi \in \mathbb{R}^3.$$

With multi-indices (α, β) , we come to define our weight function with some constant k . For $-3 \leq \gamma \leq 1$, we choose the weight function $w(\alpha, \beta)$ as

$$w(\alpha, \beta) = \begin{cases} \langle v \rangle^{k-p|\alpha|-q|\beta|+r}, & q = 6s - 3(\gamma-1), \quad p = q + \gamma - 1, r = 2q \quad \text{for Boltzmann case,} \\ \langle v \rangle^{k-p|\alpha|-q|\beta|+r}, & q = 3 - (\gamma-1), \quad p = 3, r = 2q, \quad \text{for Landau case.} \end{cases} \quad (1.15)$$

Note that $w(\alpha, \beta) \geq 1$ for any $|\alpha| + |\beta| \leq 2$ and any $k \geq 0$. For brevity, we write $w(|\alpha|, |\beta|) = w(\alpha, \beta)$ throughout the paper. Noticing $w^2(\alpha, \beta)$ is still of the form $\langle v \rangle^k$ for some $k > 0$, we have

$$\nabla_v(w^2(\alpha, \beta)) = A_{\alpha, \beta} \frac{v}{\langle v \rangle^2} w^2(\alpha, \beta), \quad (1.16)$$

here $A_{\alpha, \beta}$ depends on $\alpha, \beta, k, \gamma, s$. In this work, we will apply a useful space-velocity weight

$$e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \quad (1.17)$$

to eliminate the dissipation loss term. Next, we define some useful norms in our analysis. For this, we denote constants by

$$C_{|\alpha|, |\beta|} \gg C_{|\alpha|, |\beta_1|}, \quad \text{if } |\beta_1| < |\beta|, \quad C_{|\alpha|+1, |\beta|-1} \gg C_{|\alpha|, |\beta|}, \quad (1.18)$$

for any multi-indices α, β . Then for both the Boltzmann case and Landau case, we denote energy norms as

$$\|f\|_{X_k}^2 = \sum_{|\alpha|+|\beta| \leq 2} C_{|\alpha|, |\beta|} \|e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} w(\alpha, \beta) \partial_\beta^\alpha f\|_{L_x^2 L_v^2}^2, \quad (1.19)$$

$$\|f\|_{Y_k}^2 = \sum_{|\alpha|+|\beta| \leq 2} C_{|\alpha|, |\beta|} \|w(\alpha, \beta) \partial_\beta^\alpha f\|_{L_x^2 L_D^2}^2. \quad (1.20)$$

We further define the ‘‘instant energy functional’’ $\mathcal{E}_k(t)$ and ‘‘dissipation energy functional’’ $\mathcal{D}_k(t)$ as

$$\mathcal{E}_k(t) \approx \|f\|_{X_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2, \quad (1.21)$$

$$\mathcal{D}_k(t) := \|f\|_{Y_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2, \quad (1.22)$$

where the explicit definition of $\mathcal{E}_k(t)$ is given in (3.90).

We fix the weight index k_0 by assuming $k_0 \geq 14$ for the Boltzmann case, and $k_0 \geq 7$ for the Landau case.

1.5. Main results.

We may now state our main results.

Theorem 1.2 (Global existence, uniqueness and large-time decay). *Consider the Cauchy problem (1.3) for Vlasov-Poisson-Boltzmann/Landau system. Suppose $\gamma \in (-3, 1]$, $s \in [\frac{1}{2}, 1)$, $\gamma + 2s > -1$ for the Boltzmann case and $\gamma \in [-3, 1]$ for the Landau case. Let $l = 0$ for the hard potential case and $l > \frac{|\gamma|}{2}$ for the soft potential case. Then there exist constants $\lambda > 0$, $M > 0$ (small) such that, for any*

$$k \geq k_0 + 4 + l \quad \text{for Landau case and} \quad k \geq k_0 + 4 + l \quad \text{for Boltzmann case.}$$

Remind $\mathcal{E}_k(t)$ is given in (1.21). If the initial data f_0 satisfies $F_0(x, v) = \mu + f_0(x, v) \geq 0$, conservation laws (1.9) for the case of torus (or (1.14) for the case of union of cubes) and

$$\mathcal{E}_{k_0+l}(0) \leq M \quad \text{and} \quad \mathcal{E}_k(0) < +\infty, \quad (1.23)$$

then there exists a unique solution $f(t, x, v)$ to (1.3) satisfying $F(t, x, v) = \mu + f(t, x, v) \geq 0$ such that $\sup_{0 \leq t \leq T} \mathcal{E}_k(t) \leq 2\mathcal{E}_k(0)$ for any $T > 0$. Moreover, we have the following large-time asymptotic behavior:

$$\mathcal{E}_k(t) \leq e^{-\lambda t} \mathcal{E}_k(0) \quad \text{for } \gamma \geq 0, \quad \mathcal{E}_{k-l}(t) \lesssim (1+t)^{-\frac{2l}{|\gamma|}} \mathcal{E}_k(0) \quad \text{for } \gamma < 0.$$

Remark 1.3. (1) *It is crucial to emphasize that for global solutions with polynomial weight, the requirement for smallness of initial data is only in relation to a specific norm. In this context, we solely need X_{k_0+l} to be sufficiently small, rather than necessitating the general X_k to be small.*

(2) *Notably, the cases involving the torus and the union of cubes exhibit similarities. In the case of the union of cubes, as defined in (1.10), we exercise meticulous care when performing integration by parts. Conversely, integration by parts in the torus case is a straightforward process.*

(3) *In the scenario of a soft potential with $\gamma < 0$ and considering the non-cutoff Boltzmann equation, the work presented in [8] demonstrates that the optimal decay rate for the non-cutoff Boltzmann equation cannot exceed $k/|\gamma|$. Consequently, when the value of k is sufficiently large, the decay rate articulated in Theorem 1.2 can be regarded as approximately optimal.*

(4) *Our proof methodology can also be effectively employed to establish results for the high regularity space $H_{x,v}^N(\langle v \rangle^k)$.*

1.6. Strategies and ideas of the proof. In this subsection, we provide a brief overview of relevant literature and outline the key strategies employed in proving our results.

We begin by revisiting established findings in the context of the Landau and Boltzmann equations, with a particular focus on the aspects central to this paper—namely, the global existence and long-term behavior of solutions to spatially inhomogeneous equations within the perturbation framework. To shed light on the global solutions of the renormalized equation, particularly in the presence of substantial initial data, we draw attention to classic works such as [5, 20–22, 56, 66, 67].

For insights into the smoothing effects in the Boltzmann equation without cut-off, we refer to [13, 14]. Additionally, when considering the stability of vacuum states, we consult references [12, 40, 57], which cover scenarios involving the Landau equation, the cutoff Boltzmann equation, and the non-cutoff Boltzmann equation, with a focus on moderate soft potentials.

These references provide the foundation upon which we build our exploration of spatially inhomogeneous equations and perturbation techniques to address the questions at hand.

In the context of the non-cutoff Boltzmann equation, prior research efforts like [50–54, 60] have achieved global regularity and elucidated long-term behavior under remarkably broad assumptions. They have demonstrated that global regularity and long-term behavior can be established by merely assuming uniform bounds in both time (t) and space (x), characterized by:

$$0 < m_0 \leq M(t, x) \leq M_0, \quad E(t, x) \leq E_0, \quad H(t, x) \leq H_0,$$

where m_0 , M_0 , E_0 , and H_0 are positive constants, and $M(t, x)$, $E(t, x)$, and $H(t, x)$ are defined as:

$$M(t, x) = \int_{\mathbb{R}^3} f(t, x, v) dv, \quad E(t, x) = \int_{\mathbb{R}^3} f(t, x, v) |v|^2 dv, \quad H(t, x) = \int_{\mathbb{R}^3} f(t, x, v) \ln f(t, x, v) dv.$$

Furthermore, in the case of the Landau equation, local Hölder estimates are established in [32], while higher regularity properties of solutions are explored in [49] through the application of kinetic variants of Schauder estimates.

Subsequently, we turn our attention to the outcomes achieved within the perturbation framework. In the vicinity of the Maxwellian distribution, we find substantial progress in demonstrating the global existence and long-term behavior of solutions to spatially inhomogeneous equations. These achievements are well-documented, with [37, 38, 63, 64] presenting results for the cutoff Boltzmann equation and [36] addressing the Landau equation within this context.

For the non-cutoff Boltzmann equation, substantial contributions can be found in [1–4, 33, 34]. A more recent perspective on this topic can be found in [27], where recent developments are explored. Additionally, for insights into the theory of existence in bounded domains, references [43, 45, 46] are valuable resources.

Furthermore, the non-cutoff case within the union of cubes is examined in [17]. Importantly, all these works are founded on the following decomposition:

$$\partial_t f + v \cdot \nabla_x f = L_\mu f + \Gamma(f, f), \quad L_\mu f = \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu} f, \mu) + \frac{1}{\sqrt{\mu}} Q(\mu, \sqrt{\mu} f),$$

where

$$\Gamma(g, f) = \frac{1}{2} \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu} g, \sqrt{\mu} f) + \frac{1}{2} \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu} f, \sqrt{\mu} g),$$

signifying that the solution is constructed within a $\mu^{-1/2}$ weighted space.

In the realm of inhomogeneous Boltzmann/Landau equations with polynomial weight perturbations near the Maxwellian distribution, Gualdani-Mischler-Mouhot (GMM) laid the foundation by first proving global existence and delineating the large-time behavior of solutions featuring polynomial velocity weights. Their groundbreaking work began with the cutoff Boltzmann equation involving a hard potential, as reported in [35]. This method was subsequently extended to encompass the Landau equation, with significant contributions from [10, 11]. For the non-cutoff Boltzmann equation, the achievements in the hard potential scenario were documented in [6, 48], while the soft potential case was addressed in [8].

Moreover, it is noteworthy to acknowledge prior research on the Vlasov-Poisson/Maxwell-Boltzmann/Landau systems in proximity to Maxwellian distributions. For the cutoff hard sphere case, the Vlasov-Poisson/Maxwell-Boltzmann system has been proven in [41, 42, 62], with [29] further exploring the optimal convergence rate. Various other cases under the cutoff assumption have been studied and can be found in [24, 30, 31]. For the Landau equation, pioneering results were established in [44] for the torus case and in [65, 68] for the entire space scenario, with additional insights provided by [25].

In the non-cutoff setting, research endeavors have been directed towards the Vlasov-Poisson-Boltzmann system [26, 69], the Vlasov-Maxwell-Boltzmann system [28], and investigations into the regularizing effect [18]. The implications of bounded domains are discussed in [9, 19, 23]. Importantly, all these endeavors are rooted in the following decomposition:

$$\partial_t f_{\pm} + v \cdot \nabla_x f_{\pm} \mp \nabla_x \phi \cdot \nabla_v f_{\pm} \pm \nabla_x \phi \cdot v \sqrt{\mu} \pm \frac{1}{2} \nabla_x \phi \cdot v f = L_{\mu, \pm} f + \Gamma_{\pm}(f, f),$$

where

$$L_{\mu, \pm} f = \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu}(f_{\pm} + f_{\mp}), \mu) + 2 \frac{1}{\sqrt{\mu}} Q(\mu, \sqrt{\mu} f_{\pm}), \quad (1.24)$$

and

$$\Gamma_{\pm}(f, f) = \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu} f_{\pm}, \sqrt{\mu} f_{\pm}) + \frac{1}{\sqrt{\mu}} Q(\sqrt{\mu} f_{\mp}, \sqrt{\mu} f_{\pm}),$$

demonstrating that these studies are conducted within a $\mu^{-1/2}$ weighted space. To the best of our knowledge, our research represents the inaugural effort to address these questions within the framework of polynomial-weighted spaces.

The foundation of our results lies in the utilization of the semigroup method. This method, originally introduced by Gualdani-Mischler-Mouhot in [35], forms the basis of our work and has been further developed and expanded upon, particularly with respect to the macroscopic component $\mathbf{P}f$. The central concept of this method can be succinctly summarized as follows: Consider the case where $\gamma = 0$ and the function space $H_x^2 L_k^2$. Drawing from [58], we arrive at the following inequality:

$$\sum_{\pm} (\mathcal{L}_{\pm} f, f_{\pm})_{H_x^2 L^2(\mu^{-1/2})} \leq -\lambda \|f\|_{H_x^2 L^2(\mu^{-1/2})}^2, \quad \text{if } \mathbf{P}f = 0,$$

where $\lambda > 0$ serves as a crucial constant. Together with the macroscopic estimates from [19, 33] for $\mathbf{P}f$, we can deduce that $\|S_{\mathcal{L}}(t)f_0\|_{H_x^2 L^2(\mu^{-1/2})} \leq e^{-\lambda t} \|f_0\|_{H_x^2 L^2(\mu^{-1/2})}^2$. Next, for some $M, R > 0$, define

$$A_{\pm} = -v \cdot \nabla_x + L_{\pm} - M\chi_R, \quad K_1 = M\chi_R, \quad B_{\pm} = -v \cdot \nabla_x + L_{\pm}, \quad K_2 = \pm \mu v \cdot \nabla_x \phi,$$

where $\chi \in D(\mathbb{R})$ is the truncation function satisfying $1_{[-1,1]} \leq \chi \leq 1_{[-2,2]}$ and we denote $\chi_R(\cdot) := \chi(\cdot/R)$ for $R > 0$. By the results in [8] we have

$$\sum_{\pm} (L_{\pm} f, f_{\pm})_{H_x^2 L_k^2} \leq -C \|f\|_{H_x^2 H_k^s}^2 + C_k \|f\|_{H_x^2 L_k^v}^2,$$

Take $M, R > 0$ large, we have $(Af, f)_{H_x^2 L_k^2} \leq -C \|f\|_{H_x^2 L_k^2}^2$, which implies $\|S_A(t)f\|_{H_x^2 L_k^2} \leq e^{-\lambda t} \|f_0\|_{H_x^2 L_k^2}$. By Duhamel's formula $S_B = S_A + S_B * K_1 S_A$ and $S_{\mathcal{L}} = S_B + S_{\mathcal{L}} * K_2 S_B$, roughly speaking, we have

$$\begin{aligned} \|S_B(t)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} &\leq \|S_A(t)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} + \int_0^t \|S_B(s)\|_{H_x^2 L^2(\mu^{-1/2}) \rightarrow H_x^2 L^2(\mu^{-1/2})} \\ &\quad \times \|K_1\|_{H_x^2 L_k^2 \rightarrow H_x^2 L^2(\mu^{-1/2})} \|S_A(t-s)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} ds \leq C e^{-\lambda t}. \end{aligned}$$

and hence

$$\begin{aligned} \|S_{\mathcal{L}}(t)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} &\leq \|S_B(t)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} + \int_0^t \|S_{\mathcal{L}}(s)\|_{H_x^2 L^2(\mu^{-1/2}) \rightarrow H_x^2 L^2(\mu^{-1/2})} \\ &\quad \times \|K_2\|_{H_x^2 L_k^2 \rightarrow H_x^2 L^2(\mu^{-1/2})} \|S_B(t-s)\|_{H_x^2 L_k^2 \rightarrow H_x^2 L_k^2} ds \leq C e^{-\lambda t}. \end{aligned}$$

Thus, the rate of convergence for the linear operator \mathcal{L} is established. To estimate the nonlinear part, we need to define a scalar product by

$$((f, g))_k := (f, g)_{L_k^2} + \eta \int_0^{+\infty} (S_{\mathcal{L}}(\tau)f, S_{\mathcal{L}}(\tau)g)_{L_k^2} d\tau.$$

Due to the fact that

$$\int_0^{+\infty} (S_{\mathcal{L}}(\tau)\mathcal{L}f, S_{\mathcal{L}}(\tau)f) d\tau = \int_0^{+\infty} \frac{d}{d\tau} \|S_{\mathcal{L}}(\tau)f\|_{L^2}^2 d\tau = -\|f\|_{L^2}^2,$$

so after choosing proper η , we deduce that

$$((\mathcal{L}f, f))_k = (\mathcal{L}f, f)_{L_k^2} + \eta \int_0^{+\infty} (S_{\mathcal{L}}(\tau)\mathcal{L}f, S_{\mathcal{L}}(\tau)f)_{L_k^2} d\tau \sim \|f\|_{H_k^s}^2.$$

Therefore, it is important to note that the linear operator \mathcal{L} can exhibit non-negativity properties within an appropriate function space. The estimate of \mathcal{L} within this specific function space empowers us to establish global well-posedness by combining the nonlinear estimates.

The semigroup method serves as a valuable tool in addressing lower order terms. However, it introduces new challenging terms that require estimation, and this approach needs an assumption of $s \geq 1/2$. Precisely, when applying the semigroup method to analyze the Vlasov-Poisson term $\nabla_x \phi(x) \cdot \nabla_v f$, we encounter a departure from the usual case, where integration by parts yields the expected result

$$(\nabla_x \phi(x) \cdot \nabla_v f, f) = 0.$$

For the semigroup term, we have

$$\int_0^\infty (S_{\mathcal{L}}(\tau) \nabla_x \phi(x) \cdot \nabla_v f, S_{\mathcal{L}}(\tau) f) d\tau.$$

Since the operators $S_{\mathcal{L}}(t)$ and ∇_v are non-commutative, integration by parts with respect to ∇_v is not allowed and we must use upper bound to constrain this term. By [8] and Duhamel's formula, we have

$$\|S_B(t)f\|_{H_x^2 L_v^2} \leq t^{-1/2} e^{-\lambda t} \|f\|_{H_x^2 H_{12}^{-s}}, \quad \text{and} \quad \|S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2} \leq t^{-1/2} e^{-\lambda t} \|f\|_{H_x^2 H_{12}^{-s}}.$$

Due to the extra ∇_v term, we need to use

$$\|S_{\mathcal{L}}(t) \nabla_x \phi(x) \cdot \nabla_v f\|_{H_x^2 L_v^2} \lesssim t^{-1/2} e^{-\lambda t} \|\nabla_x \phi(x)\|_{H_x^2} \|\nabla_v f\|_{H_x^2 H_{12}^{-s}} \lesssim t^{-1/2} e^{-\lambda t} \|\nabla_x \phi(x)\|_{H_x^2} \|f\|_{H_x^2 H_{12}^s},$$

which requires $s \geq 1/2$.

1.6.1. *Comments on the weight (1.17).* The polynomial perturbation can be regarded as a change of variable $F = \mu + \langle v \rangle^{-k} f_{\pm}$ for (1.1) for some constant $k > 0$. Then we have

$$\partial_t f_{\pm} + v \cdot \nabla_x f_{\pm} \mp \nabla_x \phi \cdot \nabla_v f_{\pm} \pm \nabla_x \phi \cdot v \mu \langle v \rangle^{2k} \pm k \nabla_x \phi \cdot \frac{v}{\langle v \rangle^2} f = L_{k,\pm} f + \Gamma_{k,\pm}(f, f),$$

with

$$L_{k,\pm} f = \langle v \rangle^k Q(\langle v \rangle^{-k} (f_{\pm} + f_{\mp}), \mu) + 2 \langle v \rangle^k Q(\mu, \langle v \rangle^{-k} f_{\pm}), \quad \Gamma_{k,\pm}(f, f) = \langle v \rangle^k Q(\langle v \rangle^{-k} (f_{\pm} + f_{\mp}), \langle v \rangle^{-k} f_{\pm}).$$

In order to deal with the extra term $\pm k \nabla_x \phi \cdot \frac{v}{\langle v \rangle^2} f$, we follow the idea of [44] and introduce the corresponding weight function

$$\exp \left\{ \frac{\pm k \phi}{\langle v \rangle^2} \right\}, \tag{1.25}$$

which satisfies

$$v \cdot \nabla_x (e^{\frac{\pm k \phi}{\langle v \rangle^2}} f) = e^{\frac{\pm k \phi}{\langle v \rangle^2}} (v \cdot \nabla_x f \pm k \nabla_x \phi \cdot \frac{v}{\langle v \rangle^2} f).$$

Such choice of weight function allows us to absorb the extra term by taking integration by parts as

$$\left(\pm k \nabla_x \phi \cdot \frac{v}{\langle v \rangle^2} f, e^{\frac{\pm k \phi}{\langle v \rangle^2}} f \right)_{L_x^2 L_v^2} + \left(v \cdot \nabla_x f_{\pm}, e^{\frac{\pm k \phi}{\langle v \rangle^2}} f \right)_{L_x^2 L_v^2} = 0.$$

In contrast to the approach in [44], where the weight function was defined solely as $e^{-\phi}$, our weight function potential (1.25) exhibits dependence on the velocity variable as well. Fortunately, the newly introduced error contributions take the form of $(\nabla_x \phi \cdot v \mu, (e^{\frac{\pm k \phi}{\langle v \rangle^2}} - 1) \langle v \rangle^{2k} f)$. These contributions can be effectively managed when ϕ is sufficiently small, and the velocity derivative of ϕ remains under control, particularly if ϕ is bounded.

1.6.2. *Difference between the polynomial case and the exponential weight case.* The primary distinction lies in the symmetry properties of $L_{\mu} f$ and $\Gamma(f, f)$ as defined in (1.24). In the exponential case, these operators exhibit symmetry, whereas in the polynomial case, they do not. This lack of symmetry gives rise to two significant challenges. The first issue arises due to the introduction of an additional $\mathbf{P}f$ term in the linearized equation. To elaborate further, if we denote

$$\mathbf{P}_{\mu} f = a_{\pm}(t, x) \sqrt{\mu} + b(t, x) \cdot v \sqrt{\mu} + c(t, x) |v|^2 \sqrt{\mu}.$$

Then by $L_{\mu} \mathbf{P}_{\mu} f = 0$ and L_{μ} is symmetric, we have

$$(L_{\mu} f, f) = (L_{\mu} (\mathbf{P}_{\mu} f + \{\mathbf{I} - \mathbf{P}_{\mu}\} f), \mathbf{P}_{\mu} f + \{\mathbf{I} - \mathbf{P}_{\mu}\} f) = (L_{\mu} (\{\mathbf{I} - \mathbf{P}_{\mu}\} f), \{\mathbf{I} - \mathbf{P}_{\mu}\} f).$$

Only the term with $\{\mathbf{I} - \mathbf{P}_{\mu}\} f$ remains. While for the polynomial weight case, since L is not symmetric, one merely has

$$(L f, f) = (L \{\mathbf{I} - \mathbf{P}\} f, f) = (L \{\mathbf{I} - \mathbf{P}\} f, \mathbf{P} f) + (L \{\mathbf{I} - \mathbf{P}\} f, \{\mathbf{I} - \mathbf{P}\} f).$$

An extra term $(L\{\mathbf{I} - \mathbf{P}\}f, \mathbf{P}f)$ occurs. Fortunately, we can bound it by

$$(L\{\mathbf{I} - \mathbf{P}\}f, \mathbf{P}f) \leq \epsilon \|\mathbf{P}f\|_{L^2_{x,v}}^2 + \epsilon \|\{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_k}^2 + C_\epsilon \|\{\mathbf{I} - \mathbf{P}\}f\|_{L^2_{x,v}}.$$

Estimating the $\mathbf{P}f$ term can be accomplished through macroscopic estimates, while the last term can be assessed using the semigroup method.

However, the second challenge arises when we attempt to extend our results to $x \in \mathbb{R}^3$ using existing techniques. In previous works such as [62, 65], applied to the entire space $x \in \mathbb{R}^3$, the following fact is relied upon: $(\Gamma(g, f), \mathbf{P}_\mu h) = 0$ due to the symmetry of Γ . Consequently, in [62, 65], the dissipation rate functional $D(t)$ is defined as follows:

$$D(t) = \sum_{|\alpha| \leq K} \|\nabla_x \partial^\alpha \phi\|_{L^2_x}^2 + \sum_{1 \leq |\alpha| \leq K} \|\mathbf{P}_\mu \partial^\alpha f\|_{L^2_{x,v}}^2 + \sum_{0 \leq |\alpha| + |\beta| \leq K} \|w(\alpha, \beta) \{\mathbf{I} - \mathbf{P}_\mu\} \partial^\alpha f \partial^\beta f\|_{Y_k}^2,$$

In the exponential case, the estimation of $\mathbf{P}_\mu f$ conveniently avoids complications. However, in the polynomial case, a departure from this simplicity occurs. This is because, in the polynomial case, the relationship $(Q(f, g), \mu \phi(v)) \neq 0$ holds for functions $\phi(v)$ such as $\phi(v) = 1, v, |v|^2$. Consequently, the methods employed in [62, 65] are no longer applicable in the polynomial scenario. To address this, we necessitate the assumption that Ω represents a bounded domain and Poincaré inequality to estimate $\mathbf{P}f$ via $\nabla_x(\mathbf{P}f)$. However, extending our results to the entire space case remains an open challenge.

1.7. Organization of the paper. In Section 2, we first recall some basic properties of the non-cutoff Boltzmann/Landau collision operator and then compute the basic estimates for the VPB/VPL system. We obtain corresponding global results and decay of solutions in Section 3.

2. BASIC ESTIMATES

In this section, we recall some critical results on the non-cutoff Boltzmann equation and Landau equation and prove several weighted estimates for the VPB/VPL equation.

2.1. Preliminaries on the Boltzmann/Landau equation. Before introducing the upper and lower bound for the non-cutoff Boltzmann and Landau collision operator, we first state the following lemmas.

Lemma 2.1. ([7], Lemma 2.5) *Let $\gamma \in (-2, 0)$. Then for any function f , we have*

$$\sup_{v \in \mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^\gamma |f(v_*)| dv_* \lesssim \|f\|_{L^1}^{1+\frac{2\gamma}{3}} \|f\|_{L^2}^{-\frac{2\gamma}{3}} \quad \text{when } \gamma \in (-\frac{3}{2}, 0), \quad (2.1)$$

and

$$\sup_{v \in \mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^\gamma |f(v_*)| dv_* \lesssim \|f\|_{L^1}^{1+\frac{\gamma}{2}} \|f\|_{L^3}^{-\frac{\gamma}{2}} \quad \text{when } \gamma \in (-2, 0). \quad (2.2)$$

Lemma 2.2. *For any $\gamma \in [-3, 1]$ and any functions f, g, h , we have*

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma+2} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^2_5} \|g\|_{L^2} \|h\|_{L^2_3}, \quad (2.3)$$

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma+1} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^2_5} \|g\|_{L^2} \|h\|_{H^1_3}, \quad (2.4)$$

and

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma+1} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^2_5} \|g\|_{H^1_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}}. \quad (2.5)$$

If $\gamma \in (-3, 1]$, then we have

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^\gamma f(v_*) g^2(v) dv_* dv \lesssim \|f\|_{L^2_5} \|g\|_{H^1_{\gamma/2}}^2. \quad (2.6)$$

Proof. If $-2 \leq \gamma \leq 1$, we deduce that

$$\int_{\mathbb{R}^3} |v - v_*|^{\gamma+2} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^1_3} \|g\|_{L^2} \|h\|_{L^2_3} \lesssim \|f\|_{L^2_5} \|g\|_{L^2} \|h\|_{L^2_3}.$$

If $-3 \leq \gamma < -2$, by (2.1), we have $\sup_{v \in \mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma+2} f(v_*) dv_* \lesssim \|f\|_{L^2_5}$, which implies that

$$\int_{\mathbb{R}^3} |v - v_*|^{\gamma+2} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^2_5} \int_{\mathbb{R}^3} |g(v) h(v)| dv \lesssim \|f\|_{L^2_5} \|g\|_{L^2} \|h\|_{L^2}.$$

This proves (2.3). For (2.4), if $\gamma \geq -2$, then it reduce to the (2.3). So we only need to consider the case $\gamma \in [-3, -2)$, in which case we have $-3 < \gamma + \frac{1}{2} < 0$ and $-2 < \gamma + \frac{3}{2} < 0$. Then we have

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + \frac{1}{2}} \frac{1}{\langle v_* \rangle^4} |g(v)|^2 dv dv_* \lesssim \|g\|_{L^2}^2.$$

Applying (2.2) to h , we have

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + \frac{3}{2}} \langle v_* \rangle^4 |f(v_*)|^2 |h(v)|^2 dv dv_* \lesssim \|f\|_{L^2}^2 \|h\|_{L^1}^{1 - \frac{\gamma + 3/2}{2}} \|h\|_{L^3}^{\frac{\gamma + 3/2}{2}} \lesssim \|f\|_{L^2}^2 \|h\|_{H^1}^2.$$

Here we apply Sobolev embedding $\|\cdot\|_{L^6} \lesssim \|\cdot\|_{H^1}$. Thus,

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + 1} |f(v_*)| |g(v)| |h(v)| dv dv_* \\ & \leq \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + \frac{3}{2}} \langle v_* \rangle^4 |f(v_*)|^2 |h(v)|^2 dv dv_* \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|v - v_*|^{\gamma + \frac{1}{2}}}{\langle v_* \rangle^4} |g(v)|^2 dv dv_* \right)^{\frac{1}{2}} \\ & \lesssim \|f\|_{L^2} \|g\|_{L^2} \|h\|_{H^1}. \end{aligned}$$

The proof of (2.4) is then finished. Now we come to (2.5). If $-1 \leq \gamma \leq 1$, we deduce that

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + 1} f(v_*) g(v) h(v) dv_* dv \lesssim \|f\|_{L^1_{\gamma+1}} \int_{\mathbb{R}^3} \langle v \rangle^{\gamma+1} |g(v) h(v)| dv \lesssim \|f\|_{L^2_4} \|g\|_{L^2_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}}.$$

If $-3 \leq \gamma < -1$, we have $\langle v \rangle^{-(\gamma+1)} \lesssim \langle v_* \rangle^{-(\gamma+1)} \langle v - v_* \rangle^{-(\gamma+1)}$, which implies

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^{\gamma + 1} f(v_*) g(v) h(v) dv_* dv \\ & \lesssim \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \langle v - v_* \rangle^{-(\gamma+1)} |v - v_*|^{\gamma + 1} f(v_*) \langle v_* \rangle^{-(\gamma+1)} |g(v) h(v)| \langle v \rangle^{\gamma+1} dv_* dv \\ & \lesssim \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} (1 + |v - v_*|^{\gamma+1}) f(v_*) \langle v_* \rangle^{-(\gamma+1)} |g(v) h(v)| \langle v \rangle^{\gamma+1} dv_* dv := T_1 + T_2. \end{aligned}$$

It is easily seen that $T_1 \lesssim \|f\|_{L^1_{-(\gamma+1)}} \|g\|_{L^2_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}} \lesssim \|f\|_{L^2_5} \|g\|_{L^2_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}}$. If $\gamma \in [-2, -1)$, we have $\gamma + 1 \in (-\frac{3}{2}, 0)$ and by (2.1),

$$T_2 \lesssim \sup_{v \in \mathbb{R}^d} \int_{\mathbb{R}^3} |v - v_*|^{\gamma+1} |f(v_*)| \langle v_* \rangle^{-(\gamma+1)} dv_* \int_{\mathbb{R}^3} |g(v) h(v)| \langle v \rangle^{\gamma+1} dv \lesssim \|f\|_{L^2_5} \|g\|_{H^1_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}}.$$

If $\gamma \in [-3, -2)$, by Hardy-Littlewood-Sobolev (HLS) inequality (cf. [61, Theorem 1.1, pp. 119]) and Hölder's inequality, we have

$$\begin{aligned} T_2 & \lesssim \left\| \int_{\mathbb{R}^3} |v - v_*|^{\gamma+1} f(v_*) \langle v_* \rangle^{-(\gamma+1)} dv_* \right\|_{L^6_v} \|hg \langle v \rangle^{\gamma+1}\|_{L^{\frac{6}{5}}} \\ & \lesssim \|f\|_{L^p_{-(\gamma+1)}} \|g\|_{L^3_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}} \lesssim \|f\|_{L^2_5} \|g\|_{H^1_{\gamma/2+1}} \|h\|_{L^2_{\gamma/2}}, \end{aligned}$$

with $p = \frac{6}{9+2\gamma} \in [1, 2]$, and (2.5) is proved. Finally, we come to prove (2.6). For $\gamma \in [-2, 1]$, (2.6) can be seen as a special case of (2.5). We focus on the case $\gamma \in (-3, -2)$. Similarly to (2.1), we have

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - v_*|^\gamma |f(v_*)| |g(v)|^2 dv_* dv \lesssim \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} (1 + |v - v_*|^\gamma) |f(v_*)| \langle v_* \rangle^{-\gamma} |g(v)|^2 \langle v \rangle^\gamma dv_* dv := I_1 + I_2.$$

It is seen that $I_1 \lesssim \|f\|_{L^1_{-\gamma}} \|g\|_{L^2_{\gamma/2}}^2 \lesssim \|f\|_{L^2_5} \|g\|_{L^2_{\gamma/2}}^2$. For I_2 , by HLS inequality as above, we have

$$I_2 \lesssim \|f\|_{L^2_{-\gamma}} \left\| \int_{\mathbb{R}^3} |v - v_*|^\gamma |g(v)|^2 \langle v \rangle^\gamma dv \right\|_{L^2_{v_*}} \lesssim \|f\|_{L^2_{-\gamma}} \| |g|^2 \langle v \rangle^\gamma \|_{L^p} \lesssim \|f\|_{L^2_5} \|g\|_{L^2_{\gamma/2}}^{2p} \lesssim \|f\|_{L^2_5} \|g\|_{H^1_{\gamma/2}}^2,$$

where $p = \frac{6}{9+2\gamma}$ satisfies $2p \in [2, 6]$. This gives (2.6) and the proof of Theorem 2.2 is finished. \square

We also recall some basic interpolation on x .

Lemma 2.3. *For any function f and constant $k \in \mathbb{R}$, we have $\|fg\|_{H^2_x} \lesssim \min_{c+d=2} \{\|f\|_{H^c_x} \|g\|_{H^d_x}\}$. More precisely,*

$$\|f\|_{L^\infty_x L^2_v} \lesssim \|f \langle v \rangle^k\|_{H^1_x L^2_v} + \|f \langle v \rangle^{-k}\|_{H^2_x L^2_v}, \quad (2.7)$$

$$\|f\|_{L^3_x L^2_v} \lesssim \|f \langle v \rangle^k\|_{L^2_x L^2_v} + \|f \langle v \rangle^{-k}\|_{H^1_x L^2_v}, \quad (2.8)$$

and for any constant $s \in (0, 1)$

$$\|f\|_{H_v^1} \lesssim \|f\langle v \rangle^k\|_{H_v^s} + \|f\langle v \rangle^{-ks/(1-s)}\|_{H_v^{1+s}}. \quad (2.9)$$

Proof. Firstly, we use extension theorem [61, Theorem VI.5, pp. 181] to extend function f in domain Ω with Lipschitz boundary to a function Ef in \mathbb{R}^3 such that $Ef = f$ in Ω and $\|Ef\|_{H^k(\mathbb{R}^3)} \lesssim \|f\|_{H^k(\Omega)}$ for any $k \geq 0$. By Gagliardo-Nirenberg interpolation inequality on \mathbb{R} and \mathbb{R}^2 (cf. [55, Theorem 12.83] and [59, Page 125]), we obtain

$$\|Ef\|_{L_{x_1}^\infty(\mathbb{R})} \lesssim \|\partial_{x_1} Ef\|_{L_{x_1}^2(\mathbb{R})}^{1/2} \|Ef\|_{L_{x_1}^2(\mathbb{R})}^{1/2}, \quad \text{and} \quad \|Ef\|_{L_{x_2, x_3}^\infty(\mathbb{R}^2)} \lesssim \|\nabla_{x_2, x_3} Ef\|_{L_{x_2, x_3}^2(\mathbb{R}^2)}.$$

Combining the above estimates, we have $\|f\|_{L_x^\infty(\Omega)} \lesssim \|Ef\|_{L_{x_1}^\infty(\mathbb{R})} \|Ef\|_{L_{x_2, x_3}^\infty(\mathbb{R}^2)} \lesssim \|f\|_{H_x^2(\Omega)}^{1/2} \|f\|_{H_x^1(\Omega)}^{1/2}$, which implies (2.7). Also, by Gagliardo-Nirenberg interpolation inequality on \mathbb{R}^3 , we have

$$\|Ef\|_{L_x^3(\mathbb{R}^3)} \lesssim \|Ef\|_{L_x^2(\mathbb{R}^3)}^{1/2} \|\nabla_x Ef\|_{L_x^2(\mathbb{R}^3)}^{1/2} \lesssim \|f\|_{L_x^2(\Omega)}^{1/2} \|f\|_{H_x^1(\Omega)}^{1/2},$$

which gives (2.8). For (2.9), by Young's inequality, $\langle \eta \rangle \lesssim \langle \eta \rangle^s \langle v \rangle^k + \langle \eta \rangle^{1+s} \langle v \rangle^{-\frac{ks}{1-s}}$. Hence $\langle \eta \rangle$ is a symbol in $S(\langle \eta \rangle^s \langle v \rangle^k + \langle \eta \rangle^{1+s} \langle v \rangle^{-\frac{ks}{1-s}})$ (cf. [16]), where η is the Fourier variable of v . Then by [16, Lemma 2.3 and Corollary 2.5], we have $\|f\|_{H_v^1} \lesssim \|f\langle v \rangle^k\|_{H_v^s} + \|f\langle v \rangle^{-ks/(1-s)}\|_{H_v^{1+s}}$. This completes the Lemma. \square

2.1.1. *Upper and lower bound for the non-cutoff Boltzmann operator.* In this subsection, we consider the Boltzmann collision operator $Q(f, g)$.

Lemma 2.4 ([47], Theorem 1.1). *Suppose $\gamma \in (-3, 1], s \in (0, 1), \gamma + 2s > -1$. Let $w_1, w_2 \in \mathbb{R}, a, b \in [0, 2s]$ with $w_1 + w_2 = \gamma + 2s$ and $a + b = 2s$. Then for any functions f, g, h we have*

- (1) if $\gamma + 2s > 0$, then $|(Q(g, h), f)_{L_v^2}| \lesssim (\|g\|_{L_{\gamma+2s+(-w_1)^+ + (-w_2)^+}}^1 + \|g\|_{L^2}) \|h\|_{H_{w_1}^a} \|f\|_{H_{w_2}^b}$,
- (2) if $\gamma + 2s = 0$, then $|(Q(g, h), f)_{L_v^2}| \lesssim (\|g\|_{L_{w_3}^1} + \|g\|_{L^2}) \|h\|_{H_{w_1}^a} \|f\|_{H_{w_2}^b}$, where $w_3 = \max\{\delta, (-w_1)^+ + (-w_2)^+\}$, with $\delta > 0$ sufficiently small.
- (3) if $-1 < \gamma + 2s < 0$, then $|(Q(g, h), f)_{L_v^2}| \lesssim (\|g\|_{L_{w_4}^1} + \|g\|_{L_{-(\gamma+2s)}^2}) \|h\|_{H_{w_1}^a} \|f\|_{H_{w_2}^b}$, where $w_4 = \max\{-\gamma + 2s, \gamma + 2s + (-w_1)^+ + (-w_2)^+\}$.

As a first application, we have the following corollaries.

Corollary 2.5. *Suppose $\gamma \in (-3, 1], s \in (0, 1), \gamma + 2s > -1$. For any multi-indices $|\beta| \leq 2$, any constant $k \geq 0$ and any functions f, g , there exists some constant $C_k > 0$, such that*

$$(Q(\partial_\beta \mu, f), g\langle v \rangle^{2k}) \leq C_k \|f\|_{H_{k+\gamma/2+2s}^s} \|g\|_{H_{k+\gamma/2}^s}.$$

Lemma 2.6 ([8], Lemma 3.3). *Suppose that $-3 < \gamma \leq 1$. For any $k \geq 14$, and functions g, h , we have*

$$\begin{aligned} |(Q(h, \mu), g\langle v \rangle^{2k})| &\leq \|b(\cos \theta) \sin^{k-\frac{3+\gamma}{2}}\|_{L_\theta^1} \|h\|_{L_{k+\gamma/2, *}^2} \|g\|_{L_{k+\gamma/2, *}^2} + C_k \|h\|_{L_{k+\gamma/2-1/2}^2} \|g\|_{L_{k+\gamma/2-1/2}^2} \\ &\leq \|b(\cos \theta) \sin^{k-2}\|_{L_\theta^1} \|h\|_{L_{k+\gamma/2, *}^2} \|g\|_{L_{k+\gamma/2, *}^2} + C_k \|h\|_{L_{k+\gamma/2-1/2}^2} \|g\|_{L_{k+\gamma/2-1/2}^2}, \end{aligned} \quad (2.10)$$

for some constant $C_k > 0$. Moreover, for any $|\beta| \leq 2$ we have

$$|(Q(h, \partial_\beta \mu), g\langle v \rangle^{2k})| \leq C_k \|h\|_{L_{k+\gamma/2}^2} \|g\|_{L_{k+\gamma/2}^2}. \quad (2.11)$$

Remark 2.7. *In [8, Lemma 3.3], the authors only prove the first statement of Lemma 2.6, but the second statement can be proved in the same way by just replacing μ by $\partial_\beta \mu$.*

Theorem 2.8 ([8], Theorem 3.1). *Suppose that $-3 < \gamma \leq 1, s \in (0, 1), \gamma + 2s > -1, k \geq 14$ and $G = \mu + g \geq 0$. If there exists $A_1, A_2 > 0$ such that*

$$G \geq 0, \quad \|G\|_{L^1} \geq A_1, \quad \|G\|_{L^{\frac{1}{2}}} + \|G\|_{L \log L} \leq A_2,$$

then there exist some constants $\gamma_1, C_k > 0$, such that

$$\begin{aligned} (Q(G, f), f\langle v \rangle^{2k}) &\leq -\frac{1}{8} \|b(\cos \theta) \sin^2\|_{L_\theta^1} \|f\|_{L_{k+\gamma/2, *}^2}^2 - \gamma_1 \|f\|_{H_{k+\gamma/2}^s}^2 + C_k \|f\|_{L^2}^2 \\ &\quad + C_k \|f\|_{L_{14}^2} \|g\|_{H_{k+\gamma/2}^s} \|f\|_{H_{k+\gamma/2}^s} + C_k \|g\|_{L_{14}^2} \|f\|_{H_{k+\gamma/2}^s}^2. \end{aligned} \quad (2.12)$$

Remark 2.9. In [8, Theorem 3.1], the authors explored both exponential weight perturbation and polynomial weight perturbation cases. In the exponential weight scenario, the estimation involves an infinite sum, necessitating $k \geq 22$ for convergence. However, for the polynomial case, it has been demonstrated that $k \geq 14$ is sufficient, employing the same technique. Note that whenever $\|g\|_{L_v^1}$ is small, we have $\|\mu + g\|_{L_v^1} \geq 1 - \|g\|_{L_v^1} > A_1$. Moreover, we couldn't obtain a better estimate for the term $(Q(f, g), g\langle v \rangle^{2k})$ as the Landau case (see (2.19)) and thus we should put the term $\mu + g \geq 0$ together.

Lemma 2.10 ([8], Lemma 2.4 + Lemma 2.4). Suppose $\gamma \in (-3, 1]$, $s \in (0, 1)$, $\gamma + 2s > -1$. For any functions f, g, h and $k \geq 14$, we have

$$(Q(f, g), h\langle v \rangle^{2k}) \lesssim \|f\|_{L_{14}^2} \min\{\|g\|_{H_{k+\gamma/2}^s} \|h\|_{H_{k+\gamma/2+2s}^s}, \|g\|_{H_{k+\gamma/2+2s}^s} \|h\|_{H_{k+\gamma/2}^s}\} + \|g\|_{L_{14}^2} \|f\|_{H_{k+\gamma/2}^s} \|h\|_{H_{k+\gamma/2}^s}. \quad (2.13)$$

In particular, by duality we have

$$\|Q(f, g)\|_{H_{k-\gamma/2}^{-s}} \lesssim \|f\|_{L_{14}^2} \|g\|_{H_{k+\gamma/2+2s}^s} + \|g\|_{L_{14}^2} \|f\|_{H_{k+\gamma/2}^s}. \quad (2.14)$$

2.1.2. *Upper and lower bound for Landau operator.* Next, we give some results about the Landau collision operator $Q(f, g)$.

Lemma 2.11. Denote $m = \langle v \rangle^k$ with $k \geq 7$. For any $-3 \leq \gamma \leq 1$, $|\beta| \leq 2$ and functions f, g , we have

$$|(Q(f, \partial_\beta \mu), g\langle v \rangle^{2k})| \leq C_k \|f\|_{L_5^2} \|g\|_{L_5^2}, \quad (2.15)$$

and

$$(Q(\mu, f), f\langle v \rangle^{2k}) \leq -\gamma_1 \|f\|_{L_{D(m)}^2}^2 + C_k \|f\|_{L^2}^2, \quad (2.16)$$

for some constants $\gamma_1, C_k > 0$.

Proof. The first estimate follows from [11, Lemma 2.5] and in [10, Lemma 2.12] with μ replacing by $\partial_\beta \mu$ if necessary. The second estimate follows from [11, Lemma 2.3] and [10, Lemma 2.7]. Note that the work [10] is for the case $\gamma \in [-2, 1]$ and [11] is for $\gamma \in [-3, -2]$. \square

Lemma 2.12 ([10], Lemma 3.5 and [11], Lemma 4.3). For any $-3 \leq \gamma \leq 1$, $k \geq 7$ and any functions f, g, h , we have

$$|(Q(f, g), h\langle v \rangle^{2k})| \leq C \|f\|_{L_7^2} \min\{\|g\|_{L_{D,k+1}^2} \|h\|_{L_{D,k}^2}, \|g\|_{L_{D,k}^2} \|h\|_{L_{D,k+1}^2}\}. \quad (2.17)$$

By duality, we have

$$\|Q(f, g)\|_{H_{k-\gamma/2}^{-1}} \lesssim \|f\|_{L_7^2} \|g\|_{L_{D,k+1}^2}. \quad (2.18)$$

When $h = g$, we have a better estimate:

$$(Q(f, g), g\langle v \rangle^{2k}) \leq C \|f\|_{L_7^2} \|g\|_{L_{D,k}^2}^2. \quad (2.19)$$

Proof. The proof is similar to [10, Lemma 3.5] and [11, Lemma 4.3] with a little modification. Denoting $m = \langle v \rangle^k$, we have

$$\begin{aligned} (Q(f, g), hm^2) &= \int_{\mathbb{R}^3} \partial_j \{(\phi^{ij} * f) \partial_i g - (\partial_i \phi^{ij} * f) g\} hm^2 dv \\ &= - \int_{\mathbb{R}^3} (\phi^{ij} * f) \partial_i g \partial_j hm^2 dv - \int_{\mathbb{R}^3} (\phi^{ij} * f) \partial_i gh \partial_j m^2 dv \\ &\quad + \int_{\mathbb{R}^3} (\partial_i \phi^{ij} * f) g \partial_j hm^2 dv + \int_{\mathbb{R}^3} (\partial_i \phi^{ij} * f) gh \partial_j m^2 dv := T_1 + T_2 + T_3 + T_4. \end{aligned}$$

For the T_1, T_2 term, by [10, Lemma 3.5] and [11, Lemma 4.3], we have $|T_1| + |T_2| \leq C \|f\|_{L_7^2} \|g\|_{L_{D,k}^2} \|h\|_{L_{D,k}^2}$. Now we give a better estimate for the T_3, T_4 term. Since $\partial_i \phi^{ij}(v - v_*) \lesssim |v - v_*|^{\gamma+1}$, $\partial_j m^2 \leq C \langle v \rangle^{-1} m^2$, for the term T_3, T_4 , by (2.5), we have

$$|T_3| \leq C \|f\|_{L_7^2} \min\{\|g\|_{L_{D,k+1}^2} \|h\|_{L_{D,k}^2}, \|g\|_{L_{D,k}^2} \|h\|_{L_{D,k+1}^2}\}, \quad |T_4| \leq C \|f\|_{L_7^2} \|g\|_{L_{D,k}^2} \|h\|_{L_{D,k}^2},$$

and (2.17) is thus proved. For the case $g = h$, we give a better estimate for T_3 . In fact,

$$T_3 = \int_{\mathbb{R}^3} (\partial_i \phi^{ij} * f) g \partial_j gm^2 dv = -\frac{1}{2} \int_{\mathbb{R}^3} (\partial_{ij} \phi^{ij} * f) g^2 m^2 dv - \frac{1}{2} \int_{\mathbb{R}^3} (\partial_i \phi^{ij} * f) \partial_j m^2 g^2 dv := T_{31} + T_{32}.$$

For the T_{32} term, similar to the T_4 term, by (2.5), we have $T_{32} \leq C\|f\|_{L^2_\gamma}\|g\|_{L^2_{D,k}}^2$. For the T_{31} term, if $\gamma = -3$, then $\partial_{ij}\phi^{ij} = -8\pi\delta_0$. It is easily seen that

$$T_{31} \leq C \int_{\mathbb{R}^3} |f|g^2m^2dv \leq C\|f\|_{L^2_{-\gamma}}\|g^2\|_{L^2_{2k+\gamma}} \leq C\|f\|_{L^2_\gamma}\|g\|_{L^4_{k+\gamma/2}}^2 \leq C\|f\|_{L^2_\gamma}\|g\|_{L^2_{D,k}}^2.$$

For the case $\gamma \in [0, 1]$, we have

$$T_{31} \leq C \int_{\mathbb{R}^3} |v - v_*|^\gamma |f_*|g^2m^2dv dv_* \leq C\|f\|_{L^2_\gamma}\|g\|_{L^2_{k+\gamma/2}}^2 \leq C\|f\|_{L^2_\gamma}\|g\|_{L^2_{D,k}}^2.$$

The case $\gamma \in (-3, 1]$ follows from (2.6). Collecting the above estimates on T_{31} and T_{32} , we obtain (2.19) and complete the proof of Lemma 2.12. \square

For the Landau operator, we need another upper bound which writes

Lemma 2.13. *For the Landau operator Q , for any $\gamma \in [-3, 1]$ and any function f, g, h , we have*

$$|(Q(f, g), h)| \leq \|f\|_{L^2_\gamma}\|g\|_{L^2}\|h\|_{H^2_\gamma}, \quad |(Q(f, g), h)| \leq \|f\|_{L^2_\gamma}\|g\|_{L^2_{10}}\|h\|_{H^2_{-5}},$$

Proof. By using integration by parts, we easily have

$$(Q(f, g), h) = \int_{\mathbb{R}^3} \left(-(\phi^{ij} * f)\partial_j g \partial_i h + (\partial_j \phi^{ij} * f)g \partial_i h \right) dv = \int_{\mathbb{R}^3} \left((\phi^{ij} * f)g \partial_i \partial_j h + 2(\partial_j \phi^{ij} * f)g \partial_i h \right) dv,$$

then by the homogeneity of ϕ^{ij} and $\partial_j \phi^{ij}$, we have $|\phi^{ij}(v-v_*)| \lesssim |v-v_*|^{\gamma+2}$, $|\partial_j \phi^{ij}(v-v_*)| \lesssim |v-v_*|^{\gamma+1}$. So the theorem follows directly from (2.3) and (2.4). \square

2.2. High-order specular boundary conditions. When Ω is the union of cubes given by (1.10), we give the high-order compatible specular boundary condition from [19, Lemma 3.1 and 3.2]. Although the proof in [19] is given for exponential decay perturbations $F = \mu + \mu^{1/2}f$, similar calculations can be applied to polynomial perturbations $F = \mu + f$.

Lemma 2.14 ([19, Lemma 3.1]). *Let (f, ϕ) be the solution to (1.3) with boundary conditions (1.12) and (1.13). Fix $i \in \{1, 2, 3\}$, $x \in \Gamma_i$ and $v \cdot n(x) \neq 0$. Then we have the following identities on boundary Γ_i :*

$$\partial^\alpha f(x, v) = (-1)^{\alpha_i} \partial^\alpha f(x, R_x v), \quad (2.20)$$

for any $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3$.

As a corollary, by definition (1.6), we have the following boundary values for $[a_\pm, b, c]$.

Lemma 2.15 ([19, Lemma 3.2]). *Let (f, ϕ) be the solution to (1.3) with boundary conditions (1.12) and (1.13). Define $[a_\pm, b, c]$ by (1.6). For $i = 1, 2, 3$, $j \neq i$ and any $x \in \Gamma_i$, we have*

$$\partial_{x_i} c(x) = \partial_{x_i} a_\pm(x) = \partial_{x_i} b_j(x) = \partial_{x_i x_i} b_i(x) = b_i(x) = 0. \quad (2.21)$$

2.3. Estimates on the weight. Recall the weight function $w = w(|\alpha|, |\beta|) = w(\alpha, \beta)$ is defined in (1.15). We have the following properties of w .

Lemma 2.16. *Assume $-3 < \gamma \leq 1$, $1/2 \leq s < 1$, $\gamma + 2s > -1$ for Boltzmann case and $-3 \leq \gamma \leq 1$ for Landau case. For any multi-indices α, β and $k \in \mathbb{N}$, $w(\alpha, \beta)$ satisfies the following properties.*

• For $|\alpha_1| < |\alpha|$, $|\beta_1| < |\beta|$, we have

$$w(\alpha, \beta) \langle v \rangle^{6s} \leq w(\alpha, \beta_1), \quad w(\alpha, \beta) \langle v \rangle^{6s} \leq w(\alpha_1, \beta) \quad \text{for the Boltzmann case,} \quad (2.22)$$

$$w(\alpha, \beta) \langle v \rangle^3 \leq w(\alpha, \beta_1), \quad w(\alpha, \beta) \langle v \rangle^3 \leq w(\alpha_1, \beta) \quad \text{for the Landau case.} \quad (2.23)$$

• For both Boltzmann and Landau case, and any $|\alpha| \geq 0$, $|\beta| \geq 1$, we have

$$w(\alpha, \beta) \leq \langle v \rangle^{\gamma-1} w(|\alpha| + 1, |\beta| - 1). \quad (2.24)$$

• For the Boltzmann case and any $|\alpha| \geq 1$, $|\beta| \geq 0$, we have

$$w(\alpha, \beta) \leq w(|\alpha| - 1, |\beta|)^s w(|\alpha| - 1, |\beta| + 1)^{1-s} \langle v \rangle^\gamma. \quad (2.25)$$

Proof. Recall that $q = 6s - 3(\gamma - 1)$, $p = q + \gamma - 1$, $r = 2q + 6$, for Boltzmann case, and $q = 3 - (\gamma - 1)$, $p = 3$, $r = 2q + 6$ for the Landau case. (2.22) and (2.23) are just from the fact that $-3 \leq \gamma \leq 1$ and

$$p = -3(\gamma - 1) + 6s + \gamma - 1 \geq 6s, \quad q \geq 6s,$$

for the Boltzmann case and $p, q \geq 3$ for the Landau case. The estimate (2.24) follows from the fact that

$$k - p|\alpha| - q|\beta| + r \leq k - p(|\alpha| + 1) - q(|\beta| - 1) + r + \gamma - 1 \iff q - p + \gamma - 1 \geq 0,$$

and we conclude (2.24) from the definition of p, q . (2.25) is equivalent to

$$k - p|\alpha| - q|\beta| + r \leq (k - p(|\alpha| - 1) - q|\beta| + r)s + (k - p(|\alpha| - 1) - q(|\beta| + 1) + r)(1 - s) + \gamma,$$

which is equivalent to

$$0 \leq ps + p(1 - s) - q(1 - s) + \gamma \iff 0 \leq qs + (p - q) + \gamma = (6s - 3(\gamma - 1))s + \gamma - 1 + \gamma.$$

We thus conclude (2.25) from the definition of q , $1/2 \leq s < 1$ and $-3 < \gamma \leq 1$. \square

We can directly deduce the following Corollary from Lemma 2.16.

Corollary 2.17. *Assume the same conditions as in Lemma 2.16. Then we have*

$$\max_{|\alpha|+|\beta|=1} w^2(\alpha, \beta) \langle v \rangle^\kappa \leq w(0, 0)w(1, 0), \quad (2.26)$$

$$\max_{|\alpha|+|\beta|=2} w^2(\alpha, \beta) \langle v \rangle^\kappa \leq w(1, 0)w(2, 0), \quad \max_{\substack{|\alpha|+|\beta|=2 \\ |\beta| \geq 1}} w^2(\alpha, \beta) \langle v \rangle^\kappa \leq w(0, 1)w(1, 1), \quad (2.27)$$

$$\max_{|\alpha|+|\beta|=2} w^2(\alpha, \beta) \langle v \rangle^\kappa \leq w^2(2, 0) \langle v \rangle^\kappa \leq w(1, 0)^{4/5} w(2, 0)^{6/5}, \quad (2.28)$$

where $\kappa = 4s$ for Boltzmann case, $\kappa = 2$ for Landau case.

2.4. Weighted estimates. Next, we come to the nonlinear term for the Vlasov-Poisson term. Noticing $w^2(\alpha, \beta)$ is still of the form $\langle v \rangle^k$ for some $k > 0$, we have

$$\nabla_v(w^2(\alpha, \beta)) = A_{\alpha, \beta} \frac{v}{\langle v \rangle^2} w^2(\alpha, \beta), \quad (2.29)$$

for some constant $A_{\alpha, \beta}$ depending on α, β, k, γ . Then we can apply the space-velocity weight $e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}}$. To the end of this section, we will assume

$$\|\phi\|_{L_x^\infty} \leq C < +\infty, \quad \|\psi\|_{L_x^\infty} \leq C < +\infty. \quad (2.30)$$

and hence,

$$\left| e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \right| \leq C < \infty, \quad \left| e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \right| \leq C < \infty. \quad (2.31)$$

Lemma 2.18. *Suppose that $-3 \leq \gamma \leq 1$ for Landau case and $-3 < \gamma \leq 1$ for Boltzmann case. For $|\alpha| + |\beta| \leq 2$ and any functions f_\pm , for any function ϕ satisfies (2.30), we have*

$$\left| (v \cdot \nabla_x \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} \mp (\nabla_x \phi \cdot \nabla_v \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} \right| \lesssim \|\nabla_x \phi\|_{H_x^2} \|f\|_{Y_k}^2, \quad (2.32)$$

and for $|\beta| > 0$,

$$\sum_{|\beta_1|=1} |(\partial_{\beta_1} v \cdot \nabla_x \partial_{\beta-\beta_1}^\alpha f, e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}| \leq C \sum_{\substack{|\beta_2|=|\beta|-1 \\ |\alpha_2|=|\alpha|+1}} \|\partial_{\beta_2}^{\alpha_2} f w(\alpha_2, \beta_2)\|_{L_x^2 L_{\gamma/2}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}^2}. \quad (2.33)$$

For $|\alpha_1| \geq 1$, we have

$$|(\nabla_x \partial^{\alpha_1} \phi \cdot \nabla_v \partial_\beta^{\alpha-\alpha_1} f, e^{\frac{\pm A_{\alpha, \beta} \phi}{\langle v \rangle^2}} \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}| \lesssim \|\nabla_x \phi\|_{H_x^3} \|f\|_{Y_k}^2, \quad (2.34)$$

where $A_{\alpha, \beta}$ is given by (2.29), for some constant $C_k > 0$ for both Landau and Boltzmann case.

Proof. For (2.32), taking integration by parts with respect to ∇_x , we have

$$\begin{aligned} & (v \cdot \nabla_x \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} \pm \frac{A_{\alpha,\beta}}{2} \left(\frac{v \cdot \nabla_x \phi}{\langle v \rangle^2} \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta) \right)_{L_{x,v}^2} \\ &= \int_{\partial\Omega} \int_{\mathbb{R}^3} v \cdot n(x) |e^{\frac{\pm A_{\alpha,\beta}\phi}{2\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w(\alpha, \beta)|^2 dv dS(x). \end{aligned}$$

For the torus case, the right hand side is equal to zero. For the case of union of cubes, we apply Lemma 2.14 to obtain $|\partial_\beta^\alpha f_\pm(v)| = |\partial_\beta^\alpha f_\pm(R_x v)|$ for $x \in \partial\Omega$ and R_x is given by (1.11). Thus, by change of variable $v \mapsto R_x v$,

$$\begin{aligned} & \int_{\partial\Omega} \int_{\mathbb{R}^3} v \cdot n(x) |e^{\frac{\pm A_{\alpha,\beta}\phi}{2\langle v \rangle^2}} \partial_\beta^\alpha f_\pm(v) w(\alpha, \beta)|^2 dv dS(x) \\ &= \int_{\partial\Omega} \int_{\mathbb{R}^3} R_x v \cdot n(x) |e^{\frac{\pm A_{\alpha,\beta}\phi}{2\langle R_x v \rangle^2}} \partial_\beta^\alpha f_\pm(R_x v) w(\alpha, \beta)|^2 dv dS(x) \\ &= - \int_{\partial\Omega} \int_{\mathbb{R}^3} v \cdot n(x) |e^{\frac{\pm A_{\alpha,\beta}\phi}{2\langle v \rangle^2}} \partial_\beta^\alpha f_\pm(v) w(\alpha, \beta)|^2 dv dS(x) = 0, \end{aligned} \quad (2.35)$$

where $R_x v \cdot n(x) = -v \cdot n(x)$. Taking integration by parts with respect to ∇_v and using (2.29), we have

$$\begin{aligned} \mp (\nabla_x \phi \cdot \nabla_v \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} &= \mp \left(\frac{A_{\alpha,\beta} v \cdot \nabla_x \phi \phi}{\langle v \rangle^4} \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta) \right)_{L_{x,v}^2} \\ &\quad \pm \frac{1}{2} \left(\frac{A_{\alpha,\beta} v \cdot \nabla_x \phi}{\langle v \rangle^2} \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta) \right)_{L_{x,v}^2}. \end{aligned}$$

Noticing $|\frac{v}{\langle v \rangle^4}| \leq \langle v \rangle^\gamma$ for $-3 \leq \gamma \leq 1$ and taking summation of the above two estimates, we have

$$\begin{aligned} & \left| (v \cdot \nabla_x \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} \mp (\nabla_x \phi \cdot \nabla_v \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2} \right| \\ & \lesssim \|\nabla_x \phi\|_{L_x^\infty} \|\langle v \rangle^{\frac{\gamma}{2}} \partial_\beta^\alpha f_\pm w(\alpha, \beta)\|_{L_{x,v}^2} \|\langle v \rangle^{\frac{\gamma}{2}} \partial_\beta^\alpha f_\pm w(\alpha, \beta)\|_{L_{x,v}^2} \lesssim \|\nabla_x \phi\|_{H_x^2} \|f\|_{Y_k}^2. \end{aligned}$$

This yields (2.32). Here we used the fact (2.30) and its consequence (2.31).

Notice that $\partial_i v \cdot \nabla_x \partial_{\beta-e_i}^\alpha f = \partial_{\beta-e_i}^{\alpha+e_i} f$, $\forall i = 1, 2, 3$, where e_i is the unit vector with i -th component being 1. Then (2.33) follows from (2.24) and (2.31).

When $1 \leq |\alpha_1| \leq 2$, we have

$$\begin{aligned} & (\nabla_x \partial^{\alpha_1} \phi \cdot \nabla_v \partial_\beta^{\alpha-\alpha_1} f, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \lesssim \left(\sum_{|\alpha_1|=1} \|\nabla_x \partial^{\alpha_1} \phi\|_{L_x^\infty} \|\langle v \rangle^{-\frac{\gamma}{2}} \nabla_v \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_v^2} \right. \\ & \quad \left. + \sum_{|\alpha_1|=2} \|\nabla_x \partial^{\alpha_1} \phi\|_{L_x^3} \|\langle v \rangle^{-\frac{\gamma}{2}} \nabla_v \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^6 L_v^2} \right) \|\langle v \rangle^{\frac{\gamma}{2}} \partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_v^2} \\ & \lesssim \|\nabla_x \phi\|_{H_x^3} \sum_{|\alpha_1|=1} \|\langle v \rangle^{-\frac{\gamma}{2}} \nabla_v \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_v^2} \|f\|_{Y_k}. \end{aligned} \quad (2.36)$$

For the Boltzmann case, by (2.25) and (2.9), we have for $|\alpha_1| = 1$ that

$$\begin{aligned} & \|\langle v \rangle^{-\frac{\gamma}{2}} \nabla_v \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_v^2} \leq \|\langle v \rangle^{-\frac{\gamma}{2}} \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 H_v^1} \\ & \lesssim \|\partial_\beta^{\alpha-\alpha_1} f w(|\alpha| - 1, \beta)\|_{L_x^2 H_{\gamma/2}^s} + \|\partial_\beta^{\alpha-\alpha_1} f w(|\alpha| - 1, |\beta| + 1)\|_{L_x^2 H_{\gamma/2}^{1+s}} \lesssim \|f\|_{Y_k}. \end{aligned}$$

For the Landau case, by (2.23) we have

$$\|\langle v \rangle^{-\frac{\gamma}{2}} \nabla_v \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_v^2} \leq \|\langle v \rangle^{-\frac{\gamma}{2}} \partial_\beta^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 H_v^1} \lesssim \|\partial_\beta^{\alpha-\alpha_1} f w(|\alpha| - 1, \beta)\|_{L_x^2 L_D^2} \lesssim \|f\|_{Y_k}.$$

The proof of (2.34) is finished by plugging the above estimates into (2.36). \square

Next, we compute the linear part for the Poisson term.

Lemma 2.19. *For $|\alpha| + |\beta| \leq 2$, any functions f_\pm, ϕ satisfying (1.3) and any ψ satisfies (2.30), there exists $C_k > 0$ such that*

$$|(\partial_\beta^\alpha (\nabla_x \phi \cdot v \mu), e^{\frac{\pm A_{\alpha,\beta}\psi}{\langle v \rangle^2}} \partial_\beta^\alpha f_\pm w^2(\alpha, \beta))_{L_{x,v}^2}| \leq C_k \|\nabla_x \phi\|_{H_x^2} \|f\|_{H_x^2 L_v^2}. \quad (2.37)$$

When $|\beta| = 0$, there exists some $C_\alpha > 0$, such that

$$\begin{aligned} \sum_{\pm} \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial^\alpha f_{\pm} w^2(\alpha, \beta))_{L^2_{x,v}} &\geq C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L^2_x}^2 \\ &\quad - C \|\partial^\alpha \nabla_x \phi\|_{L^2_x} \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L^2_x L^2_v} - C \|\partial^\alpha \nabla_x \phi\|_{L^2_x} \|\nabla_x \psi\|_{H^1_x} \|f\|_{Y_k}. \end{aligned} \quad (2.38)$$

Proof. From integration by parts about ∂_β , we have

$$\begin{aligned} (\partial_\beta^\alpha (\nabla_x \phi \cdot v \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha f w^2(\alpha, \beta))_{L^2_{x,v}} &= (\nabla_x \partial^\alpha \phi \cdot \partial_\beta (v \mu) w^2(\alpha, \beta), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha f)_{L^2_{x,v}} \\ &= ((-1)^{|\beta|} \nabla_x \partial^\alpha \phi \cdot \partial_\beta (\partial_\beta (v \mu) e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} w^2(\alpha, \beta)), \partial^\alpha f)_{L^2_{x,v}}. \end{aligned} \quad (2.39)$$

By Fourier transform, Cauchy-Schwarz inequality, and using the exponential decay of μ , it is direct to show that (2.39) is bounded by

$$\|\nabla_x \partial^\alpha \phi\|_{L^2_x} \left\| \int_{\mathbb{R}^d} \partial^\alpha f \partial_\beta (\partial_\beta (v \mu) e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} w^2(\alpha, \beta)) dv \right\|_{L^2_x} \lesssim \|\nabla_x \phi\|_{H^1_x} \|f\|_{H^2_x L^2_v}.$$

When $\beta = 0$, we temporarily define the projection

$$P f = \left(\int_{\mathbb{R}^3} f(u) du + v \cdot \int_{\mathbb{R}^3} u f(u) du + (|v|^2 - 3) \int_{\mathbb{R}^3} \frac{|u|^2 - 3}{6} f(u) du \right) \mu.$$

Then we split $e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} = 1 + e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1$ and $f_{\pm} = P f + \{I - P\} f$ to obtain

$$\begin{aligned} \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_{\pm})_{L^2_{x,v}} &= \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, w^2(\alpha, 0) P \partial^\alpha f_{\pm})_{L^2_{x,v}} \\ &\quad \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, w^2(\alpha, 0) \{I - P\} \partial^\alpha f_{\pm})_{L^2_{x,v}} \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) w^2(\alpha, 0) \partial^\alpha f_{\pm})_{L^2_{x,v}}. \end{aligned} \quad (2.40)$$

Then one can check that

$$\{I - P\} f_{\pm} = \{I - P\} \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f. \quad (2.41)$$

For the first right hand term of (2.40), we have from (1.7) and (1.3)₂ that

$$\begin{aligned} \sum_{\pm} \pm (\partial^\alpha \nabla_x \phi \cdot v \mu, w^2(\alpha, 0) P \partial^\alpha f_{\pm})_{L^2_{x,v}} &= \sum_{i=1}^3 \left(\partial^\alpha \nabla_x \phi \cdot v \mu, w^2(\alpha, 0) v_i \mu \int_{\mathbb{R}^3} \frac{1}{2} v'_i \partial^\alpha (f_+ - f_-)(v') dv' \right)_{L^2_{x,v}} \\ &= 2C_\alpha \int_{\Omega} \int_{\mathbb{R}^3} \nabla_x \partial^\alpha \phi \cdot v \partial^\alpha (f_+ - f_-)(v) v dv dx = -2C_\alpha \int_{\Omega} \partial^\alpha \phi \nabla_x \cdot \int_{\mathbb{R}^3} v \partial^\alpha (f_+ - f_-)(v) v dv dx \\ &= -2C_\alpha \int_{\Omega} \partial^\alpha \phi \partial_t \Delta_x \phi dx = C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L^2_x}^2, \end{aligned} \quad (2.42)$$

for some constant $C_{\alpha, \beta}$ depending only on α, β . Here we can directly take integration by parts for the case of torus. When Ω is the union of cubes, we need to verify the zero boundary values as the following. Fix $i = 1, 2, 3$. If $\alpha_i = 1$, then $\partial_{x_i} \phi = 0$ on Γ_i . If $\alpha_i = 0, 2$, then by Lemma 2.14 and change of variable $v \mapsto R_x v$, we have on Γ_i that

$$\int_{\mathbb{R}^3} v_i \partial^\alpha (f_+ - f_-)(v) dv = \int_{\mathbb{R}^3} (R_x v)_i \partial^\alpha (f_+ - f_-)(R_x v) dv = - \int_{\mathbb{R}^3} v_i \partial^\alpha (f_+ - f_-)(v) dv = 0.$$

Note that tangent derivatives don't affect the zero boundary values. This completes the integration by parts in (2.42). For the second right hand term of (2.40), noticing exponentially velocity decay in μ and using (2.41), we estimate it by

$$\left| (\partial^\alpha \nabla_x \phi \cdot v \mu, w^2(\alpha, \beta) \{I - P\} \partial^\alpha f_{\pm})_{L^2_{x,v}} \right| \lesssim \|\partial^\alpha \nabla_x \phi\|_{L^2_x} \|\partial^\alpha \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{L^2_x L^2_v}.$$

For the third term in (2.40), noticing $|e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1| \lesssim A_{\alpha, \beta} \|\psi\|_{L^\infty} \lesssim A_{\alpha, \beta} \|\nabla_x \psi\|_{H^1_x}$, we have

$$\left| (\partial^\alpha \nabla_x \phi \cdot v \mu, (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) w^2(\alpha, \beta) \partial^\alpha f_{\pm})_{L^2_{x,v}} \right| \lesssim \|\partial^\alpha \nabla_x \phi\|_{L^2_x} \|\nabla_x \psi\|_{H^1_x} \|f\|_{Y_k}.$$

The above four estimates imply (2.38) and we conclude Lemma 2.19. \square

In [8], one only needs to compute x derivative term, but for our Vlasov-Poisson system we also need estimates about v derivative term.

Lemma 2.20. *Suppose that $\gamma \in (-3, 1]$, $s \in (0, 1)$, $\gamma + 2s > -1$ for Boltzmann case and $\gamma \in [-3, 1]$ for Landau case. For any $|\alpha| \geq 0$, $|\beta| \leq 2$, $k \geq 7$, there exist constants $C_k > 0$ such that, for any functions f, g, ψ , we have for the Boltzmann case,*

$$\begin{aligned} & |(\partial_\beta^\alpha Q(f, \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}| \leq \|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}} \\ & + C_k \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}} + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 L_{\gamma/2}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}}, \end{aligned}$$

and for the Landau case,

$$|(\partial_\beta^\alpha Q(f, \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}| \leq C_k \sum_{\beta_1 \leq \beta} \|\partial_{\beta_1}^\alpha f\|_{L_x^2 L_5^2} \|\partial_\beta^\alpha g\|_{L_x^2 L_5^2}.$$

Proof. For both cases, it's easily seen that

$$(\partial_\beta^\alpha Q(f, \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2} = \sum_{\alpha_1 \leq \alpha} (Q(\partial_{\beta_1}^\alpha f, \partial_{\beta-\beta_1} \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}.$$

For the Boltzmann case, we split it into two cases: $\beta_1 = \beta$ and $\beta_1 < \beta$. For the case $\beta_1 = \beta$, by (2.10) and integration about x , we have

$$\begin{aligned} |(Q(\partial_\beta^\alpha f, \mu), \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}| & \leq \|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}} \\ & + C_k \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}}. \end{aligned}$$

Notice that

$$\left| e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1 \right| \leq A_{\alpha, \beta} \|\psi\|_{L_x^\infty} \langle v \rangle^{-2} \lesssim A_{\alpha, \beta} \|\nabla_x \psi\|_{H_x^1} \langle v \rangle^{-2}. \quad (2.43)$$

By (2.11) we have

$$\begin{aligned} & |(Q(\partial_\beta^\alpha f, \mu), (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}| \\ & \leq C_k \|\partial_\beta^\alpha f w(\alpha, \beta) \langle v \rangle^{-1}\|_{L_x^2 L_{\gamma/2}} \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_\beta^\alpha g w(\alpha, \beta) \langle v \rangle\|_{L_x^2 L_{\gamma/2}} \\ & \leq C_k \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2-1/2}}. \end{aligned}$$

For $\beta_1 < \beta$, by (2.11), (2.31) and integration about x , we have

$$|(Q(\partial_{\beta_1}^\alpha f, \partial_{\beta-\beta_1} \mu), e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} \partial_\beta^\alpha g w^2(\alpha, \beta))_{L_{x,v}^2}| \leq C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}}.$$

So the proof for the Boltzmann case is completed by gathering the three terms. The Landau case can be proved similarly by taking integration in (2.15). \square

The next two coercive estimates play a key role. For the Boltzmann case, we have the following.

Lemma 2.21. *Remind $\|\cdot\|_{Y_k}$ is given in (1.20). Suppose that $-3 < \gamma \leq 1$, $s \in (0, 1)$, $\gamma + 2s > -1$ and $k \geq 14$. For any function f, g , let $G = \mu + g \geq 0$ satisfies $\|G\|_{L^1} \geq A_1$, $\|G\|_{L^2} + \|G\|_{L \log L} \leq A_2$ for some generic constants A_1, A_2 . Then for Q as the Boltzmann collision operator, and any $|\alpha| + |\beta| \leq 2$,*

$$\begin{aligned} (\partial_\beta^\alpha Q(\mu + g, f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} & \leq -\frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}}^2 - \gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}^2 \\ & + C_k \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|g\|_{Y_k} \|f\|_{Y_k} + C_k \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|f\|_{Y_k}^2 + C_k \|\partial_\beta^\alpha f\|_{L_{x,v}^2}^2 \\ & + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{H_{\gamma/2}^s}, \end{aligned} \quad (2.44)$$

for some constants $\gamma_1, C_k \geq 0$. Moreover, suppose ψ satisfies (2.30), we have

$$\begin{aligned} |(\partial_\beta^\alpha Q(g, f), (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2}| & \leq C_k \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \|h\|_{Y_k} \\ & + C_k \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|g\|_{Y_k} \|h\|_{Y_k}, \end{aligned} \quad (2.45)$$

and

$$\begin{aligned}
& |(\partial_\beta^\alpha Q(g, f), \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2}| + |(\partial_\beta^\alpha Q(g, f), e^{\frac{\pm A_{\alpha,\beta}\psi}{\langle v \rangle^2}} \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2}| \\
& \leq C_k \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \min\{\|\langle v \rangle^{2s} f\|_{Y_k}, \|h\|_{Y_k}, \|f\|_{Y_k}\|\langle v \rangle^{2s} h\|_{Y_k}\} + C_k \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|g\|_{Y_k} \|h\|_{Y_k}. \quad (2.46)
\end{aligned}$$

Remark 2.22. By assuming $\|g\|_{L_v^1}$ small, we can check $\|G\|_{L_v^1} = \|\mu + g\|_{L_v^1} \geq A_1$ to apply Lemma 2.21.

Proof. First we have $(\partial_\beta^\alpha Q(G, f), \partial^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} = \sum_{\alpha_1 \leq \alpha, \beta_1 \leq \beta} (Q(\partial_{\beta_1}^{\alpha_1} G, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}$. We split it into several cases. For $\alpha_1 = \beta_1 = 0$, after integrating about x in Theorem 2.8, we have

$$\begin{aligned}
& (Q(G, \partial_\beta^\alpha f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \\
& \leq -\frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2,*}^2}^2 - \gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}^2 + C_k \|\partial_\beta^\alpha f\|_{L_{x,v}^2}^2 \\
& + \int_\Omega C_k \|\partial_\beta^\alpha f\|_{L_{14}^2} \|g w(\alpha, \beta)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} + C_k \|g\|_{L_{14}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s}^2 dx.
\end{aligned}$$

For the case $|\beta_1| > 0, \alpha_1 = 0$, using $\partial_{\beta_1} G = \partial_{\beta_1} \mu + \partial_{\beta_1} g$, we split it into two parts. By (2.22), if $|\beta_1| > 0$, we have $w(\alpha, \beta) \langle v \rangle^{2s} \leq w(\alpha, \beta - \beta_1)$. Then by Corollary 2.5, we have

$$\begin{aligned}
& |(Q(\partial_{\beta_1} \mu, \partial_{\beta-\beta_1}^\alpha f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}| \leq C_k \|\partial_{\beta-\beta_1}^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2+2s}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \\
& \leq C_k \|\partial_{\beta-\beta_1}^\alpha f w(\alpha, \beta - \beta_1)\|_{L_x^2 H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}.
\end{aligned}$$

For the $\partial_{\beta_1} g$ term, by Lemma 2.10, we have

$$\begin{aligned}
& |(Q(\partial_{\beta_1} g, \partial_{\beta-\beta_1}^\alpha f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}| \leq \int_\Omega C_k \|\partial_{\beta-\beta_1}^\alpha f\|_{L_{14}^2} \|\partial_{\beta_1} g w(\alpha, \beta)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} \\
& + C_k \|\partial_{\beta_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} dx,
\end{aligned}$$

For $|\alpha_1| > 0$, since $\partial^{\alpha_1} \mu = 0$, we have $(Q(\partial_{\beta_1}^{\alpha_1} G, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f \langle v \rangle^{2k})_{L_{x,v}^2} = (Q(\partial_{\beta_1}^{\alpha_1} g, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f \langle v \rangle^{2k})_{L_{x,v}^2}$. By Lemma 2.10, we have

$$\begin{aligned}
& |(Q(\partial_{\beta_1}^{\alpha_1} g, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f \langle v \rangle^{2k})_{L_{x,v}^2}| \leq \int_\Omega C_k \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f\|_{L_{14}^2} \|\partial_{\beta_1}^{\alpha_1} g w(\alpha, \beta)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_{\gamma/2}^2} dx \\
& + C_k \int_\Omega \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} dx.
\end{aligned}$$

Gathering all the terms, we have

$$\begin{aligned}
& (\partial_\beta^\alpha Q(\mu + g, f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \leq -\frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2,*}^2}^2 \\
& - \gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}^2 + C_k \|\partial_\beta^\alpha f\|_{L_{x,v}^2}^2 + C_k \sum_{|\beta_1| < |\beta|} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \\
& + C_k \int_\Omega \sum_{\alpha_1 \leq \alpha, \beta_1 \leq \beta} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f\|_{L_{14}^2} \|\partial_{\beta_1}^{\alpha_1} g w(\alpha, \beta)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} dx \\
& + C_k \int_\Omega \sum_{\alpha_1 \leq \alpha, \beta_1 \leq \beta, |\alpha_1| + |\beta_1| > 0} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} dx \\
& + C_k \int_\Omega \|g\|_{L_{14}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s}^2 dx. \quad (2.47)
\end{aligned}$$

Now we only need to prove that

$$\int_\Omega \|g\|_{L_{14}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s}^2 dx \lesssim \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|f\|_{Y_k}^2, \quad (2.48)$$

and for all $\alpha_1 \leq \alpha, \beta_1 \leq \beta, |\alpha_1| + |\beta_1| > 0$,

$$\int_\Omega \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s} dx \lesssim \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|f\|_{Y_k}^2. \quad (2.49)$$

The fifth term on the right-hand side of (2.47) follows similarly by changing the order f and g . First, for the case $|\alpha_1| = |\beta_1| = 0$, we have

$$\int_\Omega \|g\|_{L_{14}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s}^2 dx \lesssim \|g\|_{L_x^\infty L_{14}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}^2 \lesssim \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|f\|_{Y_k}^2.$$

This gives (2.48). For (2.49), we split it into two cases: $|\alpha_1| + |\beta_1| = 1$ and $|\alpha_1| + |\beta_1| = 2$. For the case $|\alpha_1| + |\beta_1| = 1$, by $\|fg\|_{L_x^2} \leq \|f\|_{L_x^6} \|g\|_{L_x^3}$, we have

$$\begin{aligned} & \int_{\Omega} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_{\beta}^{\alpha} fw(\alpha, \beta)\|_{H_{\gamma/2}^s} dx \\ & \lesssim \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^6} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^3 H_{\gamma/2+2s}^s} \|\partial_{\beta}^{\alpha} fw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \\ & \lesssim \langle v \rangle^{14} g \|_{H_{x,v}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^3 H_{\gamma/2+2s}^s} \|f\|_{Y_k}. \end{aligned} \quad (2.50)$$

We again split it into two parts $|\alpha| + |\beta| = 1$ and $|\alpha| + |\beta| = 2$. For the case $|\alpha_1| + |\beta_1| = |\alpha| + |\beta| = 1$, we have $|\alpha - \alpha_1| = |\beta - \beta_1| = 0$. Then by (2.8) and (2.26), we have

$$\|fw(\alpha, \beta)\|_{L_x^3 H_{\gamma/2+2s}^s} \lesssim \|fw(0, 0)\|_{L_x^2 H_{\gamma/2}^s} + \|fw(1, 0)\|_{H_x^1 H_{\gamma/2}^s} \lesssim \|f\|_{Y_k}.$$

We then consider the case $|\alpha_1| + |\beta_1| = 1$, $|\alpha| + |\beta| = 2$. This time we have $|\alpha - \alpha_1| = 1$, $|\beta - \beta_1| = 0$ or $|\alpha - \alpha_1| = 0$, $|\beta - \beta_1| = 1$. For the first case, by (2.27),

$$\|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^3 H_{\gamma/2+2s}^s} \leq \|fw(1, 0)\|_{H_x^1 H_{\gamma/2}^s} + \|fw(2, 0)\|_{H_x^2 H_{\gamma/2}^s} \lesssim \|f\|_{Y_k}.$$

For the second case, $|\beta| \geq 1$ and by (2.27), we obtain

$$\|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^3 H_{\gamma/2+2s}^s} \leq \|fw(0, 1)\|_{L_x^2 H_{\gamma/2}^{1+s}} + \|fw(1, 1)\|_{H_x^1 H_{\gamma/2}^{1+s}} \lesssim \|f\|_{Y_k}.$$

For the case $|\alpha_1| + |\beta_1| = 2$, we have $|\alpha| + |\beta| = 2$ and $|\alpha - \alpha_1| = |\beta - \beta_1| = 0$. Thus

$$\begin{aligned} & \int_{\Omega} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{H_{\gamma/2+2s}^s} \|\partial_{\beta}^{\alpha} fw(\alpha, \beta)\|_{H_{\gamma/2}^s} dx \\ & \lesssim \|\partial_{\beta}^{\alpha} g\|_{L_x^2 L_{14}^2} \|fw(\alpha, \beta)\|_{L_x^{\infty} H_{\gamma/2+2s}^s} \|\partial_{\beta}^{\alpha} fw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \lesssim \langle v \rangle^{14} g \|_{H_{x,v}^2} \|fw(\alpha, \beta)\|_{L_x^{\infty} H_{\gamma/2+2s}^s} \|f\|_{Y_k}. \end{aligned} \quad (2.51)$$

By (2.7) with suitable k and (2.28), we have

$$\|fw(\alpha, \beta)\|_{L_x^{\infty} H_{\gamma/2+2s}^s} \leq \|fw(1, 0)\|_{H_x^1 H_{\gamma/2}^s} + \|fw(2, 0)\|_{H_x^2 H_{\gamma/2}^s} \lesssim \|f\|_{Y_k}.$$

These estimates imply (2.49). Substituting (2.48) and (2.49) into (2.47), we obtain (2.44). For (2.45), by (2.13) and (2.43), we have

$$\begin{aligned} & |(\partial_{\beta}^{\alpha} Q(g, f), (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw^2(\alpha, \beta))_{L_{x,v}^2}| \\ & \lesssim \sum_{\substack{\alpha_1 \leq \alpha \\ \beta_1 \leq \beta}} \left(\min \left\{ \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{H_{\gamma/2}^s} \|L_x^2 \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \langle v \rangle^{2s} \partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}, \right. \right. \\ & \quad \left. \left. + \|\partial_{\beta_1}^{\alpha_1} g\|_{L_{14}^2} \|\langle v \rangle^{2s} \partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{H_{\gamma/2}^s} \|L_x^2 \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \right\} \right. \\ & \quad \left. + \|\partial_{\beta_1}^{\alpha_1} f\|_{L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} gw(\alpha, \beta)\|_{H_{\gamma/2}^s} \|L_x^2 \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \right). \end{aligned} \quad (2.52)$$

Applying $L^{\infty} - L^2$ and $L^3 - L^6$ Hölder's inequality and using the first term in the minimum, the first right-hand term of (2.52) can be estimated as

$$\begin{aligned} & \left(\sum_{|\alpha_1|+|\beta_1|=0} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^{\infty} L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L^2 H_{\gamma/2}^s} + \sum_{|\alpha_1|+|\beta_1|=1} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^3 L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^6 H_{\gamma/2}^s} \right. \\ & \quad \left. + \sum_{|\alpha_1|+|\beta_1|=2} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2 L_{14}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^{\infty} H_{\gamma/2}^s} \right) A_{\alpha, \beta} \|\nabla_x \psi\|_{H_x^1} \|\partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \\ & \lesssim \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \|\partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}. \end{aligned} \quad (2.53)$$

The second right-hand term of (2.52) can be estimated similarly and then we deduce that

$$\begin{aligned} & |(\partial_{\beta}^{\alpha} Q(g, f), (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw^2(\alpha, \beta))_{L_{x,v}^2}| \lesssim \|\langle v \rangle^{14} g\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \|\partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s} \\ & \quad + \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|g\|_{Y_k} \|\partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}. \end{aligned}$$

The proof of (2.46) is similar by replacing the term $(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1)$ by 1 and $e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}}$. Note that we keep the minimum in (2.52) this time, and the proof is omitted for brevity. Thus Lemma 2.21 is proved. \square

We can also prove a similar result for the Landau case.

Lemma 2.23. *Let $-3 \leq \gamma \leq 1$, Q be the Landau collision operator and $k \geq 7$. There exists a constant $C_k > 0$ such that for any function f, g, h , and any function ψ satisfies (2.30), we have the following.*

(1) For any $|\alpha| + |\beta| \leq 2$,

$$\begin{aligned} (\partial_\beta^\alpha Q(\mu, f), \partial_\beta^\alpha f w^2(\alpha, \beta)) &\leq -\gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_D^2}^2 + C_k \|\partial_\beta^\alpha f\|_{L_x^2 L_D^2}^2 \\ &\quad + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 L_D^2} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_D^2}. \end{aligned} \quad (2.54)$$

(2) Remind the norm $\|\cdot\|_{Y_k}$ is given in (1.20). For any $|\alpha| + |\beta| \leq 2$, we have

$$\left| (\partial_\beta^\alpha Q(g, f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \right| \leq C_k \|g\langle v \rangle^7\|_{H_{x,v}^2} \|f\|_{Y_k}^2, \quad (2.55)$$

$$\left| (\partial_\beta^\alpha Q(g, f), (e^{\frac{\pm A_{\alpha,\beta}\psi}{\langle v \rangle^2}} - 1) \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2} \right| \leq C_k \|g\langle v \rangle^7\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \|h\|_{Y_k}, \quad (2.56)$$

and

$$\begin{aligned} \left| (\partial_\beta^\alpha Q(g, f), \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2} \right| &+ \left| (\partial_\beta^\alpha Q(g, f), e^{\frac{\pm A_{\alpha,\beta}\psi}{\langle v \rangle^2}} \partial_\beta^\alpha h w^2(\alpha, \beta))_{L_{x,v}^2} \right| \\ &\leq C_k \|g\langle v \rangle^7\|_{H_{x,v}^2} \min\{\|f\|_{Y_k} \|\langle v \rangle h\|_{Y_k}, \|\langle v \rangle f\|_{Y_k} \|h\|_{Y_k}\}. \end{aligned} \quad (2.57)$$

Proof. Notice that $(\partial_\beta^\alpha Q(\mu, f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} = \sum_{\alpha_1 \leq \alpha, \beta_1 \leq \beta} (Q(\partial_{\beta_1}^{\alpha_1} \mu, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}$. We split it into two cases: $\beta_1 = 0$ and $|\beta_1| > 0$. If $\beta_1 = 0$, after integrating about x in (2.16), we have

$$(Q(\mu, \partial_\beta^\alpha f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \leq -\gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_D^2}^2 + C_k \|\partial_\beta^\alpha f\|_{L_x^2 L_D^2}^2.$$

For the case $|\beta_1| > 0$, by (2.23), $w(\alpha, \beta)\langle v \rangle \leq w(\alpha, \beta - \beta_1)$. Then by Lemma 2.12 we have

$$\begin{aligned} \left| (Q(\partial_{\beta_1}^{\alpha_1} \mu, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \right| &\leq C_k \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_{D,1}^2} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_D^2} \\ &\leq C_k \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta - \beta_1)\|_{L_x^2 L_D^2} \|\partial_\beta^\alpha g w(\alpha, \beta)\|_{L_x^2 L_D^2}, \end{aligned}$$

So (2.54) is thus finished by gathering the above two terms together. Next, for (2.55), notice that

$$(\partial_\beta^\alpha Q(g, f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} = \sum_{\alpha_1 \leq \alpha, \beta_1 \leq \beta} (Q(\partial_{\beta_1}^{\alpha_1} g, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2}.$$

We again split it into two cases: $|\alpha_1| = |\beta_1| = 0$ and $|\alpha_1| + |\beta_1| > 0$. For the case $|\alpha_1| = |\beta_1| = 0$, by (2.17) and (2.43), we have

$$\begin{aligned} \left| (Q(g, \partial_\beta^\alpha f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \right| &\leq \int_{\Omega} C_k \|g\|_{L_x^2} \left\| \left(e^{\frac{\pm A_{\alpha,\beta}\psi}{\langle v \rangle^2}} - 1 \right) \langle v \rangle \partial_\beta^\alpha f w(\alpha, \beta) \right\|_{L_D^2}^2 dx \\ &\leq C_k \|g\|_{L_x^\infty L_D^2} \|\nabla_x \psi\|_{H_x^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_D^2}^2 \leq C_k \|g\langle v \rangle^7\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k}^2. \end{aligned}$$

For $\alpha_1 \leq \alpha, \beta_1 \leq \beta, |\alpha_1| + |\beta_1| > 0$, by (2.17), we have

$$\left| (Q(\partial_{\beta_1}^{\alpha_1} g, \partial_{\beta-\beta_1}^{\alpha-\alpha_1} f), \partial_\beta^\alpha f w^2(\alpha, \beta))_{L_{x,v}^2} \right| \leq \int_{\Omega} C_k \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_{D,1}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_D^2} dx.$$

We split it into two cases, $|\alpha_1| + |\beta_1| = 1$ and $|\alpha_1| + |\beta_1| = 2$. For the case $|\alpha_1| + |\beta_1| = 1$, by $\|fg\|_{L_x^2} \leq \|f\|_{L_x^6} \|g\|_{L_x^3}$ and similarly as (2.50), we have

$$\int_{\Omega} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_{D,1}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_D^2} dx \lesssim \|\langle v \rangle^7 g\|_{H_{x,v}^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^2 L_{D,1}^2} \|f\|_{Y_k}.$$

We again split it into two parts $|\alpha| + |\beta| = 1$ and $|\alpha| + |\beta| = 2$. For the case $|\alpha_1| + |\beta_1| = |\alpha| + |\beta| = 1$, we have $|\alpha - \alpha_1| = |\beta - \beta_1| = 0$. By (2.26) and (2.8) we obtain that

$$\|f w(\alpha, \beta)\|_{L_x^3 L_{D,1}^2} \lesssim \|f w(0, 0)\|_{L_x^2 L_D^2} + \|f w(1, 0)\|_{H_x^1 L_D^2} \lesssim \|f\|_{Y_k}.$$

For the case $|\alpha| + |\beta| = 2$, either $|\alpha - \alpha_1| = 1, |\beta - \beta_1| = 0$ or $|\alpha - \alpha_1| = 0, |\beta - \beta_1| = 1$. For the first case, by (2.27) and (2.8), we have

$$\|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^3 L_{D,1}^2} \leq \|f w(1, 0)\|_{H_x^1 L_D^2} + \|f w(2, 0)\|_{H_x^2 L_D^2} \lesssim \|f\|_{Y_k}.$$

For the second case, this time $|\beta| \geq 1$. By (2.27) and (2.8), we have

$$\|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} f w(\alpha, \beta)\|_{L_x^3 L_{D,1}^2} \leq \|\nabla_v f w(0, 1)\|_{L_x^2 L_D^2} + \|\nabla_v f w(1, 1)\|_{H_x^1 L_D^2} \lesssim \|f\|_{Y_k}.$$

For the case $|\alpha_1| + |\beta_1| = 2$, we have obviously $|\alpha| + |\beta| = 2$ and $|\alpha - \alpha_1| = |\beta - \beta_1| = 0$. Similar to (2.51), we have

$$\int_{\Omega} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_D^2} \|\partial_{\beta}^{\alpha} fw(\alpha, \beta)\|_{L_D^2} dx \lesssim \|\langle v \rangle^7 g\|_{H_{x,v}^2} \|fw(\alpha, \beta)\|_{L_x^{\infty} L_D^2} \|f\|_{Y_k}.$$

We obtain again from (2.7) and (2.28) that

$$\|fw(\alpha, \beta)\|_{L_x^{\infty} L_D^2} \leq \|fw(1, 0)\|_{H_x^1 L_D^2} + \|fw(2, 0)\|_{H_x^2 L_D^2} \lesssim \|f\|_{Y_k}.$$

The proof of (2.55) is done after gathering the above terms. For (2.56), by (2.17) and (2.43), we have

$$\begin{aligned} & \left| (\partial_{\beta}^{\alpha} Q(g, f), (e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw^2(\alpha, \beta)) \right|_{L_{x,v}^2} \\ & \lesssim \sum_{\alpha_1 \leq \alpha, \beta \leq \beta_1} \int_{\Omega} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2} \min \{ \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_D^2} \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \langle v \rangle \partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_D^2}, \\ & \quad \|\langle v \rangle \partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_D^2} \|(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1) \partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_D^2} \} \\ & \lesssim \left(\sum_{|\alpha_1|+|\beta_1|=0} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^{\infty} L_D^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^2 L_D^2} + \sum_{|\alpha_1|+|\beta_1|=1} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^3 L_D^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^6 L_D^2} \right. \\ & \quad \left. + \sum_{|\alpha_1|+|\beta_1|=2} \|\partial_{\beta_1}^{\alpha_1} g\|_{L_x^2 L_D^2} \|\partial_{\beta-\beta_1}^{\alpha-\alpha_1} fw(\alpha, \beta)\|_{L_x^{\infty} L_D^2} \right) A_{\alpha, \beta} \|\nabla_x \psi\|_{H_x^1} \|\partial_{\beta}^{\alpha} hw(\alpha, \beta)\|_{L_x^2 L_D^2} \\ & \lesssim \|\langle v \rangle^7 g\|_{H_{x,v}^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \|h\|_{Y_k}, \end{aligned} \quad (2.58)$$

where we apply $L^{\infty} - L^2$ and $L^3 - L^6$ Hölder's inequality. The proof of (2.57) is similar by replacing the term $(e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}} - 1)$ by 1 and $e^{\frac{\pm A_{\alpha, \beta} \psi}{\langle v \rangle^2}}$. Note that this time we keep the minimum in (2.58), and we omit the proof for brevity. This completes the proof of Lemma 2.23. \square

For the local existence, similar to [6] we have

Theorem 2.24. *Suppose that $-3 < \gamma \leq 1$ for Boltzmann case and $-3 \leq \gamma \leq 1$ for Landau case. For any $k \geq k_0 + 2$, there exist small constants $\varepsilon_0, \tau_0, T_0 > 0$, such that if $f_0 \in X_{k_0}$ satisfies $\mu + f_0 \geq 0$ and*

$$\mathcal{E}_{k_0}(0) \leq \varepsilon_0, \quad \mathcal{E}_k(0) \leq A, \quad (2.59)$$

for some $A > 0$, Then the Cauchy problem

$$\begin{cases} \partial_t f_{\pm} + v \cdot \nabla_x f_{\pm} \mp \nabla_x \phi \cdot \nabla_v f_{\pm} \pm \nabla_x \phi \cdot v \mu = Q(f_{\pm} + f_{\mp}, \mu) + Q(\mu + f_{\pm}, f_{\pm}) + Q(\mu + f_{\mp}, f_{\pm}), \\ -\Delta_x \phi = \int_{\mathbb{R}^3} (f_+ - f_-) dv, \quad \int_{\Omega} \phi(x) dx = 0, \quad f(0) = f_0, \quad E(0) = E_0, \\ f(t, x, R_x v) = f(t, x, v) \text{ on } \gamma_- \text{ and } \partial_n \phi = 0 \text{ on } \partial\Omega, \text{ if } \Omega \text{ is given by (1.10),} \end{cases} \quad (2.60)$$

admits a unique weak solution $f \in L_{T_0}^{\infty} X_k$ satisfying $\mu + f \geq 0$ and

$$\sup_{0 \leq t \leq T_0} \mathcal{E}_{k_0}(t) \leq \tau_0, \quad \sup_{0 \leq t \leq T_0} \mathcal{E}_k(t) \leq A. \quad (2.61)$$

3. GLOBAL REGULARITY

In this section, we prove the Theorem 1.2. We first establish the macroscopic estimates.

3.1. Macroscopic estimate. We recall the operator \mathbf{P} defined in (1.5). It is direct to obtain $\|a\|_{L_x^2}^2 + \|b\|_{L_x^2}^2 + \|c\|_{L_{x,v}^2}^2 \sim \|\mathbf{P}f\|_{L_{x,v}^2}^2$. We also rewrite the equation (1.3) as

$$\partial_t f_{\pm} + v \cdot \nabla_x f_{\pm} \pm \nabla_x \phi \cdot v \mu - L_{\pm} f = N_{\pm}(f), \quad -\Delta_x \phi = \int_{\mathbb{R}^3} (f_+ - f_-) dv, \quad (3.1)$$

with initial data $f(0) = f_0, E(0) = E_0$, where $L = [L_+, L_-]$ and $N = [N_+, N_-]$ are given by

$$L_{\pm} f := Q(f_{\pm} + f_{\mp}, \mu) + 2Q(\mu, f_{\pm}), \quad N_{\pm}(f) := \pm \nabla_x \phi \cdot \nabla_v f_{\pm} + Q(f_{\pm}, f_{\pm}) + Q(f_{\mp}, f_{\pm}). \quad (3.2)$$

Our next goal is to estimate $a(t, x), b(t, x), c(t, x)$ in terms of $\{\mathbf{I} - \mathbf{P}\}f$. In contrast to [39], our P is not symmetric and we can not compare the v_i, v_j terms on both sides. The following Lemma gives the macroscopic estimates.

Lemma 3.1. For both Landau and Boltzmann case, suppose f solves (3.1). For any integer $m \geq 0$, there exists function $G = G(t)$ satisfying

$$G(t) \lesssim \sum_{|\alpha| \leq 1} \|\partial^\alpha f\|_{L_x^2 L_{10}^2} \|\partial^\alpha \nabla_x \mathbf{P}f\|_{L_{x,v}^2}, \quad (3.3)$$

such that

$$\partial_t G(t) + \lambda \| [a_+, a_-, b, c] \|_{H_x^2}^2 + \lambda \|\nabla_x \phi\|_{H_x^2}^2 \lesssim \|\{\mathbf{I} - \mathbf{P}\}f\|_{H_x^2 L_{10}^2}^2 + \|N_\parallel\|_{H_x^1 L_v^2}^2 + \|\nabla_x \phi\|_{L_x^2}^4, \quad (3.4)$$

Here N_\parallel is the inner product of $N(f)$ with some linear combination of $(1, v_i, v_i v_j, v_i^2, v_i |v|^2)$ over $v \in \mathbb{R}^3$.

Proof. Here we only consider the non-cutoff Boltzmann case and the Landau case is similar. The proof follows the idea in [39, Section 6]. Splitting $f = \mathbf{P}f + \{\mathbf{I} - \mathbf{P}\}f$, we rewrite (3.1) to be

$$\begin{aligned} (\partial_t a_\pm + \partial_t b \cdot v + \partial_t c(|v|^2 - 3))\mu + v \cdot \nabla_x (a_\pm + b \cdot v + c(|v|^2 - 3))\mu \pm \nabla_x \phi \cdot v\mu \\ = -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f). \end{aligned} \quad (3.5)$$

Taking the inner product of (3.5) with $1, v_i^2, |v|^2, v_i v_j, i, j = 1, 2, 3, i \neq j$ over $v \in \mathbb{R}^3$, we have

$$\begin{aligned} \partial_t a_\pm + \nabla_x \cdot b &= -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), 1)_{L_v^2}, \\ \partial_t a_\pm + 2\partial_t c + 2\partial_{x_i} b_i + \nabla_x \cdot b &= -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), v_i^2)_{L_v^2}, \\ 3\partial_t a_\pm + 6\partial_t c + 5\nabla_x \cdot b &= -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), |v|^2)_{L_v^2}. \end{aligned} \quad (3.6)$$

and

$$\partial_{x_j} b_i + \partial_{x_i} b_j = -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f, v_i v_j)_{L_v^2} + (N_\pm(f), v_i v_j)_{L_v^2} =: \gamma_{1ij} + \gamma_{2ij}, \quad i \neq j. \quad (3.7)$$

By taking the subtraction of (3.6)₂ and (3.6)₁, we have

$$\begin{aligned} \partial_t c + \partial_{x_i} b_i &= \left(-(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f, \frac{|v_i|^2 - 1}{2} \right)_{L_v^2} + (N_\pm(f), \frac{|v_i|^2 - 1}{2})_{L_v^2} \\ &=: \gamma_{1i} + \gamma_{2i}. \end{aligned} \quad (3.8)$$

Taking the inner product of (3.5) with $v, v|v|^2$ over $v \in \mathbb{R}^3$, we have

$$\begin{aligned} \partial_t b + \nabla_x (a_\pm \pm \phi) + 2\nabla_x c &= -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), v)_{L_v^2}, \\ 5\partial_t b + 5\nabla_x (a_\pm \pm \phi) + 20\nabla_x c &= -(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), v|v|^2)_{L_v^2}. \end{aligned} \quad (3.9)$$

The two identities in (3.9) implies

$$\nabla_x c = \left(-(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), \frac{v(|v|^2 - 5)}{10} \right)_{L_v^2}, \quad (3.10)$$

and

$$\nabla_x a_\pm = -\partial_t b \mp \nabla_x \phi + \left(-(\partial_t + v \cdot \nabla_x)\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f - L_\pm\{\mathbf{I} - \mathbf{P}\}f + N_\pm(f), \frac{v(10 - |v|^2)}{5} \right)_{L_v^2}. \quad (3.11)$$

Here we use the fact that $\int_{\mathbb{R}^3} |v|^2 \mu dv = 3, \int_{\mathbb{R}^3} |v|^4 \mu dv = 15, \int_{\mathbb{R}^3} |v|^6 \mu dv = 105$. By the definitions (1.5) of $\mathbf{P}f$ and (3.2) of L , we have for $\psi(v) = 1, v_i, |v|^2$ that

$$(\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, \psi(v)) = 0, (L_\pm\{\mathbf{I} - \mathbf{P}\}f, \psi(v)) = 0, (\partial_t\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, \psi(v)) = 0, (v \cdot \nabla_x\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, 1) = 0.$$

Thus the inequalities (3.6)₁, (3.6)₃ and (3.9)₁ (i.e. inner product with $1, v_i, |v|^2$) become

$$\begin{aligned} \partial_t a_\pm + \nabla_x \cdot b &= (N_\pm(f), 1)_{L_v^2}, \\ \partial_t b + \nabla_x a_\pm + 2\nabla_x c \pm \nabla_x \phi &= -(v \cdot \nabla_x\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, v)_{L_v^2} + (N_\pm(f), v)_{L_v^2}, \\ 3\partial_t a_\pm + 6\partial_t c + 5\nabla_x \cdot b &= -(v \cdot \nabla_x\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, |v|^2)_{L_v^2} + (N_\pm(f), |v|^2)_{L_v^2}. \end{aligned} \quad (3.12)$$

Combining (3.12)₁ and (3.12)₃, we have

$$\partial_t c + \frac{1}{3}\nabla_x \cdot b = -\frac{1}{6}(v \cdot \nabla_x\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f, |v|^2)_{L_v^2} + \left(N_\pm(f), \frac{|v|^2 - 3}{6} \right)_{L_v^2}. \quad (3.13)$$

For brevity, we define

$$\xi_a = \frac{v(10 - |v|^2)}{5}, \quad \xi_{bi} = \frac{2v_i^2 - 5}{2}, \quad \xi_c = \frac{v(|v|^2 - 5)}{10}. \quad (3.14)$$

It follows that

$$|(v \cdot \nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f, \xi_a)| + |(v \cdot \nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f, \xi_{bi})| + |(v \cdot \nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f, v_i v_j)| \\ + |(v \cdot \nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f, \xi_c)| \leq \|\nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f\|_{L^2_\nu}.$$

For the Boltzmann case, by Lemma 2.4, we have

$$|(Q(\mu, f), \xi_a)| \leq \|\mu\|_{L^2_{10}} \|f\|_{L^2_{10}} \|\xi_a\|_{H^2_{-7}} \leq C \|f\|_{L^2_{10}}, \quad |(Q(f, \mu), \xi_a)| \leq \|f\|_{L^2_{10}} \|\mu\|_{L^2_{10}} \|\xi_a\|_{H^2_{-7}} \leq C \|f\|_{L^2_{10}}.$$

For the Landau case, by Lemma 2.13, we have

$$|(Q(\mu, f), \xi_a)| \leq \|\mu\|_{L^2_{10}} \|f\|_{L^2_{10}} \|\xi_a\|_{H^2_{-5}} \leq C \|f\|_{L^2_{10}}, \quad |(Q(f, \mu), \xi_a)| \leq \|f\|_{L^2_{10}} \|\mu\|_{L^2_{10}} \|\xi_a\|_{H^2_{-5}} \leq C \|f\|_{L^2_{10}}.$$

Similar arguments can be carried on inner product with ξ_{bi} , $v_i v_j$, ξ_c . Recalling the definition (3.2) of $L = [L_+, L_-]$, we have

$$|(Lf, \xi_a)_{L^2_\nu}| + |(Lf, \xi_{bi})_{L^2_\nu}| + |(Lf, v_i v_j)_{L^2_\nu}| + |(Lf, \xi_c)_{L^2_\nu}| \lesssim \|f\|_{L^2_{10}}. \quad (3.15)$$

Step 1: Computation of $\nabla_x \partial^\alpha c$. Let $|\alpha| \leq 2$. Applying ∂^α on both side of (3.10) and taking inner product with $\nabla_x \partial^\alpha c$ over $x \in \Omega$, we deduce that

$$\|\nabla_x \partial^\alpha c\|_{L^2_x}^2 \leq -\frac{d}{dt} \int_\Omega (\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f, \xi_c)_{L^2_\nu} \cdot \nabla_x \partial^\alpha c \, dx - \int_\Omega (\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \nabla_x \partial^\alpha f, \xi_c)_{L^2_\nu} \cdot \partial_t \partial^\alpha c \, dx \\ + C \|\nabla_x \partial^\alpha c\|_{L^2_x} (\|\partial^\alpha \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f\|_{H^1_x L^2_{10}} + \|\partial^\alpha N\|_{L^2_x}). \quad (3.16)$$

Here, for the torus case, one can take integration by parts directly. For the union of cubes case, we have from boundary (1.6) that $\partial_{x_i} c = 0$ on Γ_i . On the other hand, when $\alpha_i = 0, 2$, we have from Lemma 2.14 and definition (3.14) of ξ_c that for $x \in \Gamma_i$,

$$(\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f, \xi_c)_{L^2_\nu} = (\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f(R_x v), \xi_c(R_x v))_{L^2_\nu} = -(\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f(v), \xi_c(v))_{L^2_\nu} = 0.$$

Note that $\partial^\alpha \mathbf{P}_\pm f(R_x v) = \partial^\alpha \mathbf{P}_\pm f(v)$ by using (2.21). This implies the integration by parts about ∇_x in (3.16). For the term $\partial_t \partial^\alpha c$ in (3.16), it follows from (3.13) that

$$\partial_t \partial^\alpha c = \frac{1}{3} \nabla_x \partial^\alpha b - \frac{1}{6} (v \cdot \nabla_x \{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f, |v|^2)_{L^2_\nu} + \left(\partial^\alpha N_\pm(f), \frac{|v|^2 - 3}{6} \right)_{L^2_\nu}.$$

By Cauchy-Schwarz inequality, for any small $\varepsilon > 0$, we have

$$-\int_\Omega (\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \nabla_x \partial^\alpha f, \xi_c)_{L^2_\nu} \cdot \partial_t \partial^\alpha c \, dx \leq \varepsilon \|\nabla_x \partial^\alpha b\|^2 + \frac{C}{\varepsilon} \|\nabla_x \partial^\alpha \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f\|_{L^2_x L^2_{10}}^2 + \varepsilon \|\partial^\alpha N\|_{L^2_x}^2.$$

Plugging this into (3.16), we deduce that

$$\|\nabla_x \partial^\alpha c\|^2 \leq -\frac{d}{dt} \int_\Omega (\{\mathbf{I}_\pm - \mathbf{P}_\pm\} \partial^\alpha f, \xi_c)_{L^2_\nu} \cdot \nabla_x \partial^\alpha c \, dx + \varepsilon \|\nabla_x \partial^\alpha b\|^2 + \varepsilon \|\nabla_x \partial^\alpha c\|^2 \\ + \frac{C}{\varepsilon} (\|\partial^\alpha \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f\|_{H^1_x L^2_{10}}^2 + \|\partial^\alpha N\|_{L^2_x}^2). \quad (3.17)$$

Step 2: Computation of $\nabla_x \partial^\alpha b$. For fixed i , we use (3.7), (3.8) to compute

$$\Delta_x b_i = \sum_{j \neq i} \partial_{x_j x_j} b_i + \partial_{x_i x_i} b_i = \sum_{j \neq i} (-\partial_{x_i x_j} b_j + \partial_{x_j} (\gamma_{1ij} + \gamma_{2ij})) + \partial_{x_i} (\gamma_{1i} + \gamma_{2i}) - \partial_t \partial_{x_i} c \\ = \sum_{j \neq i} (\partial_t \partial_{x_i} c - \partial_{x_i} (\gamma_{1j} + \gamma_{2j})) + \sum_{j \neq i} \partial_{x_j} (\gamma_{1ij} + \gamma_{2ij}) + \partial_{x_i} (\gamma_{1i} + \gamma_{2i}) - \partial_t \partial_{x_i} c \\ = \sum_{j \neq i} (\partial_{x_j} (\gamma_{1ij} + \gamma_{2ij}) - \partial_{x_i} (\gamma_{1j} + \gamma_{2j})) - \partial_{x_i x_i} b_i + 2\partial_{x_i} (\gamma_{1i} + \gamma_{2i}). \quad (3.18)$$

We can rewrite the linear terms $\sum_{j \neq i} (\partial_{x_j} \gamma_{1ij} - \partial_{x_i} \gamma_{1j}) + 2\partial_{x_i} \gamma_{1i}$ including γ_{1i} , γ_{1j} , γ_{1ij} as the linear combination of $\partial_{x_j} (-(\partial_t + v \cdot \nabla_x) \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f, \xi_{ij})_{L^2_\nu} - (L_\pm \{\mathbf{I} - \mathbf{P}\} f, \xi_{ij})_{L^2_\nu}$, where ξ_{ij} are certain linear combinations of the basis $\{1, v_i v_j, |v_i|^2\}_{i,j=1}^3$. Similar to (3.15), we have $|(Lf, \xi_{ij})_{L^2_\nu}| \lesssim \|f\|_{L^2_{10}}$. Note that $-(\Delta_x \partial^\alpha b_i, \partial^\alpha b_i)_{L^2_x} = \|\nabla_x \partial^\alpha b_i\|_{L^2_x}^2$ and $(\partial_{x_i x_i} \partial^\alpha b_i, \partial^\alpha b_i)_{L^2_x} = -\|\partial_{x_i} \partial^\alpha b_i\|_{L^2_x}^2$, which is trivial for the torus case and follows from (2.21) for the union of cubes case. Applying ∂^α on both side of (3.18), taking inner product with $\partial^\alpha b_i$ over $x \in \Omega$ and summation over $i = 1, 2, 3$, we deduce that

$$\|\nabla_x \partial^\alpha b\|_{L^2_x}^2 = \sum_i \left(-\int_\Omega \sum_{j \neq i} (\partial_{x_j} \partial^\alpha (\gamma_{1ij} + \gamma_{2ij}) - \partial_{x_i} \partial^\alpha (\gamma_{1j} + \gamma_{2j})) \cdot \partial^\alpha b_i \, dx \right) \\ - \sum_i \left(\int_\Omega |\partial_{x_i} \partial^\alpha b_i|^2 \, dx - 2 \int_\Omega \partial_{x_i} \partial^\alpha (\gamma_{1i} + \gamma_{2i}) \cdot \partial^\alpha b_i \, dx \right)$$

$$\begin{aligned} &\leq \sum_{i,j} \frac{d}{dt} \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial_{x_j} \partial^{\alpha} f, \xi_{ij})_{L_v^2} \cdot \partial^{\alpha} b_i dx - \sum_{i,j} \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial_{x_j} \partial^{\alpha} f, \xi_{ij})_{L_v^2} \cdot \partial_t \partial^{\alpha} b_i dx \\ &\quad + C \|\nabla_x \partial^{\alpha} b\| (\|\partial^{\alpha} \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{H_x^1 L_{10}^2} + \|\partial^{\alpha} N_{\parallel}\|_{L_x^2}). \end{aligned} \quad (3.19)$$

Here we used integration by parts on ∂_{x_i} and ∂_{x_j} . For the case of the union of cubes, we need the following boundary values to complete the integration by parts. For $i = 1, 2, 3$ and $\alpha = (\alpha_1, \alpha_2, \alpha_3)$. If $\alpha_i = 0$ or 2, then it follows from (2.21) that $\partial^{\alpha} b_i = 0$ on Γ_i . If $\alpha_i = 1$, then applying Lemma 2.14 and change of variable $v \mapsto R_x v$, we can obtain from (3.8) that $\partial^{\alpha} \gamma_{1j} = 0$, on Γ_i for $j = 1, 2, 3$. Similarly, for $j \neq i$, if $\alpha_j = 1$, then $\partial^{\alpha} b_i = 0$ on Γ_j . If $\alpha_j = 0$ or 2, then it follows from (3.7) that $\partial^{\alpha} \gamma_{1ij} = 0$, on Γ_j for $j \neq i$.

Next we calculate the term $\partial_t \partial^{\alpha} b_i$. Applying ∂^{α} to (3.12)₂ yields

$$\partial_t \partial^{\alpha} b_i = -\nabla_{x_i} \partial^{\alpha} a_{\pm} - 2\nabla_{x_i} \partial^{\alpha} c \mp \nabla_{x_i} \partial^{\alpha} \phi - (v \cdot \nabla_x \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f + \partial^{\alpha} N_{\pm}(f), v_i)_{L_v^2}.$$

Then by Cauchy-Schwarz inequality, we have

$$\begin{aligned} - \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial_{x_j} \partial^{\alpha} f, \xi_{ij})_{L_v^2} \cdot \partial_t \partial^{\alpha} b_i dx &\leq \varepsilon (\|\nabla_x \partial^{\alpha} a_{\pm}\|_{L_x^2}^2 + \|\nabla_x \partial^{\alpha} c\|_{L_x^2}^2 + \|\nabla_x \partial^{\alpha} \phi\|_{L_x^2}^2) \\ &\quad + \varepsilon \|\partial^{\alpha} N_{\parallel}\|_{L_x^2}^2 + \frac{C}{\varepsilon} \|\nabla_x \partial^{\alpha} \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{L_x^2 L_{10}^2}^2, \end{aligned}$$

and (3.19) becomes

$$\begin{aligned} \|\nabla_x \partial^{\alpha} b\|^2 &\leq - \sum_{i,j} \frac{d}{dt} \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_{ij})_{L_v^2} \cdot \partial_{x_j} \partial^{\alpha} b_i dx + \varepsilon \|\nabla_x \partial^{\alpha} \phi\|_{L_x^2}^2 \\ &\quad + \varepsilon \|\nabla_x \partial^{\alpha} a_{\pm}\|_{L_x^2}^2 + \varepsilon \|\nabla_x \partial^{\alpha} c\|_{L_x^2}^2 + \frac{C}{\varepsilon} (\|\partial^{\alpha} \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{H_x^1 L_{10}^2}^2 + \|\partial^{\alpha} N_{\parallel}\|_{L_x^2}^2), \end{aligned} \quad (3.20)$$

where we choose ε small enough.

Step 3: Computation of $\nabla_x \partial^{\alpha} a$. Taking the addition and difference of (3.11) over \pm , it yields that

$$\nabla_x (a_+ + a_-) = -2\partial_t b + \sum_{\pm} (-(\partial_t + v \cdot \nabla_x) \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f - L_{\pm} \{\mathbf{I} - \mathbf{P}\} f + N_{\pm}(f), \xi_a)_{L_v^2}, \quad (3.21)$$

and

$$\nabla_x (a_+ - a_-) = -2\nabla_x \phi + \sum_{\pm} \pm (-(\partial_t + v \cdot \nabla_x) \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f - L_{\pm} \{\mathbf{I} - \mathbf{P}\} f + N_{\pm}(f), \xi_a)_{L_v^2}. \quad (3.22)$$

Applying ∂^{α} derivative on both side of (3.21) and taking inner product with $\nabla_x \partial^{\alpha} (a_+ + a_-)$ over Ω , we deduce that

$$\begin{aligned} \|\nabla_x \partial^{\alpha} (a_+ + a_-)\|_{L_x^2}^2 &= - \int_{\Omega} \partial_t (\partial^{\alpha} b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2}) \cdot \nabla_x \partial^{\alpha} (a_+ + a_-) dx \\ &\quad - \int_{\Omega} (v \cdot \nabla_x \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f + L_{\pm} \{\mathbf{I} - \mathbf{P}\} \partial^{\alpha} f + \partial^{\alpha} N_{\pm}(f), \xi_a)_{L_v^2} \cdot \nabla_x \partial^{\alpha} (a_+ + a_-) dx. \end{aligned} \quad (3.23)$$

By integration by parts about ∂_t and ∇_x , we have

$$\begin{aligned} &- \int_{\Omega} \partial_t (\partial^{\alpha} b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2}) \cdot \nabla_x \partial^{\alpha} (a_+ + a_-) dx \\ &= - \frac{d}{dt} \int_{\Omega} (\partial^{\alpha} b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2}) \cdot \nabla_x \partial^{\alpha} (a_+ + a_-) dx \\ &\quad - \int_{\Omega} \nabla_x \cdot (\partial^{\alpha} b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2}) \partial_t \partial^{\alpha} (a_+ + a_-) dx. \end{aligned} \quad (3.24)$$

To apply the integration by parts for the union of cubes case, we write the following boundary values by using (2.21). If $\alpha_i = 1$, then $\partial^{\alpha} (a_+ + a_-) = 0$ on Γ_i . If $\alpha_i = 0, 2$, then $\partial^{\alpha} b_i = 0$ on Γ_i and by change of variable $v \mapsto R_x v$ and the definition (3.14) of ξ_a , we have

$$(\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2} = (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f(R_x v), \xi_a(R_x v))_{L_v^2} = -(\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^{\alpha} f, \xi_a)_{L_v^2} = 0.$$

The time derivative and tangent derivatives don't affect the zero boundary values. This completes the integration by parts in (3.24). In view of (3.24) and (3.23), applying (3.12)₁, (3.15) and Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|\nabla_x \partial^\alpha (a_+ + a_-)\|^2 + \frac{d}{dt} \sum_{\pm} \int_{\Omega} \left(\partial^\alpha b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_a)_{L_v^2} \right) \cdot \nabla_x \partial^\alpha (a_+ + a_-) dx \\ \lesssim \|\partial^\alpha \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{H_x^1 L_{10}^2} + \|\partial^\alpha N_{\parallel}\|_{L_x^2}^2 + \|\nabla_x \partial^\alpha b\|_{L_x^2}^2. \end{aligned} \quad (3.25)$$

Similarly, noticing that $(\nabla_x \partial^\alpha \phi, \nabla_x \partial^\alpha (a_+ - a_-))_{L_x^2} = -(\Delta_x \partial^\alpha \phi, \partial^\alpha (a_+ - a_-))_{L_x^2} = \|\Delta_x \partial^\alpha \phi\|_{L_x^2}^2$, we apply ∂^α to (3.22) and take inner product with $\nabla_x \partial^\alpha (a_+ - a_-)$ over Ω to deduce that

$$\begin{aligned} \|\nabla_x \partial^\alpha (a_+ - a_-)\|_{L_x^2}^2 + 2\|\Delta_x \partial^\alpha \phi\|_{L_x^2}^2 + \frac{d}{dt} \int_{\Omega} \sum_{\pm} \mp (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_a)_{L_v^2} \cdot \nabla_x \partial^\alpha (a_+ - a_-) dx \\ \lesssim \|\partial^\alpha \{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} f\|_{H_x^1 L_{10}^2} + \|\partial^\alpha N_{\parallel}\|_{L_x^2}^2 + \|\nabla_x \partial^\alpha b\|_{L_x^2}^2. \end{aligned} \quad (3.26)$$

Notice that $|a_+ + a_-|^2 + |a_+ - a_-|^2 = 2|a_+|^2 + 2|a_-|^2$. For the torus case, by Sobolev inequality, we have

$$\|\nabla_x \phi\|^2 \lesssim \|\nabla_x^2 \phi\|^2 = \|\Delta_x \phi\|_{L_x^2}^2. \quad (3.27)$$

For the zeroth order, we apply the Poincaré inequality to a_{\pm}, b, c in x and use the conservation law (1.9) to deduce that

$$\begin{aligned} \|a_{\pm}\|_{L_x^2}^2 &\leq C \left(\int_{\Omega} a_{\pm}(t, x) dx \right)^2 + C \|\nabla_x a_{\pm}\|_{L_x^2}^2 \lesssim \|\nabla_x a_{\pm}\|_{L_x^2}^2, \\ \|b\|_{L_x^2}^2 &\leq C \left(\int_{\Omega} b(t, x) dx \right)^2 + C \|\nabla_x b\|_{L_x^2}^2 \lesssim \|\nabla_x b\|_{L_x^2}^2, \\ \|c\|_{L_x^2}^2 &\leq C \left(\int_{\Omega} c(t, x) dx \right)^2 + C \|\nabla_x c\|_{L_x^2}^2 \lesssim \|\nabla_x \phi\|_{L_x^2}^4 + \|\nabla_x c\|_{L_x^2}^2. \end{aligned} \quad (3.28)$$

For the union of cubes case, it follows from (1.13) that $\partial_{x_i} \phi = 0$ on Γ_i . Then by Sobolev embedding (cf. [15, Theorem 6.7-5]), we also have (3.27). The second equality in (3.27) can be obtained by using integration by parts. For the zeroth order, the conservation laws (1.14) give the conservation on mass and energy and hence, the estimates for a_{\pm}, c in (3.28) still hold. For the estimate on b , we have from (2.21) that $b_i = 0$ on Γ_i for $i = 1, 2, 3$. Then by Sobolev embedding (cf. [15, Theorem 6.7-5]), we obtain $\|b\|_{L_x^2} \lesssim \|\nabla_x b\|_{L_x^2}$. In view of these facts, we take linear combination $\lambda \times (3.26) + \lambda \times (3.25) + (3.17) + (3.20)$ and summation over $|\alpha| \leq 1$ and \pm with sufficiently small $\lambda, \varepsilon > 0$ to deduce that

$$\partial_t G(t) + \lambda \sum_{|\alpha| \leq 2} \|\partial^\alpha [a_+, a_-, b, c]\|_{L_x^2}^2 + \lambda \|\nabla_x \phi\|_{H_x^2}^2 \lesssim \|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_{10}^2}^2 + \|N_{\parallel}\|_{H_x^1}^2 + \|\nabla_x \phi\|_{L_x^2}^4,$$

where

$$\begin{aligned} G &= \sum_{\pm} \sum_{|\alpha| \leq 1} \left(\int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_c)_{L_v^2} \cdot \nabla_x \partial^\alpha c dx + \sum_{i,j} \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_{ij})_{L_v^2} \cdot \partial_{x_j} \partial^\alpha b_i dx \right) \\ &+ \lambda \int_{\Omega} \left(\partial^\alpha b + (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_a)_{L_v^2} \right) \cdot \nabla_x \partial^\alpha (a_+ + a_-) dx \\ &\mp \lambda \int_{\Omega} (\{\mathbf{I}_{\pm} - \mathbf{P}_{\pm}\} \partial^\alpha f, \xi_a)_{L_v^2} \cdot \nabla_x \partial^\alpha (a_+ - a_-) dx. \end{aligned}$$

This implies (3.4) and the estimate (3.3) can be directly deduced by Cauchy-Schwarz inequality. \square

3.2. Estimates on the energy. In this subsection, we give the energy estimates on (1.3).

Lemma 3.2. *Let $T > 0, \eta > 0$ and $k \geq k_0$. Suppose $[f, \phi]$ is the solution to equation (1.3) satisfying $\mu + f \geq 0$ and $\mathcal{E}(t) \leq C$. Then there exists functional $\mathcal{E}_1(t)$ satisfying*

$$\mathcal{E}_1(t) \approx \|f\|_{X_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2, \quad (3.29)$$

such that, for any $T, \eta > 0$ and $k \geq k_0$, we have

$$\begin{aligned} \partial_t \mathcal{E}_1(t) + \lambda \eta \|f\|_{Y_k}^2 + \lambda \sum_{|\alpha| \leq 2} \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_D^2}^2 \lesssim \|f\|_{X_{k_0}} \|f\|_{X_k}^2 \\ + C_{\eta, k} \|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_v^2}^2 + \eta \|\nabla_x \phi\|_{H_x^2}^2 + \eta \|\mathbf{P} f\|_{H_x^2 L_v^2}^2 + C(\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t). \end{aligned} \quad (3.30)$$

for some generic constant $\lambda > 0$ and any sufficiently small $\eta > 0$.

Proof. We proceed by considering spatial derivatives and mixed derivatives. Notice that $\|\phi\|_{L_x^\infty} \lesssim \|\nabla_x \phi\|_{H_x^1} \lesssim \mathcal{E}(t) \leq C$, which verify (2.30) and we can apply estimates in Section 2.4.

Step 1. Estimates with mixed derivatives. Recall that $A_{\alpha,\beta}$ is given in (1.16). For any $|\alpha|+|\beta| \leq 2$, we apply ∂_β^α to (1.3), and obtain

$$\begin{aligned} \partial_t \partial_\beta^\alpha f_\pm + \partial_\beta^\alpha (v \cdot \nabla_x f_\pm) \mp \partial_\beta^\alpha (\nabla_x \phi \cdot \nabla_v f_\pm) \pm \partial_\beta^\alpha (\nabla_x \phi \cdot v \mu) \\ = \partial_\beta^\alpha Q(f_\pm + f_\mp, \mu) + \partial_\beta^\alpha Q(\mu + f_\pm, f_\pm) + \partial_\beta^\alpha Q(\mu + f_\mp, f_\pm). \end{aligned} \quad (3.31)$$

Thus, taking inner product of (3.31) with $e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm$ over $\Omega \times \mathbb{R}^3$, one has

$$\begin{aligned} \frac{1}{2} \partial_t \|e^{\frac{\pm A_{\alpha,\beta}\phi}{2\langle v \rangle^2}} \partial_\beta^\alpha f_\pm\|_{L_{x,v}^2}^2 + (A_{\alpha,\beta} \partial_t \phi \langle v \rangle^{-2} \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \\ + (v \cdot \nabla_x \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} + \sum_{|\beta_1|=1} (\partial_{\beta_1} v \cdot \nabla_x \partial_{\beta-\beta_1}^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \\ \mp (\nabla_x \phi \cdot \nabla_v \partial_\beta^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \mp \sum_{\substack{\alpha_1 \leq \alpha \\ |\alpha_1| \geq 1}} (\partial^{\alpha_1} \nabla_x \phi \cdot \nabla_v \partial_{\beta-\alpha_1}^\alpha f_\pm, e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \\ \pm (\partial_\beta^\alpha (\nabla_x \phi \cdot v \mu), e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} = (\partial_\beta^\alpha Q(f_\pm + f_\mp, \mu), e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \\ + (\partial_\beta^\alpha Q(\mu + f_\pm, f_\pm) + \partial_\beta^\alpha Q(\mu + f_\mp, f_\pm), w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2} \\ + (\partial_\beta^\alpha Q(\mu + f_\pm, f_\pm) + \partial_\beta^\alpha Q(\mu + f_\mp, f_\pm), (e^{\frac{\pm A_{\alpha,\beta}\phi}{\langle v \rangle^2}} - 1) w^2(\alpha, \beta) \partial_\beta^\alpha f_\pm)_{L_{x,v}^2}. \end{aligned} \quad (3.32)$$

We denote the second to tenth terms in (3.32) by I_1 to I_9 and estimate them one by one. For I_1 , we make a rough estimate:

$$|I_1| \lesssim \|\partial_t \phi\|_{L_x^\infty} \|f\|_{X_k}^2. \quad (3.33)$$

Notice that $\|\phi\|_{L_x^\infty} \lesssim \|\nabla_x \phi\|_{H_x^1} \leq C$, which verify (2.30) and we can apply estimates in Section 2.4. By (2.32), we deduce that $|I_2 + I_4| \lesssim \|\nabla_x \phi\|_{H_x^2} \|f\|_{Y_k}^2$. Next, after applying (2.33), when $|\beta| \geq 1$, we have

$$|I_3| \lesssim C_\eta \sum_{\substack{|\beta_2|=|\beta|-1 \\ |\alpha_2|=|\alpha|+1}} \|\partial_{\beta_2}^{\alpha_2} f w(\alpha_2, \beta_2)\|_{L_x^2 L_{\gamma/2}^2}^2 + \eta \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}^2}^2,$$

for any $\eta > 0$. By (2.34), we have $|I_5| \lesssim \|\nabla_x \phi\|_{H_x^3} \|f\|_{Y_k}^2$. When $|\beta| = 0$, using (2.38), we have

$$\begin{aligned} \sum_{\pm} I_6 &\geq C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - C \|\partial^\alpha \nabla_x \phi\|_{L_x^2} \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_\gamma^2} - C \|\partial^\alpha \nabla_x \phi\|_{L_x^2} \|\nabla_x \psi\|_{H_x^1} \|f\|_{Y_k} \\ &\geq C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - \eta \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - \eta \|w(\alpha, 0) \partial^\alpha f\|_{L_x^2 L_{\gamma/2}^2}^2 \\ &\quad - C_\eta \|\partial^\alpha f\|_{L_{x,v}^2}^2 - C \|\partial^\alpha \nabla_x \phi\|_{L_x^2} \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}, \end{aligned}$$

for any $\eta > 0$. When $|\beta| \geq 1$, we have from (2.37) that

$$\sum_{\pm} |I_6| \leq C_k \|\nabla_x \phi\|_{H_x^2} \|f\|_{H_x^2 L_v^2}. \quad (3.34)$$

For the Boltzmann case and for convenience, we define

$$\|f\|_{Y_k} := \sum_{|\alpha|+|\beta| \leq 2} C_{|\alpha|,|\beta|} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2,*}^2}^2. \quad (3.35)$$

For I_7 , since $\theta \in [0, \frac{\pi}{2}]$, we have $\sin^2 \frac{\theta}{2} \leq \frac{1}{2}$ and hence if $k \geq 14$ we have

$$\|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L_\theta^1} \|f\|_{Y_k}^2 \leq \frac{1}{16} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|f\|_{Y_k}^2. \quad (3.36)$$

For Boltzmann case, applying Lemma 2.20, we have for any $\eta > 0$ that

$$\begin{aligned} \sum_{\pm} |I_7| &\leq \|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2,*}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2,*}^2} \\ &\quad + \eta \|f\|_{Y_k}^2 + C_\eta \|f\|_{H_{x,v}^2}^2 + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 L_{\gamma/2}^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}^2} \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{16} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}}^2 + \eta \|f\|_{Y_k}^2 + C_\eta \|f\|_{H_{x,v}^2}^2 \\ &\quad + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 L_{\gamma/2}} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2}}. \end{aligned}$$

For the Landau case, it follows from Lemma 2.20 that

$$\sum_{\pm} |I_7| \leq C_k \sum_{|\alpha|+|\beta| \leq 2} \|\partial_\beta^\alpha f\|_{L_x^2 L_v^2}^2 \leq \eta \|f\|_{Y_k}^2 + C_\eta \|f\|_{H_{x,v}^2}^2,$$

For I_8 , for Boltzmann case, we deduce from (2.44) that

$$\begin{aligned} \sum_{\pm} I_8 &\leq -\frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_{\gamma/2, *}}^2 - \gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 H_{\gamma/2}^s}^2 + C_k \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|f\|_{Y_k}^2 \\ &\quad + C_k \|\partial_\beta^\alpha f\|_{L_{x,v}^2}^2 + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{H_{\gamma/2}^s} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{H_{\gamma/2}^s}, \end{aligned}$$

For Landau case, note that and from (2.54) and (2.55),

$$\begin{aligned} \sum_{\pm} I_8 &\leq -\gamma_1 \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_D^2}^2 + C_k \|\partial_\beta^\alpha f\|_{L_x^2 L_v^2}^2 \\ &\quad + C_k \sum_{\beta_1 < \beta} \|\partial_{\beta_1}^\alpha f w(\alpha, \beta_1)\|_{L_x^2 L_D^2} \|\partial_\beta^\alpha f w(\alpha, \beta)\|_{L_x^2 L_D^2} + C_k \|f \langle v \rangle^7\|_{H_{x,v}^2} \|f\|_{Y_k}^2, \end{aligned}$$

for Landau case, with some constants $\gamma_1, C_k > 0$. For I_9 , for Boltzmann case, we have from (2.45) that

$$|I_9| \leq C_k (\|\langle v \rangle^{14} f\|_{H_{x,v}^2} + C) \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}^2 + \|\langle v \rangle^{14} f\|_{H_{x,v}^2} \|\nabla_x \phi\|_{H_x^1} (\|f\|_{Y_k} + C) \|f\|_{Y_k},$$

and for Landau case, from (2.56) we deduce

$$|I_9| \leq C_k (\|\langle v \rangle^7 f\|_{H_{x,v}^2} + C) \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}^2.$$

Combining the above estimates, we take summation $\sum_{|\alpha|+|\beta| \leq 2, \pm} C_{|\alpha|, |\beta|} \times (3.32)$ with $0 < \eta \ll 1$ and constants $C_{|\alpha|, |\beta|}$ satisfying (1.18) to deduce that for the Boltzmann case:

$$\begin{aligned} &\frac{1}{2} \partial_t \|f\|_{X_k}^2 + \partial_t \sum_{|\alpha| \leq 2} C_\alpha C_{|\alpha|, 0} \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 + \frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|f\|_{Y_k}^2 + \gamma_1 \|f\|_{Y_k}^2 \\ &\leq \frac{1}{16} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1} \|f\|_{Y_k}^2 + \eta \|f\|_{Y_k}^2 + C_{\eta, k} \|f\|_{H_{x,v}^2}^2 \\ &\quad + C \|\partial_t \phi\|_{L_x^\infty} \|f\|_{X_k}^2 + C_k \left(\|\nabla_x \phi\|_{H_x^3} \|f\|_{Y_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2 + \|f\|_{X_{14}} \|f\|_{Y_k}^2 \right) \\ &\quad + (\|f\|_{X_{14}} + C) \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}^2 + \|f\|_{X_{14}} \|\nabla_x \phi\|_{H_x^1} (\|f\|_{Y_k} + C) \|f\|_{Y_k}, \end{aligned} \quad (3.37)$$

and for the Landau case:

$$\begin{aligned} &\frac{1}{2} \partial_t \|f\|_{X_k}^2 + \partial_t \sum_{|\alpha| \leq 2} C_\alpha C_{|\alpha|, 0} \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 + \gamma_1 \|f\|_{Y_k}^2 \\ &\leq \eta \|f\|_{Y_k}^2 + C_{\eta, k} \|f\|_{H_{x,v}^2}^2 + C \|\partial_t \phi\|_{L_x^\infty} \|f\|_{X_k}^2 + C_k \left(\|\nabla_x \phi\|_{H_x^3} \|f\|_{Y_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2 \right. \\ &\quad \left. + \|f\|_{X_7} \|f\|_{Y_k}^2 + (\|f\|_{X_7} + C) \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}^2 + \|f\|_{X_7} \|\nabla_x \phi\|_{H_x^1} (\|f\|_{Y_k} + C) \|f\|_{Y_k} \right), \end{aligned} \quad (3.38)$$

for some generic constant γ_1 independent of k , where we let $k \geq 14$. Next notice that

$$\|\nabla_x^3 \phi\|_{L_x^2}^2 \lesssim \|\nabla_x \Delta_x \phi\|_{L_x^2}^2 \lesssim \|\nabla_x f\|_{L_x^2 L_4^2}^2 \lesssim \|f\|_{X_{k_0}}^2. \quad (3.39)$$

For union of cubes case, we apply embedding [15, Theorem 6.7-5] with boundary values $\partial_{x_i} \phi = 0$ on Γ_i to obtain (3.39). Taking difference of (1.7) over \pm , we have $\partial_t(a_+ - a_-) = -\nabla_x \cdot \int_{\mathbb{R}^3} v(f_+ - f_-) dv$. So

$$\begin{aligned} \|\partial_t \phi\|_{L_x^\infty} &\lesssim \|\nabla_x \partial_t \phi\|_{L_x^2}^{1/2} \|\nabla_x \partial_t \phi\|_{H_x^1}^{1/2} \lesssim \|\nabla_x^2 \partial_t \phi\|_{L_x^2} \lesssim \|\Delta_x \partial_t \phi\|_{L_x^2} \\ &\lesssim \|\partial_t(a_+ - a_-)\|_{L_x^2} \lesssim \|\nabla_x f\|_{L_x^2 L_3^2} \lesssim \|f\|_{X_{k_0}}. \end{aligned} \quad (3.40)$$

For the Boltzmann case, applying (3.36), (3.39), (3.40) and choosing $\eta < \frac{\gamma_1}{4}$, we have

$$\partial_t \left(\frac{1}{2} \|f\|_{X_k}^2 + \sum_{|\alpha| \leq 2} C_\alpha C_{|\alpha|, 0} \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 \right) + \frac{\gamma_1}{2} \|f\|_{Y_k}^2$$

$$\begin{aligned} &\leq C_k \|f\|_{H_{x,v}^2}^2 + C \|f\|_{X_{k_0}} \|f\|_{X_k}^2 + C_k \left(\|\nabla_x \phi\|_{H_x^3} \|f\|_{Y_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2 + \|f\|_{X_{k_0}} \|f\|_{Y_k}^2 \right. \\ &\quad \left. + (\|f\|_{X_{k_0}} + C) \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k}^2 + \|f\|_{X_{k_0}} \|\nabla_x \phi\|_{H_x^1} (\|f\|_{Y_k} + C) \|f\|_{Y_k} \right), \end{aligned} \quad (3.41)$$

for some generic constant γ_1 independent of k . Here C_α and $C_{|\alpha|,0}$ are given in (2.38) and (1.18) respectively. If $|\beta| \geq 1$, by interpolation we have

$$C_k \|\partial_\beta^\alpha f\|_{L_{x,v}^2}^2 \leq C \|\partial^\alpha f\|_{L_x^2 H_v^{|\beta|}}^2 \leq \frac{\gamma_1}{24} \|\partial^\alpha f\|_{L_x^2 H_v^{|\beta|+s}}^2 + C_k \|\partial^\alpha f\|_{L_{x,v}^2}^2 \leq \frac{\gamma_1}{24} \|f\|_{Y_k}^2 + C_k \|\partial^\alpha f\|_{L_{x,v}^2}^2. \quad (3.42)$$

Also, by Sobolev embedding or Poincaré's inequality,

$$\|\nabla_x \phi\|_{H_x^2} \lesssim \|\nabla_x^2 \phi\|_{H_x^1} = \|\Delta_x \phi\|_{H_x^1} \lesssim \|f\|_{H_x^1 L_x^2} \lesssim \eta \|f\|_{Y_k} + C_\eta \|f\|_{H_x^1 L_x^2}.$$

In view of the above two estimates, choosing $\eta > 0$ small enough, (3.41) implies that

$$\begin{aligned} &\partial_t \left(\frac{1}{2} \|f\|_{X_k}^2 + \sum_{|\alpha| \leq 2} C_\alpha C_{|\alpha|,0} \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 \right) + \frac{\gamma_1}{4} \|f\|_{Y_k}^2 \\ &\quad \lesssim \|f\|_{H_x^2 L_x^2}^2 + C \|f\|_{X_{k_0}} \|f\|_{X_k}^2 + \|\nabla_x \phi\|_{H_x^2}^2 + (\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t). \end{aligned} \quad (3.43)$$

For the Landau case, one can apply a similar calculation as (3.38) instead of (3.37) to derive (3.43) by using the same technique and we omit the details for brevity.

Step 2. Estimates with spatial derivatives. For any $|\alpha| \leq 2$, we apply ∂^α to (1.3), take inner product with $e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm$ over $\Omega \times \mathbb{R}^3$ to deduce that

$$\begin{aligned} &\frac{1}{2} \partial_t \|e^{\frac{\pm A_{\alpha,0}\phi}{2\langle v \rangle^2}} \partial^\alpha f_\pm\|_{L_{x,v}^2}^2 + (A_{\alpha,0} \partial_t \phi \langle v \rangle^{-2} \partial^\alpha f_\pm, e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &\quad + (v \cdot \nabla_x \partial^\alpha f_\pm, e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \mp (\nabla_x \phi \cdot \nabla_v \partial^\alpha f_\pm, e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &\quad \mp \sum_{\substack{|\alpha| \leq 2 \\ |\alpha_1| \geq 1}} (\partial^{\alpha_1} \nabla_x \phi \cdot \nabla_v \partial^{\alpha-\alpha_1} f_\pm, e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \pm (\partial^\alpha (\nabla_x \phi \cdot v \mu), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &\quad = (\partial^\alpha L_\pm f + \partial^\alpha Q(f_\pm + f_\mp, f_\pm), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2}, \end{aligned} \quad (3.44)$$

where L_\pm is given in (3.2). We denote the second to seventh terms in (3.44) by J_1 to J_6 and estimate them one by one. Similar to the calculation of I_1 to I_6 from (3.33) to (3.34), one can obtain the following estimates by using (2.32) and (2.34):

$$\begin{aligned} |J_1 + J_2 + J_3 + J_4| &\lesssim (\|\partial_t \phi\|_{L_x^\infty} + \|\nabla_x \phi\|_{H_x^3}) \|f\|_{Y_k}^2 \\ &\lesssim C \|\partial_t \phi\|_{L_x^\infty} \|f\|_{X_k}^2 + \|\nabla_x \phi\|_{H_x^2} \|f\|_{Y_k}^2 \lesssim \|f\|_{X_{k_0}} \|f\|_{X_k}^2 + \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t), \end{aligned}$$

where we apply (3.39) and (3.40) for the terms $\|\nabla_x^3 \phi\|_{L_x^2}$ and $\|\partial_t \phi\|_{L_x^\infty}$ respectively. For J_5 , using (2.38), we have

$$\begin{aligned} J_5 &\geq C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - C \|\partial^\alpha \nabla_x \phi\|_{L_x^2} \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2} - C \|\partial^\alpha \nabla_x \phi\|_{L_x^2} \|\nabla_x \phi\|_{H_x^1} \|f\|_{Y_k} \\ &\geq C_\alpha \partial_t \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - \eta \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 - \eta \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2}^2 \\ &\quad - C_\eta \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2}^2 - C \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t), \end{aligned}$$

for any $\eta > 0$. Here C_α is given in (2.38). For the term J_6 , we need some different estimates. Noticing $L_\pm(\partial^\alpha \mathbf{P}_\pm f) = 0$, we split J_6 into

$$\begin{aligned} J_6 &= (\partial^\alpha Q(2\mu + f_\pm + f_\mp, \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &\quad + (\partial^\alpha Q(f_\pm + f_\mp, \mathbf{P}_\pm f), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &\quad + (\partial^\alpha Q(\{\mathbf{I}_\pm - \mathbf{P}_\pm\} f + \{\mathbf{I}_\mp - \mathbf{P}_\mp\} f, \mu), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0) \partial^\alpha f_\pm)_{L_{x,v}^2} \\ &=: J_7 + J_8 + J_9. \end{aligned}$$

We further split J_7 as

$$J_7 = (\partial^\alpha Q(2\mu + f_\pm + f_\mp, \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f), w^2(\alpha, 0) \partial^\alpha \{\mathbf{I}_\pm - \mathbf{P}_\pm\} f)_{L_{x,v}^2}$$

$$\begin{aligned}
& + (\partial^\alpha Q(2\mu + f_\pm + f_\mp, \{\mathbf{I}_\pm - \mathbf{P}_\pm\}f), w^2(\alpha, 0)\partial^\alpha \mathbf{P}_\pm f)_{L^2_{x,v}} \\
& + (\partial^\alpha Q(2\mu + f_\pm + f_\mp, \{\mathbf{I}_\pm - \mathbf{P}_\pm\}f), (e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} - 1)w^2(\alpha, 0)\partial^\alpha f_\pm)_{L^2_{x,v}} \\
& =: J_{7,1} + J_{7,2} + J_{7,3}.
\end{aligned}$$

For Boltzmann case, we deduce from (2.44) that $J_{7,1}$ satisfies

$$\begin{aligned}
\sum_{\pm} J_{7,1} & \leq -\frac{1}{8} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L^1_\theta} \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2,*}}^2 \\
& \quad - \frac{\gamma_1}{2} \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x H^s_{\gamma/2}}^2 + C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t) + C_k \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_{x,v}}^2.
\end{aligned}$$

For $J_{7,2}$ and $J_{7,3}$, for the Boltzmann case, we deduce from Lemma 2.4 that

$$\begin{aligned}
|J_{7,2}| & \lesssim (C + \|\langle v \rangle^{14} f\|_{H^2_x L^2_v}) \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v} \|w^2(\alpha, 0)\mathbf{P}f\|_{H^2_x H^{2s}_{\gamma+2s}} \\
& \lesssim \eta \|\mathbf{P}f\|_{H^2_x L^2_v}^2 + C_\eta \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v}^2 + C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t).
\end{aligned}$$

It follows from (2.45) that

$$\begin{aligned}
|J_{7,3}| & \lesssim \|\nabla_x \phi\|_{H^1_x} \left((C + \|\langle v \rangle^{14} f\|_{H^2_x L^2_v}) \|\{\mathbf{I} - \mathbf{P}\}f\|_{Y_k} \|\mathbf{P}f\|_{Y_k} + \|\langle v \rangle^{14} \{\mathbf{I} - \mathbf{P}\}f\|_{H^2_{x,v}} (C + \|f\|_{Y_k}) \|\mathbf{P}f\|_{Y_k} \right) \\
& \lesssim (\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t).
\end{aligned}$$

For the Landau case, it follows from (2.54) and (2.55) that

$$\sum_{\pm} J_{7,1} \leq -\frac{\gamma_1}{2} \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_D}^2 + C_k \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_D}^2 + C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t),$$

with some constants $\gamma_1, C_k > 0$. Applying Lemma 2.13 and (2.56), we have

$$|J_{7,2}| \lesssim (C + \|f\|_{L^2_5}) \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v} \|\mathbf{P}f\|_{H^2_x H^2_5} \lesssim \eta \|\mathbf{P}f\|_{H^2_x L^2_v}^2 + C_\eta \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v}^2 + C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t),$$

and

$$|J_{7,3}| \lesssim (C + \|\langle v \rangle^7 f\|_{H^2_{x,v}}) \|\nabla_x \phi\|_{H^1_x} \|\{\mathbf{I} - \mathbf{P}\}f\|_{Y_k} \|f\|_{Y_k} \lesssim (\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t).$$

For J_8 , we deduce from (2.46) that for the Boltzmann case:

$$|J_8| \lesssim \|\langle v \rangle^{14} f\|_{H^2_{x,v}} \|\langle v \rangle^{2s} \mathbf{P}f\|_{Y_k} \|f\|_{Y_k} + \|\langle v \rangle^{14} \mathbf{P}f\|_{H^2_{x,v}} \|f\|_{Y_k} \|f\|_{Y_k} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t),$$

and from (2.57) that for the Landau case:

$$|J_8| \lesssim \|f\langle v \rangle^7\|_{H^2_{x,v}} \|\langle v \rangle \mathbf{P}f\|_{Y_k} \|f\|_{Y_k} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t).$$

Here we let $k \geq k_0$. For J_9 , we split it as

$$\begin{aligned}
J_9 & = (\partial^\alpha Q(\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f + \{\mathbf{I}_\mp - \mathbf{P}_\mp\}f, \mu), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0)\partial^\alpha \{\mathbf{I}_\pm - \mathbf{P}_\pm\}f f_\pm)_{L^2_{x,v}} \\
& \quad + (\partial^\alpha Q(\{\mathbf{I}_\pm - \mathbf{P}_\pm\}f + \{\mathbf{I}_\mp - \mathbf{P}_\mp\}f, \mu), e^{\frac{\pm A_{\alpha,0}\phi}{\langle v \rangle^2}} w^2(\alpha, 0)\partial^\alpha \mathbf{P}_\pm f_\pm)_{L^2_{x,v}} \\
& =: J_{9,1} + J_{9,2}.
\end{aligned}$$

Then for Boltzmann case, applying Lemma 2.20, we have for any $\eta > 0$ that

$$\begin{aligned}
|J_{9,1}| & \leq \|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L^1_\theta} \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2,*}} \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f w(\alpha, 0)\|_{L^2_x L^2_{\gamma/2,*}} \\
& \quad + C_k \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2-1/2}} \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f w(\alpha, 0)\|_{L^2_x L^2_{\gamma/2-1/2}} \\
& \leq \frac{1}{16} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L^1_\theta} \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2,*}}^2 \\
& \quad + \eta \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2}}^2 + C_{k,\eta} \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v}^2,
\end{aligned}$$

For the term $J_{9,2}$, by Lemma 2.4 we have

$$\begin{aligned}
|J_{9,2}| & \leq C \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_\gamma} \|\mu\|_{H^{2s}} \|\partial^\alpha \mathbf{P}f w(\alpha, 0)\|_{L^2_x L^2_{\gamma+2s}} \\
& \leq \eta \|\partial^\alpha \mathbf{P}f\|_{L^2_{x,v}}^2 + \eta \|w(\alpha, 0)\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_{\gamma/2}}^2 + C_{k,\eta} \|\{\mathbf{I} - \mathbf{P}\}f\|_{H^2_x L^2_v}^2,
\end{aligned}$$

For the Landau case, we have from Lemma 2.20 that

$$|J_9| \leq C_k \|\partial^\alpha \{\mathbf{I} - \mathbf{P}\}f\|_{L^2_x L^2_5} \|\partial^\alpha f\|_{L^2_x L^2_5}$$

$$\leq \eta \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2}^2 + \eta \|\partial^\alpha \mathbf{P} f\|_{L_{x,v}^2}^2 + C_\eta \|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_v^2}^2.$$

Taking summation $\sum_{\pm} \sum_{|\alpha| \leq 2}$ (3.44), combining the above estimates for J_1 to J_9 , applying (3.36) and (3.42), we deduce that for small $\eta > 0$,

$$\begin{aligned} & \partial_t \sum_{|\alpha| \leq 2} \left(\frac{1}{2} \|e^{\frac{\pm A_{\alpha,0} \phi}{2(v)^2}} \partial^\alpha f_{\pm}\|_{L_{x,v}^2}^2 + C_\alpha \|\nabla_x \phi\|_{H_x^2}^2 \right) + \frac{\gamma_1}{3} \sum_{|\alpha| \leq 2} \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 H_{\gamma/2}^s}^2 \\ & \lesssim \|f\|_{X_{k_0}} \|f\|_{X_k}^2 + C_{\eta,k} \|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_v^2}^2 + \eta \|\nabla_x \phi\|_{H_x^2}^2 + \eta \|\mathbf{P} f\|_{H_x^2 L_v^2}^2 + C(\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t). \end{aligned} \quad (3.45)$$

Here, for the Boltzmann case, we use the fact that $\|b(\cos \theta) \sin^{k-2} \frac{\theta}{2}\|_{L_\theta^1} \leq \frac{1}{16} \|b(\cos \theta) \sin^2 \frac{\theta}{2}\|_{L_\theta^1}$, which follows from $k \geq k_0$. Taking linear combination $\eta \times$ (3.43) + (3.45) and applying (3.40), we obtain

$$\begin{aligned} & \partial_t \mathcal{E}_1(t) + \lambda \eta \|f\|_{Y_k}^2 + \frac{\gamma_1}{4} \sum_{|\alpha| \leq 2} \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2}^2 \\ & \lesssim \|f\|_{X_{k_0}} \|f\|_{X_k}^2 + \eta \|f\|_{H_x^2 L_v^2}^2 + C_{\eta,k} \|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_v^2}^2 + \eta \|\nabla_x \phi\|_{H_x^2}^2 + \eta \|\mathbf{P} f\|_{H_x^2 L_v^2}^2 + C(\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t). \end{aligned} \quad (3.46)$$

where $\mathcal{E}_1(t)$ is given by

$$\mathcal{E}_1(t) = \frac{\eta}{2} \|f\|_{X_k}^2 + \eta \sum_{|\alpha| \leq 2} C_\alpha C_{|\alpha|,0} \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2 + \sum_{\pm} \sum_{|\alpha| \leq 2} \left(\frac{1}{2} \|e^{\frac{\pm A_{\alpha,0} \phi}{2(v)^2}} \partial^\alpha f_{\pm}\|_{L_{x,v}^2}^2 + C_\alpha \|\partial_t \nabla_x \phi\|_{H_x^2}^2 \right). \quad (3.47)$$

It's direct to verify that $\mathcal{E}_1(t)$ satisfies (3.29). For the second right-hand term of (3.46), we split it as

$$\eta \|f\|_{H_x^2 L_v^2}^2 \lesssim \eta \|\mathbf{P} f\|_{H_x^2 L_v^2}^2 + \eta \sum_{|\alpha| \leq 2} \|w(\alpha, 0) \partial^\alpha \{\mathbf{I} - \mathbf{P}\} f\|_{L_x^2 L_v^2}^2. \quad (3.48)$$

Choosing η sufficiently small, the second right hand term of (3.48) can be absorbed by the left hand side of (3.46) and we obtain (3.30). This completes the proof of Lemma 3.2. \square

3.3. Recover the energy from semigroup method. According to Lemma 3.2, we only need to deal with term $\|\{\mathbf{I} - \mathbf{P}\} f\|_{H_x^2 L_v^2}$ without velocity derivative on the right hand side of (3.30). In order to eliminate this term, we define the semigroup generated by \mathcal{L} given in (1.4) to be $S_{\mathcal{L}}(t)$. Then we first give some estimate on $S_{\mathcal{L}}(t)$, which is the solution operator to the equation

$$\partial_t f + v \cdot \nabla_x f \pm \nabla_x \phi \cdot v \mu = L_{\pm} f, \quad f(0) = f_0, \quad -\Delta_x \phi = \int_{\mathbb{R}^3} (f_+ - f_-) dv, \quad \int_{\Omega} \phi(t, x) dx = 0. \quad (3.49)$$

If the domain Ω is union of cubes given by (1.10), then we further assume

$$\partial_n \phi = 0 \quad \text{on } \partial\Omega. \quad (3.50)$$

To obtain the estimate of (3.49), we denote $-\Delta_x^{-1} \int_{\mathbb{R}^3} (\cdot)_+ - (\cdot)_- dv$ to be the solution operator to the second equation of (3.49). Then we define linear operators $A = [A_+, A_-]$, $B = [B_+, B_-]$ and K_1, K_2 as

$$\begin{aligned} A_{\pm} &= -v \cdot \nabla_x + L_{\pm} - M \chi_R, \quad B_{\pm} = -v \cdot \nabla_x + L_{\pm}, \quad K_1 = M \chi_R, \\ K_2 &= \pm \mu v \cdot \nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} ((\cdot)_+ - (\cdot)_-) dv, \quad \mathcal{L} = -v \cdot \nabla_x \pm \mu v \cdot \nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} ((\cdot)_+ - (\cdot)_-) dv + L_{\pm}, \end{aligned}$$

where $M > 0$ is a large constant and $\chi \in D(\mathbb{R})$ is the truncation function satisfying $1_{[-1,1]} \leq \chi \leq 1_{[-2,2]}$ and we denote $\chi_R(\cdot) := \chi(\cdot/R)$ for $R > 0$. For the case of the union of cubes, we consider the domain of these operators, i.e. A, B, \mathcal{L} , with restriction of specular-reflection boundary condition for f and Neumann boundary condition for ϕ :

$$f(R_x v) = f(v) \text{ on } \gamma_-, \quad \partial_n \Delta_x^{-1} \int_{\mathbb{R}^3} (f_+ - f_-) dv = 0 \text{ on } \partial\Omega.$$

Then we have the following lemma.

Lemma 3.3. Consider both Boltzmann and Landau cases. For $k \geq k_0$, there exists $\varepsilon_0 > 0$ such that if

$$\|f\|_{H_x^2 L_{k_0+4}^2} \leq \varepsilon_0, \quad (3.51)$$

then we have

$$\|S_{\mathcal{L}}(t)f\|_{H_x^2 L_k^2} \lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2 L_{k+4}^2}, \quad (3.52)$$

and (suppose $s = 1$ for Landau case)

$$\left(\int_0^\infty \|\langle v \rangle^k S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \lesssim \|f\|_{H_x^2 H_{k-\gamma/2}^{-s}}. \quad (3.53)$$

Proof. We will prove (3.52) and (3.53) in two steps. We only prove the Boltzmann case and the Landau case can be proved similarly.

Step 1. By Duhamel's principle, we have

$$S_B(t) = S_A(t) + \int_0^t S_B(t-s)K_1 S_A(s) ds, \quad (3.54)$$

and

$$S_{\mathcal{L}}(t) = S_B(t) + \int_0^t S_{\mathcal{L}}(t-s)K_2 S_B(s) ds. \quad (3.55)$$

Using Lemma 2.6 and Theorem 2.8 with $g = 0$ for the Boltzmann case and Lemma 2.11 for the Landau case, we deduce that for $k \geq k_0$,

$$(-Af, \langle v \rangle^{2k} f)_{L_{x,v}^2} \geq \frac{\gamma_1}{2} \|\langle v \rangle^k f\|_{L_x^2 L_v^2}^2 \gtrsim \|\langle v \rangle^{k+\frac{\gamma}{2}} f\|_{L_{x,v}^2}^2, \quad (3.56)$$

here $(v \cdot \nabla_x \partial^\alpha f, \partial^\alpha f)_{L_{x,v}^2} = 0$ by using change of variable $v \mapsto R_x v$ for the case of union of cubes. Moreover, we use Lemma 2.14 to obtain

$$(v \cdot \nabla_x f, \langle v \rangle^{2k} f)_{L_{x,v}^2} = 0. \quad (3.57)$$

For the case of the union of cubes, and by using change of variable $v \mapsto R_x v$. For the case of torus, (3.57) also holds. Then A generates a semigroup on L_k^2 such that

$$\|S_A(t)f\|_{L_x^2 L_k^2} \leq \|f\|_{L_x^2 L_k^2}. \quad (3.58)$$

Also, by definition of semigroup, we have the equation

$$\partial_t S_A(t)f - A S_A(t)f = 0. \quad (3.59)$$

Taking inner product of (3.59) with $\langle v \rangle^{2k} S_A(t)f$ over $\Omega \times \mathbb{R}^3$, we have from (3.56) that

$$\frac{1}{2} \partial_t \|\langle v \rangle^k S_A(t)f\|_{L_{x,v}^2}^2 + \lambda \|\langle v \rangle^k S_A(t)f\|_{L_x^2 L_v^2}^2 \leq 0. \quad (3.60)$$

For the hard potential case, i.e. $\gamma \geq 0$, we have

$$\|\langle v \rangle^k S_A(t)f\|_{L_x^2 L_v^2} \gtrsim \|\langle v \rangle^k S_A(t)f\|_{L_x^2 L_v^2},$$

and hence,

$$\|S_A(t)f\|_{L_x^2 L_k^2} \leq e^{-\lambda t} \|f\|_{L_x^2 L_k^2}, \quad (3.61)$$

for some $\lambda > 0$. For the soft potential case and $k \geq k_1 \geq k_0$, it follows from (3.60) that

$$\partial_t \|S_A(t)f\|_{L_x^2 L_{k_1}^2}^2 \lesssim -\|S_A(t)f\|_{L_x^2 L_{k_1+\frac{\gamma}{2}}^2}^2 \lesssim -\lambda \langle R \rangle^\gamma \|S_A(t)f\|_{L_x^2 L_{k_1}^2}^2 + \langle R \rangle^{2(k-k_1)+\gamma} \|S_A(t)f\|_{L_x^2 L_k^2}^2,$$

for some $\lambda > 0$. Solving this ODE and using (3.58), we have

$$\begin{aligned} \|S_A(t)f\|_{L_x^2 L_{k_1}^2}^2 &\lesssim e^{-\lambda \langle R \rangle^\gamma t} (\|f\|_{L_x^2 L_{k_1}^2}^2 + \langle R \rangle^{2(k-k_1)+\gamma} \int_0^t e^{\lambda \langle R \rangle^\gamma s} ds \|f\|_{L_x^2 L_k^2}^2) \\ &\lesssim e^{-\lambda \langle R \rangle^\gamma t} \|f\|_{L_x^2 L_{k_1}^2}^2 + \langle R \rangle^{2(k-k_1)} \|f\|_{L_x^2 L_k^2}^2. \end{aligned}$$

Choosing $\langle R \rangle = (\langle t \rangle [\log \langle t \rangle]^{-\frac{2(k-k_0)}{\lambda}})^{-\frac{1}{\gamma}}$, we have

$$\|S_A(t)f\|_{L_x^2 L_{k_1}^2} \lesssim \langle t \rangle^{-\frac{k-k_1-1/2}{|\gamma|}} \|f\|_{L_x^2 L_k^2}, \quad (3.62)$$

For $k \geq k_1 + 4$, since $|\gamma| \leq 3$, we know that $\langle t \rangle^{-\frac{k-k_1-1/2}{|\gamma|}} \leq \langle t \rangle^{-\frac{7}{6}}$. Together with (3.61), we deduce that

$$\|S_A(t)f\|_{L_x^2 L_{k_1}^2} \lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{L_x^2 L_k^2}, \quad (3.63)$$

For the semigroups $S_B(t)$ and $S_{\mathcal{L}}(t)$ in exponential weighted space, we have

$$\partial_t S_B(t)f - B S_B(t)f = 0, \quad \partial_t S_{\mathcal{L}}(t)f - \mathcal{L} S_{\mathcal{L}}(t)f = 0. \quad (3.64)$$

Taking $H_x^2 L_v^2$ inner product with $\mu^{-1} S_B(t)f$ and $\mu^{-1} S_{\mathcal{L}}(t)f$ over $\Omega \times \mathbb{R}^3$ respectively, we have

$$\begin{aligned} \partial_t \|\mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2 + \lambda \|\mu^{-\frac{1}{2}} \{\mathbf{I} - \mathbf{P}\mu^{\frac{1}{2}}\} (S_B(t)f)\|_{H_x^2 H_{\gamma/2}^s}^2 &\leq 0, \\ \partial_t \|\mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 + \partial_t \|\nabla_x \phi\|_{H_x^2}^2 + \lambda \|\mu^{-\frac{1}{2}} \{\mathbf{I} - \mathbf{P}\mu^{\frac{1}{2}}\} (S_{\mathcal{L}}(t)f)\|_{H_x^2 H_{\gamma/2}^s}^2 &\leq 0. \end{aligned} \quad (3.65)$$

where the dissipation rate of L_{\pm} for exponential perturbation can be found in [33, 36] and ϕ is given by

$$\phi = \Delta_x^{-1} \int_{\mathbb{R}^3} (S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_- dv. \quad (3.66)$$

In the second estimate of (3.65), we used the fact that $\partial_t \int_{\mathbb{R}^3} S_{\mathcal{L}}(t)f dv + \int_{\mathbb{R}^3} v \cdot \nabla_x S_{\mathcal{L}}(t)f dv = 0$. Thus

$$\begin{aligned} &\sum_{\pm} (\pm \mu v \cdot \nabla_x \partial^\alpha \Delta_x^{-1} \int_{\mathbb{R}^3} (S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_- dv, \mu^{-1} \partial^\alpha (S_{\mathcal{L}}(t)f)_{\pm})_{L_{x,v}^2} \\ &= (\partial^\alpha \phi, \int_{\mathbb{R}^3} v \cdot \nabla_x \partial^\alpha ((S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_-) dv)_{L_x^2} \\ &= (\partial^\alpha \phi, -\partial_t \partial^\alpha \int_{\mathbb{R}^3} ((S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_-) dv)_{L_x^2} = \frac{1}{2} \partial_t \|\partial^\alpha \nabla_x \phi\|_{L_x^2}^2. \end{aligned}$$

Here we take integration by parts on ∇_x with (3.50) and use boundary condition (1.12) to deduce $\int_{\mathbb{R}^3} v \cdot ((S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_-) dv = 0$ on $\partial\Omega$ for the case of union of cubes.

In order to derive the weighted version of (3.65), taking $H_x^2 L_v^2$ inner product of (3.64) with $\langle v \rangle^{2k} \mu^{-1} S_B(t)f$ and $\langle v \rangle^{2k} \mu^{-1} S_{\mathcal{L}}(t)f$ over $\Omega \times \mathbb{R}^3$ respectively, we have

$$\begin{aligned} \partial_t \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2 + \lambda \|\langle v \rangle^k (\mu^{-\frac{1}{2}} S_B(t)f)\|_{H_x^2 H_{\gamma/2}^s}^2 \\ \leq C_R \|\chi_R \mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2, \\ \partial_t \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 + \partial_t \|\nabla_x \phi\|_{H_x^2}^2 + \lambda \|\langle v \rangle^k (\mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f)\|_{H_x^2 H_{\gamma/2}^s}^2 \\ \leq C \|\nabla_x \phi\|_{H_x^2}^2 + C_R \|\chi_R \mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2. \end{aligned} \quad (3.67)$$

By macroscopic estimates from [19, Theorem 3.4] for the case of the union of cubes and exponential perturbation, we have

$$\begin{aligned} \partial_t \mathcal{E}_{int,B}(t) + \lambda \|\mu^{-\frac{1}{2}} \mathbf{P}\mu^{\frac{1}{2}} S_B(t)f\|_{H_x^2 H_{\gamma/2}^s}^2 &\leq C \|\mu^{-\frac{1}{2}} \{\mathbf{I} - \mathbf{P}\mu^{\frac{1}{2}}\} S_B(t)f\|_{H_x^2 L_{\gamma/2}^2}^2, \\ \partial_t \mathcal{E}_{int,\mathcal{L}}(t) + \partial_t \|\nabla_x \phi\|_{H_x^2}^2 + \lambda \|\mu^{-\frac{1}{2}} \mathbf{P}\mu^{\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 H_{\gamma/2}^s}^2 + \lambda \|\nabla_x \phi\|_{H_x^2}^2 \\ \leq C \|\mu^{-\frac{1}{2}} \{\mathbf{I} - \mathbf{P}\mu^{\frac{1}{2}}\} S_{\mathcal{L}}(t)f\|_{H_x^2 L_{\gamma/2}^2}^2 + \|\nabla_x \phi\|_{L_x^2}^4, \end{aligned} \quad (3.68)$$

where ϕ is given by (3.66) and $\mathcal{E}_{int,B}, \mathcal{E}_{int,\mathcal{L}}$ are functional satisfying

$$\mathcal{E}_{int,B}(t) \lesssim \|\mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2, \quad \mathcal{E}_{int,\mathcal{L}}(t) \lesssim \|\mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2. \quad (3.69)$$

respectively. Although the macroscopic estimates from [19] are for the union of cubes, the case of the torus can be similarly derived with simpler calculations; see also [44]. Taking linear combinations (3.65) + $\kappa^2 \times$ (3.67) + $\kappa \times$ (3.68) and assuming the *a priori* assumption

$$\|\nabla_x \phi\|_{H_x^2}^2 \leq \varepsilon, \quad (3.70)$$

with $\varepsilon > 0$ sufficiently small and ϕ given by (3.66), we have

$$\partial_t \mathcal{E}_B(t) + \lambda \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 H_{\gamma/2}^s}^2 \leq 0, \quad \partial_t \mathcal{E}_{\mathcal{L}}(t) + \lambda \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 H_{\gamma/2}^s}^2 + \lambda \|\nabla_x \phi\|_{H_x^2}^2 \leq 0, \quad (3.71)$$

where

$$\begin{aligned} \mathcal{E}_B(t) &= \|\mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2 + \kappa^2 \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_B(t)f\|_{H_x^2 L_v^2}^2 + \kappa \mathcal{E}_{int,B}, \\ \mathcal{E}_{\mathcal{L}}(t) &= \|\mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 + \kappa^2 \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 + \kappa \mathcal{E}_{int,\mathcal{L}} + \|\nabla_x \phi\|_{H_x^2}^2. \end{aligned}$$

With (3.69), it's direct to check that

$$\begin{aligned}\mathcal{E}_B(t) &\sim \|\mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_v^2}^2 + \|\langle v \rangle^k \mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_v^2}^2, \\ \mathcal{E}_{\mathcal{L}}(t) &\sim \|\mu^{-\frac{1}{2}}S_{\mathcal{L}}(t)f\|_{H_x^2L_v^2}^2 + \|\langle v \rangle^k \mu^{-\frac{1}{2}}S_{\mathcal{L}}(t)f\|_{H_x^2L_v^2}^2 + \|\nabla_x \phi\|_{H_x^2}^2.\end{aligned}$$

We calculate the first inequality of (3.71) and the second one is similar. When $\gamma \geq 0$, we have

$$\partial_t \mathcal{E}_B(t) + \lambda \|\langle v \rangle^k \mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_v^2}^2 \leq 0,$$

thus

$$\|\langle v \rangle^k \mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_v^2}^2 \leq e^{-\lambda t} \|\langle v \rangle^k \mu^{-\frac{1}{2}}f\|_{H_x^2L_v^2}^2.$$

When $\gamma < 0$, we have

$$\partial_t \mathcal{E}_B(t) + \lambda \|\langle v \rangle^{k+\frac{\gamma}{2}} \mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_v^2}^2 \leq 0.$$

Then we apply similar arguments for obtaining (3.62) to deduce that

$$\|\mu^{-\frac{1}{2}}S_B(t)f\|_{H_x^2L_{k-4}^2}^2 \lesssim \langle t \rangle^{-\frac{2(3-\frac{1}{2})}{|\gamma|}} \|\mu^{-\frac{1}{2}}f\|_{H_x^2L_k^2}^2 \lesssim \langle t \rangle^{-\frac{7}{6}} \|\mu^{-\frac{1}{2}}f\|_{H_x^2L_k^2}^2. \quad (3.72)$$

The estimate for hard potential case is also included in (3.72). Similarly, it follows from the second estimate of (3.71) that

$$\|\mu^{-\frac{1}{2}}S_{\mathcal{L}}(t)f\|_{H_x^2L_{k-4}^2}^2 + \|\nabla_x \phi\|_{H_x^2}^2 \lesssim \langle t \rangle^{-\frac{7}{6}} (\|\mu^{-\frac{1}{2}}f\|_{H_x^2L_k^2}^2 + \|\nabla_x \phi|_{t=0}\|_{H_x^2}^2). \quad (3.73)$$

Now we turn back to Duhamel's principle (3.54) and (3.55). For any f , we have from (3.54), (3.63) and (3.72) that for $k_1 \geq k_0$,

$$\begin{aligned}\|S_B(t)f\|_{H_x^2L_{k_1}^2} &\lesssim \|S_A(t)f\|_{H_x^2L_{k_1}^2} + \int_0^t \|\mu^{-\frac{1}{2}}S_B(t-s)K_1S_A(s)f\|_{H_x^2L_v^2} ds \\ &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} \|\mu^{-\frac{1}{2}}K_1S_A(s)f\|_{H_x^2L_v^2} ds \\ &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} \|S_A(s)f\|_{H_x^2L_v^2} ds \lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2}.\end{aligned} \quad (3.74)$$

Thus, recalling ϕ is given by (3.66), it follows from (3.55), (3.73) and (3.74) that for $k_1 \geq k_0$,

$$\begin{aligned}\|S_{\mathcal{L}}(t)f\|_{H_x^2L_{k_1}^2} &\lesssim \|S_B(t)f\|_{H_x^2L_{k_1}^2} + \int_0^t \|\mu^{-\frac{1}{2}}S_{\mathcal{L}}(t-s)K_2S_B(s)f\|_{H_x^2L_v^2} ds \\ &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} (\|\mu^{-\frac{1}{2}}K_2S_B(s)f\|_{H_x^2L_v^2} \\ &\quad + \|\nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} (K_2S_B(s)f)_+ - (K_2S_B(s)f)_- dv\|_{H_x^2}) ds.\end{aligned} \quad (3.75)$$

Note that by using odd property on μv , we have

$$\int_{\mathbb{R}^3} (K_2S_B(s)f)_+ - (K_2S_B(s)f)_- dv = 2 \int_{\mathbb{R}^3} \mu(v)v \cdot \nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} ((S_B(s)f)_+(u) - (S_B(s)f)_-(u)) du dv = 0.$$

Also, letting

$$\phi_1 = \Delta_x^{-1} \int_{\mathbb{R}^3} ((S_B(s)f)_+(u) - (S_B(s)f)_-(u)) du,$$

when Ω is torus, we know that

$$\|\nabla_x \phi_1\|_{L_x^2} \lesssim \|\Delta_x \phi_1\|_{L_x^2}. \quad (3.76)$$

If Ω is union of cubes given by (1.10), noticing from (3.50) that $\partial_{x_i} \phi_1 = 0$ on Γ_i , by Sobolev inequality [15, Theorem 6.7-5], we also have (3.76). Then we obtain from (3.75) and (3.63) that for $k_1 \geq k_0$,

$$\begin{aligned}\|S_{\mathcal{L}}(t)f\|_{H_x^2L_{k_1}^2} &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} \|\langle v \rangle^4 \mu^{\frac{1}{2}}(v)v \cdot \nabla_x \phi_1\|_{H_x^2L_v^2} ds \\ &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} \|\Delta_x \phi_1\|_{H_x^2L_{k_1+4}^2} ds \\ &\lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2} + \int_0^t \langle t-s \rangle^{-\frac{7}{6}} \|S_B(s)f\|_{H_x^2L_v^2} ds \lesssim \langle t \rangle^{-\frac{7}{6}} \|f\|_{H_x^2L_{k_1+4}^2}.\end{aligned} \quad (3.77)$$

Next, we check that (3.70) is fulfilled if $\varepsilon_0 > 0$ in (3.51) is small enough. Similar to (3.76), we have

$$\|\nabla_x \phi\|_{H_x^2} \lesssim \|\Delta_x \phi\|_{H_x^1} = \left\| \int_{\mathbb{R}^3} (S_{\mathcal{L}}(t)f)_+ - (S_{\mathcal{L}}(t)f)_- dv \right\|_{H_x^1}. \quad (3.78)$$

Then we can obtain that $\|\nabla_x \phi\|_{H_x^2} \lesssim \|S_{\mathcal{L}}(t)f\|_{H_x^1 L_v^2} \lesssim \|f\|_{H_x^2 L_{k_0+4}^2}$. This closes the *a priori* assumption (3.70) and completes the proof of (3.52).

Step 2. For brevity of notations, we let $s = 1$ for the Landau case. In order to obtain (3.53), we consider the dual A_l^* of $A_l = \langle v \rangle^l A(\langle v \rangle^{-l} f)$ for any $l \in \mathbb{R}$. Notice that

$$(A_l^* f, \langle v \rangle^{2k} f)_{L_x^2 L_v^2} = (\langle v \rangle^l f, A(\langle v \rangle^{2k-l} f))_{L_x^2 L_v^2} = (\langle v \rangle^{2l-2k} g, Ag)_{L_x^2 L_v^2},$$

where we let $g = \langle v \rangle^{2k-l} f$. Then the taking $L_x^2 L_v^2$ inner product of $\partial_t S_{A_l^*}(t)f - A_l^* S_{A_l^*}(t)f = 0$ with $\langle v \rangle^{2k} S_{A_l^*}(t)f$, we have

$$\partial_t \|\langle v \rangle^k S_{A_l^*}(t)f\|_{L_x^2 L_v^2}^2 + (A_l^* S_{A_l^*}(t)f, \langle v \rangle^{2k} S_{A_l^*}(t)f)_{L_x^2 L_v^2} = 0.$$

Hence for $l - k \geq k_0$,

$$\partial_t \|\langle v \rangle^{l-k} g\|_{L_x^2 L_v^2}^2 + (\langle v \rangle^{2l-2k} g, Ag)_{L_x^2 L_v^2} = 0,$$

where $g = \langle v \rangle^{2k-l} S_{A_l^*}(t)f$. Integrate over t , we have

$$\lambda \int_0^\infty \|\langle v \rangle^{l-k} g\|_{L_x^2 H_{\gamma/2}^s}^2 dt \leq \|\langle v \rangle^{l-k} g|_{t=0}\|_{L_x^2 L_v^2}^2.$$

That is, for $l - k \geq k_0$,

$$\lambda \int_0^\infty \|\langle v \rangle^k S_{A_l^*}(t)f\|_{L_x^2 H_{\gamma/2}^s}^2 dt \leq \|\langle v \rangle^k f\|_{L_x^2 L_v^2}^2. \quad (3.79)$$

Observe that if f satisfies equation $\partial_t f = Af$ then $g = \langle v \rangle^l f$ satisfies $\partial_t g = A_l g$, and also that $(Af, \langle v \rangle^{2l} f)_{L_{x,v}^2} = (A_l g, g)_{L_{x,v}^2}$. Thus, $\langle v \rangle^l S_A(t)f = S_{A_l}(\langle v \rangle^l f)$. Moreover, by duality, we have

$$(S_{A_l} f, g)_{L_{x,v}^2} = (f, S_{A_l^*} g)_{L_{x,v}^2}.$$

Therefore, for some sequence $\{\varphi_n\}$ in Schwartz space such that $\|\varphi_n\|_{L_x^2 L_v^2} \leq 1$, we have from (3.79) that

$$\begin{aligned} \int_0^\infty \|\langle v \rangle^k S_A(t)f\|_{L_x^2 L_v^2}^2 dt &= \int_0^\infty \lim_{n \rightarrow \infty} |(\langle v \rangle^{2k} S_A(t)f, \varphi_n)_{L_{x,v}^2}|^2 dt = \liminf_{n \rightarrow \infty} \int_0^\infty |(\langle v \rangle^{2k} f, S_{A_{2k}^*}(t)\varphi_n)_{L_{x,v}^2}|^2 dt \\ &\leq \liminf_{n \rightarrow \infty} \int_0^\infty \|\langle v \rangle^k f\|_{L_x^2 H_{-\gamma/2}^{-s}}^2 \|\langle v \rangle^k S_{A_{2k}^*}(t)\varphi_n\|_{L_x^2 H_{\gamma/2}^s}^2 dt \lesssim \|\langle v \rangle^{k-\gamma/2} f\|_{L_x^2 H_v^{-s}}^2, \end{aligned}$$

for $k \geq k_0$. This is the estimate for $S_A(t)$. To obtain the estimates for $S_B(t)$, it follows from Duhamel's principle (3.54) that

$$\begin{aligned} \left(\int_0^\infty \|\langle v \rangle^k S_B(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} &\lesssim \left(\int_0^\infty \|\langle v \rangle^k S_A(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \\ &\quad + \left(\int_0^\infty \left\| \langle v \rangle^k \int_0^t S_B(s) K_1 S_A(t-s)f ds \right\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}}. \end{aligned} \quad (3.80)$$

Using (3.72), the second right-hand term of (3.80) can be estimated by

$$\begin{aligned} \int_0^\infty \left(\int_s^\infty \|\mu^{-\frac{1}{2}} S_B(s) K_1 S_A(t-s)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} ds &\lesssim \int_0^\infty \left(\int_0^\infty \langle s \rangle^{-\frac{7}{6}} \|\langle v \rangle^k \mu^{-\frac{1}{2}} K_1 S_A(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} ds \\ &\lesssim \left(\int_0^\infty \|S_A(s)f\|_{H_x^2 L_v^2}^2 ds \right)^{\frac{1}{2}} \lesssim \|f\|_{H_x^2 H_{k_0-\gamma/2}^{-s}}. \end{aligned}$$

Note that ∇_x commutes with $S_A(t)$. The first term on the right-hand side of (3.80) can be estimated in the same way. Thus, for $k \geq k_0$, we have from (3.80) that

$$\left(\int_0^\infty \|\langle v \rangle^k S_B(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \lesssim \|f\|_{H_x^2 H_{k-\gamma/2}^{-s}}. \quad (3.81)$$

Next we apply (3.55), and (3.81) to derive the estimates on $S_{\mathcal{L}}(t)$:

$$\left(\int_0^\infty \|\langle v \rangle^k S_{\mathcal{L}}(t)f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}}$$

$$\begin{aligned}
& \lesssim \left(\int_0^\infty \|\langle v \rangle^k S_B(t) f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} + \left(\int_0^\infty \left\| \langle v \rangle^k \int_0^t S_{\mathcal{L}}(s) K_2 S_B(t-s) f ds \right\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \\
& \lesssim \|f\|_{H_x^2 H_{k-\gamma/2}^-} + \int_0^\infty \left(\int_s^\infty \|\langle v \rangle^k S_{\mathcal{L}}(s) K_2 S_B(t-s) f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} ds.
\end{aligned} \tag{3.82}$$

Applying (3.73), (3.76) and (3.81), the second right-hand term of (3.82) can be estimated by

$$\begin{aligned}
& \int_0^\infty \left(\int_s^\infty \|\langle v \rangle^k \mu^{-\frac{1}{2}} S_{\mathcal{L}}(s) K_2 S_B(t-s) f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} ds \\
& \lesssim \int_0^\infty \langle s \rangle^{-\frac{7}{6}} \left(\int_0^\infty \|\langle v \rangle^{k+4} \mu^{-\frac{1}{2}} K_2 S_B(t) f\|_{H_x^2 L_v^2}^2 \right. \\
& \quad \left. + \underbrace{\|\nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} (K_2 S_B(t) f)_+ - (K_2 S_B(t) f)_- dv\|_{H_x^2}^2}_{=0} dt \right)^{\frac{1}{2}} ds \\
& \lesssim \left(\int_0^\infty \|\langle v \rangle^{k+4} \mu^{-\frac{1}{2}} \mu v \cdot \nabla_x \Delta_x^{-1} \int_{\mathbb{R}^3} ((S_B(t) f)_+ - (S_B(t) f)_-) dv\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \\
& \lesssim \left(\int_0^\infty \|S_B(t) f\|_{H_x^2 L_v^2}^2 dt \right)^{\frac{1}{2}} \lesssim \|f\|_{H_x^2 H_{k-\gamma/2}^-}.
\end{aligned}$$

The term with brace is equal to zero because $\int_{\mathbb{R}^3} v \mu dv = 0$. Inserting this into (3.82), we obtain (3.53). This completes the proof of Lemma 3.3. \square

Next, we can introduce a norm

$$\|f\|_{\|\cdot\|}^2 := \sum_{|\alpha| \leq 2} \int_0^{+\infty} \|S_{\mathcal{L}}(\tau) \partial^\alpha f\|_{L_x^2 L_v^2}^2 d\tau, \tag{3.83}$$

where the associated inner product is given by

$$((f, g)) := \int_0^{+\infty} \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \partial^\alpha f, S_{\mathcal{L}}(\tau) \partial^\alpha g)_{L_x^2 L_v^2} d\tau,$$

Note that ∂^α commutes with \mathcal{L} and hence, commutes with $S_{\mathcal{L}}(t)$. Then by (3.52) and (3.53), we have

$$\|f\|_{\|\cdot\|} \lesssim \|f\|_{H_x^2 L_{k_0+4}^2}, \quad \|f\|_{\|\cdot\|} \lesssim \|f\|_{H_x^2 H_{k_0-\frac{\gamma}{2}}^-} \lesssim \|f\|_{H_x^2 H_{k_0+\frac{3}{2}}^-}, \tag{3.84}$$

where we let $s = 1$ for the Landau case. Then we can recover the loss of energy by using this norm.

Lemma 3.4. *For both the Boltzmann and Landau cases, suppose (f, ϕ) is the solution to the equation (1.3). Then we have*

$$\frac{1}{2} \partial_t \|f\|_{\|\cdot\|}^2 + \|f\|_{H_x^2 L_v^2}^2 \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t), \tag{3.85}$$

where $\|\cdot\|_{\|\cdot\|}$ is given in (3.83), $\mathcal{E}_k(t)$ and $\mathcal{D}_k(t)$ are given in (1.21) and (1.22).

Proof. Recall the first equation of (1.3):

$$\partial_t f_\pm = \underbrace{\pm \nabla_x \phi \cdot \nabla_v f}_{\text{Term 1}} + \underbrace{\mathcal{L} f}_{\text{Term 2}} + \underbrace{Q(f_\pm + f_\mp, f_\pm)}_{\text{Term 3}}, \tag{3.86}$$

where \mathcal{L} is given in (1.4). Then we take the $((\cdot, \cdot))$ inner product of (3.86) with f and compute every term separately. Since ∂_t commute with L , we know that ∂_t commute with $S_{\mathcal{L}}(\tau)$ and hence,

$$((\partial_t \partial^\alpha f, \partial^\alpha f)) = \frac{1}{2} \partial_t \| \partial^\alpha f \|_{\|\cdot\|}^2.$$

Next, we compute Term 1. Since $s \geq \frac{1}{2}$ implies $1 - s \leq s$, by Lemma 2.10 and (3.84), we have

$$\begin{aligned}
& \int_0^{+\infty} \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \partial^\alpha (\nabla_x \phi \cdot \nabla_v f), S_{\mathcal{L}}(\tau) \partial^\alpha f)_{L_x^2 L_v^2} d\tau \\
& \lesssim \sum_{|\alpha| \leq 2} \|\partial^\alpha (\nabla_x \phi \cdot \nabla_v f)\|_{L_x^2 H_{k_0+3/2}^-} \sum_{|\alpha| \leq 2} \|\partial^\alpha f\|_{L_x^2 L_{k_0+3/2}^2} \\
& \lesssim \|\nabla_x \phi\|_{H_x^2} \|f\|_{H_x^2 H_{k_0+3/2}^{1-s}} \|f\|_{H_x^2 L_{k_0+3/2}^2} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t),
\end{aligned}$$

for the non-cutoff Boltzmann case, where we let $k \geq k_0 + 3/2$. Similarly, for the Landau case, we have from (3.84) that for $k \geq k_0 + 4$,

$$\begin{aligned} \int_0^{+\infty} \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \partial^\alpha (\nabla_x \phi \cdot \nabla_v f), S_{\mathcal{L}}(\tau) \partial^\alpha f)_{L_x^2 L_v^2} d\tau &\lesssim \sum_{|\alpha| \leq 2} \|\partial^\alpha (\nabla_x \phi \cdot \nabla_v f)\|_{L_x^2 L_{k_0+4}^2} \|f\|_{H_x^2 L_{k_0+4}^2} \\ &\lesssim \|\nabla_x \phi\|_{H_x^2} \|f\|_{H_x^2 H_{k_0+4}^1} \|f\|_{H_x^2 L_{k_0+4}^2} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t). \end{aligned}$$

For the term 2 and for both Boltzmann and Landau case, since $\lim_{t \rightarrow \infty} \langle t \rangle^{-\frac{7}{6}} = 0$ in (3.52), we have

$$\int_0^\infty \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \mathcal{L} \partial^\alpha f, S_{\mathcal{L}}(\tau) \partial^\alpha f)_{L_x^2 L_v^2} d\tau = \int_0^\infty \frac{d}{d\tau} \sum_{|\alpha| \leq 2} \|S_{\mathcal{L}}(\tau) \partial^\alpha f\|_{L_x^2 L_v^2}^2 d\tau = - \sum_{|\alpha| \leq 2} \|\partial^\alpha f\|_{L_x^2 L_v^2}^2.$$

Finally, we consider the nonlinear part, i.e. Term 3. For the Boltzmann case, by (2.14), (3.84), we have

$$\begin{aligned} &\int_0^\infty \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \partial^\alpha Q(f_\pm + f_\mp, f_\pm), S_{\mathcal{L}}(\tau) \partial^\alpha f)_{L_x^2 L_v^2} d\tau \\ &\leq C_k \sum_{|\alpha| \leq 2} \sum_{\alpha_1 \leq \alpha} \|Q(\partial^{\alpha_1} (f_\pm + f_\mp), \partial^{\alpha - \alpha_1} f_\pm)\|_{L_x^2 H_{k_0 - \gamma/2}^{-s}} \|\partial^\alpha f\|_{L_x^2 L_{k_0}^2} \\ &\leq C_k (\|f\|_{H_x^2 L_{14}^2} \|f\|_{H_x^2 H_{k_0 + \gamma/2 + 2s}^s} + \|f\|_{H_x^2 L_{14}^2} \|f\|_{H_x^2 H_{k_0 + \gamma/2}^s}) \|\partial^\alpha f\|_{L_x^2 L_{k_0}^2} \leq C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t). \end{aligned}$$

Here we let $k_0 \geq 14$, $k \geq k_0 + \gamma/2 + 2s$ and apply similar discussion on α_1 as in (2.53). For the Landau case, by (2.18), (3.84), we have

$$\begin{aligned} &\int_0^\infty \sum_{|\alpha| \leq 2} (S_{\mathcal{L}}(\tau) \partial^\alpha Q(f_\pm + f_\mp, f_\pm), S_{\mathcal{L}}(\tau) \partial^\alpha f)_{L_x^2 L_v^2} d\tau \\ &\leq C_k \sum_{|\alpha| \leq 2} \sum_{\alpha_1 \leq \alpha} \|Q(\partial^{\alpha_1} f_\pm + f_\mp, \partial^{\alpha - \alpha_1} f_\pm)\|_{L_x^2 H_{k_0 + 3/2}^{-1}} \|\partial^\alpha f\|_{L_x^2 L_{k_0}^2} \\ &\leq C_k \|f\|_{H_x^2 L_{k_0 + 5/2}^2} \|f\|_{H_x^2 L_{D, k_0 + 5/2}^2} \|\partial^\alpha f\|_{L_x^2 L_{k_0 + 5/2}^2} \leq C_k \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t), \end{aligned}$$

where $k \geq k_0 + 4$ and we also apply similar discussion on α_1 as in (2.53). Combining the above estimates, the $((\cdot, \cdot))$ inner product of (3.86) with $\partial^\alpha f$ yields

$$\frac{1}{2} \partial_t \|f\|^2 + \sum_{|\alpha| \leq 2} \|\partial^\alpha f\|_{L_x^2 L_v^2}^2 \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t).$$

Then we conclude Lemma 3.4. \square

3.4. Proof of the main theorem. We start this subsection by proving the main stability theorem as below. Theorem 3.5 and 3.6 together with local existence from Theorem 2.24 will imply Theorem 1.2. We give the details as the following. To prove Theorem 1.2, we assume the *a priori* assumption as

$$\sup_{0 \leq t \leq T} \mathcal{E}_{k_0}(t) \leq 2M, \quad (3.87)$$

for the case $\gamma \in [0, 1]$ and

$$\sup_{0 \leq t \leq T} (1+t)^{\frac{2l}{|\gamma|}} \mathcal{E}_{k_0}(t) \leq CM, \quad (3.88)$$

for the case $\gamma \in [-3, 0)$, where $l > \frac{|\gamma|}{2}$ is a constant and we further assume $k \geq k_0 + l$ in this case.

Theorem 3.5. *Assume that f_0 satisfies the conservation laws (1.9) and $F_0 = \mu + f_0 \geq 0$. There exists $k_1 \geq k_0$ such that for*

$$k \geq k_0 \text{ for Landau case and } k \geq k_1 \text{ for Boltzmann case,}$$

there exists a small constant $M \geq 0$ such that if $\mathcal{E}_{k_0}(0) \leq M, \mathcal{E}_k(0) < \infty$, then there exist a unique global solution $f(t, x, v)$ to the Vlasov-Poisson-Boltzmann/Landau system (1.3) with $F = \mu + f \geq 0$ satisfying

$$\partial_t \mathcal{E}_k(t) + \lambda \mathcal{D}_k(t) \lesssim \|f\|_{X_{k_0}} \mathcal{E}_k(t), \quad (3.89)$$

for any $T > 0$, for some generic constant $\lambda > 0$.

Proof. We take linear combination $\kappa \times (3.4) + (3.30) + C_0 \times (3.85)$ with $\kappa, C_0 > 0$ to deduce that

$$\begin{aligned} & \partial_t \mathcal{E}_k(t) + \lambda \kappa \| [a_+, a_-, b, c] \|_{H_x^2}^2 + \lambda \kappa \| \nabla_x \phi \|_{H_x^2}^2 + \lambda \eta \| f \|_{Y_k}^2 + C_0 \| f \|_{H_x^2 L_v^2}^2 + \lambda \sum_{|\alpha| \leq 2} \| w(\alpha, 0) \partial^\alpha \{ \mathbf{I} - \mathbf{P} \} f \|_{L_x^2 L_v^2}^2 \\ & \lesssim \| f \|_{X_{k_0}} \| f \|_{X_k}^2 + \kappa \| \{ \mathbf{I} - \mathbf{P} \} f \|_{H_x^2 L_{10}^2}^2 + \kappa \| N_{\parallel} \|_{H_x^1 L_v^2}^2 + C_{\eta, k} \| \{ \mathbf{I} - \mathbf{P} \} f \|_{H_x^2 L_v^2}^2 \\ & \quad + \eta \| \nabla_x \phi \|_{H_x^2}^2 + \eta \| \mathbf{P} f \|_{H_x^2 L_v^2}^2 + C(\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t), \end{aligned}$$

for some generic constant $\lambda > 0$, where $\mathcal{E}_k(t)$ is given by

$$\mathcal{E}_k(t) := \kappa G(t) + \mathcal{E}_1(t) + \frac{C_0}{2} \sum_{|\alpha| \leq 2} \| \partial^\alpha f \|^2. \quad (3.90)$$

Here $G(t)$, $\mathcal{E}_1(t)$ and $\| \cdot \|$ are given in Lemma 3.1, Lemma 3.2 and (3.83) respectively. Choosing $\kappa > 0$ sufficiently small, and then $\eta > 0$ sufficiently small, and finally $C_0 > 0$ sufficiently large, we have

$$\partial_t \mathcal{E}_k(t) + \lambda \mathcal{D}_k(t) \lesssim \| f \|_{X_{k_0}} \mathcal{E}_k(t) + \kappa \| N_{\parallel} \|_{H_x^1 L_v^2}^2 + (\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t), \quad (3.91)$$

for some generic constant $\lambda > 0$, where $\mathcal{D}_k(t)$ is given by (1.22). Then one can check (1.21) by using (3.3), (3.47), (3.84) with sufficiently small κ . Note that $e^{\frac{\pm A_{\alpha, 0} \phi}{2(v)^2}} \approx 1$ as in (2.31).

Next, recall from Lemma 3.1 that

$$N_{\parallel} = (\pm \nabla_x \phi \cdot \nabla_v f_{\pm})_{\parallel} + Q(f_{\pm} + f_{\mp}, f_{\pm})_{\parallel} = (\pm \nabla_x \phi \cdot \nabla_v f_{\pm}, \xi) + (Q(f_{\pm} + f_{\mp}, f_{\pm}), \xi),$$

where ξ is some linear combination of $1, v_i, v_i v_j, v_i^2, v_i |v|^2$. For $|\alpha| \leq 1$, we have

$$\begin{aligned} \| \partial^\alpha (\nabla_x \phi \cdot \nabla_v f, \xi) \|_{L_x^2 L_v^2} & \lesssim \sum_{|\alpha_1|=0} \| \partial^{\alpha_1} \nabla_x \phi \|_{L_x^\infty} \| \partial^{\alpha - \alpha_1} \nabla_v f \|_{L_x^2 L_v^2} + \sum_{|\alpha_1|=1} \| \partial^{\alpha_1} \nabla_x \phi \|_{L_x^2} \| \partial^{\alpha - \alpha_1} \nabla_v f \|_{L_x^0 L_v^2} \\ & \lesssim \| \nabla_x \phi \|_{H_x^2} \| f \|_{H_x^1 H_v^1} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t). \end{aligned} \quad (3.92)$$

For the Boltzmann case, by Lemma 2.4, we apply similar discussion on α_1 as in (3.92) to deduce that

$$\| \partial^\alpha (Q(f_{\pm} + f_{\mp}, f_{\pm}), \xi) \|_{L_{x,v}^2} \lesssim \| f \|_{H_x^2 L_{10}^2} \| f \|_{H_x^2 L_{10}^2} \| \xi \|_{H_{-7}^{2s}} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t).$$

For the Landau case, by (2.17) we have

$$\| \partial^\alpha (Q(f_{\pm} + f_{\mp}, f_{\pm}), \xi \langle v \rangle^{-14} \langle v \rangle^{14}) \|_{L_{x,v}^2} \lesssim \| f \|_{H_x^2 L_{10}^2} \| f \|_{H_x^2 L_{D,8}^2} \| \xi \|_{L_{D,-7}^2} \lesssim \sqrt{\mathcal{E}_{k_0}(t)} \mathcal{D}_k(t).$$

Therefore, we obtain $\sum_{|\alpha| \leq 1} \| \partial^\alpha N_{\parallel} \|_{L_x^2}^2 \lesssim \mathcal{E}_{k_0}(t) \mathcal{D}_k(t)$. Then (3.91) implies

$$\partial_t \mathcal{E}_k(t) + \lambda \mathcal{D}_k(t) \lesssim \| f \|_{X_{k_0}} \mathcal{E}_k(t) + (\sqrt{\mathcal{E}_{k_0}(t)} + \mathcal{E}_{k_0}(t)) \mathcal{D}_k(t). \quad (3.93)$$

Choosing M in the *a priori* assumption (3.87) and (3.88) sufficiently small, we have from (3.93) that $\partial_t \mathcal{E}_k(t) + \lambda \mathcal{D}_k(t) \lesssim \| f \|_{X_{k_0}} \mathcal{E}_k(t)$ for some $\lambda > 0$. This concludes Theorem 3.5. \square

To conclude Theorem 1.2, we need to prove the large-time behavior as the following.

Theorem 3.6. *Let $l = 0$ for hard potential case and $l > \frac{|l|}{2}$ for soft potential case. Let $k \geq k_0 + 2 + l$ (and let k sufficiently large for Boltzmann case). If the solution (f, ϕ) of (1.3) satisfies $\mathcal{E}_{k_0+2+l}(0) \leq M$, $\mathcal{E}_k(0) < \infty$. Then there exists a constant $\lambda > 0$, such that for any $t > 0$, we have*

$$\sup_{0 \leq t \leq T} \mathcal{E}_k(t) \leq 2\mathcal{E}(0). \quad (3.94)$$

Moreover, we have $\mathcal{E}_k(t) \leq e^{-\lambda t} \mathcal{E}_k(0)$ if $0 \leq \gamma \leq 1$, $\mathcal{E}_{k-l}(t) \lesssim (1+t)^{-\frac{2l}{|\gamma|}} \mathcal{E}_k(0)$ if $\gamma < 0$.

Proof. If $0 \leq \gamma \leq 1$, noticing that $\| \cdot \|_{L_{x,v}^2} \lesssim \| \cdot \|_{L_D^2}$ and choosing M in (3.87) small enough, we have $\| f \|_{X_{k_0}} \lesssim \mathcal{E}_{k_0} \lesssim M$, and hence, by (3.89) we have $\partial_t \mathcal{E}_k(t) + \lambda \mathcal{E}_k(t) \leq 0$. Solving this ODE, we obtain $\mathcal{E}_k(t) \leq e^{-\lambda t} \mathcal{E}_k(0)$. This closes the *a priori* assumption (3.87) and concludes the case of hard potential.

Next, we assume $\gamma < 0$. For any $l > \frac{|l|}{2}$, we have from (3.88) that

$$\| f \|_{X_{k_0}} \lesssim \mathcal{E}_{k_0}(t) \lesssim M(1+t)^{-\frac{2l}{|\gamma|}}. \quad (3.95)$$

Let $p = \frac{-\gamma+2l}{2l}$ and $p' = \frac{-\gamma+2l}{-\gamma}$. Then by $L^p - L^{p'}$ Hölder's inequality, we have

$$\| w(\alpha, \beta) \partial_\beta^\alpha f \|_{L_v^2}^2 = \int_{\mathbb{R}^3} \langle v \rangle^{-\frac{2l\gamma}{-\gamma+2l}} \langle v \rangle^{-\frac{2l\gamma}{-\gamma+2l}} |w(\alpha, \beta) \partial_\beta^\alpha f|^2 dv$$

$$\leq \|\langle v \rangle^{\gamma/2} w(\alpha, \beta) \partial_\beta^\alpha f\|_{L_v^2}^{-\frac{4l}{\gamma+2l}} \|\langle v \rangle^l w(\alpha, \beta) \partial_\beta^\alpha f\|_{L_v^2}^{-\frac{2\gamma}{\gamma+2l}} \leq \mathcal{D}_k^{-\frac{2l}{\gamma+2l}} \mathcal{E}_{k+l}^{-\frac{\gamma}{\gamma+2l}}.$$

From definition (1.21) and (1.19), we have $\mathcal{E}_k^{-\frac{\gamma+2l}{2l}} \mathcal{E}_{k+l}^{\frac{\gamma}{2l}} \leq \mathcal{D}_k$. Then it follows from (3.89) and (3.95) that

$$\partial_t \mathcal{E}_k(t) + \lambda \mathcal{E}_k^{-\frac{\gamma+2l}{2l}}(t) \left(\sup_{0 \leq t \leq T} \mathcal{E}_{k+l} \right)^{\frac{\gamma}{2l}} \leq CM(1+t)^{-\frac{2l}{|\gamma|}} \mathcal{E}_k(t). \quad (3.96)$$

Neglecting the second left hand term of (3.96), we have $\partial_t \mathcal{E}_k(t) \leq CM(1+t)^{-\frac{2l}{|\gamma|}} \mathcal{E}_k(t)$. Taking integration over $t \in [0, T]$ yields

$$\sup_{0 \leq t \leq T} \mathcal{E}_k(t) \leq \mathcal{E}_k(0) + CM \int_0^T (1+t)^{-\frac{2l}{|\gamma|}} dt \sup_{0 \leq t \leq T} \mathcal{E}_k(t).$$

Since $\frac{2l}{|\gamma|} > 1$, choosing $M > 0$ sufficiently small, we have

$$\sup_{0 \leq t \leq T} \mathcal{E}_k(t) \leq 2\mathcal{E}_k(0). \quad (3.97)$$

This gives (3.94). Next, we solve (3.96) directly. It's direct to obtain that

$$\partial_t (\mathcal{E}_k^{\frac{\gamma}{2l}}(t)) = \frac{\gamma}{2l} \mathcal{E}_k^{\frac{\gamma-2l}{2l}}(t) \partial_t \mathcal{E}_k(t) \geq -\frac{\gamma\lambda}{2l} \left(\sup_{0 \leq t \leq T} \mathcal{E}_{k+l} \right)^{\frac{\gamma}{2l}} + \frac{CM\gamma}{2l} (1+t)^{-\frac{2l}{|\gamma|}} \mathcal{E}_k^{\frac{\gamma}{2l}}(t),$$

and thus

$$\partial_t \left(\exp \left\{ \frac{CM|\gamma|^2}{2l(-2l+|\gamma|)} (1+t)^{-\frac{2l}{|\gamma|}+1} \right\} \mathcal{E}_k^{\frac{\gamma}{2l}}(t) \right) \geq -\frac{\gamma\lambda}{2l} \exp \left\{ \frac{CM|\gamma|^2}{2l(-2l+|\gamma|)} (1+t)^{-\frac{2l}{|\gamma|}+1} \right\} \left(\sup_{0 \leq t \leq T} \mathcal{E}_{k+l} \right)^{\frac{\gamma}{2l}}.$$

Note that $-2l+|\gamma| < 0$. Taking integration over $t \in [0, T]$, we have

$$\exp \left\{ \frac{CM|\gamma|^2}{2l(-2l+|\gamma|)} (1+t)^{-\frac{2l}{|\gamma|}+1} \right\} \mathcal{E}_k^{\frac{\gamma}{2l}}(t) \geq C_{\gamma,l} \mathcal{E}_k^{\frac{\gamma}{2l}}(0) + C_{\gamma,l} \int_0^t 1 dt \left(\sup_{0 \leq t \leq T} \mathcal{E}_{k+l} \right)^{\frac{\gamma}{2l}},$$

and hence $\mathcal{E}_k(t) \leq C_{\gamma,l} (1+t)^{\frac{2l}{\gamma}} \sup_{0 \leq t \leq T} \mathcal{E}_{k+l}(t)$. Replacing k by $k-l \geq k_0$, we apply (3.97) to deduce

$$\mathcal{E}_{k-l}(t) \lesssim (1+t)^{\frac{2l}{\gamma}} \sup_{0 \leq t \leq T} \mathcal{E}_k(t) \lesssim (1+t)^{\frac{2l}{\gamma}} \mathcal{E}_k(0).$$

Noticing $\gamma < 0$, choosing $k-l = k_0$ and applying (1.23), we close the *a priori* assumption (3.88). Using the standard continuity arguments, we complete the proof of Theorem 3.6. \square

Acknowledgments. C.-Q. Cao has been supported by grants from Beijing Institute of Mathematical Sciences and Applications and Yau Mathematical Science Center, Tsinghua University. D.-Q. Deng was partially supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) No. RS-2023-00210484 and No. RS-2023-00212304. X.-Y. Li has been supported by grants from project ANR-17-CE40-0030 of the French National Research Agency (ANR) and CEREMADE, Université Paris Dauphine.

Conflict of Interest: The authors declare that they have no conflict of interest.

REFERENCES

- [1] R. ALEXANDRE, Y. MORIMOTO, S. UKAI, C.-J. XU, AND T. YANG, *Regularizing effect and local existence for the non-cutoff Boltzmann equation*, Arch. Ration. Mech. Anal., 198 (2010), pp. 39-123.
- [2] R. ALEXANDRE, Y. MORIMOTO, S. UKAI, C.-J. XU AND T. YANG, *The Boltzmann equation without angular cutoff in the whole space: I, Global existence for soft potential*, J. Funct. Anal., 262 (2012), pp. 915-1010.
- [3] R. ALEXANDRE, Y. MORIMOTO, S. UKAI, C.-J. XU, AND T. YANG, *The Boltzmann equation without angular cutoff in the whole space: II, Global existence for hard potential*, Anal. Appl. (Singap.), 9 (2011), pp. 113-134.
- [4] R. ALEXANDRE, Y. MORIMOTO, S. UKAI, C.-J. XU AND T. YANG, *Global existence and full regularity of the Boltzmann equation without angular cutoff*, Comm. Math. Phys., 304 (2011), pp. 513-581.
- [5] R. ALEXANDRE AND C. VILLANI, *On the Boltzmann equation for long-range interactions*, Comm. Pure Appl. Math., 55 (2022), pp. 30-70.
- [6] R. ALONSO, Y. MORIMOTO, W. SUN AND T. YANG, *Non-cutoff Boltzmann equation with polynomial decay perturbation*, Rev. Mat. Iberoam., 37 (2021), pp. 189-292.
- [7] C. CAO, *Cutoff Boltzmann equation with polynomial perturbation near Maxwellian*, J. Funct. Anal., 283 (2022), pp. 109641.
- [8] C. CAO., L-B. HE AND J.JI, *Propagation of moments and sharp convergence rate for inhomogeneous non-cutoff Boltzmann equation with soft potentials*, Preprint, [arXiv:2204.01394](https://arxiv.org/abs/2204.01394).

- [9] Y. CAO, C. KIM AND D. LEE, *Global Strong Solutions of the Vlasov–Poisson–Boltzmann System in Bounded Domains*, Arch. Ration. Mech. Anal., 233 (2019), pp. 1027–1130.
- [10] K. CARRAPATOSO, I. TRISTANI AND K.-C. WU, *Cauchy problem and exponential stability for the inhomogeneous Landau equation*, Arch. Ration. Mech. Anal. 221 (2016), pp. 363–418.
- [11] K. CARRAPATOSO AND S. MISCHLER, *Landau equation for very soft and Coulomb potentials near Maxwellians*, Ann. PDE, 3 (2017), pp. 1–65.
- [12] S. CHATURVEDI, *Stability of Vacuum for the Boltzmann Equation with Moderately Soft Potentials*, Ann. PDE, 7 (2021).
- [13] Y. CHEN AND L.-B. HE, *Smoothing estimates for Boltzmann equation with full-range interactions: Spatially homogeneous case*, Arch. Ration. Mech. Anal, 201 (2011), pp. 501–548.
- [14] Y. CHEN AND L.-B. HE, *Smoothing estimates for Boltzmann equation with full-range interactions: Spatially inhomogeneous case*, Arch. Ration. Mech. Anal, 203 (2012), pp. 343–377.
- [15] P. G. CIARLET, *Linear and Nonlinear Functional Analysis with Applications*, SIAM, 2013.
- [16] D.-Q. DENG, *Dissipation and Semigroup on H_n^k : Non-cutoff Linearized Boltzmann Operator with Soft Potential*, SIAM J. Math. Anal., 52 (2020), pp. 3093–3113.
- [17] D.-Q. DENG, *The Landau and Non-cutoff Boltzmann Equation in Union of Cubes*, Preprint, [arXiv:2106.00383](https://arxiv.org/abs/2106.00383).
- [18] D.-Q. DENG, *Regularity of the Vlasov–Poisson–Boltzmann System Without Angular Cutoff*, Comm. Math. Phys., 387 (2021), pp. 1603–1654.
- [19] D.-Q. DENG, *The non-cutoff Vlasov-Poisson-Boltzmann and Vlasov-Poisson-Landau Systems in Union of Cubes*, Preprint, [arXiv:2112.02315](https://arxiv.org/abs/2112.02315).
- [20] L. DESVILLETES AND C. VILLANI, *On the trend to global equilibrium for spatially inhomogeneous kinetic systems: the Boltzmann equation*, Invent. Math., 159 (2015), pp. 245–316.
- [21] R. J. DIPERNA AND P.-L. LIONS, *On the Fokker-Planck-Boltzmann equation*, Comm. Math. Phys., 120 (1988), pp. 1–23.
- [22] R. J. DIPERNA AND P.-L. LIONS, *On the Cauchy problem for Boltzmann equations: global existence and weak stability*, Ann. of Math., 130 (1989), pp. 321–366.
- [23] H. DONG, Y. GUO AND Z. OUYANG, *The Vlasov-Poisson-Landau System with the Specular-Reflection Boundary Condition*, Arch. Ration. Mech. Anal., 246 (2022), pp. 333–396.
- [24] R. DUAN, Y. LEI, T. YANG AND Z. ZHAO, *The Vlasov-Maxwell-Boltzmann system near Maxwellians in the whole space with very soft potentials*, Comm. Math. Phys, 351 (2017), pp. 95–153.
- [25] R. DUAN, *Global smooth dynamics of a fully ionized plasma with long-range collisions*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 31 (2014), pp. 751–778.
- [26] R. DUAN AND S. LIU, *The Vlasov-Poisson-Boltzmann system without angular cutoff*, Commun. Math. Phys. 324 (2013), pp. 1–45.
- [27] R. DUAN, S. LIU, S. SAKAMOTO AND R. STRAIN, *Global mild solutions of the Landau and non-cutoff Boltzmann equations*, Comm. Pure Appl. Math., 74 (2021), pp. 932–1020.
- [28] R. DUAN, S. LIU, T. YANG AND H. ZHAO, *Stability of the nonrelativistic Vlasov-Maxwell-Boltzmann system for angular non-cutoff potentials*, Kinet. Relat. Models, 6 (2013), pp. 159–204.
- [29] R. DUAN AND R. M. STRAIN, *Optimal large-time behavior of the Vlasov-Maxwell-Boltzmann system in the whole space*, Comm. Pure Appl. Math., 24 (2011), pp. 1497–1546.
- [30] R. DUAN, T. YANG AND H. ZHAO, *The Vlasov-Poisson-Boltzmann system in the whole space: The hard potential case*, J. Diff. Eq., 252 (2012), pp. 6356–6386.
- [31] R. DUAN, T. YANG AND H. ZHAO, *The Vlasov-Poisson-Boltzmann system for soft potentials*, Math. Models Methods Appl. Sci., 23 (2013), pp. 979–1028.
- [32] F. GOLSE, C. IMBERT, C. MOUHOT AND A. F. VASSEUR, *Harnack inequality for kinetic Fokker-Planck equations with rough coefficients and application to the Landau equation*, Ann. Sc. Norm. Super. Pisa Cl. Sci. 19 (2019), pp. 253–295.
- [33] P. T. GRESSMAN AND R. M. STRAIN, *Global classical solutions of the Boltzmann equation without angular cut-off*, J. Amer. Math. Soc., 24 (2011), pp. 771–847.
- [34] P. T. GRESSMAN AND R. M. STRAIN, *Sharp anisotropic estimates for the Boltzmann collision operator and its entropy production*, Adv. Math., 227 (2011), pp. 2349–2384.
- [35] M. P. GUALDANI, S. MISCHLER AND C. MOUHOT, *Factorization of non-symmetric operators and exponential H-theorem*, Mém. Soc. Math. Fr. (N.S.), 153 (2017).
- [36] Y. GUO, *The Landau equation in a periodic box*, Comm. Math. Phys., 231 (2002), pp. 391–434.
- [37] Y. GUO, *Classical solutions to the Boltzmann equation for molecules with an angular cutoff*, Arch. Ration. Mech. Anal., 169 (2003), pp. 305–353.
- [38] Y. GUO, *The Boltzmann equation in the whole space*, Indiana Univ. Math. J., 53 (2004), pp. 1081–1094.
- [39] Y. GUO, *Boltzmann diffusive limit beyond the Navier-Stokes approximation*, Comm. Pure Appl. Math., 59 (2006), pp. 626–687.
- [40] Y. GUO, *The Vlasov-Poisson-Boltzmann system near vacuum*, Comm. Math. Phys., 218 (2001), pp. 293–313.
- [41] Y. GUO, *The Vlasov-Poisson-Boltzmann system near Maxwellians*, Comm. Pure Appl. Math., 55 (2002), pp. 1104–1135.
- [42] Y. GUO, *The Vlasov-Maxwell-Boltzmann system near Maxwellians*, Invent. Math., 153 (2003), pp. 593–630.
- [43] Y. GUO, *Decay and Continuity of the Boltzmann Equation in Bounded Domains*, Arch. Ration. Mech. Anal., 197 (2009), pp. 713–809.
- [44] Y. GUO, *The Vlasov-Poisson-Landau system in a periodic box*, J. Amer. Math. Soc., 25 (2012), pp. 759–812.
- [45] Y. GUO, H. J. HWANG, J. W. JANG AND Z. OUYANG, *The Landau Equation with the Specular Reflection Boundary Condition*, Arch. Ration. Mech. Anal., 236 (2020), pp. 1389–1454.
- [46] Y. GUO, C. KIM, D. TONON AND A. TRESCASES, *Regularity of the Boltzmann equation in convex domains*, Invent. Math., 207 (2016), pp. 115–290.

- [47] L.-B. HE, *Sharp bounds for Boltzmann and Landau collision operators*, Ann. Sci. Éc. Norm. Supér., 51 (2018), pp. 1253-1341.
- [48] F. HÉRAU, D. TONON AND I. TRISTANI, *Regularization estimates and Cauchy Theory for inhomogeneous Boltzmann equation for hard potentials without cut-off*, Commun. Math. Phys., 377 (2020), pp. 697-771.
- [49] C. HENDERSON AND S. SNELSON, *C^∞ smoothing for weak solutions of the inhomogeneous Landau equation*, Arch. Ration. Mech. Anal., 236 (2020), pp. 113-143.
- [50] C. IMBERT, C. MOUHOT AND L. SILVESTRE, *Decay estimates for large velocities in the Boltzmann equation without cutoff*, Journal de l'École Polytechnique Mathématiques, 7 (2020), pp. 143-184.
- [51] C. IMBERT, C. MOUHOT AND L. SILVESTRE, *Gaussian lower bounds for the Boltzmann equation without cut-off*, SIAM J. Math. Anal., 52 (2020), pp. 2930-2944.
- [52] C. IMBERT AND L. SILVESTRE, *The weak Harnack inequality for the Boltzmann equation without cut-off*, J. Eur. Math. Soc., 22 (2020), pp. 507-592.
- [53] C. IMBERT AND L. SILVESTRE, *Global regularity estimates for the Boltzmann equation without cut-off*, J. Amer. Math. Soc., 35 (2022), pp. 625-703.
- [54] C. IMBERT AND L. SILVESTRE, *The Schauder estimate for kinetic integral equations*, Anal. PDE, 14 (2021), pp. 171-204.
- [55] G. LEONI, *A first course in Sobolev spaces*, American Mathematical Society, Providence, Rhode Island, 2017.
- [56] P.-L. LIONS, *On Boltzmann and Landau equations*, Philos. Trans. Roy. Soc. London Ser. A, 346 (2004), pp. 191-204.
- [57] J. LUK, *Stability of vacuum for the Landau equation with moderately soft potentials*, Ann. PDE, 5 (2019), 11.
- [58] C. MOUHOT AND R. M. STRAIN, *Spectral gap and coercivity estimates for linearized Boltzmann collision operators without angular cutoff*, J. Math. Pures Appl. 87 (2007), pp. 515-535.
- [59] L. NIRENBERG, *On elliptic partial differential equations*, Ann. Sc. Norm. Super. Pisa Cl. Sci., 13 (1959), pp. 115-162.
- [60] L. SILVESTRE, *A new regularization mechanism for the Boltzmann equation without cut-off*, Comm. Math. Phys., 348 (2016), pp. 69-100.
- [61] E. M. STEIN, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J., 1970.
- [62] R. M. STRAIN, *The Vlasov-Maxwell-Boltzmann system in the whole space*, Comm. Math. Phys., 268 (2006), pp. 543-567.
- [63] R. M. STRAIN AND Y. GUO, *Almost exponential decay near Maxwellian*, Comm. Partial Differential Equations, 31 (2006), pp. 417-429.
- [64] R. M. STRAIN AND Y. GUO, *Exponential decay for soft potentials near Maxwellian*, Arch. Ration. Mech. Anal., 187 (2008), pp. 287-339.
- [65] R. M. STRAIN AND K. ZHU, *The Vlasov-Poisson-Landau system in \mathbb{R}_x^3* , Arch. Ration. Mech. Anal., 210 (2013), pp. 615-671.
- [66] C. VILLANI, *On the Cauchy problem for Landau equation: sequential stability, global existence*, Adv. Diff. Eq., 1 (1996), pp. 793-816.
- [67] C. VILLANI, *On a new class of weak solutions to the spatially homogeneous Boltzmann and Landau equations*, Arch. Ration. Mech. Anal., 143 (1998), pp. 273-307.
- [68] Y. WANG, *Global solution and time decay of the Vlasov-Poisson-Landau system in R_x^3* , SIAM J. Math. Anal., 44 (2012), pp. 3281-3323.
- [69] Q. XIAO, L. XIONG AND H. ZHAO, *The Vlasov-Poisson-Boltzmann system for non-cutoff hard potentials*, Sci. China Math., 57 (2014), pp. 515-540.

(Chuji Cao) YAU MATHEMATICAL SCIENCE CENTER AND BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, TSINGHUA UNIVERSITY, BEIJING 100084, P. R. CHINA.

Email address: chuqicao@gmail.com

(Dingqun Deng) DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, POHANG, REPUBLIC OF KOREA (SOUTH), ORCID: 0000-0001-9678-314X

Email address: dingqun.deng@postech.ac.kr

Email address: dingqun.deng@gmail.com

(Xingyu Li) INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ DE TOULOUSE III PAUL SABATIER 118 RTE DE NARBONNE, 31400 TOULOUSE, FRANCE.

Email address: xingyuli92@gmail.com