



HAL
open science

Hyperbolic Wavelet Analysis of Classical Isotropic and Anisotropic Besov–Sobolev Spaces

Béatrice Vedel, Martin Schäfer, Tino Ullrich

► **To cite this version:**

Béatrice Vedel, Martin Schäfer, Tino Ullrich. Hyperbolic Wavelet Analysis of Classical Isotropic and Anisotropic Besov–Sobolev Spaces. *Journal of Fourier Analysis and Applications*, 2021, 27 (3), 10.1007/s00041-021-09844-z . hal-03428666

HAL Id: hal-03428666

<https://hal.science/hal-03428666v1>

Submitted on 5 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hyperbolic wavelet analysis of classical isotropic and anisotropic Besov-Sobolev spaces

Martin Schäfer^{a*}, Tino Ullrich^a, Béatrice Vedel^b

^aChemnitz Technical University, 09111 Chemnitz, Germany

^bUniversité de Bretagne Sud, UMR 6205, LMBA, 56000 Vannes, France

December 18, 2019

Abstract

In this paper we introduce new function spaces which we call anisotropic hyperbolic Besov and Triebel-Lizorkin spaces. Their definition is based on a hyperbolic Littlewood-Paley analysis involving an anisotropy vector only occurring in the smoothness weights. Such spaces provide a general and natural setting in order to understand what kind of anisotropic smoothness can be described using hyperbolic wavelets (in the literature also sometimes called tensor-product wavelets), a wavelet class which hitherto has been mainly used to characterize spaces of dominating mixed smoothness.

A centerpiece of our present work are characterizations of these new spaces based on the hyperbolic wavelet transform. Hereby we treat both, the standard approach using wavelet systems equipped with sufficient smoothness, decay, and vanishing moments, but also the very simple and basic hyperbolic Haar system.

The second major question we pursue is the relationship between the novel hyperbolic spaces and the classical anisotropic Besov-Lizorkin-Triebel scales. As our results show, in general, both approaches to resolve an anisotropy do not coincide. However, in the Sobolev range this is the case, providing a link to apply the newly obtained hyperbolic wavelet characterizations to the classical setting. In particular, this allows for detecting classical anisotropies via the coefficients of a universal hyperbolic wavelet basis, without the need of adaption of the basis or a-priori knowledge on the anisotropy.

Keywords : Hyperbolic wavelet analysis, Hyperbolic Haar wavelet, Anisotropic Sobolev and Besov-Lizorkin-Triebel spaces.

2010 Mathematics Subject Classification : 42C40, 46E35, 42B25, 41A25.

1 Introduction

With the development of wavelet analysis from the beginning of the 1980s until the present time we nowadays have several powerful tools at hand to perform signal analysis with the aim to extract important information out of a signal. The information is thereby usually coded in objects easy to compute and handle – the wavelet coefficients.

Wavelet methods have been used with the known success for the purpose of compression, denoising, inpainting, classification, etc., of data, to mention just a few. Roughly speaking, the

*Corresponding author. Email: martin.schaefer@math.tu-chemnitz.de

common underlying idea is the fact that a few wavelet coefficients contain a rather complete information of the signal to be analyzed. However, due to their construction principle (dyadic dilations and integer translates of a few basic “mother” functions) classical wavelets are not well-suited for the analysis of, say, anisotropic signals. In fact, a signal which is rather smooth in x -direction but rough in y -direction (such as layers in the earth, stripes on a shirt, etc.) can not be properly resolved by a classical multi-resolution analysis. The respective wavelet coefficients do not contain the anisotropic smoothness information, they rather resolve a certain minimal smoothness. That results in a bad decay of the sequence of wavelet coefficients or, in other words, a bad compression rate.

Anisotropy is not a rare phenomenon since it arises whenever physics does not act the same in different directions, e.g., geophysics, oceanography, hydrology, fluid mechanics, or medical image processing (see [4, 42] among others) are some of the fields where it naturally appears. For this reason wavelets have been adapted in many different ways in order to “detect” and resolve anisotropy. There is a vast amount of literature dealing with this. For instance, there are wave atoms [13] as well as curvelets [8, 9, 7], shearlets [32, 28, 34], anisets, and anisotropic wavelets [50, 51, 30]. The latter concept represents a rather flexible construction since it can be build (theoretically) for any present anisotropy. The theoretical basis of anisotropic wavelet analysis is the equivalent characterization of corresponding anisotropic function spaces, like Hölder, Besov, Sobolev and Triebel-Lizorkin spaces. The major shortcoming of the existing theory is the fact that one has to know the anisotropy in advance, i.e., one has to adapt the wavelet accordingly. In other words, if physics does not provide the anisotropy parameters of the signal we are not able to resolve the signal accordingly without “trying out” several anisotropic bases. Such a method is, of course, hardly implementable in practice.

In Abry et. al. [1] it has been shown that any anisotropic Besov space – defined with respect to the cartesian axis – can “almost” be characterized with the help of the so-called hyperbolic wavelet transform. The anisotropy of the signal can then be detected using a uniform basis and is characterized by a special weight in the wavelet coefficients. This has led to an efficient algorithm for image classification and anisotropy detection applied to both synthetic and real textures (see [41, 2]).

In this paper we further develop this idea of describing anisotropy with the help of the hyperbolic wavelet transform. For this reason we introduce a new family of anisotropic function spaces which are defined via a hyperbolic Littlewood-Paley analysis and for which we prove exact characterization with hyperbolic wavelets. The motivation behind this is to provide a general setting of anisotropic spaces characterized by one single basis of wavelets and thus to understand how one such fixed basis can help to describe anisotropic smoothness.

Concretely, we start with a hyperbolic Littlewood-Paley analysis defined as the usual tensor product

$$(1) \quad \Delta_{\vec{j}}(f) := \mathcal{F}^{-1}[\theta_{j_1} \otimes \dots \otimes \theta_{j_d} \mathcal{F}f] \quad , \quad \vec{j} = (j_1, \dots, j_d) \in \mathbb{N}_0^d.$$

This hyperbolic decomposition of the frequency space has been widely used for the Fourier analytic definition of the well-known spaces with dominating mixed smoothness, see [43, 57] and the references therein. These spaces represent a suitable framework for multivariate approximation, see [15, 48] and the recent survey article [11]. The main ideas have been developed over more than fifty years of intense research in the former Soviet Union such that it is beyond the scope of this paper to name all the relevant references (cf. [11]).

Based on the decomposition (1), we then define spaces $\hat{A}_{p,q}^{s,\vec{\alpha}}(\mathbb{R}^d)$ with $A \in \{B, F\}$ of Besov-Lizorkin-Triebel type involving an anisotropy vector $\vec{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ with $\sum_{i=1}^d \alpha_i = d$.

As a special case ($A = F$, $1 < p < \infty$, $q = 2$), these include the Sobolev type spaces $\widetilde{W}_p^{s,\bar{\alpha}}(\mathbb{R}^d) := \widetilde{F}_{p,2}^{s,\bar{\alpha}}(\mathbb{R}^d)$, where

$$\|f\|_{\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} := \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{qs\|\vec{j}/\bar{\alpha}\|_\infty} |\Delta_{\vec{j}}(f)(\cdot)|^q \right)^{1/q} \right\|_p, \quad f \in \mathcal{S}'(\mathbb{R}^d).$$

It is important to note that the anisotropy hereby only enters in the weight $2^{s\|\vec{j}/\bar{\alpha}\|_\infty}$, where we use the short-hand notation $\|\vec{j}/\bar{\alpha}\|_\infty := \max\{j_1/\alpha_1, \dots, j_d/\alpha_d\}$, but not in the choice of the Littlewood-Paley decomposition.

One of the main results of this paper is the coincidence (with respect to equivalent norms)

$$(2) \quad \widetilde{W}_p^{s,\bar{\alpha}}(\mathbb{R}^d) = W_p^{s,\bar{\alpha}}(\mathbb{R}^d), \quad \text{if } 1 < p < \infty,$$

where the space on the right-hand side represents the classical anisotropic Sobolev space defined in (3) below. This relation has already been observed for isotropic (i.e. $\bar{\alpha} = (1, \dots, 1)$) Hilbert-Sobolev spaces ($p = 2$) on the d -torus, see [27, 12], as well as on \mathbb{R}^2 in [1]. Our result extends this observation to all $1 < p < \infty$. Surprisingly, such a coincidence in the spirit of (2) is only possible in the Sobolev case. To be more precise, it holds

$$\widetilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = A_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) \quad \text{if and only if } A = F, 1 < p < \infty, \text{ and } q = 2.$$

As an important consequence of this equality (2), we can further prove that it is possible to characterize (e.g. detect and classify) classical anisotropies described by the spaces $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ via the wavelet coefficients of a universal hyperbolic wavelet basis. Compared to the classical approach using anisotropic wavelets, this approach has the advantage that one does not need a-priori knowledge on the anisotropies, otherwise required for constructing the “right” basis. In particular, our results entail that any sufficiently regular orthonormal basis $(\psi_{\vec{j},\vec{k}})_{\vec{j},\vec{k}}$ of tensorized wavelets $\psi_{\vec{j},\vec{k}} = \psi_{j_1,k_1} \otimes \dots \otimes \psi_{j_d,k_d}$ constitutes an unconditional Schauder basis for $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$, whose coefficients, measured in an appropriate corresponding sequence space, give rise to an equivalent norm on $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$, i.e. for $f \in W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ with coefficients $\langle f, \psi_{\vec{j},\vec{k}} \rangle$

$$\|f\|_{W_p^{s,\bar{\alpha}}(\mathbb{R}^d)} \asymp \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{2s\|\vec{j}/\bar{\alpha}\|_\infty} \left| \sum_{\vec{k} \in \mathbb{Z}^d} \langle f, \psi_{\vec{j},\vec{k}} \rangle \chi_{\vec{j},\vec{k}}(\cdot) \right|^2 \right)^{1/2} \right\|_p.$$

This is stated in Theorem 6.2. A similar result, see Theorem 6.3, holds true for the hyperbolic Haar system $\mathcal{H}_d = (h_{\vec{j},\vec{k}})_{\vec{j},\vec{k}}$, where $h_{\vec{j},\vec{k}} = h_{j_1,k_1} \otimes \dots \otimes h_{j_d,k_d}$, under the following restriction on the parameter s of the space $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$,

$$|s|/\alpha_{\min} < \min \left\{ \frac{1}{p}, 1 - \frac{1}{p} \right\}.$$

In this direction, we would also like to mention the new and related findings of Oswald in [37] on the Schauder basis property of the hyperbolic Haar system in the classic isotropic Besov spaces defined via first-order moduli of smoothness.

At the center of our respective proofs, we will rely on discrete characterizations provided by hyperbolic wavelets for the spaces $\widetilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$, $A \in \{B, F\}$. These characterizations are fundamental and established in separate theorems, Theorem 4.2 and Theorem 5.4, whereby we follow two paths. On the one hand, we use the usual methodology and consider orthonormal

wavelet bases for which we assume sufficient smoothness, decay, and vanishing moments. As a byproduct, we thereby significantly extend the wavelet characterizations in [57, 55] for Besov-Lizorkin-Triebel spaces with dominating mixed smoothness. On the other hand, we use a hyperbolic Haar system, which does not fulfill smoothness conditions as before but nevertheless allows for characterization in a certain restricted parameter range.

Let us remark that analysis with the Haar wavelet has a long tradition (see e.g. [25, 38, 39, 40, 3]), the Haar wavelet being the oldest and simplest orthonormal wavelet, conceived as early as 1909 [29]. Besides its elegance and simplicity, notably its connection to the Faber system [17] and other spline functions, such as e.g. the Chui-Wang wavelet [10], makes it interesting from a numerical perspective. In particular in imaging science it plays an important role in practical applications. Recently, it has attracted renewed attention with a series of publications [44, 45, 21, 22, 23, 24, 14].

The paper has the following **structure**. After having recalled in Section 2 some helpful Fourier analytic tools (in particular some classical maximal functions and associated inequalities) as well as the definition of the classical (anisotropic) function spaces $A_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$, where $A \in \{B, F\}$, we introduce in Section 3 the notion of hyperbolic Littlewood-Paley analysis and the related Besov-Lizorkin-Triebel spaces $\tilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. Wavelet characterizations of these new hyperbolic spaces are the topic of Sections 4 and 5, whereby we first resort to standard wavelets with sufficient smoothness, decay, and vanishing moments in Section 4, while in Section 5 we utilize a hyperbolic Haar basis. The relationship of the new scale to the traditional spaces is finally investigated in Sections 6 and 7. Specifically, in Section 6, we show the equality $\tilde{W}_p^{s,\bar{\alpha}}(\mathbb{R}^d) = W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$, i.e. $\tilde{F}_{p,2}^{s,\bar{\alpha}}(\mathbb{R}^d) = F_{p,2}^{s,\bar{\alpha}}(\mathbb{R}^d)$, in the range $1 < p < \infty$, from which we can then extract our main theorems concerning hyperbolic wavelet characterizations of the classical $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$.

Let us agree on the following general **notation**. As usual \mathbb{N} shall denote the natural numbers. We further put $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$, and let \mathbb{Z} denote the integers, \mathbb{R} the real numbers, and \mathbb{C} the complex numbers. By $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$ we refer to the torus identified with the interval $[0, 2\pi] \subset \mathbb{R}$. We write $\langle x, y \rangle$ or $x \cdot y$ for the Euclidean inner product in \mathbb{R}^d or \mathbb{C}^d . The letter d is hereby always reserved for the underlying dimension and by $[d]$ we mean the set $\{1, \dots, d\}$. For $0 < p \leq \infty$ and $x \in \mathbb{R}^d$ we define $\|x\|_p := (\sum_{i=1}^d |x_i|^p)^{1/p}$, with the usual modification in the case $p = \infty$. If $1 \leq p \leq \infty$ we set p' such that $1/p + 1/p' = 1$. For $0 < p, q \leq \infty$ we further denote $\sigma_{p,q} := \max\{1/p - 1, 1/q - 1, 0\}$ and $\sigma_p := \max\{1/p - 1, 0\}$. We also put $x_+ := ((x_1)_+, \dots, (x_d)_+)$, whereby $a_+ := \max\{a, 0\}$ for $a \in \mathbb{R}$. Analogously we define x_- . By $(x_1, \dots, x_d) > 0$ we shall mean that each coordinate is positive. Finally, as usual, $a \in \mathbb{R}$ is decomposed into $a = [a] + \{a\}$, where $0 \leq \{a\} < 1$ and $[a] \in \mathbb{Z}$. In case $x \in \mathbb{R}^d$, $\{x\}$ and $[x]$ are then meant component-wise. Multivariate indices are typesetted with a bar, like e.g. $\bar{k}, \bar{j}, \bar{\ell}$, or \bar{m} , to indicate the multi-index. In all the paper, the multi-index $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ thereby stands for an anisotropy and is such that $\alpha_1 + \dots + \alpha_d = d$. In addition, we here use the abbreviations $\alpha_{\min} := \min\{\alpha_1, \dots, \alpha_d\}$ and $\alpha_{\max} := \max\{\alpha_1, \dots, \alpha_d\}$. The notation $\bar{\alpha}/\bar{j}$ shall always stand for $(\alpha_1/j_1, \dots, \alpha_d/j_d)$. Given a positive real $a > 0$, we further write $a^{\bar{\alpha}}$ for the vector $(a^{\alpha_1}, \dots, a^{\alpha_d})$ and let $f(a^{\bar{\alpha}}x) := f(a^{\alpha_1}x_1, \dots, a^{\alpha_d}x_d)$ be the anisotropically scaled version of the function $f : \mathbb{R}^d \rightarrow \mathbb{C}$. For two (quasi-)normed spaces X and Y , the (quasi-)norm of an element $x \in X$ will be denoted by $\|x\|_X$. The symbol $X \hookrightarrow Y$ indicates that the identity operator is continuous. For two sequences a_n and b_n we will write $a_n \lesssim b_n$ if there exists a constant $c > 0$ such that $a_n \leq c b_n$ for all n . We will write $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $b_n \lesssim a_n$.

2 Classical spaces and tools from Fourier analysis

Let $L_p = L_p(\mathbb{R}^d)$, $0 < p \leq \infty$, be the Lebesgue space of all measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{C}$ such that

$$\|f\|_p := \left(\int_{\mathbb{R}^d} |f(x)|^p dx \right)^{1/p} < \infty,$$

with the usual modification if $p = \infty$. We will also need L_p -spaces on compact domains $\Omega \subset \mathbb{R}^d$ instead of \mathbb{R}^d and shall write $\|f\|_{L_p(\Omega)}$ for the corresponding restricted L_p -(quasi-)norms.

For $k \in \mathbb{N}_0$, we denote by $C_0^k(\mathbb{R}^d)$ the collection of all compactly supported functions φ on \mathbb{R}^d which have uniformly continuous derivatives $D^{\bar{\gamma}}\varphi$ on \mathbb{R}^d whenever $\|\bar{\gamma}\|_1 \leq k$. Additionally, we define the spaces of infinitely differentiable functions $C^\infty(\mathbb{R}^d)$ and infinitely differentiable functions with compact support $C_0^\infty(\mathbb{R}^d)$ as well as the Schwartz space $\mathcal{S} = \mathcal{S}(\mathbb{R}^d)$ of all rapidly decaying infinitely differentiable functions on \mathbb{R}^d , i.e.,

$$\mathcal{S}(\mathbb{R}^d) := \{ \varphi \in C^\infty(\mathbb{R}^d) : \|\varphi\|_{k,\ell} < \infty \text{ for all } k, \ell \in \mathbb{N} \},$$

and

$$\|\varphi\|_{k,\ell} := \left\| (1 + |\cdot|)^k \sum_{\|\bar{\gamma}\|_1 \leq \ell} |D^{\bar{\gamma}}\varphi(\cdot)| \right\|_\infty, \quad k, \ell \in \mathbb{N}.$$

The space $\mathcal{S}'(\mathbb{R}^d)$, the topological dual of $\mathcal{S}(\mathbb{R}^d)$, is also referred to as the space of tempered distributions on \mathbb{R}^d . Indeed, a linear mapping $f : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathbb{C}$ belongs to $\mathcal{S}'(\mathbb{R}^d)$ if and only if there exist numbers $k, \ell \in \mathbb{N}$ and a constant $c = c_f$ such that

$$|f(\varphi)| \leq c_f \|\varphi\|_{k,\ell}$$

for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$. Any locally integrable function f on \mathbb{R}^d belongs to $\mathcal{S}'(\mathbb{R}^d)$ in the sense that

$$f(\varphi) := \int_{\mathbb{R}^d} f(x)\varphi(x) dx, \quad \varphi \in \mathcal{S}(\mathbb{R}^d).$$

The space $\mathcal{S}'(\mathbb{R}^d)$ is equipped with the weak*-topology.

For $f \in L_1(\mathbb{R}^d)$ we define the Fourier transform

$$\mathcal{F}f(\xi) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(y)e^{-i\xi \cdot y} dy, \quad \xi \in \mathbb{R}^d,$$

and the corresponding inverse Fourier transform $\mathcal{F}^{-1}f(\xi) = \mathcal{F}f(-\xi)$. As usual, the Fourier transform can be extended to $\mathcal{S}'(\mathbb{R}^d)$ by $(\mathcal{F}f)(\varphi) := f(\mathcal{F}\varphi)$, where $f \in \mathcal{S}'(\mathbb{R}^d)$ and $\varphi \in \mathcal{S}(\mathbb{R}^d)$. The mapping $\mathcal{F} : \mathcal{S}'(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ is a bijection.

The convolution $\varphi * \psi$ of two square-integrable functions φ, ψ is defined via the integral

$$(\varphi * \psi)(x) = \int_{\mathbb{R}^d} \varphi(x-y)\psi(y) dy.$$

If $\varphi, \psi \in \mathcal{S}(\mathbb{R}^d)$ then $\varphi * \psi$ still belongs to $\mathcal{S}(\mathbb{R}^d)$. In fact, we have $\varphi * \psi \in \mathcal{S}(\mathbb{R}^d)$ even if $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and $\psi \in L_1(\mathbb{R}^d)$. The convolution can be extended to $\mathcal{S}(\mathbb{R}^d) \times \mathcal{S}'(\mathbb{R}^d)$ via $(\varphi * \psi)(x) = \psi(\varphi(x - \cdot))$, which makes sense pointwise and is a C^∞ -function on \mathbb{R}^d .

2.1 Classical (an)isotropic Littlewood-Paley analysis

Subsequently, $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ will denote an anisotropy and be such that $\alpha_1 + \dots + \alpha_d = d$. Anisotropic Besov spaces may then be introduced using an anisotropic Littlewood-Paley analysis depending on $\bar{\alpha}$. Classical isotropic spaces – as a particular case of anisotropic spaces – will thereby be obtained for $\bar{\alpha} = (1, 1, \dots, 1)$.

Let $\varphi_0^{\bar{\alpha}} \geq 0$ belong to the Schwartz class $\mathcal{S}(\mathbb{R}^d)$ and be such that, for $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$,

$$\varphi_0^{\bar{\alpha}}(\xi) = 1 \quad \text{if} \quad \sup_{i=1,2,\dots,d} |\xi_i| \leq 1,$$

$$\text{and} \quad \varphi_0^{\bar{\alpha}}(\xi) = 0 \quad \text{if} \quad \sup_{i=1,\dots,d} |2^{-\alpha_i} \xi_i| \geq 1.$$

For $j \in \mathbb{N}$, further define

$$\begin{aligned} \varphi_j^{\bar{\alpha}}(\xi) &:= \varphi_0^{\bar{\alpha}}(2^{-j\bar{\alpha}}\xi) - \varphi_0^{\bar{\alpha}}(2^{-(j-1)\bar{\alpha}}\xi) \\ &= \varphi_0^{\bar{\alpha}}(2^{-j\alpha_1}\xi_1, \dots, 2^{-j\alpha_d}\xi_d) - \varphi_0^{\bar{\alpha}}(2^{-(j-1)\alpha_1}\xi_1, \dots, 2^{-(j-1)\alpha_d}\xi_d). \end{aligned}$$

Then $\sum_{j=0}^{\infty} \varphi_j^{\bar{\alpha}} \equiv 1$, and $(\varphi_j^{\bar{\alpha}})_{j \geq 0}$ is called an *anisotropic resolution of unity*. It satisfies

$$\text{supp}(\varphi_0^{\bar{\alpha}}) \subset R_1^{\bar{\alpha}}, \quad \text{supp}(\varphi_j^{\bar{\alpha}}) \subset R_{j+1}^{\bar{\alpha}} \setminus R_{j-1}^{\bar{\alpha}},$$

$$\text{where} \quad R_j^{\bar{\alpha}} = \left\{ \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d : |\xi_i| \leq 2^{\alpha_i j} \text{ for } i \in [d] = \{1, \dots, d\} \right\}.$$

For $f \in \mathcal{S}'(\mathbb{R}^d)$, we then define

$$\Delta_j^{\bar{\alpha}} f = \mathcal{F}^{-1}(\varphi_j^{\bar{\alpha}} \mathcal{F} f).$$

The sequence $(\Delta_j^{\bar{\alpha}} f)_{j \geq 0}$ is called an *anisotropic Littlewood-Paley analysis* of f . With this tool, the anisotropic Besov spaces are now defined as follows (see [6, 5]).

Definition 2.1 For $0 < p \leq \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, the Besov space $B_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ is defined by

$$B_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \left(\sum_{j \geq 0} 2^{jsq} \|\Delta_j^{\bar{\alpha}} f\|_p^q \right)^{1/q} < \infty \right\},$$

with the usual modification for $q = \infty$.

This definition does not depend on chosen resolution of unity $\varphi_0^{\bar{\alpha}}$ and the quantity

$$\|f\|_{B_{p,q}^{s,\bar{\alpha}}} = \left(\sum_{j \geq 0} 2^{jsq} \|\Delta_j^{\bar{\alpha}} f\|_p^q \right)^{1/q}$$

is a norm (resp. quasi-norm) on $B_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ for $1 \leq p, q \leq \infty$ (resp. $0 < \min\{p, q\} < 1$) and with the usual modification if $q = \infty$.

As in the isotropic case, anisotropic Besov spaces encompass a large class of classical anisotropic function spaces (see [51] for details). For example, when $p = q = 2$, the Besov spaces coincide with the anisotropic Sobolev spaces and, when $p = q = \infty$, the spaces $B_{\infty,\infty}^{s,\bar{\alpha}}(\mathbb{R}^d)$ are called anisotropic Hölder spaces and are denoted by $\mathcal{C}^{s,\bar{\alpha}}(\mathbb{R}^d)$.

Definition 2.2 For $0 < p \leq \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, the Triebel-Lizorkin space $F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ is defined by

$$F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \left\| \left(\sum_{j \geq 0} 2^{jsq} |\Delta_j^{\bar{\alpha}} f(\cdot)|^q \right)^{1/q} \right\|_p < \infty \right\},$$

with the usual modification for $q = \infty$.

This definition does not depend on the chosen resolution of unity $\varphi_0^{\bar{\alpha}}$ and the quantity

$$\|f\|_{F_{p,q}^{s,\bar{\alpha}}} = \left\| \left(\sum_{j \geq 0} 2^{jsq} |\Delta_j^{\bar{\alpha}} f(\cdot)|^q \right)^{1/q} \right\|_p$$

is a norm (resp. quasi-norm) on $F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ for $1 \leq p < \infty$ and $1 \leq q \leq \infty$ (resp. $0 < \min\{p, q\} < 1$) and with the usual modification if $q = \infty$.

If $q = 2$ and $1 < p < \infty$, the anisotropic Triebel-Lizorkin space coincides with the anisotropic Sobolev space denoted by $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$:

$$(3) \quad W_p^{s,\bar{\alpha}} = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \left\| \mathcal{F}^{-1} \left[\left(\sum_{i=1}^d (1 + \xi_i^2)^{1/2\alpha_i} \right)^s \mathcal{F}f(\xi) \right] \right\|_p < \infty \right\}.$$

Remark 2.1 (i) As mentioned before, if $\bar{\alpha} = (1, \dots, 1)$, it is easy to check that the spaces $B_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ (resp. $F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$) coincide with the classical spaces $B_{p,q}^s(\mathbb{R}^d)$ (resp. $F_{p,q}^s(\mathbb{R}^d)$). In addition, we have $F_{p,2}^{0,\bar{\alpha}}(\mathbb{R}^d) = L_p(\mathbb{R}^d)$ in the range $1 < p < \infty$.

(ii) Our understanding of anisotropic spaces coincides with the one in Triebel [51] (see also the references therein). There are different (but related) notions of anisotropy in the Russian literature, see Nikolskij [36, Chapt. 4] or Temlyakov [48, II.3]. A consequence of our Theorem 6.1 below is the fact that in case of W -spaces the mentioned approaches coincide and lead to the same notion of anisotropy. However, in case of Hölder-Nikolskij spaces this is in general not the case as for instance Theorem 7.1 shows.

2.2 Maximal inequalities

Let us provide here the maximal inequalities for the Hardy-Littlewood and Peetre maximal functions, respectively. For further details we refer to [57, 1.2, 1.3] or [43, Chapt. 2].

For a locally integrable function $f : \mathbb{R}^d \rightarrow \mathbb{C}$ we denote by $Mf(x)$ the Hardy-Littlewood maximal function defined by

$$(4) \quad (Mf)(x) = \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f(y)| dy \quad , \quad x \in \mathbb{R}^d ,$$

where the supremum is taken over all cubes with sides parallel to the coordinate axes containing x . A vector valued generalization of the classical Hardy-Littlewood maximal inequality is due to Fefferman and Stein [18].

Theorem 2.1 ([18]) For $1 < p < \infty$ and $1 < q \leq \infty$ there exists a constant $c > 0$, such that

$$\left\| \left(\sum_{\ell \in I} |Mf_\ell|^q \right)^{1/q} \right\|_p \leq c \left\| \left(\sum_{\ell \in I} |f_\ell|^q \right)^{1/q} \right\|_p$$

holds for all sequences $\{f_\ell\}_{\ell \in I}$ of locally Lebesgue-integrable functions on \mathbb{R}^d .

We require a direction-wise version of (4)

$$(M_i f)(x) = \sup_{s>0} \frac{1}{2s} \int_{x_i-s}^{x_i+s} |f(x_1, \dots, x_{i-1}, t, x_{i+1}, \dots, x_d)| dt \quad , \quad x \in \mathbb{R}^d.$$

We denote the composition of these operators by $M_e = \prod_{i \in e} M_i$, where e is a subset of $[d] = \{1, \dots, d\}$ and $M_\ell M_k$ has to be interpreted as $M_\ell \circ M_k$. The following version of the Fefferman-Stein maximal inequality is due to Stöckert [47].

Theorem 2.2 ([47]) For $1 < p < \infty$ and $1 < q \leq \infty$ there exists a constant $c > 0$ such that for any $i \in [d]$

$$\left\| \left(\sum_{\ell \in I} |M_i f_\ell|^q \right)^{1/q} \right\|_p \leq c \left\| \left(\sum_{\ell \in I} |f_\ell|^q \right)^{1/q} \right\|_p$$

holds for all sequences $\{f_\ell\}_{\ell \in I}$ of locally Lebesgue-integrable functions on \mathbb{R}^d .

Iteration of this theorem yields a similar boundedness property for the operator $M_{[d]}$.

The following construction of a maximal function is due to Peetre, Fefferman, and Stein. Let $\bar{b} = (b_1, \dots, b_d) > 0$, $a > 0$, and $f \in L_1(\mathbb{R}^d)$ with $\mathcal{F}f$ compactly supported. We define the Peetre maximal function $P_{\bar{b}, a} f$ by

$$(5) \quad P_{\bar{b}, a} f(x) = \sup_{z \in \mathbb{R}^d} \frac{|f(x-z)|}{(1 + |b_1 z_1|)^a \cdot \dots \cdot (1 + |b_d z_d|)^a} \quad .$$

Lemma 2.1 Let $\Omega \subset \mathbb{R}^d$ be a compact set. Let further $a > 0$ and $\bar{\gamma} = (\gamma_1, \dots, \gamma_d) \in \mathbb{N}_0^d$. Then there exist two constants $c_1, c_2 > 0$ (independent of f) such that

$$(6) \quad \begin{aligned} P_{(1, \dots, 1), a}(D^{\bar{\gamma}} f)(x) &\leq c_1 P_{(1, \dots, 1), a} f(x) \\ &\leq c_2 (M_d(M_{d-1}(\dots(M_1|f|^{1/a})\dots)))^a(x) \end{aligned}$$

holds for all $f \in L_1(\mathbb{R}^d)$ with $\text{supp}(\mathcal{F}f) \subset \Omega$ and all $x \in \mathbb{R}^d$. The constants c_1, c_2 depend on Ω .

We finally give a vector-valued version of the Peetre maximal inequality which is a direct consequence of Lemma 2.1 together with Theorem 2.2.

Theorem 2.3 Let $0 < p < \infty$, $0 < q \leq \infty$ and $a > \max\{1/p, 1/q\}$. Let further $\bar{b}^\ell = (b_1^\ell, \dots, b_d^\ell) > 0$ for $\ell \in I$ and $\Omega = \{\Omega_\ell\}_{\ell \in I}$, such that

$$\Omega_\ell \subset [-b_1^\ell, b_1^\ell] \times \dots \times [-b_d^\ell, b_d^\ell]$$

is compact for $\ell \in I$. Then there is a constant $C > 0$ (independent of f and Ω) such that

$$\left\| \left(\sum_{\ell \in I} |P_{\bar{b}^\ell, a} f_\ell|^q \right)^{1/q} \right\|_p \leq C \left\| \left(\sum_{\ell \in I} |f_\ell|^q \right)^{1/q} \right\|_p$$

holds for all systems $f = \{f_\ell\}_{\ell \in I}$ with $\text{supp}(\mathcal{F}f_\ell) \subset \Omega_\ell$, $\ell \in I$.

3 Hyperbolic Littlewood-Paley analysis

Let $\theta_0 \in \mathcal{S}(\mathbb{R})$ be supported on $[-2, 2]$ with $\theta_0 = 1$ on $[-1, 1]$. For any $j \in \mathbb{N}$, let us further define

$$\theta_j = \theta_0(2^{-j}\cdot) - \theta_0(2^{-(j-1)}\cdot)$$

such that $(\theta_j)_j$ is a univariate resolution of unity, i.e., $\sum_{j \geq 0} \theta_j(\cdot) = 1$. Observe that, for any $j \geq 1$, $\text{supp}(\theta_j) \subset \{2^{j-1} \leq |\xi| \leq 2^{j+1}\}$.

Remark 3.1 *In the following, the function θ_0 can be chosen with an arbitrary compact support. It does not change the main results even if technical details of proofs and lemmas have to be adapted. This allows to choose θ_0 as the Fourier transform of a Meyer scaling function.*

Let us now come to the main concept of this paper, the hyperbolic Littlewood-Paley analysis.

Definition 3.1 (i) For any $\bar{j} = (j_1, \dots, j_d) \in \mathbb{N}_0^d$ and any $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$ set

$$\theta_{\bar{j}}(\xi_1, \dots, \xi_d) := \theta_{j_1}(\xi_1)\theta_{j_2}(\xi_2)\dots\theta_{j_d}(\xi_d).$$

The function $\theta_{\bar{j}}$ belongs to $\mathcal{S}(\mathbb{R}^d)$ for all $\bar{j} \in \mathbb{N}_0^d$ and is compactly supported on a dyadic rectangle. Further $\sum_{\bar{j} \in \mathbb{N}_0^d} \theta_{\bar{j}} \equiv 1$ and $(\theta_{\bar{j}})_{\bar{j}}$ is called a hyperbolic resolution of unity.

(ii) For $f \in \mathcal{S}'(\mathbb{R}^d)$ and $\bar{j} \in \mathbb{N}_0^d$ set

$$\Delta_{\bar{j}} f := \mathcal{F}^{-1}(\theta_{\bar{j}} \mathcal{F} f).$$

The sequence $(\Delta_{\bar{j}} f)_{\bar{j} \in \mathbb{N}_0^d}$ is called a hyperbolic Littlewood-Paley analysis of f .

We are now in the position to introduce new functional spaces called *anisotropic hyperbolic Besov spaces* and *anisotropic hyperbolic Triebel-Lizorkin spaces* defined via the hyperbolic Littlewood-Paley analysis.

Definition 3.2 For $0 < p \leq \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$ we define the anisotropic hyperbolic Besov space $\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ via

$$\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_{\infty} s q} \|\Delta_{\bar{j}} f\|_p^q \right)^{1/q} < \infty \right\},$$

with the usual modification in case $q = \infty$.

This definition does not depend on the chosen resolution of unity $(\theta_{\bar{j}})_{\bar{j}}$ and the quantity

$$\|f\|_{\tilde{B}_{p,q}^{s,\bar{\alpha}}} := \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_{\infty} s q} \|\Delta_{\bar{j}} f\|_p^q \right)^{1/q}$$

is a norm (resp. quasi-norm) on $\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ for $1 \leq p, q \leq \infty$ (resp. $0 < \min\{p, q\} < 1$) and with usual modification if $q = \infty$.

Definition 3.3 For $0 < p < \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_i \alpha_i = d$ we define the anisotropic hyperbolic Triebel-Lizorkin space via

$$\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_\infty sq} |\Delta_{\vec{j}} f(\cdot)|^q \right)^{1/q} \right\|_p < \infty \right\},$$

with the usual modification in case $q = \infty$.

This definition does not depend on the chosen resolution of unity $(\theta_{\vec{j}})_{\vec{j}}$ and the quantity

$$\|f\|_{\widetilde{F}_{p,q}^{s,\bar{\alpha}}} := \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_\infty sq} |\Delta_{\vec{j}} f(\cdot)|^q \right)^{1/q} \right\|_p$$

is a norm on $\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ for $1 \leq p < \infty$, $1 \leq q \leq \infty$ (resp. quasi-norm for $0 < \min\{p, q\} < 1$).

Remark 3.2 The above definitions of anisotropic hyperbolic Besov and Sobolev spaces include four indices: s stands for the regularity, p is the integration parameter and q the so-called fine-index. The parameter $\bar{\alpha} = (\alpha_1, \dots, \alpha_d)$ encodes the present anisotropy: the more $\alpha_{\min} = \min\{\alpha_1, \dots, \alpha_d\}$ is close to 0 and $\alpha_{\max} = \max\{\alpha_1, \dots, \alpha_d\}$ is close to d , the more we need directional smoothness in one axis compared to others. On the other hand, if $\bar{\alpha} = (1, \dots, 1)$ the anisotropy becomes an “isotropy”.

Remark 3.3 By analogy with the classical spaces, if $q = 2$ and $1 < p < \infty$, $\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ is called anisotropic hyperbolic Sobolev space and is denoted by $\widetilde{W}_p^{s,\bar{\alpha}}(\mathbb{R}^d)$. In case $\bar{\alpha} = (1, \dots, 1)$ we write $\widetilde{W}_p^s(\mathbb{R}^d)$.

Let us finally introduce classical spaces with dominating mixed smoothness in the spirit of [43, 57].

Definition 3.4 Let $r \in \mathbb{R}$, $0 < p, q \leq \infty$ ($p < \infty$ in the F -case).

(i) The Besov space with dominating mixed smoothness $S_{p,q}^r B(\mathbb{R}^d)$ is the collection of all distributions $f \in \mathcal{S}'(\mathbb{R}^d)$ such that the following (quasi-)norm

$$\|f\|_{S_{p,q}^r B(\mathbb{R}^d)} := \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{r\|j\|_1 q} \|\Delta_{\vec{j}}(f)\|_p^q \right)^{1/q} \quad \text{is finite.}$$

(ii) The Triebel-Lizorkin space with dominating mixed smoothness $S_{p,q}^r F(\mathbb{R}^d)$ is the collection of all distributions $f \in \mathcal{S}'(\mathbb{R}^d)$ such that the following (quasi-)norm

$$\|f\|_{S_{p,q}^r F(\mathbb{R}^d)} := \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{r\|j\|_1 q} |\Delta_{\vec{j}}(f)(x)|^q \right)^{1/q} \right\|_p \quad \text{is finite.}$$

(iii) If $1 < p < \infty$ and $r \in \mathbb{R}$ then the Sobolev space with dominating mixed smoothness $S_p^r W(\mathbb{R}^d)$ is the collection of all $f \in \mathcal{S}'(\mathbb{R}^d)$ such that

$$\|f\|_{S_p^r W} := \left\| \mathcal{F}^{-1} \left[\prod_{i=1}^d (1 + |\xi_i|^2)^{r/2} \mathcal{F} f \right] \right\|_p \quad \text{is finite.}$$

Let us also state a useful Fourier multiplier theorem, see [57, Thm. 1.12] or [43, p. 77].

Lemma 3.1 ([57, 43]) *Let $0 < p < \infty$, $0 < q \leq \infty$, and $r > \frac{1}{\min(p,q)} + \frac{1}{2}$. Further, let $\{\Omega_{\bar{k}}\}_{\bar{k} \in \mathbb{N}_0^d}$ be a sequence of compact subsets of \mathbb{R}^d such that*

$$\Omega_{\bar{k}} \subset \{x \in \mathbb{R}^d : |x_i| \leq 2^{k_i}, i = 1, \dots, d\}.$$

Then, there is a positive constant $C > 0$ such that

$$\left\| \left(\sum_{\bar{k} \in \mathbb{N}_0^d} \left| \mathcal{F}^{-1}[\rho_{\bar{k}} \mathcal{F} f_{\bar{k}}](\cdot) \right|^q \right)^{\frac{1}{q}} \right\|_p \leq C \left\| \left(\sum_{\bar{k} \in \mathbb{N}_0^d} |f_{\bar{k}}(\cdot)|^q \right)^{\frac{1}{q}} \right\|_p \times \sup_{\bar{\ell} \in \mathbb{N}_0^d} \|\rho_{\bar{\ell}}(2^{\ell_1} \cdot, \dots, 2^{\ell_d} \cdot)\|_{S_2^r W}$$

holds for all systems $\{f_{\bar{k}}\}_{\bar{k} \in \mathbb{N}_0^d} \in L_p(\ell_q)$ with $\text{supp}(\mathcal{F} f_{\bar{k}}) \subset \Omega_{\bar{k}}$ and all systems $\{\rho_{\bar{k}}\}_{\bar{k} \in \mathbb{N}_0^d} \subset S_2^r W(\mathbb{R}^d)$.

4 Hyperbolic wavelet analysis

In this section we prove hyperbolic wavelet characterizations of the spaces $\widetilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ defined in Definitions 3.2 and 3.3, respectively. It should be noted that the proof technique used for Theorem 4.2 below also represents a progress towards new optimal wavelet characterizations of Besov-Lizorkin-Triebel spaces with dominating mixed smoothness, which extends the results in [57, Sect. 2.4] significantly, see Remark 4.3 below.

Let us start with univariate wavelets given by a scaling function ψ_0 and a corresponding wavelet ψ . These functions are supposed to satisfy the following (minimal) conditions:

(K) It holds $\psi_0, \psi \in C^K(\mathbb{R})$. For any $M \in \mathbb{N}$ there is a constant $C_M > 0$ such that for all $0 \leq \alpha \leq K$ it holds

$$|D^\alpha \psi_0(x)| + |D^\alpha \psi(x)| \leq C_M (1 + |x|)^{-M}, \quad x \in \mathbb{R}.$$

(L) The wavelet ψ has vanishing moments up to order $L - 1$: For $L > \beta \in \mathbb{N}_0$ it holds

$$\int_{\mathbb{R}} \psi(x) x^\beta dx = 0.$$

In case $L = 0$ the condition is void.

We shall denote

$$\psi_{j,k} := \frac{1}{\sqrt{2}} \psi(2^{j-1} \cdot - k), \quad j \in \mathbb{N}, k \in \mathbb{Z},$$

and $\psi_{0,k} := \psi_0(\cdot - k)$. We set $\psi_{j,k} \equiv 0$ if $j < 0$. To obtain the hyperbolic wavelet basis in $L_2(\mathbb{R}^d)$ we tensorize over all scales and obtain

$$\psi_{\bar{j},\bar{k}}(x_1, \dots, x_d) := \psi_{j_1, k_1}(x_1) \cdot \dots \cdot \psi_{j_d, k_d}(x_d), \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d, \bar{j} \in \mathbb{Z}^d, \bar{k} \in \mathbb{Z}^d.$$

The following lemma recalls a useful convolution relation. Let us clarify the notation first. For a given univariate function Λ we will use the notation $\Lambda_j(\cdot) := 2^{j-1} \Lambda(2^{j-1} \cdot)$, $j \in \mathbb{N}$. We will further put $x_{j,m} := 2^{-j} m$ and $I_{j,m} := [2^{-j} m, 2^{-j}(m+1))$ with associated characteristic function $\chi_{j,m} := \mathbb{1}_{I_{j,m}}$.

Lemma 4.1 *Let $\Lambda_0, \Lambda \in \mathcal{S}(\mathbb{R})$ with Λ having infinitely many vanishing moments, i.e.,*

$$\int_{\mathbb{R}} \Lambda(x) x^\beta dx = 0$$

for all $\beta \in \mathbb{N}$. Let further ψ_0 and ψ satisfy (K) and (L) as above and $R > 0$ be a given real number. Then it exists a constant $C_R > 0$ such that for any $j \in \mathbb{N}_0$ and $\ell, m \in \mathbb{Z}$ the convolution relation

$$|(\Lambda_j * \psi_{j+\ell, m})(x)| \leq C_R 2^{-N_{\text{sign}(\ell)}|\ell|} (1 + 2^{\min\{j, j+\ell\}} |x - x_{j+\ell, m}|)^{-R}$$

holds true with $\text{sign}(\ell) \in \{+, -, 0\}$ and $N_0 := 0$, $N_+ := L + 1$ and $N_- := K$.

Proof. The above lemma is a special case of a more general convolution relation, see for instance [26, p. 466] for the most general version. Originally, this relation is due to Frazier, Jawerth [19, Lem. 3.3], [20, Lem. B.1, B.2]. \square

Lemma 4.1 immediately implies the following multivariate version by exploiting the tensor product structure. Similar as for the hyperbolic wavelet system, we use the notation

$$\Lambda_{\bar{j}}(x) := \Lambda_{j_1}(x_1) \cdot \dots \cdot \Lambda_{j_d}(x_d) \quad , \quad x \in \mathbb{R}^d, \bar{j} \in \mathbb{Z}^d.$$

In the sequel we will further need the notation

$$(7) \quad Q_{\bar{j}, \bar{m}} := I_{j_1, m_1} \times \dots \times I_{j_d, m_d} \quad \text{and} \quad \chi_{\bar{j}, \bar{m}}(x_1, \dots, x_d) := \chi_{j_1, m_1}(x_1) \cdot \dots \cdot \chi_{j_d, m_d}(x_d),$$

with the notation I_{j_i, m_i} and χ_{j_i, m_i} , $i \in [d] = \{1, \dots, d\}$, introduced right before Lemma 4.1.

Lemma 4.2 *Let $\Lambda, \Lambda_0, \psi_0, \psi$ as in Lemma 4.1. For any $R > 0$ there exists a constant $C_R > 0$ such that for any $\bar{j} \in \mathbb{N}_0^d$ and $\bar{\ell}, \bar{m} \in \mathbb{Z}^d$ the convolution relation*

$$(8) \quad |(\Lambda_{\bar{j}} * \psi_{\bar{j}+\bar{\ell}, \bar{m}})(x)| \leq C_R \prod_{i=1}^d 2^{-N_{\text{sign}(\ell_i)}|\ell_i|} (1 + 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m_i|)^{-R}$$

holds true with $\text{sign}(\ell_i) \in \{+, -, 0\}$ and $N_0 := 0$, $N_+ := L + 1$ and $N_- := K$.

The next proposition is also crucial and represents the ‘‘hyperbolic version’’ of [31, Lem. 3,7]. An isotropic version is originally due to Kyriazis [33, Lem. 7.1]. For the convenience of the reader we give a proof.

Proposition 4.1 *Let $0 < r \leq 1$ and $R > 1/r$. For any sequence $(\lambda_{\bar{j}})_{\bar{j} \in \mathbb{N}_0^d}$ of complex numbers and any $\bar{\ell} \in \mathbb{Z}^d, \bar{j} \in \mathbb{N}_0^d$ we have, using the notation $\bar{\ell}_+ = ((\ell_1)_+, \dots, (\ell_d)_+)$,*

$$\begin{aligned} & \sum_{\bar{m} \in \mathbb{Z}^d} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}| \prod_{i=1}^d \left(1 + 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m_i|\right)^{-R} \\ & \leq 2^{\|\bar{\ell}_+\|_1/r} \left[M \left| \sum_{\bar{m} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell}, \bar{m}} \chi_{\bar{j}+\bar{\ell}, \bar{m}} \right|^r \right]^{1/r} (x) \quad , \quad x \in \mathbb{R}^d, \end{aligned}$$

where M stands for the Hardy-Littlewood maximal operator.

Proof. We follow the proof from [31, Lem. 7]. Put $\delta = R - 1/r > 0$ and define a decomposition $\{\Omega_{\bar{k}}(x)\}_{\bar{k} \in \mathbb{N}_0^d}$ of \mathbb{Z}^d depending on $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ as follows:

$$\Omega_{\bar{k}}(x) := \Omega_{k_1}(x_1) \times \dots \times \Omega_{k_d}(x_d) \quad , \quad \bar{k} = (k_1, \dots, k_d) \in \mathbb{N}_0^d,$$

with

$$\begin{aligned} \Omega_{k_i}(x_i) &:= \{m \in \mathbb{Z} : 2^{k_i-1} < 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m| \leq 2^{k_i}\} \quad , \quad k_i \in \mathbb{N}, \\ \Omega_0(x_i) &:= \{m \in \mathbb{Z} : 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m| \leq 1\}. \end{aligned}$$

We then estimate for fixed $x = (x_1, \dots, x_d) \in \mathbb{R}^d$

$$\begin{aligned} & \sum_{\bar{m} \in \mathbb{Z}^d} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}| \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m_i|)^{-R} \\ (9) \quad &= \sum_{\bar{k} \in \mathbb{N}_0^d} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}| \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |x_i - 2^{-(j_i+\ell_i)} m_i|)^{-R} \\ &\lesssim \sum_{\bar{k} \in \mathbb{N}_0^d} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}| 2^{-\delta \|\bar{k}\|_1 - \|\bar{k}\|_1/r} \lesssim \sup_{\bar{k} \in \mathbb{N}_0^d} \left(\sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}| \right) 2^{-\|\bar{k}\|_1/r} \\ &\lesssim \left(\sup_{\bar{k} \in \mathbb{N}_0^d} 2^{-\|\bar{k}\|_1} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}|^r \right)^{1/r}. \end{aligned}$$

We further note that

$$(10) \quad 2^{-\|\bar{k}\|_1} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}|^r = 2^{-\|\bar{k}\|_1} \int_{\bigcup_{\bar{m} \in \Omega_{\bar{k}}(x)} Q_{\bar{j}+\bar{\ell}, \bar{m}}} 2^{\|\bar{j}+\bar{\ell}\|_1} \sum_{\bar{w} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{w}}|^r \chi_{\bar{j}+\bar{\ell}, \bar{w}}(y) dy$$

and observe that for $Q(x) := \bigcup_{\bar{m} \in \Omega_{\bar{k}}(x)} Q_{\bar{j}+\bar{\ell}, \bar{m}}$ we have $x \in Q(x)$ and

$$|Q(x)| \asymp 2^{\|\bar{k}\|_1 - \|\min\{\bar{j}, \bar{j}+\bar{\ell}\}\|_1}.$$

Recalling the definition of the Hardy-Littlewood maximal function in (4), we obtain

$$\begin{aligned} & 2^{-\|\bar{k}\|_1 + \|\min\{\bar{j}, \bar{j}+\bar{\ell}\}\|_1} \int_{Q(x)} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}|^r \chi_{\bar{j}+\bar{\ell}, \bar{m}}(y) dy \\ & \lesssim \frac{1}{|Q(x)|} \int_{Q(x)} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}|^r \chi_{\bar{j}+\bar{\ell}, \bar{m}}(y) dy \leq M \left| \sum_{\bar{m} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell}, \bar{m}} \chi_{\bar{j}+\bar{\ell}, \bar{m}} \right|^r(x). \end{aligned}$$

Putting this into (10), we arrive at

$$2^{-\|\bar{k}\|_1} \sum_{\bar{m} \in \Omega_{\bar{k}}(x)} |\lambda_{\bar{j}+\bar{\ell}, \bar{m}}|^r \lesssim 2^{\|\bar{\ell}\|_1} M \left| \sum_{\bar{m} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell}, \bar{m}} \chi_{\bar{j}+\bar{\ell}, \bar{m}} \right|^r(x).$$

Finally, we plug this estimate into (9) and obtain the desired assertion. \square

Before stating our main result we need a further definition.

Definition 4.1 Let $0 < q \leq \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$.

(i) If $0 < p < \infty$ we define the sequence space $\tilde{f}_{p,q}^{s,\bar{\alpha}}$ as the collection of all sequences $(\lambda_{\bar{j},\bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d} \subset \mathbb{C}$ such that the (quasi-)norm (usual modification in case $q = \infty$)

$$\|(\lambda_{\bar{j},\bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d}\|_{\tilde{f}_{p,q}^{s,\bar{\alpha}}} := \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_\infty s q} \left| \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j},\bar{k}} \chi_{\bar{j},\bar{k}}(\cdot) \right|^q \right)^{1/q} \right\|_p \quad \text{is finite.}$$

(ii) If $0 < p \leq \infty$ we define the sequence space $\tilde{b}_{p,q}^{s,\bar{\alpha}}$ as the collection of all sequences $(\lambda_{\bar{j},\bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d} \subset \mathbb{C}$ such that the (quasi-)norm (usual modification in case $q = \infty$)

$$\|(\lambda_{\bar{j},\bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d}\|_{\tilde{b}_{p,q}^{s,\bar{\alpha}}} := \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|(j_1/\alpha_1, \dots, j_d/\alpha_d)\|_\infty s q} \left\| \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j},\bar{k}} \chi_{\bar{j},\bar{k}}(\cdot) \right\|_p^q \right)^{1/q} \quad \text{is finite.}$$

Now we are ready to state the wavelet characterization of the space $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. Recall that for $0 < p, q \leq \infty$ we put

$$\sigma_{p,q} := \max\{1/p - 1, 1/q - 1, 0\} \quad \text{and} \quad \sigma_p := \max\{1/p - 1, 0\}.$$

Remark 4.1 The theorem below states the result for the F -scale of spaces $\tilde{F}_{p,q}^{a,\bar{\alpha}}(\mathbb{R}^d)$. As for the corresponding result for the Besov type spaces $\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$, we simply replace condition (11) on K, L by

$$K, L > \sigma_p + |s|/\alpha_{\min}$$

and use the corresponding sequence spaces $\tilde{b}_{p,q}^{s,\bar{\alpha}}$.

Theorem 4.2 Let $0 < p < \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ with $\sum_{i=1}^d \alpha_i = d$. Let further ψ_0, ψ be wavelets satisfying (K) and (L) above with

$$(11) \quad K, L > \sigma_{p,q} + |s|/\alpha_{\min}.$$

Then any $f \in \mathcal{S}'(\mathbb{R}^d)$ belongs to $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ if and only if it can be represented as

$$(12) \quad f = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j},\bar{k}} \psi_{\bar{j},\bar{k}}$$

with $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$ and the sum converging in $\mathcal{S}'(\mathbb{R}^d)$ with respect to some ordering. For each $f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ the convergence of the representation (12) is then even unconditional. Moreover, if $q < \infty$, the sum also converges in $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $(\psi_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ constitutes an unconditional basis in $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. The sequence of coefficients $\lambda(f) := (\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ is uniquely determined via

$$(13) \quad \lambda_{\bar{j},\bar{k}} = 2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j},\bar{k}} \rangle$$

and we have the wavelet isomorphism (equivalent (quasi-)norm)

$$\|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} \asymp \|\lambda(f)\|_{\tilde{f}_{p,q}^{s,\bar{\alpha}}} \quad , \quad f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d).$$

Remark 4.2 Following [35, Prop. 3.20], the dual pairing of $f \in \mathcal{S}'(\mathbb{R}^d)$ and $\psi_{\bar{j}, \bar{k}} \in C^K(\mathbb{R}^d)$ in (13) has to be understood in the way

$$(14) \quad \langle f, \psi \rangle := \sum_{\bar{j} \in \mathbb{N}_0^d} \langle \Theta_{\bar{j}} * f, \Lambda_{\bar{j}} * \psi \rangle_{L_2(\mathbb{R}^d)}.$$

Here we choose $\Theta_{\bar{j}} := \mathcal{F}^{-1} \theta_{\bar{j}}$ and $\Lambda_{\bar{j}} := \mathcal{F}^{-1} \lambda_{\bar{j}}$ such that $\sum_{j=0}^{\infty} \theta_j \lambda_j \equiv 1$. Using elementary estimates and the Nikolskij inequality in case $p < 1$, one can show

$$\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) \hookrightarrow \tilde{B}_{p,\infty}^{s,\bar{\alpha}}(\mathbb{R}^d) \hookrightarrow S_{\max\{p,1\},\infty}^{-|s|/\alpha_{\min}-\sigma_p} B(\mathbb{R}^d).$$

Setting $s_{\bar{\alpha},p} := |s|/\alpha_{\min} + \sigma_p$ and $\tilde{p} := \max\{p, 1\}$ we obtain

$$\begin{aligned} |\langle f, \psi \rangle| &\leq \sum_{\bar{j} \in \mathbb{N}_0^d} \|\Theta_{\bar{j}} * f\|_{\tilde{p}} \|\Lambda_{\bar{j}} * \psi\|_{\tilde{p}'} \\ &\leq \sup_{\bar{j} \in \mathbb{N}_0^d} 2^{-s_{\bar{\alpha},p} \|\bar{j}\|_1} \|\Theta_{\bar{j}} * f\|_{\tilde{p}} \cdot \sum_{\bar{j} \in \mathbb{N}_0^d} 2^{s_{\bar{\alpha},p} \|\bar{j}\|_1} \|\Lambda_{\bar{j}} * \psi\|_{\tilde{p}'} \\ &\lesssim \|f\|_{S_{\tilde{p},\infty}^{-s_{\bar{\alpha},p}} B(\mathbb{R}^d)} \cdot \|\psi\|_{S_{\tilde{p}',1}^{s_{\bar{\alpha},p}} B(\mathbb{R}^d)}, \end{aligned}$$

where the right-hand side is finite due to (11) and (8). In other words, $f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ generates a (conjugate) linear functional on the Banach space $S_{\tilde{p}',1}^{s_{\bar{\alpha},p}} B(\mathbb{R}^d)$.

Remark 4.3 As we will see below, our arguments apply as well to classical spaces of dominating mixed smoothness $S_{p,q}^r B(\mathbb{R}^d)$ and $S_{p,q}^r F(\mathbb{R}^d)$, defined in Definition 3.4 above. Examining the proof, we obtain for the relation

$$\|f\|_{S_{p,q}^r F(\mathbb{R}^d)} \lesssim \|\lambda(f)\|_{s_{p,q}^r f},$$

where $s_{p,q}^r f$ is the sequence space associated to $S_{p,q}^r F$ (for a definition see [57, Def. 2.1]), the condition

$$(15) \quad L > \sigma_{p,q} - r \quad \text{and} \quad K > r.$$

The converse relation holds under the condition

$$K > \sigma_{p,q} - r \quad \text{and} \quad L > r.$$

For the spaces $S_{p,q}^r B(\mathbb{R}^d)$ we replace $\sigma_{p,q}$ by σ_p and $s_{p,q}^r f$ by $s_{p,q}^r b$, which is the sequence space associated to $S_{p,q}^r B$ (for a definition see [57, Def. 2.1]).

Proof. [of Theorem 4.2] *Step 1.* We consider the sum

$$(16) \quad f := \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}, \bar{k}} \psi_{\bar{j}, \bar{k}}$$

with $\lambda := (\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}} \in \tilde{F}_{p,q}^{s,\bar{\alpha}}$ and show the relation

$$\|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} \lesssim \|\lambda\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}}.$$

For the issues on the convergence and uniqueness of (16) and (13) we refer to Step 3 and 4 below, where we show that under the assumption $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$ the element f is well defined, with unconditional convergence of (16) at least in $\mathcal{S}'(\mathbb{R}^d)$, which is sufficient for the subsequent considerations.

Let us consider $\Delta_{\bar{j}}(f)$ for some chosen hyperbolic Littlewood-Paley analysis. This gives for fixed $\bar{j} \in \mathbb{N}_0^d$

$$2^{\|\bar{j}/\bar{\alpha}\|_\infty s} |\Delta_{\bar{j}}(f)(x)| \leq \sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \left| \Theta_{\bar{j}} * \left(\sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \psi_{\bar{j}+\bar{\ell},\bar{k}} \right) (x) \right|,$$

where $\Theta_{\bar{j}} := \mathcal{F}^{-1} \theta_{\bar{j}}$ and $(\theta_{\bar{j}})_{\bar{j}}$ is the system from Definition 3.1. With $u := \min\{p, q, 1\}$

$$(17) \quad \begin{aligned} \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} &= \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \Delta_{\bar{j}}(f)(\cdot) \right\|_{L_p(\ell_q)} \\ &\lesssim \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \Theta_{\bar{j}} * \left(\sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \psi_{\bar{j}+\bar{\ell},\bar{k}} \right) (\cdot) \right\|_{L_p(\ell_q[\bar{j}])}^u \right)^{1/u}. \end{aligned}$$

With the help of Lemma 4.2 we are aiming for pointwise estimates first.

$$\begin{aligned} \left| \Theta_{\bar{j}} * \left(\sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \psi_{\bar{j}+\bar{\ell},\bar{k}} \right) (x) \right| &\leq \sum_{\bar{k} \in \mathbb{Z}^d} |\lambda_{\bar{j}+\bar{\ell},\bar{k}}| \cdot |(\Theta_{\bar{j}} * \psi_{\bar{j}+\bar{\ell},\bar{k}})(x)| \\ &\lesssim \left(\prod_{i=1}^d 2^{-N_{\text{sign}(\ell_i)} |\ell_i|} \right) \sum_{\bar{k} \in \mathbb{Z}^d} |\lambda_{\bar{j}+\bar{\ell},\bar{k}}| \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i + \ell_i\}} |x_i - 2^{-(j_i + \ell_i)} k_i|)^{-R}, \end{aligned}$$

where we choose $R > 1/r$ with $r < \min\{1, p, q\} = u$. Note that, due to condition (K) for the wavelet and Lemma 4.2, we can choose $R > 0$ arbitrarily large. This allows for estimating with the help of Proposition 4.1

$$\left| \Theta_{\bar{j}} * \left(\sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \psi_{\bar{j}+\bar{\ell},\bar{k}} \right) (x) \right| \lesssim 2^{-\langle \bar{N}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{\|\bar{\ell}\|_1 / r} \left[M \left| \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \chi_{\bar{j}+\bar{\ell},\bar{k}}(\cdot) \right|^r \right]^{1/r} (x).$$

Hereby we use the short-hand notation $\bar{\ell}_+ := ((\ell_1)_+, \dots, (\ell_d)_+)$ and

$$\langle \bar{N}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle := \sum_{i=1}^d N_{\text{sign}(\ell_i)} |\ell_i|.$$

Plugging this estimate into (17) gives

$$\begin{aligned} \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} &\lesssim \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-u \langle \bar{N}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{u \|\bar{\ell}_+\|_1 / r} \right. \\ &\quad \left. \left\| 2^{-\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \left[M \left| 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \chi_{\bar{j}+\bar{\ell},\bar{k}} \right|^r \right]^{1/r} (\cdot) \right\|_{L_p(\ell_q[\bar{j}])}^u \right)^{\frac{1}{u}}. \end{aligned}$$

Clearly, if $s \geq 0$ then $\|(\bar{j} + \bar{\ell})/\bar{\alpha}\|_\infty s \geq \|\bar{j}/\bar{\alpha}\|_\infty s - \|\bar{\ell}/\bar{\alpha}\|_\infty s$ and hence

$$(18) \quad 2^{-\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \leq 2^{\|\bar{\ell}/\bar{\alpha}\|_\infty |s|}.$$

In addition, if $s < 0$ we also obtain (18) via the usual ℓ_∞ -triangle inequality.

Putting this into the previous estimate, using the vector-valued Hardy-Littlewood maximal inequality (Theorem 2.1) for the space $L_{p/r}(\ell_{q/r}[\bar{j}])$, we obtain

$$\begin{aligned} \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} &\lesssim \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-u\langle \bar{N}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{u\|\bar{\ell}_+\|_1/r} 2^{u\|\bar{\ell}/\bar{\alpha}\|_\infty|s|} \right. \\ &\quad \left. \left\| 2^{(\bar{j}+\bar{\ell})/\bar{\alpha}} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}+\bar{\ell},\bar{k}} \chi_{\bar{j}+\bar{\ell},\bar{k}} \right\|_{L_p(\ell_q[\bar{j}])}^u \right)^{1/u} \\ &\lesssim \|\lambda\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-u\langle \bar{N}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{u\|\bar{\ell}_+\|_1/r} 2^{u\|\bar{\ell}/\bar{\alpha}\|_\infty|s|} \right)^{1/u}. \end{aligned}$$

The sum over $\bar{\ell}$ converges if $L+1 = N_+ > 1/r + |s|/\alpha_{\min}$ and $K = N_- > |s|/\alpha_{\min}$.

Step 2. Let us prove the converse relation $\|\lambda(f)\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} \lesssim \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)}$ with $\lambda(f) = (2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j},\bar{k}} \rangle)_{\bar{j},\bar{k}}$ and start with $f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. As already pointed out in Remark 4.2, the dual pairing $\langle f, \psi_{\bar{j},\bar{k}} \rangle$ makes sense due to condition (11). Our estimation begins as follows

$$\begin{aligned} |2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j},\bar{k}} \rangle| &\leq \sum_{\bar{\ell} \in \mathbb{Z}^d} |2^{\|\bar{j}\|_1} \langle \Theta_{\bar{j}+\bar{\ell}} * f, \Lambda_{\bar{j}+\bar{\ell}} * \psi_{\bar{j},\bar{k}} \rangle| \\ (19) \quad &= \sum_{\bar{\ell} \in \mathbb{Z}^d} \sum_{\bar{m} \in \mathbb{Z}^d} \left| \int_{Q_{\bar{j}+\bar{\ell},\bar{m}}} (\Theta_{\bar{j}+\bar{\ell}} * f)(y) 2^{\|\bar{j}\|_1} (\Lambda_{\bar{j}+\bar{\ell}} * \psi_{\bar{j},\bar{k}})(y) dy \right| \\ &\leq \sum_{\bar{\ell} \in \mathbb{Z}^d} \sum_{\bar{m} \in \mathbb{Z}^d} |\theta_{\bar{j}+\bar{\ell},\bar{m}}(f)| \int_{Q_{\bar{j}+\bar{\ell},\bar{m}}} 2^{\|\bar{j}\|_1} |(\Lambda_{\bar{j}+\bar{\ell}} * \psi_{\bar{j},\bar{k}})(y)| dy, \end{aligned}$$

where we put

$$(20) \quad \theta_{\bar{j}+\bar{\ell},\bar{m}}(f) := \sup_{y \in Q_{\bar{j}+\bar{\ell},\bar{m}}} |(\Theta_{\bar{j}+\bar{\ell}} * f)(y)|.$$

We next estimate the integral, as in Step 1 with the help of Lemma 4.2. Here we have to be particularly careful with the normalization factors. Note that, compared to (8), the signs of the components of $\bar{\ell}$ in the convolution $\Lambda_{\bar{j}+\bar{\ell}} * \psi_{\bar{j},\bar{k}}$ change the role. This is why we put this time $M_0 := 0$, $M_- := L$, $M_+ := K+1$ and $\bar{M}_{\text{sign}(\bar{\ell})} := (M_{\text{sign}(b_1)}, \dots, M_{\text{sign}(b_d)})$. We then obtain for $z = (z_1, \dots, z_d) \in Q_{\bar{j},\bar{k}}$ the estimate

$$\begin{aligned} &\int_{Q_{\bar{j}+\bar{\ell},\bar{m}}} 2^{\|\bar{j}\|_1} |(\Lambda_{\bar{j}+\bar{\ell}} * \psi_{\bar{j},\bar{k}})(y)| dy \\ &\lesssim \sup_{y \in Q_{\bar{j}+\bar{\ell},\bar{m}}} 2^{-\langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |y_i - 2^{-j_i} k_i|)^{-R} \\ &\lesssim 2^{-\langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |2^{-(j_i+\ell_i)} m_i - 2^{-j_i} k_i|)^{-R} \\ &\lesssim 2^{-\langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |2^{-(j_i+\ell_i)} m_i - z_i|)^{-R}. \end{aligned}$$

This together with (19) and Proposition 4.1 yields for $r < \min\{p, q, 1\} = u$ and $z \in \mathbb{R}^d$

$$\begin{aligned}
& \sum_{\bar{k} \in \mathbb{Z}^d} |2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j}, \bar{k}} \rangle| \chi_{\bar{j}, \bar{k}}(z) \\
& \lesssim \sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-\langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} \sum_{\bar{m} \in \mathbb{Z}^d} |\theta_{\bar{j}+\bar{\ell}, \bar{m}}(f)| \prod_{i=1}^d (1 + 2^{\min\{j_i, j_i+\ell_i\}} |2^{-(j_i+\ell_i)} m_i - z_i|)^{-R} \\
& \lesssim \sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-\langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{\|\ell_+\|_1/r} \left[M \left| \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}+\bar{\ell}, \bar{m}}(f) \chi_{\bar{j}+\bar{\ell}, \bar{m}}(\cdot) \right|^r \right]^{1/r}(z).
\end{aligned}$$

This leads to

$$\begin{aligned}
& 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \sum_{\bar{k} \in \mathbb{Z}^d} |2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j}, \bar{k}} \rangle| \chi_{\bar{j}, \bar{k}} \\
& \lesssim \sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{\|\ell_+\|_1/r - \langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle + \|\bar{j}/\bar{\alpha}\|_\infty s - \|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} \left[M \left| 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty} \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}+\bar{\ell}, \bar{m}}(f) \chi_{\bar{j}+\bar{\ell}, \bar{m}}(\cdot) \right|^r \right]^{\frac{1}{r}}.
\end{aligned}$$

Taking the $L_p(\ell_q[\bar{j}])$ -(quasi)-norm on both sides and using (18) once more, we obtain

$$\begin{aligned}
(21) \quad & \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \sum_{\bar{k} \in \mathbb{Z}^d} |2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j}, \bar{k}} \rangle| \chi_{\bar{j}, \bar{k}} \right\|_{L_p(\ell_q[\bar{j}])} \\
& \lesssim \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-u \langle \bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}| \rangle} 2^{u\|\ell_+\|_1/r} 2^{u\|\bar{\ell}/\bar{\alpha}\|_\infty |s|} \times \right. \\
& \quad \left. \times \left\| \left[M \left| 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty} \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}+\bar{\ell}, \bar{m}}(f) \chi_{\bar{j}+\bar{\ell}, \bar{m}}(\cdot) \right|^r \right]^{1/r} \right\|_{L_p(\ell_q[\bar{j}])}^u \right)^{1/u}.
\end{aligned}$$

Due to $r < \min\{1, p, q\} = u$ we can apply the Hardy-Littlewood maximal inequality (Theorem 2.1) and obtain

$$\left\| \left[M \left| 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty} \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}+\bar{\ell}, \bar{m}}(f) \chi_{\bar{j}+\bar{\ell}, \bar{m}} \right|^r \right]^{1/r} \right\|_{L_p(\ell_q[\bar{j}])} \lesssim \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}, \bar{m}}(f) \chi_{\bar{j}, \bar{m}} \right\|_{L_p(\ell_q)}.$$

From (20) we obtain for any $z \in Q_{\bar{j}, \bar{m}}$ and any $a > 0$

$$|\theta_{\bar{j}, \bar{m}}(f)| = \sup_{y \in Q_{\bar{j}, \bar{m}}} |(\Theta_{\bar{j}} * f)(y)| \lesssim \sup_{y \in \mathbb{R}^d} \frac{|(\Theta_{\bar{j}} * f)(y)|}{\prod_{i=1}^d (1 + 2^{j_i} |z_i - y_i|)^a} = P_{2^{\bar{j}}, a}(\Theta_{\bar{j}} * f)(z),$$

where we used the definition of the Peetre maximal function in (5). Choosing $a > \max\{\frac{1}{p}, \frac{1}{q}\}$ with the corresponding maximal inequality in Theorem 2.3 then yields the relation

$$\begin{aligned}
\left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \sum_{\bar{m} \in \mathbb{Z}^d} \theta_{\bar{j}, \bar{m}}(f) \chi_{\bar{j}, \bar{m}} \right\|_{L_p(\ell_q)} & \lesssim \|P_{2^{\bar{j}}, a}(2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \Theta_{\bar{j}} * f)\|_{L_p(\ell_q)} \\
& \lesssim \|2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \Theta_{\bar{j}} * f\|_{L_p(\ell_q)} \asymp \|f\|_{\tilde{F}_{p, q}^{s, \bar{\alpha}}(\mathbb{R}^d)}.
\end{aligned}$$

Returning to (21), we have seen

$$\begin{aligned} & \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} \sum_{\bar{k} \in \mathbb{Z}^d} |2^{\|\bar{j}\|_1} \langle f, \psi_{\bar{j}, \bar{k}} \rangle | \chi_{\bar{j}, \bar{k}} \right\|_{L_p(\ell_q[\bar{j}])} \\ & \lesssim \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} \left(\sum_{\bar{\ell} \in \mathbb{Z}^d} 2^{-u(\bar{M}_{\text{sign}(\bar{\ell})}, |\bar{\ell}|)} 2^{u\|\ell_+\|_1/r} 2^{u\|\bar{\ell}/\bar{\alpha}\|_\infty |s|} \right)^{1/u}. \end{aligned}$$

It remains to discuss the sum over $\bar{\ell}$. It is easy to see that it converges if $K + 1 = M_+ > 1/r + |s|/\alpha_{\min}$ and $L = M_- > |s|/\alpha_{\min}$. Recall that r is chosen such that $r < \min\{1, p, q\}$.

Step 3. Let us now clarify the convergence issues in (12) in case $q < \infty$. The arguments in Step 1 above show in particular for a finite partial summation of (12) that

$$\left\| \sum_{\bar{j}} \sum_{\bar{k}} \lambda_{\bar{j}, \bar{k}} \psi_{\bar{j}, \bar{k}} \right\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} \lesssim \left\| \left(\sum_{\bar{j}} 2^{\|\bar{j}/\bar{\alpha}\|_\infty s q} \left| \sum_{\bar{k}} \lambda_{\bar{j}, \bar{k}} \chi_{\bar{j}, \bar{k}} \right|^q \right)^{1/q} \right\|_p.$$

If $q < \infty$ (note that $p < \infty$ anyway) we use Lebesgue's dominated convergence theorem to conclude the unconditional convergence of (12) in $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. The required majorant is thereby given by

$$\|(\lambda_{\bar{j}, \bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d}\|_{\tilde{f}_{p,q}^{s,\bar{\alpha}}} < \infty.$$

In case $q = \infty$ we use the observation in Remark 4.3. From a simple application of Hölder's inequality (with respect to the sum over \bar{j}) we first obtain for any $\varepsilon > 0$ the relation

$$(22) \quad \|(\lambda_{\bar{j}, \bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d}\|_{S_{p,1}^{r(s,\bar{\alpha})-\varepsilon}} f \lesssim \|(\lambda_{\bar{j}, \bar{k}})_{\bar{j} \in \mathbb{N}_0^d, \bar{k} \in \mathbb{Z}^d}\|_{\tilde{f}_{p,\infty}^{s,\bar{\alpha}}} \quad \text{with } r(s, \bar{\alpha}) := -|s|/\alpha_{\min}.$$

Choosing $\varepsilon > 0$ small enough, we then obtain from (11) that condition (15) in Remark 4.3 is satisfied. Hence, for a finite partial summation of (12) we have

$$\left\| \sum_{\bar{j}} \sum_{\bar{k}} \lambda_{\bar{j}, \bar{k}} \psi_{\bar{j}, \bar{k}} \right\|_{S_{p,1}^{r(s,\bar{\alpha})-\varepsilon} F(\mathbb{R}^d)} \lesssim \left\| \sum_{\bar{j}} 2^{(r(s,\bar{\alpha})-\varepsilon)\|\bar{j}\|_1} \sum_{\bar{k}} |\lambda_{\bar{j}, \bar{k}}| \chi_{\bar{j}, \bar{k}} \right\|_p.$$

Again, by Lebesgue's dominated convergence theorem (the majorant given by (22)) we see the unconditional convergence of (12) in the space $S_{p,1}^{r(s,\bar{\alpha})-\varepsilon} F(\mathbb{R}^d)$. Taking the embedding $S_{p,1}^{r(s,\bar{\alpha})-\varepsilon} F(\mathbb{R}^d) \hookrightarrow \mathcal{S}'(\mathbb{R}^d)$ into account, we actually proved more than stated in the theorem.

Step 4. It remains to prove (12) for $f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and coefficients $\lambda_{\bar{j}, \bar{k}}(f)$ chosen as in (13). From Steps 1, 2, 3 above we have learned that $\{\lambda_{\bar{j}, \bar{k}}(f)\}_{\bar{j}, \bar{k}} \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$, which implies that the sum

$$\sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}, \bar{k}}(f) \psi_{\bar{j}, \bar{k}}$$

converges (at least in) $\mathcal{S}'(\mathbb{R}^d)$ to an element $g \in \mathcal{S}'(\mathbb{R}^d)$. We now prove that $f(\varphi) = g(\varphi)$ for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$. Fix $\varphi \in \mathcal{S}(\mathbb{R}^d)$, then clearly $\bar{\varphi} \in L_2(\mathbb{R}^d)$ and we have

$$(23) \quad \bar{\varphi} = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \langle \bar{\varphi}, \psi_{\bar{j}, \bar{k}} \rangle \psi_{\bar{j}, \bar{k}}$$

with convergence in $L_2(\mathbb{R}^d)$. Since $\bar{\varphi} \in S_{\bar{p}',1}^{s\bar{\alpha},p} B(\mathbb{R}^d)$ we have by Step 1, 2, 3 above that the right-hand side of (23) converges in $S_{\bar{p}',1}^{s\bar{\alpha},p} B(\mathbb{R}^d)$ to some $\eta \in S_{\bar{p}',1}^{s\bar{\alpha},p} B(\mathbb{R}^d)$. Hence, we have

$\bar{\varphi} = \eta$ in $\mathcal{S}'(\mathbb{R}^d)$ which finally gives $\bar{\varphi} = \eta$ almost everywhere and, in other words, (23) holds true in $S_{\bar{p}',1}^{s,\bar{\alpha}}B(\mathbb{R}^d)$. Then $f(\varphi)$ can be rewritten as follows, using the continuity of $\langle f, \cdot \rangle$ (see Remark 4.2),

$$\begin{aligned} f(\varphi) &= \langle f, \bar{\varphi} \rangle = \left\langle f, \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \langle \bar{\varphi}, \psi_{\bar{j},\bar{k}} \rangle \psi_{\bar{j},\bar{k}} \right\rangle \\ &= \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \overline{\langle \bar{\varphi}, \psi_{\bar{j},\bar{k}} \rangle} \langle f, \psi_{\bar{j},\bar{k}} \rangle = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \langle \varphi, \overline{\psi_{\bar{j},\bar{k}}} \rangle \langle f, \psi_{\bar{j},\bar{k}} \rangle. \end{aligned}$$

On the other hand,

$$g(\varphi) = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j},\bar{k}}(f) \psi_{\bar{j},\bar{k}}(\varphi) = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \langle \varphi, \overline{\psi_{\bar{j},\bar{k}}} \rangle \langle f, \psi_{\bar{j},\bar{k}} \rangle = f(\varphi),$$

which finishes the proof. \square

5 Hyperbolic Haar characterization

We next utilize a hyperbolic Haar basis for the characterization of the spaces $\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ from Definitions 3.2 and 3.3, the main result being Theorem 5.4. It will show that Haar characterizations are possible in a certain restricted range of parameters, although the Haar wavelet does not fulfill smoothness requirements (K) as assumed for the derivation of Theorem 4.2 in the previous section. Hence, for the proof of Theorem 5.4 a different methodology is needed than in Section 4. We follow the technique used in [23], exploiting the special structure of the Haar wavelet.

We begin by fixing a convenient inhomogeneous Haar system on the real line, namely

$$\mathcal{H}_1 := \{h_{j,k} : k \in \mathbb{Z}, j \in \mathbb{N}_0\},$$

where for $j \in \mathbb{N}$, $k \in \mathbb{Z}$, the functions $h_{j,k}$ are scaled Haar functions of the form

$$h_{j,k}(x) := \frac{1}{\sqrt{2}} h(2^{j-1}x - k) \quad , \quad \text{where} \quad h(x) := \mathbb{1}_{I_{0,0}^+}(x) - \mathbb{1}_{I_{0,0}^-}(x).$$

The intervals $I_{j,k}^+ = [2^{-j}k, 2^{-j}(k+1/2))$ and $I_{j,k}^- = [2^{-j}(k+1/2), 2^{-j}(k+1))$ thereby represent the dyadic children of the standard dyadic intervals $I_{j,k} = [2^{-j}k, 2^{-j}(k+1))$. At the lowest scale $j = 0$ the ordinary Haar functions $\mathbb{1}_{I_{0,k}^+} - \mathbb{1}_{I_{0,k}^-}$ are replaced by the characteristic functions $h_{0,k} := \mathbb{1}_{I_{0,k}}$. Further, we set $h_{j,k} \equiv 0$ if $j < 0$. Defined like this, the structure of the system \mathcal{H}_1 fits closely to the wavelet systems considered in Section 4. The inhomogeneous scale is at $j = 0$ (and not the usual standard $j = -1$ for Haar systems).

For dimension $d \in \mathbb{N}$ we derive a corresponding hyperbolic d -variate Haar system by the following tensorization procedure,

$$(24) \quad \mathcal{H}_d := \{h_{\bar{j},\bar{k}} := h_{j_1,k_1} \otimes \cdots \otimes h_{j_d,k_d} : \bar{k} = (k_1, \dots, k_d) \in \mathbb{Z}^d, \bar{j} = (j_1, \dots, j_d) \in \mathbb{N}_0^d\}.$$

Note that for $\bar{j} \in \mathbb{N}^d$ and $\bar{k} \in \mathbb{Z}^d$ the cube

$$Q_{\bar{j},\bar{k}} := [2^{-j_1}k_1, 2^{-j_1}(k_1+1)) \times \cdots \times [2^{-j_d}k_d, 2^{-j_d}(k_d+1)),$$

whose characteristic function will subsequently be denoted by $\chi_{\bar{j},\bar{k}}$ as already earlier in (7), corresponds to the strict support of the Haar function $h_{\bar{j},\bar{k}}$. At each fixed “scale” \bar{j} these cubes represent a partition of the d -dimensional domain \mathbb{R}^d .

Proposition 5.1 *Let $1 < p, q < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Under the condition*

$$|s|/\alpha_{\min} < \min \left\{ 1 - \frac{1}{p}, 1 - \frac{1}{q} \right\}$$

we have for $f \in S'(\mathbb{R}^d)$ (with the dual pairing $\langle f, h_{\bar{j},\bar{k}} \rangle$ defined as in (14) in Remark 4.2)

$$(25) \quad \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_{\infty} s q} \left| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle \chi_{\bar{j},\bar{k}}(x) \right|^q \right)^{1/q} \right\|_p \lesssim \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}},$$

whenever the left-hand side is defined. In case $|s|/\alpha_{\min} < 1 - \frac{1}{p}$ we have

$$(26) \quad \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_{\infty} s q} \left\| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle \chi_{\bar{j},\bar{k}}(\cdot) \right\|_p^q \right)^{1/q} \lesssim \|f\|_{\tilde{B}_{p,q}^{s,\bar{\alpha}}}.$$

Proof. For the proof, we first build a suitable decomposition of unity adapted to the hyperbolic tiling of the frequency domain. For a respective construction, we start with univariate functions $\phi_0, \phi \in S(\mathbb{R})$ and $\lambda_0, \lambda \in S(\mathbb{R})$ such that

$$\lambda_0 \phi_0 + \sum_{j \in \mathbb{N}} \lambda(2^{-j} \cdot) \phi(2^{-j} \cdot) = \sum_{j \in \mathbb{N}_0} \lambda_j \phi_j \equiv 1,$$

where $\phi_j := \phi(2^{-j} \cdot)$ and $\lambda_j := \lambda(2^{-j} \cdot)$ for $j \in \mathbb{N}$. The functions ϕ_0 and ϕ_1 shall thereby, as usual, be compactly supported with

$$\text{supp}(\phi_0) \subset \{|x| \leq 2\varepsilon\} \quad , \quad \text{supp}(\phi) \subset \{\varepsilon/2 \leq |x| \leq 2\varepsilon\}$$

for some $\varepsilon > 0$. As a consequence, their inverse Fourier transforms however, namely $\Phi_0 := \mathcal{F}^{-1} \phi_0$ and $\Phi := \mathcal{F}^{-1} \phi$, cannot have compact supports.

The functions λ_0, λ , on the other hand are chosen such that the supports of $\Lambda_0 := \mathcal{F}^{-1} \lambda_0$ and $\Lambda := \mathcal{F}^{-1} \lambda$ are compact. Further, they are assumed to fulfill the Tauberian conditions

$$|\lambda_0(x)| > 0 \quad \text{on } \{|x| \leq 2\varepsilon\} \quad , \quad |\lambda(x)| > 0 \quad \text{on } \{\varepsilon/2 \leq |x| \leq 2\varepsilon\}$$

with the same $\varepsilon > 0$ as above and furthermore $D^{\bar{\gamma}} \lambda(0) = 0$ for multi-indices $\bar{\gamma} \in \mathbb{N}_0^d$ with $\|\bar{\gamma}\|_1 \leq 1$. Such a construction is indeed possible, see [53, Lem. 3.6] for example.

For the subsequent proof, it is convenient to also define the functions $\Phi_j := \mathcal{F}^{-1} \phi_j$ and $\Lambda_j := \mathcal{F}^{-1} \lambda_j$ for $j \in \mathbb{N}$. They fulfill the scaling relations $\Phi_j = 2^j \Phi(2^j \cdot)$ and $\Lambda_j = 2^j \Lambda(2^j \cdot)$.

Next, we put $\Phi_j := 0$ and $\Lambda_j := 0$ for $j \in \mathbb{Z}$ with $j < 0$ and build the tensor products

$$\Phi_{\bar{\ell}} := \bigotimes_{i \in \{1, \dots, d\}} \Phi_{\ell_i} \quad \text{and} \quad \Lambda_{\bar{\ell}} := \bigotimes_{i \in \{1, \dots, d\}} \Lambda_{\ell_i} \quad \text{for } \bar{\ell} \in \mathbb{Z}^d.$$

Then we have the decomposition, which in fact is a discrete version of Calderón's reproducing formula,

$$f = \sum_{\bar{\ell} \in \mathbb{Z}^d} \Phi_{\bar{\ell}} * \Lambda_{\bar{\ell}} * f \quad \text{for every } f \in S'(\mathbb{R}^d),$$

enabling a component-wise evaluation of the scalar product $\langle f, h_{\bar{j}, \bar{k}} \rangle$. Each Haar coefficient can in this way be understood in the following sense (see also Remark 4.2),

$$\langle f, h_{\bar{j}, \bar{k}} \rangle = \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * \Lambda_{\bar{\ell}} * f, h_{\bar{j}, \bar{k}} \rangle = \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * f, \Lambda_{\bar{\ell}}(\cdot) * h_{\bar{j}, \bar{k}} \rangle,$$

whenever the right-hand sum converges.

If we further assume that $\Lambda_{\bar{\ell}}$ is even, we arrive at the estimate

$$|2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle| \lesssim \sum_{\bar{\ell} \in \mathbb{Z}^d} \left| \int_{\mathbb{R}^d} 2^{\|\bar{j}\|_1} (\Phi_{\bar{j}+\bar{\ell}} * f)(y) (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}})(y) dy \right|.$$

Let us investigate the integral on the right-hand side, and for this let us define

$$S_{\bar{j}, \bar{k}, \bar{\ell}} := \text{supp} (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}}).$$

If $\min_{i \in [d]} \{j_i + \ell_i\} < 0$ we have $S_{\bar{j}, \bar{k}, \bar{\ell}} = \emptyset$ and the integral vanishes. Otherwise, when $\min_{i \in [d]} \{j_i + \ell_i\} \geq 0$, we fix $a > 0$ and $x \in \mathbb{R}^d$ and obtain the estimate

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} 2^{\|\bar{j}\|_1} (\Phi_{\bar{j}+\bar{\ell}} * f)(y) (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}})(y) dy \right| \\ & \leq P_{2^{\bar{j}+\bar{\ell}}, a}(\Phi_{\bar{j}+\bar{\ell}} * f)(x) \cdot \sup_{y \in S_{\bar{j}, \bar{k}, \bar{\ell}}} \left[\prod_{i=1}^d (1 + 2^{j_i + \ell_i} |x_i - y_i|)^a \right] \cdot \left| \int_{\mathbb{R}^d} 2^{\|\bar{j}\|_1} (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}})(z) dz \right|, \end{aligned}$$

where $P_{2^{\bar{j}+\bar{\ell}}, a}(\Phi_{\bar{j}+\bar{\ell}} * f)$ denotes the Peetre maximal function (see (5))

$$P_{2^{\bar{j}+\bar{\ell}}, a}(\Phi_{\bar{j}+\bar{\ell}} * f)(x) = \sup_{y \in \mathbb{R}^d} \frac{|(\Phi_{\bar{j}+\bar{\ell}} * f)(y)|}{(1 + 2^{j_1 + \ell_1} |x_1 - y_1|)^a \cdots (1 + 2^{j_d + \ell_d} |x_d - y_d|)^a}.$$

The integral term splits into

$$\int_{\mathbb{R}^d} 2^{\|\bar{j}\|_1} (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}})(z) dz = 2^{\|\bar{j}\|_1} \int_{\mathbb{R}} (\Lambda_{j_1 + \ell_1} * h_{j_1, k_1})(t) dt \cdots \int_{\mathbb{R}} (\Lambda_{j_d + \ell_d} * h_{j_d, k_d})(t) dt$$

according to the relation

$$\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}} = (\Lambda_{j_1 + \ell_1} * h_{j_1, k_1}) \otimes \cdots \otimes (\Lambda_{j_d + \ell_d} * h_{j_d, k_d}).$$

For fixed $i \in \{1, \dots, d\}$, assuming $\ell_i < 0$, we can then further estimate

$$(27) \quad \int_{\mathbb{R}} (\Lambda_{j_i + \ell_i} * h_{j_i, k_i})(y) dy \lesssim 2^{-j_i + \ell_i}$$

since $|\text{supp}(\Lambda_{j_i+\ell_i} * h_{j_i,k_i})| \asymp 2^{-(j_i+\ell_i)}$ and $\|\Lambda_{j_i+\ell_i} * h_{j_i,k_i}\|_\infty \lesssim 2^{2\ell_i}$. For the latter of these two inequalities the first order vanishing moment of the Haar wavelet comes into play. Note here that indeed $j_i > 0$ due to $\min_{i \in [d]} \{j_i + \ell_i\} \geq 0$, allowing for the estimate

$$\begin{aligned} \|\Lambda_{j_i+\ell_i} * h_{j_i,k_i}\|_\infty &= \|(\Lambda_{j_i+\ell_i} - \Lambda_{j_i+\ell_i}(2^{-j_i}k_i)) * h_{j_i,k_i}\|_\infty \\ &\leq \|\Lambda_{j_i+\ell_i} - \Lambda_{j_i+\ell_i}(2^{-j_i}k_i)\|_\infty \|h_{j_i,k_i}\|_1 \lesssim 2^{2\ell_i}. \end{aligned}$$

In case $\ell_i \geq 0$ we obtain a different estimate than (27), namely

$$\int_{\mathbb{R}} (\Lambda_{j_i+\ell_i} * h_{j_i,k_i})(y) dy \lesssim 2^{-(j_i+\ell_i)}.$$

Here we use the fact that the integrand is bounded by a constant together with the observation that its support is contained in at most three intervals of length $\asymp 2^{-(j_i+\ell_i)}$. Indeed, as a consequence of the L_1 - resp. L_∞ -normalization of $\Lambda_{j_i+\ell_i}$ and h_{j_i,k_i} , we have $\|\Lambda_{j_i+\ell_i} * h_{j_i,k_i}\|_\infty \leq \|\Lambda_{j_i+\ell_i}\|_1 \|h_{j_i,k_i}\|_\infty \lesssim 1$. Furthermore, due to the vanishing moment properties of $\Lambda_{j_i+\ell_i}$, the support of the convolution merely stems from the either two or three discontinuities of the function h_{j_i,k_i} .

Now, let us turn our attention to the factor

$$\sup_{y \in S_{\bar{j}, \bar{k}, \bar{\ell}}} \left[\prod_{i=1}^d (1 + 2^{j_i+\ell_i} |x_i - y_i|)^a \right].$$

Here, we have with $x_i \in Q_{j_i,k_i}$ and $y_i \in \text{supp}(\Lambda_{j_i+\ell_i} * h_{j_i,k_i}) \subset \text{supp}(\Lambda_{j_i+\ell_i}) + \text{supp}(h_{j_i,k_i})$ (and therefore $|x_i - y_i| \lesssim 2^{-j_i}$ if $\ell_i \geq 0$ and $|x_i - y_i| \lesssim 2^{-(j_i+\ell_i)}$ if $\ell_i < 0$)

$$(1 + 2^{j_i+\ell_i} |x_i - y_i|)^a \lesssim 1$$

if $\ell_i < 0$. Otherwise, if $\ell_i \geq 0$, we estimate

$$(1 + 2^{j_i+\ell_i} |x_i - y_i|)^a \lesssim 2^{\ell_i a}.$$

Putting all together, this yields for $x \in Q_{\bar{j}, \bar{k}}$

$$\left| \int_{\mathbb{R}^d} 2^{|\bar{j}|} (\Phi_{\bar{j}+\bar{\ell}} * f)(y) (\Lambda_{\bar{j}+\bar{\ell}} * h_{\bar{j}, \bar{k}})(y) dy \right| \lesssim A(\bar{\ell}, a) P_{2^{\bar{j}+\bar{\ell}}, a}(\Phi_{\bar{j}+\bar{\ell}} * f)(x),$$

where

$$A(\bar{\ell}, a) := \prod_{i \in \{1, \dots, d\}} A(\ell_i, a) \quad \text{with} \quad A(\ell_i, a) := \begin{cases} 2^{\ell_i} & , \ell_i < 0 \\ 2^{(a-1)\ell_i} & , \ell_i \geq 0 \end{cases}.$$

Hence, we obtain uniformly in $x \in \mathbb{R}^d$ and for fixed $\bar{j} \in \mathbb{N}_0^d$

$$\sum_{\bar{k} \in \mathbb{Z}^d} |2^{|\bar{j}|} \langle f, h_{\bar{j}, \bar{k}} \rangle| \chi_{\bar{j}, \bar{k}}(x) \lesssim \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) P_{2^{\bar{j}+\bar{\ell}}, a}(\Phi_{\bar{j}+\bar{\ell}} * f)(x).$$

Finally, we can turn to the proof of (25). We estimate

$$\begin{aligned}
& \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty sq} \left| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \chi_{\bar{j}, \bar{k}}(x) \right|^q \right)^{1/q} \right\|_p \\
& \lesssim \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty sq} \left| \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) P_{2^{\bar{j}+\bar{\ell}}, a} (\Phi_{\bar{j}+\bar{\ell}} * f)(x) \right|^q \right)^{1/q} \right\|_p \\
& = \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \left| \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{(\|\bar{j}/\bar{\alpha}\|_\infty - \|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty) s} 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} P_{2^{\bar{j}+\bar{\ell}}, a} (\Phi_{\bar{j}+\bar{\ell}} * f)(x) \right|^q \right)^{1/q} \right\|_p.
\end{aligned}$$

According to (18) it holds $2^{(\|\bar{j}/\bar{\alpha}\|_\infty - \|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty) s} \leq 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|}$ for $s \in \mathbb{R}$, and hence

$$\begin{aligned}
& \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty sq} \left| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \chi_{\bar{j}, \bar{k}}(x) \right|^q \right)^{1/q} \right\|_p \\
& \lesssim \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \left| \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|} 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} P_{2^{\bar{j}+\bar{\ell}}, a} (\Phi_{\bar{j}+\bar{\ell}} * f)(x) \right|^q \right)^{1/q} \right\|_p \\
& \leq \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|} \cdot \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \left| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} P_{2^{\bar{j}}, a} (\Phi_{\bar{j}} * f)(x) \right|^q \right)^{1/q} \right\|_p,
\end{aligned}$$

where Young's convolution inequality was used in the last step.

For (26) we argue analogously, namely

$$\begin{aligned}
& \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty sq} \left\| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \chi_{\bar{j}, \bar{k}}(\cdot) \right\|_p^q \right)^{1/q} \\
& \lesssim \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_\infty sq} \left\| \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) P_{2^{\bar{j}+\bar{\ell}}, a} (\Phi_{\bar{j}+\bar{\ell}} * f)(\cdot) \right\|_p^q \right)^{1/q} \\
& \leq \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \left\| \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|} 2^{\|(\bar{j}+\bar{\ell})/\bar{\alpha}\|_\infty s} P_{2^{\bar{j}+\bar{\ell}}, a} (\Phi_{\bar{j}+\bar{\ell}} * f)(\cdot) \right\|_p^q \right)^{1/q} \\
& \leq \sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|} \cdot \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \left\| 2^{\|\bar{j}/\bar{\alpha}\|_\infty s} P_{2^{\bar{j}}, a} (\Phi_{\bar{j}} * f)(\cdot) \right\|_p^q \right)^{1/q}.
\end{aligned}$$

Choosing $a > \max\{1/p, 1/q\}$ in the F-case and $a > 1/p$ in the B-case, to ensure the boundedness of the Peetre maximal operator (see Theorem 2.3, also compare e.g. [54, Thm. 2.6]), as well as $|s| < \min_{i \in \{1, \dots, d\}} \{\alpha_i\} (1-a)$ we get

$$\sum_{\bar{\ell} \in \mathbb{Z}^d} A(\bar{\ell}, a) 2^{|\bar{\ell}/\bar{\alpha}|_\infty |s|} < \infty$$

and thus (25) and (26), respectively. \square

Using a duality argument, we can deduce an immediate companion result.

Proposition 5.2 *Let $1 < p, q < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Under the condition*

$$|s|/\alpha_{\min} < \min \left\{ \frac{1}{p}, \frac{1}{q} \right\}$$

we have for all $f \in S'(\mathbb{R}^d)$ (with the dual pairing $\langle f, h_{\bar{j}, \bar{k}} \rangle$ defined as in (14) in Remark 4.2)

$$(28) \quad \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} \lesssim \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{|\bar{j}|/\bar{\alpha}} \|\cdot\|_{\infty}^{sq} \left| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{|\bar{j}|\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \chi_{\bar{j}, \bar{k}}(x) \right|^q \right)^{1/q} \right\|_p,$$

whenever the right-hand side is defined. In case $|s|/\alpha_{\min} < 1/p$ we have

$$(29) \quad \|f\|_{\tilde{B}_{p,q}^{s,\bar{\alpha}}} \lesssim \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{|\bar{j}|/\bar{\alpha}} \|\cdot\|_{\infty}^{sq} \left\| \sum_{\bar{k} \in \mathbb{Z}^d} 2^{|\bar{j}|\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \chi_{\bar{j}, \bar{k}}(\cdot) \right\|_p^q \right)^{1/q}.$$

Proof. We showed in Proposition 5.1 (i) that the linear operator

$$A : \tilde{F}_{p,q}^{s,\bar{\alpha}} \rightarrow \tilde{f}_{p,q}^{s,\bar{\alpha}}, \quad f \mapsto (2^{|\bar{j}|\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle)_{\bar{j}, \bar{k}},$$

is well-defined and bounded in the parameter range

$$|s| < \min_{i \in \{1, \dots, d\}} \{\alpha_i\} \min \left\{ 1 - \frac{1}{p}, 1 - \frac{1}{q} \right\}.$$

Consequently, in this range, the dual operator

$$A' : (\tilde{f}_{p,q}^{s,\bar{\alpha}})' \rightarrow (\tilde{F}_{p,q}^{s,\bar{\alpha}})'$$

is also well-defined and bounded. Identifying $(\tilde{f}_{p,q}^{s,\bar{\alpha}})'$ with $\tilde{f}_{p',q'}^{-s,\bar{\alpha}}$ with respect to the non-standard duality product

$$(30) \quad \langle (\lambda_{\bar{j}, \bar{k}}), (\mu_{\bar{j}, \bar{k}}) \rangle := \sum_{\bar{j} \in \mathbb{N}_0^d} 2^{-|\bar{j}|\|_1} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}, \bar{k}} \mu_{\bar{j}, \bar{k}},$$

which is possible according to Theorem 5.5 (i) below, it can be represented in the form

$$A' : (\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}} \mapsto \sum_{\bar{j}, \bar{k}} \lambda_{\bar{j}, \bar{k}} h_{\bar{j}, \bar{k}},$$

where the convergence is weak*ly in $(\tilde{F}_{p,q}^{s,\bar{\alpha}})'$. This is a consequence of the relation

$$\begin{aligned} \langle Af, g' \rangle_{Y \times Y'} &= \sum_{\bar{j} \in \mathbb{N}_0^d} 2^{-|\bar{j}|\|_1} \sum_{\bar{k} \in \mathbb{Z}^d} 2^{|\bar{j}|\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \cdot \lambda_{\bar{j}, \bar{k}} \\ &= \int_{\mathbb{R}^d} f(x) \cdot \left(\sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}, \bar{k}} h_{\bar{j}, \bar{k}}(x) \right) dx \\ &= \langle f, A'g' \rangle_{X \times X'}, \end{aligned}$$

where we used the short-hand notation $Y = \tilde{f}_{p,q}^{s,\bar{\alpha}}$ and $X = \tilde{F}_{p,q}^{s,\bar{\alpha}}$.

Invoking Theorem 5.5(i) another time, X' can be identified with $\tilde{F}_{p',q'}^{-s,\bar{\alpha}}$ (and X'' with $\tilde{F}_{p,q}^{s,\bar{\alpha}}$). Then for $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p',q'}^{-s,\bar{\alpha}}$ and some enumeration

$$(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \rightsquigarrow (\lambda_n)_{n \in \mathbb{N}}$$

we have $\|(\lambda_n)_{n \geq N}\|_{\tilde{f}_{p',q'}^{-s,\bar{\alpha}}} \rightarrow 0$ for $N \rightarrow \infty$. We estimate

$$\begin{aligned} \|A'((\lambda_n)_{n \geq N})\|_{\tilde{F}_{p',q'}^{-s,\bar{\alpha}}} &= \sup_{\|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}}=1} |\langle A'((\lambda_n)_{n \geq N}), f \rangle| = \sup_{\|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}}=1} |\langle (\lambda_n)_{n \geq N}, Af \rangle| \\ &\leq \sup_{\|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}}=1} \|(\lambda_n)_{n \geq N}\|_{\tilde{f}_{p',q'}^{-s,\bar{\alpha}}} \|Af\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} \lesssim \|(\lambda_n)_{n \geq N}\|_{\tilde{f}_{p',q'}^{-s,\bar{\alpha}}} \rightarrow 0 \quad (N \rightarrow \infty). \end{aligned}$$

Hence, $A'((\lambda_n)_{n \leq N}) \rightarrow A'((\lambda_n)_{n \in \mathbb{N}})$ strongly and unconditionally in $\tilde{F}_{p',q'}^{-s,\bar{\alpha}}$.

In other words, we have shown that if $|s| < \min_{i \in \{1, \dots, d\}} \{\alpha_i\} \min\{1/p, 1/q\}$ and $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$ then $\sum_{\bar{j},\bar{k}} \lambda_{\bar{j},\bar{k}} h_{\bar{j},\bar{k}}$ converges strongly and unconditionally in $\tilde{F}_{p,q}^{s,\bar{\alpha}}$ and

$$(31) \quad \left\| \sum_{\bar{j},\bar{k}} \lambda_{\bar{j},\bar{k}} h_{\bar{j},\bar{k}} \right\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} \lesssim \|(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}\|_{\tilde{f}_{p,q}^{s,\bar{\alpha}}}.$$

Hence, choosing $\lambda_{\bar{j},\bar{k}} := 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle$ and assuming a finite sequence norm $\|(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}\|_{\tilde{f}_{p,q}^{s,\bar{\alpha}}}$, then $\sum_{\bar{j},\bar{k}} \lambda_{\bar{j},\bar{k}} h_{\bar{j},\bar{k}} = \sum_{\bar{j},\bar{k}} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle h_{\bar{j},\bar{k}}$ converges strongly to some limit $\tilde{f} \in \tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. Since this sum also converges (weak*ly) in $S'(\mathbb{R})$ to f we have $f = \tilde{f} = \sum_{\bar{j},\bar{k}} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle h_{\bar{j},\bar{k}}$ in $\tilde{F}_{p,q}^{s,\bar{\alpha}}$. Now (28) follows from (31).

The proof of (29) works the same, using Proposition 5.1 (ii) and Theorem 5.5 (ii). \square

Combining both, Proposition 5.1 and Proposition 5.2, we arrive at the following proposition, whereby we now concentrate on the F-case.

Proposition 5.3 *Let $1 < p, q < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Under the condition*

$$|s|/\alpha_{\min} < \min \left\{ \frac{1}{p}, \frac{1}{q}, 1 - \frac{1}{p}, 1 - \frac{1}{q} \right\}.$$

we have for all $f \in S'(\mathbb{R}^d)$ (with the dual pairing $\langle f, h_{\bar{j},\bar{k}} \rangle$ defined as in (14) in Remark 4.2)

$$(32) \quad \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j},\bar{k}} \rangle \chi_{\bar{j},\bar{k}}(x) \right)^q \right\|_p^{1/q} \asymp \|f\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}},$$

whenever the left-hand side is defined.

As a direct consequence of this result, we can finally formulate the main theorem of this section which corresponds to Theorem 4.2.

Theorem 5.4 *Let $1 < p, q < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Further, assume*

$$(33) \quad |s|/\alpha_{\min} < \min \left\{ \frac{1}{p}, \frac{1}{q}, 1 - \frac{1}{p}, 1 - \frac{1}{q} \right\}.$$

Then the Haar system $\mathcal{H}_d = (h_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ defined in (24) constitutes an unconditional basis of $\tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d)$ with associated sequence space $\tilde{f}_{p,q}^{s, \bar{\alpha}}$. The unique sequence of basis coefficients for $f \in \tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d)$ is determined by $\lambda := \lambda(f) = (\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ with

$$(34) \quad \lambda_{\bar{j}, \bar{k}} := \lambda_{\bar{j}, \bar{k}}(f) = 2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle.$$

Further, we have the wavelet isomorphism (equivalent norm)

$$\|f\|_{\tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d)} \asymp \|\lambda(f)\|_{\tilde{f}_{p,q}^{s, \bar{\alpha}}} \quad , \quad f \in \tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d).$$

In addition, we can use \mathcal{H}_d to distinguish those elements of $\mathcal{S}'(\mathbb{R}^d)$ that belong to $\tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d)$. Those are characterized by either of the following two criteria:

(i) *f can be represented as a sum*

$$(35) \quad f = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j}, \bar{k}} h_{\bar{j}, \bar{k}} \quad \text{converging (weak*ly) in } \mathcal{S}'(\mathbb{R}^d)$$

with coefficients $(\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}} \in \tilde{f}_{p,q}^{s, \bar{\alpha}}$ (with respect to some chosen ordering).

(ii) *With $\lambda(f)$ being defined as in (34), it holds*

$$\lambda(f) = (\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}} \in \tilde{f}_{p,q}^{s, \bar{\alpha}}.$$

In both cases, the sequence $(\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ is necessarily the sequence of basis coefficients and the representation (35) converges unconditionally to f in $\tilde{F}_{p,q}^{s, \bar{\alpha}}(\mathbb{R}^d)$.

Proof. As a direct consequence of the equivalence (32) proved in Proposition 5.3 the analysis operator

$$\lambda : f \mapsto \left(2^{\|\bar{j}\|_1} \langle f, h_{\bar{j}, \bar{k}} \rangle \right)_{\bar{j}, \bar{k}}$$

is well-defined and bounded from $\tilde{F}_{p,q}^{s, \bar{\alpha}}$ to $\tilde{f}_{p,q}^{s, \bar{\alpha}}$. Moreover, it is injective and we have the equivalence of norms $\|f\|_{\tilde{F}_{p,q}^{s, \bar{\alpha}}} \asymp \|\lambda(f)\|_{\tilde{f}_{p,q}^{s, \bar{\alpha}}}$. Further, for $f \in \mathcal{S}'(\mathbb{R}^d)$ we have $f \in \tilde{F}_{p,q}^{s, \bar{\alpha}}$ if and only if $\lambda(f) \in \tilde{f}_{p,q}^{s, \bar{\alpha}}$ (whenever $\lambda(f)$ is defined).

Now, let us have a look at the synthesis operator

$$(36) \quad S : (\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}} \mapsto \sum_{\bar{j}, \bar{k}} \lambda_{\bar{j}, \bar{k}} h_{\bar{j}, \bar{k}}.$$

Clearly, for every finite sequence the assignment S defines an element in $\tilde{F}_{p,q}^{s, \bar{\alpha}}$. By completion, using (32) and the fact that the finite sequences lie dense in $\tilde{f}_{p,q}^{s, \bar{\alpha}}$, this synthesis further extends to all sequences of $\tilde{f}_{p,q}^{s, \bar{\alpha}}$, with unconditional and strong convergence of (36) in $\tilde{F}_{p,q}^{s, \bar{\alpha}}$. Hence, S

is a well-defined bounded linear operator from $\tilde{f}_{p,q}^{s,\bar{\alpha}}$ to $\tilde{F}_{p,q}^{s,\bar{\alpha}}$, which, as another consequence of (32), is also injective.

Next, turning to the composition $\lambda \circ S$ operating from $\tilde{f}_{p,q}^{s,\bar{\alpha}}$ to $\tilde{f}_{p,q}^{s,\bar{\alpha}}$, we deduce for each fixed $\bar{j}_* \in \mathbb{N}_0^d$ and $\bar{k}_* \in \mathbb{Z}^d$

$$\left\langle \sum_{\bar{j}, \bar{k}} 2^{|\bar{j}|} \lambda_{\bar{j}, \bar{k}} h_{\bar{j}, \bar{k}}, h_{\bar{j}_*, \bar{k}_*} \right\rangle = \sum_{\bar{j}, \bar{k}} 2^{|\bar{j}|} \lambda_{\bar{j}, \bar{k}} \langle h_{\bar{j}, \bar{k}}, h_{\bar{j}_*, \bar{k}_*} \rangle = \lambda_{\bar{j}_*, \bar{k}_*},$$

using the orthogonality of the system $(h_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ in $L_2(\mathbb{R}^d)$. We obtain $Id_{\tilde{f}_{p,q}^{s,\bar{\alpha}}} = \lambda \circ S$ and in turn $\lambda \circ S \circ \lambda = \lambda$. Due to the injectivity of λ , the latter equality further implies $Id_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} = S \circ \lambda$. In particular, λ and S are thus bijections and every $f \in \tilde{F}_{p,q}^{s,\bar{\alpha}}$ allows for a representation (35).

To see that the representing coefficients $(\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ are unique, under the assumption of strong convergence of the sum, let $\lambda^* = (\lambda_{\bar{j}, \bar{k}}^*)_{\bar{j}, \bar{k}}$ be some sequence which satisfies (35) in a strong sense for some special ordering of the sum. Then again (32) together with a completion argument yields $\lambda^* \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$, and thus $\lambda^* = \lambda(f)$ by the injectivity of S . Hence, the expansion coefficients in (35) are unique and it follows that \mathcal{H}_d is a basis. Its unconditionality is due to the fact that the convergence of (35) for sequences $\lambda^* \in \tilde{f}_{p,q}^{s,\bar{\alpha}}$ is always unconditional.

For the proof of criterion (i) we just remark that for sequences $(\lambda_{\bar{j}, \bar{k}})_{\bar{j}, \bar{k}}$ in $\tilde{f}_{p,q}^{s,\bar{\alpha}}$ with weak*-convergence of (35) in $\mathcal{S}'(\mathbb{R}^d)$ the convergence is automatically in the stronger sense of $\tilde{F}_{p,q}^{s,\bar{\alpha}}$. \square

Remark 5.1 *For brevity, the above theorem was only stated for the F-case. There also exists a B-version, which reads precisely the same apart from condition (33) which is replaced by*

$$|s|/\alpha_{min} < \min \left\{ \frac{1}{p}, 1 - \frac{1}{p} \right\}.$$

In the proof of Proposition 5.2 we utilized isomorphisms $(\tilde{F}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{F}_{p',q'}^{-s,\bar{\alpha}}$ and $(\tilde{f}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{f}_{p',q'}^{-s,\bar{\alpha}}$ as well as $(\tilde{B}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{B}_{p',q'}^{-s,\bar{\alpha}}$ and $(\tilde{b}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{b}_{p',q'}^{-s,\bar{\alpha}}$. Hence, for the completeness of our exposition, it remains to establish those.

Theorem 5.5 *Let $1 < p, q < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ with $\sum_{i=1}^d \alpha_i = d$. Then*

$$\left(\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) \right)' \cong \tilde{F}_{p',q'}^{-s,\bar{\alpha}}(\mathbb{R}^d) \quad \text{and} \quad \left(\tilde{f}_{p,q}^{s,\bar{\alpha}} \right)' \cong \tilde{f}_{p',q'}^{-s,\bar{\alpha}},$$

whereby the second of these isomorphies has to be understood with respect to the non-standard pairing (30). In the Besov case, we have the analogous relations

$$\left(\tilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) \right)' \cong \tilde{B}_{p',q'}^{-s,\bar{\alpha}}(\mathbb{R}^d) \quad \text{and} \quad \left(\tilde{b}_{p,q}^{s,\bar{\alpha}} \right)' \cong \tilde{b}_{p',q'}^{-s,\bar{\alpha}}.$$

The proof of this theorem is based on two auxiliary propositions. The first one of these is stated without proof, since it is a straightforward generalization of the classic identifications $(L_p(\mathbb{R}^d))' \cong L_{p'}(\mathbb{R}^d)$ and $(\ell_p(I))' \cong \ell_{p'}(I)$ when $1 < p < \infty$ (see e.g. [16]).

Proposition 5.6 *Let I be an arbitrary countable index set. Then*

$$\left(\ell_q(I, L_p(\mathbb{R}^d)) \right)' \cong \ell_{q'}(I, L_{p'}(\mathbb{R}^d)) \quad \text{and} \quad \left(L_p(\mathbb{R}^d, \ell_q(I)) \right)' \cong L_{p'}(\mathbb{R}^d, \ell_{q'}(I))$$

in the sense that there exist isomorphisms $f \mapsto (f_i)_{i \in I}$ such that

$$\langle f, g \rangle_{Y' \times Y} = \sum_{i \in I} \int_{\mathbb{R}^d} f_i(x) g_i(x) dx$$

for the respective cases $Y = \ell_q(I, L_p(\mathbb{R}^d))$ and $Y = L_p(\mathbb{R}^d, \ell_q(I))$.

The second proposition provides an alternative way to characterize functions in $\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $\widetilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$. Its counterpart in the classical setting of Triebel-Lizorkin spaces is Proposition 1 in [52, Sec. 2.3.4].

Proposition 5.7 *Assume $1 < p, q < \infty$, $s \in \mathbb{R}$, $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$, and $\sum_{i=1}^d \alpha_i = d$.*

(i) *Then $f \in S'(\mathbb{R}^d)$ belongs to $\widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ if and only if there exists a family $\{f_{\bar{j}}\}_{\bar{j} \in \mathbb{N}_0^d} \subset L_p(\mathbb{R}^d)$ such that*

$$f = \sum_{\bar{j}} \Delta_{\bar{j}} f_{\bar{j}} \text{ in } S'(\mathbb{R}^d) \quad \text{and} \quad \|2^{|\bar{j}|/\bar{\alpha}} f_{\bar{j}}\|_{L_p(\mathbb{R}^d, \ell_q)} < \infty.$$

(ii) *Then $f \in S'(\mathbb{R}^d)$ belongs to $\widetilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ if and only if there exists a family $\{f_{\bar{j}}\}_{\bar{j} \in \mathbb{N}_0^d} \subset L_p(\mathbb{R}^d)$ such that*

$$(37) \quad f = \sum_{\bar{j}} \Delta_{\bar{j}} f_{\bar{j}} \text{ in } S'(\mathbb{R}^d) \quad \text{and} \quad \|2^{|\bar{j}|/\bar{\alpha}} f_{\bar{j}}\|_{\ell_q(L_p(\mathbb{R}^d))} < \infty.$$

Proof. (i) Adaption of proof of Proposition 1 in [52, Sec. 2.3.4].

(ii) Let $(\phi_{\bar{j}})_{\bar{j}}$ be a hyperbolic resolution of unity as introduced in Definition 3.1 with associated hyperbolic Littlewood-Paley analysis $(\Delta_{\bar{j}})_{\bar{j}}$. Further, take $f \in \widetilde{B}_{p,q}^{s,\bar{\alpha}}$ and put $f_{\bar{j}} := \sum_{\bar{r} \in \{-1,0,1\}^d} \mathcal{F}^{-1} \phi_{\bar{j}+\bar{r}} \mathcal{F} f$ for $\bar{j} \in \mathbb{N}_0^d$, whereby we let $\phi_{-1} := 0$. Then $\{f_{\bar{j}}\}_{\bar{j}} \subset L_p$ and

$$\sum_{\bar{j} \in \mathbb{N}_0^d} \mathcal{F}^{-1} \phi_{\bar{j}} \mathcal{F} f_{\bar{j}} = \sum_{\bar{j} \in \mathbb{N}_0^d} \mathcal{F}^{-1} \phi_{\bar{j}} \left(\sum_{\bar{r} \in \{-1,0,1\}^d} \phi_{\bar{j}+\bar{r}} \right) \mathcal{F} f = \sum_{\bar{j} \in \mathbb{N}_0^d} \mathcal{F}^{-1} \phi_{\bar{j}} \mathcal{F} f = \sum_{\bar{j} \in \mathbb{N}_0^d} \Delta_{\bar{j}} f = f$$

as well as

$$\begin{aligned} \|2^{|\bar{j}|/\bar{\alpha}} f_{\bar{j}}\|_{\ell_q(L_p)} &= \left\| 2^{|\bar{j}|/\bar{\alpha}} \sum_{\bar{r} \in \{-1,0,1\}^d} \mathcal{F}^{-1} \phi_{\bar{j}+\bar{r}} \mathcal{F} f \right\|_{\ell_q(L_p)} \\ &\lesssim \sum_{\bar{r} \in \{-1,0,1\}^d} \left\| 2^{|\bar{j}|/\bar{\alpha}} \mathcal{F}^{-1} \phi_{\bar{j}+\bar{r}} \mathcal{F} f \right\|_{\ell_q(L_p)} \asymp \|f\|_{\widetilde{B}_{p,q}^{s,\bar{\alpha}}} < \infty. \end{aligned}$$

This settles one direction of the assertion. For the other direction, let $f \in S'(\mathbb{R}^d)$ satisfy (37) with associated $\{f_{\bar{j}}\}_{\bar{j}} \subset L_p$. In view of $f = \sum_{\bar{j}} \Delta_{\bar{j}} f_{\bar{j}}$, we can estimate

$$\begin{aligned} \|2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} f\|_{\ell_q(L_p)} &= \left\| 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} \sum_{\bar{r} \in \{-1,0,1\}^d} \Delta_{\bar{j}+\bar{r}} f_{\bar{j}+\bar{r}} \right\|_{\ell_q(L_p)} \\ &\leq \sum_{\bar{r} \in \{-1,0,1\}^d} \left\| 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} \Delta_{\bar{j}+\bar{r}} f_{\bar{j}+\bar{r}} \right\|_{\ell_q(L_p)} \lesssim \sum_{\bar{r} \in \{-1,0,1\}^d} \left\| 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}+\bar{r}} f_{\bar{j}+\bar{r}} \right\|_{\ell_q(L_p)} \\ &\asymp \left\| 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} f_{\bar{j}} \right\|_{\ell_q(L_p)} \lesssim \left\| 2^{|\bar{j}|/\bar{\alpha}} f_{\bar{j}} \right\|_{\ell_q(L_p)} < \infty, \end{aligned}$$

where the last two lines are due to the Hörmander-Mikhlin multiplier theorem, which is applied twice. This estimate shows $f \in \widetilde{B}_{p,q}^{s,\bar{\alpha}}$, finishing the proof. \square

Now we are ready to give a thorough proof of the duality relations stated in Theorem 5.5.

Proof. [of Theorem 5.5] We restrict to the F-case and begin with the more involved relation $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})' \cong \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$. The subsequent proof is thereby an adaption of the proof of the classical theorem in [52, Sec. 2.11.2] to the setting of hyperbolic spaces.

It is essential to note that, since $S(\mathbb{R}^d)$ lies dense in $\widetilde{F}_{p,q}^{s,\bar{\alpha}}$, there is a natural embedding

$$(38) \quad \kappa : \left(\widetilde{F}_{p,q}^{s,\bar{\alpha}}\right)' \hookrightarrow S'(\mathbb{R}^d).$$

Hence, both $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$ and $\widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$ can be interpreted as subspaces of $S'(\mathbb{R}^d)$, simplifying the following considerations.

Let us first assume that $f \in S'(\mathbb{R}^d)$ is an element of $\widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$ and take $\Phi_{\bar{\ell}}$ and $\Lambda_{\bar{\ell}}$ as in the proof of Proposition 5.1, for instance. Then f defines an element of $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$ via

$$(39) \quad \langle f, g \rangle_* := \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * f, \overline{\Lambda_{\bar{\ell}} * g} \rangle, \quad \text{where } g \in \widetilde{F}_{p,q}^{s,\bar{\alpha}},$$

as can be seen by the following estimate,

$$\begin{aligned} |\langle f, g \rangle_*| &= \left| \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * f, \overline{\Lambda_{\bar{\ell}} * g} \rangle \right| = \left| \sum_{\bar{\ell} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} (\Phi_{\bar{\ell}} * f)(y) \cdot (\Lambda_{\bar{\ell}} * g)(y) dy \right| \\ &= \left| \int_{\mathbb{R}^d} \sum_{\bar{j} \in \mathbb{N}_0^d} (\Phi_{\bar{j}} * f)(y) \cdot (\Lambda_{\bar{j}} * g)(y) dy \right| \\ &\leq \int_{\mathbb{R}^d} \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{-\|\bar{j}/\bar{\alpha}\|_{\infty} s q'} |\Phi_{\bar{j}} * f(y)|^{q'} \right)^{1/q'} \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_{\infty} s q} |\Lambda_{\bar{j}} * g(y)|^q \right)^{1/q} dy \\ &\leq \left(\int_{\mathbb{R}^d} \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{-\|\bar{j}/\bar{\alpha}\|_{\infty} s q'} |\Phi_{\bar{j}} * f(y)|^{q'} \right)^{p'/q'} dy \right)^{1/p'} \\ &\quad \times \left(\int_{\mathbb{R}^d} \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{\|\bar{j}/\bar{\alpha}\|_{\infty} s q} |\Lambda_{\bar{j}} * g(y)|^q \right)^{p/q} dy \right)^{1/p} \\ &\lesssim \|f\|_{\widetilde{F}_{p',q'}^{-s,\bar{\alpha}}} \cdot \|g\|_{\widetilde{F}_{p,q}^{s,\bar{\alpha}}}. \end{aligned}$$

Hereby, we applied Hölder's inequality and used $\Phi_{\bar{\ell}} = \Lambda_{\bar{\ell}} = 0$ if $\bar{\ell} \notin \mathbb{N}_0^d$.

The duality product $\langle \cdot, \cdot \rangle_*$ thus yields an embedding $\iota : \widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \rightarrow (\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$. Further, we have natural embeddings $\nu : \widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \hookrightarrow S'(\mathbb{R}^d)$ and $\kappa : (\widetilde{F}_{p,q}^{s,\bar{\alpha}})' \hookrightarrow S'(\mathbb{R}^d)$ (see (38)). To establish a bridge between κ , ι , and ν , we now consider the special case of a Schwartz function $g = \phi \in S(\mathbb{R}^d)$ in (39). We obtain

$$\langle f, \phi \rangle_* = \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * f, \Lambda_{\bar{\ell}} * \phi \rangle_{S' \times S} = \sum_{\bar{\ell} \in \mathbb{Z}^d} \langle \Phi_{\bar{\ell}} * \Lambda_{\bar{\ell}} * f, \phi \rangle_{S' \times S} = \langle f, \phi \rangle_{S' \times S}.$$

Hence, the above (somewhat artificially) defined operation of f on $\widetilde{F}_{p,q}^{s,\bar{\alpha}}$ via $\langle \cdot, \cdot \rangle_*$ is compatible with the operation of f as an element of $S'(\mathbb{R}^d)$ on $S(\mathbb{R}^d)$. This proves $\nu = \kappa \circ \iota$ and thus $\widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \subset (\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$, considered as subsets of $S'(\mathbb{R}^d)$.

It remains to prove the converse inclusion $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})' \subset \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$. For this, let $f \in S'(\mathbb{R}^d)$ be an element of $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$. We will show that this implies $f \in \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$ and to this end start with a construction of an isometric embedding

$$(40) \quad \mu : \left(\widetilde{F}_{p,q}^{s,\bar{\alpha}}\right)' \rightarrow L_{p'}(\mathbb{R}^d, \ell_{q'}), \quad f \mapsto (f_{\bar{j}})_{\bar{j}}.$$

Thereby, we build upon the observation that the assignment $g \mapsto (2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} g)_{\bar{j}}$ maps $\widetilde{F}_{p,q}^{s,\bar{\alpha}}$ isometrically to a closed subspace of $L_p(\mathbb{R}^d, \ell_q)$. Via this assignment and the Hahn-Banach extension theorem, it is therefore possible to identify each functional $f \in (\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$ with a functional on $L_p(\mathbb{R}^d, \ell_q)$ having the same norm. Invoking Proposition 5.6 (i), this then yields an associated family $(f_{\bar{j}})_{\bar{j}} \in L_{p'}(\mathbb{R}^d, \ell_{q'})$ with $\|(f_{\bar{j}})_{\bar{j}}\|_{L_{p'}(\mathbb{R}^d, \ell_{q'})} = \|f\|_{(\widetilde{F}_{p,q}^{s,\bar{\alpha}})'}$ and $\langle f, g \rangle = \sum_{\bar{j}} \langle f_{\bar{j}}, 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} g \rangle$, establishing (40).

In particular, for every $\phi \in S(\mathbb{R}^d)$

$$\langle f, \phi \rangle = \sum_{\bar{j}} \langle f_{\bar{j}}, 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} \phi \rangle = \sum_{\bar{j}} \langle \Delta_{\bar{j}} \tilde{f}_{\bar{j}}, \phi \rangle,$$

with $\tilde{f}_{\bar{j}} := 2^{|\bar{j}|/\bar{\alpha}} f_{\bar{j}}$. Hence, we have

$$f = \sum_{\bar{j}} 2^{|\bar{j}|/\bar{\alpha}} \Delta_{\bar{j}} \tilde{f}_{\bar{j}} = \sum_{\bar{j}} \Delta_{\bar{j}} \tilde{f}_{\bar{j}} \quad \text{weak*ly in } S'(\mathbb{R}^d).$$

Further, it holds $\|(2^{-|\bar{j}|/\bar{\alpha}} \tilde{f}_{\bar{j}})_{\bar{j}}\|_{\ell_{q'}(L_{p'})} = \|(f_{\bar{j}})_{\bar{j}}\|_{\ell_{q'}(L_{p'})} = \|f\|_{(\widetilde{F}_{p,q}^{s,\bar{\alpha}})'}$. In view of Proposition 5.7 (i), this shows $f \in \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$ and finishes the proof of $(\widetilde{F}_{p,q}^{s,\bar{\alpha}})' \cong \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$.

We next establish $(\tilde{f}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{f}_{p',q'}^{-s,\bar{\alpha}}$, which can be elegantly done using the previous result together with the wavelet isomorphism $\lambda : \widetilde{F}_{p,q}^{s,\bar{\alpha}} \rightarrow \tilde{f}_{p,q}^{s,\bar{\alpha}}$ established in Theorem 4.2. For this, we first verify that λ preserves the duality structure of $\widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \times \widetilde{F}_{p,q}^{s,\bar{\alpha}}$. Indeed, for $f \in \widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$ and $g \in \widetilde{F}_{p,q}^{s,\bar{\alpha}}$ we have

$$\begin{aligned} \langle f, g \rangle_{\widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \times \widetilde{F}_{p,q}^{s,\bar{\alpha}}} &= \left\langle \sum_{\bar{j}, \bar{k}} 2^{|\bar{j}|} \langle f, \psi_{\bar{j}, \bar{k}} \rangle \psi_{\bar{j}, \bar{k}}, g \right\rangle_{\widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \times \widetilde{F}_{p,q}^{s,\bar{\alpha}}} = \sum_{\bar{j}, \bar{k}} 2^{|\bar{j}|} \langle f, \psi_{\bar{j}, \bar{k}} \rangle \langle \psi_{\bar{j}, \bar{k}}, g \rangle_{\widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \times \widetilde{F}_{p,q}^{s,\bar{\alpha}}} \\ &= \sum_{\bar{j}, \bar{k}} 2^{-|\bar{j}|} (2^{|\bar{j}|} \langle f, \psi_{\bar{j}, \bar{k}} \rangle) (2^{|\bar{j}|} \langle g, \psi_{\bar{j}, \bar{k}} \rangle) = \langle \lambda(f), \lambda(g) \rangle_{\tilde{f}_{p',q'}^{-s,\bar{\alpha}} \times \tilde{f}_{p,q}^{s,\bar{\alpha}}}. \end{aligned}$$

Note that hereby we relied on the strong convergence of the wavelet expansion in the space $\widetilde{F}_{p',q'}^{-s,\bar{\alpha}}$. Next we recall the isomorphism $\iota : \widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \rightarrow (\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$ established above and let $\lambda' : (\tilde{f}_{p,q}^{s,\bar{\alpha}})' \rightarrow (\widetilde{F}_{p,q}^{s,\bar{\alpha}})'$ denote the dual map of λ , which is also an isomorphism. Then we can read $(\tilde{f}_{p,q}^{s,\bar{\alpha}})' \cong \tilde{f}_{p',q'}^{-s,\bar{\alpha}}$ directly from the following chain of isomorphisms

$$(\tilde{f}_{p,q}^{s,\bar{\alpha}})' \times \tilde{f}_{p,q}^{s,\bar{\alpha}} \xrightarrow{\lambda' \times \lambda^{-1}} (\widetilde{F}_{p,q}^{s,\bar{\alpha}})' \times \widetilde{F}_{p,q}^{s,\bar{\alpha}} \xrightarrow{\iota^{-1} \times Id} \widetilde{F}_{p',q'}^{-s,\bar{\alpha}} \times \widetilde{F}_{p,q}^{s,\bar{\alpha}} \xrightarrow{\lambda \times \lambda} \tilde{f}_{p',q'}^{-s,\bar{\alpha}} \times \tilde{f}_{p,q}^{s,\bar{\alpha}}.$$

□

6 Hyperbolic and classical (anisotropic) Sobolev spaces

In the remaining two sections we will analyze the relationship between the newly introduced hyperbolic scale of spaces $\widetilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ from Section 3, where $A \in \{B, F\}$, and the classical scale of anisotropic spaces $A_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$, which was recalled in Section 2. Our first result shows that, surprisingly, for Sobolev spaces (i.e. the case $A = F$, $1 < p < \infty$, $q = 2$) both scales coincide.

Theorem 6.1 *Let $1 < p < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} > 0$ be an anisotropy vector as above. Then*

$$\widetilde{W}_p^{s,\bar{\alpha}}(\mathbb{R}^d) = W_p^{s,\bar{\alpha}}(\mathbb{R}^d) \quad (\text{in the sense of equivalent norms}).$$

Proof. The proof is divided into two steps. For convenience, we will thereby abbreviate by $m_{\bar{\alpha},s} := \left(\sum_{i=1}^d (1 + \xi_i^2)^{1/(2\alpha_i)} \right)^s$ the function which appears in the definition (3) of $\widetilde{W}_p^{s,\bar{\alpha}}$.

Step 1. In the first step we prove $\|f\|_{W_p^{s,\bar{\alpha}}} \lesssim \|f\|_{\widetilde{W}_p^{s,\bar{\alpha}}}$. For $\bar{j} = (j_1, \dots, j_d) \in \mathbb{N}_0^d$, let $(\varphi_{\bar{j}})_{\bar{j}}$ denote a fixed hyperbolic resolution of unity as introduced in Definition 3.1, with corresponding univariate family $(\varphi_j)_j$ where $\text{supp}(\varphi_0) \subset [-2, 2]$. In addition, let us also construct a second hyperbolic resolution of unity $(\psi_{\bar{j}})_{\bar{j}}$ such that $\psi_{\bar{j}}\varphi_{\bar{j}} = \varphi_{\bar{j}}$ for every $\bar{j} \in \mathbb{N}_0^d$. Hereby, it is not possible for $(\psi_{\bar{j}})_{\bar{j}}$ to obey the same strict building law as formulated in Definition 3.1. We define functions

$$\psi_0^* := \varphi_0 + \varphi_1, \quad \psi_1^* := \varphi_0 + \varphi_1 + \varphi_2, \quad \psi_2^* := \varphi_0 + \varphi_1 + \varphi_2 + \varphi_3, \quad \psi_j^* := \sum_{r=-1}^1 \varphi_{j+r} \quad (j \geq 3)$$

and then put $\psi_j := \psi_j^*/3$ for $j \in \mathbb{N}_0$. Then clearly $\sum_j \psi_j = 1$ and $\psi_j\varphi_j = \varphi_j$. Finally, we set

$$\psi_{\bar{j}} := \psi_{j_1} \otimes \dots \otimes \psi_{j_d} \quad \text{to obtain} \quad (\psi_{\bar{j}})_{\bar{j}}.$$

By construction, $\psi_0 = \varphi_0(2^{-1}\cdot)/3$, $\psi_{j+1} = \psi_j(2^{-1}\cdot)$ for $j \in \mathbb{N}_0 \setminus \{2\}$, and $\psi_3 = \psi_0(2^{-3}\cdot) - \psi_0$. As a consequence, we can record $\text{supp} \psi_j(2^j\cdot) \subset [-4, 4]$ for all $j \in \mathbb{N}_0$.

Utilizing $(\psi_{\bar{j}})_{\bar{j}}$, we then first rewrite the $W_p^{s,\bar{\alpha}}$ -norm as follows,

$$\begin{aligned} \|f\|_{W_p^{s,\bar{\alpha}}} &\asymp \left\| \mathcal{F}^{-1}[m_{\bar{\alpha},s}\mathcal{F}f] \right\|_p \\ &\asymp \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} |\mathcal{F}^{-1}\varphi_{\bar{j}}m_{\bar{\alpha},s}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p \\ &= \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} |2^{s\|\bar{j}/\bar{\alpha}\|_\infty} \mathcal{F}^{-1}(2^{-s\|\bar{j}/\bar{\alpha}\|_\infty} \psi_{\bar{j}}m_{\bar{\alpha},s})\varphi_{\bar{j}}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p. \end{aligned}$$

After this, we denote $M_{\bar{\alpha},s,\bar{j}} := 2^{-s\|\bar{j}/\bar{\alpha}\|_\infty} \psi_{\bar{j}}m_{\bar{\alpha},s}$ and apply the Fourier multiplier lemma 3.1 with $\rho_{\bar{j}} := M_{\bar{\alpha},s,\bar{j}}$ and $f_{\bar{j}} := 2^{s\|\bar{j}/\bar{\alpha}\|_\infty} \mathcal{F}^{-1}\varphi_{\bar{j}}\mathcal{F}f$. Due to $p > 1$ and $q = 2$ we can thereby choose $r := 2$. This leads to

$$\left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{2s\|\bar{j}/\bar{\alpha}\|_\infty} |\mathcal{F}^{-1}M_{\bar{\alpha},s,\bar{j}}\varphi_{\bar{j}}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p \lesssim \sup_{\bar{j} \in \mathbb{N}_0^d} \|M_{\bar{\alpha},s,\bar{j}}(2^{j_1}\cdot, \dots, 2^{j_d}\cdot)\|_{S_2^2 W} \|f\|_{\widetilde{W}_p^s}.$$

To finish Step 1, it now merely remains to check whether $\sup_{\bar{j} \in \mathbb{N}_0^d} \|M_{\bar{\alpha}, s, \bar{j}}(2^{j_1 \cdot}, \dots, 2^{j_d \cdot})\|_{S_2^2 W}$ is finite. But, defining $\tilde{\psi}_0 := \psi_0$ and $\tilde{\psi}_1 := \psi_3(2^{\cdot})$ and letting $\tilde{\psi}_{\bar{r}} := \tilde{\psi}_{r_1} \otimes \dots \otimes \tilde{\psi}_{r_d}$ for $\bar{r} \in \{0, 1\}^d$, we have

$$M_{\bar{\alpha}, s, \bar{j}}(2^{j_1} \xi_1, \dots, 2^{j_d} \xi_d) = 2^{-s \|\bar{j}/\bar{\alpha}\|_\infty} \tilde{\psi}_{s(\bar{j})}(\xi_1, \dots, \xi_d) m_{\bar{\alpha}, s}(2^{j_1} \xi_1, \dots, 2^{j_d} \xi_d),$$

where $s(\bar{j}) := (\text{sgn}(j_1 - 2), \dots, \text{sgn}(j_d - 2))_+$. And for each $\bar{j} \in \mathbb{N}_0^d$, the function $\tilde{\psi}_{s(\bar{j})}$ belongs to $\mathcal{S}(\mathbb{R}^d)$ and is supported on $[-4, 4]^d$. Hence, it is sufficient to verify that the second derivatives (up to order 2) in every component of

$$F_{\bar{\alpha}, s, \bar{j}} : (\xi_1, \dots, \xi_d) \mapsto 2^{-s \|\bar{j}/\bar{\alpha}\|_\infty} m_{\bar{\alpha}, s}(2^{j_1} \xi_1, \dots, 2^{j_d} \xi_d) = \left(\sum_{i=1}^d 2^{-\|\bar{j}/\bar{\alpha}\|_\infty} (1 + 2^{2j_i} \xi_i^2)^{1/(2\alpha_i)} \right)^s$$

are uniformly bounded over $\bar{j} \in \mathbb{N}_0^d$ and $\xi \in [-4, 4]^d$. For this, we first observe that

$$\begin{aligned} F_{\bar{\alpha}, s, \bar{j}}(\xi) &= \left(\sum_{i=1}^d 2^{-\|\bar{j}/\bar{\alpha}\|_\infty} (1 + 2^{2j_i} \xi_i^2)^{1/(2\alpha_i)} \right)^s \\ &= \left(\sum_{i=1}^d \left(2^{-2\alpha_i \|\bar{j}/\bar{\alpha}\|_\infty} + 2^{-2\alpha_i (\|\bar{j}/\bar{\alpha}\|_\infty - j_i/\alpha_i)} \xi_i^2 \right)^{1/(2\alpha_i)} \right)^s, \end{aligned}$$

with quantities $2^{-2\alpha_i \|\bar{j}/\bar{\alpha}\|_\infty}$ and $2^{-2\alpha_i (\|\bar{j}/\bar{\alpha}\|_\infty - j_i/\alpha_i)}$ all positive and never larger than one. This immediately implies

$$(41) \quad \sup_{\substack{\bar{j} \in \mathbb{N}_0^d \\ \xi \in [-4, 4]^d}} |F_{\bar{\alpha}, s, \bar{j}}(\xi)| \lesssim 1.$$

Next, we determine the partial derivatives of $m_{\bar{\alpha}, s}$. They are given by

$$\begin{aligned} \partial_\ell m_{\bar{\alpha}, s}(\xi) &= \frac{s}{\alpha_\ell} m_{\bar{\alpha}, s-1}(\xi) \xi_\ell \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 2}, \\ \partial_\ell^2 m_{\bar{\alpha}, s}(\xi) &= \frac{s(s-1)}{\alpha_\ell^2} m_{\bar{\alpha}, s-2}(\xi) \xi_\ell^2 \langle \xi_\ell \rangle^{\frac{2}{\alpha_\ell} - 4} + \frac{s(s-1)}{\alpha_\ell} m_{\bar{\alpha}, s-1}(\xi) \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 4} \left(\frac{1}{\alpha_\ell} - 1 \right) \xi_\ell^2 + 1, \end{aligned}$$

where we use the abbreviation $\langle \xi_\ell \rangle$ for $(1 + \xi_\ell^2)^{1/2}$. We deduce the estimates

$$\begin{aligned} |\partial_\ell m_{\bar{\alpha}, s}(\xi)| &\lesssim |m_{\bar{\alpha}, s-1}(\xi)| \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 1}, \\ |\partial_\ell^2 m_{\bar{\alpha}, s}(\xi)| &\lesssim |m_{\bar{\alpha}, s-2}(\xi)| \langle \xi_\ell \rangle^{\frac{2}{\alpha_\ell} - 2} + |m_{\bar{\alpha}, s-1}(\xi)| \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 2}, \end{aligned}$$

and thus obtain, using $\langle 2^{j_\ell} \xi_\ell \rangle \leq 2^{j_\ell} \langle \xi_\ell \rangle$,

$$\begin{aligned} |\partial_\ell F_{\bar{\alpha}, s, \bar{j}}(\xi)| &\lesssim 2^{-s \|\bar{j}/\bar{\alpha}\|_\infty} 2^{j_\ell} |m_{\bar{\alpha}, s-1}(2^{\bar{j}} \xi)| \langle 2^{j_\ell} \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 1} \lesssim 2^{-\|\bar{j}/\bar{\alpha}\|_\infty} 2^{j_\ell/\alpha_\ell} |F_{\bar{\alpha}, s-1, \bar{j}}(\xi)| \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 1}, \\ |\partial_\ell^2 F_{\bar{\alpha}, s, \bar{j}}(\xi)| &\lesssim 2^{-s \|\bar{j}/\bar{\alpha}\|_\infty} 2^{2j_\ell} \left(|m_{\bar{\alpha}, s-2}(2^{\bar{j}} \xi)| \langle 2^{j_\ell} \xi_\ell \rangle^{\frac{2}{\alpha_\ell} - 2} + |m_{\bar{\alpha}, s-1}(2^{\bar{j}} \xi)| \langle 2^{j_\ell} \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 2} \right) \\ &\lesssim 2^{-2 \|\bar{j}/\bar{\alpha}\|_\infty} 2^{2j_\ell/\alpha_\ell} |F_{\bar{\alpha}, s-2, \bar{j}}(\xi)| \langle \xi_\ell \rangle^{\frac{2}{\alpha_\ell} - 2} + 2^{-\|\bar{j}/\bar{\alpha}\|_\infty} 2^{j_\ell/\alpha_\ell} |F_{\bar{\alpha}, s-1, \bar{j}}(\xi)| \langle \xi_\ell \rangle^{\frac{1}{\alpha_\ell} - 2}. \end{aligned}$$

Taking (41) into account, we realize that the term $|F_{\bar{\alpha},s-2,\bar{j}}(\xi)|\langle\xi_\ell\rangle^{\frac{2}{\alpha_\ell}-2}$ and the term $|F_{\bar{\alpha},s-1,\bar{j}}(\xi)|\langle\xi_\ell\rangle^{\frac{1}{\alpha_\ell}-2}$ are uniformly bounded in the range $\xi \in [-4, 4]^d$ with respect to $\bar{j} \in \mathbb{N}_0^d$. Since further $2^{-2\|\bar{j}/\bar{\alpha}\|_\infty}2^{2j_\ell/\alpha_\ell} \leq 1$ and $2^{-\|\bar{j}/\bar{\alpha}\|_\infty}2^{j_\ell/\alpha_\ell} \leq 1$, Step 1 is finished.

Step 2. For the proof of the converse inequality $\|f\|_{\widetilde{W}_p^{s,\bar{\alpha}}} \lesssim \|f\|_{W_p^{s,\bar{\alpha}}}$ we argue analogously to Step 1 and use this time the multiplier

$$\widetilde{M}_{\bar{\alpha},s,\bar{j}}(\xi) := \frac{\psi_{\bar{j}}(\xi)2^{s\|\bar{j}/\bar{\alpha}\|_\infty}}{m_{\bar{\alpha},s}(\xi)}.$$

It is well-defined since $m_{\bar{\alpha},s} > 0$, and we have, using the same notation as in Step 1,

$$\widetilde{M}_{\bar{\alpha},s,\bar{j}}(2^{j_1}\xi_1, \dots, 2^{j_d}\xi_d) = \frac{\widetilde{\psi}_{s(\bar{j})}(\xi_1, \dots, \xi_d)}{2^{-s\|\bar{j}/\bar{\alpha}\|_\infty}m_{\bar{\alpha},s}(2^{j_1}\xi_1, \dots, 2^{j_d}\xi_d)}.$$

Again, it is not difficult to check that for every component the second derivatives are bounded on $[-4, 4]^d$ independently of \bar{j} .

By Lemma 3.1, applied with $\rho_{\bar{j}} := \widetilde{M}_{\bar{\alpha},s,\bar{j}}$, $f_{\bar{j}} := \mathcal{F}^{-1}m_{\bar{\alpha},s}\varphi_{\bar{j}}\mathcal{F}f$, and $r := 2$, we get

$$\begin{aligned} \|f\|_{\widetilde{W}_p^{s,\bar{\alpha}}} &\asymp \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{2s\|\bar{j}/\bar{\alpha}\|_\infty} |\mathcal{F}^{-1}\varphi_{\bar{j}}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p \\ &= \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} 2^{2s\|\bar{j}/\bar{\alpha}\|_\infty} |\mathcal{F}^{-1}\psi_{\bar{j}}m_{\bar{\alpha},s}^{-1}m_{\bar{\alpha},s}\varphi_{\bar{j}}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p \\ &= \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} |\mathcal{F}^{-1}\widetilde{M}_{\bar{\alpha},s,\bar{j}}m_{\bar{\alpha},s}\varphi_{\bar{j}}\mathcal{F}f|^2 \right)^{\frac{1}{2}} \right\|_p \\ &\lesssim \left(\sup_{\bar{j} \in \mathbb{N}_0^d} \|\widetilde{M}_{\bar{\alpha},s,\bar{j}}(2^{j_1}\cdot, \dots, 2^{j_d}\cdot)\|_{S_2^2W} \right) \cdot \left\| \left(\sum_{\bar{j} \in \mathbb{N}_0^d} |(\mathcal{F}^{-1}[\varphi_{\bar{j}}m_{\bar{\alpha},s}\mathcal{F}f])|^2 \right)^{\frac{1}{2}} \right\|_p \\ &\lesssim \left\| \mathcal{F}^{-1}[m_{\bar{\alpha},s}\mathcal{F}f] \right\|_p \asymp \|f\|_{W_p^{s,\bar{\alpha}}}. \end{aligned}$$

□

Remark 6.1 *We mention that, in contrast to this result, in case $A = B$ we only have coincidence when $p = q = 2$. A proof can be found in [1].*

As a direct consequence of Theorem 6.1 and Theorem 4.2, we obtain new characterizations of classical Sobolev spaces via hyperbolic wavelets.

Theorem 6.2 *Let $1 < p < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Let further ψ_0, ψ be wavelets satisfying (K) and (L) with*

$$K, L > \sigma_{p,2} + |s|/\alpha_{\min}.$$

Then any $f \in \mathcal{S}'(\mathbb{R}^d)$ belongs to $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ if and only if it can be represented as

$$(42) \quad f = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{k \in \mathbb{Z}^d} \lambda_{\bar{j},k} \psi_{\bar{j},k}$$

with $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,2}^{s,\bar{\alpha}}$. The representation (42) converges unconditionally in $\mathcal{S}'(\mathbb{R}^d)$ and in $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$. In addition, $(\psi_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ is an unconditional basis in $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$. The sequence of coefficients $\lambda := \lambda(f) = (\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ is uniquely determined via

$$\lambda_{\bar{j},\bar{k}} := \lambda_{\bar{j},\bar{k}}(f) = 2^{|\bar{j}|_1} \langle f, \psi_{\bar{j},\bar{k}} \rangle$$

and we have the wavelet isomorphism (equivalent norm)

$$\|f\|_{W_p^{s,\bar{\alpha}}(\mathbb{R}^d)} \asymp \|\lambda(f)\|_{\tilde{f}_{p,2}^{s,\bar{\alpha}}} \quad , \quad f \in W_p^{s,\bar{\alpha}}(\mathbb{R}^d).$$

Analogously, combining Theorem 6.1 with Theorem 5.4, we also derive new characterizations of Sobolev spaces with the hyperbolic Haar system \mathcal{H}_d from (24).

Theorem 6.3 *Let $1 < p < \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ such that $\sum_{i=1}^d \alpha_i = d$. Further, assume*

$$|s|/\alpha_{\min} < \min \left\{ \frac{1}{p}, 1 - \frac{1}{p} \right\}.$$

Then the Haar system $\mathcal{H}_d = (h_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ from (24) constitutes an unconditional basis of $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ with associated sequence space $\tilde{f}_{p,2}^{s,\bar{\alpha}}$. The unique sequence of basis coefficients for $f \in W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ is determined by $\lambda := \lambda(f) = (\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ with

$$(43) \quad \lambda_{\bar{j},\bar{k}} := \lambda_{\bar{j},\bar{k}}(f) = 2^{|\bar{j}|_1} \langle f, h_{\bar{j},\bar{k}} \rangle.$$

Further, we have the wavelet isomorphism (equivalent norm)

$$\|f\|_{W_p^{s,\bar{\alpha}}(\mathbb{R}^d)} \asymp \|\lambda(f)\|_{\tilde{f}_{p,2}^{s,\bar{\alpha}}} \quad , \quad f \in W_p^{s,\bar{\alpha}}(\mathbb{R}^d).$$

In addition, those elements of $\mathcal{S}'(\mathbb{R}^d)$ belonging to $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$ are characterized by either of the following two criteria:

(i) f can be represented as a sum

$$(44) \quad f = \sum_{\bar{j} \in \mathbb{N}_0^d} \sum_{\bar{k} \in \mathbb{Z}^d} \lambda_{\bar{j},\bar{k}} h_{\bar{j},\bar{k}} \quad \text{converging (weak*ly) in } \mathcal{S}'(\mathbb{R}^d)$$

with coefficients $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,2}^{s,\bar{\alpha}}$ (with respect to some chosen ordering).

(ii) With $\lambda(f)$ being defined as in (43), it holds

$$\lambda(f) = (\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}} \in \tilde{f}_{p,2}^{s,\bar{\alpha}}.$$

In both cases, the sequence $(\lambda_{\bar{j},\bar{k}})_{\bar{j},\bar{k}}$ is necessarily the sequence of basis coefficients and the representation (44) converges unconditionally to f in $W_p^{s,\bar{\alpha}}(\mathbb{R}^d)$.

7 Hyperbolic and classical (anisotropic) BLT spaces

The next and final theorem of this paper complements the statement of Theorem 6.1, showing that in general the spaces $\widetilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $A_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$, with $A \in \{B, F\}$, do not coincide.

Theorem 7.1 *Let $0 < p, q \leq \infty$, $s \in \mathbb{R}$, and $\bar{\alpha} = (\alpha_1, \dots, \alpha_d) > 0$ with $\sum_{i=1}^d \alpha_i = d$.*

(i) *If $B_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \widetilde{B}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ then $p = q = 2$.*

(ii) *In the range $0 < p < \infty$: If $F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \widetilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ then $q = 2$ and $1 < p < \infty$.*

Remark 7.1 *The Besov result (i) follows directly from the very general investigations on embeddings between decomposition spaces conducted in [56]. Even more, the findings there allow to strengthen the statement to more general independent parameters, namely*

$$B_{p_1,q_1}^{s_1,\bar{\alpha}_1}(\mathbb{R}^d) = \widetilde{B}_{p_2,q_2}^{s_2,\bar{\alpha}_2}(\mathbb{R}^d) \quad \Leftrightarrow \quad p_1 = p_2 = q_1 = q_2 = 2 \text{ and } \bar{\alpha}_1 = \bar{\alpha}_2 \text{ and } s_1 = s_2.$$

The results of [56], however, are not applicable in the proof of (ii) since Triebel-Lizorkin spaces do not fit into the decomposition space framework. In the sequel, we will therefore give a proof for the F -case (ii) which by slight modifications would also provide a direct way to establish the B -case (i).

Before we start, let us remind ourselves that the converse statement of (ii), the coincidence of $F_{p,q}^{s,\bar{\alpha}}$ and $\widetilde{F}_{p,q}^{s,\bar{\alpha}}$ when $1 < p < \infty$ and $q = 2$, is given by Theorem 6.1. The coincidence of $B_{p,q}^{s,\bar{\alpha}}$ and $\widetilde{B}_{p,q}^{s,\bar{\alpha}}$ when $p = q = 2$, the converse of (i), is further observed in Remark 6.1.

Proof. [of Theorem 7.1 (ii)] *Step 1: Preparation.* Fix an anisotropy vector $\bar{\alpha} = (\alpha_1, \dots, \alpha_d)$ and consider a univariate resolution of unity $(\theta_j)_{j \in \mathbb{N}_0}$ of the following form:

The generator $\theta_0 \in \mathcal{S}(\mathbb{R})$ shall satisfy

$$\text{supp } \theta_0 \subset [-2^{\alpha_{\min}/3}, 2^{\alpha_{\min}/3}] \quad \text{and} \quad \theta_0 = 1 \quad \text{on} \quad [-1, 1],$$

and the functions θ_j for $j \in \mathbb{N}$ shall be obtained via $\theta_j(\cdot) := \theta_0(2^{-j}\cdot) - \theta_0(2^{-(j-1)}\cdot)$.

Using $(\theta_j)_{j \in \mathbb{N}_0}$, we can then construct two multivariate resolutions of unity on \mathbb{R}^d . First, via simple tensorization, we get the hyperbolic resolution $(\theta_{\bar{j}})_{\bar{j} \in \mathbb{N}_0^d}$ with

$$\theta_{\bar{j}} := \theta_{j_1} \otimes \dots \otimes \theta_{j_d}, \quad \bar{j} = (j_1, \dots, j_d) \in \mathbb{N}_0^d.$$

It clearly fulfills all the specifications formulated at the beginning of Section 3.

Second, putting $\varphi_0^{\bar{\alpha}} := \theta_0 \otimes \dots \otimes \theta_0$ and

$$\varphi_j^{\bar{\alpha}} := \varphi_0^{\bar{\alpha}}(2^{-j\bar{\alpha}}\cdot) - \varphi_0^{\bar{\alpha}}(2^{-(j-1)\bar{\alpha}}\cdot) \quad \text{for } j \in \mathbb{N},$$

we obtain $(\varphi_j^{\bar{\alpha}})_{j \in \mathbb{N}_0}$, which is a classical anisotropic resolution of unity in compliance with the definition from Subsection 2.1.

For parameters $\alpha > 0$ and $\ell \in \mathbb{N}$, let us next introduce the intervals

$$I_\ell^\alpha := 2^{(\ell-1)\alpha} \cdot [2^{\alpha_{\min}/3}, 2^\alpha] \quad \text{and} \quad J_\ell^\alpha := [-2^{\ell\alpha}, 2^{\ell\alpha}].$$

Then $\theta_0(2^{-\ell\alpha}\cdot) = 1$ on J_ℓ^α and $\theta_0(2^{-\ell\alpha}\cdot) - \theta_0(2^{-(\ell-1)\alpha}\cdot) = 1$ on I_ℓ^α . In particular, $\theta_j = 1$ on I_j^1 for every $j \in \mathbb{N}$ and thus $\theta_{\bar{j}} = 1$ on $I_{j_1}^1 \times \dots \times I_{j_d}^1$. Further, we have $\varphi_0^{\bar{\alpha}}(2^{-j\bar{\alpha}}\cdot) = 1$ on

$J_j^{\alpha_1} \times \dots \times J_j^{\alpha_d}$ and as a consequence $\varphi_j^{\bar{\alpha}} = 1$ on $(J_j^{\alpha_1} \times \dots \times J_j^{\alpha_d}) \setminus (J_{j-1}^{\alpha_1} \times \dots \times J_{j-1}^{\alpha_d})$. This, in turn, implies $\varphi_j^{\bar{\alpha}} = 1$ on the subset $I_j^{\alpha_1} \times \dots \times I_j^{\alpha_{d-1}} \times J_j^{\alpha_d}$.

Observe now that for every $\ell \in \mathbb{N}$ and every $i \in \{1, \dots, d\}$ either $I_\ell^{\alpha_i} \cap I_{[\ell\alpha_i]}^1$ or $I_\ell^{\alpha_i} \cap I_{[\ell\alpha_i]+1}^1$ is a nonempty interval of nonzero length. This is due to the fact that always

$$(45) \quad 2^\gamma \cdot \mathcal{L}(I_\ell^{\alpha_i}) \leq \mathcal{R}(I_{[\ell\alpha_i]}^1) \leq \mathcal{R}(I_\ell^{\alpha_i}) \quad \text{or} \quad 2^\gamma \cdot \mathcal{L}(I_{[\ell\alpha_i]+1}^1) \leq \mathcal{R}(I_\ell^{\alpha_i}) \leq \mathcal{R}(I_{[\ell\alpha_i]+1}^1),$$

where $\mathcal{L}(I)$ and $\mathcal{R}(I)$ denote the left resp. right endpoint of a given interval $I = [a, b]$ and $\gamma = \alpha_{\min}^2/8$. The verification of this fact is postponed to Step 3 at the end of this proof.

As a consequence, for each $i \in \{1, \dots, d\}$ and each $\ell \in \mathbb{N}$, we may pick one of those intersections with nonvanishing interior and denote it by $\tilde{I}_\ell^{(i)}$. Depending on our choice, we then either have

$$(46) \quad \tilde{I}_\ell^{(i)} = I_\ell^{\alpha_i} \cap I_{[\ell\alpha_i]}^1 \quad \text{or} \quad \tilde{I}_\ell^{(i)} = I_\ell^{\alpha_i} \cap I_{[\ell\alpha_i]+1}^1.$$

Due to the nonvanishing interior of $\tilde{I}_\ell^{(i)}$ we can further fix nontrivial functions

$$h_\ell^{[i]} \in \mathcal{S}(\mathbb{R}) \quad \text{with} \quad \text{supp } h_\ell^{[i]} \subset \tilde{I}_\ell^{(i)}, \quad i \in \{1, \dots, d-1\}.$$

With this preparation we are finally ready for the main argumentation.

Step 2: Main Proof. For $\ell \in \mathbb{N}$ let us consider $g_\ell : \mathbb{R} \rightarrow \mathbb{C}$ with the property

$$(47) \quad \text{supp } (\mathcal{F}g_\ell) \subset J_\ell^{\alpha_d} = [-2^{\ell\alpha_d}, 2^{\ell\alpha_d}]$$

and associate a multivariate function $f_\ell : \mathbb{R}^d \rightarrow \mathbb{C}$ defined by its Fourier transform

$$\mathcal{F}f_\ell(\xi_1, \dots, \xi_d) := h_\ell^{[1]}(\xi_1)h_\ell^{[2]}(\xi_2) \cdots h_\ell^{[d-1]}(\xi_{d-1})\mathcal{F}g_\ell(\xi_d),$$

where $h_\ell^{[i]}$ are the functions introduced at the end of Step 1.

Since $\mathcal{F}f_\ell$ is supported inside $I_\ell^{\alpha_1} \times \dots \times I_\ell^{\alpha_{d-1}} \times J_\ell^{\alpha_d}$, on which $\varphi_\ell^{\bar{\alpha}} = 1$ according to our considerations in Step 1, we can easily compute the classical anisotropic Triebel-Lizorkin (quasi-)norm of f_ℓ . Denoting by $(\Delta_j^\varphi)_{j \in \mathbb{N}_0}$ the Littlewood-Paley analysis associated to $(\varphi_j^{\bar{\alpha}})_{j \in \mathbb{N}_0}$, we have

$$\|f_\ell\|_{F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)} = \left\| \left(\sum_{j \geq 0} 2^{jsq} |\Delta_j^\varphi f_\ell|^q \right)^{\frac{1}{q}} \right\|_p = 2^{\ell s} \|\Delta_\ell^\varphi f_\ell\|_p = 2^{\ell s} \|f_\ell\|_p.$$

Moreover, as f_ℓ is a tensor product, we can compute

$$\|f_\ell\|_p = \|\mathcal{F}^{-1}(h_\ell^{[1]})\|_p \cdots \|\mathcal{F}^{-1}(h_\ell^{[d-1]})\|_p \|g_\ell\|_p = C_\ell \|g_\ell\|_p$$

with $C_\ell := C_\ell^{(1)} \cdots C_\ell^{(d-1)}$ and $C_\ell^{(i)} := \|\mathcal{F}^{-1}(h_\ell^{[i]})\|_p$ for $i \in \{1, \dots, d-1\}$. Altogether, we end up with

$$\|f_\ell\|_{F_{p,q}^{s,\bar{\alpha}}} \asymp 2^{\ell s} C_\ell \|g_\ell\|_p.$$

We proceed with the computation of the hyperbolic Triebel-Lizorkin (quasi-)norm of f_ℓ . It follows right from the definition of the intervals $\tilde{I}_\ell^{(i)}$ from (46) that there exist numbers $k_i(\ell) \in \mathbb{N}$, either taking the value $[\ell\alpha_i]$ or the value $[\ell\alpha_i] + 1$, such that $\tilde{I}_\ell^{(i)} \subset I_{k_i(\ell)}^1$.

Hence, due to $\text{supp } \mathcal{F}f_\ell \subset \tilde{I}_\ell^{(1)} \times \cdots \times \tilde{I}_\ell^{(d-1)} \times J_\ell^{\alpha_d}$ the function $\mathcal{F}f_\ell$ is supported inside $I_{k_1(\ell)}^1 \times \cdots \times I_{k_{d-1}(\ell)}^1 \times J_\ell^{\alpha_d}$.

Let now $(\Delta_{\vec{j}}^\theta)_{\vec{j} \in \mathbb{N}_0^d}$ denote the Littlewood-Paley analysis corresponding to $(\theta_{\vec{j}})_{\vec{j} \in \mathbb{N}_0^d}$ and let us abbreviate $k_i(\ell)$ by k_i and the vector $(k_1, \dots, k_{d-1}, j_d)$ by $\bar{\ell}_{j_d}$. Then, since

$$\sum_{j_d=0}^{[\ell\alpha_d]+1} \theta_{\bar{\ell}_{j_d}} = 1 \quad \text{on} \quad I_{k_1}^1 \times \cdots \times I_{k_{d-1}}^1 \times J_\ell^{\alpha_d},$$

we calculate for the hyperbolic Triebel-Lizorkin (quasi-)norm of f_ℓ

$$\begin{aligned} \|f_\ell\|_{\tilde{F}_{p,q}^{s,\bar{\alpha}}} &= \left\| \left(\sum_{\vec{j} \in \mathbb{N}_0^d} 2^{\max\{j_1/\alpha_1, \dots, j_d/\alpha_d\}sq} |\Delta_{\vec{j}}^\theta f_\ell(\cdot)|^q \right)^{1/q} \right\|_p \\ &= \left\| \left(\sum_{j_d=0}^{[\ell\alpha_d]+1} 2^{\max\{k_1/\alpha_1, \dots, k_{d-1}/\alpha_{d-1}, j_d/\alpha_d\}sq} |\Delta_{\bar{\ell}_{j_d}}^\theta f_\ell(\cdot)|^q \right)^{1/q} \right\|_p \\ &\asymp 2^{\ell s} \left\| \left(\sum_{j_d=0}^{[\ell\alpha_d]+1} |\Delta_{\bar{\ell}_{j_d}}^\theta f_\ell(\cdot)|^q \right)^{1/q} \right\|_p = 2^{\ell s} C_\ell \|g_\ell\|_{F_{p,q}^0} \end{aligned}$$

with the same constant C_ℓ as obtained before in the computation of $\|f_\ell\|_{F_{p,q}^{s,\bar{\alpha}}}$.

Now we come to the core argument. Assuming that the spaces $F_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ and $\tilde{F}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d)$ coincide, the associated (quasi-)norms would be equivalent. By our calculations, this would imply that $\|g_\ell\|_{F_{p,q}^0(\mathbb{R})}$ is equivalent to $\|g_\ell\|_{L_p(\mathbb{R})}$ for any band-limited function g_ℓ with frequency support as in (47). Moreover, since the proof holds true for all $\ell \in \mathbb{N}$ this equivalence remains valid for any band-limited function g on \mathbb{R} .

But, as a consequence of Lemma 7.1(iii), since the sequence $(f_N^{(3)})_N$ constructed in its proof consists of band-limited functions, this is only possible in the range $1 < p < \infty$. Furthermore, if $1 < p < \infty$ the band-limited functions are dense in $L_p(\mathbb{R})$ as well as $F_{p,q}^0(\mathbb{R})$. Hence, by Lemma 7.1(i) also $q = 2$ is a necessary condition. It now only remains to verify (45).

Step 3: Proof of (45). We distinguish two cases depending on the size of the quantity $\delta := \ell\alpha_i - [\ell\alpha_i] \in [0, 1)$. Let us subsequently abbreviate $\rho := \frac{\alpha_{\min}}{4+\alpha_{\min}}$ and $\sigma := \frac{8\alpha_{\min}}{3\alpha_{\min}+12} = \frac{2}{3}(1-\rho)\alpha_{\min}$. Recalling that $\alpha_{\min} = \min_{i \in \{1, \dots, d\}} \{\alpha_i\} \in (0, 1]$, we note that $\rho \in (0, \frac{1}{3}]$ and $\sigma \in (0, \frac{2}{3}]$.

In case $\delta \in [0, \sigma)$ we have $\delta \leq (1-\rho)(\alpha_i - \alpha_{\min}/3)$ and thus

$$\begin{aligned} \log_2 \mathcal{R}(I_{[\ell\alpha_i]}^1) &= [\ell\alpha_i] = \ell\alpha_i - \delta \geq \ell\alpha_i - (1-\rho)(\alpha_i - \alpha_{\min}/3) \\ &= (\ell-1)\alpha_i + \alpha_{\min}/3 + \rho(\alpha_i - \alpha_{\min}/3) \geq \log_2 \mathcal{L}(I_\ell^{\alpha_i}) + \frac{2}{3}\rho\alpha_{\min}, \end{aligned}$$

where we used $\mathcal{L}(I_\ell^{\alpha_i}) = 2^{(\ell-1)\alpha_i + \alpha_{\min}/3}$ and $\mathcal{R}(I_{[\ell\alpha_i]}^1) = 2^{[\ell\alpha_i]}$. In view of $\gamma = \alpha_{\min}^2/8 < \frac{2}{3}\rho\alpha_{\min}$ and since we always have $[\ell\alpha_i] \leq \ell\alpha_i \leq [\ell\alpha_i] + 1$, i.e.

$$(48) \quad \log_2 \mathcal{R}(I_{[\ell\alpha_i]}^1) \leq \log_2 \mathcal{R}(I_\ell^{\alpha_i}) \leq \log_2 \mathcal{R}(I_{[\ell\alpha_i]+1}^1),$$

the left inequality in (45) is hence valid in the respective range of δ .

In case $\delta \in [\sigma, 1)$ we first estimate

$$\delta > \frac{7\alpha_{\min}}{3\alpha_{\min} + 1} = (1 - \rho) + \rho\alpha_{\min}/3 \geq (1 - \rho)\alpha_{\min} + \rho\alpha_{\min}/3,$$

from which we deduce, using $\ell\alpha_i = \lfloor \ell\alpha_i \rfloor + \delta$ and $\mathcal{L}(I_{\lfloor \ell\alpha_i \rfloor + 1}^1) = 2^{\lfloor \ell\alpha_i \rfloor + \alpha_{\min}/3}$,

$$\log_2 \mathcal{R}(I_\ell^{\alpha_i}) = \ell\alpha_i > \lfloor \ell\alpha_i \rfloor + (1 - \rho)\alpha_{\min} + \rho\alpha_{\min}/3 = \log_2 \mathcal{L}(I_{\lfloor \ell\alpha_i \rfloor + 1}^1) + \frac{2}{3}(1 - \rho)\alpha_{\min}.$$

This time, again taking into account (48) and $\gamma = \alpha_{\min}^2/8 < \frac{2}{3}(1 - \rho)\alpha_{\min}$, the right inequality in (45) holds true. Altogether, the proof of (45) is thus finished. \square

The behavior of the L_p -(quasi-)norms in relation to the $A_{p,q}^0$ -(quasi-)norms is crucial for the proof of Theorem 7.1. Concretely, we have shown for $A \in \{B, F\}$

$$(49) \quad A_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) = \widetilde{A}_{p,q}^{s,\bar{\alpha}}(\mathbb{R}^d) \iff \|f\|_{A_{p,q}^0(\mathbb{R})} \asymp \|f\|_{L_p(\mathbb{R})} \quad \text{for band-limited functions } f.$$

On the right-hand side, the (quasi-)norms are thereby all classical and only the univariate case matters. Using known embedding theorems, the exact parameters for equality could therefore be determined (see [49] Section 2.3.2 or [46] Theorem 3.1.1., for example).

Preferring a direct and shorter route, the following lemma provides a simple and quantitative argument for what we need. It investigates the behavior of the respective (quasi-)norms for certain sequences of test functions. As a consequence of statement (i), we extract the necessity $p = q = 2$ for equality in (49). From (ii) we further obtain $p = q$ in the B-case. Statement (iii) yields $1 < p < \infty$ in the F-case. Altogether, this shows that the Sobolev spaces in Theorem 6.1 are precisely those, where equality holds true.

Lemma 7.1 *Assume $0 < p < \infty$, $0 < q \leq \infty$, $A \in \{B, F\}$. There are sequences $(f_N^{(i)})_{N \in \mathbb{N}}$, $i \in \{1, 2, 3\}$, of functions on \mathbb{R} such that*

$$(i) \quad \|f_N^{(1)}\|_p \asymp N^{1/2} \quad \text{and} \quad \|f_N^{(1)}\|_{A_{p,q}^0(\mathbb{R})} \asymp N^{1/q},$$

$$(ii) \quad \|f_N^{(2)}\|_p \asymp N^{1/p} \quad \text{and} \quad \|f_N^{(2)}\|_{B_{p,q}^0(\mathbb{R})} \asymp N^{1/q},$$

$$(iii) \quad \|f_N^{(3)}\|_p \asymp 2^{N(1-1/p)} \quad \text{and} \quad \|f_N^{(3)}\|_{F_{p,q}^0(\mathbb{R})} \gtrsim \begin{cases} 1 & , p < 1, \\ N^{1/p} & , p \geq 1. \end{cases}$$

Remark 7.2 *In case $q = \infty$ we need to interpret $N^{1/q} \asymp 1$. Further, the case $A = B$ with $p = \infty$ is not considered in Lemma 7.1. By an analogous argument, one can show however that (ii) holds true also for $p = \infty$. So, for $B_{\infty,q}^0(\mathbb{R})$ we have the necessary condition $q = \infty$ to be equivalent to $L_\infty(\mathbb{R})$. It is further not difficult to show that the sequence $(f_N^{(1)})_{N \in \mathbb{N}_0}$ from (i) fulfills $\|f_N^{(1)}\|_\infty \nearrow \infty$ whereas $\|f_N^{(1)}\|_{B_{\infty,\infty}^0(\mathbb{R})} \asymp 1$. Hence, $B_{\infty,\infty}^0(\mathbb{R}) \neq L_\infty(\mathbb{R})$.*

Proof. ad (i): We provide the proof for $q < \infty$. Let $\bar{\varepsilon} = (\varepsilon_0, \varepsilon_1, \dots) \in \{-1, 1\}^{\mathbb{N}_0}$ and define

$$f_{N,\bar{\varepsilon}} := \sum_{j=0}^N \varepsilon_j \sum_{k=0}^{2^j} \psi_{j,k},$$

where $(\psi_{j,k})_{j,k}$ shall be a compactly supported, orthogonal, and L_∞ -normalized wavelet system with sufficient vanishing moments and smoothness to characterize the space $A_{p,q}^0(\mathbb{R})$. Further, for each $j \in \mathbb{N}_0$ and $k \in \{0, \dots, 2^j\}$, we assume the support condition $\text{supp}(\psi_{j,k}) \subset [0, 1]$.

Now we note that in a univariate setting, as considered here, we have the coincidence $\tilde{A}_{p,q}^0(\mathbb{R}) = A_{p,q}^0(\mathbb{R})$. Hence, using the wavelet isomorphism established by Theorem 4.2 for the F-scale and taking into account Remark 4.1 for the B-scale, we immediately obtain

$$\begin{aligned} \|f_{N,\bar{\varepsilon}}\|_{B_{p,q}^0(\mathbb{R})} &\asymp \left(\sum_{j=0}^N \left\| \sum_{k=0}^{2^j} \chi_{j,k} \right\|_p^q \right)^{1/q} \asymp N^{1/q}, \\ \|f_{N,\bar{\varepsilon}}\|_{F_{p,q}^0(\mathbb{R})} &\asymp \left\| \left(\sum_{j=0}^N \left| \sum_{k=0}^{2^j} \chi_{j,k} \right|^q \right)^{1/q} \right\|_p \asymp N^{1/q}, \end{aligned}$$

whereby the (quasi-)norms on the left-hand side do not depend on the choice of $\bar{\varepsilon}$.

From here we proceed with a probabilistic argument and interpret $\bar{\varepsilon}$ as a Rademacher random variable. Then, for the expectation of the L_p -(quasi-)norms over $\bar{\varepsilon}$,

$$\mathbb{E}_{\bar{\varepsilon}}(\|f_{N,\bar{\varepsilon}}\|_p^p) = \int_0^1 \int_{\mathbb{R}} \left| \sum_{j=0}^N r_j(t) \sum_{k=0}^{2^j} \psi_{j,k}(x) \right|^p dx dt,$$

where $r_j(t) := \text{sgn}(\sin 2^j \pi t)$ is the j -th Rademacher function. Applying Khintchine's inequality, we obtain from this

$$\begin{aligned} \mathbb{E}_{\bar{\varepsilon}}(\|f_{N,\bar{\varepsilon}}\|_p^p) &= \int_0^1 \int_0^1 \left| \sum_{j=0}^N r_j(t) \sum_{k=0}^{2^j} \psi_{j,k}(x) \right|^p dt dx \\ &\asymp \int_0^1 \left(\sum_{j=0}^N \left| \sum_{k=0}^{2^j} \psi_{j,k}(x) \right|^2 \right)^{p/2} dx. \end{aligned}$$

Next, we define for $j \in \mathbb{N}_0$ the auxiliary functions

$$F_j := \frac{1}{M} \left| \sum_{k=0}^{2^j} \psi_{j,k}(x) \right|^2 \quad \text{with } M > 0 \text{ such that } \|F_j\|_\infty \leq 1.$$

Observe that $0 \leq F_j \leq 1$. Then, in case $0 < p \leq 2$,

$$N^{-p/2} \cdot \mathbb{E}_{\bar{\varepsilon}}(\|f_{N,\bar{\varepsilon}}\|_p^p) \asymp \int_0^1 \left(\frac{1}{N} \sum_{j=0}^N F_j(x) \right)^{p/2} dx \geq \frac{1}{N} \int_0^1 \sum_{j=0}^N F_j(x) dx,$$

and $\int_0^1 F_j(x) dx \asymp 1$. Hence $\mathbb{E}_{\bar{\varepsilon}}(\|f_{N,\bar{\varepsilon}}\|_p^p) \gtrsim N^{p/2}$. Also, since $2/p \geq 1$, with Hölder

$$\mathbb{E}_{\bar{\varepsilon}}(\|f_{N,\bar{\varepsilon}}\|_p^p) \asymp \int_0^1 \left(\sum_{j=0}^N F_j(x) \right)^{p/2} dx \lesssim \left(\int_0^1 \sum_{j=0}^N F_j(x) dx \right)^{p/2} \asymp N^{p/2}.$$

In case $2 < p < \infty$, we again argue with Hölder

$$\mathbb{E}_{\varepsilon}(\|f_{N,\varepsilon}\|_p^p) \asymp \int_0^1 \left(\sum_{j=0}^N F_j(x) \right)^{p/2} dx \gtrsim \left(\int_0^1 \sum_{j=0}^N F_j(x) dx \right)^{p/2} \asymp N^{p/2}.$$

Further, since $2/p < 1$,

$$N^{-p/2} \cdot \mathbb{E}_{\varepsilon}(\|f_{N,\varepsilon}\|_p^p) \asymp \int_0^1 \left(\frac{1}{N} \sum_{j=0}^N F_j(x) \right)^{p/2} dx \lesssim \frac{1}{N} \int_0^1 \sum_{j=0}^N F_j(x) dx \asymp 1.$$

Altogether, these estimates show $\mathbb{E}_{\varepsilon}(\|f_{N,\varepsilon}\|_p^p) \asymp N^{p/2}$. As a consequence, we can choose $f_N := f_{N,\varepsilon(N)}$ such that $\|f_N\|_p^p \asymp N^{p/2}$, or equivalently $\|f_N\|_p \asymp \sqrt{N}$.

ad (ii): With the same wavelet system $(\psi_{j,k})_{j,k}$ as before, L_{∞} -normalized, define

$$f_N := \sum_{j=0}^N 2^{j/p} \psi_{j,k(j)},$$

where $k(j)$ is chosen such that the (spatial) support of the wavelets is mutually disjoint. Then, using again the wavelet isomorphism from Theorem 4.2 and Remark 4.1, we deduce

$$\|f_N\|_{B_{p,q}^0(\mathbb{R})} = \|f_N\|_{\tilde{B}_{p,q}^0(\mathbb{R})} \asymp \left(\sum_{j=0}^N \left\| 2^{j/p} \chi_{j,k(j)} \right\|_p^q \right)^{1/q} = \left(\sum_{j=0}^N (2^{j/p} 2^{-j/p})^q \right)^{1/q} \asymp N^{1/q}.$$

For the L_p -(quasi-)norm we obtain, due to the disjoint support,

$$\|f_N\|_{L_p(\mathbb{R})} = \left\| \sum_{j=0}^N 2^{j/p} \psi_{j,k(j)} \right\|_p = \left(\sum_{j=0}^N 2^j \|\psi_{j,k(j)}\|_p^p \right)^{1/p} \asymp N^{1/p}.$$

ad (iii): Finally, let $(\varphi_j)_j$ be a (standard) dyadic resolution of unity, with $\varphi_0 = 1$ in a neighborhood of 0 and $\varphi_1 = \varphi_0(\cdot/2) - \varphi_0$, and put

$$f_N := \mathcal{F}^{-1} \varphi_0(2^{-N} \cdot).$$

Then $f_N = 2^N F_1(2^N \cdot)$. For the L_p -(quasi-)norm we thus compute

$$\|f_N\|_{L_p(\mathbb{R})} = 2^N \|F_1(2^N \cdot)\|_p \asymp 2^{N(1-1/p)}.$$

Turning to the $F_{p,q}^0(\mathbb{R})$ -(quasi-)norm, for $N \geq 2$, we calculate, writing $\Phi_0 := \mathcal{F}^{-1} \varphi_0$ and $\Phi_1 := \mathcal{F}^{-1} \varphi_1$,

$$\begin{aligned} \|f_N\|_{F_{p,q}^0(\mathbb{R})} &= \left\| \left(\sum_{j=0}^{\infty} \left| (\mathcal{F}^{-1}(\varphi_j \cdot \varphi_0(2^{-N} \cdot))) (\cdot) \right|^q \right)^{1/q} \right\|_p \\ &\geq \left\| \left(\sum_{j=0}^{N-1} \left| (\mathcal{F}^{-1} \varphi_j) (\cdot) \right|^q \right)^{1/q} \right\|_p \\ &= \left\| \left(|\Phi_0|^q + \sum_{j=1}^{N-1} \left| 2^{j-1} \Phi_1(2^{j-1} \cdot) \right|^q \right)^{1/q} \right\|_p \\ &= \left\| \left(|\Phi_0|^q + \sum_{j=0}^{N-2} \left| 2^j \Phi_1(2^j \cdot) \right|^q \right)^{1/q} \right\|_p. \end{aligned}$$

Note that Φ_1 has (infinitely many) vanishing moments and is thus oscillatory. Assuming w.l.o.g. $|\Phi_1| > \delta$ on a set $I \subset [1, 2)$, with $\delta > 0$ being some fixed constant, we can proceed

$$\begin{aligned} \|f_N\|_{F_{p,q}^0(\mathbb{R})} &\gtrsim \left\| \left(\sum_{j=0}^{N-2} \left| 2^j \chi_I(2^j \cdot) \right|^q \right)^{1/q} \right\|_p \\ &= \left\| \sum_{j=0}^{N-2} 2^j \chi_I(2^j \cdot) \right\|_p = \left(\sum_{j=0}^{N-2} 2^{jp} \int_{\mathbb{R}} \chi_I(2^j x) dx \right)^{1/p} \\ &\asymp \left(\sum_{j=0}^{N-2} 2^{j(p-1)} \right)^{1/p} \gtrsim \begin{cases} 1 & , p < 1, \\ N^{1/p} & , p \geq 1. \end{cases} \end{aligned}$$

□

Acknowledgment. We warmly thank Glenn Byrenheid, Nadi'ia Derevianko, Dinh Dũng, Dorothee Haroske, Peter Oswald, Andreas Seeger, Winfried Sickel, Vladimir N. Temlyakov, Hans Triebel, Felix Voigtländer, and Stephane Jaffard for stimulating discussions and comments. This work has been partially supported by the ANR grant *AMATIS* (ANR2011 BS01 011 02) and the CNRS, Groupe de Recherche *Analyse Multifractale*. In addition we gratefully acknowledge support by the German Research Foundation (DFG) and the Emmy-Noether programme, UI-403/1-1.

References

- [1] P. Abry, M. Clausel, S. Jaffard, S. G. Roux, and B. Vedel. The hyperbolic wavelet transform: an efficient tool for multifractal analysis of anisotropic textures. *Rev. Mat. Iberoam.*, 1:1–36, 2015.
- [2] P. Abry, S. Jaffard, S. G. Roux, B. Vedel, J. Coddington, L. Daffner, P. Borgnat, N. Tremblay, A. Klein, H. Wendt, and P. Messier. Multiscale anisotropic texture analysis and classification of photographic prints. *IEEE Signal Process. Mag.*, 32(4):18–27, 2015.
- [3] A. V. Andrianov. Approximation of MH_q^r -class functions by Haar polynomials. *Math. Notes*, 66(3):261–270, 1999.
- [4] A. Bonami and A. Estrade. Anisotropic analysis of some Gaussian models. *J. Fourier Anal. Appl.*, 9:215–236, 2003.
- [5] M. Bownik. Atomic and molecular decomposition of anisotropic Besov spaces. *Math. Z.*, 250:539–571, 2005.
- [6] M. Bownik and K. P. Ho. Atomic and molecular decomposition of anisotropic Triebel-Lizorkin spaces. *Trans. Amer. Math. Soc.*, 385(4):1469–1510, 2005.
- [7] E. J. Cands and L. Demanet. The curvelet representation of wave propagators is optimally sparse. *Comm. Pure Appl. Math.*, 58:1472–1508, 2005.
- [8] E. J. Cands and D. L. Donoho. Curvelets - a surprisingly effective nonadaptive representation for objects with edges. In C. Rabut, A. Cohen, and L. Schumaker, editors, *Curves and Surfaces*, pages 105–120. Vanderbilt University Press, 2000.

- [9] E. J. Cands and D. L. Donoho. New tight frames of curvelets and optimal representations of objects with piecewise C^2 singularities. *Comm. Pure Appl. Math.*, 56:219–266, 2002.
- [10] C. K. Chui and J.-Z. Wang. On compactly supported spline wavelets and a duality principle. *Trans. Amer. Math. Soc.*, 330(2):903–915, 1992.
- [11] D. Dũng, V. N. Temlyakov, and T. Ullrich. Hyperbolic Cross Approximation. *arXiv e-prints*, 2015. arXiv:1601.03978 [math.NA].
- [12] D. Dũng and T. Ullrich. N-widths and ε -dimensions for high-dimensional approximations. *Found. Comput. Math.*, 13:965–1003, 2013.
- [13] L. Demanet and L. Ying. Wave atoms and sparsity of oscillatory patterns. *Appl. Comput. Harmon. Anal.*, 23(3):368–387, 2007.
- [14] N. Derevianko and T. Ullrich. A higher order Faber spline basis for sampling discretization of functions. *arXiv e-prints*, pages 1–34, 2019.
- [15] R. A. DeVore, S. V. Konyagin, and V. N. Temlyakov. Hyperbolic wavelet approximation. *Constr. Approx.*, 14:1–26, 1998.
- [16] R. E. Edwards. *Functional analysis: Theory and applications*. Athena series. Holt, Rinehart and Winston, 1965.
- [17] G. Faber. Über die Orthogonalfunktionen des Herrn Haar. *Jahresber. Deutsch. Math. Verein.*, 19:104–112, 1910.
- [18] C. Fefferman and E. M. Stein. Some maximal inequalities. *Amer. J. Math.*, 93:107–115, 1971.
- [19] M. Frazier and B. Jawerth. Decomposition of Besov spaces. *Indiana Univ. Math. J.*, 34(4):777–799, 1985.
- [20] M. Frazier and B. Jawerth. A discrete transform and decompositions of distribution spaces. *J. Funct. Anal.*, 93(1):34–170, 1990.
- [21] G. Garrigós, A. Seeger, and T. Ullrich. On uniform boundedness of dyadic averaging operators in spaces of Hardy–Sobolev type. *Anal. Math.*, 43(2):267–278, 2017.
- [22] G. Garrigós, A. Seeger, and T. Ullrich. Basis properties of the Haar system in limiting Besov spaces. *arXiv e-prints*, 2019. arXiv:1901.09117 [math.CA].
- [23] G. Garrigós, A. Seeger, and T. Ullrich. Haar frames and Haar multipliers in the absence of unconditionality. 2019. preprint.
- [24] G. Garrigós, A. Seeger, and T. Ullrich. The Haar system in Triebel-Lizorkin spaces: Endpoint results. *arXiv e-prints*, 40 pages, 2019.
- [25] B. I. Golubov. Best approximations of functions in the L_p metric by Haar and Walsh polynomials. *Math. USSR Sb.*, 16(2):265–285, 1972.
- [26] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Grad. Texts in Math.* Springer, New York, second edition, 2008.

- [27] M. Griebel and S. Knapek. Optimized general sparse grid approximation spaces for operator equations. *Math. Comp.*, 78(268):2223–2257, 2009. Also available as SFB611 preprint No 402.
- [28] K. Guo, G. Kutyniok, and D. Labate. Sparse multidimensional representations using anisotropic dilation and shear operators. In *Wavelets and Splines (Athens, GA, 2005)*, volume 14, pages 189–201. Nashboro Press, Nashville, TN, 2006.
- [29] A. Haar. Zur Theorie der orthogonalen Funktionensysteme. *Math. Ann.*, 71(1):38–53, 1911.
- [30] R. Hochmuth. Wavelet characterizations for anisotropic Besov spaces. *Appl. Comput. Harmon. Anal.*, 12:179–208, 2002.
- [31] H. Kempka. Atomic, molecular and wavelet decomposition of generalized 2-microlocal Besov spaces. *J. Funct. Spaces Appl.*, 8(2):129–165, 2010.
- [32] G. Kutyniok, D. Labate, W.-Q. Lim, and G. Weiss. Sparse multidimensional representation using shearlets. In *Wavelets XI (San Diego, CA)*, volume 5914 of *SPIE Proc.*, pages 254–262. SPIE, Bellingham, WA, 2005.
- [33] G. Kyriazis. Decomposition systems for function spaces. *Studia Math.*, 157(2):133–169, 2003.
- [34] P. Lakhonchai, J. Sampo, and S. Sumetkijakan. Shearlet transforms and directional regularities. *Int. J. Wavelets Multiresolut. Inf. Process.*, 8(5):743–771, 2010.
- [35] Y. Liang, D. Yang, W. Yuan, Y. Sawano, and T. Ullrich. A new framework for generalized Besov-type and Triebel-Lizorkin-type spaces. *Dissertationes Math. (Rozprawy Mat.)*, 489:1–114, 2013.
- [36] S. M. Nikol’skij. *Approximation of functions of several variables and imbedding theorems*, volume 205 of *Grundlehren Math. Wiss.* Springer Berlin, Heidelberg, 1975.
- [37] P. Oswald. The Haar system in Besov function spaces. 2019. preprint.
- [38] V. S. Romanyuk. Constructive characteristic of Hölder classes and m-term approximations in the multiple Haar basis. *Ukrainian Math. J.*, 66(3):391–403, 2014.
- [39] V. S. Romanyuk. Multiple Haar basis and m-term approximations for functions from the Besov classes. I. *Ukrainian Math. J.*, 68(4):625–637, 2016.
- [40] V. S. Romanyuk. Multiple Haar basis and m-term approximations of functions from the Besov classes. II. *Ukrainian Math. J.*, 68(6):928–939, 2016.
- [41] S. G. Roux, M. Clausel, B. Vedel, S. Jaffard, and P. Abry. Self-similar anisotropic texture analysis: the hyperbolic wavelet transform contribution. *IEEE Trans. Image Process.*, 22(11):4353–4363, 2013.
- [42] D. Schertzer and S. Lovejoy. Physically based rain and cloud modeling by anisotropic, multiplicative turbulent cascades. *J. Geophys. Res.*, 92:9693–9714, 1987.

- [43] H.-J. Schmeisser and H. Triebel. *Topics in Fourier analysis and function spaces*. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1987.
- [44] A. Seeger and T. Ullrich. Haar projection numbers and failure of unconditional convergence in Sobolev spaces. *Math. Z.*, 285:91 – 119, 2017.
- [45] A. Seeger and T. Ullrich. Lower bounds for Haar projections: deterministic examples. *Constr. Approx.*, 46(2):227242, 2017.
- [46] W. Sickel and H. Triebel. Hölder inequalities and sharp embeddings in function spaces of $B_{p,q}^s$ and $F_{p,q}^s$ type. *Z. Anal. Anwend.*, 14(1):105–140, 1995.
- [47] B. Stöckert. Ungleichungen vom Plancherel-Pólya-Nikol'skij-Typ in gewichteten L_p^Ω -Räumen mit gemischten Normen. *Math. Nachr.*, 86:19–32, 1978.
- [48] V. N. Temlyakov. *Approximation of periodic functions*. Computational Mathematics and Analysis Series. Nova Science Publishers, Inc., Commack, NY, 1993.
- [49] H. Triebel. *Interpolation theory, function spaces, differential operators*. Amsterdam, North-Holland, 1978.
- [50] H. Triebel. Wavelet bases in anisotropic function spaces. *Function Spaces, Differential Operators and Nonlinear Analysis, FSDONA-04. Milovy, Czech Republic.*, 18:529–550, 2004.
- [51] H. Triebel. *Theory of Function Spaces III*, volume 100 of *Monogr. Math.* Birkhäuser Verlag, Basel, 2006.
- [52] H. Triebel. *Theory of Function Spaces*. Modern Birkhäuser Classics. Springer Basel, 2010.
- [53] M. Ullrich and T. Ullrich. The role of Frolov's cubature formula for functions with bounded mixed derivative. *SIAM J. Numer. Anal.*, 54:969–993, 2016.
- [54] T. Ullrich. Continuous characterizations of Besov-Lizorkin-Triebel spaces and new interpretations as coorbits. *J. Funct. Spaces Appl.*, 2012:Article ID 163213, 47 pages, 2012.
- [55] T. Ullrich and H. Rauhut. Generalized coorbit space theory and inhomogeneous function spaces of Besov-Lizorkin-Triebel type. *J. Funct. Anal.*, 260(11):3299–3362, 2011.
- [56] F. Voigtlaender. Embeddings of decomposition spaces. *arXiv e-prints*, 187 pages, 2016.
- [57] J. Vybírál. Function spaces with dominating mixed smoothness. *Dissertationes Math. (Rozprawy Mat.)*, 436:1–73, 2006.