



**HAL**  
open science

## Copper–Ligand Cooperativity in H<sub>2</sub> Activation Enables the Synthesis of Copper Hydride Complexes

Alicia Aloisi, Étienne Crochet, Emmanuel Nicolas, Jean-Claude Berthet,  
Camille Lescot, Pierre Thuéry, Thibault Cantat

► **To cite this version:**

Alicia Aloisi, Étienne Crochet, Emmanuel Nicolas, Jean-Claude Berthet, Camille Lescot, et al.. Copper–Ligand Cooperativity in H<sub>2</sub> Activation Enables the Synthesis of Copper Hydride Complexes. *Organometallics*, 2021, 40 (13), pp.2064-2069. 10.1021/acs.organomet.1c00212 . hal-03432183

**HAL Id: hal-03432183**

**<https://hal.science/hal-03432183>**

Submitted on 17 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Copper–Ligand Cooperativity in H<sub>2</sub> Activation Enables the Synthesis of Copper Hydride Complexes

Alicia Aloisi, Étienne Crochet, Emmanuel Nicolas, Jean-Claude Berthet, Camille Lescot, Pierre Thuéry, Thibault Cantat,\*

Université Paris-Saclay, CEA, CNRS, NIMBE, 91191 Gif-sur-Yvette, France.  
E-mail: thibault.cantat@cea.fr

## Abstract

Copper(I) complexes of a new participative triphosphane ligand (**2<sup>H</sup>**) have been prepared and structurally characterized, in particular [Cu(**2<sup>H</sup>**)I] and [Cu(**2**)<sub>2</sub>]. Hydrogenation of the latter species afforded the trimetallic hydride species [Cu<sub>3</sub>(**2**)<sub>2</sub>(μ-H)] or in the presence of BEt<sub>3</sub>, [Cu(**2<sup>H</sup>**)(HBEt<sub>3</sub>)]. Their formation evidences transient formation of [Cu(**2<sup>H</sup>**)H] formed by hydrogenolysis of the Cu–N bond of [Cu(**2**)<sub>2</sub>]. [Cu(**2<sup>H</sup>**)(HBEt<sub>3</sub>)] behaves like a hydride complex and inserts CO<sub>2</sub> to yield the formate product [Cu(**2<sup>H</sup>**)(O<sub>2</sub>CH)]. QTAIM (Quantum Theory of Atoms in Molecules) analysis of the Cu–H–BEt<sub>3</sub> interaction indicates a hydride complex stabilized by the Lewis acid BEt<sub>3</sub>.

## Introduction


The development of efficient catalysts for the hydrogenation and cleavage of C–O bonds is an active area of research, in particular for the production of fuels and chemicals from renewable feedstocks such as CO<sub>2</sub> or biomass.<sup>1,2</sup> While complexes of noble metals (Ru, Ir...) favor such hydrogenation reactions, economic and environmental reasons incite the use of catalysts involving base metals (Fe, Co, Mo, Cu...).<sup>3</sup> Among them, solid materials based on copper are well known catalysts for hydrogenation of C=O bonds in carboxylic acids, esters and CO<sub>2</sub> but proceed under harsh conditions and with low selectivity.<sup>2</sup> A handful of molecular copper complexes were reported to activate H<sub>2</sub> and catalyze the hydrogenation of more reactive substrates such as ketones or aldehydes.<sup>4–7</sup> Two of them, [Cu(triphos)(MeCN)]<sup>+</sup> and Cu(OAc)<sub>2</sub>, catalyzed the hydrogenation of CO<sub>2</sub> to formic acid in the presence of an external base, thus evidencing the potential of copper species in catalytic hydrogenation reactions.<sup>8,9</sup>

The generally proposed mechanism for the hydrogenation of C=O bonds involves a transient metal-hydride species generated from the activation of H<sub>2</sub>. In the case of copper, however, while a number of Cu–H intermediates were proposed, for example in hydroelementation reactions as described by Buchwald,<sup>4,8,10,11</sup> only a few have been fully characterized.<sup>12,13</sup> The synthesis of the hexameric complex [(Ph<sub>3</sub>P)CuH]<sub>6</sub> by Osborn in 1971 led to the characterization of a number of polymetallic hydrido clusters or complexes,<sup>14–17</sup> mostly obtained through the reaction of copper halide or alkoxide complexes with borohydrides, boranes or silanes.<sup>18–24</sup> The groups of Caulton, Ruccolo, Appel, Guan and Broere reported efficient procedures that involve the hydrogenolysis of Cu–OR bonds (Scheme 1).<sup>25–31</sup> No terminal copper hydride has ever been isolated in the solid state but Bertrand *et al.* demonstrated that, by using a highly congested carbene ligand, a monomeric copper hydride could exist in solution, in equilibrium with its dimeric form.<sup>13</sup> The same group used a cyclic alkylaminocarbene to stabilize a terminal copper hydride together with a Lewis acid, which was active in CO<sub>2</sub> hydrogenation.<sup>32</sup> More recently,


Sollogoub *et al.* used NHC-capped cyclodextrins to observe a terminal copper hydride species in solution.<sup>33</sup>

**Previous work:**


Caulton *et al.*, 1981, 1985 and 1986




Ruccolo *et al.*, 2013




Appel *et al.*, 2016



**This work:**



R = Tol or NMe<sub>2</sub>




Scheme 1. Copper(I) hydride complexes obtained by hydrogenation methods

In recent years, metal-ligand cooperation in amino phosphine pincer complexes has led to the successful synthesis of many hydride complexes with Ru, Ir, Fe and Mn ions, through heterolytic cleavage of H<sub>2</sub>, and was associated with increased hydrogenation performances.<sup>34-43</sup> Yet, this strategy has never been applied to copper. As tripodal phosphine ligands are well suited to stabilize four-coordinate copper(I) complexes,<sup>8, 44</sup> we have designed a novel triphosphine P<sub>3</sub>NH ligand (**2<sup>H</sup>**) featuring an aminophosphine moiety, as possible participative group (schemes 1 and 2). We report here the preparation of reactive copper(I) complexes of the ligands **2<sup>H</sup>** and **2**, their use in the activation of H<sub>2</sub> and further reactivity with CO<sub>2</sub> and HCO<sub>2</sub>H. These results demonstrate, for the first time, how a participative ligand can facilitate the splitting of H<sub>2</sub> in the coordination sphere of copper.


## Results and Discussion

Syntheses of ligand **2<sup>H</sup>** and its potassium derivative **2<sup>K</sup>** are described in Scheme 2. Lithium halogen exchange between (2-bromophenyl)dicyclohexylphosphine and <sup>n</sup>BuLi at -78 °C in diethylether gave [Cy<sub>2</sub>PC<sub>6</sub>H<sub>5</sub>]<sup>-</sup>Li<sup>+</sup> (non-isolated) which upon addition of 0.5 equivalent of phosphorus trichloride afforded **1** in 93% yield. Addition of benzylamine to a toluene solution of **1** in the presence of an excess of triethylamine led to the formation of the targeted aminophosphine **2<sup>H</sup>** isolated as a white solid in 69 % yield. Compound **2<sup>H</sup>** was characterized by X-ray diffraction (see SI for details) and by multinuclear NMR experiments, exhibiting <sup>31</sup>P NMR shifts at 32.6 (dd, <sup>3</sup>J<sub>PP</sub> = 145, 149 Hz) and -12.6 ppm (d, <sup>3</sup>J<sub>PP</sub> = 147 Hz, average of two

overlapping doublets) for the P–NH and the two Cy<sub>2</sub>P fragments, respectively. While the NH signal is masked by the cyclohexyl protons in benzene, it appears as a multiplet at 2.62 ppm in d<sub>8</sub>-THF. However, the multiplicity of the CH<sub>2</sub> signal of the benzylic N–CH<sub>2</sub>Ph group can be used to probe the protonation of the N center. Indeed, in the presence of an NH functionality, the CH<sub>2</sub> benzylic signal is a pseudo-triplet as observed in **2<sup>H</sup>**, which displays a signal at 4.27 ppm (<sup>3</sup>J<sub>HP</sub> ≈ <sup>3</sup>J<sub>HH</sub> = 6.5 Hz). In compound **2<sup>K</sup>**, obtained from the reaction between **2<sup>H</sup>** and one equivalent of PhCH<sub>2</sub>K, deprotonation of the amine changed the CH<sub>2</sub> signal to a doublet at 4.02 ppm with a unique coupling constant <sup>3</sup>J<sub>HP</sub> = 11.9 Hz.




**Scheme 2.** Syntheses of ligand **2<sup>H</sup>**, **2<sup>K</sup>**, and of the iodide complex **3**



**Scheme 3.** Synthesis of the copper complexes

Reaction of **2<sup>H</sup>** with one equivalent of copper(I) iodide in THF led to the quantitative formation of the yellow complex [(**2<sup>H</sup>**)CuI] (**3**) isolated in 95% yield after washing with pentane and

crystallized as the orange-yellow solvate **3**·C<sub>6</sub>H<sub>6</sub> in refluxing benzene. Complex **3** was characterized by X-ray diffraction and multinuclear NMR (see ESI and Fig. 1). The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum displayed two signals at 26.1 and 5.8 ppm, broadened due to coupling to the copper nucleus.




**Figure 1.** ORTEP views for complexes **3-6**. Displacement ellipsoids are drawn at the 40% probability level. Solvent molecules and carbon bound hydrogen atoms are omitted. The cyclohexyl rings in **4** and **5** are omitted for clarity. For **6**, only one position of the disordered aromatic ring is represented. Selected bond distances (Å) and angles (°): **3**: Cu1–I1 2.6098(3), < Cu–P > 2.27(1); **4**: Cu1–N1 2.0634(16), < Cu–P > 2.31(1); **5**: Cu1–H1 1.66(3), Cu2–H1 1.61(3), Cu3–N1 1.862(3), Cu3–N2 1.858(3), < Cu1–P > 2.31(6), < Cu2–P > 2.27(1); Cu1–H1–Cu2 106.5(14), N1–Cu3–N2 176.60(12); **6**: Cu1–H1 1.66(3), H1–B1 1.35(3), < Cu–P > 2.30(1), Cu1–H1–B1 159(2).

In order to form either a [Cu(2<sup>H</sup>)H] (**6'**) hydride species or the amido derivative [Cu(2)], reaction of **3** with potassium hydride was attempted in THF. Deprotonation of **3** and loss of KI occurred with formation of the dimeric amido species [Cu(2)]<sub>2</sub> (**4**) which was isolated after extraction in diethyl ether as an orange powder (73% yield) while it crystallized as yellow platelets upon slow diffusion of pentane into a THF solution. The dimeric nature of **4** was revealed by single crystal X-ray diffraction. The doublet for the N–CH<sub>2</sub> group at 5.27 ppm (d, <sup>3</sup>J<sub>HP</sub> = 25 Hz) in the <sup>1</sup>H NMR spectrum confirms deprotonation of the amine. The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum evidenced three broad signals at 55.2, –1.9 and –4.3 ppm for the P–N and the Cy<sub>2</sub>P moieties, respectively. The structure of **4** possesses a crystallographic C<sub>2</sub> symmetry axis passing at the center of the (CuPN)<sub>2</sub> ring found in a boat configuration with the two N atoms outside the plane. The ligand is bridging in a μ-κ<sup>2</sup>-P,P-κ<sup>2</sup>-P,N mode and bound by two P atoms (of the P–N and one of the PCy<sub>2</sub> moieties) to a Cu<sup>+</sup> ion and by the second PCy<sub>2</sub> unit and the N atom to another Cu<sup>+</sup> ion so that the copper atom adopts a pseudo-tetrahedral arrangement (Figure 1). The Cu–P and Cu–N can be compared to those found in the other rare amido

triphosphine complexes such as  $[\text{Cu}\{\text{PN}\}(\text{PMe}_3)_2]$  ( $\{\text{PN}\} = p\text{-PhN-C}_6\text{H}_4\text{-P}^i\text{Pr}_2$ ), 2.25(2) and 2.086(1) Å.<sup>45</sup> The P1–N1 bond length of 1.6370(17) is notably shorter than the P–NH bond in **3** (1.6849(19) Å) or in the free ligand **2<sup>H</sup>** (1.691(2) Å) (See ESI).

Complex **4** can be viewed as a relevant precursor to the targeted monometallic copper(I) hydride  $[\text{Cu}(\mathbf{2}^{\text{H}})(\text{H})]$  (**6'**) through hydrogenolysis of the Cu–N bond. Exposure of **4** to 1 atm of  $\text{H}_2$  in benzene or toluene led, after 24 h at 60 °C, to the formation of **5** and free **2<sup>H</sup>** as the major species with 70 % conversion. Complex **5** (Figure 1) was unambiguously identified as the trimetallic monohydride  $[\text{Cu}_3(\mathbf{2})_2(\mu\text{-H})]$  by X-ray diffraction on orange-yellow crystalline platelets of **5** (and its solvate **5'**·2  $\text{C}_6\text{H}_6$ , see ESI) grown in refluxing benzene. The  $^{31}\text{P}\{^1\text{H}\}$  and  $^1\text{H}$  NMR spectra of **5** present a characteristic Cu–H signal at –1.99 ppm and the two magnetically inequivalent protons in the  $\text{PNCH}_2$  moieties appear as two pseudo triplets at 4.62 and 4.37 ppm ( $^2J_{\text{HH}} \approx ^3J_{\text{HP}} = 14.7$  Hz). The negative chemical shift of the hydride resembles that of –1.46 ppm in the 4-coordinate phosphine complex  $[\{\text{Cu}(\kappa^3\text{-triphos})\}_2(\mu^2\text{-H})]^+$  which has a linear cationic  $[\text{Cu}(\mu^2\text{-H})\text{Cu}]^+$  core distinct from that in **5**.<sup>29</sup> The structure of **5** is composed of a triangular  $\text{Cu}_2\text{H}$  core with Cu1, Cu2 and a symmetric  $\mu^2$ -bridging hydride (Figure 1, bottom left). Each of these Cu atoms is coordinated in a  $\kappa^3\text{-P,P,P}$  fashion to a ligand **2**, while the two nitrogen atoms of the amide groups bind the third, two-coordinate Cu3 copper ion in trans positions. The Cu–H distances compare well with those reported for neutral or cationic copper complexes containing bridging  $\mu^2$ -hydride.<sup>12, 13, 19, 22, 27, 28</sup> Treatment of a chilled toluene solution of **3** with 2 equiv.  $\text{KO}^t\text{Bu}$ , followed by addition of  $\text{H}_2$  (1 atm.) similarly led to the quantitative formation of **5** and free **2<sup>H</sup>** after only 1 hour at room temperature. This is the most straightforward way to obtain **5** which was isolated in good yield (69%) after washing with pentane to eliminate free **2<sup>H</sup>**.




**Scheme 4.** Formation of **5** through activation of  $\text{H}_2$  on **4** and putative generation of a copper hydride intermediate

In the hydrogenolysis of dimer **4**, the concomitant formation of a copper hydride entity and free protonated ligand strongly suggests formation of the putative  $[\text{Cu}(\mathbf{2}^{\text{H}})(\text{H})]$  species that would be immediately trapped by the precursor **4** to give **5** with loss of one equivalent of **2<sup>H</sup>** (Scheme 4). To further support the transient formation of  $[\text{Cu}(\mathbf{2}^{\text{H}})(\text{H})]$  in the above reactions, we sought to deliberately trap this species with a Lewis acid, e.g.  $\text{BEt}_3$ , in order to bring kinetic stability as previously reported for the stabilization of a Ca–H bond.<sup>46</sup> Hydrogenation (1 atm.) of a THF solution of **4** in the presence of one equivalent of  $\text{BEt}_3$  cleanly afforded, after 24h at room temperature, the new complex **6**. The conversion never exceeded 20% yield, but no side reaction was observed. Complex **6** was characterized in  $^{31}\text{P}\{^1\text{H}\}$  NMR by two broad signals (at 33.7 and 10.8 ppm) and in the  $^1\text{H}$  NMR spectrum by a signal at –1.39 ppm typical for a Cu–H bond as well as a pseudo-triplet for the benzylic  $\text{CH}_2$  at 4.25 ppm ( $^3J_{\text{HH}} \approx ^3J_{\text{HP}} = 7.3$  Hz). The

$^{11}\text{B}\{^1\text{H}\}$  NMR spectroscopy in benzene revealed a very broad singlet at  $-15.4$  ppm, in the usual region for a  $[\text{HBEt}_3]^-$  ligand.<sup>47</sup> Complex **6** was formed alternatively by the direct reaction of **3** with 1 equivalent of potassium triethylborohydride for 1 hour in toluene. After evaporation of the solvent and dissolution in  $\text{Et}_2\text{O}$ , **6** could be isolated as a yellow powder. Complex **6** is the second example of an isolable triethyl-borohydride copper complex after  $[\text{Cu}(\text{NHC})(\mu\text{-H-BEt}_3)]$  ( $\text{NHC} = \text{N-N}'(2,4,6\text{-C}_6\text{H}_2\text{Me}_3)_2\text{C}_3\text{H}_6\text{N}_2\text{C}$ ).<sup>48</sup> The latter complex features notably a downfield shift for the hydride compared to that in **6**,  $-2.6$  ppm. Such stable species are very unusual because of their extreme reactivity and propensity to lose  $\text{BEt}_3$ . Only a handful of such compounds have been described, mostly with strongly electropositive metals.<sup>49-54</sup> When compared to  $[\text{Cu}(\text{NHC})(\text{HBEt}_3)]$ , which decomposes above  $243$  K with loss of borane, **6** is stable and doesn't degrade after five days at RT.

Yellow crystals of the adduct **6**, obtained in refluxing ether and suitable for X-ray diffraction, definitively evidenced formation of the borate  $[\text{Cu}(\mathbf{2}^{\text{H}})(\mu\text{-H-BEt}_3)]$ . Its structure (Figure 1, bottom right) is very similar to that of **3** and reveals a tetrahedral  $\text{Cu}^+$  center  $\kappa^3$ -coordinated by the tridentate ligand  $\mathbf{2}^{\text{H}}$  and by a monodentate  $[\text{Et}_3\text{BH}]^-$  anion through the H atom. The Cu–H and B–H distances, respectively  $1.66(3)$  Å and  $1.35(3)$  Å, are longer than in the two-coordinate  $[\text{Cu}(\text{NHC})(\mu\text{-HBEt}_3)]$  complex ( $1.56(3)$  and  $1.22(3)$  Å) in line with the different coordination numbers of the complexes (triphosphane vs NHC). The angle Cu–H–B is also more linear ( $159(2)^\circ$  vs  $110.2(18)^\circ$ ).

Copper hydrides are reactive species in catalytic and stoichiometric reduction reactions of unsaturated oxygenated substrates. We thus investigated the behavior of **6** as a stable substitute of the putative hydride  $[\text{Cu}(\mathbf{2}^{\text{H}})(\text{H})]$  in the reduction of  $\text{CO}_2$ .<sup>55,56</sup> Exposure of **6** to 1 atm of  $\text{CO}_2$  led to the formation of the corresponding formate complex **7** with complete conversion after 1 hour at room temperature. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **7** shows two broad signals at  $26.9$  and  $7.3$  ppm while the  $^1\text{H}$  NMR spectrum reveals the presence of a  $\text{HCOO}^-$  ligand at  $8.72$  ppm and the characteristic triplet for the  $\text{CH}_2\text{-(NH)}$  group at  $4.08$  ppm. A peak at  $167.6$  ppm in the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum was also observed and assigned to the formate moiety. Complex **7** was also immediately obtained from the reaction of **4** with two equivalents of formic acid in THF, and isolated in 89 % yield. In the infra-red spectrum, the  $\nu(\text{CO}_2)_{\text{asym}}$  and  $\nu(\text{CO}_2)_{\text{sym}}$  frequencies are found at  $1595$  and  $1377$   $\text{cm}^{-1}$ , respectively.<sup>57,58</sup>



**Scheme 5.** Analysis of the bond enthalpies and energies in **6**. Calculations were performed with Gaussian09, using: PBE0 as a functional; Def2TZVP for Cu, 6-311+G(d,p) for the bridging H, and 6-31G(d) for other atoms as a basis set. Solvent effects in benzene were accounted for with the SMD model.

Table 1. Results of QTAIM Analysis.

	$\rho_{\text{BCP}}$	$\nabla^2\rho_{\text{BCP}}$	$H_{\text{BCP}}$	$G_{\text{BCP}}/\rho_{\text{BCP}}$
Cu–H in <b>6'</b>	0.090	0.126	-0.104	0.809
Cu–H in <b>6</b>	0.081	0.175	-0.116	0.897
B–H in <b>6</b>	0.095	-0.064	-0.030	0.485
B–H in $\text{HBEt}_3^-$	0.116	-0.206	0.006	0.389

BCP: Bond Critical Point. Decreasing  $\rho$  values indicates lower interaction.  $\nabla^2\rho_{\text{BCP}}$  is more positive for greater ionic character.  $G_{\text{BCP}}$  is the kinetic energy density and  $H_{\text{BCP}}$  the local energy density.

In order to gain a deeper understanding of the nature of **6**, and especially of its hydridic character, its electronic structure was studied by QTAIM analysis. The topology of the electron density and its gradient was thus interpreted in terms of chemical bonding, to map the chemical bonds and their critical points (BCP), that can be characterized thanks to the values of the density itself  $\rho_{\text{BCP}}$  and its Laplacian  $\nabla^2\rho_{\text{BCP}}$ .<sup>59</sup> Further analysis of these values through an Abramov gradient decomposition then leads to two values,  $G_{\text{BCP}}$  and  $H_{\text{BCP}}$ , that can be interpreted as the kinetic energy density and the local energy density respectively.<sup>60</sup> The interpretation of both values then allows creating a classification of bonds in covalent, ionic or dative groups. This analysis was performed on **6** and the putative terminal hydride complex  $[(2^{\text{H}})\text{CuH}]$  (**6'**) as well as on the anion  $\text{HBEt}_3^-$ , the results are displayed Table 1. As expected, the Cu–H and B–H bonds in **6** display an ionic character ( $\nabla^2\rho_{\text{BCP}}$  more positive, smaller  $\rho_{\text{BCP}}$ ) greater than in **6'** and free  $\text{HBEt}_3^-$ , which is coherent with the presence of a three-center-two-electrons bond in **6**. Comparing the  $G/\rho$  and  $H$  values at the BCPs suggests that the Cu–H bond in **6** retains the character of a terminal Cu–H bond, as in **6'**. On the other hand, the B–H bond in **6** is significantly disturbed compared to the  $\text{HBEt}_3^-$  borohydride. The change in sign of  $H$  is particularly indicative, as well as the relative differences in the values of  $G/\rho$ . To further investigate the nature of **6**, we calculated the energy required to cleave the Cu–H and the B–H bonds in **6** (Scheme 5). While the loss of  $\text{BEt}_3$  from **6** is slightly endothermic ( $15.1 \text{ kcal.mol}^{-1}$ ), splitting the Cu–H bond to form **6''** requires  $42.5 \text{ kcal.mol}^{-1}$ , pointing to a much weaker B–H bond. This is also reflected in the free energy values for these reactions:  $+1.6$  vs.  $+28.8 \text{ kcal.mol}^{-1}$  for the formation of **6'** vs. **6''**, respectively. This is in line with the energies computed for  $[\text{Cu}(\text{NHC})(\text{HBEt}_3)]$ , (for which dissociation of  $\text{BEt}_3$  has a  $\Delta G$  of  $+10.8 \text{ kcal.mol}^{-1}$ ) and which describe a weaker Cu–H bond compared to **6**.<sup>48</sup> Overall, these data show that complex **6** features an electronic structure closer to a borane-stabilized copper hydride complex that may behave similarly to the terminal hydride complex.



## Conclusion

In conclusion, we have reported the synthesis and characterization of the new P<sub>3</sub>N neutral and anionic ligands **2<sup>H</sup>** and **2<sup>K</sup>** and their copper(I) complexes **3** and **4**. Due to the close proximity to the metal center of the amide group in **4**, metal-ligand cooperativity was exploited for the first time with copper, as a new route to prepare hydride species by H<sub>2</sub> activation. Complex **4** reacted with H<sub>2</sub>, in smooth conditions, to give either the trimetallic monohydride **5** or the borate complex **6**, when BEt<sub>3</sub> is initially present in the reaction mixture. Complex **6**, a rare example of a stable M(μ-H-BE<sub>3</sub>) complex, behaves like a hydride complex with CO<sub>2</sub> and has an electronic structure close to the hydride complex [(**2<sup>H</sup>**)Cu-H⋯(BE<sub>3</sub>)], where the terminal hydride copper complex is stabilized by a Lewis acid.

## Acknowledgements

For financial support of this work, we acknowledge the CEA, CNRS, the University of Paris-Saclay and the European Research Council (ERC Consolidator Grant Agreement no. 818260). We thank GENCI for the allowance of computer time (Project No. A0080806494). T.C. thanks the Fondation Louis D. – Institut de France for its major support.

## References

1. Smith, A. M.; Whyman, R., Review of Methods for the Catalytic Hydrogenation of Carboxamides. *Chem. Rev.* **2014**, *114*, 5477-5510.
2. Pritchard, J.; Filonenko, G. A.; van Putten, R.; Hensen, E. J. M.; Pidko, E. A., Heterogeneous and homogeneous catalysis for the hydrogenation of carboxylic acid derivatives: history, advances and future directions. *Chem. Soc. Rev.* **2015**, *44*, 3808-3833.
3. Alig, L.; Fritz, M.; Schneider, S., First-Row Transition Metal (De)Hydrogenation Catalysis Based On Functional Pincer Ligands. *Chem. Rev.* **2019**, *119*, 2681-2751.
4. Chen, J.-X.; Daeuble, J. F.; Brestensky, D. M.; Stryker, J. M., Highly Chemoselective Catalytic Hydrogenation of Unsaturated Ketones and Aldehydes to Unsaturated Alcohols Using Phosphine-Stabilized Copper(I) Hydride Complexes. *Tetrahedron* **2000**, *56*, 2153-2166.
5. Shimizu, H.; Igarashi, D.; Kuriyama, W.; Yusa, Y.; Sayo, N.; Saito, T., Asymmetric Hydrogenation of Aryl Ketones Mediated by a Copper Catalyst. *Org. Lett.* **2007**, *9*, 1655-1657.
6. Shimizu, H.; Sayo, N.; Saito, T., A Practical and Highly Chemoselective Hydrogenation of Aldehydes with a Copper Catalyst. *Synlett* **2009**, *2009*, 1295-1298.
7. Junge, K.; Wendt, B.; Addis, D.; Zhou, S.; Das, S.; Fleischer, S.; Beller, M., Copper-Catalyzed Enantioselective Hydrogenation of Ketones. *Chem. Eur. J.* **2011**, *17*, 101-105.
8. Zall, C. M.; Linehan, J. C.; Appel, A. M., A Molecular Copper Catalyst for Hydrogenation of CO<sub>2</sub> to Formate. *ACS Catalysis* **2015**, *5*, 5301-5305.
9. Watari, R.; Kayaki, Y.; Hirano, S.-i.; Matsumoto, N.; Ikariya, T., Hydrogenation of Carbon Dioxide to Formate Catalyzed by a Copper/1,8-Diazabicyclo[5.4.0]undec-7-ene System. *Adv. Synth. Catal.* **2015**, *357*, 1369-1373.
10. Shi, S.-L.; Wong, Z. L.; Buchwald, S. L., Copper-catalysed enantioselective stereodivergent synthesis of amino alcohols. *Nature* **2016**, *532*, 353-356.
11. Yang, Y.; Perry, I. B.; Lu, G.; Liu, P.; Buchwald, S. L., Copper-catalyzed asymmetric addition of olefin-derived nucleophiles to ketones. *Science* **2016**, *353*, 144.
12. Jordan, A. J.; Lalic, G.; Sadighi, J. P., Coinage Metal Hydrides: Synthesis, Characterization, and Reactivity. *Chem. Rev.* **2016**, *116*, 8318-8372.
13. Romero, E. A.; Olsen, P. M.; Jazzar, R.; Soleilhavoup, M.; Gembicky, M.; Bertrand, G., Spectroscopic Evidence for a Monomeric Copper(I) Hydride and Crystallographic Characterization of a Monomeric Silver(I) Hydride. *Angew. Chem. Int. Ed.* **2017**, *56*, 4024-4027.
14. Bezman, S. A.; Churchill, M. R.; Osborn, J. A.; Wormald, J., Preparation and crystallographic characterization of a hexameric triphenylphosphinecopper hydride cluster. *J. Am. Chem. Soc.* **1971**, *93*, 2063-2065.
15. Liao, P.-K.; Fang, C.-S.; Edwards, A. J.; Kahlal, S.; Saillard, J.-Y.; Liu, C. W., Hydrido Copper Clusters Supported by Dithiocarbamates: Oxidative Hydride Removal and Neutron Diffraction Analysis of [Cu<sub>7</sub>(H){S<sub>2</sub>C(aza-15-crown-5)}<sub>6</sub>]. *Inorg. Chem.* **2012**, *51*, 6577-6591.
16. Edwards, A. J.; Dhayal, R. S.; Liao, P.-K.; Liao, J.-H.; Chiang, M.-H.; Piltz, R. O.; Kahlal, S.; Saillard, J.-Y.; Liu, C. W., Chinese Puzzle Molecule: A 15 Hydride, 28 Copper Atom Nanoball. *Angew. Chem. Int. Ed.* **2014**, *53*, 7214-7218.
17. Dhayal, R. S.; Liao, J.-H.; Kahlal, S.; Wang, X.; Liu, Y.-C.; Chiang, M.-H.; van Zyl, W. E.; Saillard, J.-Y.; Liu, C. W., [Cu<sub>32</sub>(H)<sub>20</sub>{S<sub>2</sub>P(OiPr)<sub>2</sub>}<sub>12</sub>]: The Largest Number of Hydrides Recorded in a Molecular Nanocluster by Neutron Diffraction. *Chem. Eur. J.* **2015**, *21*, 8369-8374.

18. Albert, C. F.; Healy, P. C.; Kildea, J. D.; Raston, C. L.; Skelton, B. W.; White, A. H., Lewis-base adducts of Group 11 metal(I) compounds. 49. Structural characterization of hexameric and pentameric (triphenylphosphine)copper(I) hydrides. *Inorg. Chem.* **1989**, *28*, 1300-1306.
19. Mankad, N. P.; Laitar, D. S.; Sadighi, J. P., Synthesis, Structure, and Alkyne Reactivity of a Dimeric (Carbene)copper(I) Hydride. *Organometallics* **2004**, *23*, 3369-3371.
20. Lipshutz, B. H.; Frieman, B. A., CuH in a Bottle: A Convenient Reagent for Asymmetric Hydrosilylations. *Angew. Chem. Int. Ed.* **2005**, *44*, 6345-6348.
21. Baker, B. A.; Bošković, Ž. V.; Lipshutz, B. H., (BDP)CuH: A "Hot" Stryker's Reagent for Use in Achiral Conjugate Reductions. *Org. Lett.* **2008**, *10*, 289-292.
22. Frey, G. D.; Donnadiou, B.; Soleilhavoup, M.; Bertrand, G., Synthesis of a Room-Temperature-Stable Dimeric Copper(I) Hydride. *Chem. Asian. J.* **2011**, *6*, 402-405.
23. Wyss, C. M.; Tate, B. K.; Bacsá, J.; Gray, T. G.; Sadighi, J. P., Bonding and Reactivity of a  $\mu$ -Hydrido Dicopper Cation. *Angew. Chem. Int. Ed.* **2013**, *52*, 12920-12923.
24. Nakamae, K.; Kure, B.; Nakajima, T.; Ura, Y.; Tanase, T., Facile Insertion of Carbon Dioxide into Cu<sub>2</sub>( $\mu$ -H) Dinuclear Units Supported by Tetrakisphosphine Ligands. *Chem. Asian. J.* **2014**, *9*, 3106-3110.
25. Goeden, G. V.; Caulton, K. G., Soluble copper hydrides: solution behavior and reactions related to carbon monoxide hydrogenation. *J. Am. Chem. Soc.* **1981**, *103*, 7354-7355.
26. Lemmen, T. H.; Foltling, K.; Huffman, J. C.; Caulton, K. G., Copper polyhydrides. *J. Am. Chem. Soc.* **1985**, *107*, 7774-7775.
27. Goeden, G. V.; Huffman, J. C.; Caulton, K. G., A copper-( $\mu^2$ -hydrogen) bond can be stronger than an intramolecular phosphorus  $\rightarrow$  copper bond. Synthesis and structure of di- $\mu$ -hydrido-bis[ $\eta^2$ -1,1,1-tris(diphenylphosphinomethyl)ethane]dicopper. *Inorg. Chem.* **1986**, *25*, 2484-2485.
28. Eberhart, M. S.; Norton, J. R.; Zuzek, A.; Sattler, W.; Rucolo, S., Electron Transfer from Hexameric Copper Hydrides. *J. Am. Chem. Soc.* **2013**, *135*, 17262-17265.
29. Zall, C. M.; Linehan, J. C.; Appel, A. M., Triphosphine-Ligated Copper Hydrides for CO<sub>2</sub> Hydrogenation: Structure, Reactivity, and Thermodynamic Studies. *J. Am. Chem. Soc.* **2016**, *138*, 9968-9977.
30. Kounalis, E.; Lutz, M.; Broere, D. L. J., Cooperative H<sub>2</sub> Activation on Dicopper(I) Facilitated by Reversible Dearomatization of an "Expanded PNNP Pincer" Ligand. *Chem. Eur. J.* **2019**, *25*, 13280-13284.
31. Ekanayake, D. A.; Chakraborty, A.; Krause, J. A.; Guan, H., Steric Effects of HN(CH<sub>2</sub>CH<sub>2</sub>PR<sub>2</sub>)<sub>2</sub> on the Nuclearity of Copper Hydrides. *Inorg. Chem.* **2020**, *59*, 12817-12828.
32. Romero, E. A.; Zhao, T.; Nakano, R.; Hu, X.; Wu, Y.; Jazzar, R.; Bertrand, G., Tandem copper hydride-Lewis pair catalysed reduction of carbon dioxide into formate with dihydrogen. *Nat. Catal.* **2018**, *1*, 743-747.
33. Xu, G.; Leloux, S.; Zhang, P.; Mejjide Suárez, J.; Zhang, Y.; Derat, E.; Ménand, M.; Bistri-Aslanoff, O.; Roland, S.; Leyssens, T.; Riant, O.; Sollogoub, M., Capturing the Monomeric (L)CuH in NHC-Capped Cyclodextrin: Cavity-Controlled Chemoselective Hydrosilylation of  $\alpha,\beta$ -Unsaturated Ketones. *Angew. Chem. Int. Ed.* **2020**, *59*, 7591-7597.
34. Ohkuma, T.; Ooka, H.; Hashiguchi, S.; Ikariya, T.; Noyori, R., Practical Enantioselective Hydrogenation of Aromatic Ketones. *J. Am. Chem. Soc.* **1995**, *117*, 2675-2676.
35. Käß, M.; Friedrich, A.; Drees, M.; Schneider, S., Ruthenium Complexes with Cooperative PNP Ligands: Bifunctional Catalysts for the Dehydrogenation of Ammonia-Borane. *Angew. Chem. Int. Ed.* **2009**, *48*, 905-907.
36. Filonenko, G. A.; Conley, M. P.; Copéret, C.; Lutz, M.; Hensen, E. J. M.; Pidko, E. A., The impact of Metal-Ligand Cooperation in Hydrogenation of Carbon Dioxide Catalyzed by Ruthenium PNP Pincer. *ACS Catalysis* **2013**, *3*, 2522-2526.
37. Zuo, W.; Lough, A. J.; Li, Y. F.; Morris, R. H., Amine(imine)diphosphine Iron Catalysts for Asymmetric Transfer Hydrogenation of Ketones and Imines. *Science* **2013**, *342*, 1080.
38. Gunanathan, C.; Milstein, D., Bond Activation and Catalysis by Ruthenium Pincer Complexes. *Chem. Rev.* **2014**, *114*, 12024-12087.
39. Zell, T.; Milstein, D., Hydrogenation and Dehydrogenation Iron Pincer Catalysts Capable of Metal-Ligand Cooperation by Aromatization/De aromatization. *Acc. Chem. Res.* **2015**, *48*, 1979-1994.
40. Wang, D.; Astruc, D., The Golden Age of Transfer Hydrogenation. *Chem. Rev.* **2015**, *115*, 6621-6686.
41. Dub, P. A.; Gordon, J. C., The mechanism of enantioselective ketone reduction with Noyori and Noyori-Ikariya bifunctional catalysts. *Dalton Transactions* **2016**, *45*, 6756-6781.
42. Elangovan, S.; Garbe, M.; Jiao, H.; Spannenberg, A.; Junge, K.; Beller, M., Hydrogenation of Esters to Alcohols Catalyzed by Defined Manganese Pincer Complexes. *Angew. Chem. Int. Ed.* **2016**, *55*, 15364-15368.
43. Kumar, A.; Janes, T.; Espinosa-Jalapa, N. A.; Milstein, D., Manganese Catalyzed Hydrogenation of Organic Carbonates to Methanol and Alcohols. *Angew. Chem. Int. Ed.* **2018**, *57*, 12076-12080.
44. Phanopoulos, A.; Long, N. J.; Miller, P. W., Triphosphine Ligands: Coordination Chemistry and Recent Catalytic Applications. In *The Chemical Bond III: 100 years old and getting stronger*, Mingos, D. M. P., Ed. Springer International Publishing: Cham, 2017; pp 31-61.
45. Miller, A. J. M.; Dempsey, J. L.; Peters, J. C., Long-Lived and Efficient Emission from Mononuclear Amidophosphine Complexes of Copper. *Inorg. Chem.* **2007**, *46*, 7244-7246.
46. Kriek, S.; Görls, H.; Westerhausen, M., Synthesis and characterization of rare examples of stable potassium and arylcalcium triethylboranate complexes. *Inorg. Chem. Commun.* **2010**, *13*, 1466-1469.
47. Brown, H. C.; Kramer, G. W.; Hubbard, J. L.; Krishnamurthy, S., Addition compounds of alkali metal hydrides: XVIII. Reaction of trialkylboranes with t-butyllithium. A general, convenient method for the preparation of lithium trialkylborohydrides. *J. Organomet. Chem.* **1980**, *188*, 1-10.
48. Collins, L. R.; Rajabi, N. A.; Macgregor, S. A.; Mahon, M. F.; Whittlesey, M. K., Experimental and Computational Studies of the Copper Borate Complexes [(NHC)Cu(HBEt<sub>3</sub>)] and [(NHC)Cu(HB(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>)]. *Angew. Chem. Int. Ed.* **2016**, *55*, 15539-15543.
49. Cook, K. S.; Piers, W. E.; Rettig, S. J., Reactions of Bis(pentafluorophenyl)borane with Cp<sub>2</sub>Ta(CH<sub>2</sub>)CH<sub>3</sub>. *Organometallics* **1999**, *18*, 1575-1577.
50. Evans, W. J.; Forrestal, K. J.; Ansari, M. A.; Ziller, J. W., Reactions of Olefin Polymerization Activators with Complexed Pentamethylcyclopentadienyl Ligands: Abstraction of Tetramethylfulvalene. *J. Am. Chem. Soc.* **1998**, *120*, 2180-2181.
51. Evans, W. J.; Perotti, J. M.; Ziller, J. W., Trialkylboron/Lanthanide Metallocene Hydride Chemistry: Polydentate Bridging of (HBEt<sub>3</sub>)<sub>3</sub> to Lanthanum. *Inorg. Chem.* **2005**, *44*, 5820-5825.
52. Harvey, M. J.; Hanusa, T. P.; Pink, M., Unusual stability of the coordinated triethylborohydride anion in an alkaline-earth metal complex: crystallographic characterization of [Ca(HBEt<sub>3</sub>)<sub>3</sub>]{1,2,4-C<sub>5</sub>(SiMe<sub>3</sub>)<sub>3</sub>H<sub>2</sub>}(thf)<sub>2</sub>}. *Chem. Commun.* **2000**, 489-490.

53. Janes, T.; Annibale, V. T.; Song, D., Reactivity of Ru(II) and V(III) complexes of diazafluorene derivatives towards B–H bonds. *J. Organomet. Chem.* **2018**, *872*, 79-86.
54. Yan, K.; Upton, B. M.; Ellern, A.; Sadow, A. D., Lewis Acid-Mediated  $\beta$ -Hydride Abstraction Reactions of Divalent  $M(\text{C}(\text{SiHMe}_2)_3)_2(\text{THF})_2$  ( $M = \text{Ca}, \text{Yb}$ ). *J. Am. Chem. Soc.* **2009**, *131*, 15110-15111.
55. Sordakis, K.; Tang, C.; Vogt, L. K.; Junge, H.; Dyson, P. J.; Beller, M.; Laurenczy, G., Homogeneous Catalysis for Sustainable Hydrogen Storage in Formic Acid and Alcohols. *Chem. Rev.* **2018**, *118*, 372-433.
56. Wang, W.-H.; Himeda, Y.; Muckerman, J. T.; Manbeck, G. F.; Fujita, E.,  $\text{CO}_2$  Hydrogenation to Formate and Methanol as an Alternative to Photo- and Electrochemical  $\text{CO}_2$  Reduction. *Chem. Rev.* **2015**, *115*, 12936-12973.
57. Marsich, N.; Camus, A.; Nardin, G., Reaction of carbon dioxide with arylcopper(I) complexes containing tertiary phosphines. *J. Organomet. Chem.* **1982**, *239*, 429-437.
58. Bowmaker, G. A.; Effendy, Hanna, J. V.; Healy, P. C.; Reid, J. C.; Rickard, C. E. F.; White, A. H., Crystal structures and vibrational and solid-state (CPMAS) NMR spectroscopic studies in the tris(triphenylphosphine)-copper(I) and -silver(I) formate systems. *J. Chem. Soc., Dalton Trans.* **2000**, 753-762.
59. Popelier, P. L. A., The QTAIM Perspective of Chemical Bonding. In *The Chemical Bond*, Frenking, G.; Shaik, S., Eds. 2014; pp 271-308.
60. Abramov, Y., On the Possibility of Kinetic Energy Density Evaluation from the Experimental Electron-Density Distribution. *Acta Cryst. A* **1997**, *53*, 264-272.