



HAL
open science

Foraging plasticity of yellow-eyed penguins (*Megadyptes antipodes*) in their subantarctic range

C. G. Muller, B. L. Chilvers, A. Chiaradia, R. K. French, Akiko Kato, Yan Ropert-coudert, Phil F. Battley

► To cite this version:

C. G. Muller, B. L. Chilvers, A. Chiaradia, R. K. French, Akiko Kato, et al.. Foraging plasticity of yellow-eyed penguins (*Megadyptes antipodes*) in their subantarctic range. *Marine Ecology Progress Series*, 2021, 679, pp.149-162. 10.3354/meps13911 . hal-03432359

HAL Id: hal-03432359

<https://hal.science/hal-03432359>

Submitted on 17 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Title page**

2

3 **Title**

4

5 Foraging plasticity of yellow-eyed penguins (*Megadyptes antipodes*) in their subantarctic range.

6

7 **Running page head**

8

9 Variable foraging in yellow-eyed penguins

10

11 **Keywords**

12

13 Foraging behaviour, GPS, VHF, TDR, subantarctic Auckland Islands, pelagic, benthic, climate change,

14 El Nino Southern Oscillation (ENSO), El Niño, La Niña

15

16 **Authors**

17

18 Muller, C. G.; ^{1,2,5}

19 Chilvers, B. L.; ¹

20 Chiaradia, A.; ³

21 French, R. K.; ²

22 Kato, A.; ⁴

23 Ropert-Coudert, Y.; ⁴

24 Battley, P. F.; ²

25

1 ¹ Wildbase, School of Veterinary Sciences, Massey University, Palmerston North 4442, New Zealand

2 ² Wildlife and Ecology Group, School of Agriculture and Environment, Massey University, Palmerston
3 North 4442, New Zealand

4 ³ Conservation Department, Phillip Island Nature Parks, PO Box 97 Cowes, Victoria 3922 Australia

5 ⁴ Centre d'Etudes Biologiques de Chizé, UMR 7372 CNRS - La Rochelle Université, 79360 Villiers-en-
6 Bois, France

7 ⁵ Corresponding author. Email: cmuller@technologist.com

8

9 **Author Statement**

10

11 CGM conceived and designed the experiments, carried out the fieldwork, analysed the data, and wrote
12 the paper. RKF assisted with fieldwork and data collection, and data analysis. AC, AK, and YR-C
13 supplied GPS devices and gave advice on data analysis. BLC and PFB provided general oversight, and
14 assisted with ecological aspects of the project. All authors reviewed the manuscript and gave final
15 approval for publication.

16

17 **ABSTRACT**

18

19 Foraging behaviour is crucial to the breeding success for marine predators, including seabirds. Yellow-
20 eyed penguins (*Megadyptes antipodes*) are central-place predominantly benthic foragers around
21 mainland New Zealand. The mainland population of this iconic species is declining, with changes in
22 the marine environment a suspected cause, in particular, warming water and poorer foraging success.
23 Here, we undertook a detailed foraging study of the ancestral subantarctic population which is
24 genetically separate from the northern population. Over two breeding seasons, we collected GPS logs

1 from 94 deployments on 75 breeding yellow-eyed penguins foraging from Enderby Island, Auckland
2 Islands, New Zealand. Birds foraged up to 47 km from shore, more than twice as far as yellow-eyed
3 penguins in many areas around mainland New Zealand. Foraging area size and maximum range
4 (distance from shore) were larger in a year of greater breeding effort (2016), and also for females, and
5 birds undertaking pelagic foraging trips. Differences in foraging behaviour are likely influenced by
6 local bathymetry, environmental conditions, and individual preference. Despite comparable bathymetry
7 in some areas, the southern population shows greater foraging plasticity than seen in the northern
8 population, implying foraging conditions may be less restricted in the subantarctic. However, previous
9 studies have shown breeding success at the Auckland Islands is inconsistent, and differing foraging
10 behaviour between years indicates foraging conditions are likely variable. Prey availability and
11 foraging success are also expected to be affected by warming water, with implications for future
12 breeding success particularly from the effects of climate change and El Niño Southern Oscillations
13 (ENSO).

14 **1. INTRODUCTION**

15
16 Many marine predators are central-place foragers, such as seabirds and pinnipeds that breed on land but
17 forage at sea, returning ashore to provide for their young (Boyd et al. 1994, Boersma & Rebstock 2009,
18 Jones et al. 2020). Finding food is essential for breeding success and population viability for long-lived
19 marine predators (Villegas-Amtmann et al. 2008, Chilvers & Wilkinson 2009), including seabirds
20 (Sandvik et al. 2005, Catry et al. 2013). Seabird foraging success can vary between years, as oceanic
21 conditions, reflected in indices such as the El Niño Southern Oscillation (ENSO) can change and may
22 affect prey abundance and distribution (Boyd et al. 1994, Miller & Sydeman 2004, Grémillet &
23 Boulinier 2009, Ropert-Coudert et al. 2009, Catry et al. 2013, Agnew et al. 2015, Ropert-Coudert et al.
24 2015, Poupart et al. 2017). Benthic prey are less influenced by oceanographic perturbations such as

1 ENSO and they may be a more reliable food source than pelagic prey over time (Costa et al. 2004) .
2 However, other changes to the benthos can affect benthic penguin foraging (Browne et al. 2011). One
3 approach to classify benthic and pelagic dives is by determining dive shape and depth. Benthic dives to
4 the seafloor are characterised by a “U-” or “square-shaped” dive profile with a uniform maximum
5 depth limited by bathymetry (Wilson 1995, Tremblay & Cherel 2000, Pütz & Cherel 2005, Bost et al.
6 2007). Conversely, pelagic dives are in mid-water and display a “V-” or “W-shaped” dive profile with
7 a more variable maximum depth between dives, with the former sometimes considered exploratory
8 behaviour, and the latter linked with prey pursuit activity (Wilson 1995, Ropert-Coudert et al. 2000,
9 Pütz & Cherel 2005). These dives tend to have a shorter bottom time given that prey could be
10 encountered anywhere in the water column (Wilson 1995, Tremblay & Cherel 2000, Pütz & Cherel
11 2005, Bost et al. 2007). When foraging, most penguin species rely on pelagic feeding dives, catching
12 prey within the water column (Ratcliffe & Trathan 2011), although some species such as southern
13 rockhopper penguins (*Eudyptes chrysocome filholi*) have a mixed strategy incorporating both pelagic
14 and benthic diving (Tremblay & Cherel 2000).

15
16 Yellow-eyed penguins (hōiho, *Megadyptes antipodes*) are endangered (Couch-Lewis et al. 2016, IUCN
17 2020) and endemic to New Zealand. Their distribution is restricted to the south-east of the South
18 Island, Stewart and Codfish Islands (the northern population), and subantarctic Auckland and Campbell
19 Islands (the southern population, Figure 1) (Seddon et al. 2013). The northern population is undergoing
20 a severe decline due to successive poor breeding seasons and high adult mortality, thought to be
21 primarily a result of threats at sea including poor foraging success, fisheries interactions, pollution and
22 human disturbance (Couch-Lewis et al. 2016, Mattern et al. 2017, Mattern & Wilson 2018). Over 60%
23 of the total yellow-eyed penguin population breeds in the subantarctic (Couch-Lewis et al. 2016,
24 Muller et al. 2020b), which is considered the stronghold for the species. The subantarctic population

1 appears stable at present, but with evidence of wide fluctuations and a possible decline since the 1980s
2 (Moore 1992, Muller et al. 2020b). Analysis of ancient DNA has shown that the original endemic
3 mainland species *M. waitaha* went extinct soon after Polynesian settlement of New Zealand c.1280 CE
4 (Boessenkool et al. 2009a, Collins et al. 2014). Yellow-eyed penguins from the subantarctic expanded
5 their range into this vacant niche and colonised the mainland prior to the increase of European settlers
6 in the late 1800s (Boessenkool et al. 2009a). However, there is currently almost no migration (<2%)
7 between the mainland and subantarctic, meaning these areas represent separate populations and
8 management units (Boessenkool et al. 2009b). Despite the importance of the southern population to the
9 species, there is little information available on yellow-eyed penguin foraging behaviour in the
10 subantarctic, and whether foraging differs from the northern population.

11
12 During breeding, yellow-eyed penguins are central-place foragers, and the northern population feeds in
13 shallow coastal waters adjacent to their breeding area, and over mid-shelf areas further offshore where
14 they have access to a large shelf area (Moore 1999, Mattern et al. 2007, Mattern et al. 2013). Around
15 mainland New Zealand, foraging trips are typically up to 25 km from shore and over mid-shelf areas,
16 as confirmed by VHF and GPS tracking studies (Moore 1999, Mattern et al. 2013), although the mean
17 foraging range can be as short as 6.2 km offshore at some locations (Mattern et al. 2007). While diet
18 studies in the 1990s showed occasional indications of pelagic foraging (van Heezik 1990, Moore et al.
19 1995), more recent dive data had demonstrated a predominantly benthic foraging strategy for the
20 mainland population (Mattern et al. 2007, Mattern et al. 2013, Chilvers et al. 2014). In contrast, birds in
21 the southern population utilise a mixed strategy incorporating varying amounts of pelagic foraging at
22 the subantarctic Auckland Islands, including solely pelagic foraging trips (Muller et al. 2020a).
23 Moreover, subantarctic yellow-eyed penguins forage at greater depths than in many northern areas,

1 with a maximum depth of 134 m for benthic dive bouts, and 115 m for pelagic dives (Muller et al.
2 2020a).

3
4 Yellow-eyed penguin foraging is negatively influenced by warmer water in the northern population
5 (Young 2014, Mattern & Ellenberg 2018). La Niña conditions result in warmer water and more
6 stochastic weather and wind patterns in the New Zealand region, with an adverse effect on yellow-eyed
7 penguin breeding success (Young 2014). El Niño conditions result in more variable breeding success
8 than in neutral ENSO conditions (Peacock et al. 2000, Darby 2003). There is no information about the
9 effects of ENSO and climate variability on foraging and breeding success in subantarctic yellow-eyed
10 penguins, and whether this may vary from the northern population.

11
12 The southern population displays widely variable breeding success which is likely linked to foraging
13 success (Moore 1992, Muller et al. 2020b). Foraging parameters are a product of the physical
14 environment, changing environmental parameters, or individuals' preference to target specific prey in
15 particular habitats. The subantarctic populations forage in deeper water and over greater distances than
16 many mainland birds (Muller et al. 2020a) and may be expected to expend more energy foraging.
17 Therefore, in light of this variability, and the importance of the subantarctic population to the species, a
18 study of foraging parameters was urgently needed in the southern population.

19
20 This study determined the size and location of foraging areas used by yellow-eyed penguins breeding at
21 Enderby Island in the New Zealand subantarctic (Figure 1). We compared foraging habitats to
22 published data for the northern population. Given the importance of the southern population to the
23 species, the knowledge of foraging area use by subantarctic yellow-eyed penguins is vital to
24 complement research on diving behaviour and breeding success in the subantarctic. This information

1 could assist with future conservation management of the species and marine-based threats in these
2 isolated subantarctic areas.

3 **2. MATERIALS & METHODS**

4 **2.2. Fieldwork and Equipment**

5
6 Fieldwork was carried out on Enderby Island, Auckland Islands, in the New Zealand subantarctic
7 (50°29'45"S 166°17'44"E, Figure 1) for three breeding seasons; 2015 (Nov 2015–Feb 2016), 2016
8 (Nov 2016–Feb 2017), and 2017 (Nov 2017–Jan 2018). GPS data were collected in 2016 and 2017,
9 while dive and breeding success data were collected in all three years in parallel studies (Muller et al.
10 2020a, Muller et al. 2020b). Nests were located using manual ground searches, ground-based VHF
11 telemetry, and an Unmanned Aerial Vehicle (UAV) equipped with a Very High Frequency (VHF)
12 radio receiver (Muller et al. 2019). Adult yellow-eyed penguins were captured by hand as they returned
13 from sea and placed in a capture bag for processing and collection of morphometric data using a spring
14 balance and callipers. Birds were marked with a microchip (Allflex, Palmerston North, New Zealand)
15 for permanent identification (Muller et al. 2020b) and sex was determined using the relationship
16 between head plus beak length and foot length (Setiawan et al. 2004), or the relative sizes between
17 breeding partners with males assumed to be the larger individual (Setiawan et al. 2004).

18
19 Data loggers were deployed during late November and December, corresponding to the guard phase of
20 breeding. GPS loggers were attached using waterproof tape (TESA, Beiersdorf, Germany) (Muller et
21 al. 2020a) to the lower back to optimise streamlining and orientation to the sky during the typical
22 posture adopted during swimming or brooding. Other data loggers were taped to the upper back on the
23 midline to maximise streamlining (Figure 2). GPS loggers were customised CatTraQ™ GPS loggers,
24 14 × 35 × 70 mm, weight ~30 g (Catnip Technologies, USA) modified for underwater use with the

1 addition of a magnetic on/off switch, and a moulded resin housing (Pelletier et al. 2014). GPS loggers
2 were programmed to record a fix every 3 minutes, providing a battery life of approximately 4–5 days.
3 After programming, each unit was waterproofed with heat-shrink tubing (TE Connectivity,
4 Schaffhausen, Switzerland) before deployment. Time Depth Recorders (TDR) and VHF transmitters
5 were also attached following a similar protocol (Muller et al. 2020a). Where possible, loggers were
6 deployed for one foraging trip before being recovered to minimise potential disturbance. All
7 deployments in 2016 consisted of a GPS and TDR logger (Table 1). Of the total deployments in 2017, a
8 subset of 31 birds received two logger deployments of either a GPS + TDR, or a TDR only, followed
9 by the alternative in a subsequent trip, with the deployment order randomised (Muller et al. 2020a).
10 The remaining unpaired trips in 2017 included 10 GPS + TDR deployments, and two TDR only
11 deployments.

12 **2.3. Data Analysis**

13 GPS data files were downloaded and filtered by deleting any duplicate records (those with a distance of
14 0 m between subsequent positions), then interpolated to account for missed GPS fixes (at 3 min
15 intervals) when the unit was underwater during a scheduled fix attempt. Interpolation assumed a
16 constant heading and velocity between recorded locations and used an automated script in Python 3.5.2
17 (Python Software Foundation, Beaverton USA, available at www.python.org). Where multiple foraging
18 trips were recorded in one deployment, the data were considered as separate trips (Muller et al. 2020a).
19 Data were displayed in ArcGIS 10.2.2 (ESRI), with NIWA New Zealand region bathymetry data at 50
20 m depth contours overlaid for comparison (Mitchell et al. 2016). Points on land were deleted using a
21 spatial selection tool in ArcGIS. Data were projected in the NZTM coordinate system, and geodesic
22 distance calculations between points were automated in a Python script using the GeoPy library.

23

1 The foraging range (maximum straight-line distance away from the shore, measured from the sea
2 access point) and the total trip distance (cumulative distance travelled between all points in a foraging
3 trip) were calculated from interpolated data. Summary data were calculated from these distances (mean
4 + SD). When calculating mean distances, short trips (<10 km foraging range, or <25 km total distance)
5 and with low GPS fix success (<20%) were excluded to avoid biasing distance measurements, since
6 these may not have represented the furthest distance travelled. However, all GPS fixes were included
7 for area calculations and other analyses.

8
9 Foraging range areas were analysed using GME 0.7.3.0 (www.spatial ecology.com) and ArcGIS
10 functions (Beyer 2012, Locher & Lindenberg 2016). Kernel Density Estimates (KDE) were determined
11 with smoothed cross-validation bandwidth which provided the best visually-estimated match with data
12 point concentrations, and a cell size of 50 m. Values were calculated using 95% and 50% kernel
13 contours to represent the home range and core foraging areas, respectively (Hamer et al. 2007), and
14 isopleth and polygon features were generated for spatial analysis in ArcGIS. Foraging area calculations
15 were determined using the 95% confidence interval. The intersection between kernel density polygons
16 was used to compare the percentage overlap of foraging activity location between groups of interest
17 representing different years, sexes, and dive types (i.e. benthic or pelagic diving). Dive data were
18 categorised separately using Bayesian analysis (see Muller et al. (2020a)). Individual dives were
19 classified as benthic if they displayed an inter-dive depth change of less than 2.9% from the previous
20 AND following dive. Foraging trips were also classified, with benthic trips having more than 3.6%
21 benthic dives, which ensured that remaining trips classified as pelagic contained entirely pelagic diving
22 bouts (Muller et al. 2020a). Dive analysis included all dives greater than 2 m depth which likely
23 included some travelling dives to and from the foraging area. Polygon areas were calculated in ArcGIS,
24 along with percentage of spatial overlap between different foraging areas. Only GPS data were used to

1 determine foraging areas to avoid skewing habitat use, i.e. excluding interpolated positions which were
2 more likely to occur during transits to and from foraging locations when penguins swim quickly and
3 surface only briefly (Mattern et al. 2007).

4
5 Statistical analyses were performed in R Studio version 1.1.456 running R version 3.5.1 (R. Core Team
6 2017), and using the lme4 package (Bates et al. 2015). Linear mixed effects models were used to
7 compare maximum foraging range and total trip distance with year and forage type (as fixed variables),
8 and with bird ID as a random effect (since some birds made more than one trip). ANOVA tests were
9 used to determine the significance of these effects in each model. Graphs were generated in R,
10 including the ggplot2 package (Wickham 2010).

11 **3. RESULTS**

12 **3.1. Foraging Area**

13 A total of 87 GPS foraging tracks were collected (52 in 2016, and 35 in 2017), from 69 individual birds
14 (Table S1). These data included 48 trips made by 38 females and 38 trips by 30 males, plus one trip by
15 one bird of unknown sex. Yellow-eyed penguins foraged on an offshore continental shelf plateau
16 approximately 30 – 40 km south-east of Enderby Island (Figure 3), where the water depth is
17 predominantly 50 – 100 m, with some spill-over into deeper water up to 150 m deep. Of all tracks, 32
18 corresponded to benthic and 52 corresponded to pelagic foraging trips. Dive type could not be
19 determined for three trips where there were no corresponding dive records. A small subset of eleven
20 birds (seven in 2016 and four in 2017) travelled to the northwest to forage off the northern coast of
21 Auckland Island (Figure 3), with nine of these (82%) conducting pelagic foraging trips.
22 Birds travelled over a more extensive range in 2016 compared to 2017 (Figure 3A, upper and middle
23 panels), with an estimated total foraging area size of 801 vs 462 km², respectively (Table 2). Analysis

1 of the intersection (overlap) between years (Figure 3A, bottom panel) showed only 37% of birds
2 foraged in the overlap area in 2016, compared to 64% of birds in 2017 (Table 2). Benthic foraging trips
3 covered a smaller area than pelagic trips (Figure 3B, upper and middle panels), with estimated total
4 foraging areas of 571 vs 985 km², respectively (Table 2). Across all years the benthic foraging area
5 than was smaller than for pelagic foraging. The shared home range area had a 91% overlap with the
6 foraging area used by individuals undertaking benthic foraging (Figure 3B, bottom panel), compared to
7 only 53% overlap with the area used by pelagic foragers (Table 2). Females foraged over a much larger
8 range than males (Figure 3C, upper and middle panels), with foraging areas of 963 vs 585 km²,
9 respectively (Table 2) and 85% of males foraged in this overlap area (Figure 3C, bottom panel),
10 compared to only 52% of females (Table 2).

11 **3.1. Foraging Distances**

12 The largest foraging range was 46.7 km offshore, with a mean of 27.8 ± 7.5 km. The longest total trip
13 distance was 151.9 km, with a mean value of 69.2 ± 26.7 km (Table 3). Linear mixed-effects models
14 showed that the foraging range was significantly greater in 2016 than 2017 ($\chi^2=19.78$, $p<0.0001$) and
15 was also significantly greater for benthic foragers than for pelagic ($\chi^2=8.53$, $p=0.014$). Although
16 pelagic foragers used a greater geographical area including travelling furthest from shore (Figure 3B)
17 their shorter mean foraging range can be explained by the wide variation (Figure S2A), with the mean
18 value of the core foraging range closer to shore than for benthic foragers (50% contours, Figure 3B).
19 The total trip distance was also significantly greater in 2016 compared to 2017 ($\chi^2=13.82$ $p<0.001$) but
20 there was no significant difference between benthic or pelagic foraging trip types ($\chi^2=4.86$, $p=0.088$).
21 Sex was not significant for foraging range ($p=0.36$), or total trip distance ($p=0.187$).

22 **3.2. Foraging Changes**

23 In 2016, individuals foraged over a larger area and had a smaller overlap of shared areas indicating
24 birds were foraging more widely than in 2017, when the foraging home range area reduced in size by

1 340 km² or 42% (Figure 3A). Birds also foraged further offshore in 2016 (Table 3), and on average
2 foraging range was greater for benthic foragers than pelagic foragers in both years (Figure S2A).
3 However, pelagic foragers had a much wider spread in their foraging range including travelling the
4 greatest distance offshore (Table S1, Figure S2), and had a much larger foraging home range at the
5 95% CI (Figure 3B). The total trip distance showed a similar trend (Figure S2B) although this was not
6 significantly different between benthic and pelagic foraging.

7 **4. DISCUSSION**

8 Foraging by breeding yellow-eyed penguins was concentrated over a plateau to the east of Enderby
9 Island where the maximum water depth is up to 150 m (Mitchell et al. 2016), and the substrate is a
10 mixture of coarse sand, broken shells, coral, and pebbles (Tidey & Hulbe 2019, LINZ 2020). The
11 foraging range of Enderby birds averaged up to 30.5 km offshore in 2016, with a maximum of 46.7 km
12 (Table 3), but was less in other years. This is larger than ranges reported for the northern population
13 which typically forage a maximum of 25 km from shore around mainland New Zealand (Moore 1999,
14 Mattern et al. 2007, Mattern et al. 2013). The total distance travelled per trip by Enderby Island birds
15 was also greater than at many northern population locations, where penguins typically swam around 31
16 ± 10 km per trip with extremes of 55 ± 12 km recorded (Mattern et al. 2013). The northern population
17 are considered to be predominantly benthic foragers with only benthic dives published in studies using
18 dive loggers (Mattern et al. 2007, Mattern et al. 2013, Chilvers et al. 2014), although there is possible
19 evidence for occasional pelagic foraging from diet and other studies (van Heezik 1990, Moore et al.
20 1995, Mattern et al. 2018). Conversely, birds in the subantarctic Auckland Islands show a much greater
21 degree of diving plasticity. They demonstrate a mixed diving strategy incorporating varying amounts of
22 pelagic foraging between and within seasons, including solely pelagic foraging trips (Muller et al.
23 2020a). Changes in diving behaviour between years also corresponded with changes in foraging

1 behaviour, including home range size and distance travelled. Foraging trip duration in the subantarctic
2 also changes between years, with trips in 2017 significantly shorter than trips in
3 2015 and 2016, but there is no difference between the sexes (Muller et al. 2020a).

4 **4.1. Changes in foraging behaviour**

5 Individuals foraged over a larger area with a smaller overlap in 2016 than in 2017. Mean foraging
6 range was greater for benthic foragers than pelagic foragers in both years. This difference suggests that
7 while benthic foragers may have travelled further offshore on average to reach their foraging area,
8 pelagic foragers generally covered similar distances while searching for prey although they utilised a
9 much larger and more variable area. Benthic diving, especially in deep water can use more energy than
10 other types of diving (Costa et al. 2004, Chilvers & Wilkinson 2009), so this may represent a greater
11 energy expenditure by birds conducting benthic foraging. In all years benthic foraging took place over
12 a smaller area than pelagic foraging. Given that both benthic and pelagic diving occurred together in
13 some locations (the overlap areas) this indicates that dive type was not governed by bathymetry or
14 water depth in these shared areas.

15
16 Dive logger data from the same study published elsewhere showed the proportion of pelagic foraging
17 trips increased each year from none in 2015, to 79% in 2017 (Muller et al. 2020a). Pelagic foraging
18 was associated with a larger foraging area size in our data (Figure 3B) which might be expected to
19 result in increasing foraging area size over the same period. However, while the highest proportion of
20 pelagic diving was in 2017, the foraging area was actually smaller in 2017 than in 2016 (Figure 3A),
21 although sample size was also smaller in 2017. Foraging area use is therefore likely affected by
22 additional complexity in the type of prey species and their spatial distribution each year. During 2017 a
23 total of 28% of birds changed their diving behaviour on a subsequent trip in the same year, and 56%
24 changed their behaviour between different years (Muller et al. 2020a). This indicates diving plasticity is

1 present in the subantarctic, and given the link between diving behaviour and foraging this implies
2 plasticity in foraging area use also. There is no difference in diving behaviour between male and female
3 birds in the subantarctic (Muller et al. 2020a), although females may have been foraging over a wider
4 area during the guard phase (Figure 3C), with only 51.6% overlap with the shared foraging area,
5 compared to 85.1% overlap for males.

6
7 In comparison, mean foraging ranges for mainland New Zealand yellow-eyed penguins ranged from
8 6.2 ± 0.8 (SD) km to 23.3 ± 11.2 (SD) km (Table 3), although these included data collected using
9 different methods, and during different breeding phases and years where birds may have foraged more
10 widely (Moore 1999, Mattern et al. 2007, Mattern et al. 2013). Mean and maximum mainland foraging
11 ranges were closer to shore than in the subantarctic, however statistical comparison between these data
12 sets was not possible. The northern population displays a generally consistent benthic foraging strategy
13 utilising the same foraging areas consistently over different years (Mattern et al. 2007). However, while
14 mainland birds tend to be either inshore or offshore foragers, some could switch strategies (Moore
15 1999), and foraging areas (Moore et al. 1995, Moore 1999), indicating some foraging plasticity in the
16 northern population as well. Differences are likely due to the local environment and prey availability
17 (Muller et al. 2020a), rather than any inherent behavioural differences between these genetically similar
18 populations (Boessenkool et al. 2009a).

19
20 This study reports on the foraging area used by breeding penguins during the guard phase when
21 parental attendance at the nest is high (Richdale 1957, Darby et al. 1990). However, many penguin
22 species forage over considerably larger areas during incubation and post-guard phases of the breeding
23 season compared to the guard phase (Jouventin et al. 1994), including an area over five times larger for
24 little penguins (Sánchez et al. 2018). In little penguins, foraging closer to the colony during chick-

1 rearing was also associated with a diet switch to higher trophic level prey (Poupart et al. 2017), and the
2 mean maximum foraging range in winter was significantly larger (up to eight times greater) than during
3 the breeding season (Hoskins et al. 2008, McCutcheon et al. 2011). Yellow-eyed penguins in the
4 northern population travel further from the breeding area during incubation and post-guard stages
5 (Moore 1999) and in winter (M. Young pers. comm.). Therefore, given that no foraging data are
6 available for other breeding phases in the subantarctic, the foraging areas described here should be
7 regarded as a minimum estimate of the total area used by yellow-eyed penguins from Enderby Island,
8 which may be considerably larger. Our data also tended to show that larger datasets with more
9 positions represented larger foraging areas, so more data may reveal additional use areas. However,
10 these sample sizes were considerably larger than for comparable studies on the northern population,
11 and increasing the sample size is likely impractical due to logistics and ethical concerns for this
12 endangered species.

13
14 It is possible that carrying the larger GPS loggers may have influenced behaviour. However, Muller et
15 al (2020a) showed that the type of loggers deployed (TDR only, or TDR + GPS) did not have a
16 biologically significant effect on diving behaviour or deployment order (the difference was 0.55 m,
17 which was less than the error margin of the loggers).

19 **4.3. Foraging and breeding success**

20 In the Auckland Islands, there was a much greater number of breeding birds and a higher overall
21 breeding success (total number of fledged chicks) in 2016 than in 2015 or 2017 (Muller et al. 2020b).
22 This also coincided with an increase in pelagic foraging from 2015 to 2017 (Muller et al. 2020a). A
23 foraging strategy's success will depend on the trip duration, distance travelled and the return for effort,
24 which will be affected by the distribution and quality of prey, and the effort required for capture (Costa

1 2015). Therefore, the switch from predominantly benthic foraging behaviour in 2015 indicates that
2 during 2016 and 2017 pelagic foraging likely provided greater returns that outweighed benthic
3 foraging. A favourable pelagic food source may have provided nutritional benefits outweighing the
4 energy requirement of foraging over a larger area to obtain it. However, while the breeding population
5 and breeding success both declined in 2017 (Muller et al. 2020b), the proportion of pelagic foraging
6 trips continued to increase compared to previous years (Muller et al. 2020a). This difference suggests a
7 more complex interaction between foraging and breeding success, suggesting that conditions for all
8 types of foraging were less favourable in 2017.

9
10 The 2015 breeding season corresponded with severe El Niño conditions (Null 2019), which have been
11 linked to a local decline of prey species and poorer foraging and breeding outcomes for yellow-eyed
12 penguins than during neutral ENSO conditions (Peacock et al. 2000, Darby 2003). The 2016 and 2017
13 seasons both had weak La Niña conditions (Null 2019) which have a stronger negative effect on
14 northern yellow-eyed penguin breeding success than El Niño conditions (Young 2014). Benthic
15 foraging generally took place over a smaller area (Figure 3B), and the proportion of benthic foraging
16 was greatest during the El Niño season in 2015 (Muller et al. 2020a). Therefore, the 95% CI of the
17 foraging home range would also be expected closer to shore during El Niño conditions. While we do
18 not have GPS data from 2015 to confirm this, trip times in 2015 were significantly shorter than in 2016
19 (Muller et al. 2020a), implying a more compact foraging area, and possibly closer to shore. It is not
20 known whether prey assemblages are consistent in this area during different ENSO conditions, so 2015
21 may not have been representative of all El Niño seasons. Yellow-eyed penguin prey species in the
22 subantarctic are unknown, as is their distribution in space and time, so this would require further
23 research to confirm.

24

1 Travelling greater distances can result in longer times at sea, although this is not always the case. For
2 subantarctic yellow-eyed penguins, pelagic foraging trips are not significantly different in time duration
3 from benthic trips (Muller et al. 2020a), or in total trip length (Figure S2B), although individual trip
4 distances could vary. Trips in 2017 were of significantly shorter time than in 2016 (Muller et al.
5 2020a), which is consistent with the smaller foraging area utilised (Figure 3A). Longer foraging trips in
6 seabirds including Magellanic (*Spheniscus magellanicus*), Adélie (*Pygoscelis adeliae*), and little
7 penguins (*Eudyptula minor*) are directly related to lower breeding success (Chiaradia & Nisbet 2006,
8 Boersma & Rebstock 2009). Changes in prey availability, particularly the distance travelled to obtain
9 it, will affect the effort required (Miller & Sydeman 2004). This, in turn, affects both adult energetics
10 and chick provisioning; longer travel or search times by foraging parents can result in less-frequent
11 feeding of chicks, and may result in lower growth rates and fledging weights for chicks (Kitaysky et al.
12 2000, Davoren & Montevecchi 2003, Pinaud et al. 2005). Longer foraging trips may also result in more
13 food digestion, with less available for transfer to offspring (Weimerskirch et al. 1994, Ropert-Coudert
14 et al. 2004). Increased energy expenditure by foraging adults may lead to a reduction in their body
15 condition (Arnould et al. 1996, Shaffer et al. 2003), as well as breeding success (Inchausti et al. 2003),
16 thereby influencing long-term survival and evolutionary fitness of breeders. Therefore, any factors
17 affecting the type and distribution of prey which may require travelling greater distances or spending
18 more time at sea, could have a negative effect on future yellow-eyed penguin breeding success.

19 **4.4. Foraging and conservation management**

20 Enderby Island holds over 50% of the breeding population for the Auckland Islands archipelago, and
21 will likely continue to be the main breeding location in the future unless predators are removed from
22 Auckland Island (Muller et al. 2020b). Only part of the foraging area used by Enderby Island birds is
23 protected from fisheries interactions by the Auckland Islands Motu Maha Marine Reserve which
24 extends 12 nm from shore (Figure 3) Although there was no mortality of yellow-eyed penguins

1 reported as bycatch in the trawl fishery around the Auckland Islands reserve during the period of this
2 study (Ministry of Primary Industries 2018), fisheries activities such as bottom-trawling modify the
3 benthos and may affect penguin foraging (Browne et al. 2011), and indirect competition with fisheries
4 has been linked to declines in some mainland yellow-eyed penguin populations (Ellenberg & Mattern
5 2012). Therefore, research into the direct and indirect impacts of fishing activities on yellow-eyed
6 penguins in the Auckland Islands area would be crucial for their long term conservation. The core
7 foraging areas (represented by the 50% isopleths) were contained within the marine reserve boundary,
8 and the home range of all penguins we examined (represented by the 95% isopleths) contained 595 km²
9 (81%) of foraging area within the marine reserve. However, the areas presented here represent
10 minimum estimates of the habitat used, and therefore should be regarded as the minimum areas where
11 protection should be considered. Additionally, birds breeding at Carnley Harbour in the south may have
12 a smaller foraging area available, as water depths drop to 150 m within only 11 km of the harbour
13 entrance.

14

15 **CONCLUSIONS**

16 This study found foraging trips of longer distance and the use of a larger foraging area by the southern
17 population. These differences are likely a result of local conditions, rather than any inherent differences
18 between these two genetically similar populations. Yellow-eyed penguins have many factors currently
19 impacting their survival as a species including land- and sea-based mortality and have been classified
20 as endangered since 2000, with the population continuing to decline (IUCN 2020). These impacts
21 should receive ongoing monitoring and mitigation to ensure they do not combine with poor foraging
22 seasons to exacerbate population declines in both locations. For the southern population, removing
23 introduced mammalian predators from Auckland Island should be an essential step for future
24 management of subantarctic yellow-eyed penguins. The foraging area size and distance varied at

1 Enderby Island between years, likely due to changing climate effects bringing changes in prey
2 availability, with considerable variability in breeding success between years. Poor breeding outcomes
3 could increase in frequency and become more pronounced in the future due to increasing water
4 temperature associated with ENSO oscillations or climate change. Therefore, this variable breeding
5 success remains a cause for concern for subantarctic populations, as successive poor breeding seasons
6 in the future could lead to similar declines to those observed in the northern population.

7 **ACKNOWLEDGEMENTS**

8 Fieldwork was carried out under DOC Wildlife Act and Massey University Animal Ethics Permits
9 (MUAEC 14/67), and with the approval of Ngai Tahu iwi. Thanks to Mel Young and Thomas Mattern
10 for information on mainland diving, and to Danielle Sijbranda for assistance in the field. Funding was
11 provided by a Massey University PhD Scholarship, Massey University Research Funding, and Institute
12 of Veterinary Animal and Biomedical Sciences / School of Veterinary Science Post-graduate studies
13 grant, Birds New Zealand Research Fund, and Freemasons Charity. Thanks also to the following for
14 supplying equipment, logistics, and other assistance: Jo Hiscock, DOC Southland; Wildlife and
15 Ecology Group Massey University; Yellow-eyed Penguin Trust; Blue Planet Marine; Phillip Island
16 Nature Parks; Sirtrack; Ponant Expeditions. Thanks to Steve Kafka and crew of *Evohe* for safe passage
17 to the subantarctic and back.

18

19

LITERATURE CITED

- 1
- 2
- 3 Agnew P, Lalas C, Wright J, Dawson S (2015) Variation in breeding success and survival of little
4 penguins *Eudyptula minor* in response to environmental variation. *Marine Ecology Progress*
5 *Series* 541:219-229
- 6 Arnould JPY, Boyd IL, Speakman JR (1996) The relationship between foraging behaviour and energy
7 expenditure in Antarctic fur seals. *Journal of Zoology* 239(4):769-782
- 8 Bates D, Maechler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4.
9 *Journal of Statistical Software* 67:1-48
- 10 Beyer HL (2012) Geospatial Modelling Environment. www.spatialecology.com/gme.
- 11 Boersma PD, Rebstock GA (2009) Foraging distance affects reproductive success in Magellanic
12 penguins. *Marine Ecology Progress Series* 375:263-275
- 13 Boessenkool S, Austin JJ, Worthy TH, Scofield RP, Cooper A, Seddon PJ, Waters JM (2009a) Relict
14 or colonizer? Extinction and range expansion of penguins in southern New Zealand.
15 *Proceedings of the Royal Society B: Biological Sciences* 276(9):815-821
- 16 Boessenkool S, Star B, Waters JM, Seddon PJ (2009b) Multilocus assignment analyses reveal
17 multiple units and rare migration events in the recently expanded yellow-eyed penguin
18 (*Megadyptes antipodes*). *Molecular Ecology* 18(11):2390-2400
- 19 Bost CA, Handrich Y, Butler PJ, Fahlman A, Halsey LG, Woakes AJ, Ropert-Coudert Y (2007)
20 Changes in dive profiles as an indicator of feeding success in king and Adélie penguins. *Deep*
21 *Sea Research Part II: Topical Studies in Oceanography* 54(3):248-255
- 22 Boyd IL, Arnould JPY, Barton T, Croxall JP (1994) Foraging behaviour of Antarctic fur seals during
23 periods of contrasting prey abundance. *Journal of Animal Ecology* 63(3):703-713
- 24 Browne T, Lalas C, Mattern T, Van Heezik Y (2011) Chick starvation in yellow-eyed penguins:
25 Evidence for poor diet quality and selective provisioning of chicks from conventional diet
26 analysis and stable isotopes. *Austral Ecology* 36(1):99-108
- 27 Catry T, Ramos JA, Catry I, Monticelli D, Granadeiro JP (2013) Inter-annual variability in the breeding
28 performance of six tropical seabird species: influence of life-history traits and relationship with
29 oceanographic parameters. *Marine Biology* 160(5):1189-1201
- 30 Chiaradia A, Nisbet IC (2006) Plasticity in parental provisioning and chick growth in little penguins
31 *Eudyptula minor* in years of high and low breeding success. *Ardea* 94(2):257
- 32 Chilvers BL, Dobbins ML, Edmonds HK (2014) Diving behaviour of yellow-eyed penguins, Port
33 Pegasus/Pikihati, Stewart Island/Rakiura, New Zealand. *New Zealand Journal of Zoology*
34 41(3):161-170
- 35 Chilvers BL, Wilkinson IS (2009) Diverse foraging strategies in lactating New Zealand sea lions.
36 *Marine Ecology Progress Series* 378:299-308
- 37 Collins CJ, Rawlence NJ, Prost S, Anderson CNK, Knapp M, Scofield RP, Robertson BC, Smith I,
38 Matisoo-Smith EA, Chilvers BL, Waters JM (2014) Extinction and recolonization of coastal
39 megafauna following human arrival in New Zealand. *Proceedings of the Royal Society B:*
40 *Biological Sciences* 281:20140097
- 41 Costa DP (2015) Reproductive and foraging energetics of high latitude penguins, albatrosses and
42 pinnipeds: Implications for life history patterns. *American Zoologist* 31(1):111-130
- 43 Costa DP, Kuhn CE, Weise MJ, Shaffer SA, Arnould JPY (2004) When does physiology limit the
44 foraging behaviour of freely diving mammals? *International Congress Series* 1275:359-366
- 45 Couch-Lewis Y, McKinlay B, Murray S, Edge Hill K-A. (2016) Yellow-eyed penguin stock-take report -
46 He pūrongo mō te hoiho - A report of progress against the hoiho recovery plan (Department of
47 Conservation, 2000) objectives and actions. *Terrestrial Ecosystems Unit, Dunedin, New*
48 *Zealand*.
- 49 Darby JT (2003) The yellow-eyed penguin (*Megadyptes antipodes*) on Stewart and Codfish Islands.
50 *Notornis* 50(3):148-154

- 1 Darby JT, Seddon PJ, Davis LS (1990) Breeding biology of yellow-eyed penguins (*Megadyptes*
2 *antipodes*). In: Davis LS, Darby JT (eds) Penguin Biology, Book Penguin Biology. Academic
3 Press, London, UK
- 4 Davoren GK, Montevecchi WA (2003) Consequences of foraging trip duration on provisioning
5 behaviour and fledging condition of common murre *Uria aalga*. *Journal of Avian Biology*
6 34(1):44-53
- 7 Ellenberg U, Mattern T (2012) Yellow-eyed penguin - review of population information. Department of
8 Conservation, Wellington, New Zealand.
- 9 Grémillet D, Boulinier T (2009) Spatial ecology and conservation of seabirds facing global climate
10 change: a review. *Marine Ecology Progress Series* 391:121-137
- 11 Hamer K, Humphreys E, Garthe S, Hennicke J, Peters G, Grémillet D, Phillips R, Harris M, Wanless S
12 (2007) Annual variation in diets, feeding locations and foraging behaviour of gannets in the
13 North Sea: flexibility, consistency and constraint. *Marine Ecology Progress Series* 338:295-305
- 14 Hoskins AJ, Dann P, Ropert-Coudert Y, Kato A, Chiaradia A, Costa DP, Arnould JPY (2008) Foraging
15 behaviour and habitat selection of the little penguin *Eudyptula minor* during early chick rearing
16 in Bass Strait, Australia. *Marine Ecology Progress Series* 366:293-303
- 17 Inchausti P, Guinet C, Koudil M, Durbec J-P, Barbraud C, Weimerskirch H, Cherel Y, Jouventin P
18 (2003) Inter-annual variability in the breeding performance of seabirds in relation to
19 oceanographic anomalies that affect the Crozet and the Kerguelen sectors of the Southern
20 Ocean. *Journal of Avian Biology* 34(2):170-176
- 21 IUCN (2020) *Megadyptes antipodes* (Yellow-eyed Penguin). IUCN Red List of Threatened Species.
22 Accessed 24 Feb 2021. <http://www.iucnredlist.org>
- 23 Jones KA, Ratcliffe N, Votier SC, Newton J, Forcada J, Dickens J, Stowasser G, Staniland IJ (2020)
24 Intra-specific niche partitioning in Antarctic fur seals, *Arctocephalus gazella*. *Scientific reports*
25 10(1):3238
- 26 Jouventin P, Capdeville D, Cuenot-Chaillet F, Boiteau C (1994) Exploitation of pelagic resources by a
27 non-flying seabird: satellite tracking of the king penguin throughout the breeding cycle. *Marine*
28 *Ecology-Progress Series* 106:11-11
- 29 Kitaysky AS, Hunt GL, Jr., Flint EN, Rubega MA, Decker MB (2000) Resource allocation in breeding
30 seabirds: responses to fluctuations in their food supply. *Marine Ecology Progress Series*
31 206:283-296
- 32 LINZ (2020) Chart NZ 286 Auckland Islands. Land Information New Zealand (LINZ), Wellington, New
33 Zealand.
- 34 Locher A, Lindenbergh M (2016) Home range creation and analysis using Geospatial Modeling
35 Environment and ArcGIS software. Grand Valley State University, Biology Department,
36 Allendale, Michigan.
- 37 Mattern T, Ellenberg U (2018) Yellow-eyed penguin diet and indirect effects affecting prey composition
38 - Collation of biological information. Department of Conservation, Wellington, New Zealand.
39 DOI 10.13140/RG.2.2.23828.81284
- 40 Mattern T, Ellenberg U, Houston DM, Davis LS (2007) Consistent foraging routes and benthic foraging
41 behaviour in yellow-eyed penguins. *Marine Ecology Progress Series* 343:295-306
- 42 Mattern T, Ellenberg U, Houston DM, Lamare M, Davis LS, Van Heezik Y, Seddon PJ (2013) Straight
43 line foraging in yellow-eyed penguins: New insights into cascading fisheries effects and
44 orientation capabilities of marine predators. *PLoS ONE* 8(12):e84381
- 45 Mattern T, McPherson MD, Ellenberg U, van Heezik Y, Seddon PJ (2018) High definition video
46 loggers provide new insights into behaviour, physiology, and the oceanic habitat of a marine
47 predator, the yellow-eyed penguin. *PeerJ* 6:e5459
- 48 Mattern T, Meyer S, Ellenberg U, Houston DM, Darby JT, Young M, van Heezik Y, Seddon PJ (2017)
49 Quantifying climate change impacts emphasises the importance of managing regional threats
50 in the endangered Yellow-eyed penguin. *PeerJ* 5:e3272
- 51 Mattern T, Wilson KJ (2018) New Zealand penguins - current knowledge and research priorities. A
52 report compiled for Birds New Zealand, Dunedin, New Zealand.

- 1 McCutcheon C, Dann P, Salton M, Renwick L, Hoskins AJ, Gormley AM, Arnould JPY (2011) The
2 foraging range of Little Penguins (*Eudyptula minor*) during winter. *Emu* 111(4):321-329
- 3 Miller AK, Sydeman WJ (2004) Rockfish response to low-frequency ocean climate change as revealed
4 by the diet of a marine bird over multiple time scales. *Marine Ecology Progress Series*
5 281:207-216
- 6 Ministry of Primary Industries (2018) Non-fish bycatch events within fishing area SQU6T dataset.
7 Wellington, New Zealand
- 8 Mitchell JS, Mackay KA, Neil HL, Mackay EJ, Pallentin A, Notman P (2016) New Zealand region
9 bathymetry, 1:4 000 000, 2nd edition. . In: NIWA (ed) NIWA Chart, Miscellaneous Series No
10 92. NIWA, Wellington, NZ
- 11 Moore PJ (1992) Yellow-eyed penguin population estimates on Campbell and Auckland Islands 1987-
12 90. *Notornis* 39(1):1-15
- 13 Moore PJ (1999) Foraging range of the yellow-eyed penguin *Megadyptes antipodes*. *Marine*
14 *Ornithology* 27(1-2):56-58
- 15 Moore PJ, Wakelin M, Douglas ME, McKinlay B, Nelson D, Murphy B (1995) Yellow-eyed penguin
16 foraging study, south-eastern New Zealand, 1991-1993. Department of Conservation,
17 Wellington, New Zealand.
- 18 Muller CG, Chilvers BL, Barker Z, Barnsdale KP, Battley PF, French RK, McCullough J, Samandari F
19 (2019) Aerial VHF tracking of wildlife using an unmanned aerial vehicle (UAV): comparing
20 efficiency of yellow-eyed penguin (*Megadyptes antipodes*) nest location methods. *Wildlife*
21 *Research* 46(2):145-153
- 22 Muller CG, Chilvers BL, French RK, Battley PF (2020a) Diving plasticity in the ancestral range of the
23 yellow-eyed penguin, *Megadyptes antipodes*, an endangered marine predator. *Marine Ecology*
24 *Progress Series* 648:191-205
- 25 Muller CG, Chilvers BL, French RK, Hiscock JA, Battley PF (2020b) Population estimate for yellow-
26 eyed penguins (*Megadyptes antipodes*) in the subantarctic Auckland Islands, New Zealand.
27 *Notornis* 67:299–319
- 28 Null J (2019) El Niño and La Niña years and intensities. Accessed 28 September.
29 <https://ggweather.com/enso/oni.htm>
- 30 Peacock L, Paulin M, Darby J (2000) Investigations into climate influence on population dynamics of
31 yellow-eyed penguins *Megadyptes antipodes*. *New Zealand Journal of Zoology* 27(4):317-325
- 32 Pelletier L, Chiaradia A, Kato A, Ropert-Coudert Y (2014) Fine-scale spatial age segregation in the
33 limited foraging area of an inshore seabird species, the little penguin. *Oecologia* 176(2):399-
34 408
- 35 Pinaud D, Chérel Y, Weimerskirch H (2005) Effect of environmental variability on habitat selection,
36 diet, provisioning behaviour and chick growth in yellow-nosed albatrosses. *Marine Ecology*
37 *Progress Series* 298:295-304
- 38 Poupart TA, Waugh SM, Bost C, Bost C-A, Dennis T, Lane R, Rogers K, Sugishita J, Taylor GA,
39 Wilson K-J, Zhang J, Arnould JPY (2017) Variability in the foraging range of *Eudyptula minor*
40 across breeding sites in central New Zealand. *New Zealand Journal of Zoology* 44(3):225-244
- 41 Pütz K, Chérel Y (2005) The diving behaviour of brooding king penguins (*Aptenodytes patagonicus*)
42 from the Falkland Islands: variation in dive profiles and synchronous underwater swimming
43 provide new insights into their foraging strategies. *Marine Biology* 147(2):281-290
- 44 R. Core Team (2017) R: a language and environment for statistical computing. [https://www.R-](https://www.R-project.org/)
45 [project.org/](https://www.R-project.org/)
- 46 Ratcliffe N, Trathan P (2011) A review of the diet and foraging movements of penguins breeding within
47 the CCAMLR area. *CCAMLR Science* 18:75-114
- 48 Richdale LE (1957) A population study of penguins. Clarendon Press, Oxford
- 49 Ropert-Coudert Y, Kato A, Chiaradia A (2009) Impact of small-scale environmental perturbations on
50 local marine food resources: a case study of a predator, the little penguin. *Proceedings*
51 *Biological sciences / The Royal Society* 276(1676):4105-4109

- 1 Ropert-Coudert Y, Kato A, Meyer X, Pellé M, Macintosh AJJ, Angelier F, Chastel O, Widmann M,
2 Arthur B, Raymond B, Raclot T (2015) A complete breeding failure in an Adélie penguin colony
3 correlates with unusual and extreme environmental events. *Ecography* 38(2):111-113
- 4 Ropert-Coudert Y, Sato K, Kato A, Charrassin J-B, Bost C-A, Maho YI, Naito Y (2000) Preliminary
5 investigations of prey pursuit and capture by king penguins at sea.
- 6 Ropert-Coudert Y, Wilson RP, Daunt F, Kato A (2004) Patterns of energy acquisition by a central
7 place forager: benefits of alternating short and long foraging trips. *Behavioral Ecology*
8 15(5):824-830
- 9 Sánchez S, Reina RD, Kato A, Ropert-Coudert Y, Cavallo C, Hays GC, Chiaradia A (2018) Within-
10 colony spatial segregation leads to foraging behaviour variation in a seabird. *Marine Ecology*
11 *Progress Series* 606:215-230
- 12 Sandvik H, Erikstad KE, Barrett RT, Yoccoz NG (2005) The effect of climate on adult survival in five
13 species of North Atlantic seabirds. *Journal of Animal Ecology* 74(5):817-831
- 14 Seddon P, Ellenberg U, Van Heezik Y (2013) The Yellow-eyed Penguin. In: Borboroglu GP, Boersma
15 D (eds) *Penguins: Natural History and Conservation, Book 1*. University of Washington Press,
16 Seattle and London
- 17 Setiawan AN, Darby JT, Lambert DM (2004) The use of morphometric measurements to sex yellow-
18 eyed penguins. *Waterbirds* 27(1):96-101
- 19 Shaffer SA, Costa DP, Weimerskirch H (2003) Foraging effort in relation to the constraints of
20 reproduction in free-ranging albatrosses. *Functional Ecology* 17(1):66-74
- 21 Tidey EJ, Hulbe CL (2019) Sub-Antarctic Auckland Islands seafloor mapping investigations using
22 legacy data. *Geosciences* 9(2):56
- 23 Tremblay Y, Cherel Y (2000) Benthic and pelagic dives: a new foraging behaviour in Rockhopper
24 Penguins. *Marine Ecology Progress Series* 204:257-267
- 25 van Heezik Y (1990) Diets of yellow-eyed, Fiordland crested, and little blue penguins breeding
26 sympatrically on Codfish Island, New Zealand. *New Zealand Journal of Zoology* 17(4):543-548
- 27 Villegas-Amtmann S, Costa DP, Tremblay Y, Salazar S, Auriolles-Gamboa D (2008) Multiple foraging
28 strategies in a marine apex predator, the Galapagos sea lion *Zalophus wollebaeki*. *Marine*
29 *Ecology Progress Series* 363:299-309
- 30 Weimerskirch H, Chastel O, Ackermann L, Chaurand T, Cuenot-Chaillet F, Hindermeier X, Judas J
31 (1994) Alternate long and short foraging trips in pelagic seabird parents. *Animal Behaviour*
32 47(2):472-476
- 33 Wickham H (2010) ggplot2: elegant graphics for data analysis. *J Stat Softw* 35(1):65-88
- 34 Wilson R (1995) Foraging ecology. In: *The penguins Spheniscidae*. Oxford University Press, Oxford
- 35 Young MJ (2014) Determining the drivers of yellow-eyed penguin productivity. MSc thesis, University
36 of Otago, New Zealand
- 37

38

1 **FIGURES & TABLES**

2

3 Table 1: Data logger deployments on breeding yellow-eyed penguins from Enderby Island, showing the
4 number of trips with loggers of each type in each year of the study. In 2015 only TDRs were deployed,
5 and in 2016 all birds received both types of logger. In 2017 a subset of 31 birds carried data loggers on
6 two separate occasions (one with GPS + TDR, and one with TDR only) with the order randomised.
7 Additional single deployments of GPS + TDR loggers were also made on eight other individuals.
8 Where multiple foraging trips were recorded in one deployment, the data were divided into separate
9 trips.

10

Year	TDR only	GPS + TDR	TOTAL
2015	13	0	13
2016	0	51	51
2017	31	39	70
TOTAL	44	90	134

11

12

13 Table 2: Combined foraging area size of breeding yellow-eyed penguins from Enderby Island,
14 comparing different parameters (year, dive type, and sex). Areas were calculated from GPS data with
15 the 95% confidence interval of kernel density estimates representing the combined home range area
16 used by all birds, and the 50% confidence interval representing the combined core foraging area. For
17 each comparison, Intersect shows the size of the spatial overlap indicating the shared area common to
18 both parameters, and overlap shows the percentage overlap of the shared intersect area with each
19 parameter. Spatial representations of the areas for all parameters are shown on separate maps (Figure
20 3).

21

22

Comparison	Parameter	Home range (95% CI)		Core use (50% CI)	
		Area (km ²)	Overlap	Area (km ²)	Overlap
Year	2016	801	37%	196	29%
Year	2017	462	65%	91	63%
Year	Intersect	299	-	57	-
Dive Type	Benthic	571	91%	134	64%
Dive Type	Pelagic	985	52%	203	42%
Dive Type	Intersect	517	-	86	-
Sex	Females	963	52%	191	48%
Sex	Males	585	85%	132	69%
Sex	Intersect	497	-	91	-

1

2



3 Table 3: Foraging range (maximum distance from shore) and total trip distance (cumulative distance
4 travelled) of breeding yellow-eyed penguins from Enderby Island in different years. Short trips (<10
5 km foraging range, or <25 km total distance) and with low GPS fix success (<20%) were ignored for
6 distance calculations to avoid bias. Also shown are the number of individual foraging trips, and the
7 number of birds which returned with valid GPS logs each season. Grey highlights show data from
8 mainland sites for comparison.

9

Reference	Year	Location	Site	Year	Phase	Foraging range (km)			Total distance (km)			No. Birds	No. Trips
						Mean	SD	Max	Mean	SD	Max		
This study	2016	Subantarctic	Enderby Is	2016	guard	30.5	5.7	46.7	74.6	25.6	151.9	52	42
	2017	Subantarctic	Enderby Is	2017	guard	19.2	5.9	30.0	50.1	22.1	98.9	29	13
	2016-17	Subantarctic	Enderby Is	2016-17	guard	27.8	7.5	46.7	69.2	26.7	151.9	81	55
Mattern	2013	Mainland	Boulder Beach, Otago	2004	guard	21.1	5.9		54.5	12.0		8	
		Mainland	Boulder Beach, Otago	2005	guard	11.0	3.1		30.5	10.3		4	
		Mainland	Boulder Beach, Otago	2012	guard, post-guard	10.8	6.2		33.6	18.9		11	
Mattern	2007	Mainland	Bushy Beach, Oamaru	2003	guard, post-guard	6.2	0.8		15.9	1.2		5	
		Mainland	Bushy Beach, Oamaru	2003	guard, post-guard	17.5	2.5		47.5	1.8		5	
		Mainland	Bushy Beach, Oamaru	2004	guard	18.2	1.1		46.0	3.0		4	
Moore	1999	Mainland	Boulder Beach, Otago	1990	post-guard	14.4	7.2					6	
		Mainland	Boulder Beach, Otago	1991	incubation	23.3	11.2					13	
		Mainland	Boulder Beach, Otago	1991	guard	13.4	6.1					10	
		Mainland	Boulder Beach, Otago	1991	post-guard	15.5	8.7					10	
		Mainland	Boulder Beach, Otago	1992	incubation	14.0	8.6					14	
		Mainland	Boulder Beach, Otago	1992	guard	14.4	5.8					10	
		Mainland	Boulder Beach, Otago	1992	post-guard	12.4	6.1					10	
		Mainland	Long Point, Catlins	1991	post-guard	11.1	7.2					9	

1

Mainland	Long Point, Catlins	1992	post-guard	9.4	5.3	10
----------	------------------------	------	------------	-----	-----	----



1

2 **Figure 1:** (A) The breeding range of yellow-eyed penguins (green highlights) around the New Zealand mainland (above the dashed line) and
3 in the subantarctic (below the dashed line). (B) The Auckland Islands archipelago with Enderby Island to the NE. Selected depth contours
4 are shown in blue, from Mitchell et al. (2016). (C) A close-up of Enderby Island showing the area where breeding birds were sampled (green
5 ellipse). Modified from Muller et al (2020a), Fig. 1.

6

7

8


9



1

2

- 1 **Figure 2:** Nesting yellow-eyed penguin with a Time-Depth Recorder and VHF transmitter attached to the upper back, and GPS logger
- 2 attached to the lower back. Enderby Island, Auckland Islands, New Zealand subantarctic. Photo credit: CG Muller



1 **Figure 3:** Kernel density estimates of yellow-eyed penguin foraging trips to compare different
2 parameters showing comparison by A) year, B) dive type, and C) sex. In all cases 95% contours (outer
3 polygon) indicate combined home range use, and 50% contours (inner polygon) indicate combined core
4 foraging area use. The spatial intersection of kernel density estimates (lower panels) shows the overlap
5 area common to both parameters (upper and middle panels). 95% contours (orange) indicate shared
6 home range use and 50% contours (red) indicate shared core foraging use. Also shown on all maps are
7 50 m depth contours (light grey) with selected depth values per Figure 1, and the extent of the marine
8 reserve 12 nm from shore (black line).

9

1 **Supplementary materials**

2

3 Table S1: Foraging trips by breeding yellow-eyed penguins from Enderby Island. Data were derived
4 from simultaneous GPS and TDR deployments, and are ordered chronologically. Where multiple
5 foraging trips were recorded in one deployment the data were separated into individual trips. Trip times
6 are shown from GPS logs. Foraging type was determined from dive logs, where available (Muller et al.
7 2020a). GPS fix success was calculated from the theoretical number of possible fixes between the first
8 and last GPS position times per trip.

9


BirdID	Sex	Year	Deploy				GPS Fix Success	Foraging Range (km)	Total Distance (km)	Forage Type	
			No.	Start Date	Start Time	End Date					End Time
bird186	F	2016	1	5/12/2016	05:27:04	5/12/201	20:46:59	6%	6.41	9.89	Benthic
bird187	F	2016	1	7/12/2016	05:19:04	8/12/201	11:55:09	4%	1.95	1.77	Benthic
bird001	M	2016	1	8/12/2016	20:10:04	10/12/20	08:03:59	19%	31.18	52.21	Benthic
bird052	F	2016	1	9/12/2016	15:03:04	10/12/20	14:06:59	11%	32.76	63.96	Benthic
bird053	F	2016	1	9/12/2016	17:48:04	10/12/20	19:22:59	13%	25.72	31.18	Benthic
bird086	M	2016	1	10/12/201	05:32:04	10/12/20	15:01:59	8%	6.47	11.57	Pelagic
bird030	M	2016	1	10/12/201	05:58:04	12/12/20	16:07:59	21%	30.67	72.75	Benthic
bird019	M	2016	1	10/12/201	07:05:04	11/12/20	20:41:59	6%	24.89	56.24	Benthic
bird143	F	2016	1	10/12/201	07:25:04	11/12/20	20:29:59	12%	34.09	95.59	Benthic
bird117	M	2016	1	10/12/201	15:09:04	11/12/20	16:44:59	8%	26.94	59.63	Benthic
bird142	F	2016	1	12/12/201	14:05:04	15/12/20	19:16:59	11%	33.36	81.02	Pelagic
bird042	F	2016	1	12/12/201	16:45:04	14/12/20	14:46:59	3%	2.58	4.67	Pelagic
bird106	M	2016	1	12/12/201	17:59:04	13/12/20	17:38:59	11%	25.01	56.79	Benthic
bird107	M	2016	1	13/12/201	14:47:04	15/12/20	10:55:59	6%	25.60	53.40	Benthic
bird175	F	2016	1	13/12/201	16:35:04	15/12/20	22:21:59	10%	29.18	72.86	Benthic
bird132	M	2016	1	14/12/201	05:46:04	16/12/20	00:30:59	4%	3.56	5.45	Pelagic
bird189	M	2016	1	14/12/201	15:01:04	14/12/20	19:49:59	7%	1.31	0.89	Pelagic
bird022	F	2016	1	14/12/201	18:15:04	18/12/20	14:25:59	5%	35.72	143.37	Pelagic
bird145	F	2016	1	15/12/201	05:19:04	17/12/20	16:03:59	10%	27.84	85.48	Benthic
bird111	F	2016	1	15/12/201	05:27:04	16/12/20	18:52:59	9%	37.18	87.35	Pelagic
bird101	F	2016	1	16/12/201	18:19:04	17/12/20	17:30:59	11%	23.08	56.19	Pelagic
bird192	M	2016	1	17/12/201	16:41:04	18/12/20	19:34:59	14%	22.03	70.89	Benthic
bird172	M	2016	1	17/12/201	18:17:04	19/12/20	11:06:59	5%	8.66	20.98	Pelagic

bird171	M	2016	1	17/12/201	22:48:04	19/12/20	10:08:59	4%	7.18	12.44	Pelagic
bird191	M	2016	1	17/12/201	23:58:04	19/12/20	20:41:59	12%	36.41	83.15	Benthic
bird170	F	2016	1	18/12/201	06:07:04	18/12/20	22:00:59	3%	7.12	14.84	Pelagic
bird164	F	2016	1	24/12/201	13:25:04	26/12/20	08:00:59	27%	35.71	70.43	Pelagic
bird155	M	2016	1	24/12/201	16:01:04	25/12/20	16:23:59	30%	33.92	67.14	Benthic
bird154	M	2016	1	24/12/201	17:03:04	25/12/20	15:03:59	33%	32.30	64.08	Benthic
bird161	F	2016	1	24/12/201	17:36:04	25/12/20	17:47:59	40%	35.71	76.51	Pelagic
bird159	M	2016	1	24/12/201	20:00:04	25/12/20	20:44:59	29%	36.57	73.41	Benthic
bird205	F	2016	1	25/12/201	16:47:04	27/12/20	08:48:59	30%	33.23	67.19	Pelagic
bird207	F	2016	1	25/12/201	16:59:04	26/12/20	17:26:59	45%	25.62	62.15	Benthic
bird210	F	2016	1	25/12/201	17:00:04	28/12/20	07:30:44	16%	32.10	122.78	Benthic
bird206	M	2016	1	25/12/201	17:40:04	26/12/20	21:07:59	43%	27.60	66.17	Benthic
bird120	F	2016	1	25/12/201	17:40:04	28/12/20	03:57:59	34%	33.64	96.97	Benthic
bird160	M	2016	1	25/12/201	18:46:04	27/12/20	03:36:59	30%	29.32	66.42	Benthic
bird208	F	2016	1	25/12/201	20:40:04	26/12/20	20:51:59	29%	24.53	53.08	Benthic
bird149	F	2016	1	25/12/201	22:04:04	27/12/20	05:41:59	29%	33.07	69.76	Pelagic
bird162	F	2016	1	25/12/201	22:52:04	27/12/20	22:21:59	28%	34.51	133.11	Pelagic
bird032	F	2016	1	26/12/201	05:21:04	28/12/20	12:16:59	30%	46.73	110.35	Pelagic
bird183	F	2016	1	26/12/201	18:31:04	27/12/20	19:18:59	25%	17.21	36.61	Pelagic
bird064	?	2016	1	26/12/201	19:37:04	28/12/20	23:20:59	11%	38.78	151.88	Benthic
bird209	M	2016	1	27/12/201	05:25:04	28/12/20	03:54:59	23%	31.50	62.94	Benthic
bird215	F	2016	1	30/12/201	03:47:04	1/01/201	09:42:59	7%	28.14	57.90	Benthic
bird213	F	2016	1	30/12/201	05:17:04	31/12/20	18:13:59	45%	36.97	79.56	Pelagic
bird217	M	2016	1	30/12/201	15:39:04	1/01/201	13:22:59	22%	30.32	70.55	Benthic
bird214	F	2016	1	30/12/201	16:00:04	31/12/20	18:21:59	16%	8.97	18.43	Pelagic
bird216	M	2016	1	31/12/201	18:08:04	1/01/201	18:09:59	31%	27.13	65.80	Benthic
bird065	F	2016	1	16/12/201	06:42:48	19/12/20	19:09:53	3%	21.02	53.93	unkno
bird204	M	2016	1	25/12/201	19:27:20	26/12/20	20:27:47	29%	23.09	60.49	unkno
bird193	M	2016	1	18/12/201	04:03:08	20/12/20	14:50:08	3%	30.84	72.70	unkno
bird249	M	2017	1	4/12/2017	01:14:04	5/12/201	21:37:59	3%	1.17	0.61	Pelagic
bird250	F	2017	1	4/12/2017	02:29:04	6/12/201	23:32:59	23%	29.95	64.59	Benthic
bird247	F	2017	1	4/12/2017	05:44:04	4/12/201	19:54:59	5%	4.04	5.24	Pelagic
bird233	F	2017	1	4/12/2017	06:01:04	4/12/201	17:54:59	3%	3.33	3.61	Pelagic
bird244	M	2017	1	4/12/2017	18:33:04	5/12/201	18:16:59	7%	6.45	19.89	Pelagic
bird117	M	2017	1	10/12/201	15:16:04	12/12/20	23:15:59	8%	27.25	49.57	Benthic
bird068	M	2017	1	10/12/201	15:37:04	11/12/20	15:52:59	4%	12.82	23.76	Benthic
bird142	F	2017	1	11/12/201	05:19:04	13/12/20	20:52:59	2%	5.53	17.38	Pelagic
bird095	F	2017	1	11/12/201	15:55:04	12/12/20	13:30:59	38%	18.36	26.94	Pelagic
bird053	F	2017	1	13/12/201	15:39:04	15/12/20	11:57:59	13%	11.62	31.90	Pelagic
bird035	M	2017	1	13/12/201	17:30:04	15/12/20	18:11:59	2%	9.83	18.60	Pelagic
bird013	M	2017	1	14/12/201	14:00:04	16/12/20	23:18:59	16%	22.87	98.95	Pelagic
bird042	F	2017	1	15/12/201	05:33:04	15/12/20	15:42:59	4%	2.39	2.72	Pelagic
bird052	F	2017	1	15/12/201	05:51:04	18/12/20	13:47:59	31%	23.61	42.32	Pelagic
bird155	M	2017	1	16/12/201	04:49:04	16/12/20	20:13:59	5%	6.19	9.91	Pelagic

bird245	F	2017	1	16/12/201	06:29:04	16/12/20	20:12:59	2%	1.58	1.06	Pelagic
bird244	M	2017	2	16/12/201	11:09:04	17/12/20	00:57:59	4%	4.16	6.67	Pelagic
bird254	F	2017	1	16/12/201	15:23:04	17/12/20	10:02:59	26%	20.14	53.81	Pelagic
bird249	M	2017	2	17/12/201	05:34:04	17/12/20	20:41:59	1%	4.29	5.63	Pelagic
bird247	F	2017	2	17/12/201	05:34:04	18/12/20	16:12:59	3%	5.59	11.77	Pelagic
bird086	M	2017	1	17/12/201	05:49:04	17/12/20	19:48:59	16%	8.97	17.12	Pelagic
bird233	F	2017	2	18/12/201	05:50:04	18/12/20	17:13:59	9%	22.58	65.33	Pelagic
bird025	F	2017	1	19/12/201	16:06:04	20/12/20	12:46:59	12%	5.11	8.79	Pelagic
bird106	M	2017	1	21/12/201	15:47:04	22/12/20	22:09:59	2%	1.65	1.17	Pelagic
bird160	M	2017	1	23/12/201	06:00:04	23/12/20	16:47:59	25%	15.49	28.90	Pelagic
bird161	F	2017	1	23/12/201	18:17:04	25/12/20	16:40:59	15%	18.08	71.96	Pelagic
bird101	F	2017	1	24/12/201	05:46:04	24/12/20	18:26:59	5%	5.79	8.55	Pelagic
bird160	M	2017	2	24/12/201	13:22:04	24/12/20	21:59:59	9%	4.72	7.40	Pelagic
bird050	M	2017	1	24/12/201	19:52:04	25/12/20	16:52:59	9%	2.65	3.29	Pelagic
bird161	F	2017	2	26/12/201	05:28:04	26/12/20	11:34:59	4%	1.62	3.24	Pelagic
bird022	F	2017	1	27/12/201	05:55:04	27/12/20	13:16:59	6%	3.65	5.70	Pelagic
bird267	F	2017	1	2/01/2018	07:04:04	2/01/201	19:18:59	33%	12.68	25.30	Pelagic
bird270	F	2017	1	2/01/2018	13:03:04	2/01/201	21:00:59	17%	7.22	14.03	Pelagic
bird268	M	2017	1	3/01/2018	13:33:04	3/01/201	21:40:59	15%	7.51	12.18	Pelagic
bird269	F	2017	1	3/01/2018	14:58:04	4/01/201	15:03:59	49%	13.58	41.15	Pelagic

1

2



3

4 Figure S2: A) Foraging range (maximum distance travelled offshore), and B) Total distance

5 (cumulative distance travelled per foraging trip) by foraging type for breeding yellow-eyed penguins

1 from Enderby Island. Distance was determined from GPS logs and foraging type was determined from
2 analysis of simultaneous dive (TDR) logs (Muller et al. 2020a).

3

4