



HAL
open science

AP2/ERF transcription factors orchestrate very long chain fatty acid biosynthesis during Arabidopsis lateral root development

Soazig Guyomarc'H, Yohann Boutté, Laurent Laplaze

► To cite this version:

Soazig Guyomarc'H, Yohann Boutté, Laurent Laplaze. AP2/ERF transcription factors orchestrate very long chain fatty acid biosynthesis during Arabidopsis lateral root development. *Molecular Plant*, 2021, 14 (2), pp.205-207. 10.1016/j.molp.2021.01.004 . hal-03432678

HAL Id: hal-03432678

<https://hal.science/hal-03432678>

Submitted on 25 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

1 **AP2/ERF transcription factors orchestrate Very Long Chain Fatty** 2 **Acids biosynthesis during Arabidopsis lateral root development**

3
4 Soazig Guyomarc'h¹, Yohann Boutté² and Laurent Laplaze^{1,*}

5 ¹UMR DIADE, Univ. Montpellier, IRD, Montpellier, France

6 ²LBM, Bordeaux, France

7 *Correspondence: Laurent Laplaze (laurent.laplaze@ird.fr)

8
9 One key factor defining root system architecture is root branching i.e. the process of the formation of new root
10 axes, called lateral roots (LR), from an existing root. In the model plant *Arabidopsis thaliana*, LR originate
11 from pre-selected pericycle cells that divide to create a LR primordium (LRP) that grows and eventually forms
12 a new root apical meristem (Banda et al., 2019). During this developmental process, cell fate acquisition is
13 very plastic and depends on positional information rather than on cell lineage (Banda et al., 2019). Importantly,
14 LR formation occurs inside the parent root and the LRP must grow through overlying tissues before emerging
15 into the soil. This lateral root emergence process is controlled by dynamic intercellular signaling and involves
16 a tight coordination of cellular responses in the LRP and surrounding tissues (Banda et al., 2019). In particular,
17 the growing LRP rapidly pushes against the overlying endodermis, a layer of cells mechanically reinforced by
18 the Casparian strip. The crossing of the endodermis by the LRP between stage IV and V has been proposed as
19 an important developmental checkpoint during lateral root development (Banda et al., 2019).

20 21 **The AP2/ERF transcription factors PUCHI and ERF13 regulate LR development**

22 The APETALA2/ETHYLENE-RESPONSIVE FACTOR (AP2/ERF) transcription factors (TFs) form a large
23 group of TFs (147 genes in Arabidopsis) that are mainly found in plants (Feng et al., 2020). One AP2/ERF TF,
24 named PUCHI, was shown to be an important regulator of LR development in Arabidopsis (Hirota et al., 2007).
25 The *puchi-1* loss-of-function mutant exhibits higher LR initiation rate and forms LRPs with an abnormal shape
26 that are delayed in development and emergence (Hirota et al., 2007; Trinh et al., 2019). Interestingly, an
27 increased proportion of LRPs are blocked before the endodermis crossing checkpoint in *puchi* compared to
28 wild type (Trinh et al., 2019). Altogether, all published data indicate that the AP2/ERF TF PUCHI coordinates
29 cell proliferation during LR development, while the downstream targets of this TF were not known.

30 Recently, Lv et al. (Lv et al., 2020) reported that another Arabidopsis AP2/ERF TF, ERF13, played an important
31 role in LR development. Auxin, a key regulator of different steps of LR development (Banda et al., 2019),

32 activates the MITOGEN-ACTIVATED PROTEIN KINASE 14 (MPK14; (Lv et al., 2020)). A loss-of-function
33 mutation in *MPK14* or treatment with a map kinase kinase inhibitor (U0126) disturbed the progression of LRP
34 development and in particular the transition between stage IV and V (Lv et al., 2020). Using different *in vitro*
35 and *in vivo* assays, they demonstrated that MPK14 phosphorylates ERF13 leading to its degradation (Lv et al.,
36 2020). While an *erf13* loss-of-function mutant did not show any LR phenotype, *ERF13* overexpression led to
37 reduced LR development and an increased proportion of arrested LRP at the stage IV to V transition,
38 reminiscent of the *mpk14* phenotype (Lv et al., 2020). This indicates that ERF13 is a negative regulator of LR
39 development that is degraded upon auxin perception through a MPK14-dependent pathway to allow LRP
40 development to proceed through the stage IV to V developmental checkpoint (Figure 1A).

41

42 **AP2/ERF as master regulators of VLCFA biosynthesis during LR development**

43 Recent studies revealed that both AP2/ERF TFs PUCHI and ERF13 control LR development, at least in part,
44 through regulation of Very Long Chain Fatty Acid (VLCFA) biosynthesis. VLCFAs are fatty acids of at least
45 20 carbon atoms (C20) that can either be present as free CoA-VLCFAs or incorporated into lipid pools, mainly
46 into sphingolipids (SLs) and phosphatidylserine (PS) for membrane lipids, and into suberin and cuticle/cutin
47 for extracellular lipids (Li-Beisson et al., 2013). VLCFAs are synthesized by the elongase complex through a
48 4-step mechanism: a condensation by a β -ketoacyl-CoA synthase (KCS, 21 genes in *Arabidopsis*), a reduction
49 by a β -ketoacyl-CoA reductase (2 genes *KCR1* and *KCR2* in *Arabidopsis*), a dehydration catalyzed by the β -
50 hydroxyacyl-CoA dehydratase HCD/PAS2 and a reduction catalyzed by the enoyl-CoA reductase ECR/CER10.
51 While KCR, PAS2 and CER10 have a broad range of substrates, KCS enzymes display specificities towards
52 the acyl-chain length of their substrate (Haslam and Kunst, 2013).

53 Using a systems biology approach, Trinh et al. (Trinh et al., 2019) identified a subset of genes, including *KCS1*,
54 *KCS2*, *KCS20*, whose expression was induced in a PUCHI-dependent manner in developing LRP (Figure 1).
55 Accordingly, lipidomic profiling revealed that *puchi-1* display reduced content in VLCFA compared to WT.
56 Moreover, mutants perturbed in VLCFA biosynthesis showed similar lateral root developmental defects as
57 *puchi-1*. Hence, PUCHI activates a subset of VLCFA biosynthetic genes to regulate cell proliferation and
58 allows correct organogenesis during lateral root development.

59 Similarly, using transcriptomics to compare gene expression in WT and *ERF13* loss and gain-of-function lines,
60 Lv et al. (Lv et al., 2020) found that expression of 3 *KCS* genes (*KCS8*, *KCS16*, and *KCS18*) was negatively
61 regulated downstream of ERF13 (Figure 1). Accordingly, VLCFA content was reduced in *35S:ERF13* lines
62 compared to WT. Furthermore, a *kcs16* loss-of-function mutant showed similar LR development defects to
63 *35S:ERF13* lines and overexpression of *KCS16* or treatment with VLCFA could rescue the LR development
64 phenotype of *35S:ERF13* lines (Lv et al., 2020).

65 Thus, the AP2/ERF TFs ERF13 and PUCHI have opposite effects on LR development but both act, at least in

66 part, through regulation of VLCFA biosynthesis via the transcriptional control of multiple *KCS* genes (Figure
67 1A). In both cases, VLCFA biosynthetic genes were indirect targets of *PUCHI* and *ERF13*.

68 69 **Distinct pools of VLCFAs might have different roles in LR development**

70 This raises the question of the role of VLCFAs during LR development. VLCFAs can be incorporated into
71 extracellular lipids, including suberin and cuticle, and integrated into the cell wall. In *Arabidopsis*, an atypical
72 cutin layer covers the root apical meristem and developing LRP (Berhin et al., 2019). Perturbing the deposition
73 of the cuticle-like layer over the LRP leads to delay in LRP emergence and deformation of LRP, possibly due
74 to mechanical interactions between the LRP and cortical cells (Berhin et al., 2019). Thus, the cuticle-like layer
75 covering LRP may act as a “lubricant” to facilitate emergence through cortical and epidermal cell layers.
76 However, this process may not primarily involve VLCFAs, given that the main components are C18 fatty acids
77 (Berhin et al., 2019).

78 VLCFAs are also included into suberin, which is mainly present in the root endodermis. Suberin biosynthesis
79 and degradation play a role in LR formation (Li et al., 2017). However, *ERF13* loss and gain-of-function lines
80 or *puchi* loss-of-function do not seem to alter suberin formation (Trinh et al., 2019; Lv et al., 2020) suggesting
81 that VLCFA-suberin biosynthesis is not the main target of *ERF13* and *PUCHI* during LR formation.

82 Membrane lipids are other potential targets especially given that SLs, which contain VLCFA, are known to act
83 in LR formation (Roudier et al., 2010). Importantly, different profiles of VLCFAs incorporated into
84 endomembranes and plasma membranes could impact their structural and functional properties, e.g. fluidity or
85 integration of transmembrane proteins (Wattelet-Boyer et al., 2016).

86 87 **Future directions**

88 Published results suggest that *PUCHI* and *ERF13* do not regulate the same *KCS* gene sets during LR
89 development (Trinh et al., 2019; Lv et al., 2020) thus hinting that different pools of VLCFA-containing lipids
90 are produced in a very dynamic and localized manner during LR development. Indeed, membrane or
91 extracellular VLCFA-containing lipids differ in fatty acid composition, for instance for chain length or
92 modification of the chain by hydroxylation. Thus, a metabolic channeling might direct specific VLCFA
93 signature to different pools of lipids at precise domains in LRP. The identification of the direct regulators of
94 VLCFA biosynthetic genes could reveal if specific VLCFA signatures are differentially triggered during LR
95 development. Moreover, the characterization of the distinct lipid pools containing VLCFA signatures, as well
96 as their localization in LRP and their impact in cell membrane composition and properties constitute another
97 important target.

98

99 **Funding**

100 Research in the authors's lab is supported by the ANR projects NewRoot (n°ANR-17-CE13-0004 to SG) and
101 caLIPSO (ANR-18-CE13-0025 to YB).

102

103 **Acknowledgements**

104 The authors thank Dr Timothy TRANBARGER (IRD) for critical reading of the manuscript. No conflict of
105 interest is declared.

106

107 **References**

108 **Banda, J., Bellande, K., von Wangenheim, D., Goh, T., Guyomarc'h, S., Laplaze, L., and Bennett, M. J.**
109 (2019). Lateral Root Formation in Arabidopsis: A Well-Ordered LReXit. *Trends in Plant Science*
110 **24**:826–839.

111 **Berhin, A., Bellis, D. de, Franke, R. B., Buono, R. A., Nowack, M. K., and Nawrath, C.** (2019). The Root
112 Cap Cuticle: A Cell Wall Structure for Seedling Establishment and Lateral Root Formation. *Cell*
113 **176**:1367-1378.e8.

114 **Feng, K., Hou, X.-L., Xing, G.-M., Liu, J.-X., Duan, A.-Q., Xu, Z.-S., Li, M.-Y., Zhuang, J., and Xiong,**
115 **A.-S.** (2020). Advances in AP2/ERF super-family transcription factors in plant. *Critical Reviews in*
116 *Biotechnology* **40**:750–776.

117 **Goh, T., Toyokura, K., Yamaguchi, N., Okamoto, Y., Uehara, T., Kaneko, S., Takebayashi, Y., Kasahara,**
118 **H., Ikeyama, Y., Okushima, Y., et al.** (2019). Lateral root initiation requires the sequential induction
119 of transcription factors LBD16 and PUCHI in Arabidopsis thaliana. *New Phytol* **224**:749–760.

120 **Haslam, T. M., and Kunst, L.** (2013). Extending the story of very-long-chain fatty acid elongation. *Plant*
121 *Science* **210**:93–107.

122 **Hirota, A., Kato, T., Fukaki, H., Aida, M., and Tasaka, M.** (2007). The Auxin-Regulated AP2/EREBP Gene
123 PUCHI Is Required for Morphogenesis in the Early Lateral Root Primordium of Arabidopsis. *Plant*
124 *Cell* **19**:2156–2168.

125 **Li, B., Kamiya, T., Kalmbach, L., Yamagami, M., Yamaguchi, K., Shigenobu, S., Sawa, S., Danku, J. M.**
126 **C., Salt, D. E., Geldner, N., et al.** (2017). Role of LOTR1 in Nutrient Transport through Organization
127 of Spatial Distribution of Root Endodermal Barriers. *Current Biology* **27**:758–765.

- 128 **Ly, B., Wei, K., Hu, K., Tian, T., Zhang, F., Yu, Z., Zhang, D., Su, Y., Sang, Y., Zhang, X., et al. (2020).**
129 MPK14-mediated auxin signaling controls lateral root development via ERF13 regulated very-long-
130 chain fatty acids (VLCFAs) biosynthesis. *Molecular Plant* **0**.
- 131 **Roudier, F., Gissot, L., Beaudoin, F., Haslam, R., Michaelson, L., Marion, J., Molino, D., Lima, A., Bach,**
132 **L., Morin, H., et al. (2010).** Very-Long-Chain Fatty Acids Are Involved in Polar Auxin Transport and
133 Developmental Patterning in Arabidopsis. *Plant Cell* **22**:364–375.
- 134 **Trinh, D.-C., Lavenus, J., Goh, T., Boutté, Y., Drogue, Q., Vaissayre, V., Tellier, F., Lucas, M., Voß, U.,**
135 **Gantet, P., et al. (2019).** PUCHI regulates very long chain fatty acid biosynthesis during lateral root
136 and callus formation. *PNAS* **116**:14325–14330.
- 137 **Wattelet-Boyer, V., Brocard, L., Jonsson, K., Esnay, N., Joubès, J., Domergue, F., Mongrand, S., Raikhel,**
138 **N., Bhalerao, R. P., Moreau, P., et al. (2016).** Enrichment of hydroxylated C24- and C26-acyl-chain
139 sphingolipids mediates PIN2 apical sorting at trans-Golgi network subdomains. *Nat Commun* **7**:12788.
- 140
- 141 **Ly, B., Wei, K., Hu, K., Tian, T., Zhang, F., Yu, Z., Zhang, D., Su, Y., Sang, Y., Zhang, X., et al. (2020).**
142 MPK14-mediated auxin signaling controls lateral root development via ERF13 regulated very-long-
143 chain fatty acids (VLCFAs) biosynthesis. *Molecular Plant*, in press. Published November 19, 2020,
144 <https://doi.org/10.1016/j.molp.2020.11.011>

145
146

147 **Figure legend**

148 **Figure 1.** Expression of sets of VLCFA biosynthetic genes is regulated by AP2/ERF transcription factors during
149 lateral root development.

150 (A) The AP2/ERF transcription factor PUCHI induces *KCSI*, *KCS2*, *KCS20* expression upon auxin-dependent
151 stimulation by LBD16 (Goh et al., 2019; Trinh et al., 2019). Auxin-dependent MPK-mediated phosphorylation
152 and subsequent degradation of the AP2/ERF transcription factor ERF13 results in induction of *KCS16*, *KCS8*
153 and *KCS18* expression in developing LRP (Lv et al., 2020). Expression of the different KCSs leads to the
154 production of VLCFAs.


155 (B) PUCHI and ERF13 regulate the expression of *KCS* genes in developing LRP. In both cases the controlled
156 induction of VLCFA biosynthesis is necessary for correct LRP development and especially breaking through
157 the endodermis layer and proceeding to LRP emergence. Synthesized VLCFAs may incorporate into
158 hydrophobic cell wall materials (cuticle, suberin) and/or participate to endomembrane and plasma membrane

159 structures and functional dynamics in the LRP.

160

161

A



B

