



Chapter X Alternative DNA repair pathways to handle complex DNA damage generated by oxidative stress and anticancer drugs

E Matta, U Aliyaskarova, A A Kuznetsova, B T Matkarimov, O S Fedorova, N A Kuznetsov, A A Ishchenko, M Saparbaev

► To cite this version:

E Matta, U Aliyaskarova, A A Kuznetsova, B T Matkarimov, O S Fedorova, et al.. Chapter X Alternative DNA repair pathways to handle complex DNA damage generated by oxidative stress and anticancer drugs. DNA Damage, DNA Repair and Disease, 1, Royal Society of Chemistry, 2020, Chemical Biology, 978-1-78801-889-0. <10.1039/9781839160769>. <hal-03432737>

HAL Id: hal-03432737

<https://hal.science/hal-03432737v1>

Submitted on 17 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Chapter X

Alternative DNA repair pathways to handle complex DNA damage generated by oxidative stress and anticancer drugs

E. Matta^{a,1}, U. Aliyaskarova^{a,1}, A. A. Kuznetsova^{b,1}, B. T. Matkarimov^c, O. S. Fedorova^b, N. A. Kuznetsov^{b,*}, A. A. Ishchenko^{a,*} and M. Saparbaev^{a,*}

^aGroupe «Réparation de l'ADN», Equipe Labellisée LIGUE Contre le Cancer 2016, CNRS UMR9019, Université Paris-Sud, Gustave Roussy Cancer Campus, F-94805 Villejuif Cedex, France. ^bInstitute of Chemical Biology and Fundamental Medicine, Siberian Branch of Russian Academy of Sciences, Novosibirsk 630090, Russia. ^cNational laboratory Astana, Nazarbayev University, Astana 010000, Kazakhstan.

*corresponding email addresses: murat.saparbaev@gustaveroussy.fr; Nikita.Kuznetsov@niboch.nsc.ru; Alexander.ISHCENKO@gustaveroussy.fr

¹These authors contributed equally to this work.

ABSTRACT

The clinical features of inherited human DNA repair deficient disorders such as Cockayne syndrome and Fanconi anemia point to the complex nature of endogenous oxidative DNA damage, which may include bulky adducts, inter-strand DNA crosslinks (ICLs) and clustered lesions. Conversely, severe biological effects of DNA crosslinking agents and ionizing radiation correlated with formation of ICLs and double-strand breaks in DNA, respectively. These complex DNA damages are postulated to be critical because they are more difficult to repair than singular lesions. It anticipated that the removal of ICLs and clustered oxidized bases on both strands would, if not tightly regulated, either inhibit certain steps of repair or produce persistent chromosome breaks and thus be lethal for the cells. Genetic and biochemical data indicate that the elimination of complex damages requires several distinct DNA repair pathways including: base excision repair, nucleotide incision repair, Poly(ADP-ribose) polymerases-mediated DNA strand break repair, global genome and transcription coupled nucleotide excision repair, mismatch repair, homologous recombination, non-homologous end joining, and translesion DNA synthesis pathways. In this review, we describe the role of recently discovered alternative DNA repair pathways in the removal of complex DNA lesions.

X.1 Chemical nature of complex DNA damage

Cellular DNA constantly undergoes chemical alterations, resulting in DNA lesions that are cytotoxic, miscoding or both and are believed to be at the origin of mutagenesis and cell lethality.^{1,2} DNA damage can be classified by their nature: spontaneous *versus* induced; by their structure: bulky *versus* nonbulky, base *versus* sugar damage, singular *versus* complex damage (Figure 1). Complex DNA damage (CDD) such as bulky DNA adducts, inter-strand DNA crosslinks (ICLs) and clustered lesions (including double-strand breaks, DSBs) have complex character/structure as compared to singular lesions such as randomly distributed abasic sites, deaminated, alkylated and oxidized DNA bases.

[Insert Figure X.1 here]

Two of the most important features of CDD are their bulky character and presence of more than one modification within one turn of DNA helix. Although CDD, such as ICLs and DSBs, typically constitute relatively small fraction of the total DNA damage induced by DNA crosslinking agents and ionizing radiation, respectively, they are expected to play disproportionately important biological roles, since if not repaired, these CDD could lead to a number of serious consequences, such as gross chromosomal rearrangements and genome instability.

X.1.2. Repair-resistant bulky DNA adducts

X.1.2.1. 8,5'-Cyclopurine-2'-deoxynucleosides

Reactive oxygen species (ROS), produced during oxidative cellular metabolism and by ionizing radiation, damage cellular DNA leading to formation of CDD. Bulky DNA adducts such as diastereoisomeric (5'*S*)- and (5'*R*)-8,5'-*cyclo*-2'-deoxyadenosine

Figure 1




Figure 1. Schematic presentation of singular and complex DNA damage. **(A)** Chemical structures of some singular non-bulky DNA base lesions. **(B)** Chemical structures of complex DNA lesions including cisplatin-ICL and bulky repair-resistant DNA base lesions.

(cdA) and 8,5'-*cyclo*-2'-deoxyguanosine (cdG) are generated by hydroxyl radical attack at C5' sugar by H-abstraction resulting in the formation of a covalent bond between the C5'- and C8-positions of the purine nucleoside³ (Figure 1B). These DNA adducts are helix-distorting lesions and strong blocks for both DNA replication and transcription.^{4,5} It was demonstrated that the global genome nucleotide excision repair (GG-NER) pathway can remove cdA adducts with an efficiency comparable to that of UV-induced cyclobutane pyrimidine dimers, but exhibits higher activity in excising the *R*-isomer.^{4,6} Indeed, cdG and cdA adducts accumulate in keratinocytes from NER-deficient xeroderma pigmentosum (XP) complementation group C (XPC) and Cockayne syndrome (CS) complementation group A (CSA) patients exposed to X-rays and potassium bromate (KBrO₃)^{7,8} and also in organs of CSB^{-/-} knockout mice.⁹ Noteworthy, cdA and cdG lesions accumulate in age dependent manner in nuclear DNA of wild-type mice, implying the difficulties in the removal of these complex DNA lesions.¹⁰ Also, the *S*-cdA diastereoisomer present at higher levels in organs from control non-exposed mice because they are removed by GG-NER much less efficiently than the corresponding the *R*-cdA diastereomer.^{10,11}

X.1.2.2. Bulky DNA adducts induced by environmental carcinogens

Bulky DNA adducts that are generated by exposure to environmental carcinogens cause mutations that can drive malignant transformation of affected cells. As we described above, bulky DNA adducts are removed in the GG-NER pathway; however, certain bulky lesions that cause a minimal decrease, or an enhancement in the stabilities of the DNA duplex, can be very resistant to the repair machinery. Benzo[*a*]pyrene (B[*a*]P) is a polycyclic aromatic hydrocarbon (PAH) and well known carcinogen, which upon metabolic activation to diol epoxide can interact with guanines in DNA to form bulky B[*a*]P-*N*²-dG adducts. In contrast, dibenzo[*a,h*]pyrene

(DB[a,l]P), another PAH that is present in about ten-fold lower concentration in the environment, is 100 times more tumorigenic than B[a]P in rodent model systems.¹²⁻¹⁷ The reactive diol epoxide intermediates of DB[a,l]P react with either *N*⁶-adenine or *N*²-guanine in DNA to form stable DB[a,l]P-*N*⁶-dA and DB[a,l]P-*N*²-dG adducts¹⁸⁻²⁰ (Figure 1B). Remarkably, Kropachev et al., demonstrated that the *S* and *R* diastereoisomers of DB[a,l]P-*N*²-dG adduct are 15 and 35 times more efficiently removed by GG-NER, respectively, as compared to the stereochemically identical DB[a,l]P- *N*⁶-dA adducts.²¹ This observation suggests that the higher genotoxic activity of DB[a,l]P, as compared to B[a]P, might be due to the generation of repair-resistant and thus persistent DB[a,l]P-derived adenine adducts in exposed cells.

Other bulky DNA lesions, that cause little or no distortion and small thermodynamic destabilization of DNA duplex, also exhibit NER resistance including aristolactam-DNA adducts generated by the exposure to aristolochic acid^{22,23} (Figure 1B), 2-acetylaminofluorene AAF-*N*²-dG adduct,^{24,25} 3-benzanthrone derived DNA adducts,²⁶ and stable Fapy-dG adducts derived from aflatoxin B1.²⁷ Thus, despite their bulky character, certain highly mutagenic DNA lesions can escape DNA damage surveillance and repair. These repair-resistant DNA damage present a challenge for the cell since their can persist in the genome and lead to the transcription and replication blockages and mutations.

X.1.3. Formation and chemical nature of DNA crosslinks

Bifunctional alkylating agents, such as nitrogen mustard (HN2), psoralen and mitomycin C (MMC), can generate a covalent bond between nucleotides on opposite strands of a DNA duplex resulting in formation of ICLs. ICLs are highly lethal DNA lesions that block DNA replication, transcription and recombination by preventing

strand separation. Due to their high cytotoxicity, ICL-inducing agents are widely used against hyperplastic diseases, such as cancer and psoriasis. 8-methoxypsoralen (8-MOP) is a planar, tricyclic compound that intercalates into DNA duplex preferentially at 5'-TpA sites. Upon photoactivation, 8-MOP primarily photoalkylates DNA by cycloaddition to the 5,6-double bond of a thymine generating monoadducts (MA) and ICLs.^{28,29} Although, the yield of psoralen MAs to pyrimidine bases is three-fold higher than that of ICLs, the latter class of damage appears to have more severe biological effect.³⁰ Derivatives of HN2 induce a variety of DNA lesions, and only 5% among them constitute cross-links between the N7 position of guanine bases on opposite DNA strands.³¹ The platinum compounds such as cis-diamminedichloroplatinum (II), also known as cisplatin, reacts with guanines and induces mainly DNA diadducts: 65% d(GpG) intra-strand cross-links, 25% d(ApG) intra-strand cross-links and 5–8% ICLs between the guanines in the sequence d(GpC). The structures of HN2 and cisplatin-induced ICLs revealed that these agents induce dramatic distortions to the DNA helix,³² whereas MMC and psoralen-induced ICLs cause relatively minor distortions.³³ Endogenous sources such as aldehydes formed as products of lipid peroxidation and nitrous acid can also form ICLs in non-treated cells.³⁴ In addition to ICLs, formaldehyde, ionizing and UV radiations can crosslink proteins to an undisrupted DNA strand generating a DNA-protein crosslink (DPC).³⁵

X. 2 DNA glycosylase-mediated repair of complex DNA lesions

X. 2. 1. DNA glycosylase-mediated removal of bulky DNA lesions

Bulky DNA adducts and ICLs are highly toxic for cells and accumulation of endogenously occurring ICLs have been linked to premature aging.^{36,37} Although the GG-NER pathway is thought to be main pathway for the removal of ICLs and bulky

134 lesions that disrupt DNA helix conformation,^{38,39} most of singular DNA lesions
135 generated by oxidative stress are eliminated by two distinct pathways: base excision
136 and nucleotide incision repair (BER) and (NIR), respectively.^{40,41}

137 BER is initiated by a DNA glycosylase that recognizes and excises damaged
138 bases by hydrolysis of the glycosidic bond. In general, substrate specificities of DNA
139 glycosylases and AP endonucleases are limited to structurally small non-bulky DNA
140 base damage that have little impact on thermodynamic stability of DNA duplex.

141 Majority of DNA glycosylases bind to damaged DNA and flip out of the duplex the
142 aberrant nucleotide into the enzyme's active site pocket, while inserting residues in
143 the void left in the helix to stabilize the DNA substrate conformation.⁴²⁻⁴⁴ In general,
144 nucleobase binding pockets are too small to accommodate large size DNA adducts
145 and ICLs; however, several lines of evidence argue that DNA glycosylases are able
146 to accommodate bulky DNA adducts despite steric constraints to fit these lesions
147 into their active sites. Specifically, it was demonstrated that DNA glycosylases Fpg,
148 T4 endonuclease V (T4 pyrimidine dimer glycosylase) and NEIL1 can recognize and
149 efficiently excise very bulky DNA adducts such as imidazole ring opened form of
150 guanine-C8-N-hydroxy-2-aminofluorene adduct, cyclobutane dimer, psoralen-
151 thymine monoadduct, and aflatoxin-Fapy-guanine adducts (AFB₁-FapyGua),
152 respectively.⁴⁵⁻⁴⁸ Structural studies revealed that T4 endonuclease V kinks the DNA
153 helix by about 60° and flips out the opposing adenine base complementary to the 5'
154 pyrimidine of the thymine dimer out of the DNA base stack, thus avoiding steric
155 problems.⁴⁶ Whereas, Fpg flips bulky *N*7-substituted Fapy-dG derivatives guanine
156 lesion out of the DNA helix to the binding pocket, but enables *N*7-bulky group to stay
157 outside.⁴⁹ This latter mechanism might be also employed by human NEIL1 DNA

glycosylase when acting upon large voluminous DNA lesions such as psoralen-thymine, protein-DNA crosslinks and AFB₁-FapyGua.^{47,48,50}

Recent studies by Eichmann laboratory revealed a new family of DNA glycosylases, which utilizes a non-base-flipping mechanism to recognize bulky DNA base damage and a certain type of ICL.⁵¹⁻⁵³ DNA glycosylases AlkC and AlkD originally discovered in *Bacillus cereus*⁵⁴ and constituting a distinct DNA glycosylases superfamily⁵⁵ can excise positively charged N³- and N7-alkylpurines with high specificity without inserting a damaged base into active site pocket.⁵¹ AlkD traps damaged bases in a sheared, base-stacked conformation, directly contacting the deoxyribose with catalytic active site residues, but not the nucleobase like other DNA glycosylases. AlkD is capable of excising bulky DNA adducts such as pyridyloxobutyl (POB) adducts, arising in DNA as a result of cigarette-smoke carcinogen – nitrosamine ketone, and N3-yatakemycinyladenine (YTMA) produced by extremely cytotoxic alkylating product yatakemycin (YTM).^{51,56,57} The ability of AlkD to resolve bulky YTM and POB DNA modifications located in any grooves is presumed to be due to the absence of steric restraint. Nevertheless, the presence of other classic alkylpurine DNA glycosylases in these bacteria points to the fact that catalytic activity of AlkD is limited to positively charged DNA lesions that can be easily removed naturally.^{51,58} AlkC DNA glycosylase contains immunoglobulin (Ig)-like domain that was not found in the known repair proteins, which along with HEAT-like repeats (HLR), allows the enzyme to almost completely wrap around DNA duplex and excise damaged nucleobases. AlkC acts through non-base-flipping mechanism similar to AlkD; however, it generates sharp kink in DNA duplex to gain access to the modified base through inserting its catalytic residues into the exposed space.⁵⁵

Azinomycin B (AZB) is a genotoxic non-ribosomal peptide-polyketide secondary metabolite produced by soil-dwelling bacteria including *Streptomyces sahachiroi* and *Streptomyces griseofuscus*. AZB generates ICLs in DNA duplexes at 5'-GNC or 5'-GNT sequence contexts by nucleophilic addition at the N7 positions of purines in a major groove.⁵⁹ The classical repair mechanisms used to resolve ICLs in bacterial cells consists of coordinated function of NER machinery and homologous recombination (HR).⁶⁰ Nevertheless, recently it has been found that AlkZ DNA glycosylase of *Streptomyces sahachiroi* is capable of unhooking AZB-induced ICLs through the cleavage of N-glycosidic bond on both sides of the complementary DNA strands.^{58,61,62} The existence of unusual alternative DNA repair mechanisms in toxin-producing cells may provide insight for potential mechanisms implicated in drug resistance in cancer cells.⁵³

X. 2. 2. DNA glycosylases mediated repair of inter-strand DNA crosslinks

Genetic and biochemical evidences suggest that in vertebrates, the repair of ICLs proceeds in DNA replication-dependent manner and coordinated by Fanconi anemia (FA) proteins. The removal of an ICL goes through formation of a DSB as a result of the unhooking via dual incisions on both sides of the lesion by the scaffolding protein SLX4 and structure-specific endonuclease XPF/ERCC1.^{63,64} The resulting unhooked ICL is processed by translesion synthesis (TLS) specific DNA polymerases yielding a three-stranded DNA repair intermediate composed of a short oligomer covalently bound to the duplex. It thought that this crosslinked fragment or unhooked ICL is removed by the classic GG-NER pathway.^{38,39} However, Couve et al., revealed that the oxidative DNA glycosylases *Escherichia coli* (*E. coli*) Nei and human NEIL1 excise with high efficiency the psoralen-induced bulky unhooked ICL fragment within a three-stranded DNA structure⁶⁵ (Figure 2A). Three Nei-like (NEIL) DNA

Figure 2




Figure 2. DNA replication-coupled repair of inter-strand DNA crosslinks. **(A)** NEIL1 excises unhooked crosslinked oligomer in three-stranded DNA repair intermediate. **(B)** NEIL3- and **(C)** Fanconi anemia-mediated repair of ICL. **(D)** The hypothetical mechanism of the FANCM-mediated DNA replication fork bypass of ICL and NEILs-mediated repair of three- and four stranded crosslinked DNA structures.

glycosylases are present in mammalian cells; these proteins show structural
homology to the Fpg and Nei proteins of *E. coli* and remove oxidised bases from
DNA.⁶⁶ While NEIL1 and NEIL3 appear to be cell cycle regulated, with expression
peaking in S phase and late S/G2 respectively, NEIL2 is constitutively expressed
throughout the cell cycle.^{67,68} All three proteins have DNA glycosylase and β or β/δ -
lyase activities, with an unusual preference for single-stranded DNA and other DNA
open structures generated during DNA replication and transcription.

[Insert Figure X.2 here]

Reconstitution of the repair of plasmids containing a single ICL in *Xenopus*
eggs extracts showed that the removal of ICL was coupled to DNA replication and
involved convergence of two replication forks on the lesion, with the formation of an
X-shaped DNA structure⁶⁹ (Figure 2B). Subsequently, it has been proposed that ICL
repair requires the convergence of two forks on the lesion since, when only one fork
was stalled at the ICL in egg extracts, no ICL repair took place.⁷⁰ On the other side,
Huang et al., investigated the collision of replication forks with fluorescently labelled
psoralen-ICLs in mammalian cells using DNA combing.⁷¹ They observed that during
S phase the majority of psoralen ICLs (around 60%) were processed through a
replication-traverse pathway, in which the ICLs are left unrepaired, but traversed by
the replication machinery to allow DNA synthesis to resume on the other side. In
these scenarios of fork traverse and the former one of dual fork convergence, a
similar X-shaped DNA structure is generated around the ICL, which is critical to
initiate ICL repair.⁷² Noteworthy, these unrepaired psoralen-induced ICLs are
subsequently removed during a post-replication repair without the generation of
DSBs.⁷¹ In the following work, Semlow et al., have demonstrated that NEIL3 from
Xenopus laevis can excise psoralen and abasic site ICLs in DNA in X-shaped

double-stranded DNA structures via an incision-independent repair mechanism, suggesting that this activity may be a principal role of NEIL3 in rapidly dividing cells⁷³ (Figure 2B). More recent biochemical studies by Martin et al., showed that human NEIL3 cleaves psoralen-induced ICL in single-stranded, three-stranded and four-stranded DNA substrates to generate unhooked DNA fragments containing either an abasic site or a psoralen-thymine monoadduct. In addition, Nei and NEIL1 also excised a psoralen-induced four-stranded DNA substrate to generate two unhooked DNA duplexes with a nick, while NEIL3 targeted both DNA strands in the ICL substrate without generating single-strand breaks. Based on these observations, Martin et al., proposed a model for the mechanism of ICL repair in mammalian cells that implicates the FANCM-catalysed replication traverse of an ICL site, without preceding DNA incision and the generation of either three- or four-stranded DNA structures composed of the duplex cross-linked either to single-stranded or duplex DNA⁷⁴ (Figure 2C). These putative ICL lesion bypass products can then be repaired by the Nei-like DNA glycosylases-initiated BER pathway without generation of highly toxic DSB (Figure 2C).

Thus, the ICL repair in eukaryotic cells is coupled to DNA replication and proceeds *via* two alternative mechanisms: (i) Fanconi anemia system incision-dependent pathway, which engenders replication fork collapse and generation of DSBs; and, (ii) the incision-independent pathway in which NEIL1 and NEIL3 DNA glycosylases mediate ICL unhooking without formation of highly genotoxic DNA repair intermediates. Although, an ICL covalently links the two strands of the DNA, it is important to highlight that the level of distortion of the DNA helix and the choice of the DNA repair pathway depends on the chemical nature of ICL. Cisplatin- and nitrogen mustard-generated ICLs are preferentially processed via the incision-

dependent Fanconi anemia coordinated network,^{39,69} whereas, psoralen and abasic site induced ICLs processed rather via NEIL3-catalyzed BER pathway.^{73,74} Nonetheless, NEIL3^{-/-} knockout mouse embryonic fibroblasts (MEF) exhibit modest sensitivity to cisplatin,⁷⁵ implying that some cisplatin-ICLs might be removed in the BER pathway, which can serve as a back-up repair system for FA and NER. Importantly, it has been proposed that NEIL3-mediated repair of ICLs would be more mutagenic as compared to that of Fanconi anemia.⁷³

X. 2. 3. Aberrant repair of interstrand DNA crosslinks

Cisplatin [*cis*-diamminedichloroplatinum(II)] is well known for its capacity to trigger a strong DNA damage response signal, which may eventually result in irreversible apoptosis. Studies of the roles of BER enzymes in the repair of ICLs have produced conflicting results, suggesting involvement of multiple pathways and mechanisms depending on the cellular and structural contexts. For example, Kothandapani et al., have shown that inhibition of BER by methoxyamine (MX), a specific inhibitor of APE1-catalyzed cleavage of AP sites, along with the depletion of DNA polymerase β (Pol β) and uracil-DNA glycosylase (UNG) enhances the resistance of cells to cisplatin.⁷⁶ Using synthetic oligonucleotides with a single site-specific cisplatin-ICL, the authors demonstrated that the accelerated and preferential deamination of flipped out cytosine residues next to cross-linked guanines, can induce formation of uracil residues adjacent to ICL. These uracils are excised by UNG to generate AP sites, which then are cleaved by APE1, with following Pol β -catalyzed gap-filling DNA repair synthesis. The authors proposed that initiation of BER adjacent to cisplatin ICL sites could interfere with the NER and HR pathways and lead to the inhibition of ICL removal and increasing cisplatin cytotoxicity. Both repair and redox domains of APE1 appear to require for mediating cisplatin toxicity in breast cancer.⁷⁷ Several small

inhibitors developed against APE1, Pol β and PARP1 are currently studying in anti-cancer therapy [see (see Chapters X,X,X (Kelley, Curtin, Madhusudan))]. Thus, understanding the roles of BER proteins in the repair of complex DNA lesions might be important to counteract the resistance of cancer cells.⁷⁸

X.3. The apurinic/aprimidinic (AP) endonuclease–initiated nucleotide incision repair pathway for oxidative DNA damage

X.3.1. Substrate specificity of AP endonucleases as multifunctional enzymes

Recognition and removal of non-bulky damage of nitrogenous bases proceed via the BER pathway initiated by DNA glycosylases.⁷⁹⁻⁸¹ In general, the BER pathway includes sequential actions of two enzymes for DNA incision: a DNA glycosylase and an AP endonuclease (Figure 3, pathways 1-3). Noteworthy, DNA glycosylases generate highly genotoxic intermediates in DNA, e.g. AP-sites and blocking 3'-end groups, which should be removed in order to initiate the DNA repair synthesis and ligation. AP endonucleases are key participants in the BER process, since all three types of products generated by DNA glycosylases^{82,83} are repaired by these enzymes. An AP endonuclease hydrolyses the phosphodiester bond located 5' to the AP-site and introduces a break into the deoxyribosephosphate backbone (Figure 3, pathway 1), in addition, it removes the remaining 3'-blocking groups: either the 3' phospho α,β -unsaturated aldehyde (Figure 3, pathway 2) or the 3'-terminal phosphate group (Figure 3, pathway 3).

[Insert Figure X.3 here]

Previously, it was thought that the main biological function of AP endonucleases was the hydrolytic cleavage 5' at AP-site and subsequent formation of SSB with a 5'-

Figure 3




Figure 3. The scheme of BER and NIR pathways.

deoxyribophosphate and a 3'-hydroxyl group.^{84,85} However, certain AP endonucleases can recognise not only AP-sites, but also damaged nucleotides containing a modified base (Figure 3, pathway 4).⁸⁶ This DNA glycosylase-independent removal of damaged bases was named the NIR pathway.⁴¹ Afterwards, a number of studies have shown that AP endonucleases recognise a variety of structurally diverse types of lesions from bulky benzene-derived DNA adducts⁸⁷ up to small oxidatively damaged pyrimidines.⁸⁸ Noteworthy, the genetic dissection of BER and NIR functions of an AP endonuclease revealed that the NIR pathway is essential for protecting cells from potentially lethal oxidative DNA lesions, which cannot be removed by BER.⁸⁹

A comparison of amino acid sequences among AP endonucleases suggests that there are two structural families, judging by similarities to two prokaryotic enzymes: exonuclease III (Exo III, also referred as Xth)⁹⁰ and endonuclease IV (Endo IV, also referred as Nfo).⁹¹ Mammalian cells possess two AP endonucleases (APE1 and APE2) that are homologues of the bacterial Xth protein.⁹²⁻⁹⁴ APE1 contains a highly efficient AP endonuclease activity; however, its 3'-end cleansing including 3'→5' exonuclease activities is substantially lower than that of Xth.^{84,93,95-98} In addition to AP-sites, APE1 recognises a variety of damaged DNA bases, including α -anomers of 2'-deoxynucleosides (α A, α C and α T),⁹⁹ oxidized pyrimidines,⁸⁸ and uracil.¹⁰⁰ Contrary to APE1, the 3'→5' exonuclease activity of APE2 is higher than its AP endonuclease activity and may represent the main biological function of this enzyme.¹⁰¹ Of note, several other Xth-like AP endonucleases such as prokaryotic enzymes (*Mycobacterium tuberculosis* MtbXthA and *Neisseria meningitidis* NApe and NExo) and a plant enzyme (*Arabidopsis thaliana* atAPE1L) lack the NIR function.¹⁰²⁻¹⁰⁴

Furthermore, recently, a novel AP endonuclease, referred as endonuclease Q (EndoQ), which does not belong to Xth and Nfo-family, was identified in the hyperthermophilic euryarchaeon *Pyrococcus furiosus*. EndoQ is involved in the NIR pathway for uracil and hypoxanthine residues in DNA.^{105,106} Noteworthy, amino acid sequence analysis indicates that putative EndoQ homologs are present in most of Archaea, but only in a few groups in bacteria.¹⁰⁵ A homolog of EndoQ in the gram-positive bacterium *Bacillus pumilus*, exhibits NIR activity towards uracil residues.¹⁰⁷ EndoQ enzymes generate a nick immediately 5' to uracil, hypoxanthine, xanthine, and AP site in DNA.^{105,107} Notably, the endonuclease activity of *P. furiosus* EndoQ towards a uracil is identical to that of *E. coli* Xth acting upon AP-site.¹⁰⁶

It should be noted that the same catalytic mechanism of 5'-phosphodiester bond hydrolysis was proposed for members of both the Xth family^{92,108-111} and Nfo family enzymes.^{91,112,113} Despite differences in the structure and metal requirements among these AP endonucleases, a comparison of their active-site structures surprisingly reveals strong geometric conservation of the catalytic reaction.^{114,115} Nonetheless, there is still a debate in the literature about the functional role of certain amino acid residues in the active site of enzyme.¹¹⁶⁻¹¹⁹ In addition, it has been shown that during the endoribonuclease action of human APE1, the Mg²⁺ ion is not required, thus pointing to an alternative catalytic mechanism in the case of RNA substrates.¹²⁰

X. 3. 2. A putative physiological role of APE1-catalyzed NIR and 3'→5' exonuclease functions

Both human APE1 and its homolog APE2 contain 3'→5' exonuclease activity^{101,121} and this function is evolutionary conserved in Xth- and Nfo-families of AP endonucleases. Importantly, previous studies have established that *E. coli* Nfo and

human APE1-catalyzed NIR and 3'→5' exonuclease activities are functionally and genetically coupled and that the repair of oxidatively damaged base in the DNA glycosylase-independent NIR pathway can generate a small single-stranded (ss) gap 5' to the lesion in duplex DNA.^{88,122,123} On the other hand, it has been shown that *Xenopus* and human APE2 can exonucleolytically process single-strand DNA break (SSB) in the 3'→5' direction to generate an extended ssDNA gap, which in turn activates ATR-Chk1 DNA damage response (DDR) pathway.^{124,125} More recently, using *Xenopus* egg extracts Lin et al., demonstrated that APE1's exonuclease function is required for APE2-mediated 3'-resection of SSB and ATR-Chk1 signalling.¹²⁶ Strikingly, *Xenopus* APE1-D306A mutant, which corresponds to human NIR-deficient APE1-D308A variant,¹²³ is unable to trigger ATR-Chk1 DDR in egg extracts,¹²⁶ implying the role of NIR function in the generation of extended ssDNA gaps at oxidized DNA bases. Taken together, these observations led us to propose a schematic model in which the removal of oxidized bases in DNA glycosylase-independent NIR pathway generates extended ssDNA gap via coupled action of APE1 and APE2/PCNA complex (Figure 3, pathways 5-6). The resulting gapped ssDNA structure bound with RPA, triggers ATR-Chk1 DDR pathway in the replication-independent manner, which in turn enables the PCNA-directed gap filling by high-fidelity DNA polymerases (Figure 3, left panel). It is tempting to speculate that the NIR pathway may take place in actively transcribed open chromatin regions to ensure high-fidelity DNA repair synthesis, whereas, the classic BER pathway occurs in heterochromatin and proceeds mainly via DNA polymerase β -dependent short-patch pathway (Figure 3, pathways 2 and 3).

X. 3. 3. Conformational dynamics of enzyme–substrate complexes

Despite the active research into functional features of AP endonucleases, it is currently unknown how the active site of one enzyme can recognise substantially different (by structure and by nature) damaged and undamaged nucleotides. Certain important features of catalytic complex formation are still unclear: the sequence of the enzymatic events; the nature of interactions at the initial stage of substrate recognition; formation of which contacts leads to the discrimination between a substrate and 'non-substrate'; and which interactions ensure substrate specificity of the enzyme. Therefore, a major contribution to the understanding of the mechanisms underlying specific enzyme–substrate interactions has been done in studies utilizing pre-steady-state kinetics and thermodynamics of the enzymatic process with registration of conformational transformations of the interacting molecules.

Conformational transformations of AP endonucleases were recorded by means of a change in the intrinsic fluorescence of their tryptophan residues.¹²⁷⁻¹³² In nucleic acids, conformational transitions can be detected using fluorescent analogues of heterocyclic bases.^{119,133-141} It has been reported that the interaction of APE1 with a substrate includes at least two stages: DNA binding and recognition of an AP-site, which result in the formation of a catalytically-competent complex. The irreversible stage of catalytic hydrolysis of the 5'-phosphodiester bond at the AP-site takes place in this complex. The last stage of the kinetic mechanism is characterised by an equilibrium process of enzymatic-complex dissociation from the reaction product. Thermodynamic analysis of the fast stages of recognition and cleavage of DNA containing an abasic site¹³² has shown that there are two steps, DNA binding and F-site recognition that cause the formation of a catalytically competent complex.

X. 3. 4. The mechanism of substrate specificity towards damaged nucleotides, with human APE1 as an example

During the last three decades, extensive characterization of DNA repair enzymes provided insight into how damaged DNA bases were located and recognised among numerous unmodified bases.^{142,143} Structural data indicate that conformational changes in the enzyme and DNA play an important part in the recognition of specific substrates and may control the substrate specificity of enzymes.¹⁴⁴⁻¹⁵⁴

Nevertheless, a recent report uncovered substantial differences in the mechanisms of specific substrate recognition by many well-studied DNA glycosylases and by human APE1.¹⁵⁵ This study clarified the key steps of the mechanism underlying APE1–DNA interaction that enable high-specificity recognition of structurally different damaged DNA containing ϵ A, α A, 5,6-dihydrouridine, and an F-site. Accordingly,¹⁵⁵ the substrate specificity of APE1 should depend on the efficiency of damaged-nucleotide eversion from the duplex during enzyme-induced DNA bending. The ability of an AP site or damaged nucleotide to flip out of DNA and to penetrate the enzyme pocket during the formation of contacts between the enzyme and DNA may be the main contributing factor for substrate specificity of APE1. This conclusion means that the model of a mutual enzyme–substrate induced fit, as determined for DNA glycosylases,¹⁵⁶ is now transformed into the model of conformational selection in the case of APE1 ([Figure 4](#)). In this model, any nucleotide, which everts from the duplex in the response to a disturbance induced by the enzyme and enters into the active site, will be a substrate for the enzyme. This model of substrate specificity is well consistent with the ability of APE1 to act as an exonuclease and endoribonuclease because of the structural features of RNA that help with the eversion of intact nucleotides at the junction of the single-stranded parts, hairpins, duplexes, and loops. Moreover, it is reasonable to propose that non-B-form DNA and some structures formed due to DNA sequence flexibility can bind to APE1 and lead

to undamaged-nucleotide recognition with subsequent topology-specific DNA cleavage.

[Insert Figure X.4 here]

X. 4. Poly(ADP-ribose) polymerase catalysed covalent modification of DNA strand break extremities and its role in DNA repair

Endogenous and exogenous DNA damage are rapidly sensed and activate evolutionarily-conserved signaling pathways, known collectively as DDR, whose components can be separated into four functional groups: damage sensors, signal transducers, repair effectors, and arrest or death effectors.¹⁵⁷ Single-stranded, double-stranded and complex closely spaced clustered DNA strand breaks that represent either an initial DNA damage or DNA repair intermediates generated during DNA damage processing by cellular DNA repair enzymes are one of the fastest and most strong inducers of DDR. DNA break-sensing molecules, poly(ADP-ribose) polymerases (PARPs) are the key element of the DDR in mammalian cells. PARPs catalyze transfer of mono- or poly-ADP-ribose (MAR or PAR, respectively) moieties on proteins,¹⁵⁸ DNA^{159,160} and RNA¹⁶¹ by using nicotinamide adenine dinucleotide (NAD⁺) as a substrate. It has been shown that these post-translational, post-replicative and post-transcriptional modifications play roles not only in DNA damage repair,¹⁶² but also in a large array of cellular processes including regulation of gene expression, apoptosis and autophagy (reviewed in¹⁶³⁻¹⁶⁶).

The PAR polymer designates a chain of ADP-ribose (ADPr) units, where each unit is linked to other one through glycosidic ribose–ribose 1''→2' bonds.¹⁶⁷ This polymer can vary in length from a few to 200 – 400 ADPr units *in vivo* and *in vitro*.¹⁶⁸ The PAR can be branched at a frequency of 1 branch per 20-50 subunits of the linear

Figure 4




Figure 4. Mechanisms of DNA damage recognition by DNA glycosylases,^{90,104} **(A)** and AP endonucleases^{78,102} **(B)**.

polymer. Branched PAR polymer can have a very complex structure, including helicoidal secondary structures which are, at some extent, similar to RNA and DNA.¹⁶⁹ PAR (free or attached to proteins) that is synthesized upon genotoxic stress may undergo biphasic decay, with most (~85%) of PAR having a half-life of about 40 s and the remaining catabolized within approximately 6 min. In contrast, the constitutive fraction of PAR have a much longer half-life (~7.7 h).¹⁷⁰ A tight regulation of the level of the polymer in the cell by ADPr-protein hydrolases, that reverse the reaction by hydrolyzing the protein–ADPr bond or the bonds between two ADPr units of PAR is important, since its accumulation have strong cytotoxic effects.¹⁷¹

Family of PARPs, also known as diphtheria toxin-like ADP-ribosyltransferases includes 18 known members¹⁶⁸ that are divided into 4 sub-families as DNA dependent PARPs, Tankyrases, CCCH PARPs and macroPARPs.¹⁷² Among PARPs only PARP1, PARP2 and PARP3 are activated in DNA dependent manner, which signals the presence of DNA strand breaks.¹⁷³ PARP1-3 structures consist of multiple independent domains connected by flexible linkers.¹⁷⁴ Binding and recognition of sites of DNA strand breaks induce the reorganization of PARPs domains structures, promoting extensive inter-domain contacts and allosteric activation, by unfolding of an auto-inhibitory helical domain (HD), which then allows NAD⁺ binding necessary for ADP-ribosylation activity.¹⁷⁴⁻¹⁷⁷ PARP1 is the most ubiquitous and abundant PARP, which responsible for majority of the PARylation activity in a cell in response to DNA damage and involved in multiple DNA repair pathways, including BER, HR, NHEJ and NER.^{166,178,179} PARP2 plays partially redundant functions with PARP1 and accounts for about 10% of DNA damage-induced cellular PARylation activities, which are essential for cell survival. Depletion of PARP1 or PARP2 results in hypersensitivity to ionizing radiation, oxidative stress

and alkylating agents, whereas double-knockout mice show early embryonic lethality.¹⁸⁰ PARP3 is characterized by PARylation activity and act synergistically with PARP1 in response to genotoxic drugs and X-irradiation generating DSBs.¹⁸¹⁻¹⁸³ Importantly, PAR attached to a protein rapidly degraded by PAR glycohydrolase (PARG), except for the last PAR moiety, which remains bound to the protein. PARG specifically hydrolyses the ribose–ribose bonds and it is the main enzyme in mammalian cells. Disruption of PARG gene in mice results in embryonic lethality and PARG-deficient cells exhibited increased cell death and impaired repair of DNA base damage and strand breaks,¹⁸⁴ indicating that accumulation of the PARylated macromolecules is highly toxic to the cell.

The widespread presence of PARP proteins in eukaryotes and their protein modification activity are possibly due to the fact that in eukaryotic cells, DNA is tightly packed into a composite of DNA and proteins referred to as chromatin. Chromatin structure imposes restriction upon DNA-protein interactions, and several studies have demonstrated that tight chromatin packaging restrict the ability of DNA repair machinery to access the sites of DNA damage.^{185,186} PARPs mediated ADP-ribosylation of histones and other nuclear proteins induce recruitment and modification of chromatin remodeling factors, including chromatin insulator protein CTCF (CCCTC-binding factor) and chromodomain-helicase-DNA-binding protein 1-like (ALC1), and provide an effective chromatin remodeling mechanism necessary for the recruitment of repair proteins to DNA damage sites and efficient repair of DNA strand breaks (reviewed in^{164,179,187,188}). Moreover, covalently attached ADP-ribose polymer with a complex branched structure confers negative charge to PARPs and histones, resulting in decrease in DNA binding and electrostatic repulsion of these proteins from DNA.^{189,190} However, in the case of severe

genotoxic stress, hyperactivation of PARP results in strong reduction of NAD⁺ levels, which in turn, induce release of apoptosis-inducing factor (AIF) from mitochondria and lead to apoptosis in a caspase-independent manner.¹⁹¹

Recently, it has been demonstrated that *in vitro* PARP1-3 can reversibly ADP-ribosylate termini of DNA molecules containing multiple closely spaced DNA strand breaks (DSB and SSB), producing a covalent PAR/MAR–DNA adducts, pointing to the underlying complex nature of the PARP-dependent DNA strand break signaling.^{159,192,193} Mechanistic studies showed that PARP1 and PARP2 can covalently modify DNA duplexes by PARylation of 3'- and 5'-terminal phosphates and 2'-OH termini of modified nucleotides at DNA strand breaks.¹⁵⁹ Contrary to ADP-ribosylation of proteins, PAR and MAR DNA modifications can be effectively and completely removed by PARG.^{159,192,193} MACROD2, TARG1 and ARH3 hydrolases can also remove the MAR adduct from DNA, leaving phosphorylated termini.¹⁹³ PARP2 and PARP3 can ADP-ribosylate DNA termini in long linear and circular DNA duplexes, which more closely resemble the cellular DNA context.¹⁹² Most importantly, under certain configuration of strand breaks in duplex DNA, PARPs can switch their substrate specificity from auto-ADP-ribosylation to DNA-ADP-ribosylation.¹⁹² DNA PARylation experiments in cell-free extracts and immunoblotting experiments with purified genomic DNA after genotoxic treatment suggest that certain types of complex DNA strand breaks in chromosomal DNA can be ADP-ribosylated by PARPs in the cellular response to DNA damage.¹⁹²

The three-dimensional structures of PARP1 bound to DNA with a SSB revealed that PARP1 binds and bends DNA duplex at an SSB site through cooperative interactions of two N-terminal zinc finger domains (F1 and F2) with exposed DNA bases and adjacent minor grooves and then drives stepwise assembly

Royal Society of Chemistry – Book Chapter Template

of the remaining F3, WGR (Trp-Gly-Arg) and catalytic (CAT) domains leading to PARP1 activation.^{174,176} In contrast, PARP2 and PARP3 use only short N-terminal and WGR domains for binding to damaged DNA.^{181,194} However, the resolved crystal structures of assembled PARP-DNA complexes still do not provide sufficient insight into the observed PARPs specificity for the phosphorylated DNA termini. In these complexes, the DNA-binding site of PARP1 is far from its catalytic (CAT) domain in activated PARP1-DNA complex and sterically protected from ADP-ribosylation by the PARP1 protein itself. However, another proximal DNA strand breakage site in the same PARP-DNA complex might interact with the CAT domain taking into account high flexibility of domain linkers. A putative mechanistic model of DNA strand break-oriented ADP-ribosylation of DNA termini by PARP1-3 shown in [Figure 5](#) has been proposed based on the observed dependence of DNA termini covalent modification on the distance between the phosphorylated DNA terminus as an acceptor and the DNA strand break as an activator.¹⁹² The following rules have been established for the efficient ADP-ribosylation of 5' or 3' upstream DNA termini in duplex DNA containing multiple strand breaks: (i) the distance between phosphate residue and SSB should not exceed ~2 helix turns of DNA duplex; and (ii) the relative position of the acceptor phosphate on DNA helix to the DNA-bound and activated PARPs.¹⁹²

[Insert Figure X.5 here]

Specific requirements for closely spaced or clustered DNA strand breaks in the PARP-dependent DNA ADP-ribosylation open an important question: whether such DNA strand breaks can be formed for example by ionizing radiation, and if there are some DNA processing pathways that may generate strand breaks in a non-random manner? It is quite possible that DNA duplexes containing DSB and proximal SSB

Figure 5




Figure 5. The mechanistic model of DNA strand break-oriented DNA termini ADP-ribosylation by PARP1, PARP2 and PARP3.

can be generated during DSB repair by HR and NHEJ repair pathways. Intriguingly, it has been found that the MRN–CtIP complex generated an internal nick located 20 nucleotides downstream of 5'-termini of a DSB,¹⁹⁵ and that this configuration of closely-spaced DNA strand breaks is prone to covalent DNA ADP-ribosylation by the PARP1-3 proteins.

Importantly, not only DNA strand breaks activate PARPs catalyzed ADP-ribosylation. PARP1 can also be activated by DNA hairpins, cruciform structures, stalled replication forks, stably unpaired regions and other non-B-conformations of DNA.^{196,197} Thus, the presence of a single SSB or DSB near to some of non-B DNA structures or specific chromatin configurations could potentially trigger ADP-ribosylation of DNA strand breaks termini. Moreover, the pool of PARPs activators has been enlarged during different studies showing that PARPs can be activated independently of DNA by protein-protein interactions, post-translational modifications and by specific type of RNA structures (summarized in [Table 1](#)).^{163,172,198-200} In addition to PARP1 and Tankyrase, a new PARPs family member has been shown to be activated by a DNA damage independent way. During genotoxic stress, PARP2 is activated by binding to RNA via its SAP domain, in nucleoli, and this stimulates PAR formation.²⁰¹ On the other hand, PARP1 inactivation was observed upon binding to the histone variant macroH2A1.1,²⁰² also the deacetylation and sumoylation of PARP1 inhibit its self-PARYlation.¹⁹⁹

[Insert Table x.1 here]

The phenomenon of ADP-ribosylation of a terminal phosphate group at the DNA strand breaks opens important question about possible functional role of this new modifying activity of PARPs. It is generally accepted that covalently attached PAR

Table 1. Mediators of PARP1-3 activation

Type of activation	Activator	Activated PARP	References
Nucleic acids binding	non-B DNA structures (hairpins, cruciforms, and loops)	PARP1	197,204
	DNA breaks: nicks, gaps, flaps, DSB	PARP1-3	174-177,205,206
	small nucleolar RNA	PARP1	207
	Short rRNA and other single-stranded RNAs	PARP2	201
	PAR polymer	PARP2	208
Protein-Protein interactions	Nucleosomes and histones	PARP1	209,210
	Phosphorylated extracellular signal-regulated kinases, ERK		211
	Nicotinamide mononucleotide adenylyltransferase 1, NMNAT1		212
	CTCF		213
	TET1		214
	TOPO2		215
	Bromodomain adjacent to zinc finger domain 2A, BAZ2A/TIP5		216
	Human tyrosyl tRNA synthetase, TyrRS		217

Post-translational modifications	phosphorylation at Ser372 and Thr373 (by extracellular signal-regulated kinases, ERK1/2)	PARP1	218
	phosphorylation at Tyr907 (by c-Jun N-terminal kinase, JNK1/ MAPK8)		219
	phosphorylation at Ser782, Ser785, and Ser786 (by cyclin-dependent kinase, CDK2)		220
	acetylation at Lys498, Lys505, Lys508, Lys521, and Lys524 (by E1A binding protein, p300/ CREB-binding protein CBP)		221
	acetylation of 1-214 and 477-525 aa regions (by P300/CBP-associated factor, PCAF)		222
	MARylation by PARP3		223
	MARylation at Lys 521 (by Sirtuin 6, SIRT6)		224
	phosphorylation at Thr420, Thr622, Thr656 (by checkpoint kinase 2, CHK2)		225
	methylation at K508 (by Histone-lysine N-methyltransferase, SET7/9)		226

950

951

952

polymer confers negative charge to PARPs, histones and other PARylated proteins, resulting in decrease in DNA binding and electrostatic repulsion of these proteins from DNA.¹⁹⁰ Thus, PARylated proteins together with PAR-recruited complexes of DNA repair factors might be also withdrawn together from the site of DNA damage. In contrast, the PAR chain covalently attached to DNA termini should stably recruit those factors directly to the site of DNA damage. Importantly, it has been recently demonstrated that DNA PARylated at 5'-phosphate at 1 nt gap is recognized as 5'-adenylated DNA substrate and can be ligated in the absence of ATP by DNA ligase I, IIIa and by other DNA ligases, producing ligated double-stranded DNA molecule with an aberrant AP site. This AP site can be further processed by BER-associated APE1.¹⁶⁰ In line with these results, it has been suggested that PARP2 and PARP3 are involved in the final ligation step of NHEJ, because 5' phosphorylated nicks are particularly efficient activators of auto-ADP-ribosylation activity of PARP2 and PARP3, but not PARP1. We speculate that DNA ADP-ribosylation can promote retention of the DSB ends until either the full repair complex put in place or required ATP concentration for DNA ligation reached. Similarly, in the case of SSB repair PARP-mediated ADP-ribosylation can promote ligation of a gap without a polymerase synthesis and ATP. Of note, PARP-mediated extensive PAR synthesis can block the glycolysis and lead to ATP loss and the inhibition of hexokinase 1 activity.²⁰³

We hypothesize that the role of the ADP-ribosylation of DNA strand break termini is different from that of proteins (Figure 6) since it: (i) can be used as a temporary block of the processing of strand breaks and by this protect them from non-specific degradation and aberrant error-prone end joining; (ii) inhibits the NHEJ via inhibition of the binding of Ku to DSB ends; (iii) promotes direct aberrant ligation of ADP-

Figure 6




Figure 6. Putative and confirmed roles of the PARPs-catalyzed ADP-ribosylation in response to DNA damage.

ribosylated DNA breaks during bioenergetic collapse in order to avoid degradation and de novo formation of toxic DSBs; (iv) initiates and stimulates DNA strand resection by MRN complex to promote the alternative DNA repair pathways; (v) enables an efficient apoptotic signal if not removed; (vi) triggers relocation of the damaged DNA loci to the special nuclear compartments; (vii) provides precise PAR/MAR-guided recruitment and assembly of the DNA repair factors; and (viii) coordinates the choice of the DNA breaks signaling, processing and repair pathways.

[Insert Figure X.6 here]

Acknowledgments

This research was supported by grants to Murat Saparbaev from la Ligue National Contre le Cancer "Equipe Labellisee LIGUE 2016" and Electricité de France (RB 2017); to Alexander Ishchenko from French National Research Agency (ANR-18-CE44-0008) and Fondation ARC (PJA-20181208015); to Alexandra Kuznetsova the Russian Science Foundation grant No. 19-74-10034 (substrate specificity of AP endonucleases); to Bakhyt T. Matkarimov from Nazarbayev University Oak Ridge Associated Universities (ORAU) grant 091019CRP2111. E.M. and U.A. were supported by doctoral fellowships from ABELA FRERES and from MESR (Ministère de l'Enseignement Supérieur et de la Recherche), respectively.

624 References

- 625 1. T. Lindahl, *Nature*, 1993, **362**, 709.
- 626 2. J. A. Imlay and S. Linn, *Science*, 1988, **240**, 1302.
- 627 3. P. J. Brooks, *DNA Repair (Amst)*, 2008, **7**, 1168.
- 628 4. I. Kuraoka, C. Bender, A. Romieu, J. Cadet, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*, 2000,
- 629 **97**, 3832.
- 630 5. C. Marietta and P. J. Brooks, *EMBO Rep.*, 2007, **8**, 388.
- 631 6. P. J. Brooks, D. S. Wise, D. A. Berry, J. V. Kosmoski, *et al.*, *J. Biol. Chem.*, 2000, **275**,
- 632 22355.
- 633 7. M. D'Errico, E. Parlanti, M. Teson, B. M. de Jesus, *et al.*, *Embo J.*, 2006, **25**, 4305.
- 634 8. M. D'Errico, E. Parlanti, M. Teson, P. Degan, *et al.*, *Oncogene*, 2007, **26**, 4336.
- 635 9. G. Kirkali, N. C. de Souza-Pinto, P. Jaruga, V. A. Bohr, *et al.*, *DNA Repair (Amst)*, 2009,
- 636 **8**, 274.
- 637 10. J. Wang, C. L. Clauson, P. D. Robbins, L. J. Niedernhofer, *et al.*, *Aging Cell*, 2012, **11**,
- 638 714.
- 639 11. C. Chatgililoglu, C. Ferreri, N. E. Geacintov, M. G. Krokidis, *et al.*, *Cells*, 2019, **8**.
- 640 12. A. Luch, *Exs*, 2009, **99**, 151.
- 641 13. S. Amin, D. Desai, W. Dai, R. G. Harvey, *et al.*, *Carcinogenesis*, 1995, **16**, 2813.
- 642 14. S. Amin, J. Krzeminski, A. Rivenson, C. Kurtzke, *et al.*, *Carcinogenesis*, 1995, **16**, 1971.
- 643 15. A. K. Prahalad, J. A. Ross, G. B. Nelson, B. C. Roop, *et al.*, *Carcinogenesis*, 1997, **18**,
- 644 1955.
- 645 16. E. L. Cavalieri, S. Higginbotham, N. V. RamaKrishna, P. D. Devanesan, *et al.*,
- 646 *Carcinogenesis*, 1991, **12**, 1939.
- 647 17. S. M. Zhang, K. M. Chen, C. Aliaga, Y. W. Sun, *et al.*, *Chem. Res. Toxicol.*, 2011, **24**,
- 648 1297.
- 649 18. K. M. Li, M. George, M. L. Gross, C. H. Lin, *et al.*, *Chem. Res. Toxicol.*, 1999, **12**, 778.
- 650 19. K. M. Li, J. Byun, M. L. Gross, D. Zamzow, *et al.*, *Chem. Res. Toxicol.*, 1999, **12**, 749.
- 651 20. H. Yagi, H. Frank, A. Seidel and D. M. Jerina, *Chem. Res. Toxicol.*, 2008, **21**, 2379.
- 652 21. K. Kropachev, M. Kolbanovskiy, Z. Liu, Y. Cai, *et al.*, *Chem. Res. Toxicol.*, 2013, **26**,
- 653 783.
- 654 22. V. S. Sidorenko, J. E. Yeo, R. R. Bonala, F. Johnson, *et al.*, *Nucleic Acids Res.*, 2012, **40**,
- 655 2494.
- 656 23. M. Lukin, T. Zaliznyak, F. Johnson and C. de los Santos, *Nucleic Acids Res.*, 2012, **40**,
- 657 2759.
- 658 24. T. Zaliznyak, R. Bonala, F. Johnson and C. de Los Santos, *Chem. Res. Toxicol.*, 2006,
- 659 **19**, 745.
- 660 25. X. S. Cui, L. C. Eriksson and L. Moller, *Mutat. Res.*, 1999, **442**, 9.
- 661 26. C. A. Bieler, M. G. Cornelius, M. Stiborova, V. M. Arlt, *et al.*, *Carcinogenesis*, 2007, **28**,
- 662 1117.
- 663 27. H. Mao, Z. Deng, F. Wang, T. M. Harris, *et al.*, *Biochemistry*, 1998, **37**, 4374.
- 664 28. R. S. Cole, *Biochim. Biophys. Acta.*, 1971, **254**, 30.
- 665 29. B. H. Johnston and J. E. Hearst, *Biochemistry*, 1981, **20**, 739.
- 666 30. G. D. Cimino, H. B. Gamper, S. T. Isaacs and J. E. Hearst, *Annu. Rev. Biochem.*, 1985,
- 667 **54**, 1151.
- 668 31. K. W. Kohn, C. L. Spears and P. Doty, *J. Mol. Biol.*, 1966, **19**, 266.

- 669 32. F. Coste, J. M. Malinge, L. Serre, W. Shepard, *et al.*, *Nucleic Acids Res.*, 1999, **27**,
670 1837.
- 671 33. M. Tomasz, *Chem. Biol.*, 1995, **2**, 575.
- 672 34. I. D. Kozekov, L. V. Nechev, M. S. Moseley, C. M. Harris, *et al.*, *J. Am. Chem. Soc.*,
673 2003, **125**, 50.
- 674 35. S. Barker, M. Weinfeld and D. Murray, *Mutat. Res.*, 2005, **589**, 111.
- 675 36. J. Grillari, H. Katinger and R. Voglauer, *Nucleic Acids Res.*, 2007, **35**, 7566.
- 676 37. W. P. Vermeij, J. H. Hoeijmakers and J. Pothof, *Curr. Opin. Genet. Dev.*, 2014, **26C**,
677 124.
- 678 38. A. Sancar, L. A. Lindsey-Boltz, K. Unsal-Kacmaz and S. Linn, *Annu. Rev. Biochem.*,
679 2004, **73**, 39.
- 680 39. L. Cipak, N. Watanabe and T. Bessho, *Nat. Struct. Mol. Biol.*, 2006, **13**, 729.
- 681 40. H. E. Krokan and M. Bjoras, *Cold Spring Harb. Perspect. Biol.*, 2013, **5**, a012583.
- 682 41. A. A. Ischenko and M. K. Saparbaev, *Nature*, 2002, **415**, 183.
- 683 42. S. Klimasauskas, S. Kumar, R. J. Roberts and X. Cheng, *Cell*, 1994, **76**, 357.
- 684 43. J. T. Stivers, *Prog. Nucleic Acid Res. Mol. Biol.*, 2004, **77**, 37.
- 685 44. K. Hitomi, S. Iwai and J. A. Tainer, *DNA Repair (Amst)*, 2007, **6**, 410.
- 686 45. S. Boiteux, M. Bichara, R. P. Fuchs and J. Laval, *Carcinogenesis*, 1989, **10**, 1905.
- 687 46. D. G. Vassilyev, T. Kashiwagi, Y. Mikami, M. Ariyoshi, *et al.*, *Cell*, 1995, **83**, 773.
- 688 47. S. Couve-Privat, G. Mace, F. Rosselli and M. K. Saparbaev, *Nucleic Acids Res.*, 2007,
689 **35**, 5672.
- 690 48. V. Vartanian, I. G. Minko, S. Chawanthayatham, P. A. Egner, *et al.*, *Proc. Natl. Acad.*
691 *Sci. U. S. A.*, 2017, **114**, 4207.
- 692 49. F. Coste, M. Ober, Y. V. Le Bihan, M. A. Izquierdo, *et al.*, *Chem. Biol.*, 2008, **15**, 706.
- 693 50. P. L. McKibbin, A. M. Fleming, M. A. Towheed, B. Van Houten, *et al.*, *J. Am. Chem.*
694 *Soc.*, 2013, **135**, 13851.
- 695 51. E. A. Mullins, R. Shi, Z. D. Parsons, P. K. Yuen, *et al.*, *Nature*, 2015, **527**, 254.
- 696 52. E. A. Mullins, R. Shi, L. A. Kotsch and B. F. Eichman, *PLoS One*, 2015, **10**, e0127733.
- 697 53. E. A. Mullins, G. M. Warren, N. P. Bradley and B. F. Eichman, *Proc. Natl. Acad. Sci. U.*
698 *S. A.*, 2017, **114**, 4400.
- 699 54. I. Alseth, T. Rognes, T. Lindbäck, I. Solberg, *et al.*, *Mol Microbiol*, 2006, **59**, 1602.
- 700 55. R. Shi, E. A. Mullins, X. X. Shen, K. T. Lay, *et al.*, *Embo j*, 2018, **37**, 63.
- 701 56. E. A. Mullins, R. Shi and B. F. Eichman, *Nat. Chem. Biol.*, 2017, **13**, 1002.
- 702 57. H. Xu, W. Huang, Q. L. He, Z. X. Zhao, *et al.*, *Angew. Chem. Int. Ed. Engl.*, 2012, **51**,
703 10532.
- 704 58. E. A. Mullins, A. A. Rodriguez, N. P. Bradley and B. F. Eichman, *Trends Biochem. Sci.*,
705 2019, **44**, 765.
- 706 59. S. Wang, K. Liu, L. Xiao, L. Yang, *et al.*, *Nucleic Acids Res.*, 2016, **44**, 187.
- 707 60. S. Cheng, A. Sancar and J. E. Hearst, *Nucleic Acids Res.*, 1991, **19**, 657.
- 708 61. D. M. Noll, T. M. Mason and P. S. Miller, *Chem. Rev.*, 2006, **106**, 277.
- 709 62. Y. Huang and L. Li, *Transl. Cancer Res.*, 2013, **2**, 144.
- 710 63. D. Klein Douwel, R. A. Boonen, D. T. Long, A. A. Szypowska, *et al.*, *Mol. Cell*, 2014, **54**,
711 460.
- 712 64. M. R. Hodskinson, J. Silhan, G. P. Crossan, J. I. Garaycochea, *et al.*, *Mol. Cell*, 2014,
713 **54**, 472.
- 714 65. S. Couve, G. Mace-Aime, F. Rosselli and M. K. Saparbaev, *J. Biol. Chem.*, 2009, **284**,
715 11963.

- 716 66. M. Liu, S. Doublet and S. S. Wallace, *Mutat. Res.*, 2013, **743-744**, 4.
- 717 67. C. G. Neurauter, L. Luna and M. Bjoras, *DNA Repair (Amst)*, 2012, **11**, 401.
- 718 68. M. L. Hegde, P. M. Hegde, L. J. Bellot, S. M. Mandal, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, E3090.
- 719
- 720 69. M. Raschle, P. Knipsheer, M. Enoiu, T. Angelov, *et al.*, *Cell*, 2008, **134**, 969.
- 721 70. J. Zhang, J. M. Dewar, M. Budzowska, A. Motnenko, *et al.*, *Nat. Struct. Mol. Biol.*,
- 722 2015, **22**, 242.
- 723 71. J. Huang, S. Liu, M. A. Bellani, A. K. Thazhathveetil, *et al.*, *Mol. Cell*, 2013, **52**, 434.
- 724 72. J. Zhang and J. C. Walter, *DNA Repair (Amst)*, 2014, **19**, 135.
- 725 73. D. R. Semlow, J. Zhang, M. Budzowska, A. C. Drohat, *et al.*, *Cell*, 2016, **167**, 498.
- 726 74. P. R. Martin, S. Couve, C. Zutterling, M. S. Albelazi, *et al.*, *Sci. Rep.*, 2017, **7**, 17438.
- 727 75. V. Rolseth, S. Z. Krokeide, D. Kunke, C. G. Neurauter, *et al.*, *Biochim. Biophys. Acta*,
- 728 2013, **1833**, 1157.
- 729 76. A. Kothandapani, V. S. Dangeti, A. R. Brown, L. A. Banze, *et al.*, *J. Biol. Chem.*, 2011,
- 730 **286**, 14564.
- 731 77. A. Sawant, A. M. Floyd, M. Dangeti, W. Lei, *et al.*, *DNA Repair (Amst)*, 2017, **51**, 46.
- 732 78. A. Kothandapani and S. M. Patrick, *Mutat. Res.*, 2013, **743-744**, 44.
- 733 79. M. L. Dodson, M. L. Michaels and R. S. Lloyd, *J. Biol. Chem.*, 1994, **269**, 32709.
- 734 80. M. L. Dodson and R. S. Lloyd, *Free Radic. Biol. Med.*, 2002, **32**, 678.
- 735 81. L. Gros, M. K. Saparbaev and J. Laval, *Oncogene*, 2002, **21**, 8905.
- 736 82. H. E. Krokan, R. Standal and G. Slupphaug, *Biochem. J.*, 1997, **325 (Pt 1)**, 1.
- 737 83. A. L. Jacobs and P. Schar, *Chromosoma*, 2012, **121**, 1.
- 738 84. D. M. Wilson III and D. Barsky, *Mutat. Res.*, 2001, **485**, 283.
- 739 85. B. Dimple and J.-S. Sung, *DNA Repair (Amst)*, 2005, **4**, 1442.
- 740 86. H. Ide, K. Tedzuka, H. Shimzu, Y. Kimura, *et al.*, *Biochemistry*, 1994, **33**, 7842.
- 741 87. A. B. Guliaev, B. Hang and B. Singer, *Nucleic Acids Res.*, 2004, **32**, 2844.
- 742 88. S. Daviet, S. Couve-Privat, L. Gros, K. Shinozuka, *et al.*, *DNA Repair (Amst)*, 2007, **6**, 8.
- 743 89. A. A. Ishchenko, E. Deprez, A. Maksimenko, J. C. Brochon, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*,
- 744 2006, **103**, 2564.
- 745 90. C. D. Mol, C. F. Kuo, M. M. Thayer, R. P. Cunningham, *et al.*, *Nature*, 1995, **374**, 381.
- 746 91. D. J. Hosfield, Y. Guan, B. J. Haas, R. P. Cunningham, *et al.*, *Cell*, 1999, **98**, 397.
- 747 92. C. D. Mol, T. Izumi, S. Mitra and J. A. Tainer, *Nature*, 2000, **403**, 451.
- 748 93. B. Dimple and L. Harrison, *Annu. Rev. Biochem.*, 1994, **63**, 915.
- 749 94. G. Barzilay and I. D. Hickson, *Bioessays*, 1995, **17**, 713.
- 750 95. H. E. Krokan, H. Nilsen, F. Skorpen, M. Otterlei, *et al.*, *FEBS Lett.*, 2000, **476**, 73.
- 751 96. D. M. Wilson, 3rd, *J. Mol. Biol.*, 2003, **330**, 1027.
- 752 97. K. M. Chou and Y. C. Cheng, *Nature*, 2002, **415**, 655.
- 753 98. J. L. Parsons, Dianova, II and G. L. Dianov, *Nucleic Acids Res.*, 2005, **33**, 2204.
- 754 99. L. Gros, A. A. Ishchenko, H. Ide, R. H. Elder, *et al.*, *Nucleic Acids Res.*, 2004, **32**, 73.
- 755 100. P. Prorok, D. Alili, C. Saint-Pierre, D. Gasparutto, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*,
- 756 2013, **110**, E3695.
- 757 101. P. Burkovics, V. Szukacsov, I. Unk and L. Haracska, *Nucleic Acids Res.*, 2006, **34**, 2508.
- 758 102. B. Joldybayeva, P. Prorok, I. R. Grin, D. O. Zharkov, *et al.*, *PLoS One*, 2014, **9**, e92963.
- 759 103. M. Redrejo-Rodriguez, A. Vigouroux, A. Mursalimov, I. Grin, *et al.*, *Biochimie*, 2016,
- 760 **128-129**, 20.
- 761 104. S. Abeldenov, I. Talhaoui, D. O. Zharkov, A. A. Ishchenko, *et al.*, *DNA Repair (Amst)*,
- 762 2015, **33**, 1.

- 763 105. M. Shiraishi, S. Ishino, T. Yamagami, Y. Egashira, *et al.*, *Nucleic Acids Res.*, 2015, **43**,
764 2853.
- 765 106. M. Shiraishi and S. Iwai, *J. Bacteriol.*, 2020, **202**.
- 766 107. M. Shiraishi, S. Ishino, I. Cann and Y. Ishino, *Biosci. Biotechnol. Biochem.*, 2017, **81**,
767 931.
- 768 108. C. D. Mol, D. J. Hosfield and J. A. Tainer, *Mutat. Res.*, 2000, **460**, 211.
- 769 109. J. P. Erzberger and D. M. Wilson, 3rd, *J. Mol. Biol.*, 1999, **290**, 447.
- 770 110. B. D. Freudenthal, W. A. Beard, M. J. Cuneo, N. S. Dyrkheeva, *et al.*, *Nat. Struct. Mol.*
771 *Biol.*, 2015, **22**, 924.
- 772 111. P. T. Beernink, B. W. Segelke, M. Z. Hadi, J. P. Erzberger, *et al.*, *J. Mol. Biol.*, 2001,
773 **307**, 1023.
- 774 112. E. D. Garcin, D. J. Hosfield, S. A. Desai, B. J. Haas, *et al.*, *Nat. Struct. Mol. Biol.*, 2008,
775 **15**, 515.
- 776 113. I. Ivanov, J. A. Tainer and J. A. McCammon, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**,
777 1465.
- 778 114. S. E. Tsutakawa, D. S. Shin, C. D. Mol, T. Izumi, *et al.*, *J. Biol. Chem.*, 2013, **288**, 8445.
- 779 115. A. Mazouzi, A. Vigouroux, B. Aikeshv, P. J. Brooks, *et al.*, *Proc. Natl. Acad. Sci. U. S.*
780 *A.*, 2013, **110**, E3071.
- 781 116. S. T. Mundle, M. H. Fattal, L. F. Melo, J. D. Coriolan, *et al.*, *DNA Repair (Amst)*, 2004,
782 **3**, 1447.
- 783 117. S. T. Mundle, J. C. Delaney, J. M. Essigmann and P. R. Strauss, *Biochemistry*, 2009, **48**,
784 19.
- 785 118. M. M. Aboelnga and S. D. Wetmore, *J. Am. Chem. Soc.*, 2019, **141**, 8646.
- 786 119. I. V. Alekseeva, A. S. Bakman, Y. N. Vorobjev, O. S. Fedorova, *et al.*, *J. Phys. Chem. B*,
787 2019, **123**, 9546.
- 788 120. W. C. Kim, B. R. Berquist, M. Chohan, C. Uy, *et al.*, *J. Mol. Biol.*, 2011, **411**, 960.
- 789 121. D. M. Wilson, 3rd, M. Takeshita, A. P. Grollman and B. Demple, *J. Biol. Chem.*, 1995,
790 **270**, 16002.
- 791 122. G. Golan, A. A. Ishchenko, B. Khassenov, G. Shoham, *et al.*, *Mutat. Res.*, 2010, **685**,
792 70.
- 793 123. A. Gelin, M. Redrejo-Rodriguez, J. Laval, O. S. Fedorova, *et al.*, *PLoS One*, 2010, **5**,
794 e12241.
- 795 124. J. Willis, Y. Patel, B. L. Lentz and S. Yan, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**,
796 10592.
- 797 125. Y. Lin, L. Bai, S. Cupello, M. A. Hossain, *et al.*, *Nucleic Acids Res.*, 2018, **46**, 2479.
- 798 126. Y. Lin, J. Raj, J. Li, A. Ha, *et al.*, *Nucleic Acids Res.*, 2019, doi: 10.1093/nar/gkz1175.
- 799 127. J. R. Lakowicz, *Principles of Fluorescence Spectroscopy*, Springer, Berlin., 2006.
- 800 128. L. Y. Kanazhevskaya, V. V. Koval, D. O. Zharkov, P. R. Strauss, *et al.*, *Biochemistry*,
801 2010, **49**, 6451.
- 802 129. N. A. Timofeyeva, V. V. Koval, D. G. Knorre, D. O. Zharkov, *et al.*, *J. Biomol. Struct.*
803 *Dyn.*, 2009, **26**, 637.
- 804 130. N. A. Timofeyeva, V. V. Koval, A. A. Ishchenko, M. K. Saparbaev, *et al.*, *PLoS One*,
805 2011, **6**, e24063.
- 806 131. N. A. Timofeyeva, V. V. Koval, A. A. Ishchenko, M. K. Saparbaev, *et al.*, *Biochemistry*
807 *(Mosc)*, 2011, **76**, 273.
- 808 132. A. D. Miroshnikova, A. A. Kuznetsova, N. A. Kuznetsov and O. S. Fedorova, *Acta*
809 *Naturae*, 2016, **8**, 103.

- 810 133. R. W. Sinkeldam, N. J. Greco and Y. Tor, *Chem. Rev.*, 2010, **110**, 2579.
- 811 134. L. M. Wilhelmsson, *Q. Rev. Biophys.*, 2010, **43**, 159.
- 812 135. N. A. Timofeyeva and O. S. Fedorova, *Mol. Biosyst.*, 2016, **12**, 3435.
- 813 136. L. Y. Kanazhevskaya, V. V. Koval, Y. N. Vorobjev and O. S. Fedorova, *Biochemistry*,
814 2012, **51**, 1306.
- 815 137. L. Y. Kanazhevskaya, V. V. Koval, A. A. Lomzov and O. S. Fedorova, *DNA Repair*, 2014,
816 **21**, 43.
- 817 138. I. V. Alekseeva, A. T. Davletgildeeva, O. V. Arkova, N. A. Kuznetsov, *et al.*, *Biochimie*,
818 2019, **163**, 73.
- 819 139. E. S. Dyakonova, V. V. Koval, A. A. Lomzov, A. A. Ishchenko, *et al.*, *Biochimie*, 2018,
820 **152**, 53.
- 821 140. E. S. Dyakonova, V. V. Koval, A. A. Ishchenko, M. K. Saparbaev, *et al.*, *Biochemistry*
822 *(Mosc)*, 2012, **77**, 1162.
- 823 141. A. D. Miroshnikova, A. A. Kuznetsova, Y. N. Vorobjev, N. A. Kuznetsov, *et al.*, *Mol.*
824 *Biosyst.*, 2016, **12**, 1527.
- 825 142. S. C. Brooks, S. Adhikary, E. H. Robinson and B. F. Eichman, *Biochim. Biophys. Acta*,
826 2013, **1834**, 247.
- 827 143. N. A. Kuznetsov and O. S. Fedorova, *Biochemistry (Mosc)*, 2016, **81**, 1136.
- 828 144. V. M. Marathias, B. Jerkovic and P. H. Bolton, *Nucleic Acids Res.*, 1999, **27**, 1854.
- 829 145. A. R. Dunn, N. M. Kad, S. R. Nelson, D. M. Warshaw, *et al.*, *Nucleic Acids Res.*, 2011,
830 **39**, 7487.
- 831 146. A. Prakash, S. Doublie and S. S. Wallace, *Prog. Mol. Biol. Transl. Sci.*, 2012, **110**, 71.
- 832 147. S. R. Nelson, A. R. Dunn, S. D. Kathe, D. M. Warshaw, *et al.*, *Proc. Natl. Acad. Sci. U. S.*
833 *A.*, 2014, **111**, E2091.
- 834 148. V. V. Koval, N. A. Kuznetsov, A. A. Ishchenko, M. K. Saparbaev, *et al.*, *Mutat. Res.*,
835 2010, **685**, 3.
- 836 149. N. A. Kuznetsov, O. A. Kladova, A. A. Kuznetsova, A. A. Ishchenko, *et al.*, *J. Biol.*
837 *Chem.*, 2015, **290**, 14338.
- 838 150. N. A. Kuznetsov, C. Bergonzo, A. J. Campbell, H. Li, *et al.*, *Nucleic Acids Res.*, 2015, **43**,
839 272.
- 840 151. A. J. Lee and S. S. Wallace, *Radiat. Phys. Chem. Oxf. Engl. 1993*, 2016, **128**, 126.
- 841 152. A. J. Lee and S. S. Wallace, *Free Radic. Biol. Med.*, 2017, **107**, 170.
- 842 153. O. A. Kladova, A. A. Kuznetsova, O. S. Fedorova and N. A. Kuznetsov, *Genes (Basel)*,
843 2017, **8**, 1.
- 844 154. A. A. Kuznetsova, D. A. Iakovlev, I. V. Misovets, A. A. Ishchenko, *et al.*, *Mol. Biosyst.*,
845 2017, **13**, 2638.
- 846 155. A. A. Kuznetsova, A. G. Matveeva, A. D. Milov, Y. N. Vorobjev, *et al.*, *Nucleic Acids*
847 *Res.*, 2018, **46**, 11454.
- 848 156. N. A. Kuznetsov and O. S. Fedorova, *Adv. Exp. Biol. Med.*, 2020, in press.
- 849 157. C. Blanpain, M. Mohrin, P. A. Sotiropoulou and E. Passegue, *Cell Stem Cell*, 2011, **8**,
850 16.
- 851 158. A. Burkle, *FEBS J.*, 2005, **272**, 4576.
- 852 159. I. Talhaoui, N. A. Lebedeva, G. Zarkovic, C. Saint-Pierre, *et al.*, *Nucleic Acids Res.*,
853 2016, **44**, 9279.
- 854 160. E. A. Belousova, C. Ishchenko capital A and O. I. Lavrik, *Sci. Rep.*, 2018, **8**, 4176.
- 855 161. D. Munnur, E. Bartlett, P. Mikolcevic, I. T. Kirby, *et al.*, *Nucleic Acids Res.*, 2019, **47**,
856 5658.

- 857 162. C. Beck, I. Robert, B. Reina-San-Martin, V. Schreiber, *et al.*, *Exp. Cell. Res.*, 2014, **329**,
858 18.
- 859 163. M. O. Hottiger, *Annu. Rev. Biochem.*, 2015, **84**, 227.
- 860 164. M. Posavec Marjanovic, K. Crawford and I. Ahel, *Semin. Cell Dev. Biol.*, 2017, **63**, 102.
- 861 165. J. M. Rodriguez-Vargas, F. J. Oliver-Pozo and F. Dantzer, *Oxid. Med. Cell Longev.*,
862 2019, **2019**, 2641712.
- 863 166. R. Gupte, Z. Liu and W. L. Kraus, *Genes Dev.*, 2017, **31**, 101.
- 864 167. L. Palazzo, O. Leidecker, E. Prokhorova, H. Dauben, *et al.*, *Elife*, 2018, **7**.
- 865 168. M. O. Hottiger, P. O. Hassa, B. Luscher, H. Schuler, *et al.*, *Trends Biochem. Sci.*, 2010,
866 **35**, 208.
- 867 169. T. Minaga and E. Kun, *J. Biol. Chem.*, 1983, **258**, 5726.
- 868 170. R. Alvarez-Gonzalez and F. R. Althaus, *Mutat. Res.*, 1989, **218**, 67.
- 869 171. L. Davidovic, M. Vodenicharov, E. B. Affar and G. G. Poirier, *Exp. Cell. Res.*, 2001, **268**,
870 7.
- 871 172. B. A. Gibson and W. L. Kraus, *Nat. Rev. Mol. Cell Biol.*, 2012, **13**, 411.
- 872 173. F. G. Sousa, R. Matuo, D. G. Soares, A. E. Escargueil, *et al.*, *Carcinogenesis*, 2012, **33**,
873 1433.
- 874 174. S. Eustermann, W. F. Wu, M. F. Langelier, J. C. Yang, *et al.*, *Mol. Cell*, 2015, **60**, 742.
- 875 175. M. F. Langelier, J. L. Planck, S. Roy and J. M. Pascal, *Science*, 2012, **336**, 728.
- 876 176. J. M. Dawicki-McKenna, M. F. Langelier, J. E. DeNizio, A. A. Riccio, *et al.*, *Mol. Cell*,
877 2015, **60**, 755.
- 878 177. J. D. Steffen, M. M. McCauley and J. M. Pascal, *Nucleic Acids Res.*, 2016, **44**, 9771.
- 879 178. W. M. Shieh, J. C. Ame, M. V. Wilson, Z. Q. Wang, *et al.*, *J. Biol. Chem.*, 1998, **273**,
880 30069.
- 881 179. K. Martin-Hernandez, J. M. Rodriguez-Vargas, V. Schreiber and F. Dantzer, *Semin. Cell*
882 *Dev. Biol.*, 2017, **63**, 92.
- 883 180. J. Menissier de Murcia, M. Ricoul, L. Tartier, C. Niedergang, *et al.*, *Embo J.*, 2003, **22**,
884 2255.
- 885 181. G. J. Grundy, L. M. Polo, Z. Zeng, S. L. Rulten, *et al.*, *Nat. Commun.*, 2016, **7**, 12404.
- 886 182. C. Boehler, L. R. Gauthier, O. Mortusewicz, D. S. Biard, *et al.*, *Proc. Natl. Acad. Sci. U.*
887 *S. A.*, 2011, **108**, 2783.
- 888 183. C. Beck, C. Boehler, J. Guirouilh Barbat, M. E. Bonnet, *et al.*, *Nucleic Acids Res.*, 2014,
889 **42**, 5616.
- 890 184. H. Shirai, A. R. Poetsch, A. Gunji, D. Maeda, *et al.*, *Cell Death Dis.*, 2013, **4**, e656.
- 891 185. J. M. Hinz, Y. Rodriguez and M. J. Smerdon, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**,
892 4646.
- 893 186. I. D. Odell, J. E. Barbour, D. L. Murphy, J. A. Della-Maria, *et al.*, *Mol. Cell. Biol.*, 2011,
894 **31**, 4623.
- 895 187. J. Abplanalp and M. O. Hottiger, *Semin. Cell Dev. Biol.*, 2017, **63**, 114.
- 896 188. F. Ciccarone, M. Zampieri and P. Caiafa, *Semin. Cell Dev. Biol.*, 2017, **63**, 123.
- 897 189. Y. Tanaka, K. Yoshihara, A. Itaya, T. Kamiya, *et al.*, *J. Biol. Chem.*, 1984, **259**, 6579.
- 898 190. M. S. Satoh, G. G. Poirier and T. Lindahl, *Biochemistry*, 1994, **33**, 7099.
- 899 191. S. W. Yu, H. Wang, M. F. Poitras, C. Coombs, *et al.*, *Science*, 2002, **297**, 259.
- 900 192. G. Zarkovic, E. A. Belousova, I. Talhaoui, C. Saint-Pierre, *et al.*, *Nucleic Acids Res.*,
901 2018, **46**, 2417.
- 902 193. D. Munnur and I. Ahel, *Febs J.*, 2017, **284**, 4002.
- 903 194. A. A. Riccio, G. Cingolani and J. M. Pascal, *Nucleic Acids Res.*, 2016, **44**, 1691.

- 904 195. R. Anand, L. Ranjha, E. Cannavo and P. Cejka, *Mol. Cell*, 2016, **64**, 940.
- 905 196. H. E. Bryant, E. Petermann, N. Schultz, A. S. Jemth, *et al.*, *Embo J.*, 2009, **28**, 2601.
- 906 197. I. Lonskaya, V. N. Potaman, L. S. Shlyakhtenko, E. A. Oussatcheva, *et al.*, *J. Biol.*
907 *Chem.*, 2005, **280**, 17076.
- 908 198. A. Hakme, H. K. Wong, F. Dantzer and V. Schreiber, *EMBO Rep.*, 2008, **9**, 1094.
- 909 199. P. Bai, *Mol. Cell*, 2015, **58**, 947.
- 910 200. X. Luo and W. L. Kraus, *Genes Dev.*, 2012, **26**, 417.
- 911 201. K. Leger, D. Bar, N. Savic, R. Santoro, *et al.*, *Nucleic Acids Res.*, 2014, **42**, 5072.
- 912 202. K. Ouararhni, R. Hadj-Slimane, S. Ait-Si-Ali, P. Robin, *et al.*, *Genes Dev.*, 2006, **20**,
913 3324.
- 914 203. E. Fouquerel, E. M. Goellner, Z. Yu, J. P. Gagne, *et al.*, *Cell Rep.*, 2014, **8**, 1819.
- 915 204. M. Y. Kim, S. Mauro, N. Gevry, J. T. Lis, *et al.*, *Cell*, 2004, **119**, 803.
- 916 205. M. F. Langelier, A. A. Riccio and J. M. Pascal, *Nucleic Acids Res.*, 2014, **42**, 7762.
- 917 206. M. Audebert, B. Salles and P. Calsou, *J. Biol. Chem.*, 2004, **279**, 55117.
- 918 207. D. S. Kim, C. V. Camacho, A. Nagari, V. S. Malladi, *et al.*, *Mol. Cell*, 2019, **75**, 1270.
- 919 208. Q. Chen, M. A. Kassab, F. Dantzer and X. Yu, *Nat. Commun.*, 2018, **9**, 3233.
- 920 209. N. J. Clark, M. Kramer, U. M. Muthurajan and K. Luger, *J. Biol. Chem.*, 2012, **287**,
921 32430.
- 922 210. C. Thomas, Y. Ji, C. Wu, H. Datz, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*, 2019, **116**, 9941.
- 923 211. M. Cohen-Armon, L. Visochek, D. Rozensal, A. Kalal, *et al.*, *Mol. Cell*, 2007, **25**, 297.
- 924 212. F. Berger, C. Lau and M. Ziegler, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 3765.
- 925 213. T. Guastafierro, B. Cecchinelli, M. Zampieri, A. Reale, *et al.*, *J. Biol. Chem.*, 2008, **283**,
926 21873.
- 927 214. F. Ciccarone, E. Valentini, M. Zampieri and P. Caiafa, *Oncotarget*, 2015, **6**, 24333.
- 928 215. S. Erener, M. Hesse, R. Kostadinova and M. O. Hottiger, *Mol. Endocrinol.*, 2012, **26**,
929 79.
- 930 216. C. Guetg and R. Santoro, *Epigenetics*, 2012, **7**, 811.
- 931 217. M. Sajish and P. Schimmel, *Nature*, 2015, **519**, 370.
- 932 218. T. M. Kauppinen, W. Y. Chan, S. W. Suh, A. K. Wiggins, *et al.*, *Proc. Natl. Acad. Sci. U.*
933 *S. A.*, 2006, **103**, 7136.
- 934 219. S. Zhang, Y. Lin, Y. S. Kim, M. P. Hande, *et al.*, *Cell Death Differ.*, 2007, **14**, 1001.
- 935 220. R. H. Wright, G. Castellano, J. Bonet, F. Le Dily, *et al.*, *Genes Dev.*, 2012, **26**, 1972.
- 936 221. P. O. Hassa, S. S. Haenni, C. Buerki, N. I. Meier, *et al.*, *J. Biol. Chem.*, 2005, **280**,
937 40450.
- 938 222. S. B. Rajamohan, V. B. Pillai, M. Gupta, N. R. Sundaresan, *et al.*, *Mol. Cell. Biol.*, 2009,
939 **29**, 4116.
- 940 223. O. Loseva, A. S. Jemth, H. E. Bryant, H. Schuler, *et al.*, *J. Biol. Chem.*, 2010, **285**, 8054.
- 941 224. M. Van Meter, Z. Mao, V. Gorbunova and A. Seluanov, *Cell Cycle*, 2011, **10**, 3153.
- 942 225. P. C. Hsu, R. K. Gopinath, Y. A. Hsueh and S. Y. Shieh, *Oncogene*, 2019, **38**, 1166.
- 943 226. I. Kassner, A. Andersson, M. Fey, M. Tomas, *et al.*, *Open Biol.*, 2013, **3**, 120173.

944

FIGURE AND TABLE CAPTIONS

Figure 1. Schematic presentation of singular and complex DNA damage. (A) Chemical structures of some singular non-bulky DNA base lesions. (B) Chemical structures of complex DNA lesions including cisplatin-ICL and bulky repair-resistant DNA base lesions.

Figure 2. DNA replication-coupled repair of inter-strand DNA crosslinks. (A) NEIL1 excises unhooked crosslinked oligomer in three-stranded DNA repair intermediate. (B) NEIL3- and (C) Fanconi anemia-mediated repair of ICL. (D) The hypothetical mechanism of the FANCM-mediated DNA replication fork bypass of ICL and NEILs-mediated repair of three- and four stranded crosslinked DNA structures.

Figure 3. The scheme of BER and NIR pathways.

Figure 4. Mechanisms of DNA damage recognition by DNA glycosylases^{90,104} (A) and AP endonucleases^{78,102} (B).

Figure 5. The mechanistic model of DNA strand break-oriented DNA termini ADP-ribosylation by PARP1, PARP2 and PARP3.

Figure 6. Putative and confirmed roles of the PARPs-catalyzed ADP-ribosylation in response to DNA damage.

Table 1. Mediators of PARP1-3 activation