



HAL
open science

Subchronic exposure to high-density polyethylene microplastics alone or in combination with chlortoluron significantly affected valve activity and daily growth of the Pacific oyster, *Crassostrea gigas*

Arno Bringer, H el ene Thomas, Emmanuel Dubillot, St ephane Le Floch,
Justine Receveur, J er ome Cachot, Damien Tran

► To cite this version:

Arno Bringer, H el ene Thomas, Emmanuel Dubillot, St ephane Le Floch, Justine Receveur, et al.. Subchronic exposure to high-density polyethylene microplastics alone or in combination with chlortoluron significantly affected valve activity and daily growth of the Pacific oyster, *Crassostrea gigas*. *Aquatic Toxicology*, 2021, 237, pp.105880. 10.1016/j.aquatox.2021.105880 . hal-03434008

HAL Id: hal-03434008

<https://hal.science/hal-03434008v1>

Submitted on 23 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 Subchronic exposure to high-density polyethylene microplastics alone or in combination
2 with chlortoluron significantly affected valve activity and daily growth of *Crassostrea gigas*.

3

4 Arno Bringer^{1*}, H  l  ne Thomas¹⁺, Emmanuel Dubillot¹, St  phane Le Floch³, Justine Receveur³, J  r  me
5 Cachot²⁺, Damien Tran²

6

7 ¹ *Littoral Environnement et Soci  t  s (LIENSs), UMR 7266, CNRS-Universit   de La Rochelle, 2 rue Olympe de*
8 *Gouges, F-17042 La Rochelle Cedex 01, France.*

9 ² *Universit   de Bordeaux, CNRS, EPOC, EPHE, UMR 5805, F-33600 Pessac, France.*

10 ³ *Cedre, 715 Rue Alain Colas, 29218 Brest Cedex 2, France.*

11 ⁺: *equal contributions*

12 ^{*}: *Corresponding author: Arno Bringer: Tel: +33 645299739 (FR), Email: arno.bringer1@univ-lr.fr*

13


14 **Abstract**

15 Nowadays, pesticides and microplastics (MPs) are commonly found in coastal waters worldwide. Due
16 to their widespread use, their persistence and toxicity, they may induce adverse effects on
17 physiology and behaviour of marine organisms such as the Pacific oyster (*Crassostrea gigas*). This
18 study explored the growth and valve activity of juvenile oysters exposed for 24 days to two
19 frequently detected pollutants in the Pertuis Charentais (South-West France): an herbicide
20 (chlortoluron, 30 $\mu\text{g}\cdot\text{L}^{-1}$) and high-density polyethylene microparticles (HDPE 20-25 μm , 10 $\mu\text{g MP}\cdot\text{L}^{-1}$)
21 alone or in combinaison. The characterisation of the valve activity of juvenile oysters, recorded by
22 using a High Frequency and Non-Invasive valvometer (HFNI) depended on three parameters: the
23 number of valve micro-closures (VMC), the Valve Opening Amplitude (VOA) and the Valve Opening
24 Duration (VOD). Additionally, daily shell growth and the oyster daily rhythm were assessed. The
25 exposure to MPs of oysters led to a significant, increase of VMC and a decrease of VOD and shell
26 growth. The exposure to chlortoluron showed a significant, increase of VOA and a decrease of VMC.

27 In mixture with MPs, chlortoluron still increased VOA and decreased VMC but also reduced the shell
28 growth. Chronobiological analysis did not reveal any effects on the daily rhythm of both
29 contaminants. High experimental concentrations of MP and chlortoluron were tested. This work has
30 underlined their selective impacts on the behavioral and physiological parameters studied.

31

32 Graphical abstract



33

34

35 **Keywords** Bivalve molluscs, Experimental biomonitoring, Microplastics, Pesticide, Valve behaviour,
36 Growth.

37

38 1. Introduction

39 With a production of 135,000 tons of bivalves in 2017, including 81,000 tons of oysters, France, is the
40 2nd most important producer of bivalves in Europe (France Agrimer, 2020). Shellfish farming turnover
41 was estimated at 774 million euros in 2019 (CNC, 2020). The Pertuis Charentais area (South-West,
42 Atlantic coast, France) represents 23 % of the national oyster production (CNC, 2016). Since 2010,
43 juvenile and adult oysters are facing excess mortality rates, reaching 70 to 80 % of oyster farms
44 located in the Atlantic coast (Soletchnik *et al.*, 2018 ; Lucasson, 2018 and Girard & Agundez, 2014).
45 The quality of the coastal waters, where bivalves are grown, is a source of concern today. The
46 increased use of plastics (Phuong, 2018) in aquaculture generates MPs pollution, which could affect
47 physiology of bivalves (*e.g.* growth, immune status, reproduction) (Sussarellu *et al.*, 2016; Paul-Pont
48 *et al.*, 2018; Gardon *et al.*, 2020). Scientific studies are working to offer more and more toxicological
49 tests on emerging contaminants such as MPs and pesticides, in order to better understand the
50 modes of action of these pollutants (Wang *et al.*, 2020; Bhagat *et al.*, 2020).

51 Plastics are widely used polymers, which came as a revolution in the past decades (Xu *et al.*, 2017). In
52 2019, nearly 370 million tons of plastics were produced in the world and in Europe, plastics
53 production almost reached 58 million tons (PlasticsEurope, 2020). In Europe, 80 % of the plastic
54 production resorts to six main polymers (PlasticsEurope, 2016): polypropylene (PP), high and low-
55 density polyethylene (HDPE and LDPE), polyvinyl chloride (PVC), polyurethane (PUR), polyethylene
56 terephthalate (PET) and polystyrene (PS). It is estimated that rivers carry between 70 % and 80 % of
57 plastic waste, most of which ends up in oceans (Horton *et al.*, 2017). Through wave activities, UV
58 degradation and physical abrasion, larger plastic pieces in the marine environment are fragmented,
59 resulting in producing MPs and nanoplastics (NPs) (Born & Schüttrumpf, 2019; Díaz-Mendoza *et al.*,
60 2020). MPs are plastic particles ranging from 1 µm to 5 mm (Cole *et al.*, 2011; Wagner *et al.*, 2014).
61 MPs ingestion was observed in numerous aquatic species, notably in filter-feeding bivalves at
62 different life stages (Rist *et al.*, 2019; Ward *et al.*, 2019; Betsill *et al.*, 2019; Bringer *et al.*, 2020b). It
63 can cause several harmful effects at the physiological and tissue levels, affecting the growth and
64 reproduction of bivalves (Lee *et al.*, 2013; Sussarellu *et al.*, 2016; Lo and Chan, 2018), as well as

65 physical injuries (Gall & Thompson, 2015) and reduced eating behavior (Cole *et al.*, 2015). At the
66 individual level, this can lead to the depletion of energy reserves (Wright *et al.*, 2013 and Xu *et al.*,
67 2017), larval development leading to delays and malformations (Tallec *et al.*, 2018), and swimming
68 activity (Bringer *et al.*, 2020a). Toxic effects are mainly observed using high laboratory exposures
69 (and not *in situ*) compared to environmental concentrations reported in the literature (Paul-Pont *et*
70 *al.*, 2018). The environmental concentrations mainly consider size classes of MP > 333 μm (Song *et*
71 *al.*, 2015; Zhao *et al.*, 2014; Baldwin *et al.*, 2016) and therefore contain a significant bias, with respect
72 to the small sizes used in the laboratory. Environmental concentrations ranging from 0.002 to 0.5 mg
73 $\text{MP} \cdot \text{L}^{-1}$ have been detected in surface water in different areas of the world (Eriksen *et al.*, 2013;
74 Enders *et al.*, 2015).

75 Chlortoluron that belongs to substituted urea family is one of the most commonly used herbicides in
76 wheat and barley crops. Chlortoluron inhibits photosynthesis and affects phytoplankton (Valiente
77 Moro *et al.*, 2012). In 2017, a maximum value of 0.18 $\mu\text{g} \cdot \text{L}^{-1}$ was detected at the mouth of a river
78 (surface water) flowing into the Pertuis Charentais (Leonard, 2017). A concentration of 0.14 $\mu\text{g} \cdot \text{L}^{-1}$ of
79 chlortoluron was detected in 2018, at the mouth of the Charente (Pertuis Charentais), close to oyster
80 farms (Action program, 2018). The herbicide family has already been studied on oysters, showing
81 effects at the early stages of development on survival and development (Mottier *et al.*, 2013; Gamain
82 *et al.*, 2017; Behrens *et al.*, 2016; Mai *et al.*, 2013), during gametogenesis with genotoxic effects
83 (Akcha *et al.*, 2020), as well as by modulating molecular and biochemical parameters (Lee *et al.*,
84 2017).

85 The valve activity on the oyster was measured using an HFNI (High Frequency Non-Invasive)
86 valvometer biosensor (Andrade *et al.*, 2016). This biosensor has made possible to monitor
87 behavioural responses of bivalves when exposed to stress events, such as, contaminants. Assessing
88 the valve activity behaviour through an HFNI valvometer has already been used to study
89 contaminants, including traces of metals (Tran *et al.*, 2004a, 2007), radionuclides (Tran *et al.*, 2004b,
90 2005) and harmful algae (Tran *et al.*, 2010, 2015; Haberkorn *et al.*, 2011). To our knowledge, no

91 valvometric behavioral study, under laboratory conditions, has been carried out to assess the
92 consequences of exposure to pesticides and MPs. However, in recent years, several articles have
93 underlined the relevance of modelling and analysing the behaviour of marine animals under
94 experimental conditions mimicking environmental conditions (Mat *et al.*, 2014; Ahmed *et al.*, 2015).
95 In the present study, under controlled conditions and to characterize the behaviour of the oyster
96 valve, three parameters were recorded: VMC, VOA and VOD. In addition, a chronobiological analysis
97 of the daily rhythm of the valve was carried out as well as a measurement of the daily growth. The
98 aim of this study was to better understand the potentially disrupting effects of emerging
99 contaminants (MPs and pesticides) alone or in cocktail on the daily growth and the valve activity of
100 Pacific juveniles oysters (*C. gigas*).

101

102 **2. Material and Methods**

103 *2.1. Individuals and acclimatization*

104

105 Pacific oysters, *Crassostrea gigas* (Bayne *et al.*, 2019; Bayne *et al.*, 2017), were kindly provided by a
106 hatchery (™France Naissain, France). This experiment was conducted on 12-month-old juvenile
107 oysters (59.5 ± 1.1 mm, 22.6 ± 1 g total mass) during autumn season (after the period of
108 gametogenesis and fertilization). Four homogenous groups of diploid oysters ($n = 8/\text{condition}$) were
109 chosen. Each group was distributed in a 37.5 L tank. Each tank was continuously filled with
110 oxygenated seawater coming from a large buffer tank. For the experiment, oysters were acclimated
111 for 7 days in UV-treated running ($78.3 \text{ L}\cdot\text{h}^{-1}$) seawater before being exposed to pollutants for 24 days.
112 Seawater was taken from the Pertuis Breton (South-West, France), which is directly connected to the
113 experimental area, and filtered using a 50 μm and 10 μm membrane in order to eliminate debris.
114 Filtered seawater (FSW) was used for experimental exposure tanks. The FSW temperature in tanks
115 was of 16.8 ± 0.4 °C and the salinity of 35.7 ± 0.2 ‰. The temperature value and salinity were daily
116 measured. Throughout the experiment, nitrates, nitrites, chlorine and pH (7.3 ± 1.0) were regularly

117 measured to ensure consistency. A continuous bubbling and a brewing system were set up.
118 Throughout the duration of the experiment, a natural photoperiod was established. In addition, the
119 oysters were nourished thanks to the natural presence of phytoplankton in the seawater used.
120 During the acclimatization and experimental periods, no mortality events were recorded.
121 Nevertheless, failures of the valvometric signal recording, due to an electrical leak at electrodes level
122 during the study, reduced the number of oysters studied to respectively $n = 7$ in the control, $n = 6$ in
123 the chlortoluron, $n = 8$ in the MP and $n = 5$ under cocktail conditions.

124

125 *2.2 Experimental procedure*

126

127 MPs were used ([™]HDPE 20-25 μm MPP-1241, density 0.96, Micropowders Inc. USA) and chlortoluron
128 ([™]Pestanal, analytical standard, 45400-250MG-R reference) was purchased from Sigma-Aldrich
129 Chemical (St. Quentin Fallavier, France). The solutions containing MPs and chlortoluron were
130 prepared beforehand in test glass tubes. Stock solutions, prepared in pure Mili-Q-water, showed a
131 MPs concentration of 10 mg MP.L^{-1} and a chlortoluron concentration of 3 mg.L^{-1} . They were kept cool
132 in the dark in order to prevent bacterial growth. The stored solutions were then diluted in FSW to
133 lower the concentrations to $10 \mu\text{g MP.L}^{-1}$ and $30 \mu\text{g.L}^{-1}$ respectively. After the 7-day acclimatization
134 period, the pollutants are added to the experimental system, in the buffer tanks. Every two days, 30
135 % of the FSW in the buffer tank was renewed, for the well-health of the oysters. Upstream, FSW
136 were stored in separate tanks, with the tested pollutants, at a temperature similar to that of tanks
137 containing oysters. They were used to supply the buffer tanks during the renewal of FSW. Water
138 stirrers have been installed in each buffer tanks in order to homogenize the plastic microparticles as
139 much as possible in the seawater. In addition, the cytometric analyzes made it possible to determine
140 a certain homogeneity in the exposure tanks. Polystyrene foams protected tanks and equipment to
141 reduce vibrations from electrical equipment. In addition, capacitance chambers were installed to

142 reduce vibrations from the air pipes bringing oxygen into FSW, along with behavioral analyzes (Fig.
143 S1).

144

145 *2.3 HFNI valvometry analysis*

146

147 To assess the pollutant impact (MPs and chlortoluron) on the oyster's valve behaviour, the valve
148 activity was recorded using a High Frequency Non-Invasive (HFNI) valvometer. The oysters were
149 equipped with light electromagnetic electrodes coated in resin (≈ 1 g), which were glued at the same
150 spot on both shells. The electrodes were attached to free covered stranded copper wires (diameter:
151 0.98 mm, length: 120 cm), sending data to a device. The induced voltage, measured by biosensors,
152 varied according to the distance between the electromagnetic electrodes. For each oyster, a
153 measure was taken every 4.8 sec., *i.e.* 18 000 data/oyster/day. For more details, see Tran *et al.*
154 (2015) and Andrade *et al.* (2016). Bivalve behaviour was monitored 24/24 hours and 7/7 days (Tran
155 *et al.*, 2004a and Haberkorn *et al.*, 2011). The biosensor used consists of very sensitive electrodes,
156 resulting in minimal experimental constraints. The valve activity was recorded non-stop, both for the
157 7-day acclimatisation and the 24-day experimental periods. Then, the data were analysed using the
158 LabView 8.6 software ([™]National Instruments, Austin, TX, USA).

159

160 Three behavioural parameters were assessed: daily and hourly VMC, VOA and VOD. Micro-closures
161 (VMC) designate partial and rapid closures followed by a re-opening of the valves. A VOA equal to
162 100 % and 0 % meant that the valves were opened at maximal and minimal amplitudes, respectively,
163 during the studied duration. Finally, A VOD equal to 100 % and 0 % meant that the valves were
164 opened or closed, respectively, during the studied duration. Mean daily VMC, VOA and VOD were the
165 average results of the hourly values recorded for each oyster. The HFNI biosensor enabled to
166 individually measure the daily growth of shells by calculating the minimal distance between the
167 electrodes when the oyster valves were closed. For instance, in bivalve molluscs, calcification

168 appears at the shell internal surface. Thus, the cumulative daily shell length growth was calculated as
169 indicated: minimal distance between electrodes (day (n) – day (1)) x 100 % (n/24), n representing the
170 number of the day. To obtain the actual shell length growth, manual measurements (digital caliper, ±
171 0.1 mm) were performed at the start (D_0) and at the end (D_{24}) of the experiment. The difference in
172 mm equals 100 % growth. Growth rates have been measured using the HFNI valvometer based on
173 the fact that calcification in bivalves occurs in the mantle cavity, all over the shell's internal structure
174 (Figure 4). When daily growth layers are deposited, the minimum distance between the electrodes
175 glued to the shells increases providing a good proxy of growth (Schwartzmann *et al.*, 2011; Berge *et*
176 *al.*, 2015).

177 To study the daily biological rhythm ($\tau = 24$ h) of the valve activity, a statistical chronobiological
178 analysis was conducted using the software Time Series Analysis Seriel Cosinor 6.3
179 (<http://www.euroestech.com>). Several steps were required to validate a significant biological rhythm
180 (Gouthiere *et al.*, 2005). The first step included verifying the data quality of biological and physical
181 phenomena. The absence of random distribution in data was assessed using an autocorrelation
182 diagram, whereas the absence of a stationary phenomenon was calculated with a partial
183 autocorrelation function (Box *et al.*, 2015). The second step involved searching significant periodicity
184 ($p = 0.95$), using the Lomb and Scargle periodogram (Scargle, 1982). The third step added a modelling
185 and statistical validation using the Cosinor model, which is a cosine function included in a regression
186 model (Nelson *et al.*, 1979). Two tests validated the model and, thus, the existence of a biological
187 rhythm. The elliptic test (Bingham *et al.*, 1982) had to be rejected, leading to a probability of null
188 amplitude hypothesis of < 0.05 . When conditions were met, a chronobiometric parameter computed
189 the Percent Rhythm (PR, %), which is the percentage of cyclic behaviour determined by the Cosinor
190 model.

191

192 *2.4 Contaminant analysis*

193

194 Chlortoluron and MPs concentrations were measured twice: on day 5 (D_5 , 5 days after the start of
195 the experiment) and day 24 (D_{24} , the last day of the experiment). Water samples ($V = 100$
196 mL/replicate with $n = 3$ replicates) and oyster tissues ($n = 7$ /condition) exposed to both chlortoluron
197 and the cocktail solution were assayed using the QuEChERS method (Quick Easy Cheap Efficient
198 Rugged and Safe, Anastassiades *et al.*, 2003) and the SBSE analysis (Stir Bar Sorptive Extraction,
199 Baltussen *et al.*, 1999) (Pfannkoch *et al.*, 2010 and from the AOAC 2007.01 method of Lehotay,
200 2007).

201 To analyse the oyster tissue and determine chlortoluron concentrations, 3 g of oyster tissue was
202 spiked with an internal standard solution. Milli-Q water was added to the solution to reach 15 g.
203 After being vortexed once, samples were added with 15 ml of acetonitrile (Pestnorm, VWR
204 Chemicals), before being vortexed a second time (1 min, 160 rpm). Then, 6 g of magnesium sulphate
205 (anhydrous, BioReagent, Sigma-Aldrich) and 1.5 g of sodium acetate (anhydrous, >99 %, Sigma-
206 Aldrich) were added to the samples, which were then vortexed again (10 min, 250 rpm) and
207 centrifuged (5 min, 4700 rpm, 5 °C). Two phases were formed and supernatant containing the
208 organic phase with acetonitrile was sampled. Then, 10 mL of the supernatant were collected in the
209 previous step and were taken and placed in a 125 mL flask, where 40 ml of sodium carbonate (0.1 M,
210 ≥ 99.7 %, Sigma-Aldrich) was added. A Twister (magnetic stir bar coated with polydiméthylsiloxane)
211 was inserted into the bottle. SBSE extraction was performed (16 h, 700 rpm), in the dark). After the
212 completion of the extraction, the bars were rinsed off with Milli-Q water before being dried on a
213 fabric.

214 To analyse the chlortoluron seawater exposure samples, 10 mL of methanol (Pestnorm, VWR
215 Chemicals) was put in a bottle containing the internal standard solution. 100 mL of seawater and a
216 Twister were then added. Protected from light, the solution was stirred for 16 hours at 700 rpm.
217 Chlortoluron was quantified by gas chromatography (HP 7890N equipped with a Combipal MPS2
218 multifunction injection system, 300 °C, Gerstel, Switzerland) and mass spectrometry (GC/MS),
219 coupled with a TDU (Thermal Desorption Unit, Gerstel: 50 °C (0.5 min) to 280 °C (6 min) at 15 °C/min)

220 and a temperature-programmed injector (Cooled Injection System, Gerstel: -10 °C (0.05 min) to 300
221 °C (10 min) at 12 °C/s). The temperature was programmed from 70 °C (5 min) to 150 °C increasing by
222 20 °C/min, then to 320 °C (5 min) increasing by 7 °C/min. The carrier gas, helium, was at constant
223 flow rate (1 mL/min). The capillary column was a 5-ms RXi (Restek, Bellefonte, USA): 30m x 0.25mm
224 ID x 0.25µm (film thickness) column. Through tandem mass spectrometry (Agilent 7000 Triple Quad),
225 a chromatograph and a detector were coupled together. The quantitative analysis of chlortoluron
226 was performed in MRM mode by internal calibration. Two transitions for each compound were set
227 up: the first to quantify the molecules and the second to confirm their nature. The second step
228 defined the nature by calculating the qualifier/quantifier ratio and comparing them to the reference
229 values of pure compounds. The acquisition frequency of each fragment was of 2 cycles/s.

230

231 Due to the small sizes of particles (20-25 µm), the MPs analysis in oyster tissues was not performed.
232 According to the protocol described in Bringer *et al.* (2020a & 2020b), flow cytometer (Attune
233 Acoustic Focusing Cytometer) enabled a quantitative analysis of MPs in the 10 µg MP.L⁻¹ condition. 2
234 mL samples were prepared with the MPs solution (n=8/condition). The samples were vortexed
235 (StarLab Vortex IR, 12,000 rpm for 20 s) before being transferred into the cytometer to homogenise
236 the solution. 300 mL were taken to carry out a flow cytometry analysis. Calibration was conducted to
237 reach an analysis rate of 500 mL.min⁻¹ and a saturation of maximum 10,000 particles detected. The
238 blank obtained enabled to remove background particles (naturally present in seawater). Using
239 seawater filtered at 0.2 µm, a first calibration step was conducted to select the < 200-time detected
240 particles.

241

242 2.5 Statistics

243

244 All data were expressed as means ± standard error of the mean (SEM). Treatment differences were
245 determined using a one-way analysis of variance (ANOVA), when both criteria of homogeneity of

246 variance (Levene's test) and normality of distribution (Shapiro-Wilk) were met. When these
247 requirements were not met, a statistical analysis was performed using the non-parametric Kruskal-
248 Wallis test, followed by the Nemenyi Post-hoc test for pair comparisons. Statistical comparisons were
249 made on the means of each parameter according to the exposure conditions. Significant difference
250 was accepted when p -value < 0.05. The statistical analysis was conducted using R.

251

252 **3. Results**

253 *3.1 Pollutant exposure*

254

255 It is noted, that no significant difference was observed concerning the assays carried out for
256 chlortoluron (FSW and total tissues), nor for MP (FSW). Chlortoluron concentrations were
257 determined in exposure waters (Table 1.a). Under control, water contamination increased over time.
258 On D_{24} (condition chlortoluron vs cocktail), an increase in the concentration of chlortoluron was
259 noted. The concentration of chlortoluron in the cocktail condition was slightly higher than for the
260 condition of chlortoluron alone (at D_5 and D_{24}). In both cases, the measured concentrations were well
261 above the theoretical concentration of $30 \mu\text{g.L}^{-1}$. The concentrations of chlortoluron measured on the
262 soft body of the oyster are shown in Table 1.a. No trace of herbicide was detected in control
263 individuals. While in exposed individuals, tissue concentrations of chlortoluron were measured at
264 both sampling times D_5 and D_{24} .

265

266 The concentrations of MP in FSW (Table 1.b) were above the desired theoretical concentration of 10
267 $\mu\text{g of MP.L}^{-1}$. The approximate actual concentrations calculated in this study were determined using
268 additional data from a previous study (Bringer *et al.*, 2020a), which used and tested the same
269 commercial MP (HDPE, 20-25 μm), at different concentrations. The additional data allowed for
270 calibration and validation of the current results. The concentrations of MP were stable over time (D_5

271 and D₂₄) and similar between the exposure conditions (MP alone vs cocktail). In addition, an analysis
 272 of the particles in the control FSW was performed and no MP particles were detected.

273

274 **Table 1. (a)** Theoretical and measured chlortoluron (herbicide) concentrations in oyster tissues (dry weight)
 275 and seawater for three conditions: control, chlortoluron and cocktail (MPs + chlortoluron). Values are mean ±
 276 SEM for n=7/condition for oyster tissues and n=3/condition for water samples. **(b)** Theoretical, measured MPs
 277 and approximate concentrations concentrations in three conditions: control (n=12), MPs (n=8) and cocktail
 278 (MPs + chlortoluron, n=8). Values are mean ± SEM.

279 **a**

Conditions	Samples time	Theoretical ($\mu\text{g.L}^{-1}$)	Oyster tissues (ng.g^{-1})	Water exposure ($\mu\text{g.L}^{-1}$)
control	Day 5	0	0	0.06 ± 0.0
	Day 24	0	0	1.5 ± 0.9
chlortoluron	Day 5	30	226.9 ± 69.5	58.3 ± 25.1
	Day 24	30	229.5 ± 92.0	85.5 ± 17.0
cocktail	Day 5	30	366.0 ± 132.7	74.6 ± 4.9
	Day 24	30	242.3 ± 199.6	96.6 ± 25.8

280 **b**

Conditions	Samples time	Theoretical ($\mu\text{g MP.L}^{-1}$)	Measured (MP.mL ⁻¹) <i>20-25 μm size</i>	Approximate concentrations ($\mu\text{g MP.L}^{-1}$)
control	Day 5	0	0	0
	Day 24	0	0	0
MPs	Day 5	10	93.9 ± 31.2	47
	Day 24	10	111.7 ± 11.5	56
cocktail	Day 5	10	124.4 ± 30.3	60
	Day 24	10	107.6 ± 49.3	54

281

282 3.2 Valve activity of oysters


283 3.2.1 Valve Micro-Closures (VMC)

284

285 The number of VMC per day and per individual is shown on Fig. 1a. Compared to the other
 286 conditions, the daily VMC in MPs over the 24-day experiment highlighted a significant increase ($p <$
 287 0.001). The mean VMC for MPs was 2.9 times higher than for the control condition (Fig. 1b). Fig. 1b
 288 clearly showed a significant decrease in the chlortoluron and cocktail VMC, compared to the control

289 condition VMC. The mean VMC for the conditions of exposure to chlortoluron and cocktail were
290 respectively 40.3 % and 58 % lower compared to the control condition.

291



292

293 **Fig. 1. (a)** Daily VMC showing the 7-day acclimatisation and the 24-day exposure periods. **(b)** Histograms of
294 mean VCM \pm SEM for the 24-day exposure period. The different conditions are: control (white, n = 7),
295 chlortoluron alone (green, n = 6), MPs alone (blue, n = 8) and cocktail (red, n = 5). Different letters indicate
296 significant differences between concentrations ($p < 0.05$).

297


298 3.2.2 Valve Opening Amplitude (VOA)

299

300 In terms of VOA, the 24-day experiment underlined contrasting results between the MPs condition
301 and the two other conditions containing herbicides (Fig. 2a, b, c). Although the VOA was similar for
302 the MPs and control conditions (with no significant differences), it was significantly higher in
303 chlortoluron and cocktail conditions compared to control (both $p < 0.001$)

304 Over the entire experiment, the mean VOA displayed major differences between the four conditions
305 ($p < 0.001$). Compared to the control condition, a significant increase in mean VOA of 18.2 % and 22.3
306 % were respectively noted for the chlortoluron alone and cocktail conditions (Fig. 2d).

307



308

309

310 **Fig. 2.** Valve Opening Amplitude (VOA) in percentage for each condition: **(a)** chlortoluron alone (green), **(b)** MPs
 311 alone (blue) and **(c)** cocktail (red), in comparison with the control condition (white). **(d)** Histograms of VOA (%)
 312 for the entire exposure and all conditions. Values are mean \pm SEM for control ($n = 7$), chlortoluron ($n = 6$), MPs
 313 ($n = 8$) and cocktail ($n = 5$) conditions. Different letters indicate significant differences between concentrations
 314 ($p < 0.05$).


315

316 3.2.3 Valve Opening Duration (VOD)

317

318 Over the 24-day experiment (Fig. 3 a, b, c), the VOD for both the chlortoluron and cocktail conditions
 319 remained unchanged. In contrast, the VOD for the MPs condition significantly decreased compared
 320 to the control condition. Indeed, over the entire experiment, the mean VOD highlighted a significant
 321 decrease ($p < 0.001$) of 5.1 % for the MPs condition, compared to the control one (Fig. 3d).

322



323

324

325 **Fig. 3.** Valve Opening Duration (VOD) in percentage for each condition: **(a)** chlortoluron alone (green), **(b)** MPs
 326 alone (blue) and **(c)** cocktail (red), in comparison with the control condition (white). **(d)** Histograms of VOA (%)
 327 for the entire exposure and all conditions. Values are mean \pm SEM for control (n = 7), chlortoluron (n = 6), MPs
 328 (n = 8) and cocktail (n = 5) conditions. Different letters indicate significant differences between concentrations
 329 ($p < 0.05$).

330

331 3.2.4 Chronobiological analysis

332

333 Through a statistical chronobiological approach, a daily rhythm analysis was performed using the
 334 Cosinor method. When comparing the three conditions to the control one, no significant disruption
 335 of VOA rhythm was observed. Figure S2 displays for each condition, the spectral analysis of the valve
 336 activity (Lomb and Scargle periodogram). It highlighted a significant periodicity in the valve
 337 behaviour. The daily mean VOA for all conditions were: $\tau = 24.0 \pm 0.1$ (control); 23.8 ± 0.1

338 (chlortoluron); 23.9 ± 0.1 (MPs); 23.9 ± 0.1 h (cocktail). Periodicities were observed using the Cosinor
 339 model, which validated the presence of a daily rhythm for all conditions ($p < 0.0001$ for all
 340 conditions). The percent rhythms (PR) between conditions were quite close, except for chlortoluron,
 341 which had a lower PR. PR were 46.4 % (control), 32.3 % (chlortoluron), 42.8 % (MPs) and 45 %
 342 (cocktail).


343

344 3.2.5 Shell Growth

345

346 The Fig. 4a displays the cumulative daily shell growth, monitored for 24 days. No significant
 347 difference was calculated between the control and chlortoluron conditions ($p = 0.07$) but this p-value
 348 is close to the statistical significance and, therefore express a trend. In contrast, over 24 days, oysters
 349 exposed to the MPs and cocktail conditions underlined a lower shell growth. Indeed, after the 24-day
 350 period, the mean shell length growth for the MPs and cocktail conditions respectively was reduced of
 351 50.4 % ($p < 0.0001$) and 42.7 % ($p = 0.021$), in comparison with the control one (Fig. 4b). The oyster
 352 shell growth observed for the MPs and cocktail conditions were also lower than observed for the
 353 exposure to chlortoluron alone.

354



355

356 **Fig. 4. (a)** Daily shell length growth monitored for all conditions (mm). **(b)** After the completion of the 24-day

357 experiment, histograms displaying the mean shell growth (mm) of the oysters equipped with electrodes. The

358 different conditions are: control (no pollutants, white), chlortoluron alone (green), MPs alone (blue) and
359 cocktail (red). Values are mean \pm SEM for control (n = 7), chlortoluron (n = 6), MPs (n = 8) and cocktail (n = 5)
360 conditions. Different letters indicate significant differences between concentrations ($p < 0.05$).

361

362 3.2.6 Summary of the main results

363

364 The actual concentrations measured on D₂₄, in the FSW exposure (for MP and chlortoluron alone and
365 cocktail), are presented in Table 2, as well as in the remainder of this present study. In our work, the
366 analysis validated the existence of a daily rhythm for all tested conditions. No significant differences
367 in chronobiological rhythms were noted between the four conditions tested. Regarding the
368 behavioural parameters, our results suggest that, when exposed to chlortoluron, the oyster valve
369 activity was significantly disrupted (Table 2). Indeed, in comparison with the control condition, a
370 decrease in the number of VMC and an increase in VOA were noted. In terms of MPs exposure, a
371 significant increase in VMC, and a decrease in VOD and growth were observed. Concerning the
372 cocktail condition, a decrease in VMC and growth, and an increase in VOA were highlighted.
373 Depending on the exposure conditions, the valve behaviour in oysters responded differently.

374

375 **Table 2.** Summary of the effects on the Pacific oyster valve activity induced after a 24-day exposure to
376 chlortoluron alone (85 $\mu\text{g}\cdot\text{L}^{-1}$), 20-25 μm HDPE MPs alone (112 MP. mL^{-1}) and a combination of both (cocktail,
377 97 $\mu\text{g}\cdot\text{L}^{-1}$ of chlortoluron + 108 MP. mL^{-1}). Measured parameters are VMC (Valve Micro-Closures), VOA (Valve
378 Opening Amplitude), VOD (Valve Opening Duration), daily rhythm (chronobiological analysis) and shell growth (
379 \uparrow : significant increase compared to the control condition, \downarrow : significant decrease compared to the control
380 condition and - : absence of significant effects compared to the control condition).

381

Conditions	VMC	VOA	VOD	Daily rhythm	Shell growth
Chlortoluron alone	↓	↑	-	-	-
MPs alone	↑	-	↓	-	↓
Cocktail	↓	↑	-	-	↓

382

383 4. Discussion

384 4.1 Experimental concentrations versus natural environment

385

386 In our FSW exposures, concentrations between 58 and 96 $\mu\text{g.L}^{-1}$ were measured The study by
387 Valiente Moro *et al.* (2012) reported an IC_{50} (half-maximum inhibitory concentration) of 50 $\mu\text{g.L}^{-1}$ on
388 the freshwater phytoplankton *Ankistrodesmus fusiformis*. Although this concentration is higher than
389 those commonly found in the Pertuis Charentais, it is quite close to the maximum concentrations
390 detected in some European rivers. Indeed, concentrations of herbicide (chlortoluron in particular) of
391 20 $\mu\text{g.L}^{-1}$ and more (Beitz *et al.*, 1994) were determined. For herbicide exposure, control FSW
392 highlighted a slight contamination to chlortoluron. Being volatile, it could easily evaporate in the
393 atmosphere and contaminate the other tanks (Ineris, 2013). Nevertheless, the concentrations of
394 chlortoluron measured in the control FSW samples were satisfactory, to validate the study. An
395 increase of chlortoluron concentrations was observed between the start (D_5) and the end of the
396 experiment (D_{24}). This increase could be explained by the initial presence of chlortoluron in natural
397 seawater (0.14 $\mu\text{g.L}^{-1}$ in 2018, Action program, 2018) and regular water changes during the
398 experiment. In this present study, the theoretical concentration of HDPE MPs (20-25 μm) was set at
399 10 $\mu\text{g MP.L}^{-1}$, where other studies have tested these same concentrations (Beiras *et al.*, 2018;
400 Constant, 2018; Lei *et al.*, 2018 and Dris *et al.*, 2016). To assess MPs/NPs in water, flow cytometry
401 was performed, as used in some studies (Long *et al.*, 2017; Caputo *et al.*, 2021; Kaile *et al.*, 2020).
402 Instead of being 10 $\mu\text{g MP.L}^{-1}$, the average concentration for MPs-contained solutions was
403 approximately of 50 $\mu\text{g MP.L}^{-1}$ at the two sampling periods. However, the assays by cytometry, we

404 confirm contents between 93.9 and 107.6 MP.mL⁻¹. The highest environmental concentration of MP
405 detected in Artic sea ice cores is around 13.7 MP.mL⁻¹ (Peeken *et al.*, 2018). However, the size classes
406 of MP detected in the natural environment are larger than MP tested in the laboratory (Paul-Pont *et*
407 *al.*, 2018). Our present study could be defined as realistic in terms of size tested but not in terms of
408 concentration. According to a study on the characterization of MP on the Atlantic coast, the majority
409 of MP (64 %) were < 40 µm. In addition, about half (42 %) of them were PE (Enders *et al.*, 2015).
410 These results justify the use of HDPE-MP (20-25 µm) in this study. In addition, some equipment used
411 in aquaculture for bivalves consists mainly of PE (Lusher *et al.*, 2017 and Bringer *et al.*, 2021). The
412 differences observed between the theoretical and measured MP concentrations could come from
413 the aggregation of MP (Alimi *et al.*, 2018; Michels *et al.*, 2018) on the walls of the exposure tanks.
414 Long *et al.* (2017) clearly showed that micro-PS may attach to glassware, form homo-aggregates and
415 hetero-aggregates. In addition, regular water changes, for the welfare of oysters, could also have
416 played a role in increasing theoretical concentrations.

417

418 *4.2 Behavioural responses of juveniles' oysters (C. gigas) to chlortoluron exposure*

419

420 Chlortoluron is a photosynthesis-disrupting herbicide, and more specifically as an inhibitor of
421 photosystem II (Faggiano *et al.*, 2010). Phytoplankton communities have been shown to be very
422 sensitive to herbicides (Solomon *et al.*, 1996). Indeed, effects on phytoplankton growth and
423 development have been observed (El-Sheekh *et al.*, 1994), and could indirectly affect filter feeders
424 such as oysters when they feed. Chlortoluron showed a significant decrease in VMC in number
425 compared to condition control. An equivalent result has already been demonstrated for *Mytilus*
426 *galloprovincialis* exposed to cypermethrin (insecticide) with a reduced the valve closure in a time-
427 dependent manner (Ayad *et al.*, 2011). Higher VOA, at condition control, was also detected following
428 exposure to chlortoluron. An apparent increase in the rate of opening of the shell under
429 contamination with an herbicide (lenacil) was reported in an earlier study (Chmist *et al.*, 2019).

430 However, this parameter has been little studied in order to obtain relevant interpretations (Bae &
431 Park, 2014). Chlortoluron may have a specific reducing effect on some distinctive phytoplankton
432 species important for the physiology, development and behaviour of oysters (Stachowski-Haberkorn
433 *et al.* 2008). Additional studies should be carried out in order to better understand the effect of this
434 herbicide on phytoplankton populations and on oyster nutrition. **Moreover**, combined effect of VMC
435 decrease and VOA increased may suggest that the herbicide lead to an apparent relaxed behaviour
436 of **valve activity**. Next investigations should focus on **putative** neurotoxic effects of chlortoluron on
437 oyster valve activity. Indeed, it would be of interest to study if the chlortoluron may block conduction
438 of action potential at the voltage-gated sodium channels level, which plays a crucial role in
439 membrane excitability in nerve cell, and lead to a decrease of neuro-muscular response. **This**
440 **neurotoxic effect has already been shown** for many neurotoxins produced by animals or plants, such
441 as saxitoxin and tetrodotoxin (Boulot *et al.*, 2017).

442

443 *4.3 Behavioural responses of juveniles' oysters (C. gigas) to microplastics exposure*

444

445 Oysters exposed to MP showed a higher number of VMC compared to the others conditions. VMC is
446 defined as a marker of stress in oysters (Tran *et al.*, 2010). The presence of MP in the seawater,
447 which could be detected by oysters, can induce mechanical stress (Sow *et al.*, 2011). The presence of
448 harmful substances could activate chemoreceptors on the edge of the oyster mantle or directly
449 affect the gills when the pollutants are in direct contact with the tissue of the pallial cavity,
450 responsible for the rapid increase in the frequency of VMC. According to the work of Castrec *et al.*
451 (2018), a similar hypothesis could be made, implying that the increase in the frequency of VMC is a
452 protective behavioral response to accelerate interval water turnover and minimize potential contacts
453 with MP, irritating the gills (Zhu *et al.*, 2020). A significant decrease in VOD was also measured.
454 Likewise, a freshwater bivalve (*Corbicula fluminea*) had a behavioral closure reaction - as a protective
455 strategy - when exposed to a metallic contaminant (Tran *et al.*, 2004a). The branchial cavities and the

456 labial palps (Ward *et al.*, 1998) determine the different modes of selection and sorting of particles,
457 which could slow down the filtration of nutrients of exposed oysters, affect their feeding activity,
458 their behaviour, and reduce their metabolism and their growth. Indeed, several studies have noted
459 that toxins and foreign bodies can modify the nutritional activity of oysters through behavioral
460 modifications (Tran *et al.*, 2010; Haberkorn *et al.*, 2010). Oysters exposed to MP grow more slowly
461 than control oysters. Some previous work has shown non-effects of MP on oysters during subchronic
462 exposure (Green, 2016; Revel *et al.*, 2020). However larger sizes of MP (103 μm and 300 μm) were
463 tested contrary to this present study (20-25 μm). The study by Gardon *et al.* (2018) showed a
464 decrease in the scope for growth from 25 $\mu\text{g MP.L}^{-1}$ of PS-MP in *P. margaritifera*. These results
465 indicate that the exposed oysters must have drawn their energy from the reserves. In addition, the
466 assimilation of microalgae was disturbed by the presence of micro-PS, and the energy provided by
467 food intake was lower for exposed oysters (Gardon *et al.*, 2018). In the study conducted by Thomas
468 *et al.* (2020), the condition index at the highest concentration (106 MP.L^{-1}) decreased significantly
469 over time. The concentration tested is similar to our concentrations measured in MP exposure
470 waters.

471

472 *4.4 Effects of co-contamination by microplastics and chlortoluron on juveniles' oysters (C.* 473 *gigas)*

474

475 For oysters exposed to the cocktail, results similar to the chlortoluron condition were noted, on VMC
476 and VOA. No effect on VOD was noted in contrast to its decrease for exposure to MP. Chlortoluron
477 and MP may have antagonistic effects. A previous study demonstrated antagonistic effects between
478 NP and glyphosate on the microalgae *Microcystis aeruginosa* (Zhang *et al.*, 2018). Combined
479 exposure to diclofop-methyl and silver nanoparticles has been shown to antagonize the growth
480 inhibition of *A. Thaliana* (Li *et al.*, 2018). In our study, growth decreased under the cocktail condition
481 as for the MP condition. Studies have pointed out that MP can make hydrophobic organic chemicals

482 bioavailable by acting as an absorption vector (Horton *et al.*, 2018). However, our results do not
483 show significantly different effects for the cocktail, compared to MP and therefore we cannot
484 conclude on a potential vectorization of chlortoluron on MP. The results showed that the synergistic
485 effect of chlortoluron and MPs is not significant, compared to the contamination conditions alone.
486 Their common ecotoxic effects are affected by physical and chemical factors such as particle size,
487 color, composition, function (Yu *et al.*, 2021). Due to their high filtration rate, marine bivalves, like
488 oysters, are particularly sensitive to chemical contamination, which can be a major source of stress
489 and behavioral alteration (Sokolova and Lannig, 2008; Islam and Tanaka, 2004).

490

491 **5. Conclusions**

492 On a larger scale, valvometric technology is a tool for monitoring water quality, using oysters as
493 biosensors (Ahmed *et al.*, 2015). This study provided valuable data on the behavior of bivalves
494 exposed to pollutants potentially present in the marine environment. Exposure to chlortoluron had
495 little impact on their behavior and none on their growth. This study provided insight into the
496 evidence that MP could potentially affect the physiology and behavior of *C. gigas*. MPs tended to act
497 as stressors in bivalves, triggering urgent behavioral responses. In addition, MP could affect the
498 filtration capacity of bivalves (nutrient system) and, indirectly, their growth. The combination of the
499 two pollutants showed antagonistic responses to the conditions of contamination alone. Rarely used
500 in studies, valvometric technology is an innovative, integrated and comprehensive marker that helps
501 to understand the effects of environmental pollution. Future studies on the effects of MP on bivalves
502 could use HFNI valvometry, mimicking environmental conditions (*e.g.* size range, concentrations,
503 pollutant cocktail).

504

505 **Acknowledgements**

506

507 A. Bringer received a PhD grant (Comité Régional de la Conchyliculture de la Charente Maritime
508 CRC17) as well as financial support from the Région Nouvelle Aquitaine and Comité Départemental
509 de la Charente Maritime (CD17) to develop his research. The AQUAECOs project (**Amélioration de la**
510 **QUALité Environnementale sur les zones CO**nchylicoles des Pertuis Charentais), funded by a
511 partnership with CRC17, and the AFB (Agence Française de la Biodiversité, *Parc naturel marin de*
512 *l'estuaire de la Gironde et de la mer des Pertuis*) contributed financially to this study. The funding was
513 partly supported by the University of La Rochelle, the University of Bordeaux, the Centre National de
514 la Recherche Scientifique (France). The authors thank the France Naissain hatchery for providing the
515 oyster specimens as well as our collaborators for their work and help.

516

517 **References**

518

519 Action program, Coastal water quality in shellfish farming areas, Qualyse, CD17, 2018-2020 (accessed
520 October 2020).

521

522 Ahmed, H., Ushirobira, R., Efimov, D., Tran, D., & Massabuau, J. C. (2015). Velocity estimation of
523 valve movement in oysters for water quality surveillance. *IFAC-PapersOnLine*, 48(11), 333-338.

524

525 Akcha, F., Barranger, A., & Bachère, E. (2020). Genotoxic and epigenetic effects of diuron in the
526 Pacific oyster: in vitro evidence of interaction between DNA damage and DNA
527 methylation. *Environmental Science and Pollution Research*, 1-15.

528

529 Alimi, O. S., Farnier Budarz, J., Hernandez, L. M., & Tufenkji, N. (2018). Microplastics and nanoplastics
530 in aquatic environments: aggregation, deposition, and enhanced contaminant transport.

531 *Environmental science & technology*, 52(4), 1704-1724.

532

533 Andrade, H., Massabuau, J. C., Cochrane, S., Ciret, P., Tran, D., Sow, M., & Camus, L. (2016). High
534 frequency non-invasive (HFNI) bio-sensors as a potential tool for marine monitoring and
535 assessments. *Frontiers in Marine Science*, 3, 187.

536

537 Ayad, M. A., Fdil, M. A., & Mouabad, A. (2011). Effects of cypermethrin (pyrethroid insecticide) on
538 the valve activity behavior, byssal thread formation, and survival in air of the marine mussel *Mytilus*
539 *galloprovincialis*. *Archives of environmental contamination and toxicology*, 60(3), 462-470.

540

541 Bae, M. J., & Park, Y. S. (2014). Biological early warning system based on the responses of aquatic
542 organisms to disturbances: a review. *Science of the Total Environment*, 466, 635-649.

543

544 Baldwin, A. K., Corsi, S. R., & Mason, S. A. (2016). Plastic debris in 29 Great Lakes tributaries: relations
545 to watershed attributes and hydrology. *Environmental Science & Technology*, 50(19), 10377-10385.

546

547 Bayne, B.L., Ahrens, M., Allen, S.K., D'auriac, M.A., Backeljau, T., Beninger, P., Bohn, R., Boudry, P.,
548 Davis, J., Green, T., Guo, X., Hedgecock, D., Ibarra, A., Kingsley- Smith, P., Krause, M., Langdon, C.,
549 Lape_gue, S., Li, C., Manahan, D., Mann, R., Perez-Paralle, L., Powell, E.N., Rawson, P.D., Speiser, D.,
550 Sanchez, J.-L., Shumway, S., Wang, H., 2017. The proposed dropping of the genus *Crassostrea* for all
551 pacific cupped oysters and its replacement by a new genus *Magallana*: a dissenting view. *J. Shellfish*
552 *Res.* 36 (3), 545e548.

553

554 Bayne, B., d'Auriac, M.A., Backeljau, T., Beninger, P., Boudry, P., Carnegie, R., Langdon, C., 2019. A
555 scientific name for Pacific oysters. *Aquaculture* 499, 373.

556

557 Behrens, D., Rouxel, J., Burgeot, T., & Akcha, F. (2016). Comparative embryotoxicity and genotoxicity
558 of the herbicide diuron and its metabolites in early life stages of *Crassostrea gigas*: implication of
559 reactive oxygen species production. *Aquatic Toxicology*, 175, 249-259.

560

561 Beiras, R., Bellas, J., Cachot, J., Cormier, B., Coussin, X., Engwall, M., Gambardella, C., Garaventa, F.,
562 Keiter, S., Le Bihanic, F., Lopez-Ibanez, S., Piazza, V., Rial, D., Tato, T., Vidal-Linan, L. (2018). Ingestion
563 and contact with polyethylene microplastics does not cause acute toxicity on marine zooplankton.
564 *Journal of Hazardous Materials*, 360, 452-460.

565

566 Beitz, H., Schmidt, H., & Herzel, F. (1994). Occurrence, toxicological and ecotoxicological significance
567 of pesticides in groundwater and surface water. In *Pesticides in ground and surface water* (pp. 1-56).
568 Springer, Berlin, Heidelberg.

569

570 Berge, J., Daase, M., Renaud, P. E., Ambrose, W. G. Jr., Darnis, G., Last, K. S., et al. (2015). Unexpected
571 levels of biological activity during the polar night offer new perspectives on a warming arctic. *Curr.*
572 *Biol.* 25, 2555–2561.

573

574 Betsill, M., Gonzalez, J., & Woods, A. (2019). The Impact of Microplastic Ingestion on the Bivalve
575 Filtration Efficiency of the Hooked Mussel (*Ischadium recurvum*) from the Chesapeake Bay. Research
576 Proposal for UH 1604: Honors Research Practices at Virginia Tech.

577

578 Bhagat, J., Nishimura, N., & Shimada, Y. (2020). Toxicological interactions of
579 microplastics/nanoplastics and environmental contaminants: Current knowledge and future
580 perspectives. *Journal of Hazardous Materials*, 123913.

581

582 Bingham, C., Arbogast, B., Cornélissen, G., Lee, J.-K., Halberg, F. (1982). Inferential statistical methods
583 for estimating and comparing cosinor parameters. *Chronobiologia*, 9, 397–439.

584

585 Born, M. P., & Schüttrumpf, H. (2019, September). Microplastic in Coastal Areas-Impact of Waves,
586 Sediments and Saltwater on the Degradation Behaviour. In *International Conference on Microplastic
587 Pollution in the Mediterranean Sea* (pp. 158-163). Springer, Cham.

588

589 Boullot, F., Castrec, J., Bidault, A., Dantas, N., Payton, L., Perrigault, M., **Tran, D.**, Amzil, Z., Boudry, P.,
590 Soudant, P., Hégaret, H., Fabioux, C. (2017) Molecular Characterization of Voltage-Gated Sodium
591 Channels and Their Relations with Paralytic Shellfish Toxin Bioaccumulation in the Pacific Oyster
592 *Crassostrea gigas*. *Mar. Drugs*. 15(1): 21.

593

594 Box, G. E., Jenkins, G. M., Reinsel, G. C., & Ljung, G. M. (2015). *Time series analysis: forecasting and
595 control*. John Wiley & Sons.

596

597 Bringer, A., Thomas, H., Prunier, G., Dubillot, E., Bossut, N., Churlaud, C., Clérandeau, C., Le Bihanic,
598 F., Cachot, J. (2020a). High density polyethylene (HDPE) microplastics impair development and
599 swimming activity of Pacific oyster D-larvae, *Crassostrea gigas*, depending on particle
600 size. *Environmental Pollution*, 113978.

601

602 Bringer, A., Cachot, J., Prunier, G., Dubillot, E., Clérandeau, C., & Thomas, H. (2020b). Experimental
603 ingestion of fluorescent microplastics by pacific oysters, *Crassostrea gigas*, and their effects on the
604 behaviour and development at early stages. *Chemosphere*, 126793.

605

606 Bringer, A., Le Floch, S., Kerstan, A., & Thomas, H. (2021). Coastal ecosystem inventory with
607 characterization and identification of plastic contamination and additives from aquaculture
608 materials. *Marine Pollution Bulletin*, 167, 112286.

609

610 Caputo, F., Vogel, R., Savage, J., Vella, G., Law, A., Della Camera, G., Hannon, G., Peacock, B., Mehn,
611 D., Ponti, J., Geiss, O., Aubert, D., Prina-Mello, A., Calzolari, L. (2021). Measuring particle size
612 distribution and mass concentration of nanoplastics and microplastics: addressing some analytical
613 challenges in the sub-micron size range. *Journal of Colloid and Interface Science*, 588, 401-417.

614

615 Castrec, J., Soudant, P., Payton, L., Tran, D., Miner, P., Lambert, C., Goïc, N. Le, Huvet, A., Quillien, V.,
616 Boullot, F., Amzil, Z., Hégaret, H., Fabioux, C., 2018. Bioactive extracellular compounds produced by
617 the dinoflagellate *Alexandrium minutum* are highly detrimental for oysters. *Aquatic Toxicology*. 199,
618 188–198.

619

620 Chmist, J., Szoszkiewicz, K., & Drożdżyński, D. (2019). Behavioural responses of *Unio tumidus*
621 freshwater mussels to pesticide contamination. *Archives of environmental contamination and*
622 *toxicology*, 77(3), 432-442.

623

624 CNC (Comité National de la Conchyliculture), 2016. [http://www.cnc-france.com/La-Production-](http://www.cnc-france.com/La-Production-francaise.aspx)
625 [francaise.aspx](http://www.cnc-france.com/La-Production-francaise.aspx) (accessed July 2019).

626

627 CNC (Comité National de la Conchyliculture), 2020 ; Les statistiques. www.cnc-france.com (accessed
628 October 2020).

629

630 Cole, M., Lindeque, P., Halsband, C., Galloway, T.S. (2011). Microplastics as contaminants in the
631 marine environment: a review. *Marine Pollution Bulletin*, 62, 2588-2597.

632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657

Cole, M., Galloway, T.S. (2015). Ingestion of nanoplastics and microplastics by Pacific oyster larvae. *Environmental Sciences Technology*, 49(24), 14625-14632.

Constant, M. (2018). Source, Transfert et Devenir des microplastiques (MP) en MerMediterranee Nord-Occidentale. *PhD thesis. University of Perpignan-CEFREM*, pp277.

Díaz-Mendoza, C., Mouthon-Bello, J., Pérez-Herrera, N. L., & Escobar-Díaz, S. M. (2020). Plastics and microplastics, effects on marine coastal areas: a review. *Environmental Science and Pollution Research*, 1-10.

Dris, R., Lahens, L., Rocher, V., Gasperi, J., & Tassin, B. (2016). Premières investigations sur la contamination en microplastiques d'une zone urbaine. Cas de l'agglomération parisienne, vol. 12.

El-Sheekh, M. M., Ktkat, H. M., & Hammouda, O. H. (1994). Effect of atrazine herbicide on growth, photosynthesis, protein synthesis, and fatty acid composition in the unicellular green alga *Chlorella kessleri*. *Ecotoxicology and environmental safety*, 29(3), 349-358.

Enders, K., Lenz, R., Stedmon, C. A., & Nielsen, T. G. (2015). Abundance, size and polymer composition of marine microplastics $\geq 10 \mu\text{m}$ in the Atlantic Ocean and their modelled vertical distribution. *Marine pollution bulletin*, 100(1), 70-81.

Eriksen, M., Maximenko, N., Thiel, M., Cummins, A., Lattin, G., Wilson, S., Hafner, J., Zellers, A., & Rifman, S. (2013). Plastic pollution in the South Pacific subtropical gyre. *Marine pollution bulletin*, 68(1-2), 71-76.

658 Faggiano, L., de Zwart, D., García-Berthou, E., Lek, S., & Gevrey, M. (2010). Patterning ecological risk
659 of pesticide contamination at the river basin scale. *Science of the Total Environment*, 408(11), 2319-
660 2326.

661

662 France Agrimer, 2020. Chiffres clés des filières pêche et aquaculture en France, en 2020 (access in
663 February).
664 [https://www.franceagrimer.fr/fam/content/download/64767/document/CC%20pêche%20aqua%20](https://www.franceagrimer.fr/fam/content/download/64767/document/CC%20pêche%20aqua%20FR.pdf?version=22021)
665 [FR.pdf?version=22021](https://www.franceagrimer.fr/fam/content/download/64767/document/CC%20pêche%20aqua%20FR.pdf?version=22021)).

666

667 Gall, S. C., & Thompson, R. C. (2015). The impact of debris on marine life. *Marine pollution*
668 *bulletin*, 92(1-2), 170-179.

669

670 Gamain, P., Gonzalez, P., Cachot, J., Clérandeau, C., Mazzella, N., Gourves, P. Y., & Morin, B. (2017).
671 Combined effects of temperature and copper and S-metolachlor on embryo-larval development of
672 the Pacific oyster, *Crassostrea gigas*. *Marine pollution bulletin*, 115(1-2), 201-210.

673

674 Gardon, T., Reisser, C., Soyez, C., Quillien, V., & Le Moullac, G. (2018). Microplastics affect energy
675 balance and gametogenesis in the pearl oyster *Pinctada margaritifera*. *Environmental science &*
676 *technology*, 52(9), 5277-5286.

677

678 Gardon, T., Morvan, L., Huvet, A., Quillien, V., Soyez, C., Le Moullac, G., & Le Luyer, J. (2020).
679 Microplastics induce dose-specific transcriptomic disruptions in energy metabolism and immunity of
680 the pearl oyster *Pinctada margaritifera*. *Environmental Pollution*, 266, 115180.

681

682 Girard, S., & Agúndez, J. A. P. (2014). The effects of the oyster mortality crisis on the economics of
683 the shellfish farming sector: Preliminary review and prospects from a case study in Marennes-Oleron
684 Bay (France). *Marine Policy*, 48, 142-151.

685

686 Gouthiere, L., Mauvieux, B., Davenne, D., & Waterhouse, J. (2005). Complementary methodology in
687 the analysis of rhythmic data, using examples from a complex situation, the rythmicity of
688 temperature in night shift workers. *Biological Rhythm Research*, 36(3), 177-193.

689

690 Green, D. S. (2016). Effects of microplastics on European flat oysters, *Ostrea edulis* and their
691 associated benthic communities. *Environmental pollution*, 216, 95-103.

692

693 Haberkorn, H., Lambert, C., Le Goïc, N., Moal, J., Suquet, M., Guéguen, M., Sunila, I & Soudant, P.
694 (2010). Effects of *Alexandrium minutum* exposure on nutrition-related processes and reproductive
695 output in oysters *Crassostrea gigas*. *Harmful algae*, 9(5), 427-439.

696

697 Haberkorn, H., Tran, D., Massabuau, J. C., Ciret, P., Savar, V., & Soudant, P. (2011). Relationship
698 between valve activity, microalgae concentration in the water and toxin accumulation in the
699 digestive gland of the Pacific oyster *Crassostrea gigas* exposed to *Alexandrium minutum*. *Marine*
700 *pollution bulletin*, 62(6), 1191-1197.

701

702 Horton, A. A., Svendsen, C., Williams, R. J., Spurgeon, D. J., & Lahive, E. (2017). Large microplastic
703 particles in sediments of tributaries of the River Thames, UK—Abundance, sources and methods for
704 effective quantification. *Marine pollution bulletin*, 114(1), 218-226.

705

706 Horton, A.A., Vijver, M.G., Lahive, E., Spurgeon, D.J., Svendsen, C., Heutink, R., van Bodegom, P.M.,
707 Baas, J., 2018. Acute toxicity of organic pesticides to *Daphnia magna* is unchanged by co-exposure to
708 polystyrene microplastics. *Ecotoxicol. Environ. Saf.* 166, 26-34.
709
710 Ineris. (2013). Valeur guide environnementale: Chlortoluron (15545-48-9)
711 <https://substances.ineris.fr/fr/substance/cas/15545-48-9/3>
712
713 Islam, M. S., & Tanaka, M. (2004). Impacts of pollution on coastal and marine ecosystems including
714 coastal and marine fisheries and approach for management: a review and synthesis. *Marine pollution*
715 *bulletin*, 48(7-8), 624-649.
716
717 Kaile, N., Lindivat, M., Elio, J., Thuestad, G., Crowley, Q. G., & Hoell, I. A. (2020). Preliminary results
718 from detection of microplastics in liquid samples using flow cytometry. *Frontiers in Marine Science*, 7,
719 856.
720
721 Lee, K.W., Shim, W.J., Kwon, O.Y., Kang, J.H., 2013. Size-dependent effects of micro polystyrene
722 particles in the marine copepod *Tigriopus japonicus*. *Environmental Sciences Technology*, 47 (19),
723 11278-11283.
724
725 Lee, D. H., Rhee, Y. J., Choi, K. S., Nam, S. E., Eom, H. J., & Rhee, J. S. (2017). Sublethal concentrations
726 of atrazine promote molecular and biochemical changes in the digestive gland of the Pacific oyster
727 *Crassostrea gigas*. *Toxicology and Environmental Health Sciences*, 9(1), 50-58.
728
729 Lehotay SJ (2007). Determination of pesticide residues in foods by acetonitrile extraction and
730 partitioning with magnesium sulfate: collaborative study. *J AOAC Int* 90:485–520
731

732 Lei, L., Wu, S., Lu, S., Liu, M., Song, Y., Fu, Z., ... & He, D. (2018). Microplastic particles cause intestinal
733 damage and other adverse effects in zebrafish *Danio rerio* and nematode *Caenorhabditis*
734 *elegans*. *Science of the Total Environment*, 619, 1-8.

735

736 Leonard, S. Dispersion des pesticides vers les eaux - Bassin de la Sèvre Niortaise et Marais Poitevin
737 Période 2004-2014, 2017 - [http://www.sevre-niortaise.fr/wp-](http://www.sevre-niortaise.fr/wp-content/uploads/2017/04/etude_pesticides_SAGE_SNMP_31032017.pdf)
738 [content/uploads/2017/04/etude_pesticides_SAGE_SNMP_31032017.pdf](http://www.sevre-niortaise.fr/wp-content/uploads/2017/04/etude_pesticides_SAGE_SNMP_31032017.pdf) (accessed January 2020).

739

740 Li, X., Ke, M., Zhang, M., Peijnenburg, W. J. G. M., Fan, X., Xu, J., Zhang, Z., Lu, T., Fu, Z., & Qian, H.
741 (2018). The interactive effects of diclofop-methyl and silver nanoparticles on *Arabidopsis thaliana*:
742 growth, photosynthesis and antioxidant system. *Environmental pollution*, 232, 212-219.

743

744 Long, M., Paul-Pont, I., Hegaret, H., Moriceau, B., Lambert, C., Huvet, A., Soudant, P., 2017.
745 Interactions between polystyrene microplastics and marine phytoplankton lead to species-specific
746 hetero-aggregation. *Environmental Pollution*, 228, 454-463.

747

748 Lucasson, A. (2018). Characterization and diversity of the mechanisms of the mortality syndrome
749 affecting juveniles of *Crassostrea gigas*. *PhD thesis. University of Montpellier*, pp165.

750

751 Lusher, A., Hollman, P., & Mendoza-Hill, J. (2017). Microplastics in fisheries and aquaculture. *FAO*
752 *Fisheries and Aquaculture Technical Paper (FAO) eng no. 615*.

753

754 Mai, H., Morin, B., Pardon, P., Gonzalez, P., Budzinski, H., & Cachot, J. (2013). Environmental
755 concentrations of irgarol, diuron and S-metolachlor induce deleterious effects on gametes and
756 embryos of the Pacific oyster, *Crassostrea gigas*. *Marine environmental research*, 89, 1-8.

757

758 Mat, A.M., Massabuau, J.C., Ciret, P., Tran, D. (2014) Looking for the clock mechanism responsible for
759 the circatidal behavior in the oyster *Crassostrea gigas*. *Mar. Biol.* 161: 89-99.
760

761 Michels, J., Stippkugel, A., Lenz, M., Wirtz, K., & Engel, A. (2018). Rapid aggregation of biofilm-
762 covered microplastics with marine biogenic particles. *Proceedings of the Royal Society B*, 285(1885),
763 20181203.
764

765 Mottier, A., Kientz-Bouchart, V., Serpentine, A., Lebel, J. M., Jha, A. N., & Costil, K. (2013). Effects of
766 glyphosate-based herbicides on embryo-larval development and metamorphosis in the Pacific oyster,
767 *Crassostrea gigas*. *Aquatic toxicology*, 128, 67-78.
768

769 Nelson, W., Tong, Y.L., Lee, J.K., Halberg, F., 1979. Methods for cosinor-rhythmometry.
770 *Chronobiologia*, 6, 305–323.
771

772 Paul-Pont, I., Tallec, K., Gonzalez-Fernandez, C., Lambert, C., Vincent, D., Mazurais, D., Zambonino-
773 Infante, J.L., Brotons, G., Lagarde, F., Fabioux, C., Soudant, P., & Huvet, A. (2018). Constraints and
774 priorities for conducting experimental exposures of marine organisms to microplastics. *Frontiers in*
775 *Marine Science*, 5, 252.
776

777 Peeken, I., Primpke, S., Beyer, B., Gütermann, J., Katlein, C., Krumpfen, T., Bergmann, M., Hehemann,
778 L., & Gerdt, G. (2018). Arctic sea ice is an important temporal sink and means of transport for
779 microplastic. *Nature communications*, 9(1), 1-12.
780

781 Pfannkoch, E. A., Stuff, J. R., Whitecavage, J. A., & Moran, J. H. (2010). High Throughput Method for
782 the Determination of PAHs in Seafood by QuEChERS-SBSE-GC-MS.
783

784 Phuong, N. N. (2018). Analytical developments for the characterization and quantification of
785 microplastic contamination of sedimentary and biological matrices: application to shellfish farming
786 areas in Pays de la Loire. *PhD thesis. University of Nantes*, pp232.

787

788 PlasticsEurope (2016) Plastics – the Facts 2016. URL [http://](http://www.plasticseurope.de/Document/plastics—the-facts-2016-15787.aspx)
789 www.plasticseurope.de/Document/plastics—the-facts-2016-15787.aspx

790

791 PlasticsEurope. (2020). Plastics - the facts 2020: an analysis of European plastics production, demand
792 and waste data. PlasticsEurope Internal Report 4312 (64pp.).
793 <https://www.plasticseurope.org/en/resources/publications/4312-plastics-facts-2020>.

794

795 Revel, M., Châtel, A., Perrein-Ettajani, H., Bruneau, M., Akcha, F., Sussarellu, R., Rouxel, J., Costil, K.,
796 Decottignies, P., Cognie, B., Lagarde, F., & Mouneyrac, C. (2020). Realistic environmental exposure to
797 microplastics does not induce biological effects in the Pacific oyster *Crassostrea gigas*. *Marine*
798 *pollution bulletin*, 150, 110627.

799

800 Rist, S., Baun, A., Almeda, R., & Hartmann, N. B. (2019). Ingestion and effects of micro-and
801 nanoplastics in blue mussel (*Mytilus edulis*) larvae. *Marine pollution bulletin*, 140, 423-430.

802

803 Scargle, J. D. (1982). Studies in astronomical time series analysis. II-Statistical aspects of spectral
804 analysis of unevenly spaced data. *The Astrophysical Journal*, 263, 835-853.

805

806 Schwartzmann, C., Durrieu, G., Sow, M., Ciret, P., Lazareth, C. E., and Massabuau, J.-C. (2011). *In situ*
807 giant clam growth rate behavior in relation to temperature: a one-year coupled study of high-
808 frequency noninvasive valvometry and sclerochronology. *Limnol. Oceanogr.* 56, 1940–1951.

809

810 Sokolova, I. M., & Lannig, G. (2008). Interactive effects of metal pollution and temperature on
811 metabolism in aquatic ectotherms: implications of global climate change. *Climate Research*, 37(2-3),
812 181-201.

813

814 Soletchnik, P., Bouquet, A. L., & Mille, D. (2018). Flash Info Maline 2008-2018, ten years already... of
815 communication on the mortalities of oysters (*Crassostrea gigas*) and mussels (*Mytilus edulis*) in the
816 Pertuis Charentais. *Ifremer report*, pp18.

817

818 Solomon, K.R., Baker, D.B., Richards, R.P., Dixon, D.R., Klaine, S.J., LaPoint, T.W., 1996. Ecological risk
819 assessment of atrazine in North American surface waters. *Environmental Toxicology and Chemistry*
820 15 (1), 31-74.

821

822 Song, Y. K., Hong, S. H., Jang, M., Han, G. M., & Shim, W. J. (2015). Occurrence and distribution of
823 microplastics in the sea surface microlayer in Jinhae Bay, South Korea. *Archives of environmental*
824 *contamination and toxicology*, 69(3), 279-287.

825

826 Sow, M., Durrieu, G., Briollais, L., Ciret, P., & Massabuau, J. C. (2011). Water quality assessment by
827 means of HFNI valvometry and high-frequency data modeling. *Environmental monitoring and*
828 *assessment*, 182(1-4), 155-170.

829

830 Stachowski-Haberkorn, S., Quiniou, F., Nedelec, M., Robert, R., Limon, G., & de La Broise, D. (2008).
831 In-situ microcosms, a tool for assessment of pesticide impacts on oyster spat (*Crassostrea*
832 *gigas*). *Ecotoxicology*, 17(4), 235-245.

833

834 Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M.E.J., Goïc, N.L., Quillien, V.,
835 Mingant, C., Epelboin, Y., Charlotte, C., Julien, G., Johan, R., Ika, P.-P., Philippe, S., Arnaud, H., 2016.

836 Oyster reproduction is affected by exposure to polystyrene microplastics. Proc. Natl. Acad. Sci. U. S.
837 A. 113, 2430-2435.

838

839 Tallec, K., Huvet, A., Di Poi, C., González-Fernández, C., Lambert, C., Petton, B., Le Goic, N., Berchel,
840 M., Soudant, P., & Paul-Pont, I. (2018). Nanoplastics impaired oyster free living stages, gametes and
841 embryos. *Environmental pollution*, 242, 1226-1235.

842

843 Thomas, M., Jon, B., Craig, S., Edward, R., Ruth, H., John, B., Dick, V.A., Heather, L.A., & Matthew, S.
844 (2020). The world is your oyster: low-dose, long-term microplastic exposure of juvenile oysters.
845 *Heliyon*, 6(1), e03103.

846

847 Tran D., Fournier E., Durrieu G. and Massabuau J-C. (2004a) Copper detection in the asiatic clam
848 *Corbicula fluminea*: optimum valve closure response. *Aquatic Toxicology*, 66(3): 333-343.

849

850 Tran D., Massabuau J-C and Garnier-Laplace J. (2004b) The effect of carbon dioxide on uranium
851 bioaccumulation in the freshwater clam *Corbicula fluminea*. *Environmental Toxicology and Chemistry*,
852 23(3): 739-747.

853

854 Tran D., Bourdineaud J-P, Massabuau J-C and Garnier-Laplace J. (2005) Modulation of uranium
855 bioaccumulation by hypoxia in the freshwater clam *Corbicula fluminea*: induction of multixenobiotic
856 resistance protein and heat shock protein 60 in gill tissues. *Environmental Toxicology and Chemistry*,
857 24(9): 2278-2284

858

859 Tran D., Fournier E., Durrieu G. and Massabuau J-C. (2007) *Corbicula fluminea* valve closure response
860 as sensitivity threshold to inorganic mercury contamination. *Environmental Toxicology and*
861 *Chemistry*, 26(7): 1545-1551.

862

863 Tran, D., Haberkorn, H., Soudant, P., Ciret, P., & Massabuau, J. C. (2010). Behavioral responses of
864 *Crassostrea gigas* exposed to the harmful algae *Alexandrium minutum*. *Aquaculture*, 298(3-4), 338-
865 345.

866

867 Tran, D., Ciutat, A., Mat, A., Massabuau, J.C., Hégaret, H., Lambert, C., Le Goic, N., Soudant, P, (2015).
868 The toxic dinoflagellate *Alexandrium minutum* disrupts daily rhythmic activities at gene transcription,
869 physiological and behavioral levels in the oyster *Crassostrea gigas*. *Aquatic Toxicology*, 158, 41–49.

870

871 Valiente Moro, C., Bricheux, G., Portelli, C., & Bohatier, J. (2012). Comparative effects of the
872 herbicides chlortoluron and mesotrione on freshwater microalgae. *Environmental toxicology and*
873 *chemistry*, 31(4), 778-786.

874

875 Wagner, M., Scherer, C., Alvarez-Munoz, D., Brennholt, N., Bourrain, X., Buchinger, S., Fries, E.,
876 Grosbois, C., Klasmeier, J., Marti, T., Rodriguez-Mozaz, S., Urbatzka, R., Vethaak, A., Winther-Nielsen,
877 M., Reifferscheid, G., 2014. Microplastics in freshwater ecosystems: what we know and what we
878 need to know. *Environ. Sci. Eur.* 26 (1), 12.

879

880 Wang, F., Gao, J., Zhai, W., Liu, D., Zhou, Z., & Wang, P. (2020). The influence of polyethylene
881 microplastics on pesticide residue and degradation in the aquatic environment. *Journal of hazardous*
882 *materials*, 394, 122517.

883 Ward, J. E.; Levinton, J.; Shumway, S.; Cucci, T. Particle sorting in bivalves: *in vivo* determination of
884 the pallial organs of selection. *Mar. Biol.* 1998, 131 (2), 283–292.

885 Ward, J. E., Zhao, S., Holohan, B. A., Mladinich, K. M., Griffin, T. W., Wozniak, J., & Shumway, S. E.
886 (2019). Selective Ingestion and Egestion of Plastic Particles by the Blue Mussel (*Mytilus edulis*) and

887 Eastern Oyster (*Crassostrea virginica*): Implications for Using Bivalves as Bioindicators of Microplastic
888 Pollution. *Environmental science & technology*, 53(15), 8776-8784.

889

890 Wright, S.L., Rowe, D., Thompson, R.C., Galloway, T.S., 2013. Microplastic ingestion decreases energy
891 reserves in marine worms. *Current Biology* 23 (23), R1031–R1033.

892

893 Xu, X. Y., Lee, W. T., Chan, A. K. Y., Lo, H. S., Shin, P. K. S., & Cheung, S. G. (2017). Microplastic
894 ingestion reduces energy intake in the clam *Atactodea striata*. *Marine pollution bulletin*, 124(2), 798-
895 802.

896

897 Yu, H., Peng, J., Cao, X., Wang, Y., Zhang, Z., Xu, Y., & Qi, W. (2021). Effects of microplastics and
898 glyphosate on growth rate, morphological plasticity, photosynthesis, and oxidative stress in the
899 aquatic species *Salvinia cucullata*. *Environmental Pollution*, 279, 116900.

900

901 Zhang, Q., Qu, Q., Lu, T., Ke, M., Zhu, Y., Zhang, M., Zhang, Z., Du, B., Pan, X., Sun, L., & Qian, H.
902 (2018). The combined toxicity effect of nanoplastics and glyphosate on *Microcystis aeruginosa*
903 growth. *Environmental Pollution*, 243, 1106-1112.

904


905 Zhao, S., Zhu, L., Wang, T., & Li, D. (2014). Suspended microplastics in the surface water of the
906 Yangtze Estuary System, China: first observations on occurrence, distribution. *Marine pollution*
907 *bulletin*, 86(1-2), 562-568.

908

909 Zhu, X., Qiang, L., Shi, H., & Cheng, J. (2020). Bioaccumulation of microplastics and its in vivo
910 interactions with trace metals in edible oysters. *Marine Pollution Bulletin*, 154, 111079.

911 **Supplementary data**

912




913

914

915 **Fig. S1.** Experimental design located in an experimental laboratory (LIENSs La Rochelle – CNRS). The numbers
916 correspond to: **(1)** experimental batch of exposed oysters equipped with valvometric electrodes (zoom
917 picture), **(2)** second experimental batch of exposed oysters, **(3)** buffer tank, **(4)** reduced vibrations thanks to the
918 capacitance chamber and **(5)** HFNI valvometer (High Frequency Non-Invasive) connected to the electrodes
919 stuck on oysters.

920

Chronobiological analysis



921

922 **Fig. S2.** Rhythmic valve behaviour of *C. gigas*. For each condition, the daily periodicity (τ) was determined by

923 spectral analysis (Lomb and Scargle periodogram). The X axis indicates the test duration in order to determine a

924 significant period. The Y axis displays the power (arbitrary units) of the determined period. The red dotted line

925 highlights a significant period for p -value = 0.95.