



Hg concentrations and stable isotope variations in tropical fish species of a gold-mining-impacted watershed in French Guiana

Laure Laffont, Johanna Menges, Sylvaine Goix, Sophie Gentès, Régine Maury-Brachet, Jeroen Sonke, Alexia Legeay, Patrice Gonzalez, Raphaëlle Rinaldo, Laurence Maurice

► To cite this version:

Laure Laffont, Johanna Menges, Sylvaine Goix, Sophie Gentès, Régine Maury-Brachet, et al.. Hg concentrations and stable isotope variations in tropical fish species of a gold-mining-impacted watershed in French Guiana. *Environmental Science and Pollution Research*, 2021, 28 (43), pp.60609-60621. <10.1007/s11356-021-14858-7>. <hal-03434104>

HAL Id: hal-03434104

<https://hal.science/hal-03434104v1>

Submitted on 25 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Hg concentrations and stable isotopes variations in tropical fish species of a gold-mining impacted watershed in French Guiana

Laure Laffont^{1,*}, Johanna Menges², Sylvaine Goix^{1,3}, Sophie Gentès⁴, Régine Maury-Brachet⁴, Jeroen E. Sonke¹, Alexia Legeay⁴, Patrice Gonzalez⁴, Raphaëlle Rinaldo⁵, Laurence Maurice^{1,*}

1 – Geosciences Environnement Toulouse, CNRS/IRD/CNES/Université Paul Sabatier, 14 avenue Edouard Belin, 31400 Toulouse, France

2 – Section 4.6, Geomorphology, GFZ–German Research Centre for Geosciences, 14473 Potsdam, Germany

3 – University Paul Sabatier, 14 avenue Edouard Belin, 31400 Toulouse, France

4 – EPOC, EPHE, Univ. Bordeaux, CNRS, 33120 Arcachon, France

5 – Parc Amazonien de Guyane, 1 rue Lederson 97354 Remire-Montjoly, France

*Corresponding authors: laure.laffont@get.omp.eu; laurence.maurice@ird.fr

ORCID numbers: 0000-0001-5850-7937 (L. Laffont), 0000-0001-7223-4787 (J. Menges), 0000-0002-5392-793X (S. Gentes), 0000-0001-7146-3035 (J. Sonke), 0000-0002-1052-142X (A. Legeay), 0000-0003-2628-2993 (P. Gonzalez), 0000-0003-3482-3892 (L. Maurice)

Key words: mercury, gold-mining, French Guiana, fish, stable isotopes, methylmercury

Abstract:

The aim of the study was to determine if gold mining activities could impact the mercury (Hg) concentrations and isotopic signatures in freshwater fish consumed by riparian people in French Guiana. Total Hg, MeHg concentrations and Hg stable isotopes ratios were analyzed in fish muscles from different species belonging to three feeding patterns (herbivorous, periphytophagous and

piscivorous). We compared tributaries impacted by gold-mining activities (Camopi, CR) with a pristine area upstream (Trois-Sauts, TS), along the Oyapock River. We measured $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ to examine whether Hg patterns are due to differences in trophic level. Differences in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values between both studied sites were only observed for periphytophagous fish, due to difference of CN baselines, with enriched values at TS. Total Hg concentrations and Hg stable isotopes fractionations showed that Hg accumulated in fish from both areas have undergone different biogeochemical processes. $\Delta^{199}\text{Hg}$ variation in fish (-0.5 to 0.2‰) was higher than the ecosystem baseline defined by a $\Delta^{199}\text{Hg}$ of -0.66‰ in sediments, and suggested limited aqueous photochemical MeHg degradation. Photochemistry-corrected $\delta^{202}\text{Hg}$ in fish were 0.7‰ higher than the baseline, consistent with biophysical and chemical isotope fractionations in aquatic environment. While THg concentrations in periphytophagous fish were higher in the gold-mining area than in TS, the ensemble of Hg isotope shifts in fish is affected by the difference of biotic (methylation/demethylation) and abiotic (photochemistry) processes between both areas and did therefore not allow to resolve the contribution of gold-mining related liquid Hg(0) in fish tissues. Mercury isotopes of MeHg in fish and lower trophic level organisms can be complementary to light stable isotope tracers.

Introduction

Mercury (Hg) is a global pollutant which can induce serious health effects on living organisms depending on its concentration and speciation. The key neurotoxic form of Hg is methylmercury (MeHg) which is produced naturally in aquatic environments by microorganisms (Guimarães et al., 1998; Roulet et al., 2000, 2001; Fleming et al., 2006; Kerin et al., 2006). MeHg is then bioaccumulated and biomagnified along the trophic chain (Mason et al., 1995; Maurice-Bourgoin et al., 1999; Kwon et al., 2012a, Chen et al., 2014) resulting in highly elevated MeHg concentrations in large predator fish (Roulet et al., 2000, 2001, Maury-Brachet et al., 2006). Fish consumption is a major pathway of human exposure to MeHg which can lead to severe health effects such as trembling, eyesight problems, coordination disorders, defects and ultimately death (WHO, 1990). It is therefore important to understand how MeHg is generated and incorporated into the food webs and identify its main sources to the food web.

Indigenous riparian communities with a diet heavily relying on fish are especially at risk (Maurice-Bourgoin et al., 1999 and 2000, Mergler et al., 2007). In French Guiana, the regular consumption of local fish leads to elevated Hg concentrations in blood and hair of the local population exceeding the World Health Organization (WHO, 1990) safety limit ($10 \mu\text{g}\cdot\text{g}^{-1}$) (Cordier et al., 1998, Fréry et al., 2001). Neurodevelopmental investigations among MeHg-exposed children in French Guiana also showed significant links between mercury concentrations and the presence of neurological deficiencies (Cordier et al., 2002). The main sources of Hg to the aquatic environment (and ultimately humans) in French Guiana are small-scale gold mining activities and the erosion of tropical soils. Small- and large-scale gold mining activities are the largest Hg emission source ($>55\%$) in South America (Pacyna et al., 2010). The gold mining activities have been greatly enhanced by the increase of the gold price since 1980. Over the last decade gold mining has dramatically impacted the environment by increasing deforestation, soil erosion, surface water turbidity as well as the fishing and hunting pressure (Rimbaud et al., 2017). Large parts of the gold mining operations are clandestine and not controlled by the government. The total gold exploitation in French Guiana has been estimated at 5 to 10 tons per year by the WWF (World Wildlife Fund for Nature, 2018). The mean ratio of “liquid Hg used” to “gold produced” is estimated to be 5 (AMAP/UN, 2019) leading to an estimation of 25 to 50 tons per year of liquid Hg used for the amalgamation process in this region. Even though the liquid Hg used by small-scale gold-miners is partly recovered by retorting, large quantities of elemental Hg are liberated into the environment, both as vapor emitted to the atmosphere and in its liquid form directly released into soils and sediments (Velásquez-López, P. C., 2010). These locally Hg-contaminated soils and sediments, but also natural and non-gold mining anthropogenic Hg from soils, can then be transferred to the aquatic environment (Meech et al., 1997; Roulet et al., 2000; Dominique et al., 2007a and b; Goix et al., 2019). The speciation processes of liquid Hg were investigated in a freshwater indoor micro-cosmos which revealed Hg transfer between Hg^0 droplets and water column, leading to the formation of aqueous Hg(II) and subsequently MeHg (Dominique et al., 2007a). However, there is no data on natural settings. The direct impact of liquid Hg released during gold mining activities on the aquatic food chain and subsequently on the local population remains few documented (Telmer et al., 2006, Marshall et al., 2018). Soil run-off is an important source of Hg to

the hydrosystems. In this region, soils are characterized by elevated organic matter (around 60 $\mu\text{g.g}^{-1}$ in the upper horizon in Grimaldi et al., 2008 and Guedron et al., 2006) and clay content and are enriched in Al- and Fe-oxyhydroxydes with a high storage capacity for Hg (Roulet et al., 1998, 2000; Grimaldi et al., 2008; Guedron et al., 2006, 2009). These soils accumulated atmospheric Hg deposition of natural and anthropogenic origin over thousands of years resulting in medium to high Hg concentrations ranging from 0.029 to 0.100 $\mu\text{g.g}^{-1}$ (Goix et al., 2019). Basically, these values are in the low or medium end of the concentration of contaminated soil (Guedron et al., 2006).

The variation in the natural abundances of Hg stable isotopes has become a widely used tool during the last few years (Blum, 2011, Sonke and Blum, 2013). Mercury has seven stable isotopes with nominal masses of 196, 198, 199, 200, 201, 202, and 204 and their isotopic variation can be used to identify and trace biogeochemical pathways in analogy to stable isotopes of lighter elements.

Hg isotope variations can be explained by mass-dependent fractionation (MDF) and mass-independent fractionation (MIF). MDF is proportional to the mass difference of the Hg isotopes and occurs during redox transformations, biological cycling, and volatilization of Hg (Bergquist and Blum, 2009). MIF is predominantly associated with odd numbered Hg isotopes (^{199}Hg and ^{201}Hg) and only induced by a few transformation reactions. The ratio between anomalies $\Delta^{199}\text{Hg}$ and $\Delta^{201}\text{Hg}$ (defined as the difference between measured $\delta^{199}\text{Hg}$ or $\delta^{201}\text{Hg}$ and theoretical $\delta^{199}\text{Hg}$ or $\delta^{201}\text{Hg}$ calculated from mass dependent fractionation only) appears characteristic of two known processes which result in MIF: the nuclear volume effect or the magnetic isotope effect which occurs during photochemical radical pair reactions (Estrade et al., 2009; Bergquist and Blum, 2007). However, in the environment MIF introduced by magnetic isotope effect induced by photochemical reactions is predominant relative to MIF by nuclear volume effect (Blum, 2011, Sonke, 2011). As $\Delta^{199}\text{Hg}$ and $\Delta^{201}\text{Hg}$ signatures remain unchanged during non-photochemical transport or transfer processes, they are especially useful to trace sources, when we know the Hg isotopic signatures, including MIF, of sources, and photochemical transformations of Hg in the environment. More recently, MIF on even isotope ^{200}Hg has been observed and is measured as anomaly $\Delta^{200}\text{Hg}$ (defined as the difference between measured $\delta^{200}\text{Hg}$ and theoretical $\delta^{200}\text{Hg}$ calculated from mass dependent fractionation only). It is used to

discriminate between two atmospheric sources: dry and wet deposits. First have negative $\Delta^{200}\text{Hg}$ mostly representative of GEM signature while wet deposition exhibits positive $\Delta^{200}\text{Hg}$ (Gratz et al., 2010; Demers et al., 2013 and 2015; Lepak et al., 2015; Yin et al., 2016; Fu et al., 2016 and Obrist et al., 2017).

MDF with a large range of $\delta^{202}\text{Hg}$ is observed in natural samples and in many natural samples anomalies signatures have also been reported (Blum et al., 2014). Bergquist and Blum (2007) observed large positive odd-isotopes anomalies during the experimental photochemical degradation of inorganic Hg (IHg) and MeHg as well as in natural fish samples. They ascribed the fractionation during the photochemical reaction and suggested that the complementary odd-isotopes signatures of the residual MeHg in water gets conservatively incorporated into the food web and can therefore be observed in the fish samples. As explained by Tsui et al., odd-isotopes anomalies are likely attributable to differences in diet or % MeHg (Tsui et al., 2012). In a variety of other studies, no indication of *in vivo* MIF was observed (Gantner et al., 2009; Perrot et al., 2010, 2012; Senn et al., 2010). Hg isotopes have been analyzed in high-trophic level biota to reflect the Hg isotope values of MeHg in the ambient environment as trophic fractionation of Hg isotopes in the lower trophic levels is not significant for Hg(II) and MeHg (Tsui et al., 2020). Kwon et al. (2012b) conducted two controlled experiments to evaluate MDF and MIF during trophic transfer into fish. They found no indication of isotopic fractionation and suggested that stable Hg isotope ratios in fish can be used to trace environmental sources of Hg in aquatic ecosystems. More recently, in Tsui et al. review (2020), authors suggested that Hg isotopes signature, especially odd-MIF, can increase with river size due to reduction in canopy cover which induces increase of photochemistry of Hg and MeHg (Tsui et al., 2013).

It is the case of areas impacted by goldmining activities, miners use liquid elemental mercury for the gold amalgamation with a known isotopic signature of mercury (Laffont et al., 2011; Goix et al., 2019). This includes Hg(0) vapor produced by amalgam burning which could be transported in the atmosphere, reduced and redeposited with the precipitation in the forest (foliage, soils, etc.). Actually, Grimaldi et al. 2015 showed that redeposition of Hg(0) vapor after amalgam burning in upper tropical

soils is minor relative to natural Hg. However, natural mercury from pristine sediments showed a different isotopic signature than in gold-mined area (Goix et al., 2019). Mercury inputs in sediments as well as in water column, from natural and from goldmining activities, undergo physico-bio-chemical processes in the hydrosystem (desorption/sorption of mercury between liquid and solid phase, absorption/methylation/demethylation in aquatic organisms, and photochemistry in surface water) that induce isotopes fractionation which are highly dependent of the environment; for example, the MIF signature in macroinvertebrates and fish is mediated by the effect of canopy on in-stream photodemethylation (Tsui et al., 2013). The knowledge of the fractionations induced by each of these processes can help to discriminate between the different sources of mercury for identical environmental conditions.

To understand the feeding relationships among consumers in a food web, carbon and nitrogen stable isotopes analysis are recognized as relevant tools (Layman et al. 2012). Indeed, nitrogen incorporation by organisms leads to an enrichment in $\delta^{15}\text{N}$ of approximately 3 to 5‰ between each trophic level (Cabana & Rasmussen, 1994; Zanden et al. 2001; Caut et al. 2009), allowing the trophic position of species to be inferred. In contrast, the relatively low enrichment of $\delta^{13}\text{C}$ along food chains (0 to 1‰ between two trophic levels) enables discriminating the different sources of organic carbon (Hobson et al. 2002. Neres-Lima et al. 2017).

The main objective of the present study was to determine if gold mining activities impact the Hg concentrations and isotopic signatures of tropical fish species. We studied the Oyapock river basin in French Guiana characterized by a pristine (Trois-Sauts, TS) and a gold mining (Camopi River, CR) area. Specific objectives were: i) to better understand the underlying processes of Hg stable isotopes fractionation in fish species from three different feeding patterns and ii) to assess if Hg isotopes signatures can be used to distinguish anthropogenic from natural Hg in fish muscle, mainly in biofilm-eating fish species. In the frame of the same project (RIMNES research program), Goix et al. (2019) analyzed sediments sampled at the same time and in the same locations than the fish results presented in this article. Our team showed that Hg isotopes distinguish both sources of Hg in sediments and

revealed that release of liquid mercury from gold-mining activities directly impact locally exploited rivers.

Fish samples with different feeding ecology (herbivorous, periphytophagous and piscivorous) have been analyzed for total Hg and nitrogen ($\delta^{15}\text{N}$) and carbon ($\delta^{13}\text{C}$) stable isotopes, while MeHg concentrations have been determined in periphytophagous fish and Hg stable isotopes have been measured in periphytophagous and piscivorous fish species. Samples were collected in two sub-basins of the Oyapock River, in French Guiana: the Oyapock river upstream at Trois-Sauts village, and the Camopi river, respectively in pristine and gold-mining areas.

Methods

2.1 Study area

Our studied area is located in the Oyapock basin at the eastern part of French Guiana, at the Brazilian border (Figure 1). The Oyapock River (404 km long) originates in the Tumuk Humak Mountains and runs northward towards the Atlantic Ocean. Its mean annual discharge reaches $800 \text{ m}^3 \text{ s}^{-1}$ at the hydrological gauging station close to Saint Georges. Samples were collected 50 km upstream at Camopi village and 250 km upstream at Trois Sauts villages. At the Camopi village, the Camopi River (secondary stream), one of its main tributaries, joins the Oyapock River (primary stream). The Camopi River basin is affected by artisanal and small-scale gold mining activities (ASGM) since the 19th century. Nowadays a recent rise in ASGM has been reported on small creeks upstream (tertiary streams), in the Camopi River basin, and on the Oyapock River basin downstream of the confluence with the Camopi River (Parc Amazonien de Guyane, PAG 2017; Rahm et al. 2017). The pristine region of Trois Sauts lies on the Oyapock River 110 km upstream of the Camopi confluence and no gold mining activities have been reported in this area (Gentès et al. 2019).

2.2 Sampling

Fish samples were collected in the Oyapock watershed during the low water stage, in the Camopi river in October 2013 (Hg-contaminated sites impacted by gold mining activities) and in the Oyapock river upstream in October 2012 (the pristine area of Trois Sauts), with the help of local

fishermen. Different fishing techniques were used such as nets and fish lines. Following the protocol of Maury-Brachet et al. (2006), fish were dissected on site with a clean stainless steel scalpel and Teflon dissecting forceps and flesh samples were collected. The standard length (cm) and total fresh weight (g) of each individual were measured immediately. Fish samples were frozen in the field using a generator-powered freezer and shipped to GET and EPOC laboratories under dry ice, with final storage in the laboratory at -80°C. Before further analyses, samples were freeze-dried and manually ground to homogenize in an agate mortar.

For the purpose of this study, two piscivorous (*Hoplias aimara* and *Boulengerella cuvieri*), two periphytophagous (*Pseudoancistrus barbatus* and *Hypostomus gymnorhynchus*) and two herbivorous (*Myloplus rubripinnis* and *Myloplus ternetzi*) species were selected because 1) they are well represented in the Amazon basin (Le Bail et al. 2012); 2) they are regularly consumed by the local population (Fréry et al. 2001) and therefore represent the main Hg exposure pathway; 3) they represent different trophic levels and have a well-known and specific diet (Keith et al., 2000, Le Bail et al., 2000; Durrieu et al., 2005, Maury-Brachet et al., 2006).

The feeding ecology of both piscivorous fish species are mainly fish of varying size, giving them a high trophic position in the food web (see trophic position calculation in results part). Moreover, *Hoplias aimara* was proposed as a bioindicator species for Hg (Durrieu et al., 2005; Gentès et al. 2019). Piscivorous fish were selected by length. *H. aimara* specimen with standard length less than 38 cm could be mistaken with another species (*H. malabaricus*). Small *B. cuvieri* individuals (standard length < 40 cm) are juveniles and their diet is different from that of adult fish. Consequently, we selected *H. aimara* > 38 cm and *B. cuvieri* > 40 cm for analysis.

Periphytophagous fish species are benthic primary consumers and consume exclusively periphyton or biofilms on hard substrates (like rocks or immersed tree trunks, etc.). Periphyton is an important food source for the food web, even when its biomass is low (Lamberti, 1996). Moreover, periphyton has been identified as the principal entry point for methylmercury in an Amazonian region (Roulet et al., 2000). Indeed, several studies have shown that periphyton supports high Hg methylation

rates due to microbial activity in tropical ecosystems (Guimarães et al., 1998; Acha et al., 2005, Klaus et al. 2016).

The both chosen herbivorous species feed exclusively on terrestrial materials from the river banks (leaves, flowers, fruits) and provide information about pelagic primary consumers (Planquette et al., 1996).

2.4 Chemical analysis

2.4.1 Nitrogen and carbon stable isotopes analysis

We measured nitrogen and carbon ratios in muscles of herbivorous, periphytophagous and piscivorous fish species. All samples were dried and reduced to powder and 0.5 mg were weighed and placed in tin capsules (purity 99.998%) and then analyzed using a continuous flow isotope ratio mass spectrometer in Toulouse (Elemental Microanalysis, UK) in Ecolab laboratory (France). The Isoprime 100 spectrometer is coupled in continuous flow mode using pure helium as gas vector to an elemental analyzer Micro Vario for solids combustion. A dual inlet injection system introduces the reference gases (N₂ and CO₂).

The ¹³C/¹²C (denoted δ¹³C) and ¹⁵N/¹⁴N (denoted δ¹⁵N) ratios, expressed in ‰, were calculated as the relative differences between the sample and the conventional standard following Peterson and Fry (1987):

$$\delta^{15}\text{N or } \delta^{13}\text{C (‰)} = [(R_{\text{sample}} - R_{\text{standard}}) / R_{\text{standard}}] \times 1,000 \text{ (Eq. 1)}$$

where R is ¹³C/¹²C or ¹⁵N/¹⁴N (Peterson and Fry, 1987). The international reference material standard is atmospheric N₂ for nitrogen and Vienna Pee Dee Belemnite for carbon.

Standards were run in duplicate every twenty measurements. Three internal standards of urea, alanine and acetanilid were analyzed. The obtained average values were δ¹⁵N = -0.45 ± 0.14 ‰ and δ¹³C = -36.33 ± 0.26 ‰ (n = 9, 2SD) for urea, δ¹⁵N = 8.38 ± 0.23 ‰ and δ¹³C = -23.03 ± 0.18 ‰ (n = 6, 2SD) for alanine and δ¹⁵N = -3.89 ± 0.46 ‰ and δ¹³C = -33.29 ± 0.14 ‰ (n = 6, 2SD) for acetanilid. The highest analytic precisions were 0.46 ‰ for δ¹⁵N and 0.26 ‰ for δ¹³C (2SD).

Measured C/N ratios provide an indication of the fat content in the tissue analyzed (Post et al. 2007). Because lipids are ^{13}C -depleted relative to proteins, high lipid content (with $\text{C/N} > 4$) in organisms affects measured $\delta^{13}\text{C}$ ratios. In our study, all organisms had a $\text{C/N} < 4$, thus not requiring lipid extraction.

2.4.2 Total Hg concentrations

Fish samples were immediately stored in sealed polyethylene bags and frozen at $-18\text{ }^{\circ}\text{C}$. Back to the laboratories, they were freeze-dried and analyzed a couple of months after sampling. Total Hg concentrations (THg) were measured on 108 fish muscle samples, using an Advanced Mercury Analyzer (AMA254) (GET laboratory). Each sample was combusted inside the AMA at 750°C for 150 seconds. The released Hg was trapped in a gold trap and subsequently released and measured with an integrated atomic absorption spectrometer. The accuracy of the measurements was assigned by the analysis of certified reference materials ERM-CE464 (tuna fish) and DOLT4 (dogfish liver) (for the piscivorous fish species) as well as TORT 2 (lobster hepatopancreas) for the less concentrated periphytophagous fish species.

The measured concentrations for the reference materials were in agreements with the certified values for TORT2 ($\text{THg} = 0.307 \pm 0.004\text{ }\mu\text{g.g}^{-1}$; $n = 6$; recovery: 114%) and DOLT4 ($\text{THg} = 2.385 \pm 0.003\text{ }\mu\text{g.g}^{-1}$; $n = 4$; recovery : 92%). For CE464 the recovery was $91 \pm 1\%$ ($4.757 \pm 0.208\text{ }\mu\text{g.g}^{-1}$, $n = 3$). Natural samples were all measured in duplicate. If the difference between the duplicate measurements was above 10%, a third measurement was carried out. All THg concentrations are reported on a dry weight basis.

2.4.3 Hg speciation analyses

We assumed that %MeHg in piscivorous fish is close to 100% (Maury-Brachet et al. 2006) and chose to not analyze mercury speciation of these samples. However, a number of nineteen muscle samples of periphytophagous fish and two certified reference materials, TORT2 and NIST2976, were analyzed. Speciation analyses have been made using double-spike species specific isotope dilution method with GC-ICP-MS detection at the GET laboratory (Monperrus et al. 2008). Subsamples of 200 mg were digested with 5 mL of TMAH in a microwave Discover SP-D (CEM society) with a ramp of

2 min and constant heating during 4 min at 70°C and stirring (Navarro et al. 2011). After cooling, an aliquot of the extract was transferred to glass vials and 5 ml HAc/NaAc buffer solution (pH 3.9) was added to the aliquot. Then, an appropriate amount of isotopically enriched ¹⁹⁹IHg and ²⁰¹MeHg was added to each sample (Monperrus et al, 2008). After letting the sample equilibrate for two hours, the pH was adjusted to pH 3.9 using bi-distilled HCl. The species were then propylated by adding 0.2 ml of newly prepared Sodium-tetra(n-propyl)borate (NaBPr₄) 10% (w/v). Then, 2 ml of hexane was added and each vial was manually shaken for 5 min to extract all the Hg species into the organic phase. An aliquot of the organic phase was then transferred into a GC-vial and kept frozen until analysis. The samples were analyzed using a GC (Thermo) coupled to a sector field ICP-MS (Thermo Element XR) at the GET laboratory. The measured yield of MeHg for the two reference materials, (58 % for TORT2 and 49 % for NIST2976), are consistent with the certified values of 56 % and 46 %, respectively. The average yield of the analysis was 98±8 % quantified by comparison of the sum of the MeHg and IHg concentrations measured with the described method and THg analysis with AMA254.

2.4.4 Hg stable isotope analyses

Appropriate amounts of samples (0.5 g of periphytophagous fish muscle and 0.05 g of piscivorous fish muscle) were digested in 5 ml inverse aqua regia (HNO₃ and HCl, 3:1 v:v) at 120°C for 4 h in closed glass vials on a heating plate. After cooling, 1 ml of H₂O₂ was added and samples were heated again at 120°C for at least 4 h. The average yield of the digestion was 103±6 % for periphytophagous and 104±5 % for piscivorous fish samples quantified by comparison of the results from the MC-ICP-MS and THg analysis with AMA254. In 2013-2014, our technical materials did not allow us to analyze Hg isotopes of herbivorous fish due to low THg concentration.

Hg stable isotope ratios were measured by continuous flow cold vapor generation (using Sn(II) reduction) multi collector inductively coupled plasma mass spectrometry (CV-MC-ICPMS, Thermo Finnigan Neptune, Germany) at the GET laboratory (Toulouse, France) according to previously published methods (Laffont et al., 2009, 2011; Masbou et al., 2013, 2014; Enrico et al., 2016). Instrumental mass bias was corrected by sample bracketing using the international Hg standard NIST SRM 3133 at matching concentrations (several sequences between 0.8 and 2.5 ppb depending

on initial THg of fish) and prepared in the same matrix than samples. Hg isotopic composition is reported as δ -values, which represents the deviation from the bracketing standard and expressed in permil (‰) using the following equation:

$$\delta^{\text{xxx}}\text{Hg} = \left(\frac{\left(\frac{\text{xxxHg}}{198\text{Hg}} \right)_{\text{sample}}}{\left(\frac{\text{xxxHg}}{198\text{Hg}} \right)_{\text{SRM3133}}} - 1 \right) * 1000 \text{ (Eq. 2)}$$

MIF is quantified as the δ -value deviation from the theoretical MDF using the following equation:

$$\Delta^{\text{xxx}}\text{Hg} = \delta^{\text{xxx}}\text{Hg}_{\text{sample}} - \beta * \delta^{202}\text{Hg}_{\text{sample}} \text{ (Eq. 3)}$$

Where β -values for isotopes ^{199}Hg , ^{200}Hg , ^{201}Hg and ^{204}Hg are 0.252, 0.502, 0.752, and 1.493 respectively according to the kinetic MDF law (Blum and Bergquist, 2007). Long-term reproducibility of Hg isotope measurements was assessed by analyzing two reference materials UM-Almaden and ETH in-house Fluka along with the samples at the same THg concentrations. The obtained values for UM-Almaden are $\delta^{202}\text{Hg} = -0.52 \pm 0.14$ ‰, $\Delta^{199}\text{Hg} = -0.03 \pm 0.08$ ‰ and $\Delta^{200}\text{Hg} = 0.03 \pm 0.04$ ‰ (mean \pm 2SD, n = 7) and for ETH in-house Fluka $\delta^{202}\text{Hg} = -1.53 \pm 0.14$ ‰, $\Delta^{199}\text{Hg} = -0.10 \pm 0.04$ and $\Delta^{200}\text{Hg} = 0.02 \pm 0.04$ ‰ (mean \pm 2SD, n = 14) and agree with previously published values (Blum and Bergquist, 2007; Smith et al., 2014). The certified reference materials TORT2 and ERM-CE464 were also analyzed. Measured isotopic compositions for reference materials are consistent with previously published values (Laffont et al., 2009; Masbou et al., 2013). Uncertainties applied to samples were the larger of either the 2SD on the ETH in-house Fluka or the 2SD on sample replicate when available and larger than the 2SD of ETH in-house Fluka.

2.5 Data analysis

2.5.1 Trophic position calculation for piscivorous fish

The position of an organism within a food chain can be deducted from the relatively constant $\delta^{15}\text{N}$ enrichment observed between each trophic level (Vander Zanden and Rasmussen, 1996). The trophic position (TP) of piscivorous fish was calculated based on $\delta^{15}\text{N}$ values and following the equation of Bergamino et al. (2011):

$$\text{TP}_i = [(\delta^{15}\text{N}_i - \delta^{15}\text{N}_{\text{pc}}) / 3.4] + 2 \text{ (Eq 4)}$$

where TP_i represents the average position of species i ; $\delta^{15}N_i$ the average $\delta^{15}N$ value of species i ; $\delta^{15}N_{pc}$ the average $\delta^{15}N$ value of primary consumers; 3.4 the mean $\delta^{15}N$ trophic enrichment occurring per trophic level (Post, 2002); and 2, the trophic position of the baseline organism (= primary consumer, here herbivorous fish species were selected because they present the lowest $\delta^{15}N$ values, in comparison to periphytophagous fish, see result part 3.1).

2.5.2 Statistical analysis

Statistical analyses were done using ©R (Version 3.4.4). Comparison tests were conducted with Wilcoxon (medians) as distribution is not normal and sample size low. A confidence level of 0.95 was chosen. Means or medians were considered to be significantly different when the p value was less than 0.05.

Results

These are first data of THg, MeHg concentrations and stable Hg isotopes ratios for different fish species highly consumed in French Guiana by local population. All the results are detailed in the Supplementary information (Table S1).

3.1 Nitrogen and carbon stable isotopes ratios in fish

Carbon ($\delta^{13}C$) and nitrogen ($\delta^{15}N$) stable isotopes individual values for herbivorous, periphytophagous and piscivorous fish from both sites are plotted in Figure 2. Results shown that the $\delta^{15}N$ is similar between two sites for piscivorous fish (Wilcoxon test, $W = 26$, $p > 0.05$), but not for periphytophagous fish which it is significantly different (Wilcoxon test, $W = 62$, $p = 0.013$) with higher values in TS (10.8 ± 0.6 ‰, $n = 12$) than in CR (9.8 ± 0.7 ‰, $n = 6$). We cannot calculate a Wilcoxon test for herbivorous fish as $n = 5$ at CR but the average $\delta^{15}N$ is close for both site (6.2 ± 0.3 ‰, $n = 9$, at TS and 6.2 ± 0.5 ‰, $n = 5$, at CR). Trophic positions (TP) were 3.8 ± 0.1 and 3.9 ± 0.1 for piscivorous fish for CR and TS, respectively and were not statistically different between both sites (Wilcoxon test, $W = 0.94$, $p = 0.34$).

Periphytophagous fish present a variation of $\delta^{13}C$ (Fig. 2) ranging from -30.5 to -25.8 ‰ with an average of -28.3 ± 1.3 ‰ ($n = 12$) at TS and from -24.0 to -27.9 ‰ with an average of -26.0 ± 1.3 ‰ ($n =$

6) at CR, showing a slightly more positive $\delta^{13}\text{C}$ at CR than TS (Wilcoxon test, $W = 7$, $p = 0.005$). The $\delta^{13}\text{C}$ values of herbivorous fish (-27.2 ± 0.2 ‰, $n = 9$, at TS and -26.7 ± 0.3 ‰, $n = 5$, at CR) are very close between both sites (Wilcoxon test not possible with $n = 5$). Piscivorous fish $\delta^{13}\text{C}$ values are also statistically similar between both sites: -26.2 ± 0.5 ‰ ($n = 6$) at TS and -26.2 ± 0.2 ‰ ($n = 6$) at CR (Wilcoxon test, $W = 19$, $p = 0.94$).

3.2 Total mercury concentrations in fish

THg concentrations (Fig. 3) in herbivorous species ($n = 36$) vary between 0.006 and 0.037 $\mu\text{g.g}^{-1}$ dw and are significantly higher at TS compared to CR (0.022 ± 0.006 $\mu\text{g.g}^{-1}$ at TS and 0.012 ± 0.008 $\mu\text{g.g}^{-1}$ at CR, Wilcoxon test, $W = 276$, $p = 8.1 \times 10^{-5}$). In periphytophagous species ($n = 41$), THg concentrations vary between 0.017 and 0.248 $\mu\text{g.g}^{-1}$ dw and are significantly higher at CR relatively to TS (0.13 ± 0.05 $\mu\text{g.g}^{-1}$ at TS and 0.20 ± 0.04 $\mu\text{g.g}^{-1}$ at CR, Wilcoxon test, $W = 28$, $p = 0.003$). In piscivorous species ($n = 30$), THg concentrations range from 0.216 to 6.346 $\mu\text{g.g}^{-1}$ dw and are not significantly different between CR and TS (average of 2.0 ± 0.9 $\mu\text{g.g}^{-1}$ at TS and 2.9 ± 1.5 $\mu\text{g.g}^{-1}$ at CR, Wilcoxon test, $W = 65$, $p = 0.094$). THg concentrations in all piscivorous fish samples (except one) exceeded the US EPA (2012) recommendation THg value of 0.256 $\mu\text{g.g}^{-1}$ dw for a daily consumption of 0.113 kg (ww)/day of fish for a pregnant women of 78 kg.

3.3 Methylmercury concentrations in fish

The %MeHg has only been measured in periphytophagous species because it reaches more than 95 % in piscivorous study species (Maury-Brachet et al., 2006): it varies between 92 and 95 % with a mean of 95 ± 1 % (1SD, $n = 6$) in CR and between 82 and 95 % with a mean of 89 ± 5 % (1SD, $n = 13$) in TS (Table S1). The difference of %MeHg between both locations is significant (Wilcoxon test, $W = 4$, $p = 0.0009$) with the %MeHg higher at CR. Moreover, we observe that the variance of %MeHg is higher at TS than at CR (20.9 and 1.3 respectively, variance unity is the square of %).

3.4 Mercury isotopes signatures

For periphytophagous fish, $\delta^{202}\text{Hg}$ values vary between -1.40 ‰ and -0.89 ‰ (mean of -1.14 ± 0.16 ‰ (1SD)) at TS and between -1.41 ‰ and -1.03 ‰ (mean of -1.21 ± 0.14 ‰) at CR. For

piscivorous fish, $\delta^{202}\text{Hg}$ values vary between -1.68 ‰ and -1.17 ‰ (mean of -1.42 ± 0.11 ‰) at TS and between -1.68 ‰ and -0.95 ‰ (mean of -1.29 ± 0.21 ‰) at CR (Fig. 4). Regarding the stable Hg isotopes anomalies, $\Delta^{199}\text{Hg}$ values for periphytophagous vary between -0.49 ‰ and 0.00 ‰ (mean of -0.30 ± 0.16 ‰) at TS and between -0.60 ‰ and 0.20 ‰ (mean of 0.06 ± 0.12 ‰) at CR. $\Delta^{199}\text{Hg}$ values for piscivorous vary between -0.50 ‰ and -0.03 ‰ (mean of -0.38 ± 0.11 ‰) at TS and between -0.26 ‰ and -0.06 ‰ (mean of -0.20 ± 0.06 ‰) at CR (Fig. 4). Even isotope anomaly for periphytophagous fish, $\Delta^{200}\text{Hg}$, varies between -0.09 ‰ and -0.04 ‰ (mean of -0.06 ± 0.02 ‰) at TS and between -0.09 ‰ and -0.01 ‰ (mean of -0.05 ± 0.03 ‰) at CR. $\Delta^{200}\text{Hg}$ values for piscivorous vary between -0.08 ‰ and -0.04 ‰ (mean of -0.06 ± 0.01 ‰) at TS and between -0.07 ‰ and -0.02 ‰ (mean of -0.04 ± 0.02 ‰) at CR (Table S1).

Despite the lower number of samples at CR, we investigate the $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$ regression slope for each fish feeding pattern in both pristine and gold-mining areas. Slopes of the regression lines are 1.31 ± 0.06 ($r^2 = 0.96$, $n = 16$, 1SD) and 1.37 ± 0.09 ($r^2 = 0.95$, $n = 15$, 1SD) for periphytophagous and piscivorous respectively at TS. In the gold-mining area of CR, the regression line slopes are 1.61 ± 0.34 ($r^2 = 0.85$, $n = 6$, 1SD) and 0.84 ± 0.12 ($r^2 = 0.86$, $n = 10$, 1SD) for periphytophagous and piscivorous fish respectively. Slopes for periphytophagous fish between TS and CR are not significantly different (ANCOVA test, $p = 0.22$) while slopes for piscivorous are significantly different (ANCOVA test, $p = 0.002$).

Discussion

4.1 $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ differences between both study locations

The significantly higher $\delta^{15}\text{N}$ values measured in periphytophagous fish from TS compared to CR could be explained by difference in the composition of biofilm between both locations. The significantly higher (but slight from an ecological point of view) $\delta^{15}\text{N}$ values measured in periphytophagous fish from TS compared to CR could be explained by several environmental parameters specific to each site. It could be due, for example, to the difference in the composition and age of the biofilm between both locations (chlorophyll a and a/b ratio in periphyton). A variation in periphyton composition could be explained by a difference in transparency of water due to the

occurrence of suspended particles (which could be linked to the gold mining activity or the soil use) or to the difference in canopy cover over the river (Oyapock river at TS receives less light because of a higher vegetation cover than the Camopi river at the sampling sites). Another explanation for the difference in $\delta^{15}\text{N}$ measured values in periphytophagous fish between both sites could be linked to the difference in food resources accessibility which may be due to the variation in the vegetation density or in the grazing intensity but also to the different predation pressure on these fish species.

Goix et al. (2019) observed that sediments in TS contained more organic carbon but also more MeHg (in %) than in CR. On the contrary, we observed that %MeHg in periphytophagous fish species is generally lower but more variable at TS than at CR. This suggests that there is not direct exposure route from methylmercury contained in sediments, and that periphytophagous fish contamination is mainly due to trophic transfer. Moreover, the large variation of $\delta^{13}\text{C}$ values of periphytophagous fish at both locations suggests multiple C sources which will need to be investigated.

4.2 Total mercury concentrations in fish

Average total Hg concentration in fish muscles increased according to the trophic levels. In herbivorous fish species studied, THg concentrations are 125 times lower compared with piscivorous fish species levels. Indeed, herbivorous fish eat terrestrial plants (Planquette et al. 1996) which contain low levels of Hg and MeHg. Terrestrial plants absorb Hg from substrate soil (via the roots) and the atmosphere (via the leaves) which are reservoirs of mainly inorganic Hg species. THg concentrations in the herbivorous fish studied are, thus, much lower than herbivorous fish eating aquatic plants species (Roulet and Maury-Brachet, 2001).

Periphytophagous fish have higher THg concentrations at CR than at TS. As explained in the methods section, periphytophagous fish ingest a mix of bacteria, algae and fungi contained in the biofilm. Inorganic Hg in the biofilm is methylated into MeHg subsequently absorbed and bioaccumulated by periphytophagous fish (Guimarães et al., 1998; Acha et al. 2005). The higher THg concentration measured in the periphytophagous fish could result from higher MeHg concentrations in biofilms at CR (Le Bail et al., 2000). In our previous study, we measured higher THg concentration in

sediments at CR than at TS. We hypothesize that this THg is first desorbed from sediments and then reached the water column in dissolved form, methylated into the biofilm after its sorption and then ingested by fish (Laperche et al., 2007, Goix et al., 2019).

Piscivorous fish had the same THg concentrations at CR than at TS (Wilcoxon, $W = 65$, $p = 0.094$), while CR is greatly Hg-impacted by goldmining activities. This could not be explained by different size ranges of fish samples (average of 532 ± 103 cm ($n = 19$) and 490 ± 68 cm ($n = 11$) at TS and CR, respectively), nor to differences in their trophic position between both sites. However, the distribution of prey fish depends on several factors such as hydrological characteristics (river size, turbidity and flow, season, etc.), local ecological factors and variability, and fisheries or gold-mining pressures (Maury-Brachet et al., 2020).

4.3 Hg isotope signatures in relation to the feeding pattern of the fish species

Very few Hg isotopic studies on river fish flesh have been published. In the Bolivian Amazon, Laffont et al. (2009) analyzed fish from the Beni river and its floodplain lakes. Three other studies have been conducted on river fish by Janssen et al. (2019) in Northeastern USA streams, Donovan et al. (2016) in the Yuba river (California, USA) and Tsui et al. (2012) in the Eel river (California, USA).

The mass dependent isotopic signature, $\delta^{202}\text{Hg}$, for all fish from this study varies between -1.68 ‰ and -0.89 ‰. This signature is more negative than the ones observed for river fish in literature including the one in the Bolivian Amazon basin. However, $\delta^{202}\text{Hg}$ in sediments is also more negative than the ones in the literature cited in the precedent paragraph. The average Hg isotopic signature from pristine sediments of the Oyapock R. (Goix et al., 2019) were $\delta^{202}\text{Hg} = -2.27$ ‰, $\Delta^{199}\text{Hg} = -0.66$ ‰ and $\Delta^{200}\text{Hg} = -0.04$ ‰ ($n = 27$). Negative $\delta^{202}\text{Hg}$, $\Delta^{199}\text{Hg}$ and $\Delta^{200}\text{Hg}$ of pristine sediments and soils reflected dry deposition of atmospheric mercury mostly after vegetation uptake by plants and subsequent transfer to the river system (Goix et al., 2019). The MIF of even isotopes is commonly associated with either dry or wet atmospheric deposition that display negative and positive $\Delta^{200}\text{Hg}$, respectively (Fu et al., 2016 and Obrist et al., 2017). Average $\Delta^{200}\text{Hg}$ anomaly close to -0.05 ‰ measured in fish from the Oyapock basin indicated that Hg sources for fish is atmospheric Hg^0 as observed for sediments. It could be Hg desorbed and dissolved from sediments and also from

vegetation (foliage/litter) inputs into the water column (Jiskra et al., 2017; Woerndle et al., 2018). Recent studies monitoring the export of Hg via streamflow in forested ecosystems have also shown that the majority of Hg exported was derived from previously dry deposited Hg in the forested watersheds, as their isotopic signatures resemble those of foliage/litter and forest floor rather than direct precipitation (Tsui et al., 2020).

Following Janssen et al. (2019), Tsui et al. (2012) and Donovan et al. (2016), we applied a $\delta^{202}\text{Hg}$ correction to calculate the $\delta^{202}\text{Hg}$ before isotopic fractionation induced by photochemistry. This calculation has been first made by Gehrke et al. (2011) using experimental relationships to subtract the known MDF and MIF that occurs during photochemical degradation and calculating $\Delta^{199}\text{Hg}/\delta^{202}\text{Hg}$ slope to obtain an estimated $\delta^{202}\text{Hg}$ value for prephotodegraded MeHg and IHg. Actually, we used Bergquist and Blum (2007) experimental data: photoreduction of mercury with 1 mg/L C (slope 1.15) and photodemethylation of MeHg with 1 mg/L C (slope 2.43) (average 2.7 mg.L^{-1} ($n = 16$) of DOC was measured in the study rivers (database of the ANR-RIMNES Research Program)). Assuming that mercury in water column of TS and CR came mainly from sediments, pre-photodegraded mercury has $\Delta^{199}\text{Hg}$ close to the $\Delta^{199}\text{Hg}$ of pristine sediments, the average $\delta^{202}\text{Hg}$ of pre-photodegraded MeHg is estimated at -1.54 ‰ and -1.70 ‰ for piscivorous fish at TS and CR respectively and at -1.29 ‰ and -1.51 ‰ for periphytophagous fish at TS and CR respectively. These values are much higher than the $\delta^{202}\text{Hg}$ in the pristine sediment ($\delta^{202}\text{Hg} = -2.27 \text{ ‰}$), suggesting there is a positive offset of around +0.7 ‰ (apart from periphytophagous fish from TS) between sediments and fish, consistent with Janssen et al. (2019), Gehrke et al. (2011) and Tsui et al. (2012) observations. This offset is the net result of biotic MDF during IHg methylation (biotic methylation, biotic MeHg degradation, etc.).

The odd-isotopes anomalies of Hg of periphytophagous fish reflect the isotopic signature of dissolved MeHg. Indeed, the slopes ($\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$) close to 1.3 have been interpreted as the result of partly photodegraded MeHg in river water, bioaccumulated by fish (Bergquist and Blum 2007). But the latter slope close to 1 is more likely due to photodegradation of IHg prior to its methylation and ingestion by fish (Bergquist and Blum 2007). The offset of around -0.2 ‰ observed in $\Delta^{199}\text{Hg}$ between piscivorous and periphytophagous fish for each location could indicate that sources of this MeHg are

different for the two fish diets: piscivorous fish are opportunistic hunters and integrate the Hg signature of the entire environment while periphytophagous are specific biofilm consumers.

Moreover, odd-isotopes anomalies observations point out the fact that both areas have different aquatic processes affecting Hg: $\Delta^{199}\text{Hg}$ is more positive for periphytophagous fish at CR and $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$ regression slope at CR is close to 1 for piscivorous fish. Tsui et al. (2012) observed the same slope for terrestrial predators. According to $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values which are similar between two areas for piscivorous fish, we could suppose that their food sources are quite the same. These observations can only be explained by differences in photochemistry processes between both areas induced by a different water composition, sunlight exposure (canopy coverage), river flow, and others parameters of water dynamic and chemistry.

Consequently, Hg isotopic signatures in fish are affected by the difference of biotic (methylation/demethylation) and abiotic (photochemistry) processes between both areas. We suppose that both aquatic systems are slightly different as explained before and, moreover, disturbed by gold-mining activities by inputs of suspended particles.

Conclusion

In periphytophagous fish species, THg concentrations were higher in the gold-mining (CR) than in the pristine (TS) area whereas $\delta^{13}\text{C}$ values were enriched and $\delta^{15}\text{N}$ depleted in CR compared to TS. As these fish exclusively eat biofilm settled on immersed rocks, adventive roots or plant residues, we hypothesize that this difference is due to a different composition of the biofilm between two sites. We showed that Hg isotopic signatures in fish caught in pristine and gold-mined basins have undergone different speciation processes. Variations in Hg isotopes fractionation are attributed to: i) the difference in biotic processes and MeHg sources shown by variations of $\delta^{202}\text{Hg}$, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in both locations, and ii) variation of abiotic processes such as photochemistry between both areas shown by the difference of odd-isotopes anomalies and $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$ regression slope between fish species and locations.

To conclude, Hg isotopic signatures in fish are affected by the difference of biotic (methylation/demethylation) and abiotic (photochemistry) processes between both areas and did therefore not allow to resolve the contribution of gold-mining related liquid Hg(0) in fish tissues. Mercury isotopes of MeHg in fish and lower trophic level organisms can be complementary to light stable isotope tracers.

Acknowledgements

This study was funded by the French National Research Agency (ANR-11-CESA-0013, RIMNES Program) and all the analyses were performed in two laboratories of the Midi-Pyrénées Observatory (OMP-GET and ECOLAB) in France. We would like to thank both teams of the HYDRECO laboratory and the Parc Amazonien de Guyane (PAG) for their help during sampling campaigns and the sharing of material devices. We sincerely acknowledge the indigenous communities (Wayãpi and Teko) of the study villages, the doctors from local Health Centers for their kind cooperation during the field work, and the fishermen for sharing their knowledge of the territory. We also thank Marc Pouilly and Fabrice Duponchelle from the French Research Institute for the Development (IRD) for their helpful comments on Amazonian fish behavior and diet. J. Chmeleff is thanked for expert management of the OMP mass spectrometry division in the GET laboratory.

Authors' contribution

LM was the coordinator of the RIMNES Research Program (funded by the French ANR). LM and RM-B designed the study dedicated to the ichthyologic part of the program. Formal analysis (THg, speciation and Hg stable isotopes) and geochemical lab work were performed by LL, JM and SGx under the supervision of JS. RR helped with logistics in French Guiana. Fish sampling was realized by RM-B, AL and PG. Statistical analysis were performed by LL and SGx. Funds were acquired by LM and RR. The original draft was written by LL, SGx, JM and LM and revised and edited by all authors.

Funding: Open access funding provided by the IRD (French Research Institute for Sustainable Development). This work was supported by the French National Research Agency (ANR-11-CESA-0013, RIMNES Program).

Data availability: the online version contains the raw data available in the supplementary material.

Compliance with ethical standards

Ethical approval: Not applicable.

Consent to participate: Not applicable.

Consent to publish: Not applicable.

Competing interests: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Acha D., Iñiguez V., Roulet M., Guimarães J. R. D., Luna R., Alanoca L. and Sanchez S. (2005). Sulfate-reducing bacteria in floating macrophyte rhizospheres from an Amazonian floodplain lake in Bolivia and their association with Hg methylation. *Applied and Environmental Microbiology* 71: 7531-7535

AMAP/UN Environment. (2019). Technical Background Report for the Global Mercury Assessment 2018. Arctic Monitoring and Assessment Programme, Oslo, Norway/UN Environment Programme, Chemicals and Health Branch, Geneva, Switzerland.

Bergamino, L., Lercari, D., Defeo, O. (2011). Food web structure of sandy beaches: Temporal and spatial variation using stable isotope analysis. *Estuar. Coast. Shelf Sci.* 91(4): 536-543

Bergquist B. A., and Blum J. D. (2007). Mass-Dependent and -Independent Fractionation of Hg Isotopes by Photoreduction in Aquatic Systems. *Science* 318: 417–420

Bergquist B. A., and Blum J. D. (2009). The Odds and Evens of Mercury Isotopes: Applications of Mass-Dependent and Mass-Independent Isotope Fractionation. *Elements* 5: 353–357

Blum J. D. (2011). Applications of stable mercury isotopes to biogeochemistry. In *Handbook of Environmental Isotope Geochemistry*. Springer 229–245

Blum J. D. and Bergquist B. A. (2007). Reporting of variations in the natural isotopic composition of mercury. *Anal. Bioanal. Chem.* 388: 353–359

- Blum J. D. Sherman L. S. and Johnson M. W. (2014). Mercury Isotopes in Earth and Environmental Sciences. *Annu. Rev. Earth Planet. Sci.* 42: 249–269
- Cabana G. and Rasmussen, J. B. (1994). Modelling food chain structure and contaminant bioaccumulation using stable nitrogen isotopes. *Nature* 372 (6503): 255.
- Caut S., Angulo E. and Courchamp F. (2009). Variation in discrimination factors ($\Delta^{15}\text{N}$ and $\Delta^{13}\text{C}$): the effect of diet isotopic values and applications for diet reconstruction. *Journal of Applied Ecology* 46 (2): 443–453
- Chen C. Y., Borsuk M.E., Bugge D. M., Hollweg T., Balcom P. H., Ward D. M., Williams J. and Mason R. P. (2014). Benthic and Pelagic Pathways of Methylmercury Bioaccumulation in Estuarine Food Webs of the Northeast United States. *PLoS ONE* 9, e89305.
- Cordier S., Grasmick C., Paquier-Passelaigue M., Mandereau L., Weber J.-P. and Jouan M. (1998). Mercury Exposure in French Guiana: Levels and Determinants. *Arch. Environ. Health Int. J.* 53: 299–303
- Cordier S., Garel M., Mandereau L., Morcel H., Doineau P., Gosme-Seguret S., Josse D., White R. and Amiel-Tison C. (2002). Neurodevelopmental Investigations among Methylmercury-Exposed Children in French Guiana. *Environ. Res.* 89: 1–11
- Demers J. D., Blum J. D. and Zak D. R. (2013). Mercury isotopes in a forested ecosystem: Implications for air-surface exchange dynamics and the global mercury cycle. *Global Biogeochem. Cy.* 27 (1): 222–238.
- Demers J. D., Sherman L. S., Blum J. D., Marsik F. J. and Dvornch, J. T. (2015). Coupling atmospheric mercury isotope ratios and meteorology to identify sources of mercury impacting a coastal urban/industrial region near Pensacola, Florida, USA. *Global Biogeochem. Cy.* 29 (10): 1689–1705.
- Donovan P., Blum J. D., Bliss Singer M., Marvin-DiPasquale M. and Tsui M. T. K. (2016). Isotopic composition of inorganic mercury and methylmercury downstream of a historical gold-mining region. *Environ. Sci. Technol.* 50: 1691–1702
- Dominique Y., Muresan B., Duran R., Richard S. and Boudou A. (2007). Simulation of the Chemical Fate and Bioavailability of Liquid Elemental Mercury Drops from Gold Mining in Amazonian Freshwater Systems. *Environ. Sci. Technol.* 41: 7322–7329
- Dominique Y., Maury-Brachet R., Muresan B., Vigouroux R., Richard S., Cossa D., Mariotti A. and Boudou A. (2007). Biofilm and mercury availability as key factors for mercury accumulation in fish (*Curimata cyprinoides*) from a disturbed Amazonian freshwater system. *Environmental toxicology and chemistry/SETAC* 26: 45–52.
- Dominique Y. (2008). Contamination par les différentes formes chimiques du mercure de la composante biologique du barrage hydroélectrique de Petit-Saut et des zones amont/aval du fleuve Sinnamary, en Guyane française (études in situ et approches expérimentales)

- Durrieu G., Maury-Brachet R. and Boudou A. (2005). Goldmining and mercury contamination of the piscivorous fish *Hoplias aimara* in French Guiana (Amazon basin). *Ecotoxicol. Environ. Saf.* 60: 315–323
- Enrico M., Le Roux G., Maruszczak N., Heimbürger L-E., Claustres A., Fu X., Sun R. and Sonke J. E. (2016). Atmospheric mercury transfer to peat bogs dominated by gaseous elemental mercury dry deposition. *Environ. Sci. Technol.* 50: 2405–2412
- Estrade N., Carignan J., Sonke J. E. and Donard, O. F. X. (2009). Mercury isotope fractionation during liquid–vapor evaporation experiments. *Geochim. Cosmochim. Acta* 73: 2693–2713
- Fleming E. J., Mack E. E., Green P. G. and Nelson D. C. (2006). Mercury Methylation from Unexpected Sources: Molybdate-Inhibited Freshwater Sediments and an Iron-Reducing Bacterium. *Appl. Environ. Microbiol.* 72: 457–464
- Fréry N., Maury-Brachet R., Maillot E., Deheeger M., De Merona B. and Boudou A. (2001). Gold-mining activities and mercury contamination of native Amerindian communities in French Guiana: key role of fish in dietary uptake. *Environ. Health Perspect.* 109: 449
- Fu X., Maruszczak N., Wang X., Gheusi F. and Sonke, J. E. (2016). Isotopic composition of gaseous elemental mercury in the free troposphere of the Pic du Midi observatory, France. *Environ. Sci. Technol.* 50 (11): 5641–5650
- Gantner N., Hintelmann H., Zheng W. and Muir D. C. (2009). Variations in stable isotope fractionation of Hg in food webs of Arctic lakes. *Environ. Sci. Technol.* 43: 9148–9154
- Gentès S., Coquery M., Vigouroux R., Hanquiez V., Allard L. and Maury-Brachet R. (2019). Application of the European Water Framework Directive: Identification of reference sites and bioindicator fishspecies for mercury in tropical freshwater ecosystems (FrenchGuiana). *Ecological Indicators.* 106. 105468
- Gerhke G. E., Blum J. D., Slotton D. G., Greenfield B. K. (2011). Mercury isotopes link mercury in San Francisco Bay forage fish to surface sediments. *Environ. Sci. Technol.* 45: 1264–1270
- Goix S., Maurice L., Laffont L., Rinaldo R., Lagane C., Bosq A., Chmeleff J., Menges J., Heimbürger L. E., Maury-Brachet R. and Sonke J. (2019). A Hg stable isotope study of Hg cycling in the tropical environment and quantification of the gold-mining impact on French Guiana river sediments. *Chemosphere* 219: 684–694.
- Gratz L. E., Keeler G. J., Blum J. D. and Sherman L. S. (2010). Isotopic composition and fractionation of mercury in Great Lakes precipitation and ambient air. *Environ. Sci. Technol.* 44 (20): 7764–7770
- Grimaldi C., Grimaldi M. and Guedron S. (2008). Mercury distribution in tropical soil profiles related to origin of mercury and soil processes. *Sci. Total Environ.* 401: 121–129
- Grimaldi M., Guédron S. and Grimaldi C. (2015). Impact of gold mining on mercury contamination and soil degradation in Amazonian ecosystems of French Guiana. In : Brearley F.Q. (ed.), Thomas

- A.D. (ed.) Land-use change impacts on soil processes : tropical and savannah ecosystems. Wallingford : CABI, 95-107. ISBN 978-1-78064-210-9
- Guedron S., Grimaldi C., Chauvel C., Spadini L. and Grimaldi M. (2006). Weathering versus atmospheric contributions to mercury concentrations in French Guiana soils. *Appl. Geochem.* 21: 2010–2022
- Guedron S., Grangeon S., Lanson B. and Grimaldi M. (2009). Mercury speciation in a tropical soil association; Consequence of gold mining on Hg distribution in French Guiana. *Geoderma* 153: 331–346
- Guimarães J. R. D., Meili M., Malm O. and Maria de Souza Brito E. (1998). Hg methylation in sediments and floating meadows of a tropical lake in the Pantanal floodplain, Brazil. *Sci. Total Environ.* 213: 165–175
- Hobson, K. A., Fisk, A., Karnovsky, N., Holst, M., Gagnon, J. M. and Fortier, M. (2002). A stable isotope (d13C, d15N) model for the NorthWater food web: implications for evaluating trophodynamics and the flow of energy and contaminants. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 49 (22e23): 5131e5150
- Jiskra, M., Wiederhold, J.G., Skjellberg, U., Kronberg, R.-M., Kretzschmar, R., 2017. Source tracing of natural organic matter bound mercury in boreal forest runoff with mercury stable isotopes. *Environ. Sci.: Processes Impacts* 19, 1235–1248.
- Keith, P., Persat, H., Feunteun, E. and Allardi, J. (2011). *Les Poissons d’eau douce de France*. Collection Inventaires & biodiversité. Biotope Editions, Publications scientifiques du Muséum 552 (in French)
- Kerin E. J., Gilmour C. C., Roden E., Suzuki M. T., Coates J. D. and Mason R. P. (2006). Mercury Methylation by Dissimilatory Iron-Reducing Bacteria. *Appl. Environ. Microbiol.* 72: 7919–7921
- Klaus, J. E., Hammerschmidt, C. R., Costello, D. M., Burton, G. A. (2016). Net methylmercury production in 2 contrasting stream sediments and associated accumulation and toxicity to periphyton. *Environ. Toxicol. Chem.* 35(7), 1759-1765
- Kwon S. Y., McIntyre P. B., Flecker A. S. and Campbell L. M. (2012a). Mercury biomagnification in the food web of a neotropical stream. *Sci. Total Environ.* 417-418: 92–97
- Kwon S. Y., Blum J. D., Carvan M. J., Basu N., Head J. A., Madenjian C. P. and David S. R. (2012b). Absence of Fractionation of Mercury Isotopes during Trophic Transfer of Methylmercury to Freshwater Fish in Captivity. *Environ. Sci. Technol.* 46: 7527–7534
- Laffont L., Sonke J. E., Maurice L., Hintelmann H., Pouilly M., Sánchez Bacarreza Y., Perez T. and Behra P. (2009). Anomalous mercury isotopic compositions of fish and human hair in the Bolivian Amazon. *Environ. Sci. Technol.* 43: 8985–8990

712 Laffont L., Sonke J. E., Maurice L., Monrroy S. L., Chincheros J., Amouroux D. and Behra P. (2011).
 713 Hg Speciation and Stable Isotope Signatures in Human Hair As a Tracer for Dietary and Occupational
 714 Exposure to Mercury. *Environ. Sci. Technol.* 45: 9910–9916
 715
 716 Lamberti, G. A. (1996). The role of periphyton in benthic food webs. *Algal ecology*
 717
 718 Laperche V., Maury-Brachet R., Blanchard F., Dominique Y., Durrieu G., Massabuau J. C., Bouillard
 719 H., Joseph B., Laporte P., Mesmer-Dudons N., Duflo V. et Callier L. (2007). Répartition régionale
 720 du mercure dans les sédiments et les poissons de six fleuves de Guyane: Rapport BRGM/RP55965-
 721 FR, 203.
 722
 723 Layman C. A., Araujo M. S., Boucek R., Hammerschlag- Peyer C. M., Harrison E., Jud Z. R., Matich
 724 P., Rosenblatt A. E., Vaudo J. J., Yeager L. A., Post D. M. and Bearhop S. (2012). Applying stable
 725 isotopes to examine food- web structure: an overview of analytical tools. *Biological Reviews* 87 (3):
 726 545-562
 727
 728 Le Bail, P.-Y., Covain, R., Jégu, M., Fisch-Muller, S., Vigouroux R. and Keith, P. (2012). Updated
 729 checklist of the freshwater and estuarine fishes of French Guiana. *Cybum* 36 (1): 293–319.
 730
 731 Le Bail, P. Y., Keith, P. and Planquette, P. (2000). Atlas des poissons d’eau douce de Guyane ; Tome
 732 2, fascicule II: Siluriformes. Patrimoines naturels (MNHN/SPN), 43, 307. Paris (in French).
 733
 734 Lee, C. C. (2005). *Environmental Engineering Dictionary* (Government Institutes)
 735
 736 Lepak R., Yin R., Krabbenhoft D. P., Ogorek J. M., DeWild J. F., Holsen T. M. and Hurley J. P.
 737 (2015). Use of stable isotope signatures to determine mercury sources in the Great Lakes. *Environ.*
 738 *Sci. Technol. Lett.* 2 (12): 335–341
 739
 740 Marshall B. G., Veiga M. M., Kaplan R. J., Adler Miserendino R., Schudel G., Bergquist B. A.,
 741 Guimarães J. R. D., Sobral L. G. S. and Gonzalez-Mueller C. (2018). Evidence of transboundary
 742 mercury and other pollutants in the Puyango-Tumbes River basin, Ecuador–Peru. *Environ. Sci.:*
 743 *Processes Impacts.* 20: 632-641
 744
 745 Masbou J. (2014). Etude des processus métaboliques, écologiques et biogéochimiques contrôlant le
 746 fractionnement isotopique du Hg chez les mammifères marins de l’Arctique. PhD dissertation,
 747 Université Toulouse Paul Sabatier, pp 275
 748
 749 Masbou J., Point D. and Sonke J. E. (2013). Application of a selective extraction method for
 750 methylmercury compound specific stable isotope analysis (MeHg-CSIA) in biological materials. *J.*
 751 *Anal. At. Spectrom.* 28: 1620
 752
 753 Mason R. P., Reinfelder J. R. and Morel F. M. M. (1995). Bioaccumulation of mercury and
 754 methylmercury. *Water. Air. Soil Pollut.* 80: 915–921
 755
 756 Maurice L., Goix S., Laffont L., Sonke J. E., Maury-Brachet R., Lagane C., Bosq A., Candaudap F.
 757 and Rinaldo R. (2014). Hg stable isotope as a tracer of gold mining activities in the Oyapock river
 758 basin, French Guiana. In: 17th ICHMET International Conference on Heavy Metals in the
 759 Environment, Sept. 2014, Guiyang, China.

- Maurice-Bourgoin L., Quiroga I., Chincheros J. and Courau P. (2000). Mercury distribution in waters and fishes of the upper Madeira rivers and mercury exposure in riparian Amazonian populations. *Sci. Total Environ.* 260: 73-86
- Maurice-Bourgoin L., Quiroga I., Guyot I. L. and Malm O. (1999). Mercury Pollution in the Upper Beni River Basin Bolivia. *Ambio* 28: 302–306
- Maury-Brachet R., Durrieu G., Dominique Y. and Boudou A. (2006). Mercury distribution in fish organs and food regimes: Significant relationships from twelve species collected in French Guiana (Amazonian basin). *Sci. Total Environ.* 368: 262–270
- Maury-Brachet R., Gentes S., Dassié E. P., Feurtet-Mazel A., Vigouroux R., Laperche V., Gonzalez P., Hanquiez V., Mesmer-Dudons N., Durrieu G. and Legeay A. (2020). Mercury contamination levels in the bioindicator piscivorous fish *Hoplias aimara* in French Guiana rivers: mapping for risk assessment. *Environ Sci Pollut Res* 27: 3624–3636
- Meech J. A., Veiga M. M. and Tromans D. (1997). Emission and Stability of Mercury in the Amazon. *Can. Metall. Q.* 36: 231–239
- Mergler, D.; Anderson, H. A.; Chan, L. H. M.; Mahaffey, K. R.; Murray, M.; Sakamoto, M.; Stern, A. H. (2007). Methylmercury exposure and health effects in humans: A worldwide concern. *Ambio* 36: 3-11
- Monperrus M., Rodriguez-Gonzalez P., Amouroux D., J. Garcia Alonso I. and Donard O. F. X. (2008). Evaluating the Potential and Limitations of Double-Spiking Species-Specific Isotope Dilution Analysis for the Accurate Quantification of Mercury Species in Different Environmental Matrices. *Analytical and Bioanalytical Chemistry* 390 (2): 655-666
- De Morais L. T. and Raffray J. (1999). Movements of *Hoplias aimara* during the filling phase of the Petit-Saut dam, French Guyana. *J. Fish Biol.* 54: 627–635
- Navarro P., Clémens S., Perrot V., Bolliet V., Tabouret H., Guérin T., Monperrus M. and Amouroux D. (2011). Simultaneous determination of mercury and butyltin species using a multiple species-specific isotope dilution methodology on the European, *Anguilla anguilla* glass eel and yellow eel. *International Journal of Environmental Analytical Chemistry*. 1-17
- Neres- Lima V., Machado- Silva F., Baptista D. F., Oliveira R. B. S., Andrade P. M., Oliveira A. F., Sasada- Sato C. Y., Silva- Junior E. F., Feijó- Lima R., Angelini R., Camargo P. B. and Moulton T.P. (2017). Allochthonous and autochthonous carbon flows in food webs of tropical forest streams. *Freshwater Biology* 62 (6): 1012-1023
- Obrist D., Agnan Y., Jiskra M., Olson C. L., Colegrove D. P., Hueber J., Moore C. W., Sonke J. E. and Helmig D. (2017). Tundra uptake of atmospheric elemental mercury drives Arctic mercury pollution. *Nature*. 547 (7662): 201–204

- Pacyna E. G., Pacyna J. M., Sundseth K., Munthe J., Kindbom K., Wilson S., Steenhuisen F. and Maxson P. (2010). Global emission of mercury to the atmosphere from anthropogenic sources in 2005 and projections to 2020. *Atmos. Environ.* 44: 2487–2499
- PAG, Suivi environnemental des impacts de l'orpaillage illégal. 2017 Bulletin n°5 5p.
- Perrot V., Epov V. N., Pastukhov M. V., Grebenshchikova V. I., Zouiten C., Sonke J. E., Husted S., Donard O. F. X. and Amouroux D. (2010). Tracing sources and bioaccumulation of mercury in fish of Lake Baikal- Angara River using Hg isotopic composition. *Environ. Sci. Technol.* 44: 8030–8037
- Perrot V., Pastukhov M. V., Epov V. N., Husted S., Donard O. F. X. and Amouroux D. (2012). Higher Mass-Independent Isotope Fractionation of Methylmercury in the Pelagic Food Web of Lake Baikal (Russia). *Environ. Sci. Technol.* 46: 5902–5911
- Peterson, B. J., Fry, B. (1987). Stable isotopes in ecosystem studies. *Annu. Rev. Ecol. Syst.* 18: 293-320
- Pfeiffer W. C. and de Lacerda L. D. (1988). Mercury inputs into the Amazon Region, Brazil. *Environ. Technol. Lett.* 9: 325–330
- Planquette P., Keith P. and Le Bail P-Y. (1996). Atlas des poissons d'eau douce de Guyane. Museum d'Histoire Naturelle. Paris
- Post D. M. (2002). Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83: 703–718
- Post, D. M., Layman, C. A., Arrington, D. A., Takimoto, G., Quattrochi, J., Montana, C. G. (2007). Getting to the fat of the matter: models, methods and assumption for dealing with lipids in stable isotope analyses. *Oecologia*. 152: 179-189
- Pouilly M., Rejas D., Perez T., Duprey J-L., Molina C. I., Hubas C. and Guimarães J. R. D. (2013). Trophic structure and mercury biomagnification in tropical fish assemblages, Itenez River, Bolivia. *PLoS ONE* 8(5): e65054
- Rahm M., Thibault P., Shapiro A., Smartt T., Paloeng C., Crabbe S., Farias P., Carvalho R., Joubert P. (2017). Monitoring the impact of gold mining on the forest cover and freshwater in the Guiana Shield. Reference year 2015. REDD+ for the Guiana Shield Project and WWF Guianas. Report pp.20
- Rimbaud D., Restrepo M., Louison A., Boukhari R., Ardillon V., Carles G., Lambert V. and Jolivet A. (2017). Blood lead levels and risk factors for lead exposure among pregnant women in western French Guiana: the role of manioc consumption. *Journal of Toxicology and Environmental Health, Part A*, DOI:10.1080/15287394.2017.1331490
- Roulet M., Lucotte M., Canuel R., Rheault I., Tran S., De Freitas Gog Y.G., Farella N., Souza do Vale R., Sousa Passos C. J., Jesus da Silva E., Mergler D. and Amorim M. (1998). Distribution and partition of mercury in waters of Tapajos. *Sci. Total Environ.* 213: 203–2011

- Roulet M., Lucotte M., Guimarães J. R. D. and Rheault I. (2000). Methylmercury in water, seston, and epiphyton of an Amazonian river and its floodplain, Tapajos River, Brazil. *Sci. Total Environ.* 261: 43–59
- Roulet M., Guimaraes J. R. D. and Lucotte M. (2001). Methylmercury production and accumulation in sediments and soils of an Amazonian floodplain—effect of seasonal inundation. *Water. Air. Soil Pollut.* 128: 41–60
- Roulet M. and Maury-Brachet R. (2001). Le mercure dans les organismes aquatiques amazoniens. In *Le mercure en Amazonie*. IRD Editions
- Senn D. B., Chesney E. J., Blum J. D., Bank M. S., Maage A. and Shine J. P. (2010). Stable Isotope (N, C, Hg) Study of Methylmercury Sources and Trophic Transfer in the Northern Gulf of Mexico. *Environ. Sci. Technol.* 44: 1630–1637
- Smith R. S., Wiederhold J. G., Jew A. D., Brown G. E., Bourdon B. and Kretzschmar R. (2014). Small-scale studies of roasted ore waste reveal extreme ranges of stable mercury isotope signatures. *Geochim. Cosmochim. Acta* 137: 1–17
- Sonke, J. E., (2011). A global Hg isotope box-model for mass-independent Hg isotope fractionation. . *Geochim. Cosmochim. Acta* 75: 4577-4590
- Sonke, J. E. and Blum, J. D. (2013). Advances in mercury stable isotope biogeochemistry. *Chemical Geology* 336: 1-4
- Telmer K., Costa M., Simões Angélica R., Araujo E. S. and Maurice Y. (2006). The source and fate of sediment and mercury in the Tapajós River, Pará, Brazilian Amazon: Ground- and space-based evidence. *J. Environ. Manage.* 81: 101–113
- Tsui M. T. K., Blum J. D., Kwon S. Y., Finlay J. C., Balogh S. J. and Nollet Y. H. (2012). Sources and Transfers of Methylmercury in Adjacent River and Forest Food Webs. *Environ. Sci. Technol.* 46: 10957–10964.
- Tsui, M.T.K., Blum, J.D., Finlay, J.C., Balogh, S.J., Kwon, S.Y., Nollet, Y.H., 2013. Photodegradation of methylmercury in stream ecosystems. *Limnol. Oceanogr.* 58, 13–22.
- Tsui M. T. K., Blum J. D. and Kwon S. Y. 2020. Review of stable mercury isotopes in ecology and biogeochemistry. *Science of the Total Environment.* 716: 135386
- US Environmental Protection Agency. (2012). <https://www.epa.gov/fish-tech/epa-fda-fish-advice-technical-information>
- Vander Zanden, J. and Rasmussen, J.B. (1996). A trophic position model of pelagic food webs: impact on contaminant bioaccumulation in lake trout. *Ecol. Monogr.* 66 (4): 451e477
- Velásquez-López, P. C. (2010). Mercury in artisanal and small scale gold mining: identifying strategies to reduce environmental contamination in Southern Ecuador. University of British Columbia. doi:10.14288/1.0071256

901
 902 WHO. (1990). Environmental health criteria 101.
 903 <http://www.inchem.org/documents/ehc/ehc/ehc101.htm>, 02.12.2014
 904
 905 Woerndle, G.E., Tsui, M.T.K., Sebestyen, S.D., Blum, J.D., Nie, X., Kolka, R.K., 2018. New insights
 906 on ecosystem mercury cycling revealed by stable isotopes of mercury in water flowing from a
 907 headwater peatland catchment. *Environ. Sci. Technol.* 52, 1854–1861.
 908
 909 World Wildlife Founds. (2018). Lutte contre l'Orpillage Illégal en Guyane; orientations pour une
 910 efficacité renforcée. Analyse WWF France, bureau Guyane.
 911 [https://www.wwf.fr/sites/default/files/doc-](https://www.wwf.fr/sites/default/files/doc-201806/180613_Rapport_Lutte_orpillage_illegal_Guyane.pdf)
 912 [201806/180613_Rapport_Lutte_orpillage_illegal_Guyane.pdf](https://www.wwf.fr/sites/default/files/doc-201806/180613_Rapport_Lutte_orpillage_illegal_Guyane.pdf)
 913
 914 Yin R., Feng X., Hurley J. P., Krabbenhoft D. P., Lepak R. F., Kang S., Yang H. and Li X. (2016).
 915 Historical records of mercury stable isotopes in sediments of Tibetan Lakes. *Sci. Rep.* 6: 23332