



HAL
open science

On the semi-positive definition of the product of a positive diagonal matrix and a symmetric semi-positive definite matrix

Noelie Ramuzat

► **To cite this version:**

Noelie Ramuzat. On the semi-positive definition of the product of a positive diagonal matrix and a symmetric semi-positive definite matrix. 2022. hal-03434195v2

HAL Id: hal-03434195

<https://hal.science/hal-03434195v2>

Preprint submitted on 4 Jan 2022 (v2), last revised 7 Jan 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the semi-positive definition of the product of a positive diagonal matrix and a symmetric semi-positive definite matrix

Noëlie Ramuzat - LAAS CNRS

January 2021

Problem statement: We have Λ a real symmetric semi-positive definite matrix and D a real positive diagonal matrix (thus also symmetric). The product ΛD is a square matrix but non-symmetric. We want to prove that ΛD quadratic form, $x^T \Lambda D x$, is semi-positive definite, i.e. $x^T \Lambda D x \geq 0 \forall x \in \mathbb{R}^n \setminus \{0\}$.

If the elements of the diagonal D are equal, i.e. $D = aI$ with $a \in \mathbb{N}$, then the product $\Lambda D = a\Lambda$ is a real symmetric semi-positive definite matrix. Thus, for a diagonal D with equal elements we have directly $x^T \Lambda D x \geq 0 \forall x \in \mathbb{R}^n \setminus \{0\}$.

For different elements on the diagonal, we have the following results:

- The eigenvalues of ΛD are real and non-negatives
- If $\frac{\Lambda D + (\Lambda D)^T}{2}$ has non-negative eigenvalues, then the quadratic form of ΛD is semi-positive definite

Proof. Let us recall some useful matrix definitions and properties:

Definition 1 (Real symmetric semi-positive definite matrix). *A is a real symmetric matrix (its eigenvalues are thus real): A is semi-positive definite \iff all its eigenvalues are non-negatives.*

Definition 2 (Quadratic form semi-positive definition). *A matrix A is semi-positive definite $\iff x^T A x \geq 0, \forall x \in \mathbb{R}^n \setminus \{0\}$*

Property 1. *If A is a real semi-positive definite matrix, then $B^T A B$ is semi-positive definite for any matrix B.*

Property 2 (Matrix congruent to a symmetric matrix). *Any matrix congruent to a symmetric matrix is again symmetric: If A is a symmetric matrix then so is $B^T A B$ for any matrix B.*

Proof that the eigenvalues of ΛD are real and non-negatives

Because D is diagonal and positive we can write $D = D^{\frac{1}{2}}D^{\frac{1}{2}}$, $D^{\frac{1}{2}}$ is also real positive and symmetric thus invertible. Let us reformulate the matrix ΛD :

$$\begin{aligned}\Lambda D &= \Lambda D^{\frac{1}{2}}D^{\frac{1}{2}} \\ &= D^{-\frac{1}{2}}(D^{\frac{1}{2}}\Lambda D^{\frac{1}{2}})D^{\frac{1}{2}}\end{aligned}\tag{1}$$

It corresponds to a change of basis of $D^{\frac{1}{2}}$. Because the eigenvalues (denoted λ) are invariant to change of basis we have:

$$\lambda(\Lambda D) = \lambda(D^{\frac{1}{2}}\Lambda D^{\frac{1}{2}})\tag{2}$$

Using the Properties. 1 and 2, because Λ is a real symmetric semi-positive definite matrix, $(D^{\frac{1}{2}})^T\Lambda D^{\frac{1}{2}} = D^{\frac{1}{2}}\Lambda D^{\frac{1}{2}}$ is a real symmetric semi-positive definite matrix. Thus its eigenvalues are real and non-negatives (Definition.1) and so are the ones of ΛD because of Eq.2. Then we have proven our first result: the eigenvalues of ΛD are real and non-negatives.

Semi-positive definition of ΛD

In this part we look at a way to prove the semi-positive definition of ΛD by studying its symmetric part in the Toeplitz decomposition:

Definition 3 (Toeplitz decomposition). *Every square matrix A can be decomposed uniquely as the sum of two matrices U and V , where U is symmetric and V is skew-symmetric.*

$$A = U + V = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T)\tag{3}$$

In our case $A = \Lambda D$ and $U = \frac{1}{2}(\Lambda D + (\Lambda D)^T)$ is symmetric.

We recall the fact that the quadratic form of a skew-symmetric matrix equals to zero. Indeed, by definition $V^T = -V$ and thus $x^T V x = (x^T V^T x)^T = -x^T V x$ which holds only if it equals to zero. Thus, the quadratic form of $x^T \Lambda D x$ is the same one of $x^T U x$, i.e:

$$x^T \Lambda D x = x^T \left(\frac{\Lambda D + (\Lambda D)^T}{2} \right) x\tag{4}$$

One can thus prove the semi-positive definition of the symmetric matrix U to prove the semi-positive definition of ΛD . Indeed, if $x^T U x \geq 0, \forall x \in \mathbb{R}^n \setminus \{0\}$, using Eq.4, we obtain $x^T \Lambda D x \geq 0, \forall x \in \mathbb{R}^n \setminus \{0\}$: proving of the semi-positive definition of ΛD .

One way to prove the semi-positive definition of U is to look at its eigenvalues. Because U is symmetric, if its eigenvalues are non-negatives then U is semi-positive definite (see Definition 1). This gives our second result: if U has non-negative eigenvalues, then ΛD is semi-positive definite.

On the eigenvalues of U

One may notice that we have further information on the eigenvalues of U with respect to the ones of ΛD . Using the following theorem of Fan on matrices [1] (Chapter 10, Theorem 10.28):

Theorem 1. Let A be an $n \times n$ matrix with eigenvalues $\lambda_1(A), \dots, \lambda_n(A)$ and $\mathcal{R}_e \lambda_i(A)$ their real parts. Then:

$$\sum_{i=1}^n \mathcal{R}_e \lambda_i(A) \leq \sum_{i=1}^n \lambda_i\left(\frac{A + A^T}{2}\right) \quad (5)$$

Then, because the eigenvalues of ΛD are real and non-negatives (as proven in the first part), we have using the Theorem 1:

$$0 \leq \sum_{i=1}^n \lambda_i(\Lambda D) \leq \sum_{i=1}^n \lambda_i(U) \quad (6)$$

Thus, we know that the sum of the eigenvalues of U is non-negative, however to prove the semi-positive definition of U it is needed to prove that all its eigenvalues are non-negatives. \square

References

- [1] Fuzhen Zhang. *Matrix theory: basic results and techniques*. Springer, 2011.