



# H<sup>-</sup> and D<sup>-</sup> production efficiency in a multi-dipole ECR-plasma source as a function of gas pressure

M. Mitrou, P. Svarnas, S. Béchu

## ► To cite this version:

M. Mitrou, P. Svarnas, S. Béchu. H<sup>-</sup> and D<sup>-</sup> production efficiency in a multi-dipole ECR-plasma source as a function of gas pressure. Journal of Physics: Conference Series, 2022, 2244 (1), pp.012007. 10.1088/1742-6596/2244/1/012007 . hal-03434717v1

HAL Id: hal-03434717

<https://hal.science/hal-03434717v1>

Submitted on 18 Nov 2021 (v1), last revised 17 Nov 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# H<sup>-</sup> and D<sup>-</sup> production efficiency in a multi-dipole ECR-plasma source as a function of gas pressure

M. Mitrou<sup>1,2</sup>, P. Svarnas<sup>1</sup>, S. Béchu<sup>2</sup>

<sup>1</sup>High Voltage Laboratory, Electrical and Computer Engineering Dept., University of Patras, Rion-Patras, GR-26504, Greece


<sup>2</sup>Univ. Grenoble Alpes, CNRS, Grenoble INP, LPSC-IN2P3, 38000 Grenoble, France

E-mails: [maria.mitrou@ece.upatras.gr](mailto:maria.mitrou@ece.upatras.gr), [svarnas@ece.upatras.gr](mailto:svarnas@ece.upatras.gr), [bechu@lpsc.in2p3.fr](mailto:bechu@lpsc.in2p3.fr)

## Introduction


The present work is devoted to the ECR volume source Prometheus I, extending its exploitation to D<sub>2</sub> since previously operation only with H<sub>2</sub> had been thoroughly carried out. Basic plasma properties are accessed via electrostatic probe and laser induced photo-detachment. The experiments are carried out throughout a wide range of working pressure, whereas the microwave power is maintained constant. The results are then compared and evaluated so as to identify similarities and differences between the ECR plasmas of the two isotopes.

## Experimental Setup




(a) 1/2 wave-plate on rotation mount; (b) beam splitter; (c) beam dump; (d) mirrors; (e) Ø5 mm diaphragm; (f) pyroelectric laser energy sensor


## Results




(a) Cold and (b) hot electron temperature vs. the operating gas pressure. • H<sub>2</sub>; ○ D<sub>2</sub>; 0.9 kW




(a) Cold and (b) hot electron density vs. the operating gas pressure. • H<sub>2</sub>; ○ D<sub>2</sub>; 0.9 kW



EEDFs under bi-Maxwellian assumption at (a) 0.27 Pa, (b) 1.33 Pa, and (c) 2.67 Pa. • H<sub>2</sub>; ○ D<sub>2</sub>; 0.9 kW



(a) H<sup>-</sup> & D<sup>-</sup> ion density and (b) H<sup>-</sup> & D<sup>-</sup> ion to cold electron density ratio vs. the operating gas pressure. • H<sub>2</sub>; ○ D<sub>2</sub>; 0.9 kW.



H<sup>-</sup> & D<sup>-</sup> ion density vs. (a) cold and (b) hot electron density. Data pairs are created with the pressure elimination. • H<sub>2</sub>; ○ D<sub>2</sub>; 0.9 kW.

## Conclusions

- The plasma potential, cold electron temperature, and cold electron density tend to be higher in deuterium.
- No pronounced difference in the hot electron density and temperature is found between the two plasmas.
- The two observations on the electrons are jointly mirrored on the EEDF patterns. The diverging features between the H<sub>2</sub> and D<sub>2</sub> EEDFs are more pronounced at the lower pressures.
- A similar H<sup>-</sup> and D<sup>-</sup> negative ion yield (up to  $6 \times 10^{15} \text{ m}^{-3}$ ; under the present conditions) is achieved.
- For equal plasma densities an isotope effect is exhibited showing higher H<sup>-</sup> density over the entire pressure range.
- The n<sub>H<sup>-</sup></sub> / n<sub>e</sub> ratio is constantly higher than the n<sub>D<sup>-</sup></sub> / n<sub>e</sub> one and they both peak around 1.33 Pa.

## References

- Hemsworth R S, Boilson D, Blatchford P, Dalla Palma M, Chitarin G, De Esch H P L, Geli F, Dremel M, Graceffa J, Marcuzzi D, Serianni G, Shah D, Singh M, Urbani M and Zaccaria P 2017 *New J. Phys.* **19** 025005
- Wunderlich D, Riedl R, Mano I, Mimo A, Fantz U, Heinemann B and Kraus W 2019 *Rev. Sci. Instrum.* **90** 113304
- Dougar-Jabon V D 2001 *Physica Scripta* **63** 322-325
- Friedl R, Kurutz U and Fantz U 2016 *AIP Conf. Proc.* **1869** 030022
- Kurutz U, Friedl R and Fantz U 2017 *Plasma Phys. Control. Fusion* **59** 075008
- Béchu S, Lemaire J L, Gavilan L, Aleiferis S, Shakhmatov V, Lebedev Yu A, Fombaron D, Bonny L, Menu J, Bès A, Svarnas P and de Oliveira N 2020 *J. Quant. Spectrosc. Radiat. Transf.* **257** 107325
- Aleiferis S, Tarvainen O, Svarnas P, Bacal M and Béchu S 2016 *J. Phys. D: Appl. Phys.* **49** 095203
- Aleiferis S, Svarnas P, Béchu S, Tarvainen O and Bacal M 2018 *Plasma Sources Sci. Technol.* **27** 075015
- Celiberto R, Janev R K, Laricchiuta, Capitelli M, Wadehra J M and Atems D E 2001 *At. Data Nucl. Data Tables* **77**(2) 161-213
- Janev R K, Reiter D and Samm U 2003 *Forschungszentrum Jülich JUEL* 4105
- Svarnas P, Breton J, Bacal M and Mosbach T 2006 *Rev. Sci. Instrum.* **77** 03A532