



**HAL**  
open science

## Use of complex physiological traits as ecotoxicological biomarkers in tropical freshwater fishes

Diana A Monteiro, Ana L Kalinin, F Tadeu Rantin, David Mckenzie

► **To cite this version:**

Diana A Monteiro, Ana L Kalinin, F Tadeu Rantin, David Mckenzie. Use of complex physiological traits as ecotoxicological biomarkers in tropical freshwater fishes. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*, 2021, 335 (9-10), pp.745 - 760. 10.1002/jez.2540 . hal-03438296

**HAL Id: hal-03438296**

**<https://hal.science/hal-03438296>**

Submitted on 26 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Use of complex physiological traits as ecotoxicological biomarkers in tropical**  
2 **freshwater fishes**

3

4 Diana A. Monteiro<sup>1</sup>, Ana L. Kalinin<sup>1</sup>, F. Tadeu Rantin<sup>1</sup>, David J. McKenzie<sup>1,2\*</sup>

5

6 1. Department of Physiological Sciences, Federal University of São Carlos (UFSCar),  
7 13565-905, São Carlos, São Paulo, Brazil.

8 2. UMR Marbec, Univ. Montpellier, CNRS, IRD, Ifremer, 34095 Montpellier, France

9

10

11

12

13 **Running title:** Physiological biomarkers in tropical freshwater fishes

14

15

16 \*Author to whom correspondence should be addressed: david.mckenzie@cnr.fr

17

18 **Abstract**

19 We review the use of complex physiological traits, of tolerance and performance, as  
20 biomarkers of the toxicological effects of contaminants in subtropical and tropical  
21 freshwater fishes. Such traits are growing in relevance due to climate change, as  
22 exposure to contaminants may influence the capacity of fishes to tolerate and perform  
23 in an increasingly stressful environment. We review the evidence that the critical  
24 oxygen level, a measure of hypoxia tolerance, provides a valuable biomarker of  
25 impacts of diverse classes of contaminants. When coupled with measures of  
26 cardiorespiratory variables, it can provide insight into mechanisms of toxicity. The  
27 critical thermal maximum, a simple measure of tolerance of acute warming, also  
28 provides a valuable biomarker despite a lack of understanding of its mechanistic basis.  
29 Its relative ease of application renders it useful in rapid evaluation of multiple species,  
30 and in understanding how the severity of contaminant impacts depends upon prevailing  
31 environmental temperature. The critical swimming speed is a measure of exercise  
32 performance that is widely used as a biomarker in temperate species but very few  
33 studies have been performed on subtropical or tropical fishes. Overall, the review  
34 serves to highlight a critical lack of knowledge for subtropical and tropical freshwater  
35 fishes. There is a real need to expand the knowledge base and to use physiological  
36 biomarkers in support of decision making to manage tropical freshwater fish  
37 populations and their habitats, which sustain rich biodiversity but are under relentless  
38 anthropogenic pressure.

39 **Key-words:**

40 Biomarker, critical oxygen level, critical swimming speed, critical thermal maximum,  
41 ecotoxicology, hypoxia, warming.

## 42 **Introduction**

43           Freshwater ecosystems, and their resident fish populations, are particularly at  
44 risk from negative impacts of anthropogenic global change (Mason, 2002; Costa and  
45 Barletta, 2016; Higgins et al., 2021). Freshwater ecosystems are suffering relentless  
46 contamination with a bewildering diversity of chemical hazards and toxic substances  
47 such as agricultural chemicals; industrial solvents, surfactants and flame-retardants;  
48 metals, and pharmaceutical products, among many other pollutants (Stauffer, 1998;  
49 Mason, 2002; Rantin et al., 2020). These enter waterways directly from treated and  
50 untreated industrial, agricultural and domestic wastewaters, from run-off and  
51 atmospheric deposition (including spray drift) and indirectly from leaching (Holt, 2000).  
52 The diversity of contaminants and their potential toxic effects on freshwater fishes grow  
53 incessantly (e.g. Jobling and Tyler, 2003; Pérez-Parada et al., 2018; Wagner and  
54 Lambert, 2018).

55           In aquatic organisms, biomarkers are used to provide early warning signals of  
56 exposure or effects of specific pollutants or pollutant classes (Peakall, 1994; van der  
57 Oost et al., 2003). They are defined, in a broad sense, as any biological response by  
58 an organism to exposure to environmental chemicals, or to their toxic effects. The term  
59 is most commonly used to refer to measurements in body fluids, cells or tissues, which  
60 are indicative of bioaccumulation of toxic chemicals, biochemical and cellular  
61 modifications provoked by specific toxicants (van der Oost et al., 2003; Kroon et al.,  
62 2017) but would also be pertinent for other responses, for example behavioural  
63 alterations (Little and Finger, 1990; Saaristo et al., 2018).

64           For fishes it has long been recognized that complex traits of whole animal  
65 physiology, especially traits of functional integrity and energetics, are valuable  
66 indicators of sublethal toxicological impacts of contaminants (Brett, 1958; Cairns, 1966;  
67 Sprague, 1971; McKenzie et al., 2007). They have direct ecological relevance because  
68 the survival of an organism under stressful conditions depends on its ability to balance  
69 energy demand and energy supply (Calow and Forbes, 1998), and complex traits of  
70 functional integrity and energetics will influence how effectively a fish can achieve this,  
71 and cope with the conditions and the demands of life in their habitat (Cairns, 1966;  
72 Sprague, 1971; McKenzie et al., 2007). Physiological traits are particularly useful as  
73 components of an approach combining multiple biomarkers, which can capture  
74 different response kinetics and toxicological impacts of contaminants, especially in  
75 natural environments where contaminants occur as complex mixtures (Hook et al.,  
76 2014; Dalzochio et al., 2016).

77           Traits of tolerance of environmental stressors, such as hypoxia or warming,  
78 were first proposed as biomarkers 50 years ago (Cairns, 1971; Sprague, 1971). They

79 are now gaining increased ecological relevance because of ongoing climate change,  
80 which is predicted to make freshwater environments increasingly physiologically  
81 challenging for fishes (Diaz and Breitburg, 2009; Noyes et al., 2009; Stillman, 2019).  
82 Aquatic hypoxia is expected to be ever more common as a stressor, including an  
83 increased frequency of severe hypoxic episodes (Diaz and Breitburg, 2009; Costa and  
84 Barletta, 2016). The same is true of warming where, beyond gradual rises in seasonal  
85 mean temperatures, an increased frequency, intensity and duration of seasonal  
86 heatwaves is projected to be a major source of environmental stress for fauna globally  
87 (Williams et al., 2016; Stillman, 2019). The extent to which contaminants can influence  
88 tolerance of hypoxia or warming may, therefore, be determinants of the capacity of fish  
89 populations to persist in certain habitats (Noyes et al., 2009; Noyes and Lema, 2015;  
90 Khursigara et al., 2019; Isaza, 2020). The capacity to perform exercise is also a  
91 valuable and ecologically relevant biomarker of sub-lethal toxic effects of pollutants on  
92 fishes, because activities such as foraging, escaping predators, protecting territory, and  
93 migration, all depend upon swimming (Brett, 1958; Cairns, 1966; Beamish, 1978;  
94 McKenzie et al., 2007; Domenici, 2010; Domenici and Hale, 2019). One particular  
95 advantage, of complex physiological traits of tolerance and performance as  
96 biomarkers, is that they integrate the function of multiple underlying physiological  
97 systems, and can provide insight into why fish fail to colonize some polluted habitats  
98 (McKenzie et al., 2007). It is therefore possible to investigate what causes impairment  
99 of tolerance or performance, to gain valuable understanding of the impacts of  
100 contaminants on fish health and populations (e.g. Beaumont et al., 1995, 2000; Tierney  
101 et al., 2007; Thomaz et al., 2009; Gomez Isaza et al., 2020), which is particularly useful  
102 within programmes of ecological risk assessment.

103 We review knowledge on the application of physiological biomarkers in  
104 subtropical and tropical fish species, which we take to mean those that can occupy  
105 habitats where average monthly temperatures do not fall below 18 °C. Many  
106 subtropical and tropical freshwater ecosystems suffer from pollution and limited  
107 environmental regulation, so exposure to contaminants represents a serious hazard to  
108 population health of resident fishes (Langiano et al., 2008; Cazenave et al., 2009; Braz-  
109 Mota et al., 2015; Beltrão et al., 2019; Barreto et al., 2020). At the same time, tropical  
110 ecosystems are relatively poorly studied in terms of ecotoxicology, despite harbouring  
111 important resources in terms of biodiversity (Lacher and Goldstein, 1997; Daam and  
112 Van Den Brink, 2010; Wang et al., 2019). They can also be naturally subject to  
113 stressful hypoxic and warming events, especially because many have seasonal  
114 variations in rainfall that severely modify the volume and connectivity of habitats. We  
115 consider studies that have used the critical oxygen level ( $O_{2crit}$ ) as a biomarker. For

116 these studies, we pay particular attention to how changes in  $O_{2crit}$  caused by  
117 contaminant exposure are linked to elements of cardiorespiratory physiology and  
118 performance, as these are core components of whole animal tolerance (Driedzic and  
119 Gesser, 1994; Gamperl and Driedzic, 2009). We then consider studies that have used  
120 traits of thermal tolerance as biomarkers, in particular the critical thermal maximum  
121 ( $CT_{max}$ ). Finally, we review the few studies that have used critical swimming speed  
122 ( $U_{crit}$ ) as a biomarker.

123

## 124 **Effects of contaminants on tolerance of hypoxia**

125 Hypoxic events can be very frequent in subtropical and tropical freshwater  
126 ecosystems, with dissolved oxygen (DO) fluctuations of different duration and  
127 magnitude, diurnally, spatially, and seasonally (Almeida-Val et al., 2006). Although  
128 fluctuating levels of DO can be a natural phenomenon, hypoxia caused by  
129 eutrophication and pollution is one of the most pressing and critical water problems in  
130 the world (Pollock et al., 2007). Hypoxia and contaminant exposure often co-occur in  
131 aquatic environments but their interactions remain poorly studied (Monteiro et al.,  
132 2020). There is evidence that co-exposures to hypoxia and polycyclic aromatic  
133 hydrocarbon mixtures caused several disorders in fishes, such as altered growth and  
134 sexual differentiation, reduced reproductive capacity, and developmental toxicity  
135 (Matson et al. 2008; Di Giulio and Clark, 2015; Mu et al., 2017). Rahman et al. (2018)  
136 demonstrated that co-exposure to hypoxia and polychlorinated biphenyl reduced  
137 biotransformation of organic xenobiotics by monooxygenase enzyme cytochrome  
138 P450-1A (CYP1A), but markedly augmented hepatic endothelial nitric oxide synthase  
139 (eNOS) and interleukin-1 $\beta$  (IL-1 $\beta$ ) mRNA levels, and protein carbonyl (PC) contents.

140 Hypoxia survival requires a well-coordinated response to either secure more  $O_2$   
141 from the depleted environment or to protect against the metabolic consequences of low  
142  $O_2$  levels at the mitochondria, which limit aerobic ATP production (Regan et al., 2017;  
143 Richards, 2019). The critical oxygen level ( $O_{2crit}$ ) is widely used as a quantifiable trait of  
144 hypoxia tolerance in fishes. It represents the minimum water oxygen level required to  
145 sustain standard metabolic rate (SMR, the basal metabolic rate of an ectotherm at their  
146 acclimation temperature) (Claireaux and Chabot, 2016; Rogers et al., 2016). It is  
147 determined by plotting rates of oxygen uptake ( $MO_2$ ) against water  $O_2$  tension (kPa), to  
148 identify the point at which  $MO_2$  falls below SMR, and becomes directly dependent upon  
149 water  $O_2$  tension (Rogers et al., 2016). Once below their  $O_{2crit}$ , fishes must depend on  
150 anaerobic energy metabolism and basal functions can be compromised (Pörtner,  
151 2010). If the hypoxic conditions worsen and persist, the combination of hypoxic severity  
152 and accumulation of metabolic products can threaten survival (Claireaux and Chabot,

153 2016). Although the ecological significance of the  $O_{2crit}$  threshold has been debated  
154 (Wood, 2018; Regan et al., 2019; Ultsch and Regan, 2019), it varies among species in  
155 a manner that is consistent with their overall hypoxia sensitivity, being lower in more  
156 tolerant species (Rogers et al., 2016). A low  $O_{2crit}$  value indicates a higher capacity for  
157  $O_2$  extraction by the gills and  $O_2$  delivery to the tissues as  $O_2$  availability declines,  
158 indicating a greater hypoxia tolerance (Mandic et al., 2009).

159 Figure 1 shows data for one species, matrinxã *Brycon amazonicus*, where  
160 sublethal exposures to various contaminants all increased  $O_{2crit}$  values. The degree of  
161 physiological plasticity for hypoxia tolerance is a key determinant of species  
162 performance (Rogers et al., 2016). This was associated with various impacts of the  
163 contaminants on  $MO_2$  during graded hypoxia (Table 1). This is shown in more detail for  
164 the matrinxã, where contaminants had various effects on  $O_2$  uptake such as increases  
165 and decreases in  $MO_2$  during normoxia (for Roundup Original® and Roundup  
166 Transorb®, respectively) and/or early decline in  $MO_2$  during graded hypoxia for all  
167 pollutants (Figure 2). All these effects had the consequence of impairing the capacity of  
168 the species to oxyregulate. According to McRae et al. (2018) metabolic rate is, itself, a  
169 useful integrated measurement of toxicological impact because it is a measure of  
170 energy resource associated with multiple sublethal processes and ecologically  
171 important endpoints such as reproduction, growth and survival.

172 The exposure to various contaminants markedly increased  $O_{2crit}$ , and therefore  
173 reduced tolerance of hypoxia, in four species of tropical fishes, including hypoxia-  
174 tolerant fish species such as Nile tilapia *Oreochromis niloticus* and trahira *Hoplias*  
175 *malabaricus* (Table 1). These data clearly indicate that exposure to contaminants may  
176 reduce survival chances of many species of fishes in tropical hypoxic environments.  
177 Table 1 also summarises how different pollutants can have a diverse array of effects on  
178 both ventilatory and cardiac activity in normoxia, and how such activity can be affected  
179 at given levels of hypoxia.

180 Oxyregulation is achieved by reflex cardiorespiratory responses that aim to  
181 sustain oxygen uptake and delivery, and possibly protect vital organs such as the  
182 heart. There is a profound hyperventilation, involving increases in gill ventilation volume  
183 ( $\dot{V}G$ ) achieved through increased respiratory frequency ( $f_R$ ) and amplitude, and a  
184 bradycardia (Farrell, 2007; Perry et al., 2009; Joyce et al., 2016; Rantin et al., 2020).  
185 The hyperventilation increases  $O_2$  transport to the site of gas exchange (Bergsson et  
186 al., 2019). Enhancements in  $\dot{V}G$  are required to maintain a constant  $MO_2$ , which implies  
187 that fish survival in hypoxic environments can be compromised if the ability to increase  
188  $\dot{V}G$  is affected (Martins et al., 2011). Pollutants can, indeed, impair the ability to

189 increase  $f_R$  and  $\dot{V}G$  in response to hypoxia, impairing the regulation of metabolic rate.  
190 For example, the lower  $f_R$  induced by acute exposure to the organophosphate  
191 trichlorfon in Nile tilapia *Oreochromis niloticus*, both in normoxia and hypoxia, impaired  
192 gill ventilation and compromised the hyperventilatory response in severe hypoxia  
193 (Thomaz et al., 2009). The same response pattern was observed in trahira after  
194 subchronic and trophic exposure to mercury chloride (Monteiro et al., 2013). In  
195 matrinxã, increases of  $f_R$  and  $\dot{V}G$  in response to hypoxia were significantly reduced  
196 after acute exposure to the emerging contaminant triclosan, an antimicrobial agent  
197 widely used in personal care products. In normoxia, triclosan had caused an increased  
198  $f_R$  with concomitant decreased  $\dot{V}G$  (Martins, 2018). Under hypoxia,  $\dot{V}G$  values were also  
199 considerably lower in Nile tilapia 48 h after intraperitoneal injection of microcystin-LR  
200 (Martins et al., 2011) and in matrinxã acutely exposed to glyphosate-based herbicides  
201 Roundup WG<sup>®</sup> and Roundup Transorb<sup>®</sup> (Anelli Jr., 2010), probably due to alterations of  
202 homeostatic reflexes that impaired respiratory responses to hypoxia, due to unknown  
203 toxic mechanisms. All these effects described above and presented in Table 1 can  
204 reduce the ability of tropical fish to maintain a constant O<sub>2</sub> uptake when they encounter  
205 hypoxia in contaminated environments

206 Organophosphate (OP) pesticides, mercury chloride, triclosan, and glyphosate-  
207 based herbicides inhibit acetylcholinesterase activity resulting in paralysis, mainly of  
208 the fins and respiratory muscles, uncoordinated movements, seizures, and even  
209 central depression of respiration (Suresh et al., 1992; Sancho et al., 1998; Antonijevic  
210 and Stojiljkovic, 2007; Thomaz et al., 2009; Modesto and Martinez, 2010; Menéndez-  
211 Helman et al., 2012; Pullaguri et al., 2021). Furthermore, contaminants such as  
212 microcystins and glyphosate-based herbicides cause oxidative stress and alterations in  
213 gill structure and respiratory epithelium (Gupta and Guha, 2006; Shiogiri et al., 2012;  
214 Braz-Mota et al., 2015; Martins et al., 2017) which could result in disturbances in  
215 respiratory function. Specifically for matrinxã exposed to Roundup Original<sup>®</sup>, Roundup  
216 WG<sup>®</sup> and Roundup Transorb<sup>®</sup>,  $\dot{V}G$  values were already high in normoxia (Table 1),  
217 indicating increased respiratory drive to maintain O<sub>2</sub> extraction.

218 On the other hand, contaminants can also induce rises in respiratory frequency  
219 and conspicuous increases in gill ventilation, including in hypoxia, which may indicate a  
220 desperate attempt to meet tissue oxygen demand. Increased gill ventilation was  
221 detected in matrinxã after methyl parathion and Roundup Original<sup>®</sup> exposure, mainly  
222 during hypoxia (Table 1, Olle 2007; Anelli Jr., 2010). Wang et al. (2013) evaluated the  
223 ventilatory responses of 3 tropical species, zebrafish *Danio rerio*, rare minnow  
224 *Gobiocypris rarus* and grass carp *Ctenopharyngodon idellus* to acute exposure to 4  
225 heavy metal ions (Hg<sup>2+</sup>, Cu<sup>2+</sup>, Cd<sup>2+</sup>, and Zn<sup>2+</sup>). Ventilatory frequency and amplitude



226 increased after exposure to  $\text{Hg}^{2+}$  and  $\text{Cu}^{2+}$  but markedly decreased with increasing  
227 concentrations of  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$ . This shows that respiratory activity of fishes is highly  
228 sensitive to heavy metal pollution but in complex ways (Wang et al., 2013). Differences  
229 in ventilatory responses to hypoxia could also be related to changes in blood  $\text{O}_2$ -  
230 carrying capacity (Pereira et al., 2013; Islam et al., 2019), according to the type and  
231 concentration of contaminants and the physiology of the target species. In the silver  
232 perch *Bidyanus bidyanus* from Australia, exposure to nitrate caused animals to perform  
233 aquatic surface respiration (ASR) at a higher oxygen level during progressive hypoxia  
234 (from 19 to 1 kPa), coupled with increased ventilation frequency. This was linked to  
235 lower levels of hemoglobin and hematocrit, indicating reduced blood oxygen carrying  
236 capacity (Gomez Isaza et al., 2021a). Fish exposed to pollutants often show  
237 hyperplasia and hypertrophy of gill epithelia cells with fusing and thickening of gill  
238 lamellae and mucus hypersecretion, which increases the water/blood diffusion distance  
239 and impairs  $\text{O}_2$  uptake, leading to hypoxemia (Schjolden et al., 2007; Sokolova and  
240 Lannig, 2008; Bilberg et al., 2010). In traíra, exposure to a single dose of microcystin-  
241 LR ( $100 \mu\text{g Kg}^{-1}$  of body weight, i.p.) caused hyperventilation coupled with significantly  
242 reduced  $\text{O}_2$  extraction from the ventilatory current under normoxic and hypoxic  
243 conditions, which compromised  $\text{O}_{2\text{crit}}$  and was attributed to compromised gill gas  
244 transfer (Martins et al., 2019).

245 Many pollutants (methyl parathion, trichlorfon, Roundup WG<sup>®</sup>, Roundup  
246 Transorb<sup>®</sup>, mercury chloride, microcystin) also impaired cardiac activity, causing a  
247 bradycardia that then lead to very low heart rates under hypoxic conditions (Table 1).  
248 This may indicate direct toxic effects on cardiac myocytes. Several studies have  
249 reported decreased myocardial isometric twitch force development in tropical fishes  
250 exposed to pollutants (Incardona et al., 2004; Thomaz et al., 2009; de Andrade  
251 Waldemarin et al., 2012; Monteiro et al., 2017). Reduced cardiac performance can  
252 increase the likelihood of starvation, predation vulnerability, and disease susceptibility  
253 (Incardona and Scholz, 2017). Exposure to  $1 \text{ mg L}^{-1}$  of copper sulfate for 96 h caused  
254 significant changes in Nile tilapia cardiomyocytes, including decreased myocardial  
255 contractility, cellular swelling, mitochondrial swelling, and alterations in mitochondrial  
256 inner membranes, which may impair ATP supply (de Andrade Waldemarin et al.,  
257 2012). Monteiro et al. (2020) found that cardiac responses to hypoxia were impaired in  
258 both matrinxã and traíra after exposure to inorganic mercury by different routes (water  
259 and diet) and durations (acute and sub-chronic). There were alterations in the rhythm  
260 of the cardiac pacemaker such as first-degree atrioventricular block, lengthening of the  
261 action potential (AP) plateau, and the development of cardiac arrhythmias and  
262 abnormalities (episodes of missing QRS complexes with sequential P waves,

263 extrasystoles, negative QRS complex, T wave inversion) (Monteiro et al., 2020). These  
264 disruptions of electrical conduction could underly the myocardial dysfunction and  
265 increase hypoxia sensitivity by impairing cardiac output (Rantin et al., 1995).

266 Overall, these findings demonstrate that  $O_{2crit}$  is a valuable biomarker of impacts  
267 of contaminants in tropical freshwater fishes, irrespective of the ecology and life history  
268 characteristics of a given species. In addition, analyses of effects on  $MO_2$  and  
269 cardiorespiratory variables can improve understanding of the mode of toxic action of  
270 some contaminants, which can aid in the environmental risk assessment of aquatic  
271 systems. The fact that contaminants can impair hypoxia tolerance may have significant  
272 ecological consequences, not just directly influencing survival but potentially also  
273 reducing fitness, by impairing the capacity to forage, capture and digest food, with  
274 consequences for growth and/or reproductive rates. Nonetheless, the knowledge base  
275 is critically limited and should be extended to further species and contaminants.

276

### 277 **Effects of contaminants on tolerance of warming**

278 There are complex interactions between temperature and the toxicity of  
279 contaminants in fishes, which depend upon multiple factors that are too extensive to  
280 review here (Noyes et al., 2009; Patra et al., 2015). We focus exclusively upon the use  
281 of thermal tolerance as a biomarker. In fishes, limits of thermal tolerance are measured  
282 using acute thermal ramping protocols, especially the critical thermal (CT) methodology  
283 that reveals the threshold temperature which causes loss of equilibrium (LOE). The  
284 protocol is simple and defines the temperature where survival is threatened because at  
285 LOE the fish cannot escape the conditions (Beitinger and Lutterschmidt, 2011). There  
286 are, however, quite major limitations to the critical thermal protocol as an experimental  
287 tool (Rezende et al., 2014; Blasco et al., 2020). Notably, the idea that the complexity of  
288 thermal tolerance limits can be captured by a single measure is a major  
289 oversimplification (Rezende et al., 2014; Rezende and Bozinovic, 2019; Lefevre et al.,  
290 2021). There are also methodological issues; particularly that the measured  $CT_{max}$   
291 depends upon heating rate (Lutterschmidt and Hutchison, 1997). Furthermore,  
292 although the mechanism(s) that causes LOE are presumed to involve impaired function  
293 of critical organs such as the heart and brain (Wang et al., 2014; Brijs et al., 2015;  
294 Jutfelt et al., 2019; Lefevre et al., 2021), they are not known and may differ among  
295 species and with warming rate (Currie and Schulte, 2014; Healy et al., 2018; Blasco et  
296 al., 2020). What is well-established, however, is that a given increase in acclimation (or  
297 acclimatization) temperature ( $T_{acc}$ ) is not linked to an equivalent increase in  $CT_{max}$ , both  
298 among and within fish species (Cossins and Bowler, 1987; Currie and Schulte, 2014;  
299 McKenzie et al., 2020). As a consequence, when comparing among species, tropical

300 fishes have a lower thermal safety margin (TSM) than temperate, where TSM is  
301 calculated as the net difference between  $T_{acc}$  and  $CT_{max}$  (Comte and Olden, 2017a; b).  
302 Within a species, TSM also becomes smaller as a fish is acclimated to progressively  
303 warmer temperatures across its thermal range (e.g. McDonnell et al., 2019; McKenzie  
304 et al., 2020).

305 The notion that  $CT_{max}$  could be used as a biomarker of sublethal toxic effects of  
306 contaminants dates from the 1970s (Cairns, 1971; Johnson, 1978; Dale Becker and  
307 Wolford, 1980) although the trait has only been used in this way on 11 species of  
308 subtropical or tropical fish (Table 2). As for all biomarkers that are complex  
309 physiological traits,  $CT_{max}$  can provide a sensitive indicator of functional impairment  
310 caused by a contaminant. The lack of mechanistic understanding does, however,  
311 hinder comprehension of essential questions such as why  $CT_{max}$  is modified by  
312 exposure to contaminants. Nonetheless, the  $CT_{max}$  is interesting from a toxicological  
313 viewpoint because it is an incipient lethal threshold, therefore if it is reduced by  
314 contaminants this could have clear consequences for persistence of fish populations  
315 that suffer heat stress in polluted waters. The low TSM of subtropical and tropical  
316 species acclimated (or acclimatized) to warm waters may render them particularly  
317 vulnerable to heatwaves if they are in contaminated environments, especially if they  
318 are at temperatures close to the top of their thermal range.

319 There have been various investigations of effects on  $CT_{max}$  of organic  
320 contaminants, notably pesticides, herbicides and common pollutants from industry such  
321 as phenol. Many of these molecules act as neurotoxins in vertebrates, hence they  
322 might be expected to influence  $CT_{max}$  since LOE almost certainly has some  
323 neurological component (Lutterschmidt and Hutchison, 1997; Jutfelt et al., 2019). In  
324 many cases, the studies involved first establishing lethal concentrations (LC50) for the  
325 focal species, and then evaluating effects of sublethal concentrations on  $CT_{max}$ .  
326 Johnson (1978) studied the effects of four widely-used OP insecticides on  $CT_{max}$  in the  
327 western mosquitofish *Gambusia affinis*, when acclimated to 20 °C (Table 2).  
328 *Gambusia affinis* is a subtropical North American species that is highly eurythermal.  
329 As acetylcholinesterase inhibitors, OP insecticides have neurotoxic effects on fishes.  
330 Taking concentrations that were proposed as doses for field applications, the  
331 pesticides had no significant effects on mosquitofish  $CT_{max}$  aside from chlorpyrifos,  
332 which caused a modest but significant decline in tolerance (Table 2). Takle et al. (1983)  
333 reported that sublethal concentrations of the biocide endothall, which is a protein  
334 phosphatase  $A_2$  inhibitor, had no significant effect on  $CT_{max}$  in another subtropical  
335 eurythermal north American species, the red shiner *Cyprinella lutrensis*, acclimated to  
336 20 °C. In the same species, Messaad et al. (2000) investigated effects on  $CT_{max}$  of

337 sublethal concentrations of the triazine herbicide, atrazine, of an OP pesticide,  
338 terbufos, plus a cocktail of these, comparing effects on fish acclimated to 23° or 30 °C.  
339 Toxic actions of atrazine in animals are supposedly related only to endocrine  
340 disruption.  $CT_{max}$  of control fish increased with acclimation temperature but the  
341 contaminants caused more marked discrete declines in  $CT_{max}$  at a  $T_{acc}$  of 30 °C  
342 compared to 23 °C (Table 2). In particular, the cocktail of the two pesticides caused a  
343 very profound decline in  $CT_{max}$  at 30 °C (Table 2). The mechanisms underlying the  
344 increased impact upon  $CT_{max}$  at the higher temperatures may relate to increased  
345 metabolic rates and ventilatory water flows, which would cause increased uptake of  
346 contaminants from the water (Yang et al., 2000). Such results indicate that, as  $T_{acc}$   
347 increases, fishes may be more vulnerable to heat stress when they are also exposed to  
348 certain environmental contaminants (Messaad et al., 2000).

349 Patra et al. (2007) studied the effects on  $CT_{max}$  of sublethal concentrations of  
350 three contaminants, endosulfan, an organochlorine pesticide; the OP chlorpyrifos, and  
351 an organic contaminant, phenol, in three subtropical Australian species, silver perch,  
352 eastern rainbowfish *Melanotaenia duboulayi* and western carp gudgeon *Hypseleotris*  
353 *klunzingeri*. Endosulfan is neurotoxic, a GABA-gated chloride channel antagonist, and  
354 a  $Ca^{2+}$ ,  $Mg^{2+}$  ATPase inhibitor. It is used to control insect pests but is acutely toxic to  
355 animals and is now banned in many countries. Phenol is one of the most common  
356 molecules in the chemical industry globally with a vast array of uses, it can have toxic  
357 effects on the central nervous system and heart of vertebrates. Although both  
358 endosulfan and chlorpyrifos caused significant declines in  $CT_{max}$  in all species, while  
359 phenol was without effect. Kumar et al. (2016) reported that, in a euryhaline tropical  
360 Asian species the milkfish *Chanos chanos* acclimated to 28 °C, sublethal  
361 concentrations of endosulfan reduced  $CT_{max}$  compared to a control (Table 2), an effect  
362 that was abolished by dietary supplements of vitamin B<sub>6</sub>, pyroxidine. Pyroxidine  
363 supplementation was linked to changes in tissue oxidative status and expression of the  
364 heat shock protein HSP70, which may have been related to the improved thermal  
365 tolerance (Kumar et al., 2016).

366 Zebral et al. (2018) investigated effects of the glyphosate-based herbicide,  
367 Roundup®, on thermal tolerance of embryos of an endangered annual neotropical  
368 killifish, *Austrolebias nigrofasciatus*, hatched at 20 °C. This annual species leaves  
369 eggs in diapause in dried water courses during the dry season, which hatch when rains  
370 arrive. Glyphosate is not considered to have acute toxic effects on vertebrates.  
371 Glyphosate caused, however, a significant decline in  $CT_{max}$  in hatched embryos held at  
372 21 °C, when measured as cardiac arrest followed by mortality despite being  
373 immediately returned to acclimation temperature (Table 2). Philippe et al. (2019)

374 investigated the effects, on  $CT_{max}$  of adults of the turquoise killifish *Notobranchius*  
375 *furzeri*, an annual species from Africa, of a highly persistent compound released from  
376 chlorinated aniline herbicides, 3,4-dichloroaniline (3,4-DCA). Fish were acclimated to  
377 either 24 or 28 °C and exposed to two concentrations of 3,4-DCA,  $CT_{max}$  increased with  
378  $T_{acc}$  and the contaminant only caused a significant decline in  $CT_{max}$  at the highest  
379 concentration at 28 °C (Table 2).

380         There has been some investigation of effects of sublethal concentrations of  
381 toxic heavy metals on  $CT_{max}$  in subtropical or tropical fish species. Cadmium and lead  
382 are both highly toxic to fishes, they attack various organs systems including gills,  
383 intestine and kidney. Notably, at sublethal concentrations they are neurotoxic (Green  
384 and Planchart, 2018) and, therefore, could influence thresholds for LOE and  $CT_{max}$ .  
385 Carrier and Beitinger (1988) found that sublethal concentrations of cadmium decreased  
386  $CT_{max}$  in the red shiner acclimated to 20 °C, with a severity that increased with  
387 exposure dose and duration (Table 2). By contrast, Philippe et al. (2018) found that  
388 two sublethal concentrations of cadmium had no effect on  $CT_{max}$  in the turquoise  
389 killifish acclimated to either 24 or 28 °C, although  $CT_{max}$  increased with acclimation  
390 temperature (Table 2). In the Mekong striped catfish *Pangasianodon hypophthalmus*  
391 acclimated to 34 °C, exposure to a sublethal concentration of lead caused a significant  
392 decline in  $CT_{max}$ , which was ameliorated by dietary supplementation with selenium or  
393 zinc (Kumar et al., 2017, 2018). This amelioration was concomitant with changes in  
394 redox balance of the tissues and in expression of HSPs, further circumstantial evidence  
395 that these elements of cell physiology may be involved in thermal tolerance in fishes.

396         There has also been some investigation of effects of nutrient contaminants on  
397 tolerance of warming. Nitrite is a pervasive form of nutrient pollution that enters aquatic  
398 habitats from diverse sources such as nitrogen-based fertilisers, livestock manure,  
399 sewage wastewater, and atmospheric deposits. The major toxic effect of nitrite is that it  
400 compromises blood oxygen transport in fishes, by causing methemoglobin formation  
401 (Lewis and Morris, 1986; Jensen, 2003). Watenpaugh et al. (1985) found that  
402 increasing sublethal concentrations of nitrite caused a progressive decline of  $CT_{max}$  in  
403 the subtropical eurythermal channel catfish *Ictalurus punctatus* acclimated to 20 °C.  
404 The  $CT_{max}$  was negatively correlated with blood methaemoglobin concentrations, which  
405 would seem to indicate that oxygen transport capacity is a determinant of  $CT_{max}$  in this  
406 species. Rodgers and De Boeck (2019) reported that a sub-lethal nitrite concentration  
407 caused a significant decline in  $CT_{max}$  of in a globally extremely widespread subtropical  
408 eurytherm, the common carp *Cyprinus carpio*, when acclimated to 23°C (Table 2).

409         Within the extreme boundaries delineated by  $CT_{max}$ , thermal tolerance in fishes  
410 is typically evaluated with a thermal performance curve (TPC) that measures a trait of

411 organismal performance over a range of temperatures to identify features such as a  
412 thermal optimum for performance (Schulte et al., 2011; Currie and Schulte, 2014). The  
413 most common TPC is for aerobic metabolic scope (AS), the net capacity to supply  
414 oxygen for aerobic metabolic activities beyond basal maintenance (Fry, 1971; Schulte  
415 et al., 2011; Schulte, 2015). Very few studies have investigated how contaminants  
416 might influence elements of a TPC in subtropical or tropical fishes (Rodgers and De  
417 Boeck, 2019; Gomez Isaza et al., 2020a), also because this requires very extensive  
418 experimental work.

419         The study by Rodgers and De Boeck (2019) on common carp exposed to nitrite  
420 is a rare example where changes in  $CT_{max}$  were related to aerobic performance and  
421 aerobic scope. Although nitrite reduced  $CT_{max}$  this was not related to any decline in AS,  
422 indicating that  $CT_{max}$  in the carp is not affected by capacity to provide oxygen for  
423 metabolism when warmed. The most comprehensive evaluations of how a contaminant  
424 might influence thermal tolerance and thermal sensitivity of performance in a tropical  
425 freshwater fish are studies on effects of nitrate on the silver perch (Gomez Isaza et al.,  
426 2020a, 2021b). Nitrate is the most abundant nutrient contaminant globally; although  
427 less toxic than nitrite it nonetheless shows persistently high concentrations in  
428 freshwaters due to agricultural runoff from nitrogen-based fertilisers, fossil fuel  
429 combustion and wastewater treatment plants. Nitrate also causes the formation of  
430 methaemoglobin, compromising blood oxygen carrying capacity (Gomez Isaza et al.,  
431 2020b). Gomez Isaza et al. (2020a) investigated whether nitrate exposure would  
432 increase susceptibility to elevated temperatures, measured as  $CT_{max}$ , and limit aerobic  
433 performance at warmer temperatures, measured as AS by swimming respirometry. In  
434 silver perch acclimated to 28 °C, exposure to either 50 or 100 mg l<sup>-1</sup> nitrate caused a  
435 decline in  $CT_{max}$  that was not, however, observed in fish acclimated to a warmer  
436 temperature of 32 °C, despite these latter having higher overall  $CT_{max}$  (Table 2).  
437 Furthermore, when swum at three temperatures (28, 32 or 36 °C), nitrate-exposed fish  
438 with a  $T_{acc}$  of 28 °C showed a decline in AS (and swimming performance, see below) at  
439 36 °C, whereas those with a  $T_{acc}$  of 32 °C displayed a thermally-insensitive phenotype,  
440 with no change in AS or performance across temperatures. It was subsequently  
441 demonstrated that the silver perch with a  $T_{acc}$  of 32 °C exhibit thermally-related plastic  
442 compensation of the cardiorespiratory system, including increased gill surface area and  
443 greater ventricular thickness (Gomez Isaza et al., 2021b). This demonstrates how the  
444 thermal physiology of fishes may have complex interactions with their relative  
445 sensitivity to contaminants (Gomez Isaza et al., 2020a, 2021b).

446         One major conclusion to be drawn from Table 2 is that the knowledge base for  
447 effects of contaminants on thermal tolerance in subtropical and tropical freshwater

448 fishes is extremely limited, with a majority of information for subtropical species. Given  
449 that the mechanisms that cause LOE in  $CT_{max}$  are not known, it is not possible to  
450 speculate about why some contaminants cause a decline in thermal tolerance while  
451 others do not. Some patterns can, however, be demonstrated in the limited dataset, as  
452 a basis for stimulating further research. When simply considering all  $CT_{max}$  data in  
453 Table 2, there is a very clear and highly significant decline in TSM of control animals as  
454  $T_{acc}$  increases (Fig 3). This analysis makes no effort to correct for warming rate,  
455 developmental stage (one species was studied as an embryo), but the results confirm  
456 that is known about effects of  $T_{acc}$  on TSM in fishes (Comte and Olden, 2017a; b). The  
457 question that then arises is whether exposure to contaminants will exacerbate this  
458 effect, rendering tropical species even more vulnerable to warming events when they  
459 are in contaminated habitats. For those species in Table 2 where a contaminant  
460 caused a significant decline in  $CT_{max}$ , there was also a highly significant decline in TSM  
461 with  $T_{acc}$  (Figure 3). The intercept and slope of the relationships did not, however, differ  
462 significantly from those of control animals (Fig 3). Fig 4 shows, however, the  
463 proportional extent (as a percentage) to which contaminants reduced TSM compared  
464 to control conditions, as a function of  $T_{acc}$ . There was a significant effect whereby the  
465 higher  $T_{acc}$ , the greater the extent to which a contaminant would compromise TSM (Fig  
466 4). We show this figure precisely to highlight why it is unsatisfactory, namely that there  
467 is a critical lack of information for effects of contaminants on TSMs of fishes at warm  
468 tropical temperatures. There are major reasons for caution in interpreting this figure.  
469 Firstly, the four points at 30 °C and above are from two species, red shiner and striped  
470 catfish; the three points at 30°C all come the from the study on red shiner that  
471 demonstrated how two pesticides and their cocktail caused much more profound  
472 declines in  $CT_{max}$  in fish acclimated to 30 ° compared to 23 ° C by (Messaad et al.  
473 2000). Indeed, at 30 °C the pesticide cocktail reduced the TSM by over 80% compared  
474 to uncontaminated animals (Fig 4). Secondly, as mentioned above, warmer  
475 temperatures did not increase susceptibility to nitrate in silver perch, because of plastic  
476 physiological responses by this eurythermal subtropical species (Gomez Isaza et al.,  
477 2020a, 2021b), so this data is not carried on the figure. Therefore, it is very important to  
478 extend the knowledge base on how contaminants affect thermal tolerance of tropical  
479 species, to investigate whether they are particularly vulnerable to impacts of heatwaves  
480 when exposed to common pollutants. This is highly relevant to the life cycle of many  
481 tropical species that, during dry seasons, may find themselves confined to increasingly  
482 smaller water bodies, which will tend to concentrate any pollutants as water evaporates  
483 and also to be submitted to extremes of warmth due to solar radiation (Patra et al.,  
484 2007, 2015).

485

## 486 **Critical swimming speed as a biomarker**

487 Swimming performance has clear ecological relevance for many fish species,  
488 so measures of performance can provide valuable biomarkers of toxic effects of  
489 aquatic contaminants (Cairns, 1966; Beamish, 1978; Tudorache et al., 2008; McKenzie  
490 et al. 2007, 2008; Domenici, 2010). In particular, the critical swimming speed protocol  
491 ( $U_{crit}$ , Brett, 1964) was first proposed as a sublethal physiological biomarker more than  
492 50 years ago (Cairns, 1966; Beamish, 1978). The  $U_{crit}$  protocol allows investigation of  
493 proximate mechanisms that underly impaired performance following exposure to  
494 contaminants, so gaining insight into mechanisms of toxic effects in target organs such  
495 as skeletal muscle (Beaumont et al., 1995, 2000; Tierney et al., 2007) or the heart  
496 (Khursigara et al., 2019).

497 The  $U_{crit}$  protocol exposes fish to stepwise increases in speed in a swimming  
498 respirometer, to establish the speed at which it fatigues. The knowledge base is  
499 dominated by work on temperate species and especially salmonids, while the  
500 Deepwater Horizon disaster led to extensive research on subtropical marine species.  
501 Studies have demonstrated impacts of many classes of contaminant under controlled  
502 laboratory conditions, including metals (e.g. Beaumont et al., 1995), xenobiotic  
503 organics (Wood et al., 1996; Tierney et al., 2007), crude oil itself (Stieglitz et al., 2016),  
504 waste products such as ammonia (Shingles et al., 2001), and other consequences of  
505 anthropogenic pressures, such as harmful algae (Corriere et al., 2020). Critical  
506 swimming speed has also been used to show impacts of exposure to water containing  
507 complex contaminant mixtures (e.g. Goertzen et al., 2012; Folkerts et al., 2020)  
508 including by exposing fishes in cages (McKenzie et al., 2007). The zebrafish is an  
509 important model vertebrate species for laboratory studies to demonstrate impacts of  
510 contaminants on swimming performance, to demonstrate toxic effects and their  
511 mechanisms (e.g. Thomas and Janz, 2011; Gerger et al., 2015; Lucas et al., 2016;  
512 Folkerts et al., 2017). Such studies on zebrafish have never, however, been related to  
513 conditions in their natural environment in tropical south Asia, so we will not review them  
514 here.

515 There are so few studies that, to our knowledge, have used  $U_{crit}$  as a biomarker  
516 of effects of contaminants on subtropical and tropical freshwater species (other than  
517 zebrafish), that it is almost trivial to review them. We do this, however, to stimulate  
518 further work. One necessity for such studies is, of course, access to a swim tunnels or  
519 swim flume. These are now commercially available, which opens up possibilities of  
520 using swimming performance as a valuable biomarker to support decision-making for  
521 water quality control in tropical countries. This is especially relevant for the protection of



522 economically valuable species that perform remarkable migrations through tropical  
523 river systems that are at increasing risk of contamination by multiple pollutants (Hogan  
524 et al., 2007; Duponchelle et al., 2016; Speranza et al., 2016). Other traits of  
525 performance such as the escape response (Tudorache et al., 2008; McKenzie et al.,  
526 2009) or aspects of swimming behavior (Little and Finger, 1990), which do not require  
527 a swim tunnel, can also be used as biomarkers.

528 Two studies have investigated effects of organic pollutants on  $U_{crit}$  in freshwater  
529 subtropical/tropical species. Perfluorooctane sulfonic acid (PFOS) is a fluorosurfactant  
530 with a wide range of commercial and industrial applications, which has become a  
531 pervasive and persistent pollutant of freshwaters globally (Augustsson et al., 2021). Xia  
532 et al. (2015) studied effects of a range of sublethal concentrations of PFOS on  $U_{crit}$  in  
533 qingbo *Spinibarbus sinensis*, a subtropical Asian species acclimated to two  
534 temperatures, 18 ° and 28 °C. Increasing concentrations of PFOS caused progressive  
535 declines in  $U_{crit}$ , while  $T_{acc}$  had no interaction with these impairments. Swimming  
536 respirometry showed that the decline in performance was linked to reduced swimming  
537 efficiency, in that higher rates of oxygen uptake were needed to achieve a given a  
538 swimming speed. The author suggested that this might be due to sensorimotor or  
539 muscle tissue impairments, due to toxic effects of PFOS. McKenzie et al. (2017) used  
540  $U_{crit}$  to evaluate individual variation in sub-lethal sensitivity to the OP pesticide  
541 trichlorfon in a globally widespread tropical species, the Nile tilapia. At a  $T_{acc}$  of 27 °C,  
542 trichlorfon reduced  $U_{crit}$  significantly and fish only exhibited a partial recovery 96h after  
543 return to control freshwater. The decline in  $U_{crit}$  was due to reduced swimming  
544 efficiency, which was attributed to the fact that OPs are acetylcholinesterase inhibitors  
545 that interfere with cholinergic nerve function at motor endplates and in the central  
546 nervous system of vertebrates. This would impair neuromuscular coordination in fishes,  
547 reducing their swimming efficiency (Tierney et al., 2007; McKenzie et al., 2017). This  
548 study is a rare investigation of how individuals in populations differ in their sensitivity to  
549 a pollutant, finding that intrinsically efficient swimmers were the least affected by the  
550 OP (McKenzie et al., 2017).

551 Gomez Isaza et al. (2020a) investigated effects of nitrate, the most prevalent  
552 nutrient contaminant globally, on thermal sensitivity of swimming performance in silver  
553 perch, acclimated to either 28 or 32 °C. Two sublethal exposure concentrations were  
554 evaluated at three temperatures (28, 32 and 36 °C), to reveal complex effects whereby  
555 fish acclimated to 28 °C had better overall  $U_{crit}$  performance than those acclimated to  
556 32 °C. Exposure to nitrate, however, impaired  $U_{crit}$  of the 28 °C fish, in a manner  
557 proportional to nutrient concentration, but was without effect on performance of the 32  
558 °C fish. This absence of effects of nitrate in fish acclimated to 32 °C was unexpected

559 but paralleled the findings for  $CT_{max}$  and aerobic performance that are reported above.  
560 In a study on the subtropical highly eurythermal spangled perch *Leiopotherapon*  
561 *unicolor*, Gomez Isaza et al. (2020b) investigated effects on  $U_{crit}$  of combinations of  
562 sublethal nitrate and low pH (acidification). The  $U_{crit}$  was impaired by both nitrate and  
563 acid pH, with most severe impairment being in fish at the highest nitrate concentration  
564 and low pH, although the combined effect was additive rather than synergistic. These  
565 impairments in performance were linked to a reduced capacity to raise oxygen uptake  
566 during swimming, leading to lower AAS. These proximate effects could, in turn, be  
567 attributed to measured negative effects of nitrate and low pH on blood oxygen carrying  
568 capacity (Gomez Isaza et al., 2020b). Such multifactorial studies provide excellent but  
569 challenging templates for future investigations of effects of contaminants on the  
570 environmental tolerance and physiological performance of freshwater tropical fishes.

571

## 572 **Conclusions**

573 This review demonstrates that  $O_{2crit}$ ,  $CT_{max}$  and  $U_{crit}$  can all be useful  
574 physiological biomarkers of the toxicological effects of water contaminants on tropical  
575 and subtropical freshwater fishes. Understanding the interactive effects on fish  
576 physiology of sublethal exposures to contaminants, hypoxia and heatwaves will be  
577 crucial for monitoring the health of populations and projecting their biological  
578 consequences before serious changes in population size occur. This review very  
579 clearly demonstrates the need to extend the knowledge base to more species and  
580 contaminants, to better evaluate subtropical and tropical freshwater fish vulnerability in  
581 light of ongoing climate change. Complex physiological traits should be used to support  
582 decision-making in formulating water quality criteria for fishes and conservation  
583 strategies for their habitats, to protect their rich biodiversity. They may be particularly  
584 useful in procedures to identify adverse outcome pathways (van der Oost et al., 2020).  
585 Despite the technical challenges involved, future studies should consider interactions  
586 among pollutants, hypoxia and warming in tropical fishes, as the stresses all co-occur  
587 in nature.

588

## 589 **Acknowledgments**

590 The studies performed by the authors were funded by grants from the  
591 Coordination for the Improvement of Higher Education Personnel (CAPES) and the  
592 National Council for Scientific and Technological Development (CNPq) of Brazil, and by  
593 the São Paulo Research Foundation (FAPESP).

594

## 595 **Conflict of Interest**

596           The authors declare that there are no conflicts of interests.

597

598   **Data Availability Statement**

599           The data that support the findings of this study are available from the  
600   corresponding author upon reasonable request.

601

602 **References**

- 603 Almeida-Val, V.M.F., Gomes, A.R.C., Lopes, N.P. (2006). Metabolic and physiological  
604 adjustments to low oxygen and high temperature in fishes of the Amazon. In A.L.  
605 Val, V.M.F. Almeida-Val, & D.J. Randall (Eds.), *Fish Physiology* (pp. 443-359).  
606 Elsevier, Heidelberg.
- 607 Anelli Jr., L.C. (2010). Efeitos de diferentes formulações comerciais do herbicida  
608 Roundup® sobre a função cardiorespiratória de matrinxã, *Brycon amazonicus*  
609 (Teleostei, Characidae). [Doctoral dissertation, Federal University of São Carlos].  
610 UA Campus Repository. <https://repositorio.ufscar.br/handle/ufscar/1220>
- 611 Antonijevic, B., Stojiljkovic, M.P. (2007). Unequal efficacy of pyridinium oximes in acute  
612 organophosphate poisoning. *Clinical Medicine & Research*, **5**, 71-82.
- 613 Augustsson, A., Lennqvist, T., Osbeck, C.M.G., Tibblin, P., Glynn, A., Nguyen, M.A.,  
614 Westberg, E., Vestergren, R. (2021). Consumption of freshwater fish: A variable  
615 but significant risk factor for PFOS exposure. *Environmental Research*, **192**,  
616 110284.
- 617 Barreto, L.S., Souza, A.T.D.C., Martins, C.C., Araujo, S.B.L., Oliveira Ribeiro, C.A.  
618 (2020). Urban effluents affect the early development stages of Brazilian fish  
619 species with implications for their population dynamics. *Ecotoxicology and*  
620 *Environmental Safety*, **188**, 109907.
- 621 Beamish, F.W.H. (1978). Swimming capacity. In: Hoar, W.S. & Randall, D.J. (Eds.),  
622 *Fish Physiology* (pp 101-187). New York: Academic Press.
- 623 Beaumont, M., Butler, P., Taylor, E. (1995). Plasma ammonia concentration in brown  
624 trout in soft acidic water and its relationship to decreased swimming performance.  
625 *Journal of Experimental Biology*, **198**, 2213-20.
- 626 Beaumont, M.W., Taylor, E.W., Butler, P.J. (2000). The resting membrane potential of  
627 white muscle from brown trout (*Salmo trutta*) exposed to copper in soft, acidic  
628 water. *Journal of Experimental Biology*, **203**, 2229-2236.
- 629 Beitinger, T. & Lutterschmidt, W. (2011). Temperature measures of thermal tolerance.  
630 In: A.P. Farrell (Ed.), *Encyclopedia of fish physiology: from genome to*  
631 *environment* (pp. 1695-1702). Elsevier Ltd.
- 632 Beltrão, H., Zuanon, J., Ferreira, E. (2019). Checklist of the ichthyofauna of the Rio  
633 Negro basin in Brazilian Amazon. *ZooKeys*, **881**, 53-89.
- 634 Bergsson, H., Andersen, N.R., Svendsen, M.B.S, Hansen, P.J., Steffensen, J.F.  
635 (2019). Respiratory physiology of european plaice (*Pleuronectes platessa*)  
636 exposed to *Prymnesium parvum*. *Fishes*, **4**, 32.

637 Bilberg, K., Malte, H., Wang, T., Baatrup, E. (2010). Silver nanoparticles and silver  
638 nitrate cause respiratory stress in Eurasian perch (*Perca fluviatilis*). *Aquatic*  
639 *Toxicology*, **96**, 159-165.

640 Blasco, F.R., Esbaugh, A.J., Killen, S.S., Rantin, F.T., Taylor, E.W., McKenzie, D.J.  
641 (2020). Using aerobic exercise to evaluate sub-lethal tolerance of acute warming  
642 in fishes. *Journal of Experimental Biology*, **223**, jeb218602.

643 Braz-Mota, S., Sadauskas-Henrique, H., Duarte, R.M., Val, A.L., Almeida-Val, V.M.F.  
644 (2015). Roundup® exposure promotes gills and liver impairments, DNA damage  
645 and inhibition of brain cholinergic activity in the Amazon teleost fish *Colossoma*  
646 *macropomum*. *Chemosphere*, **135**, 53-60.

647 Brett, JR. (1958). Implications and assessments of environmental stress. In: P.A.  
648 Larkin (Ed.), *The investigation of fish-power problems* (pp. 69-93). Vancouver,  
649 BC: University of British Columbia.

650 Brett, JR. (1964). The respiratory metabolism and swimming performance of young  
651 sockeye salmon. *Journal of the Fisheries Research Board of Canada*, **21**, 1183-  
652 1226.

653 Brijs, J., Jutfelt, F., Clark, T.D., Gräns, A., Ekström, A., Sandblom, E. (2015).  
654 Experimental manipulations of tissue oxygen supply do not affect warming  
655 tolerance of European perch. *Journal of Experimental Biology*, **218**, 2448-2454.

656 Cairns, J. (1966). Don't be half-safe- the current revolution in bio-assay techniques.  
657 Proc. 21 St Ind. Waste Conf. Purdue Univ. Engng Extn Ser. No. 121.

658 Cairns, Jr.J. (1971). Thermal pollution: A cause for concern on. *Water Pollution Control*  
659 *Federation*, **43**:55–66.

660 Calow, P. & Forbes, V.E. (1998). How do physiological responses to stress translate  
661 into ecological and evolutionary processes? *Comparative Biochemistry and*  
662 *Physiology Part A: Molecular & Integrative Physiology*, **120**, 11-16.

663 Carrier, R., Beiting, T.L. (1988). Reduction in thermal tolerance of *Notropis lutrensis*  
664 and *Pimephales promelas* exposed to cadmium. *Water Research*, **22**, 511-515.

665 Cazenave, J., Bacchetta, C., Parma, M.J., Scarabotti, P.A., Wunderlin, D.A. (2009).  
666 Multiple biomarkers responses in *Prochilodus lineatus* allowed assessing  
667 changes in the water quality of Salado River basin (Santa Fe, Argentina).  
668 *Environmental Pollution*, **157**, 3025-3033.

669 Claireaux, G., Chabot, D. (2016). Responses by fishes to environmental hypoxia:  
670 integration through Fry's concept of aerobic metabolic scope. *Journal of Fish*  
671 *Biology*, **88**, 232-51.

672 Comte, L., Olden, J.D. (2017a). Climatic vulnerability of the world's freshwater and  
673 marine fishes. *Nature Climate Change*, **7**, 718-722.

674 Comte, L., Olden, J.D. (2017b). Evolutionary and environmental determinants of  
675 freshwater fish thermal tolerance and plasticity. *Global Change Biology*, **23**, 728-  
676 736.

677 Corriere, M., Baptista, M., Paula, J.R., Repolho, T., Rosa, R., Costa, P.R., Soliño, L.  
678 (2020). Impaired fish swimming performance following dietary exposure to the  
679 marine phycotoxin okadaic acid. *Toxicon*, **179**, 53-59.

680 Cossins, A.R. & Bowler, K. (1987). *Temperature Biology of Animals*. London: Chapman  
681 and Hall.

682 Costa, M.F. & Barletta, M. (2016). Special challenges in the conservation of fishes and  
683 aquatic environments of South America. *Journal of Fish Biology*, **89**, 4-11.

684 Costa, M.J., Ribeiro, L.R., Salla, R.F., Gamero, F.U., Alves, L.M.L.M., Silva-Zacarin,  
685 E.C.M. (2015). Effects of the organophosphorus pesticide Folisuper 600 (methyl  
686 parathion) on the heart function of bullfrog tadpoles, *Lithobates catesbeianus*  
687 (Shaw, 1802). *Brazilian Journal of Biology*, **75**, 163-168.

688 Currie, S. & Schulte, P.M. (2014). Thermal stress. In: D.H. Evans, J. Claiborne, S.  
689 Currie (Eds.), *The physiology of fishes* (pp. 257-279). Boca Raton: CRC Press.

690 Daam, M.A. & Van Den Brink, P.J. (2010). Implications of differences between  
691 temperate and tropical freshwater ecosystems for the ecological risk assessment  
692 of pesticides. *Ecotoxicology*, **19**, 24-37.

693 Dalzochio, T., Rodrigues, G.Z.P., Petry, I.E., Gehlen, G., da Silva, L.B. (2016). The use  
694 of biomarkers to assess the health of aquatic ecosystems in Brazil: a review.  
695 *International Aquatic Research*, **8**, 283-298.

696 Dale Becker, C. & Wolford, M.G. (1980). Thermal resistance of juvenile salmonids  
697 sublethally exposed to nickel, determined by the critical thermal maximum  
698 method. *Environmental Pollution Series A, Ecological and Biological*, **21**, 181-  
699 189.

700 de Andrade Waldemarin, K.C., Alves, R.N., Beletti, M.E., Rantin, F.T., Kalinin, A.L.  
701 (2012). Copper sulfate affects Nile tilapia (*Oreochromis niloticus*) cardiomyocytes  
702 structure and contractile function. *Ecotoxicology*, **21**, 783-794.

703 De Angelis, C.F. (2017). Efeitos do surfactante nonilfenol etoxilado em peixes: função  
704 cardiorrespiratória e histopatologia em Tilápia-do-Nilo, *Oreochromis niloticus*, em  
705 condições de normóxia e hipóxia aquáticas. [Master's dissertation, Federal  
706 University of São Carlos]. UA Campus Repository.  
707 <https://repositorio.ufscar.br/handle/ufscar/9542>

708 Di Giulio, R.T. & Clark, B.W. (2015). The Elizabeth River story: A case study in  
709 evolutionary toxicology. *Journal of Toxicology and Environmental Health Part B*,  
710 **18**, 259-298.

711 Diaz, R.J. & Breitburg, D.L. (2009). Hypoxia. In: J.G. Richards, A.P. Farrell, C.J.  
712 Brauner (Eds), *Hypoxia, Fish Physiology* (vol. 27, pp. 1-23). Physiology. Elsevier.

713 Domenici, P. (2010). Context-dependent variability in the components of fish escape  
714 response: integrating locomotor performance and behavior. *Journal Experimental*  
715 *Zoology Part A: Ecological and Integrative Physiology*, **313**, 59-79.

716 Domenici, P. & Hale, M.E. (2019). Escape responses of fish: a review of the diversity in  
717 motor control, kinematics and behaviour. *Journal of Experimental Biology*, **222**,  
718 jeb166009.

719 Driedzic, W.R. & Gesser, H. (1994). Energy metabolism and contractility in ectothermic  
720 vertebrate hearts: hypoxia, acidosis, and low temperature. *Physiological*  
721 *Reviews*, **74**, 221-258.

722 Duponchelle, F., Pouilly, M., Pécheyrán, C., Hauser, M., Renno, J.F., Panfili, J.,  
723 Darnaude, A.M., García-Vasquez, A., Carvajal-Vallejos, F., García-Dávila, C.,  
724 Doria, C., Bérail, S., Donard, A., Sondag, F., Santos, R.V., Nuñez, J., Point, D.,  
725 Labonne, M., Baras, E. (2016). Trans-Amazonian natal homing in giant catfish.  
726 *Journal of Applied Ecology*, **53**, 1511-1520.

727 Farrell, A.P. (2007). Tribute to P.L. Lutz: a message from the heart--why hypoxic  
728 bradycardia in fishes? *Journal of Experimental Biology*, **210**, 1715-1725.

729 Folkerts, E.J., Blewett, T.A., He, Y., Goss, G.G. (2017). Alterations to juvenile zebrafish  
730 (*Danio rerio*) swim performance after acute embryonic exposure to sub-lethal  
731 exposures of hydraulic fracturing flowback and produced water. *Aquatic*  
732 *Toxicology*, **193**, 50–59.

733 Folkerts, E.J., Heuer, R.M., Flynn, S., Stieglitz, J.D., Benetti, D.D., Alessi, D.S., Goss,  
734 G.G., Grosell, M. (2020). Exposure to hydraulic fracturing flowback water impairs  
735 mahi-mahi (*Coryphaena hippurus*) cardiomyocyte contractile function and  
736 swimming performance. *Environmental Science & Technology*, **54**, 13579-13589.

737 Fry, F.E.J. (1971). The effect of environmental factors on the physiology of fish. In:  
738 W.S. Hoar & D.J. Randall (Eds.), *Fish physiology* (vol 6, pp. 1-98). New York:  
739 Academic Press.

740 Gamperl, A.K., Driedzic, W.R. (2009). Cardiovascular function and cardiac metabolism.  
741 In J.G. Richards, A.P. Farrell & C.J. Brauner (Eds.), *Fish Physiology* (vol. 27, pp.  
742 301-360). Academic Press.

743 Gerger, C.J., Thomas, J.K., Janz, D.M., Weber, L.P., Gerger, C.J., Thomas, J.K., Janz,  
744 A.D.M., Weber, A.L.P. (2015). Acute effects of b-naphthoflavone on  
745 cardiorespiratory function and metabolism in adult zebrafish (*Danio rerio*). *Fish*  
746 *Physiology Biochemistry*, **41**, 289-298.

747 Goertzen, M.M., Hauck, D.W., Phibbs, J., Weber, L.P., Janz, D.M. (2012). Swim  
748 performance and energy homeostasis in spottail shiner (*Notropis hudsonius*)  
749 collected downstream of a uranium mill. *Ecotoxicology and Environmental Safety*,  
750 **75**, 142-150.

751 Gomez Isaza, D.F., Cramp, R.L., Franklin, C.E. (2020a). Thermal acclimation offsets  
752 the negative effects of nitrate on aerobic scope and performance. *Journal of*  
753 *Experimental Biology*, **223**, jeb224444.

754 Gomez Isaza, D.F., Cramp, R.L., Franklin, C.E. (2020b). Simultaneous exposure to  
755 nitrate and low pH reduces the blood oxygen-carrying capacity and functional  
756 performance of a freshwater fish. *Conservation Physiology*, **8**, 1-15.

757 Gomez Isaza, D.F., Cramp, R.L., Franklin, C.E. (2021a). Exposure to nitrate increases  
758 susceptibility to hypoxia in fish. *Physiological and Biochemical Zoology*, **94**, 124-  
759 142.

760 Gomez Isaza, D.F., Cramp, R.L., Franklin, C.E. (2021b). Thermal plasticity of the  
761 cardiorespiratory system provides cross-tolerance protection to fish exposed to  
762 elevated nitrate. *Comparative Biochemistry and Physiology Part C: Toxicology &*  
763 *Pharmacology*, **240**, 108920

764 Green, A.J. & Planchart, A. (2018). The neurological toxicity of heavy metals: A fish  
765 perspective. *Comparative Biochemistry and Physiology Part C: Toxicology &*  
766 *Pharmacology*, **208**, 12-19.

767 Gupta, U., & Guha, S. (2006). Microcystin toxicity in a freshwater fish, *Heteropneustes*  
768 *fossilis* (Bloch). *Current Science*, **91**, 1261-1271.

769 Habary, A., Johansen, J.L., Nay, T.J., Steffensen, J.F., Rummer, J.L. (2017). Adapt,  
770 move or die – how will tropical coral reef fishes cope with ocean warming? *Global*  
771 *Change Biology*, **23**, 566-577.

772 Healy, T.M, Brennan, R.S., Whitehead, A., Schulte, P.M. (2018). Tolerance traits  
773 related to climate change resilience are independent and polygenic. *Global*  
774 *Change Biology*, **24**, 5348-5360.

775 Higgins, J., Zablocki, J., Newsock, A., Krolopp, A., Tabas, P., Salama, M. (2021).  
776 Durable freshwater protection: A framework for establishing and maintaining  
777 long-term protection for freshwater ecosystems and the values they sustain.  
778 *Sustainability*, **13**, 1-17.

779 Hogan, Z., Baird, I.G., Radtke, R., Vander Zanden, M.J. (2007). Long distance  
780 migration and marine habitation in the tropical Asian catfish, *Pangasius krempfi*.  
781 *Journal of Fish Biology*, **71**, 818-832.

782 Holt, M.S. (2000). Sources of chemical contaminants and routes into the freshwater  
783 environment. *Food and Chemical Toxicology*, **38**, S21-S27.



784 Hook, S.E., Gallagher, E.P., Batley, G.E. (2014). The role of biomarkers in the  
785 assessment of aquatic ecosystem health. *Integrated Environmental Assessment*  
786 *and Management*, **10**, 327-341.

787 Incardona, J.P., Collier, T.K., Scholz, N.L. (2004). Defects in cardiac function precede  
788 morphological abnormalities in fish embryos exposed to polycyclic aromatic  
789 hydrocarbons. *Toxicology and Applied Pharmacology*, **196**, 191-205.

790 Incardona, J.P. & Scholz, N.L. (2017). Environmental pollution and the fish heart In  
791 A.K. Gamperl, T.E. Gillis, A.P. Farrell & C.J. Brauner (Eds.), *Fish Physiology* (vol.  
792 36, pp. 373-433), Academic Press.

793 Isaza, D.G. (2020). Anthropogenic disturbances to freshwater taxa : Interactions  
794 between nitrate and additional stressors on various physiological traits. [PhD  
795 Thesis, School of Biological Sciences, The University of Queensland].  
796 <https://doi.org/10.14264/ea9709d>.

797 Islam, S.M.M., Rahman, M.A., Nahar, S., Uddin, M.H., Haque, M.M., Shahjahan, M.  
798 (2019). Acute toxicity of an organophosphate insecticide sumithion to striped  
799 catfish *Pangasianodon hypophthalmus*. *Toxicology Reports*, **6**, 957-962.

800 Jensen, F.B. (2003). Nitrite disrupts multiple physiological functions in aquatic animals.  
801 *Comparative Biochemistry and Physiology - A Molecular and Integrative*  
802 *Physiology*, **135**, 9-24.

803 Jobling, S. & Tyler, C.R. (2003). Endocrine disruption in wild freshwater fish. *Pure and*  
804 *Applied Chemistry*, **75**, 2219-2234.

805 Johnson, C.R. (1978). The effects of sublethal concentrations of five  
806 organophosphorus insecticides on temperature tolerance, reflexes, and  
807 orientation in *Gambusia affinis affinis* (Pisces: Poeciliidae). *Zoological Journal of*  
808 *the Linnean Society*, **64**, 63-70.

809 Joyce, W., Simonsen, M., Gesser, H., Wang, T. (2016). The effects of hypoxic  
810 bradycardia and extracellular HCO<sub>3</sub><sup>-</sup>/CO<sub>2</sub> on hypoxic performance in the eel  
811 heart. *Journal of Experimental Biology*, 219, 302–305.

812 Jutfelt, F., Roche, D.G., Clark, T.D., Norin, T., Binning, S.A., Speers-Roesch, B.,  
813 Amcoff, M., Morgan, R., Andreassen, A.H., Sundin, J. (2019). Brain cooling  
814 marginally increases acute upper thermal tolerance in Atlantic cod. *Journal of*  
815 *Experimental Biology*, **222**, jeb208249.

816 Khursigara, A.J., Ackerly, K.L., Esbaugh, A.J. (2019). Oil toxicity and implications for  
817 environmental tolerance in fish. *Comparative Biochemistry and Physiology Part*  
818 *C: Toxicology & Pharmacology*, **220**, 52-61.

819 Kroon, F., Streten, C., Harries, S. (2017). A protocol for identifying suitable biomarkers  
820 to assess fish health: A systematic review. *PLoS One*, **12**, e0174762.

821 Kumar, N., Ambasankar, K., Krishnani, K.K., Kumar, P., Akhtar, M.S., Bhushan, S.,  
822 Minhas, P.S. (2016). Dietary pyridoxine potentiates thermal tolerance, heat shock  
823 protein and protect against cellular stress of Milkfish (*Chanos chanos*) under  
824 endosulfan-induced stress. *Fish & Shellfish Immunology*, **55**, 407-414.

825 Kumar, N., Krishnani, K.K., Chandan, N.K., Singh, N.P. (2018). Dietary zinc potentiates  
826 thermal tolerance and cellular stress protection of *Pangasius hypophthalmus*  
827 reared under lead and thermal stress. *Aquaculture Research*, **49**, 1105-1115.

828 Kumar, N., Krishnani, K.K., Gupta, S.K, Singh, N.P. (2017). Selenium nanoparticles  
829 enhanced thermal tolerance and maintain cellular stress protection of *Pangasius*  
830 *hypophthalmus* reared under lead and high temperature. *Respiratory Physiology*  
831 *& Neurobiology*, **246**, 107-116.

832 Lacher, T.E. & Goldstein, M.I. (1997). Tropical ecotoxicology: Status and needs.  
833 *Environmental Toxicology and Chemistry*, **16**, 100-111.

834 Langiano, V. do C., Martinez, C.B. (2008). Toxicity and effects of a glyphosate-based  
835 herbicide on the Neotropical fish *Prochilodus lineatus*. *Comparative Biochemistry*  
836 *and Physiology Part C: Toxicology & Pharmacology*, **147**, 22-231.

837 Lefevre, S., Wang, T., McKenzie, D.J. (2021). The role of mechanistic physiology in  
838 investigating impacts of global warming on fishes. *Journal of Experimental*  
839 *Biology*, **224**, jeb238840.

840 Lewis, W.M. & Morris, D.P. (1986). Toxicity of nitrite to fish: A Review. *Transactions of*  
841 *the American Fisheries Society*, **115**, 183-195.

842 Little, E.E., Finger, S.E. (1990). Swimming behavior as an indicator of sublethal toxicity  
843 in fish. *Environ Toxicol Chem* **9**,13–19.

844 Lucas, J., Percelay, I., Larcher, T., Lefrançois, C. (2016). Effects of pyrolytic and  
845 petrogenic polycyclic aromatic hydrocarbons on swimming and metabolic  
846 performance of zebrafish contaminated by ingestion. *Ecotoxicology and*  
847 *Environmental Safety*, **132**,145-152.

848 Lutterschmidt, W.I., Hutchison, V.H. (1997). The critical thermal maximum: history and  
849 critique. *Canadian Journal of Zoology*, **75**, 1561-1574.

850 Mandic, M., Todgham, A.E., Richards, J.G. (2009) Mechanisms and evolution of  
851 hypoxia tolerance in fish. *Proceedings. Biological Sciences B*, **276**, 735-744.

852 Martins, N.D., Colvara, W., Rantin, F.T., Kalinin, A.L. (2011). Microcystin-LR: How it  
853 affects the cardio-respiratory responses to hypoxia in Nile tilapia, *Oreochromis*  
854 *niloticus*. *Chemosphere*, **84**, 154-159.

855 Martins, N.D., Yunes, J.S., McKenzie, D.J. Rantin, F.T., Kalinin, A.L., Monteiro, D.A.  
856 (2019). Microcystin-LR exposure causes cardiorespiratory impairments and

857 tissue oxidative damage in trahira, *Hoplias malabaricus*. *Ecotoxicology and*  
858 *Environmental Safety*, **173**, 436-443.

859 Martins, N.F. (2018). Efeitos do contaminante emergente triclosan sobre a função  
860 cardiorespiratória do matrinxã, *Brycon amazonicus*, em normóxia e durante  
861 hipóxia gradual. [Doctoral dissertation, Federal University of São Carlos]. UA  
862 Campus Repository. <https://repositorio.ufscar.br/handle/ufscar/10718>

863 Martins, N.D., Yunes, J.S., Monteiro, D.A., Rantin, F.T., Kalinin, A.L. (2017).  
864 Microcystin-LR leads to oxidative damage and alterations in antioxidant defense  
865 system in liver and gills of *Brycon amazonicus* (SPIX & AGASSIZ, 1829).  
866 *Toxicon*, **139**, 109-116.

867 Mason, C.F. (2002). *Biology of freshwater pollution*. London: Pearson Education.

868 Matson, C.W., Timme-Laragy, A.R., Di Giulio, R.T. (2008). Fluoranthene, but not  
869 benzo[a]pyrene, interacts with hypoxia resulting in pericardial effusion and  
870 lordosis in developing zebrafish. *Chemosphere*, **74**, 149-154.

871 Menéndez-Helman, R.J., Ferreyroa, G.V., dos Santos Afonso, M., Salibián, A. (2012).  
872 Glyphosate as an acetylcholinesterase inhibitor in *Cnesterodon decemmaculatus*.  
873 *Bulletin of Environmental Contamination and Toxicology*, **88**, 6-9.

874 McArley, T.J., Hickey, A.J.R., Herbert, N.A. (2018). Hyperoxia increases maximum  
875 oxygen consumption and aerobic scope of intertidal fish facing acutely high  
876 temperatures. *Journal Experimental Biology*, **221**, jeb189993

877 McDonnell, L.H., Reemeyer, J.E., Chapman, L.J. (2019). Independent and interactive  
878 effects of long-term exposure to hypoxia and elevated water temperature on  
879 behavior and thermal tolerance of an equatorial cichlid. *Physiological and*  
880 *Biochemical Zoology*, **92**, 253-265.

881 McKenzie, D.J., Blasco, F.R., Belão, T.C., Killen, S.S., Martins, N.D., Taylor, E.W.,  
882 Rantin, F.T. (2017). Physiological determinants of individual variation in  
883 sensitivity to an organophosphate pesticide in Nile tilapia *Oreochromis niloticus*.  
884 *Aquatic Toxicology*, **189**, 108-114.

885 McKenzie, D.J., Garofalo, E., Winter, M.J., Ceradini, S., Verweij, F., Day, N., Hayes,  
886 R., van der Oost, R., Butler, P.J., Chipman, J.K., Taylor, E.W. (2007). Complex  
887 physiological traits as biomarkers of the sub-lethal toxicological effects of  
888 pollutant exposure in fishes. *Philosophical Transactions of the Royal Society B*,  
889 **362**, 2043-2059.

890 McKenzie, D.J., Shingles, A., Claireaux, G., Domenici, P. (2009). Sublethal  
891 Concentrations of Ammonia Impair Performance of the Teleost Fast-Start Escape  
892 Response. *Physiol Biochem Zool* **82**, 353–362.

893 McKenzie, D.J., Zhang, Y., Eliason, E.J., Schulte, P.M., Blasco, F.R., Claireaux, G.,  
894 Nati, J.J.H., Farrell, A.P. (2020). Intraspecific variation in tolerance of of warming  
895 in fishes. *Journal of Fish Biology*, 1-20. <https://doi.org/10.1111/jfb.14620>

896 McRae, N.K., Gaw, S., Glover, C.N. (2018). Effects of waterborne cadmium on  
897 metabolic rate, oxidative stress, and ion regulation in the freshwater fish, inanga  
898 (*Galaxias maculatus*). *Aquatic Toxicology*, **194**, 1-9.

899 Messaad, I.A., Peters, E.J., Young, L. (2000). Thermal tolerance of red shiner  
900 (*Cyprinella lutrensis*) after exposure to atrazine, terbufos, and their mixtures.  
901 *Bulletin of Environmental Contamination and Toxicology*, **64**, 748-754.

902 Modesto, K.A. & Martinez, C.B. (2010). Effects of Roundup Transorb on fish:  
903 hematology, antioxidant defenses and acetylcholinesterase activity.  
904 *Chemosphere*, **81**, 781-787.

905 Monteiro, D.A., Taylor, E.W., McKenzie, D.J., Rantin, F.T., Kalinin, A.L. (2020).  
906 Interactive effects of mercury exposure and hypoxia on ECG patterns in two  
907 Neotropical freshwater fish species: matrinxã, *Brycon amazonicus* and traíra,  
908 *Hoplias malabaricus*. *Ecotoxicology*, **29**, 375-388.

909 Monteiro, D.A., Taylor, E.W., Rantin, F.T., Kalinin, A.L. (2017). Impact of waterborne  
910 and trophic mercury exposures on cardiac function of two ecologically distinct  
911 Neotropical freshwater fish *Brycon amazonicus* and *Hoplias malabaricus*.  
912 *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*,  
913 **201**, 26-34.

914 Monteiro, D.A., Thomaz, J.M., Rantin, F.T., Kalinin, A.L. (2013). Cardiorespiratory  
915 responses to graded hypoxia in the neotropical fish matrinxã (*Brycon*  
916 *amazonicus*) and traíra (*Hoplias malabaricus*) after waterborne or trophic  
917 exposure to inorganic mercury. *Aquatic Toxicology*, **140(41)**, 346-355.

918 Mu, J., Chernick, M., Dong, W., Di Giulio, R. T., & Hinton, D. E. (2017). Early life co-  
919 exposures to a real-world PAH mixture and hypoxia result in later life and next  
920 generation consequences in medaka (*Oryzias latipes*). *Aquatic Toxicology*, **190**,  
921 162-173.

922 Noyes, P.D., Lema, S.C. (2015). Forecasting the impacts of chemical pollution and  
923 climate change interactions on the health of wildlife. *Current Zoology*, **61**, 669-  
924 689.

925 Noyes, P.D., McElwee, M.K., Miller, H.D., Clark, B.W., Van Tiem, L.A., Walcott, K.C.,  
926 Erwin, K.N., Levin, E.D. (2009). The toxicology of climate change: Environmental  
927 contaminants in a warming world. *Environment International*, **35**, 971-986.

928 Olle, C.D. (2007). Efeito do inseticida organofosforado metilparation (Folisuper 600BR)  
929 sobre a função cardio-respiratória do peixe teleósteo matrinxã, *Brycon cephalus*.

930 [Doctoral dissertation, Federal University of São Carlos]. UA Campus Repository.  
931 <https://repositorio.ufscar.br/handle/ufscar/1207>

932 Patra RW, Chapman JC, Lim RP, Gehrke PC, Sunderam RM. 2015. Interactions  
933 between water temperature and contaminant toxicity to freshwater fish.  
934 *Environmental Toxicology and Chemistry*, **34**, 1809-1817.

935 Patra, R.W., Chapman, J.C., Lim, R.P., Gehrke, P.C. (2007). The effects of three  
936 organic chemicals on the upper thermal tolerances of four freshwater fishes.  
937 *Environmental Toxicology and Chemistry*, **26**, 1454-1459.

938 Peakall, D.B. (1994). The role of biomarkers in environmental assessment.  
939 *Ecotoxicology*, **3**, 157-160.

940 Pereira, L., Fernandes, M.N., Martinez, C.B. (2013). Hematological and biochemical  
941 alterations in the fish *Prochilodus lineatus* caused by the herbicide clomazone.  
942 *Environmental Toxicology and Pharmacology*, **36**, 1-8.

943 Pérez-Parada, A., Goyenola, G., Teixeira de Mello, F., Heinzen, H. (2018). Recent  
944 advances and open questions around pesticide dynamics and effects on  
945 freshwater fishes. *Current Opinion in Environmental Science & Health*, **4**, 38-44.

946 Perry, S.F., Jonz, M.G., Gilmour, K.M. (2009). Oxygen sensing and the hypoxic  
947 ventilatory response. In: J.G. Richards, A.P. Farrell, C.J. Brauner (Eds), *Hypoxia*,  
948 *Fish Physiology* (vol 27, pp. 194-255). Academic Press, Elsevier.

949 Philippe, C., Hautekiet, P., Grégoir, A.F., Thoré, E.S.J., Brendonck, L., De Boeck, G.,  
950 Pinceel, T. (2019). Interactive effects of 3,4-DCA and temperature on the annual  
951 killifish *Nothobranchius furzeri*. *Aquatic Toxicology*, **212**, 146-153.

952 Philippe, C., Hautekiet, P., Grégoir, A.F., Thoré, E.S.J., Pinceel, T., Stoks, R.,  
953 Brendonck, L., De Boeck, G. (2018). Combined effects of cadmium exposure and  
954 temperature on the annual killifish (*Nothobranchius furzeri*). *Environmental*  
955 *Toxicology and Chemistry*, **37**, 2361-2371.

956 Pollock, M.S., Clarke, L.M.J., Dubé, M.G. (2007). The effects of hypoxia on fishes: from  
957 ecological relevance to physiological effects. *Environmental Reviews*, **15**, 1-14.

958 Pörtner, H.O. (2010). Oxygen- and capacity-limitation of thermal tolerance: a matrix for  
959 integrating climate-related stressor effects in marine ecosystems. *Journal of*  
960 *Experimental Biology*, **213**, 881-893.

961 Pullaguri, N., Grover, P., Abhishek, S., Rajakumara, E., Bhargava, Y., Bhargava, A.  
962 (2021). Triclosan affects motor function in zebrafish larva by inhibiting ache and  
963 syn2a genes. *Chemosphere*, **266**, 128930.

964 Rahman, M.S. & Thomas, P. (2018). Interactive effects of hypoxia and PCB co-  
965 exposure on expression of CYP1A and its potential regulators in Atlantic croaker  
966 liver. *Environmental Toxicology*, **33**, 411-421.

967 Rantin, F.T., Kalinin, A., Monteiro, D.A. (2020). The cardiovascular system. In B.  
968 Baldisserotto, E.C. Urbinati, & J.E.P. Cyrino (Eds.), *Biology and physiology of*  
969 *freshwater neotropical fish* (1st ed., pp. 185-216). Academic Press, Elsevier Inc.

970 Rantin, F.T., Kalinin, A.L., Guerra C.D.R., Maricondi-Massari, M.; Verzola, R.M.M.  
971 (1995). Electrocardiographic characterization of myocardial function in normoxic  
972 and hypoxic teleosts. *Brazilian Journal of Medical and Biological Research*, **28**,  
973 1277-1289.

974 Regan, M.D., Gill, I.S., Richards, J.G. (2017). Calorespirometry reveals that goldfish  
975 prioritize aerobic metabolism over metabolic rate depression in all but near-  
976 anoxic environments. *Journal of Experimental Biology*, **220**, 564-572.

977 Regan, M.D., Mandic, M., Dhillon, R.S., Lau, G.Y., Farrell, A.P., Schulte, P.M., Seibel,  
978 B.A., Speers-Roesch, B., Ultsch, G.R., Richards, J.G. (2019). Don't throw the fish  
979 out with the respirometry water. *Journal of Experimental Biology*, **222**, 9-10.

980 Rezende, E.L, Bozinovic, F. (2019). Thermal performance across levels of biological  
981 organization. *Philosophical transactions of the Royal Society of London. Series B,*  
982 *Biological Sciences*, **374**, 20180549

983 Rezende, E.L., Castañeda, L.E., Santos, M. (2014). Tolerance landscapes in thermal  
984 ecology. *Functional Ecology*, **28**, 799-809.

985 Richards, J.G. (2009). Metabolic and molecular responses of fish to hypoxia. In J.G.  
986 Richards, A.P. Farrell & C.J. Brauner (Eds.), *Fish Physiology* (vol. 27, pp. 301-  
987 360). Academic Press.

988 Rodgers, E.M. & De Boeck, G. (2019). Nitrite-induced reductions in heat tolerance are  
989 independent of aerobic scope in a freshwater teleost. *Journal of Experimental*  
990 *Biology*, **222**, jeb212035.

991 Rogers, N.J., Urbina, M.A., Reardon, E.E., McKenzie, D.J., Wilson, R.W. (2016). A new  
992 analysis of hypoxia tolerance in fishes using a database of critical oxygen level  
993 ( $P_{crit}$ ). *Conservation Physiology*, **4**, cow012.

994 Saaristo, M., Brodin, T., Balshine, S., Bertram, M.G., Brooks, B.W., Ehlman, S.M.,  
995 McCallum, E.S., Sih, A., Sundin, J., Wong, B.B.M., Arnold, K.E. (2018). Direct  
996 and indirect effects of chemical contaminants on the behaviour, ecology and  
997 evolution of wildlife. *Proc. R. Soc. B* **285**, 20181297.

998 Sancho, E., Ferrando, M.D., Andreu, E. (1998). In vivo inhibition of AChE activity in the  
999 European eel *Anguilla anguilla* exposed to technical grade fenitrothion.  
1000 *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and*  
1001 *Endocrinology*, **120**, 389-395.

1002 Schjolden, J., Sørensen, J., Nilsson, G.E, Poléo, A.B.S. (2007). The toxicity of copper  
1003 to crucian carp (*Carassius carassius*) in soft water. *Science of the Total*  
1004 *Environment*, **384**, 239-251.

1005 Schulte PM, Healy TM, Fangue NA. 2011. Thermal performance curves, phenotypic  
1006 plasticity, and the time scales of temperature exposure. *Integrative and*  
1007 *Comparative Biology*, **51**, 691-702.

1008 Schulte, P.M. (2015). The effects of temperature on aerobic metabolism: Towards a  
1009 mechanistic understanding of the responses of ectotherms to a changing  
1010 environment. *Journal of Experimental Biology*, **218**, 1856-1866.

1011 Shingles, A., McKenzie, D.J., Taylor, E.W., Moretti, A., Butler, P.J., Ceradini, S. (2001).  
1012 Effects of sublethal ammonia exposure on swimming performance in rainbow  
1013 trout (*Oncorhynchus mykiss*). *Journal of Experimental Biology*, **204**, 2691-2698.

1014 Shiogiri, N.S., Paulino, M.G., Carraschi, S.P, Baraldi, F.G., Cruz, C., Fernandes, M.N.  
1015 (2012). Acute exposure of a glyphosate-based herbicide affects the gills and liver  
1016 of the Neotropical fish, *Piaractus mesopotamicus*. *Environmental Toxicology and*  
1017 *Pharmacology*, **34**, 388-396.

1018 Sokolova, I.M. & Lannig, G. (2008). Interactive effects of metal pollution and  
1019 temperature on metabolism in aquatic ectotherms: implications of global climate  
1020 change. *Climate Research*, **37**, 181-201.

1021 Speranza, E.D., Colombo, M., Tatone, L.M., Cappelletti, N., Migoya, M.C., Colombo,  
1022 J.C. (2016). Fatty acid alterations in the detritivorous *Prochilodus lineatus*  
1023 promoted by opportunistic feeding on sewage discharges in the Río de la Plata  
1024 estuary. *Journal of Fish Biology*, **89**, 2024-2037.

1025 Spinks RK, Munday PL, Donelson JM. 2019. Developmental effects of heatwave  
1026 conditions on the early life stages of a coral reef fish. *J Exp Biol* 222:jeb202713.

1027 Sprague, J.B. (1971). Measurement of pollutant toxicity to fish III: Sublethal effects and  
1028 "safe" concentrations. *Water Research*, **5**, 245-266.

1029 Stauffer, J. (1998). The water crisis: Constructing solutions to freshwater pollution.  
1030 London: Earthscan.

1031 Stieglitz, J.D., Mager, E.M., Hoenig, R.H., Benetti, D.D., Grosell, M. (2016). Impacts of  
1032 Deepwater Horizon crude oil exposure on adult mahi-mahi (*Coryphaena*  
1033 *hippurus*) swim performance. *Environmental Toxicology and Chemistry*, **35**,  
1034 2613-2622.

1035 Stillman, J.H. (2019). Heat waves, the new normal: Summertime temperature extremes  
1036 will impact animals, ecosystems, and human communities. *Physiology*, **34**, 86-  
1037 100.

- 1038 Suresh, A., Sivaramakrishna, B., Victoriamma, P.C., Radhakrishnaiah, K. (1992).  
1039 Comparative study on the inhibition of acetylcholinesterase activity in the  
1040 freshwater fish *Cyprinus carpio* by mercury and zinc. *Biochemistry International*,  
1041 **26**, 367-375.
- 1042 Takle, J.C.C., Beitinger, T.L., Dickson, K.L. (1983). Effect of the aquatic herbicide  
1043 endothal on the critical thermal maximum of red shiner, *Notropis lutrensis*.  
1044 *Bulletin of Environmental Contamination and Toxicology*, **31**, 512-517.
- 1045 Thomas, J.K. & Janz, D.M. (2011). Dietary selenomethionine exposure in adult  
1046 zebrafish alters swimming performance, energetics and the physiological stress  
1047 response. *Aquatic Toxicology*, **102**, 79-86.
- 1048 Thomaz, J.M., Martins, N.D., Monteiro, D.A., Rantin, F.T., Kalinin, A.L. (2009). Cardio-  
1049 respiratory function and oxidative stress biomarkers in Nile tilapia exposed to the  
1050 organophosphate insecticide trichlorfon (NEGUVON®). *Ecotoxicology*  
1051 *Environmental Safety*, **72**, 1413-1424.
- 1052 Tierney, K., Casselman, M., Takeda, S., Farrell, T., Kennedy, C. (2007). The  
1053 relationship between cholinesterase inhibition and two types of swimming  
1054 performance in chlorpyrifos-exposed coho salmon (*Oncorhynchus kisutch*).  
1055 *Environmental Toxicology and Chemistry*, **26**, 998-1004.
- 1056 Tudorache, C., Blust, R., De Boeck, G. (2008). Social interactions, predation behaviour  
1057 and fast start performance are affected by ammonia exposure in brown trout  
1058 (*Salmo trutta* L.). *Aquat. Toxicol.* **90**, 145–153.
- 1059 Ultsch, G.R., Regan, M.D. (2019). The utility and determination of Pcrit in fishes.  
1060 *Journal of Experimental Biology*, **222**, 1-9.
- 1061 van der Oost, R., Beyer, J., Vermeulen, N.P.E. (2003). Fish bioaccumulation and  
1062 biomarkers in environmental risk assessment: a review. *Environmental*  
1063 *Toxicology and Pharmacology*, **13**, 57-149.
- 1064 van der Oost, R., McKenzie, D.J., Verweij, F., Satumalay, C., van der Molen, N.,  
1065 Winter, M.J., Chipman, J.K. (2020). Identifying adverse outcome pathways (AOP)  
1066 for Amsterdam city fish by integrated field monitoring. *Environmental Toxicology*  
1067 *and Pharmacology*, **74**, 103301.
- 1068 Wagner, M. & Lambert, S. (2018). Microplastics are contaminants of emerging concern  
1069 in freshwater environments: An overview. In: M. Wagner & S. Lambert (Eds.),  
1070 *Freshwater microplastics. emerging environmental contaminants* (pp 1-24). The  
1071 Handbo. Springer Open. p 1–24.
- 1072 Wang, T., Lefevre, S., Iversen, N.K., Findorf, I., Buchanan, R., McKenzie, D.J. (2014).  
1073 Anaemia only causes a small reduction in the upper critical temperature of sea



1074 bass: is oxygen delivery the limiting factor for tolerance of acute warming in  
1075 fishes? *Journal of Experimental Biology*, **217**, 4275-4278.

1076 Wang, Z., Kwok, K.W., Leung, K.M. (2019). Comparison of temperate and tropical  
1077 freshwater species' acute sensitivities to chemicals: An update. *Integrated*  
1078 *Environmental Assessment and Management*, **15**, 352-363.

1079 Wang, H., Liang, Y., Li, S., Chang, J. (2013). Acute toxicity, respiratory reaction, and  
1080 sensitivity of three cyprinid fish species caused by exposure to four heavy metals.  
1081 *PLoS ONE*, **8**, e65282.

1082 Watenpaugh, D.E., Beiting, T.L., Huey, D.W. (1985). Temperature tolerance of nitrite-  
1083 exposed catfish. *Transactions of the American Fisheries Society*, **114**, 274-278.

1084 Williams, C.M., Buckley, L.B., Sheldon, K.S., Vickers, M., Pörtner, H.O., Dowd, W.W.,  
1085 Gunderson, A.R., Marshall, K.E., Stillman, J.H. (2016). Biological impacts of  
1086 thermal extremes: mechanisms and costs of functional responses matter.  
1087 *Integrative and Comparative Biology*, **56**, 73-84.

1088 Wood, A.W., Johnston, B.D., Farrell, A.P., Kennedy, C.J. (1996). Effects of  
1089 didecyldimethylammonium chloride (DDAC) on the swimming performance, gill  
1090 morphology, disease resistance, and biochemistry of rainbow trout  
1091 (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences*,  
1092 **53**, 2424-2432.

1093 Wood, CM. (2018). The fallacy of the Pcrit - Are there more useful alternatives? *Journal*  
1094 *of Experimental Biology*, **221**, jeb163717

1095 Xia, J.G., Nie, L.J., Mi, X.M., Wang, W.Z., Ma, Y.J., Cao, Z.D., Fu, S.J. (2015).  
1096 Behavior, metabolism and swimming physiology in juvenile *Spinibarbus sinensis*  
1097 exposed to PFOS under different temperatures. *Fish Physiology Biochemistry*,  
1098 **41**, 1293-1304.

1099 Yang, R., Brauner, C., Thurston, V., Neuman, J., Randall, D. (2000). Relationship  
1100 between toxicant transfer kinetic processes and fish oxygen consumption.  
1101 *Aquatic Toxicology*, **48**, 95-108.

1102 Zebral, Y.D., Lansini, L.R., Costa, P.G., Roza, M., Bianchini, A., Robaldo, R.B. (2018).  
1103 A glyphosate-based herbicide reduces fertility, embryonic upper thermal  
1104 tolerance and alters embryonic diapause of the threatened annual fish  
1105 *Austrolebias nigrofasciatus*. *Chemosphere*, **196**, 260-269.

1106

1107 **Figure Legends**


1108 **Figure 1.** Reduced tolerance of hypoxia, as measured by critical oxygen level ( $O_{2crit}$ ), in  
1109 matrixã *Brycon amazonicus*, after exposure to various different pollutants. Ct = control  
1110 condition, MP = methyl parathion, Hg =  $HgCl_2$ , TC = triclosan, RT = Roundup Transorb,  
1111 RO = Roundup Original, RWG = Roundup WG. (Adapted from: Olle, 2007; Anelli Jr.,  
1112 2010; Monteiro et al., 2013; Martins, 2018).

1113 **Figure 2.** Influence of ambient oxygenation on oxygen uptake rate ( $\dot{M}O_2$ ) of matrixã  
1114 *Brycon amazonicus*, after exposure to various different pollutants. Ct = control, MP =  
1115 methyl parathion, Hg =  $HgCl_2$ , TC = triclosan, RT = Roundup Transorb, RO = Roundup  
1116 Original, RWG = Roundup WG. Closed symbols denote a significant difference in relation  
1117 to the normoxic values ( $P < 0.05$ ). (Adapted from: Olle, 2007; Anelli Jr., 2010; Monteiro  
1118 et al., 2013; Martins, 2018).

1119 **Figure 3.** The relationship of acclimation temperature to thermal safety margin (TSM) in  
1120 subtropical and tropical freshwater fishes, where TSM is the difference, in °C, between  
1121 acclimation T and  $CT_{max}$ . Open symbols, control conditions for all data carried in Table  
1122 2. Grey symbols, conditions where contaminants caused a significant decline in  $CT_{max}$ .  
1123 Based upon data from 14 studies on 12 species exposed to 15 contaminants (Table 2).  
1124 Control data, dashed line,  $R^2 = 0.828$ ,  $P < 1 \times 10^{-8}$ , exposed data, dotted line,  $R^2 = 0.716$ ,  
1125  $P < 1 \times 10^{-5}$ . Slopes are not significantly different ( $P = 0.129$ , two-way T-Test).


1126 **Figure 4.** Relationship between acclimation temperature ( $T_{acc}$ ) and the extent, as a  
1127 percentage of control conditions, to which contaminants decrease thermal safety  
1128 margin (TSM) in sub-tropical and tropical freshwater fishes. Data are taken from Table  
1129 2, for all those situations where contamination caused a significant decline in  $CT_{max}$  ( $n$   
1130 = 17, from 11 studies of 11 contaminants on 10 species). Line describes a linear  
1131 regression, dotted lines are 95% confidence intervals,  $R^2 = 0.43$ ,  $P = 0.0012$ . This  
1132 figure is based upon a very limited dataset and should, therefore, be interpreted with  
1133 caution until the knowledge base is extended (see main text for details)

1134




1135

1136




1137

1138



1139

1140



1141

1142

