

Will Mesenchymal Stromal Cells become tools for immunomodulation in Myasthenia Gravis ?

Alexandra Bayer Wildberger¹, José Villegas¹, Julien Verdier¹, Mariette Giannini¹, Solène Maillard¹, Frédérique Truffault¹, Aurélien Corneau², Nadine Dragin¹, Valérie Vanneaux³, Hélène Rouard⁴, Danièle Noël⁵, Christophe Martinaud⁶, Rozen le Panse¹, Sonia Berrih-Aknin¹, Jean-Thomas Vilquin¹.

1: Sorbonne Université, INSERM UMRS 974, Centre de Recherche en Myologie, Institut de Myologie, groupe hospitalier Pitié-Salpêtrière ; Paris, France.

2: Sorbonne Université, INSERM UMRS 1135, Centre d'Immunologie et des Maladies Infectieuses (CIMI) ; Plateforme de Cytométrie (CypS), UMS30-LUMIC, Faculté de Médecine Pierre et Marie Curie, groupe hospitalier Pitié-Salpêtrière ; Paris, France.

3: Cell Therapy Unit, CIC de Biothérapies (CBT-501), INSERM U1160, Institut Universitaire d'Hématologie, Hôpital Saint-Louis, AP-HP ; University Paris Diderot, Sorbonne Paris Cité ; Paris, France.

4: Université Paris-Est-Créteil (UPEC), INSERM/IMRB-U955, laboratoire BCTS VT ; Etablissement Français du Sang d'Ile-de-France, Unité d'Ingénierie et de Thérapie Cellulaire ; Créteil, France.

5 : INSERM U1183, Institut de Médecine Régénératrice et Biothérapies (IRMB), Hôpital Saint-Eloi ; Montpellier, France.

6 : INSERM UMRS-MD 1197, Unité des Médicaments de Thérapie Innovante, Centre de Transfusion Sanguine des Armées ; Clamart, France.

Presenting author's details:

Alexandra BAYER WILDBERGER,

alexandrabayerw@gmail.com

Sorbonne Université, INSERM UMRS 974, Centre de Recherche en Myologie, Institut de Myologie, groupe hospitalier Pitié-Salpêtrière. Paris, France.

Tel : +33 1 40 77 81 16

Background and aims

Myasthenia Gravis (MG) is due to auto-antibodies directed against the neuromuscular endplate. Mesenchymal Stromal/Stem Cells (MSC) can modulate the immune system via soluble mediators and cell-cell contacts. Research grade (RG-) MSC conditioned by peripheral blood mononucleated cells (PBMC) improved the clinical outcomes in a humanized mouse model of MG (*Sudres et al., JCI Insight 2017*). In a clinical perspective, we characterized the phenotype and functional capacities of resting and conditioned MSC of clinical grade (CG-MSC), as compared to RG-MSC.

Methods

RG- and CG-MSC were derived from adipose tissue. They were conditioned by PBMC, monocytes, or activated by IFN-γ, or left untreated. Their phenotype was assessed by flow cytometry (61 antibodies), and mass cytometry (CyTOF) (31 antibodies). Immunomodulating capacities of CG-MSC were evaluated by PBMC inhibition assay, and quantifications of cytokines (IL-6, TGF-β, IDO and PGE2).

Results

(1) RG- and CG-MSC showed close phenotypes despite some differences in growth kinetics. However the strongest correlations were obtained in comparing CG cultures from different donors between them, or RG cultures between them. (2) Some markers were modulated by all conditioning treatments, while others exclusively by one of them, thus conferring

conditioning signatures. (3) Both resting and conditioned CG-MSC blocked the proliferation of activated PBMC in vitro. (4) The conditioning treatments enhanced the secretion of immunomodulatory molecules by MSC, but they differed in their molecular footprint.

Conclusion

RG- and CG-MSC show similarities and differences ; CG-MSC present immunomodulation capacities and the different treatments result in phenotypic and functional signatures.