



Hormonal and environmental signaling pathways target membrane water transport

Christophe Maurel, Colette Tournaire-Roux, Lionel Verdoucq, Véronique Santoni-Kervella

► To cite this version:

Christophe Maurel, Colette Tournaire-Roux, Lionel Verdoucq, Véronique Santoni-Kervella. Hormonal and environmental signaling pathways target membrane water transport. Plant Physiology, 2021, <10.1093/plphys/kiab373>. <hal-03440937>

HAL Id: hal-03440937

<https://hal.science/hal-03440937v1>

Submitted on 28 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Short title: Signaling pathways target membrane water transport

Corresponding author: Christophe Maurel: christophe.maurel@cnrs.fr

Hormonal and environmental signaling pathways target membrane water transport

Christophe Maurel, Colette Tournaire-Roux, Lionel Verdoucq, and Véronique Santoni

BPMP, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France.

ORCID IDs: 0000-0002-4255-6440 (CM); 0000-0002-2966-5597 (C. T.-R); 0000-0001-8238-3700 (L.V.);
0000-0002-1437-0921 (V.S.);

Materials distribution statement:

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plphys/pages/General-Instructions>) is: _
Christophe Maurel (christophe.maurel@cnrs.fr).

One-sentence summary: This paper describes the multiple signaling pathways targeting cell and tissue hydraulics to maintain the plant water status, and it reveals the commonalities of these pathways.

Author contributions: All authors contributed to the literature survey. CM wrote the article which was revised by all other authors. C. T.-R., L.V. and V.S. designed the figures and table.

Funding information: This work was supported in part by the Agence Nationale de la Recherche (ANR- 18-CE92-0055)

Introduction

As elaborate terrestrial living organisms, vascular plants have to maintain their water status as tightly as possible and throughout their life cycle (Davies and Bennett 2015; Steudle 1992). Particularly challenging processes include the expansion/growth of underground and aerial parts and the desiccation, imbibition and germination of seeds and pollen. Most often, these processes occur under variable environmental conditions, including changes to water availability and temperature in the soil and atmosphere, which have marked impacts on the plant water status. The latter is primarily maintained through tight stomatal regulation controlling transpirational water flow (Tardieu et al. 2015). Osmotic adjustment and regulation of water permeability (hydraulics), which both contribute to water potential gradients and water movement intensity, provide a complementary and finely-tuned regulation of plant water relations at the cell and tissue levels. In support for these general principles, a large body of studies has shown that plant water transport and its molecular components are responsive, across diverse time scales, to an extremely wide array of environmental and hormonal signals (Maathuis et al. 2003; Maurel et al. 2015; Chaumont et al. 1998; Aroca et al. 2012). Thus, plant water relations are under constant adjustment in virtually any physiological context.

Plant aquaporins, which are present in the plasma and intracellular membranes of most plant cells, play a central role in these processes by ensuring cell-to-cell water transport and to a lesser extent, single cell osmotic regulation (Maurel et al. 2015; Chaumont and Tyerman 2014; Tyerman et al. 1999). More generally, aquaporins have emerged as central membrane targets of environmental and hormonal signalling pathways acting on plant water relations. In addition, aquaporins are highly abundant proteins amenable to biochemical (proteomic) and cell biological approaches. Besides studies in the plant, their function and regulation can be easily studied after functional reconstitution (proteoliposomes) or heterologous expression (yeast, *Xenopus* oocytes). Thus, aquaporins are particularly relevant and convenient entry points for bottom up dissections of plant signalling cascades.

Due to the central position of aquaporins in multiple hormonal and environmental responses determining the plant water status, the first aim of the present review was to inventory corresponding signalling pathways. Beyond the diversity of these pathways, a knowledge network with common or converging paths is now emerging. Conversely, the dedicated study of signalling pathways per se seems to reveal regulation states/contexts that escaped classical physiological analyses of plant water relations. A particularly challenging question is to understand how plants are able to perceive and respond to combined stresses.

1) Water deficit, abscisic acid (ABA) and related signals

1.1 Drought.

ABA-dependent pathways.

Whereas physiological links between drought stress, ABA and plant water transport have been identified for some time, it is only recently that the signalling mechanisms involved have started to be elucidated. Studies on epidermal peels from wild-type and aquaporin knock-out *Arabidopsis* (*Arabidopsis thaliana*) plants have revealed that the plasma membrane aquaporin Plasma Membrane Intrinsic Protein 2;1 (*AtPIP2;1*) is required for ABA-induced stomatal closure, but dispensable during

light- and CO₂-induced stomatal movements (Grondin et al. 2015). Consistent with this, isolated guard cell protoplasts responded to exogenous ABA by a two-fold increase in osmotic water permeability that was strictly dependent on the functionality of AtPIP2;1. This aquaporin, which is known to be permeable to H₂O₂ (Dynowski et al. 2008), was also required for ABA-induced accumulation of H₂O₂ in the guard cell interior (Rodrigues et al. 2017). These studies establish a dual water- and H₂O₂-transport function for guard cell plasma membrane aquaporins as players and targets of signalling processes triggering stomatal closure (Grondin et al. 2015; Rodrigues et al. 2017). More specifically, *in vitro* phosphorylation assays and transgenic plant analysis led to a model whereby Snf1-related protein kinase 2.6 (SnRK2.6)/Open stomata 1 (OST1), a protein kinase that plays a pivotal role in ABA signalling, is able to phosphorylate AtPIP2;1 at a specific cytosolic Ser residue (Ser121), with this phosphorylation being necessary for proper ABA-induced stomatal closure. SnRK2.6/OST1 also activates plasma membrane NADPH oxidases, which feed superoxide dismutases for production of apoplastic H₂O₂ (Maurel et al. 2016) (Figure 1A). While the above mentioned studies were performed on epidermal peels of wild-type and *pip2;1* plants, we are concerned that intact leaves of a quadruple *pip1;1 pip1;2 pip2;1 pip2;2* mutant showed normal responses to ABA (Ceciliato et al. 2019). We hypothesize that integration of guard cells within whole leaf physiology can compensate for the defects captured in isolated guard cells or protoplasts. We also note that, in contrast to stomata, ABA can down-regulate aquaporin activity in bundle-sheaths (Shatil-Cohen et al. 2011) and roots (Rosales et al. 2019). Whereas the canonical ABA signaling machinery formed by RCAR/PYR/PYL receptors, Protein Phosphatase 2C (PP2C) co-receptors acting on Class 2 Sucrose-Non-Fermenting Protein kinases (SnRK2s) is likely to be involved, in roots in particular (Rosales et al. 2019), the exact components providing these hydraulic regulations remain as yet unknown. Aquaporins and water transport can also be under ABA control through more indirect pathways. In Arabidopsis, ABA induced the expression of the membrane protein Tryptophan-rich sensory protein/translocator (TPSO), which physically interacts with AtPIP2;7 in the early secretory pathway, to trigger its degradation through the autophagic pathway, thereby reducing its

water transport activity at the plasma membrane (Hachez et al. 2014b). In rice (*Oryza sativa*), the gene encoding a leucine-rich repeat (LRR) receptor like kinase (RLK) named Leaf Panicle 2 (LP2) is down-regulated by drought and ABA, and acts a negative regulator of drought response (Wu et al. 2015). Its action involves a molecular interaction at the plasma membrane with the drought-responsive aquaporins *OsPIP1;1*, *OsPIP1;3*, and *OsPIP2;3*.

ABA-independent pathways

Drought can also exert ABA-independent effects on aquaporin expression and function. For instance, several aquaporin gene promoters harbour Drought Responsive Elements (DRE) that promote the binding of DREB transcription factors (Rae et al. 2011). Other classes of transcription factors (*e.g.* AP2/EREBP, ASR1) also confer drought-dependent expression on aquaporin genes (Zhu et al. 2014; Ricardi et al. 2014). Furthermore, induction under severe drought of ubiquitin-conjugating (E2) enzyme UBC32 and RING-type E3 ligase, which both interact with *AtPIP2;1* and *AtPIP2;2*, leads to ubiquitination and degradation of these aquaporins, thereby preventing their negative impact on drought tolerance (Chen et al. 2021).

A functional interaction between an aquaporin (*RhPIP2;1*) and a membrane tethered MYB protein named *RhPTM* was recently described in rose (Zhang et al. 2019b). In brief, dehydration stress induced C-terminal phosphorylation of *RhPIP2;1* at Ser273, which in turn led to membrane release and nuclear activation of *RhPTM*. *RhPTM* itself acts as a transcription factor with general growth inhibitory effects. This landmark work establishes an unexpected link between hydraulics and a transcriptionally determined metabolic control of growth that promotes water deficiency tolerance. Yet, the protein kinases/phosphatases that act on Ser273 of *RhPIP2;1* and their drought-dependent activation/inhibition mechanisms remain to be identified. Similarly, some PIP aquaporins of *Arabidopsis* (*e.g.* *AtPIP2;4*) show enhanced phosphorylation as early as 5 min after exposure to a hypertonic stress (0.3 M mannitol) (Stecker et al. 2014).

Cell biological approaches are now developed to address associated regulatory mechanisms. An improved super-resolution microscopic technique called single particle tracking Photo-Activated

Localization Microscopy (sptPALM) was used to monitor the lateral mobility of proteins at the plasma membrane (LTi6a, AtPIP2;1) and tonoplast (AtTIP1;1) (Hosy et al. 2015). In contrast to the other membrane proteins, AtPIP2;1 showed a low lateral mobility, was confined to membrane nanodomains and its mobility was enhanced upon a hyperosmotic stress. sptPALM further showed how reactive oxygen species (ROS) control specific diffusion and nano-organization of membrane cargo proteins (Martinieri et al. 2019). In particular, ROS initiated clustering of AtPIP2;1 and its removal from the plasma membrane. This process is achieved in part by clathrin-mediated endocytosis. It is linked to further aquaporin internalization and transfer into the endocytic pathways (Zwiewka et al. 2015). In the longer term, the same aquaporins are subjected to ubiquitination and further degraded (Lee et al. 2009) in a phosphorylation-dependent manner (Chen et al. 2021). In addition, a role of autophagic pathways was recently uncovered in *Medicago truncatula* under water deficit (Li et al. 2020b). A dehydrin (*MtCAS31*) was shown to act as a cargo receptor for the *MtPIP2;7* aquaporin. *MtCAS31* further promoted complex formation with the autophagy-related protein ATG8a to facilitate *MtPIP2;7* degradation. This, in turn, reduced water loss under water deficit and improved drought tolerance (Li et al. 2020b). The possible relation of this mechanism with the ABA-dependent TPSO pathway (Hachez et al. 2014b) mentioned above is not yet clear.

Finally, a bottom-up approach was used to further explore the signaling mechanisms acting upstream of aquaporins in the Arabidopsis root under hyperosmotic stress. A RESPIRATORY BURST OXIDASE HOMOLOG (RBOH) pathway defined by two NADPH oxidases and an additional pathway involving apoplastic ascorbate and iron were found to mediate osmotic stress-induced ROS production (Martinieri et al. 2019). Furthermore, osmotic stress was shown to trigger interaction of a Rho GTPase, Rho-of-Plant 6 (ROP6) with the two NADPH oxidases (Smokvarska et al. 2020). This interaction occurs in specific plasma membrane nanodomains that are necessary and sufficient to transduce production of ROS and several plant adaptive responses to osmotic constraints (Figure 1B).

1.2 Salinity

Under salt stress, PIP aquaporins are subjected to cellular regulations that are reminiscent of those seen under a purely osmotic stress. In *Arabidopsis* roots for instance, a 100 mM NaCl treatment decreased C-terminal phosphorylation at Ser283 of AtPIP2;1 after 2-4 h (Prak et al. 2008). On a shorter timescale (30 min), a salt treatment also enhanced the lateral diffusion of AtPIP2;1 at the plasma membrane (Li et al. 2011) and its constitutive cycling (Luu et al. 2012). These movements lead to internalization of AtPIP2;1 from the plasma membrane, and its transfer to the pre-vacuolar and vacuolar compartments, a process that is mediated by ROS (Boursiac et al. 2008) (Figure 1B) and is dependent on clathrin, phosphatidylinositol 3-kinase (PI3K) and phosphatidylinositol 4-kinase (PI4K) (Ueda et al. 2016). These mechanisms may well be generalized: AtPIP2;7, another major aquaporin isoform in the *Arabidopsis* root, showed a combined transcriptional down-regulation and protein internalization that resulted in a dramatic decrease in cell and tissue hydraulics (Pou et al. 2016). As in the case of osmotic stress, the primary signalling events triggering salt-dependent aquaporin regulation are not elucidated. Yet, salt-dependent expression of AtPIP2;3 and AtPIP2;5 is in part mediated by SnRK2s of subclass 1 (Kawa et al. 2020). Some of these protein kinases (SnRK2.4, SnRK2.10) interact with and phosphorylate VARICOSE (VCS) and VARICOSE RELATED (VCR) (Kawa et al. 2020; Soma et al. 2017), two components of the mRNA decapping complex that catalyzes the first step of mRNA decay.

A role of PIP aquaporins in ion transport, tentatively associated to the so-called Non Selective Cation Channels described by patch clamp in many cell types, was recently proposed in *Arabidopsis* and barley (Tran et al. 2020; Byrt et al. 2017). As thoughtfully discussed by (Tyerman et al. 2021), this activity may well be central during plant response to salinity stress. The ionic conductance of AtPIP2;1 is repressed by calcium and strongly enhanced upon C-terminal phosphorylation (Qiu et al. 2020). In barley, HvPIP2;8 shows similar regulation, and, in addition, its gene is induced in shoots of salt stressed plants, specifically in a tolerant cultivar (Tran et al. 2020). The connections of these regulatory processes to upstream salt stress signaling events and, possibly, to water transport regulation remain to be investigated.

2) Other abiotic constraints and stimuli

2.1 Oxygen availability

Flooding stress, whether due to water logging or submergence, deeply impacts plant water relations by favoring a massive inflow of water through roots while transpiration is directly or indirectly impaired (Tan et al. 2018). Two pathways leading to aquaporin regulation have been uncovered in this context (Maurel and Nacry 2020). On the short term, a pronounced cytosolic acidosis can be observed under oxygen (O₂) deprivation due to the metabolic production of organic acids while a drop in cellular respiration and ATP production hampers efficient proton (H⁺) extrusion by H⁺-ATPases. This in turn induces protonation of a His residue located on the second cytosolic loop of PIPs, triggering channel gating. It is remarkable that this His residue is perfectly conserved in PIPs of all species, supporting the generality of the H⁺-dependent gating mechanism (Tournaire-Roux et al. 2003). We note that certain plant species do not show any effect of hypoxia on root hydraulic conductivity (L_p). In the case of two lupin species (*Lupinus angustifolius*, *Lupinus luteus*), hypoxia induced a marked inhibition of cortical cell hydraulic conductivity but had no substantial effect at the whole root level, suggesting the predominance of an apoplastic pathway during water uptake (Bramley et al. 2010).

The second pathway was recently uncovered through quantitative genetic analysis of Arabidopsis root hydraulics. It is under the control of *Hydraulic Conductivity of Root 1 (HCR1)* (Shahzad et al. 2016). HCR1 mediates root responses to hypoxia in relation to potassium (K⁺) availability, thereby acting as a negative regulator of L_p and root aquaporins. HCR1 encodes a RAF-type MAP3K protein kinase that specifically accumulates in the presence of K⁺ and absence of O₂. HCR1 enhances the abundance of RAP2.12, a key Ethylene Response Factor (ERF) VII transcription factor that is stabilized under O₂ starvation and triggers the core anaerobic transcriptional response. We have proposed that HCR1 phosphorylates RAP2.12 or one of its direct regulators in planta and

thereby potentiates the most early plant responses to O₂ starvation (Shahzad et al. 2016). However, the mode of aquaporin down-regulation through this pathway has not been elucidated.

2.2 Nitrogen (N)

Nitrate (NO₃⁻) is the major inorganic N source absorbed by non-wetland plants and its uptake and metabolism are tightly associated with water utilization. Whereas ammonium has a rather negative effect on water absorption, NO₃⁻ was shown to enhance *L_p* in multiple plant species. Although post-translational regulation may also be involved, the increase in *L_p* induced by NO₃⁻ was shown to correspond to up-regulation of aquaporin expression (Gorska et al. 2008; Gloser et al. 2007). It has also been disputed whether NO₃⁻ acts directly or through its assimilation products. In Arabidopsis, the conserved *L_p* response of a nitrate reductase mutant (*nir*) supports the first hypothesis (Li et al. 2016). Furthermore, characterization of various Arabidopsis genotypes, altered in NO₃⁻ transport, assimilation, or signaling and growing under various N supply conditions revealed a strong positive relationship between *L_p* and NO₃⁻ accumulation, but in shoots rather than in roots (Li et al. 2016). This suggests a control of *L_p* by putative long-distance signals generated in the shoots. Such systemic regulation of aquaporins was also observed in split-root experiments with rice plants subjected to heterogeneous NO₃⁻ treatments (Ishikawa-Sakurai et al. 2014). We also note that *L_p* regulation by transpiration or wounding (Laur and Hacke 2013; Ishikawa-Sakurai et al. 2014; Liu et al. 2014; Vandeleur et al. 2014) may also be mediated by shoot-borne signals acting on root hydraulics.

Although NRT1.1 is considered as the major NO₃⁻ sensor, the high-affinity nitrate transporter NRT2.1 is also thought to have a signaling role. In contrast with *nrt1.1* plants, *nrt2.1* plants showed a pronounced reduction in *L_p*, by up to 30% (Li et al. 2016). Surprisingly, this regulation was independent of NO₃⁻ and hence, of *NRT2.1* expression level. Thus, the signalling mechanisms involved in this newly discovered pathway remain fairly mysterious.

2.3 Cold

Both moderate cold and freezing challenge the plant water status. For instance, low temperature, which reduces L_p , due to a higher viscosity of water and aquaporin down-regulation, causes water deficit and growth arrest in leaves (Lee et al. 2012). In maize, aquaporins seems to play a key role in the recovery of L_p that is specifically observed in chilling tolerant genotypes (Aroca et al. 2005). Similarly, cold acclimation of *Arabidopsis* seedlings is accompanied by upregulation of two specific PIP genes (*AtPIP1;4*; *AtPIP2;5*) (Rahman et al. 2020), which also show mRNA accumulation in leaves under water deficit (Alexandersson et al. 2005). In rice, proper expression and function of Calcium-dependent Protein Kinase (CDPK) *OsCPK17* appears to be key for plant adaptation to low temperature stress (Almadanim et al. 2017). A comparative phosphoproteomic approach further identified *OsPIP2;1* and *OsPIP2;6* as putative targets of *OsCPK17*. A direct activation by *OsCPK17* is further supported by *in vitro* assays showing that the two aquaporins can be phosphorylated by the protein kinase in a calcium-dependent manner. Since *OsCPK17* also targets sugar metabolism, it was proposed that cold stress-activated *OsCPK17* phosphorylates *OsPIP2;1* and *OsPIP2;6* to coordinate membrane water transport with cell osmotic regulation (Almadanim et al. 2017).

2.4 Circadian control.

The circadian clock allows the fine tuning and synchronization of key physiological, metabolic and signaling pathways, in phase with daily variations of environmental parameters. While circadian regulation of tissue hydraulics has been largely documented (Henzler et al. 1999), it is only recently that the corresponding modes of aquaporin regulation have been investigated. In roots of *Arabidopsis* and maize, the control of aquaporin expression seems to be largely transcriptional and depends on well-identified clock components such as *ELF3* (Takase et al. 2011; Caldeira et al. 2014). In maize, the oscillation amplitude of aquaporin expression and plant hydraulic conductance was strongly enhanced under water deficit, while oscillating expression of the core circadian clock genes was not altered (Caldeira et al. 2014). Interestingly, this specific potentiation of the hydraulic outputs, which possibly depends on ABA, appears to be beneficial for cumulated plant growth under

water deficit whereas steady tissue hydraulic properties are more appropriate under water replete conditions. These data show how adaptive responses of plant hydraulics to specific environmental variations rely on exquisite coupling to endogenous signalling mechanisms.

In contrast with roots, the circadian control of leaf hydraulics seems to be mediated through post-translational modifications of proteins (Prado et al. 2019; Choudhary et al. 2015). In particular, the rosette hydraulic conductivity (K_{ros}) of *Arabidopsis* shows circadian oscillations with a peak at midday, in phase with quantitative variations of PIP2 aquaporin phosphorylation (Prado et al. 2019). Remarkably, this modification is necessary but not sufficient for circadian control of K_{ros} , which requires the additional contribution of 14-3-3 proteins (Prado et al. 2013; Prado et al. 2019). Two of these (GRF4, GRF10) can physically interact with AtPIP2;1 and modulate its function in *Xenopus* oocytes depending on the phosphorylation status of the aquaporin C-terminal tail, with more pronounced effects on phosphomimetic than on phosphodeficient forms (Prado et al. 2019). Much remains to be learned about the protein kinase/protein phosphatase machinery that ensures oscillations in aquaporin phosphorylation and, possibly, a synchronized activation of 14-3-3 proteins.

3) A multiplicity of endogenous signals

3.1 Auxin.

Transcriptomic analyses in *Arabidopsis* have revealed a dramatic down-regulation of aquaporin expression during lateral root formation (Péret et al. 2012). This effect is mediated in part by auxin which, more generally, down-regulates most aquaporin expression in roots within 5-10 hours, and, consequently, *Lp*. Auxin also induces, after 24h, a drop in turgor of cortical cells. Interestingly, mutant analyses have shown that all these molecular and physiological responses are mediated through Auxin Response Factor 7 (ARF7), a transcription factor that serves as a master regulator of lateral root formation and emergence. Yet, the down-stream targets of ARF7 in these responses remain to be identified. Our current understanding is that a finely tuned spatial and temporal control of *PIP* expression favours water entry into the lateral root primordium. This inflow is made at this

expense of overlaying cells, thereby reducing their mechanical resistance and facilitating lateral root emergence.

3.2 Methyl jasmonate (MeJa)

Contrasting effects of MeJa on root water transport have been reported by different groups. (Sanchez-Romera et al. 2014) observed that exogenously supplied MeJA increases L_p in bean, tomato and Arabidopsis. These effects are associated with an increased phosphorylation of PIP2s and partly depend on ABA, which accumulates in response to MeJa. In bean, the fluoridone inhibitor counteracted the effects of MeJA on both ABA accumulation and L_p . By contrast, exogenous MeJa induced a decrease in cell hydraulic conductivity in the cortex of Arabidopsis roots (Lee and Zwiazek 2019). Surprisingly, *fad3-2* and *fad7-2* plants, which are defective in MeJa endogenous biosynthesis, were insensitive to this external treatment. These contrasting results suggest that effects of MeJa on root hydraulics are cell-type specific, somewhat indirect and involve important hormonal cross-talks that remain to be uncovered.

3.3 Ethylene

The effects of ethylene on aquaporin function in various plant species and organs have revealed the contribution of multiple signalling pathways. In rose, ethylene acts as a repressor of flower opening. It negatively controls petal cell expansion through down-regulation of *RhPIP2;1* (Ma et al. 2008) and *RhPIP1;1* (Chen et al. 2013), which functionally interacts with the former. Consistent with these negative hormonal effects, silencing of ethylene receptors (*RhETRs*) which, in the absence of the hormone function as repressors of ethylene effects, resulted in an ethylene-independent decrease in *RhPIP1;1* expression (Chen et al. 2013). In Arabidopsis leaves, ethylene exerts positive effects on water transport through enhanced C-terminal phosphorylation of *AtPIP2;1* (Qing et al. 2016). These post-translational effects require the ETR-1 receptor and its interacting partner Constitutive Triple-Response 1 (CTR1). They are independent of the major transcriptional pathway involving Ethylene-Insensitive 1 (EIN1) and Ethylene-Insensitive 3-like (EIL3), which would in turn mediate the

transcriptional down-regulation of PIPs, as mentioned above. Tentative regulatory networks acting on aquaporin phosphorylation, under the control of ethylene and through balanced effects on protein kinases and phosphatases, were inferred in part from a differential phosphoproteomics approach in Col-0 and *ctr1* plants (Yang et al. 2013). Since ethylene enhances auxin accumulation, hormonal interactions will also have to be taken into account to comprehend the multiple pathways leading to ethylene-dependent aquaporin regulation (Yang et al. 2013; Qing et al. 2016)

3.4 Sugars

Sugars act as metabolites but also signalling molecules to regulate various whole plant and cellular processes such as photosynthesis and stomatal movements. With respect to tissue water transport, sucrose was shown to enhance Arabidopsis L_p , specifically in the dark, whereas it inhibited L_p in the light, as did other osmotica (Di Pietro et al. 2013). Sucrose also induced changes in AtPIP2;6 phosphorylation (Niittylä et al. 2007), thereby providing a tentative link between light, sugars and aquaporins. This link was corroborated by the high negative correlation existing between the expression of aquaporins and either exogenous glucose or the expression of Hexokinase 1 (AtHXK1) in over-expressing or knock-out transgenic Arabidopsis (Kelly et al. 2017). Protein kinases that are involved in sucrose-induced phosphorylation responses, among which RLKs (e.g. SIRK1) and receptor-like cytoplasmic kinases (RLCKs e.g. Brassinosteroid Signaling Kinase 8 (BSK8)) have been identified through typical changes in their phosphorylation status within the few minutes following sucrose resupply. Some aquaporins were subsequently identified as downstream targets of these protein kinases (Wu et al. 2019; Wu et al. 2013; Wu et al. 2014). For instance, molecular interactions of SIRK1 with AtPIP2;6 were established through pull down experiments and phosphorylation assays using an AtPIP2;6-specific peptide. The mesophyll protoplast water permeability of Arabidopsis lines showing different expression levels of active SIRK1 revealed that the RLK directly regulates aquaporins via phosphorylation in a sucrose dependent manner (Wu et al. 2013). In addition, SIRK1 forms a complex with its co-receptor QSK1. SIRK1 shows autophosphorylation, subsequently trans-phosphorylates

QSK1, which in turn enhances and stabilizes the interaction of aquaporins with the receptor kinase complex (Wu et al. 2019). Our understanding of BSK8 action on aquaporins is not as precise, but the RLCK possibly acts on another isoform, AtPIP2;7, that shows reduced phosphorylation in a *bsk8* mutant. Although still incomplete, the whole set of data suggests intricate osmotic, metabolic and signalling effects of sugars acting on aquaporins through multiple receptor pathways and time frames (Wu et al. 2014). At the whole plant level, the somewhat opposite effects of sugar on hydraulics depending on organs, timing, or environmental conditions surely reflect contrasting contexts where maintenance of growth or of water content dominate one another.

3.5 Peptidic hormones and other unknown ligands

The endodermis functions as a major barrier within the root, determining fundamental features of water and nutrient absorption from the soil (Doblas et al. 2017). These properties follow from the typical differentiation of endodermal cells, with an initial formation of Casparian strips made out of lignin and further deposition of suberin lamellas that wrap the whole cell. Casparian strips mostly block apoplastic solute movements and favor xylem loading by preventing solute backflow from the stele to the cortex. Suberin lamellas, rather, limit transcellular transport of nutrients and possibly water at the endodermis. These barriers, which are at the cross-roads of mineral nutrient and water uptake, are therefore under tight control by endogenous and environmental factors (Barberon et al. 2016). For instance, the loss of Casparian strip integrity is detected by diffusible vasculature-derived peptides CASPARIAN STRIP INTEGRITY FACTORS 1 & 2 (CIF1 & 2) which act through the SCHENGEN3 receptor-like kinase (Pfister et al. 2014, Doblas et al. 2017, Nakayama et al. 2017) to rebalance water and mineral nutrient uptake. More specifically, CIF1 and CIF2 induce a reduction in L_p , driven by the deactivation of aquaporins (Wang et al. 2019). They also limit ion leakage through deposition of suberin in endodermal cell walls. Inhibition of aquaporins seems to be a primary effect of CIFs, which in turn may promote suberin deposition. Interestingly, CIF1 and CIF2 act independently of ABA signaling, which can also be a potent regulator of aquaporins in endodermal cells.

Casparian strip deposition is guided by the local expression of a family of Casparian Strip domain Proteins (CASPs). Related proteins named CASP-like are expressed throughout the plant. Four of these (CASPL1B1, CASPL1B2, CASPL1D1, and CASPL1D2) were identified as interacting with AtPIP2;1 (Bellati et al. 2016). Although the physiological relevance of these molecular interactions is not known, CASPL1B1 was shown to specifically activate phosphorylated AtPIP2;1 (Champeyroux et al. 2019), pointing to a putative composite regulatory pathway.

4) Biotic interactions

4.1 Plant immunity

In recent years, several landmark works have reported a crucial role of aquaporins in plant defense (Zhang et al. 2019a; Li et al. 2020a; Wang et al. 2018; Tian et al. 2016; Ai et al. 2021). AtPIP1;4 was identified as one of the PIPs, the expression of which is induced by infection with *Pseudomonas syringae* pv. *Tomato* (Tian et al. 2016). The use of intracellular and extracellular ROS dyes in yeast and transgenic Arabidopsis further revealed the capacity of AtPIP1;4 to mediate the diffusion of apoplastic H₂O₂ into the cytoplasm. This process represents a major step in the signaling response of plants to pathogen-associated molecular patterns (PAMPs), which activate RBOH NADPH oxidases present on the plasma membrane. More generally, analysis of knock-out mutants and overexpressing plants revealed that AtPIP1;4 is necessary for PAMP-triggered immunity (Tian et al. 2016) and mediates systemic acquired resistance (SAR) through a parallel pathway involving activation of NONINDUCER OF PR GENES1 (NPR1). These findings establish a major signaling role of PIPs in plant defense (Figure 1C). Interestingly, a mechanism allowing bacterial effectors to counteract this aquaporin signalling function was recently uncovered (Ai et al. 2021). In brief, a Crinkler effector (CRN78) from *Phytophthora* displaying a protein kinase activity was shown to phosphorylate the C-terminal tail of several H₂O₂-transporting PIPs of soybean or *Nicotiana benthamiana*, thereby triggering their degradation through a proteasome-dependent pathway. This led to inhibition of

H₂O₂-accumulation and reduced immunity of these plants when infected by *Phytophthora* oomycetes (Ai et al. 2021).

A signalling role for aquaporins during plant defense was also uncovered for AtPIP2;1 in guard cells, with this aquaporin being necessary for flg22-induced stomatal closure. In this case, a pathway was proposed, whereby the bacterial PAMP flg22 binds to its receptor FLS2 leading to the successive activation of the BAK1 co-receptor and SnRK2.6 protein kinase. This in turn phosphorylates AtPIP2;1 at Ser121 to promote both water and H₂O₂ transport as shown in the context of ABA-induced stomatal closure (Rodrigues et al. 2017) (Figure 1A). Flagellin (flg22) also enhanced lateral diffusion of AtPIP2;1 at the guard cell plasma membrane through effects on microtubules, resulting in its internalization to the cell (Cui et al. 2021). It is not yet clear whether these effects contribute to the primary role of AtPIP2;1 in stomatal closure or reflect a feedback inhibition of its function due to flg22-induced ROS accumulation. Interestingly, AtPIP2;1 dynamics at the plasma membrane of adjacent (subsidiary) cells showed a distinct dependency on the cytoskeleton and were somewhat insensitive to flg22 (Cui et al. 2021). Thus, cell-specific regulation of aquaporins probably contributes to an integrated control of water exchange between cell types during stomatal movements.

PAMPs such as chitin are produced during plant attack by fungi and insects. Aquaporin-mediated water transport in leaf tissues such as the mesophyll and bundle sheath was shown to be down-regulated by chitin (Attia et al. 2020). This inhibition could contribute to plant defense by hampering vascular pathogen propagation in xylem vessels and by promoting stomatal closure through long-distance hydraulic signalling. Although the mode of aquaporin regulation by chitin remains as yet unknown, the contribution of two chitin receptor-like kinases, chitin elicitor receptor kinase 1 (CERK1) and lysine motif receptor kinase 5 (LYK5), could be established (Attia et al. 2020)

Recent work also points to another molecular role of aquaporin during bacterial infection. For instance, rice OsPIP1;3 and Arabidopsis AtPIP1;4 molecularly interact with the bacterial protein harpin (Hpa1) which is secreted by Gram-negative pathogenic bacteria. Thus, PIPs serve as receptors

for harpins at the plant plasma, which in turn mediate the delivery of effector proteins (such as PthXo1) into the plant cell cytoplasm (Li et al. 2019; Li et al. 2015b). Along the same lines, AtPIP2;7 was recently shown to form a molecular complex with the protease domain of the tobacco etch potyvirus NIa protein (Martinez et al. 2016). Due to its numerous additional connections to stress responses, the aquaporin seems to be a preferential target during viral infection.

The overall data show that aquaporins are at the heart of the arms race between plants and their aggressors. On the one hand, aquaporins help plants resist pathogens by contributing to defense signaling pathways and by preventing their ingress through promotion of stomatal closure (Zhang et al. 2019a; Li et al. 2020a). On the other hand, some aquaporins can be hijacked by bacterial or viral pathogenic machineries, thereby promoting pathogen propagation.

4.2 Mycorrhizae

Aquaporins also contribute to positive plant–microbe interactions whereby they play important roles in nutrient transport and abiotic stress responses. In particular, arbuscular mycorrhizal symbiosis can notably improve the plant water status. A fine dialog exists between the fungus and the host plants, leading to precise transcriptional and post-translational regulation of plant aquaporins, particularly under drought (Quiroga et al. 2019). Similarly, the sub-cellular expression of specific tonoplast aquaporins seems to be subtly targeted in N-fixing symbiotic nodules (Gavrin et al. 2014). Although strong hormonal interplays are suspected, the signaling pathways leading to these regulations are as yet unknown.

5) Commonalities of signalling pathways

The survey above provides a broad but scattered knowledge of signalling processes targeting cell and tissue hydraulics in plants (Table 1). Yet, a few main principles do emerge from this information.

Whereas transcriptional control is definitely key for long-term adjustment of plant hydraulics in response to environmental or developmental cues, post-translational mechanisms seem to dominate short term (min and hours) regulations, as these are more tailored for fast and reversible

responses to the multiple factors (including water availability) acting on the plant water status. Here, we discuss a few common players of the latter regulatory mechanisms.

Firstly, ROS play a central signalling role in such regulation of plant water transport (Figure 1). They are produced in response to numerous stimuli including osmotic, salt and cold stress, nutrient deprivation and hormones (salicylic acid, ABA). Although several ROS generating systems may operate at the plasma membrane (Martiniere et al. 2019), RBOH NADPH oxidases seem to occupy a central position in this device (Li et al. 2011; Martiniere et al. 2019). By enhancing constitutive cycling of aquaporins at the plasma membrane and shifting the equilibrium towards endocytosis, ROS promote aquaporin internalization, thereby reducing their density at the cell surface and, as a consequence, cell hydraulic conductivity (Boursiac et al. 2008; Wudick et al. 2015). As suggested by recent work in guard cells (Cui et al. 2021), this mechanism could provide a feedback regulation in contexts where aquaporins contribute to the initial steps of ROS signalling.

The sub-cellular trafficking and regulated degradation of aquaporins also appears to be a main component of plant responses to environmental stimuli. Syntaxins and other traffic proteins, such as adaptor proteins (AP), play well-established roles in these processes (Besserer et al. 2012; Hachez et al. 2014a; Pertl-Obermeyer et al. 2016). While some internalized aquaporins accumulate in pre-vacuolar compartments (Luu et al. 2012), a stress-response pathway under the control of Tryptophan-rich sensory protein/translocator (TPSO) (Hachez et al. 2014b) can relay the above mentioned trafficking to promote aquaporin degradation. The autophagic pathways described in *Medicago truncatula* under water deficit (Li et al. 2020b) may also operate under other stressing conditions. In addition, two recent works have pointed to a control by phosphorylation of ubiquitin-dependent degradation of aquaporins in plants under abiotic or biotic stress (Ai et al. 2021; Chen et al. 2021). This mechanism, which allows control of aquaporin abundance, further expands the multiple, and possibly opposite, roles of phosphorylation in controlling aquaporin function.

In relation to the central role of phosphorylation, aquaporins are known to be targeted by multiple classes of protein kinases. This feature holds promises to further decipher aquaporin

signalling pathways in bottom-up approaches (Figure 2). CDPKs have long been proposed as major players of aquaporin regulation (Johnson and Chrispeels 1992; Johansson et al. 1996). Yet, there are very few cases where their role was formally established (Ji et al. 2017; Almadanim et al. 2017). We also note that CDPKs can be involved in upstream steps and therefore indirectly act on aquaporin regulation (Li et al. 2015a). Due to their key role during stomatal movements (Yip Delormel and Boudsocq 2019), it will be worth exploring the action of CDPKs on guard cell aquaporins in closer detail.

SnRK2s represent another potentially important class of protein kinases acting on plant aquaporins. Besides SnRK2.6 (OST1), which mediates ABA- and flg22-dependent activation of AtPIP2;1 in guard cells (Grondin et al. 2015; Rodrigues et al. 2017), other SnRK2s may delineate additional ABA- or osmotic-stress dependent pathways acting on aquaporins. In support for this, functional expression of plant signalling pathways in yeast using a transcriptional readout indicated that four SnRK2s initially thought to be specific for osmotic responses can also contribute to ABA signalling (Ruschhaupt et al. 2019). Thus, the model of ABA-dependent regulation of AtPIP2;1 may be extended to additional pairs of SnRK2s and PIPs.

RLKs constitute another major a class of serine/threonine kinases that perceive environmental and extracellular developmental signals and transduce them via their intracellular kinase domain. While their role in aquaporin regulation in the endodermis (SGN3) and during sugar response (SIRK1, BSK8) can serve as models, we anticipate a role for RLKs in many other cellular contexts including water stress (see Outstanding questions). For instance, a RLK (LP2) and a Cytoplasmic Receptor-like Kinase (OsRLCK311), both from rice, were shown to interact with PIPs *in vivo* (Wu et al. 2015; Sade et al. 2020), but the functional role of these interactions was not definitively elucidated. Moreover, Arabidopsis RKL1, the activating signals of which remain to be identified, was shown to molecularly interact with AtPIP2;1 (Bellati et al. 2016). Co-expression of AtPIP2;1 with RKL1 in *Xenopus* oocytes resulted in enhanced AtPIP2;1 water transport activity by an as yet unknown mechanism (Bellati et al. 2016). RLKs may also regulate aquaporins by non-canonical

mechanisms. For instance, co-expression of FERONIA and AtPIP2;1 in *Xenopus* oocytes leads to inactivation of the latter (Bellati et al. 2016). This inactivation seems to require the protein kinase activity of FERONIA and is favored by two C-terminal phosphorylation sites of AtPIP2;1 that, in contrast, have been described as activating sites. Thus, these phosphorylated residues may favor the FERONIA-AtPIP2;1 interaction, which in turn would lead to aquaporin inhibition.

Finally, CBL-interacting protein kinases (CIPKs) represent major regulators of plant nutrition and response to abiotic stresses (Dong et al. 2020). It is rather surprising that, despite efforts from several laboratories including ours, no aquaporin has ever been identified as one of their numerous substrates. In addition, protein phosphatases, which surely play a key role in phosphorylation-dependent regulation of aquaporins, will deserve specific attention in future studies.

Lipid signalling represents another emerging component targeting aquaporins (see Outstanding questions). For instance, genetic alteration of phosphoinositides (phosphatidyl inositol [4,5] bisphosphate: PtdInsP2) in tobacco (*Nicotiana tabacum*) increased the osmotic water permeability and aquaporin activity of protoplasts isolated from suspension cells (Ma et al. 2015). The lipid effects may operate through direct interaction with aquaporins or through indirect cellular effects of PtdInsP2. For instance, interaction of PtdInsP2 with TPSO is required for interaction with AtPIP2;7 and its subsequent degradation (Jurkiewicz et al. 2020). A role for lipid signaling in PIP regulation is also suggested by the physical interaction between AtPIP2;1 and two phospholipases D (PLD δ and PLD γ 1) (Bellati et al. 2016). In plants, PLDs and their enzymatic product phosphatidic acid (PA) play key roles in cellular responses to hormonal and abiotic stimuli. Along these lines, AtPIP2;1 and AtPIP2;2 were identified as PA-binding proteins (McLoughlin et al. 2013) but the functional effects of these interactions remain to be elucidated.

In complement to these targeted studies, we believe that systems biology approaches will help uncover previously unidentified signalling pathways targeting aquaporins. For instance, protein interaction networks involving aquaporins have pointed to links with clusters associated with brassinosteroid signaling (Bellati et al. 2016). A genome-wide search for membrane-protein

complexes has revealed other clusters that are enriched in aquaporins and have potential roles in stimulus-dependent regulation of water transport (Jones et al. 2014). Finally, a close inspection of aquaporin gene position in gene regulation networks will for sure reveal other modes of developmental or environmental regulation of aquaporins (Alexandersson et al. 2010).

6) Perspectives

Whereas this overview principally deals with chemical signals, whether abiotic or hormonal, there is still much to be learned about how physical cues lead to aquaporin regulation (see Outstanding questions). Particularly challenging seems the direct link between tissue hydraulics and transpiration that may well be mediated through hydraulic signals, or in other words, pressure-dependent regulation of aquaporins (Pou et al. 2013; Laur and Hacke 2013). Such dependency may also be at the basis of wound-induced alterations in root or leaf hydraulic conductivity (Vandeleur et al. 2014; Liu et al. 2014). Plant defence is another emerging context in which aquaporin regulation will have to be investigated. While the plant water status is crucial for resilience of plants under attacks, it is also a key target for pathogens to facilitate their proliferation within the plant (Xin et al. 2016). We now have some ideas on how aquaporins serve as both signalling components and targets during innate immunity (Zhang et al. 2019a). In contrast, the action of effectors on aquaporins and possible resistance mechanisms associated to these are just emerging (Ai et al. 2021).

Beyond regulation of water transport in a strict sense, we realize that mechanisms targeting other aquaporin transport functions may have some relevance for plant water relations or share similarities with mechanisms targeting the latter. For instance, aquaporin-mediated CO₂ transport is regulated through interaction with β CA4 carbonic anhydrase (Wang et al. 2016), indicating that the various molecular functions of aquaporin interactants and their ability to associate aquaporin regulation to upstream signalling cascades still represent a gold mine of information. Also, the ion transport of AtPIP2;1, which is strongly dependent on divalent cations and pH (Byrt et al. 2017), confirms that both Ca²⁺ and H⁺ are definitely crucial second messengers acting on aquaporin activity.

Because plants are continuously exposed to combinations of environmental constraints, we realize that our analytic representation of single signalling pathways remains largely inappropriate. Although we emphasized how conserved pathways and their cross talks ultimately act on aquaporins, our knowledge of these processes remains very poor. More generally, basic information on how plant water transport is adjusted in response to combined stresses is still lacking. This large field of investigation, combining fundamental plant physiology and systems biology, is now necessary to properly comprehend the amazing capacity of plants to acclimate to a large array of stressful environments.

Advances.

- ABA-dependent and independent signalling pathways act on aquaporins in plants under water deficit.
- Aquaporin-mediated water transport is targeted by multiple abiotic, biotic, or endogenous stimuli; recently identified stimuli include peptide hormones and PAMPs.
- ROS play a central signalling role in regulating water transport in plants by acting on aquaporin subcellular dynamics, localization and/or phosphorylation.
- Plasma membrane aquaporins exhibit a dual water- and H₂O₂-transport function, acting as players in and targets of signalling processes.
- Phosphorylation of aquaporins, which interferes with their gating, trafficking, and stability, is a major target of plant signalling cascades, and multiple classes of protein kinases that act on distinct phosphorylation sites have been identified.

Outstanding questions.

- Is aquaporin-mediated H₂O₂ transport a common process in plant cell signaling?
- Which cellular signals trigger RLK-dependent aquaporin regulation?
- What is the role of lipid signaling in water transport regulation?
- What are the mechanisms for coupling physical stimuli to aquaporin regulation?
- How do aquaporins contribute to plant defense? What are their active roles in plant immunity, and how can they be hijacked by pathogens?
- What are the effects of combined stresses on plant water transport, and which signaling pathways are involved?

549 **Acknowledgments.** The authors thank their colleagues of the Aqua team for fruitful discussions. This
550 work was supported in part by the Agence Nationale de la Recherche (ANR- 18-CE92-0055).

551 **TABLE****Table 1:** Molecular signaling mechanisms involved in aquaporin regulation under abiotic and biotic stresses and by endogenous signals

Type of signal	Stimulus	Signalling intermediate	Molecular mechanism on aquaporins	Aquaporin regulation	Reference
ABIOTIC	drought (ABA-dependent)	SnRK2.6	AtPIP2;1 loop B phosphorylation	up-regulated activity	Grondin et al., 2015; Rodrigues et al., 2017
		LP2	interaction with OsPIP1;1, OsPIP1;3 and OsPIP2;3	down-regulated activity	Wu et al., 2015
	drought (ABA-independent)	RhPTM	interaction with RhPIP2;1	nd	Zhang et al., 2019
		TSPO	nd	AtPIP2;7 protein degradation	Hachez et al., 2014b
		MtCas31/MtATG8	MtCas31 as a cargo receptor for MtPIP2;7	protein degradation by autophagy	Li et al., 2020b
		nd	increased C-ter phosphorylation of AtPIP2;4	nd	Stecker et al., 2014
	osmotic	RBOHs and ROP6	membrane lateral mobility	clustering in nano-domains	Hosy et al., 2015; Martinieri et al., 2019; Smokvarska et al., 2020
		clathrin	nd	protein internalization	Zwiewka et al., 2015
		Rma1H1	AtPIP2;1 ubiquitination	protein degradation	Lee et al., 2009
		UBC32	AtPIP2;1 ubiquitination	protein degradation	Chen et al., 2021
	salinity	clathrin, PI3K, PI4K, SnRK2.4, SnRK2.10	C-ter phosphorylation of PIPs; enhanced mRNA decay	protein internalization and down-regulated activity	Prak et al., 2008; Boursiac et al., 2008; Ueda et al., 2016; Pou et al., 2016; Kawa et al., 2020
	oxygen	cytosolic protons	His protonation of PIPs	down-regulated activity	Tournaire-Roux et al., 2003
	availability	HCR1	nd	down-regulated activity	Shahzad et al., 2016
	Nitrogen	NRT2;1	nd	enhanced gene expression, protein accumulation and up-regulated activity	Li et al., 2016; Ishikawa-Sakurai et al., 2014
	cold	nd	AtPIP1;4 and AtPIP2;5 gene expression	up-regulation of gene expression	Rahman et al., 2020
		OsCPK17	OsPIP2;1 and OsPIP2;6 C-ter phosphorylation	up-regulated activity	Almadanim et al., 2017
ABIOTIC	circadian clock	14-3-3	AtPIP2;1 C-ter phosphorylation	up-regulated activity	Prado et al., 2019
		ELF3	AtPIP1;2 and AtPIP2;1 gene expression	oscillating activity	Caldeira et al., 2014; Takase et al., 2011
	flg22	BAK1 and SnRK2.6	increased AtPIP2;1 loop B phosphorylation	up-regulated activity	Rodrigues et al., 2017

BIOTIC	<i>Phytophthora sojae</i>	CRN78	<i>GmPIP2;13</i> and <i>NbPIP2;2</i> phosphorylation	protein degradation		Ai et al., 2021
	chitin	CERK1 and LYK5	nd	up-regulated activity		Attia et al., 2020
	<i>Xanthomonas oryzae</i>	Harpin	interaction with <i>OsPIP1;3</i> and <i>AtPIP1;4</i>	nd		Li et al., 2015b; Li et al., 2019
ENDOGENOUS	auxin	ARF7	<i>PIP</i> gene expression	down-regulated activity		Péret et al., 2012
	MeJa	nd	PIP2 C-ter phosphorylation	up-regulated activity		Lee and Zwiazek, 2019
	ethylene	<i>RhETR</i> s	<i>RhPIP1;1</i> and <i>RhPIP2;1</i> gene expression	down-regulated gene expression	gene	Ma et al., 2008 ; Chen et al., 2013
		ETR1 and CTR1	<i>AtPIP2;1</i> phosphorylation	C-ter up-regulated activity		Qing et al., 2016; Yang et al., 2013
	sucrose	SIRK1, QSK1	<i>AtPIP2;6</i> phosphorylation	C-ter up-regulated activity		Niittyla et al., 2007; Wu et al., 2014; Wu et al., 2019; Wu et al., 2013
		BSK8	<i>AtPIP2;7</i> phosphorylation	C-ter nd		Wu et al., 2014
	CIF peptide	SGN3	nd	down-regulated activity		Wang et al., 2019
	Other	CASPL1B1	physical interaction with <i>AtPIP2;1</i>	up-regulated activity		Champeyroux et al., 2019

Figure legends

Figure 1. Cellular responses to biotic or abiotic stimuli involving hydrogen peroxide (H_2O_2) and aquaporins. **A**, In guard cells (right), the Snf1-Related protein Kinase 2.6 (SnRK2.6) is a key component of abscisic acid (ABA)- and flagellin (flg22)-induced stomatal closure. Flg22 is a Pathogen-Associated Molecular Pattern (PAMP) recognized by the plasma membrane Flagellin-Sensitive 2 (FLS2) receptor, which acts in concert with BRI1-Associated Kinase 1 (BAK1). Upon activation, SnRK2.6 phosphorylates two NADPH oxidases, Respiratory Burst Oxidase Homolog protein D (RBOHD) and RBOHF, and the AtPIP2;1 aquaporin. The latter facilitates the influx of H_2O_2 and the efflux of water, thereby promoting stomatal closure (Grondin et al. 2015; Rodrigues et al. 2017). Flg22 also enhances the lateral mobility of AtPIP2;1 at the guard cell plasma membrane (left), thereby promoting its internalization (Cui et al. 2021). **B**, The root hydraulic conductivity (L_p) is strongly decreased in response to abiotic stresses such as osmotic and salt stress, or hormones such as salicylic acid (SA). All these stimuli act through accumulation of H_2O_2 which lowers AtPIP2;1 phosphorylation (P), alters its plasma membrane mobility and promotes its internalization (Boursiac et al. 2008; Wudick et al. 2015). In contrast, PIP phosphorylation promotes its trafficking to the plasma membrane. The increased apoplastic H_2O_2 production observed in response to an osmotic stress is mediated through the clustering in nanodomains of the Rho-related GTPase of plants 6 (Rop6) together with the RBOHD and F NADPH oxidases (Smokvarska et al. 2020; Martiniere et al. 2019). **C**, In mesophyll cells, PAMPs are perceived by plasma membrane receptors, which trigger, in combination with BAK1 co-receptor and other non-represented cellular components, the activation of RBOHB NADPH oxidase and, as a consequence, apoplastic accumulation of H_2O_2 . The AtPIP1;4 aquaporin plays a key role in facilitating the diffusion of apoplastic H_2O_2 into the cytoplasm (Tian et al. 2016). The increase in cytoplasmic H_2O_2 concentration ultimately leads to activation of key proteins (Pathogenesis-Related protein 1: PR1; Pathogenesis-Related protein 2: PR2; Plant Defensin 1.2: PDF1.2; Noninducer of PR genes 1 : NPR1; Mitogen-Activated Protein Kinase 3 : MAPK3; Glucan Synthase-like 5: GSL5) involved in Systemic Acquired Resistance (SAR) or Pathogen Triggered Immunity (PTI), as indicated. The figure also shows how the Crinkler 78 (CRN78) effector of *Phytophthora* can penetrate the plant cell where it interacts with phosphorylated NbPIP2;2, to induce its internalization and further degradation, thereby preventing its role in PAMP signaling (Ai et al., 2021). Thick blue arrows and thick red arrows indicate initial effects of stimuli and their output responses, respectively; thin dotted blue line arrows refer to diffusion of the indicated molecules; red dotted line arrows represent protein phosphorylation by indicated protein kinases; red dashed-dotted line arrows and red dashed-dotted line blunt arrows describe indirect activation and

inactivation pathways, respectively; brown line arrows indicate movements of proteins or membrane compartments.

Figure 2. Signaling pathways converging on PIP phosphorylation. The figure represents a prototypal PIP aquaporin with its transmembrane domains (1-6) and extracellular (A, C, E) or cytosolic (B, D) loops. The indicated stimuli lead to enhanced (arrows) or reduced (blunt arrow) phosphorylation of Ser residues in loop B or C-terminal tail. The inset shows three pathways acting on unknown PIP phosphorylation sites (white question marks). Signaling intermediates are indicated with the following color code: Snf1-related protein kinase 2 (SnRK2): blue; Calcium-Dependent Protein Kinase (CDPK), green; Receptor-Like Kinase (RLK), red; others, black. Note that the indicated protein kinases do not necessarily act directly on the indicated sites. Other abbreviations: Brassinosteroid Signaling Kinase 8 (BSK8); Chitin Elicitor Receptor Kinase 1 (CERK1); Calcium-dependent Protein Kinase 17 (CPK17); Constitutive Triple-Response 1 (CTR1); Leaf Panicle 2 (LP2); Lysine motif receptor Kinase 5 (LYK5); phosphatidylinositol 3-kinase (PI3K); phosphatidylinositol 4-kinase (PI4K); Qian Shou Kinase 1 (QSK1); Receptor-like Kinase 1 (RLK1), Sucrose Induced Receptor Kinase 1 (SIRK1); 14-3-3 protein (14-3-3). For more details and references see text and Table 1.

References.

- Ai G, Xia Q, Song T, Li T, Zhu H, Peng H, Liu J, Fu X, Zhang M, Jing M, Xia A, Dou D** (2021) A *Phytophthora sojae* CRN effector mediates phosphorylation and degradation of plant aquaporin proteins to suppress host immune signaling. *PLoS Pathog* **17**:e1009388
- Alexandersson E, Danielson JA, Råde J, Moparthy VK, Fontes M, Kjellbom P, Johanson U** (2010) Transcriptional regulation of aquaporins in accessions of *Arabidopsis* in response to drought stress. *Plant J* **61**:650-660
- Alexandersson E, Frayse L, Sjøvall-Larsen S, Gustavsson S, Fellert M, Karlsson M, Johanson U, Kjellbom P** (2005) Whole gene family expression and drought stress regulation of aquaporins. *Plant Mol Biol* **59**:469-484
- Almadanim MC, Alexandre BM, Rosa MTG, Sapeta H, Leitao AE, Ramalho JC, Lam TT, Negrao S, Abreu IA, Oliveira MM** (2017) Rice calcium-dependent protein kinase OsCPK17 targets plasma membrane intrinsic protein and sucrose-phosphate synthase and is required for a proper cold stress response. *Plant Cell Environ* **40**:1197-1213
- Aroca R, Amodeo G, Fernandez-Illescas S, Herman EM, Chaumont F, Chrispeels MJ** (2005) The role of aquaporins and membrane damage in chilling and hydrogen peroxide induced changes in the hydraulic conductance of maize roots. *Plant Physiol* **137**:341-353
- Aroca R, Porcel R, Ruiz-Lozano JM** (2012) Regulation of root water uptake under abiotic stress conditions. *J Exp Bot* **63**:43-57
- Attia Z, Dalal A, Moshelion M** (2020) Vascular bundle sheath and mesophyll cells modulate leaf water balance in response to chitin. *Plant J* **101**:1368-1377
- Barberon M, Vermeer JE, De Bellis D, Wang P, Naseer S, Andersen TG, Humbel BM, Nawrath C, Takano J, Salt DE, Geldner N** (2016) Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell* **164**:447-459
- Bellati J, Champeyroux C, Hem S, Rofidal V, Krouk G, Maurel C, Santoni V** (2016) Novel aquaporin regulatory mechanisms revealed by interactomics. *Mol Cell Proteomics* **15**:3473-3487
- Besserer A, Burnotte E, Bienert GP, Chevalier AS, Errachid A, Grefen C, Blatt MR, Chaumont F** (2012) Selective regulation of maize plasma membrane aquaporin trafficking and activity by the SNARE SYP121. *Plant Cell* **24**:3463-3481
- Boursiac Y, Boudet J, Postaire O, Luu DT, Tournaire-Roux C, Maurel C** (2008) Stimulus-induced downregulation of root water transport involves reactive oxygen species-activated cell signalling and plasma membrane intrinsic protein internalization. *Plant J* **56**:207-218

- Bramley H, Turner NC, Turner DW, Tyerman SD** (2010) The contrasting influence of short-term hypoxia on the hydraulic properties of cells and roots of wheat and lupin. *Function Plant Biol* **37**:183-193
- Byrt CS, Zhao M, Kourghi M, Bose J, Henderson SW, Qiu J, Gilliam M, Schultz C, Schwarz M, Ramesh SA, Yool A, Tyerman S** (2017) Non-selective cation channel activity of aquaporin AtPIP2;1 regulated by Ca^{2+} and pH. *Plant Cell Environ* **40**: 802-815
- Caldeira CF, Jeanguenin L, Chaumont F, Tardieu F** (2014) Circadian rhythms of hydraulic conductance and growth are enhanced by drought and improve plant performance. *Nat Commun* **5**:5365
- Ceciliato PHO, Zhang J, Liu Q, Shen X, Hu H, Liu C, Schaffner AR, Schroeder JI** (2019) Intact leaf gas exchange provides a robust method for measuring the kinetics of stomatal conductance responses to abscisic acid and other small molecules in *Arabidopsis* and grasses. *Plant Methods* **15**:38
- Champeyroux C, Bellati J, Barberon M, Rofidal V, Maurel C, Santoni V** (2019) Regulation of a plant aquaporin by a Casparian strip membrane domain protein-like. *Plant Cell Environ* **42**:1788-1801
- Chaumont F, Barrieu F, Herman EM, Chrispeels MJ** (1998) Characterization of a maize tonoplast aquaporin expressed in zones of cell division and elongation. *Plant Physiol* **117**:1143-1152
- Chaumont F, Tyerman SD** (2014) Aquaporins: highly regulated channels controlling plant water relations. *Plant Physiol* **164**:1600-1618
- Chen Q, Liu R, Wu Y, Wei S, Wang Q, Zheng Y, Xia R, Shang X, Yu F, Yang X, Liu L, Huang X, Wang Y, Xie Q** (2021) ERAD-related E2 and E3 enzymes modulate the drought response by regulating the stability of PIP2 aquaporins. *Plant Cell*. doi:10.1093/plcell/koab141
- Chen W, Yin X, Wang L, Tian J, Yang R, Liu D, Yu Z, Ma N, Gao J** (2013) Involvement of rose aquaporin RhPIP1;1 in ethylene-regulated petal expansion through interaction with RhPIP2;1. *Plant Mol Biol* **83**:219-233
- Choudhary MK, Nomura Y, Wang L, Nakagami H, Somers DE** (2015) Quantitative circadian phosphoproteomic analysis of *Arabidopsis* reveals extensive clock control of key components in physiological, metabolic, and signaling pathways. *Mol Cell Proteomics* **14**:2243-2260
- Cui Y, Zhao Y, Lu Y, Su X, Chen Y, Shen Y, Lin J, Li X** (2021) In vivo single-particle tracking of the aquaporin AtPIP2;1 in stomata reveals cell type-specific dynamics. *Plant Physiol* **185**:1666-1681
- Davies WJ, Bennett MJ** (2015) Achieving more crop per drop. *Nature Plants* **1**:15118
- Di Pietro M, Vialaret J, Li G, Hem S, Rossignol M, Maurel C, Santoni V** (2013) Coordinated post-translational responses of aquaporins to abiotic and nutritional stimuli in *Arabidopsis* roots. *Mol Cell Proteomics* **12**:3886-3897

- 671 **Doblas VG, Geldner N, Barberon M** (2017) The endodermis, a tightly controlled barrier for nutrients.
 672 *Curr Opin Plant Biol* **39**:136-143
- 673 **Dong Q, Bai B, Almutairi BO, Kudla J** (2020) Emerging roles of the CBL-CIPK calcium signaling
 674 network as key regulatory hub in plant nutrition. *J Plant Physiol* **257**:153335
- 675 **Dynowski M, Schaaf G, Loque D, Moran O, Ludewig U** (2008) Plant plasma membrane water
 676 channels conduct the signalling molecule H₂O₂. *Biochem J* **414**:53-61
- 677 **Gavrin A, Kaiser BN, Geiger D, Tyerman SD, Wen Z, Bisseling T, Fedorova EE** (2014) Adjustment of
 678 host cells for accommodation of symbiotic bacteria: vacuole defunctionalization, HOPS
 679 suppression, and TIP1g retargeting in *Medicago*. *Plant Cell* **26**:3809-3822
- 680 **Gloser V, Zwieniecki MA, Orians CM, Holbrook NM** (2007) Dynamic changes in root hydraulic
 681 properties in response to nitrate availability. *J Exp Bot* **58**:2409-2415
- 682 **Gorska A, Ye Q, Holbrook NM, Zwieniecki MA** (2008) Nitrate control of root hydraulic properties in
 683 plants: translating local information to whole plant response. *Plant Physiol* **148**:1159-1167
- 684 **Grondin A, Rodrigues O, Verdoucq L, Merlot S, Leonhardt N, Maurel C** (2015) Aquaporins contribute
 685 to ABA-triggered stomatal closure through OST1-mediated phosphorylation. *Plant Cell*
 686 **27**:1945-1954
- 687 **Hachez C, Laloux T, Reinhardt H, Cavez D, Degand H, Grefen C, De Rycke R, Inze D, Blatt MR,**
 688 **Russinova E, Chaumont F** (2014a) Arabidopsis SNAREs SYP61 and SYP121 coordinate the
 689 trafficking of plasma membrane aquaporin PIP2;7 to modulate the cell membrane water
 690 permeability. *Plant Cell* **26**:3132-3147
- 691 **Hachez C, Veljanovski V, Reinhardt H, Guillaumot D, Vanhee C, Chaumont F, Batoko H** (2014b) The
 692 *Arabidopsis* abiotic stress-induced TSPO-related protein reduces cell-surface expression of
 693 the aquaporin PIP2;7 through protein-protein interactions and autophagic degradation. *Plant*
 694 *Cell* **26**:4974-4990
- 695 **Henzler T, Waterhouse RN, Smyth AJ, Carvajal M, Cooke DT, Schaffner AR, Steudle E, Clarkson DT**
 696 (1999) Diurnal variations in hydraulic conductivity and root pressure can be correlated with
 697 the expression of putative aquaporins in the roots of *Lotus japonicus*. *Planta* **210**:50-60
- 698 **Hosy E, Martiniere A, Choquet D, Maurel C, Luu DT** (2015) Super-resolved and dynamic imaging of
 699 membrane proteins in plant cells reveal contrasting kinetic profiles and multiple confinement
 700 mechanisms. *Mol Plant* **8**:339-342
- 701 **Ishikawa-Sakurai J, Hayashi H, Murai-Hatano M** (2014) Nitrogen availability affects hydraulic
 702 conductivity of rice roots, possibly through changes in aquaporin gene expression. *Plant Soil*
 703 **379**:289-300

- 704 **Ji R, Zhou L, Liu J, Wang Y, Yang L, Zheng Q, Zhang C, Zhang B, Ge H, Yang Y, Zhao F, Luan S, Lan W**
 705 (2017) Calcium-dependent protein kinase CPK31 interacts with arsenic transporter AtNIP1;1
 706 and regulates arsenite uptake in *Arabidopsis thaliana*. *PLoS One* **12**:e0173681
- 707 **Johansson I, Larsson C, Ek B, Kjellbom P** (1996) The major integral proteins of spinach leaf plasma
 708 membranes are putative aquaporins and are phosphorylated in response to Ca^{2+} and
 709 apoplastic water potential. *Plant Cell* **8**:1181-1191
- 710 **Johnson KD, Chrispeels MJ** (1992) Tonoplast-bound protein kinase phosphorylates tonoplast intrinsic
 711 protein. *Plant Physiol* **100**:1787-1795
- 712 **Jones AM, Xuan Y, Xu M, Wang RS, Ho CH, Lalonde S, You CH, Sardi MI, Parsa SA, Smith-Valle E, Su**
 713 **T, Frazer KA, Pilot G, Pratelli R, Grossmann G, Acharya BR, Hu HC, Engineer C, Villiers F, Ju**
 714 **C, Takeda K, Su Z, Dong Q, Assmann SM, Chen J, Kwak JM, Schroeder JI, Albert R, Rhee SY,**
 715 **Frommer WB** (2014) Border control--a membrane-linked interactome of *Arabidopsis*. *Science*
 716 **344**:711-716
- 717 **Jurkiewicz P, Senicourt L, Ayeb H, Lequin O, Lacapere JJ, Batoko H** (2020) A Plant-Specific N-terminal
 718 Extension Reveals Evolutionary Functional Divergence within Translocator Proteins. *iScience*
 719 **23**:100889
- 720 **Kawa D, Meyer AJ, Dekker HL, Abd-El-Hallem AM, Gevaert K, Van De Slijke E, Maszkowska J,**
 721 **Bucholc M, Dobrowolska G, De Jaeger G, Schuurink RC, Haring MA, Testerink C** (2020)
 722 SnRK2 Protein Kinases and mRNA Decapping Machinery Control Root Development and
 723 Response to Salt. *Plant Physiol* **182**:361-377.
- 724 **Kelly G, Sade N, Doron-Faigenboim A, Lerner S, Shatil-Cohen A, Yeselson Y, Egbaria A, Kottapalli J,**
 725 **Schaffer AA, Moshelion M, Granot D** (2017) Sugar and hexokinase suppress expression of
 726 PIP aquaporins and reduce leaf hydraulics that preserves leaf water potential. *Plant J* **91**:325-
 727 339.
- 728 **Laur J, Hacke UG** (2013) Transpirational demand affects aquaporin expression in poplar roots. *J Exp*
 729 *Bot* **64**:2283-2293
- 730 **Lee HK, Cho SK, Son O, Xu Z, Hwang I, Kim WT** (2009) Drought stress-induced Rma1H1, a RING
 731 membrane-anchor E3 ubiquitin ligase homolog, regulates aquaporin levels via ubiquitination
 732 in transgenic *Arabidopsis* plants. *Plant Cell* **21**:622-641
- 733 **Lee SH, Chung GC, Jang JY, Ahn SJ, Zwiazek JJ** (2012) Overexpression of PIP2;5 aquaporin alleviates
 734 effects of low root temperature on cell hydraulic conductivity and growth in *Arabidopsis*.
 735 *Plant Physiol* **159**:479-488
- 736 **Lee SH, Zwiazek JJ** (2019) Regulation of water transport in *Arabidopsis* by methyl jasmonate. *Plant*
 737 *Physiol Biochem* **139**:540-547

- Li G, Boudsocq M, Hem S, Vialaret J, Rossignol M, Maurel C, Santoni V** (2015a) The calcium-dependent protein kinase CPK7 acts on root hydraulic conductivity. *Plant Cell Environ* **38**:1312-1320
- Li G, Chen T, Zhang Z, Li B, Tian S** (2020a) Roles of Aquaporins in Plant-Pathogen Interaction. *Plants* **9**:1134
- Li G, Tillard P, Gojon A, Maurel C** (2016) Dual regulation of root hydraulic conductivity and plasma membrane aquaporins by plant nitrate accumulation and high-affinity nitrate transporter NRT2.1. *Plant Cell Physiol* **57**:733-742
- Li L, Wang H, Gago J, Cui H, Qian Z, Kodama N, Ji H, Tian S, Shen D, Chen Y, Sun F, Xia Z, Ye Q, Sun W, Flexas J, Dong H** (2015b) Harpin Hpa1 Interacts with Aquaporin PIP1;4 to Promote the Substrate Transport and Photosynthesis in Arabidopsis. *Sci Rep* **5**:17207
- Li P, Zhang L, Mo X, Ji H, Bian H, Hu Y, Majid T, Long J, Pang H, Tao Y, Ma J, Dong H** (2019) Rice aquaporin PIP1;3 and harpin Hpa1 of bacterial blight pathogen cooperate in a type III effector translocation. *J Exp Bot* **70**:3057-3073
- Li X, Liu Q, Feng H, Deng J, Zhang R, Wen J, Dong J, Wang T** (2020b) Dehydrin MtCAS31 promotes autophagic degradation under drought stress. *Autophagy* **16**:862-877
- Li X, Wang X, Yang Y, Li R, He Q, Fang X, Luu DT, Maurel C, Lin J** (2011) Single-molecule analysis of PIP2;1 dynamics and partitioning reveals multiple modes of *Arabidopsis* plasma membrane aquaporin regulation. *Plant Cell* **23**:3780-3797
- Liu J, Equiza MA, Navarro-Rodenas A, Lee SH, Zwiazek JJ** (2014) Hydraulic adjustments in aspen (*Populus tremuloides*) seedlings following defoliation involve root and leaf aquaporins. *Planta* **240**:553-564
- Luu DT, Martinière A, Sorieul M, Runions J, Maurel C** (2012) Fluorescence recovery after photobleaching reveals high cycling dynamics of plasma membrane aquaporins in *Arabidopsis* roots under salt stress. *Plant J* **69**:894-905
- Ma N, Xue J, Li Y, Liu X, Dai F, Jia W, Luo Y, Gao J** (2008) *Rh-PIP2;1*, a rose aquaporin gene, is involved in ethylene-regulated petal expansion. *Plant Physiol* **148**:894-907
- Ma X, Shatil-Cohen A, Ben-Dor S, Wigoda N, Perera IY, Im YJ, Diminshtein S, Yu L, Boss WF, Moshelion M, Moran N** (2015) Do phosphoinositides regulate membrane water permeability of tobacco protoplasts by enhancing the aquaporin pathway? *Planta* **241**:741-755
- Maathuis FJ, Filatov V, Herzyk P, Krijger GC, Axelsen KB, Chen S, Green BJ, Li Y, Madagan KL, Sanchez-Fernandez R, Forde BG, Palmgren MG, Rea PA, Williams LE, Sanders D, Amtmann A** (2003) Transcriptome analysis of root transporters reveals participation of multiple gene families in the response to cation stress. *Plant J* **35**:675-692

- 772 **Martinez F, Rodrigo G, Aragonés V, Ruiz M, Lodewijk I, Fernandez U, Elena SF, Daros JA** (2016)
 773 Interaction network of tobacco etch potyvirus NIa protein with the host proteome during
 774 infection. *BMC Genomics* **17**:87
- 775 **Martiniere A, Fiche JB, Smokvarska M, Mari S, Alcon C, Dumont X, Hematy K, Jaillais Y, Nollmann**
 776 **M, Maurel C** (2019) Osmotic stress activates two reactive oxygen species pathways with
 777 distinct effects on protein nanodomains and diffusion. *Plant Physiol* **179**:1581-1593
- 778 **Maurel C, Boursiac Y, Luu DT, Santoni V, Shahzad Z, Verdoucq L** (2015) Aquaporins in plants. *Physiol*
 779 *Rev* **95**:1321-1358
- 780 **Maurel C, Nacry P** (2020) Root architecture and hydraulics converge for acclimation to changing
 781 water availability. *Nat Plants* **6**:744-749
- 782 **Maurel C, Verdoucq L, Rodrigues O** (2016) Aquaporins and plant transpiration. *Plant Cell Environ*
 783 **39**:2580-2587
- 784 **McLoughlin F, Arisz SA, Dekker HL, Kramer G, de Koster CG, Haring MA, Munnik T, Testerink C**
 785 (2013) Identification of novel candidate phosphatidic acid-binding proteins involved in the
 786 salt-stress response of *Arabidopsis thaliana* roots. *Biochem J* **450**:573-581
- 787 **Niittylä T, Fuglsang AT, Palmgren MG, Frommer WB, Schulze WX** (2007) Temporal analysis of
 788 sucrose-induced phosphorylation changes in plasma membrane proteins of *Arabidopsis*. *Mol*
 789 *Cell Proteomics* **6**:1711-1726
- 790 **Péret B, Li GW, Zhao J, Band LR, Voß U, Postaire O, Luu D-T, Da Ines O, Casimiro I, Lucas M, Wells**
 791 **DM, Lazzerini L, Nacry P, King JR, Jensen OE, Schäffner AR, Maurel C, Bennett MJ** (2012)
 792 Auxin regulates aquaporin function to facilitate lateral root emergence. *Nature Cell Biol*
 793 **14**:991-998
- 794 **Pertl-Obermeyer H, Wu XN, Schrodtt J, Mudsam C, Obermeyer G, Schulze WX** (2016) Identification
 795 of Cargo for Adaptor Protein (AP) Complexes 3 and 4 by Sucrose Gradient Profiling. *Mol Cell*
 796 *Proteomics* **15**:2877-2889
- 797 **Pou A, Jeanguenin L, Milhiet T, Batoko H, Chaumont F, Hachez C** (2016) Salinity-mediated
 798 transcriptional and post-translational regulation of the *Arabidopsis* aquaporin PIP2;7. *Plant*
 799 *Mol Biol* **92**:731-744
- 800 **Pou A, Medrano H, Flexas J, Tyerman SD** (2013) A putative role for TIP and PIP aquaporins in
 801 dynamics of leaf hydraulic and stomatal conductances in grapevine under water stress and
 802 re-watering. *Plant Cell Environ* **36**:828-8439
- 803 **Prado K, Boursiac Y, Tournaire-Roux C, Monneuse J-M, Postaire O, Da Ines O, Schäffner AR, Hem S,**
 804 **Santoni V, Maurel C** (2013) Regulation of *Arabidopsis* leaf hydraulics involves light-
 805 dependent phosphorylation of aquaporins in veins. *Plant Cell* **25**:1029-1039

- Prado K, Cotellet V, Li G, Bellati J, Tang N, Tournaire-Roux C, Martiniere A, Santoni V, Maurel C** (2019) Oscillating Aquaporin Phosphorylation and 14-3-3 Proteins Mediate the Circadian Regulation of Leaf Hydraulics. *Plant Cell* **31**:417-429
- Prak S, Hem S, Boudet J, Viennois G, Sommerer N, Rossignol M, Maurel C, Santoni V** (2008) Multiple phosphorylations in the C-terminal tail of plant plasma membrane aquaporins. Role in sub-cellular trafficking of AtPIP2;1 in response to salt stress. *Mol Cell Proteomics* **7**:1019-1030
- Qing D, Yang Z, Li M, Wong WS, Guo G, Liu S, Guo H, Li N** (2016) Quantitative and functional phosphoproteomic analysis reveals that ethylene regulates water transport via the C-terminal phosphorylation of aquaporin PIP2;1 in *Arabidopsis*. *Mol Plant* **9**:158-174
- Qiu J, McGaughey SA, Groszmann M, Tyerman SD, Byrt CS** (2020) Phosphorylation influences water and ion channel function of AtPIP2;1. *Plant Cell Environ* **43**:2428-2442
- Quiroga G, Erice G, Ding L, Chaumont F, Aroca R, Ruiz-Lozano JM** (2019) The arbuscular mycorrhizal symbiosis regulates aquaporins activity and improves root cell water permeability in maize plants subjected to water stress. *Plant Cell Environ* **42**:2274-2290
- Rae L, Lao NT, Kavanagh TA** (2011) Regulation of multiple aquaporin genes in *Arabidopsis* by a pair of recently duplicated DREB transcription factors. *Planta* **234**:429-444
- Rahman A, Kawamura Y, Maeshima M, Rahman A, Uemura M** (2020) Plasma Membrane Aquaporin Members PIPs Act in Concert to Regulate Cold Acclimation and Freezing Tolerance Responses in *Arabidopsis thaliana*. *Plant Cell Physiol* **61**:787-802
- Ricardi MM, Gonzalez RM, Zhong S, Dominguez PG, Duffy T, Turjanski PG, Salgado Salter JD, Alleva K, Carrari F, Giovannoni JJ, Estevez JM, Iusem ND** (2014) Genome-wide data (ChIP-seq) enabled identification of cell wall-related and aquaporin genes as targets of tomato ASR1, a drought stress-responsive transcription factor. *BMC Plant Biol* **14**:29
- Rodrigues O, Reshetnyak G, Grondin A, Saijo Y, Leonhardt N, Maurel C, Verdoucq L** (2017) Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proc Natl Acad Sci USA* **114**:9200-9205
- Rosales MA, Maurel C, Nacry P** (2019) Absciscic acid coordinates dose-dependent developmental and hydraulic responses of roots to water deficit. *Plant Physiol* **180**:2198-2211
- Ruschhaupt M, Mergner J, Mucha S, Papacek M, Doch I, Tischer SV, Hemmler D, Chiasson D, Edel KH, Kudla J, Schmitt-Kopplin P, Kuster B, Grill E** (2019) Rebuilding core absciscic acid signaling pathways of *Arabidopsis* in yeast. *Embo J* **38**:e101859.
- Sade N, Weng F, Tajima H, Zeron Y, Zhang L, Rubio Wilhelmi MDM, Day G, Peleg Z, Blumwald E** (2020) A Cytoplasmic Receptor-like Kinase Contributes to Salinity Tolerance. *Plants (Basel)* **9**:1383

- 840 **Sanchez-Romera B, Ruiz-Lozano JM, Li G, Luu DT, Martinez-Ballesta Mdel C, Carvajal M, Zamarrero**
841 **AM, Garcia-Mina JM, Maurel C, Aroca R** (2014) Enhancement of root hydraulic conductivity
842 by methyl jasmonate and the role of calcium and abscisic acid in this process. *Plant Cell*
843 *Environ* **37**:995-1008
- 844 **Shahzad Z, Canut M, Tournaire-Roux C, Martinière A, Boursiac Y, Loudet O, Maurel C** (2016) A
845 potassium-dependent oxygen sensing pathway regulates plant root hydraulics. *Cell* **167**:87-
846 98
- 847 **Shatil-Cohen A, Attia Z, Moshelion M** (2011) Bundle-sheath cell regulation of xylem-mesophyll water
848 transport via aquaporins under drought stress: a target of xylem-borne ABA? *Plant J* **67**:72-80
- 849 **Smokvarska M, Francis C, Platre MP, Fiche JB, Alcon C, Dumont X, Nacry P, Bayle V, Nollmann M,**
850 **Maurel C, Jaillais Y, Martiniere A** (2020) A Plasma Membrane Nanodomain Ensures Signal
851 Specificity during Osmotic Signaling in Plants. *Curr Biol* **30**:4654-4664
- 852 **Soma F, Mogami J, Yoshida T, Abekura M, Takahashi F, Kidokoro S, Mizoi J, Shinozaki K, Yamaguchi-**
853 **Shinozaki K** (2017) ABA-unresponsive SnRK2 protein kinases regulate mRNA decay under
854 osmotic stress in plants. *Nature Plants* **3**:16204
- 855 **Stecker KE, Minkoff BB, Sussman MR** (2014) Phosphoproteomic Analyses Reveal Early Signaling
856 Events in the Osmotic Stress Response. *Plant Physiol* **165**:1171-1187
- 857 **Steudle E** (1992) The biophysics of plant water: compartmentation, coupling with metabolic
858 processes, and water flow in plant roots. In: Somero GN, Osmond CB, Bolis CL (eds) *Water*
859 *and life: Comparative analysis of water relationships at the organismic, cellular,*
860 *and molecular levels.* Springer-Verlag, Berlin, pp 273-304
- 861 **Takase T, Ishikawa H, Murakami H, Kikuchi J, Sato-Nara K, Suzuki H** (2011) The circadian clock
862 modulates water dynamics and aquaporin expression in *Arabidopsis* roots. *Plant Cell Physiol*
863 **52**:373-383
- 864 **Tan X, Xu H, Khan S, Equiza MA, Lee SH, Vaziriyeganeh M, Zwiazek JJ** (2018) Plant water transport
865 and aquaporins in oxygen-deprived environments. *J Plant Physiol* **227**:20-30
- 866 **Tardieu F, Simonneau T, Parent B** (2015) Modelling the coordination of the controls of stomatal
867 aperture, transpiration, leaf growth, and abscisic acid: update and extension of the Tardieu-
868 Davies model. *J Exp Bot* **66**:2227-2237
- 869 **Tian S, Wang X, Li P, Wang H, Ji H, Xie J, Qiu Q, Shen D, Dong H** (2016) Plant Aquaporin AtPIP1;4
870 Links Apoplastic H₂O₂ Induction to Disease Immunity Pathways. *Plant Physiol* **171**:1635-1650
- 871 **Tournaire-Roux C, Sutka M, Javot H, Gout E, Gerbeau P, Luu DT, Bligny R, Maurel C** (2003) Cytosolic
872 pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature*
873 **425**:393-397

- 874 **Tran STH, Imran S, Horie T, Qiu J, McGaughey S, Byrt CS, Tyerman SD, Katsuhara M** (2020) A Survey
875 of Barley PIP Aquaporin Ionic Conductance Reveals Ca^{2+} -Sensitive HvPIP2;8 Na^+ and K^+
876 Conductance. *Int J Mol Sci* **21**:7135
- 877 **Tyerman SD, Bohnert HJ, Maurel C, Steudle E, Smith JA** (1999) Plant aquaporins: their molecular
878 biology, biophysics and significance for plant water relations. *J Exp Bot* **50**:1055-1071
- 879 **Tyerman SD, McGaughey SA, Qiu J, Yool AJ, Byrt CS** (2021) Adaptable and Multifunctional Ion-
880 Conducting Aquaporins. *Annu Rev Plant Biol.* doi:10.1146/annurev-arplant-081720-013608
- 881 **Ueda M, Tsutsumi N, Fujimoto M** (2016) Salt stress induces internalization of plasma membrane
882 aquaporin into the vacuole in *Arabidopsis thaliana*. *Biochem Biophys Res Commun* **474**:742-
883 746
- 884 **Vandeleur RK, Sullivan W, Athman A, Jordans C, Gilliam M, Kaiser BN, Tyerman SD** (2014) Rapid
885 shoot-to-root signalling regulates root hydraulic conductance via aquaporins. *Plant Cell*
886 *Environ* **37**:520-538
- 887 **Wang C, Hu H, Qin X, Zeise B, Xu D, Rappel WJ, Boron WF, Schroeder JI** (2016) Reconstitution of CO_2
888 regulation of SLAC1 anion channel and function of CO_2 -permeable PIP2;1 aquaporin as
889 CARBONIC ANHYDRASE4 interactor. *Plant Cell* **28**:568-582
- 890 **Wang P, Calvo-Polanco M, Reyt G, Barberon M, Champeyroux C, Santoni V, Maurel C, Franke RB,**
891 **Ljung K, Novak O, Geldner N, Boursiac Y, Salt DE** (2019) Surveillance of cell wall diffusion
892 barrier integrity modulates water and solute transport in plants. *Sci Rep* **9**:4227
- 893 **Wang R, Wang M, Chen K, Wang S, Mur LAJ, Guo S** (2018) Exploring the Roles of Aquaporins in
894 Plant-Microbe Interactions. *Cells* **7**:267
- 895 **Wu F, Sheng P, Tan J, Chen X, Lu G, Ma W, Heng Y, Lin Q, Zhu S, Wang J, Wang J, Guo X, Zhang X, Lei**
896 **C, Wan J** (2015) Plasma membrane receptor-like kinase leaf panicle 2 acts downstream of the
897 DROUGHT AND SALT TOLERANCE transcription factor to regulate drought sensitivity in rice. *J*
898 *Exp Bot* **66**:271-281
- 899 **Wu X, Sklodowski K, Encke B, Schulze WX** (2014) A kinase-phosphatase signaling module with BSK8
900 and BSL2 involved in regulation of sucrose-phosphate synthase. *J Proteome Res* **13**:3397-
901 3409
- 902 **Wu XN, Chu L, Xi L, Pertl-Obermeyer H, Li Z, Sklodowski K, Sanchez-Rodriguez C, Obermeyer G,**
903 **Schulze WX** (2019) Sucrose-induced Receptor Kinase 1 is Modulated by an Interacting Kinase
904 with Short Extracellular Domain. *Mol Cell Proteomics* **18**:1556-1571
- 905 **Wu XN, Sanchez Rodriguez C, Pertl-Obermeyer H, Obermeyer G, Schulze WX** (2013) Sucrose-
906 induced receptor kinase SIRK1 regulates a plasma membrane aquaporin in *Arabidopsis*. *Mol*
907 *Cell Proteomics* **12**:2856-2873

- 908 **Wudick MM, Li X, Valentini V, Geldner N, Chory J, Lin J, Maurel C, Luu DT** (2015) Subcellular
 909 redistribution of root aquaporins induced by hydrogen peroxide. *Mol Plant* **8**:1103-1114
- 910 **Xin XF, Nomura K, Aung K, Velasquez AC, Yao J, Boutrot F, Chang JH, Zipfel C, He SY** (2016) Bacteria
 911 establish an aqueous living space in plants crucial for virulence. *Nature* **539**:524-529
- 912 **Yang Z, Guo G, Zhang M, Liu CY, Hu Q, Lam H, Cheng H, Xue Y, Li J, Li N** (2013) Stable isotope
 913 metabolic labeling-based quantitative phosphoproteomic analysis of *Arabidopsis* mutants
 914 reveals ethylene-regulated time-dependent phosphoproteins and putative substrates of
 915 constitutive triple response 1 kinase. *Mol Cell Proteomics* **12**:3559-3582
- 916 **Yip Delormel T, Boudsocq M** (2019) Properties and functions of calcium-dependent protein kinases
 917 and their relatives in *Arabidopsis thaliana*. *New Phytol* **224**:585-604
- 918 **Zhang L, Chen L, Dong H** (2019a) Plant Aquaporins in Infection by and Immunity Against Pathogens -
 919 A Critical Review. *Front Plant Sci* **10**:632
- 920 **Zhang S, Feng M, Chen W, Zhou X, Lu J, Wang Y, Li Y, Jiang CZ, Gan SS, Ma N, Gao J** (2019b) In rose,
 921 transcription factor PTM balances growth and drought survival via PIP2;1 aquaporin. *Nat*
 922 *Plants* **5**:290-299
- 923 **Zhu D, Wu Z, Cao G, Li J, Wei J, Tsuge T, Gu H, Aoyama T, Qu LJ** (2014) TRANSLUCENT GREEN, an ERF
 924 family transcription factor, controls water balance in *Arabidopsis* by activating the expression
 925 of aquaporin genes. *Mol Plant* **7**:601-615
- 926 **Zwiewka M, Nodzynski T, Robert S, Vanneste S, Friml J** (2015) Osmotic Stress Modulates the Balance
 927 between Exocytosis and Clathrin-Mediated Endocytosis in *Arabidopsis thaliana*. *Mol Plant*
 928 **8**:1175-1187
- 929




Figure 1. Cellular responses to biotic or abiotic stimuli involving hydrogen peroxide (H₂O₂) and aquaporins. **A**, In guard cells (right), the Snf1-Related protein Kinase 2.6 (SnRK2.6) is a key component of abscisic acid (ABA)- and flagellin (flg22)-induced stomatal closure. Flg22 is a Pathogen-Associated Molecular Pattern (PAMP) recognized by the plasma membrane Flagellin-Sensitive 2 (FLS2) receptor, which acts in concert with BRI1-Associated Kinase 1 (BAK1). Upon activation, SnRK2.6 phosphorylates two NADPH oxidases, Respiratory Burst Oxidase Homolog protein D (RBOHD) and RBOHF, and the AtPIP2;1 aquaporin. The latter facilitates the influx of H₂O₂ and the efflux of water, thereby promoting stomatal closure (Grondin et al. 2015; Rodrigues et al. 2017). Flg22 also enhances the lateral mobility of AtPIP2;1 at the guard cell plasma membrane (left), thereby promoting its internalization (Cui et al. 2021). **B**, The root hydraulic conductivity (L_p) is strongly decreased in response to abiotic stresses such as osmotic and salt stress, or hormones such as salicylic acid (SA). All these stimuli act through accumulation of H₂O₂ which lowers AtPIP2;1 phosphorylation (P), alters its plasma membrane mobility and promotes its internalization (Boursiac et al. 2008; Wudick et al. 2015). In contrast, PIP phosphorylation promotes its trafficking to the plasma membrane. The increased apoplastic H₂O₂ production observed in response to an osmotic stress is mediated through the clustering in nanodomains of the Rho-related GTPase of plants 6 (Rop6) together with the RBOHD and F NADPH oxidases (Smokvarska et al. 2020; Martiniere et al. 2019). **C**, In mesophyll cells, PAMPs are perceived by plasma membrane receptors, which trigger, in combination with BAK1 co-receptor and other non-represented cellular components, the activation of RBOHB NADPH oxidase and, as a consequence, apoplastic accumulation of H₂O₂. The AtPIP1;4 aquaporin plays a key role in facilitating the diffusion of apoplastic H₂O₂ into the cytoplasm (Tian et al. 2016). The increase in cytoplasmic H₂O₂ concentration ultimately leads to activation of key proteins (Pathogenesis-Related protein 1: PR1; Pathogenesis-Related protein 2: PR2; Plant Defensin 1.2: PDF1.2; Noninducer of PR genes 1: NPR1; Mitogen-Activated Protein Kinase 3: MAPK3; Glucan Synthase-like 5: GSL5) involved in Systemic Acquired Resistance (SAR) and Pathogen Triggered Immunity (PTI), as indicated. The figure also shows how the Crinkler 78 (CRN78) effector of *Phytophthora* can penetrate the plant cell where it interacts with phosphorylated NbPIP2;2, to induce its internalization and further degradation, thereby preventing its role in PAMP signaling (Ai et al., 2021). Thick blue arrows and thick red arrows indicate initial effects of stimuli and their output responses, respectively; thin dotted blue line arrows refer to diffusion of the indicated molecules; red dotted line arrows represent protein phosphorylation by indicated protein kinases; red dashed-dotted line arrows and red dashed-dotted line blunt arrows describe indirect activation and inactivation pathways, respectively; brown line arrows indicate movements of proteins or membrane compartments.




Figure 2. Signaling pathways converging on PIP phosphorylation. The figure represents a prototypal PIP aquaporin with its transmembrane domains (1-6) and extracellular (A, C, E) or cytosolic (B, D) loops. The indicated stimuli lead to enhanced (arrows) or reduced (blunt arrow) phosphorylation of Ser residues in loop B or C-terminal tail. The inset shows three pathways acting on unknown PIP phosphorylation sites (white question marks). Signaling intermediates are indicated with the following color code: Snf1-related protein kinase 2 (SnRK2): blue; Calcium-Dependent Protein Kinase (CDPK), green; Receptor-Like Kinase (RLK), red; others, black. Note that the indicated protein kinases do not necessarily act directly on the indicated sites. Other abbreviations: Brassinosteroid Signaling Kinase 8 (BSK8); Chitin Elicitor Receptor Kinase 1 (CERK1); Calcium-dependent Protein Kinase 17 (CPK17); Constitutive Triple-Response 1 (CTR1); Leaf Panicle 2 (LP2); Lysine motif receptor Kinase 5 (LYK5); phosphatidylinositol 3-kinase (PI3K); phosphatidylinositol 4-kinase (PI4K); Qian Shou Kinase 1 (QSK1); Receptor-like Kinase 1 (RKL1), Sucrose Induced Receptor Kinase 1 (SIRK1); 14-3-3 protein (14-3-3). For more details and references see text and Table 1.

ADVANCES

- ABA-dependent and independent signalling pathways act on aquaporins in plants under water deficit.
- Aquaporin-mediated water transport is targeted by multiple abiotic, biotic, or endogenous stimuli; recently identified stimuli include peptide hormones and PAMPs.
- ROS play a central signalling role in regulating water transport in plants by acting on aquaporin subcellular dynamics, localization and/or phosphorylation.
- Plasma membrane aquaporins exhibit a dual water- and H₂O₂-transport function, acting as players in and targets of signalling processes.
- Phosphorylation of aquaporins, which interferes with their gating, trafficking, and stability, is a major target of plant signalling cascades, and multiple classes of protein kinases that act on distinct phosphorylation sites have been identified.

OUTSTANDING QUESTIONS

- Is aquaporin-mediated H_2O_2 transport a common process in plant cell signaling?
- Which cellular signals trigger RLK-dependent aquaporin regulation?
- What is the role of lipid signaling in water transport regulation?
- What are the mechanisms for coupling physical stimuli to aquaporin regulation?
- How do aquaporins contribute to plant defense? What are their active roles in plant immunity, and how can they be hijacked by pathogens?
- What are the effects of combined stresses on plant water transport, and which signaling pathways are involved?

Parsed Citations

- Ai G, Xia Q, Song T, Li T, Zhu H, Peng H, Liu J, Fu X, Zhang M, Jing M, Xia A, Dou D (2021) A *Phytophthora sojae* CRN effector mediates phosphorylation and degradation of plant aquaporin proteins to suppress host immune signaling. *PLoS Pathog* 17:e1009388
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Alexandersson E, Danielson JA, Råde J, Moparthi VK, Fontes M, Kjellbom P, Johanson U (2010) Transcriptional regulation of aquaporins in accessions of *Arabidopsis* in response to drought stress. *Plant J* 61:650-660
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Alexandersson E, Fraysse L, Sjoval-Larsen S, Gustavsson S, Fellert M, Karlsson M, Johanson U, Kjellbom P (2005) Whole gene family expression and drought stress regulation of aquaporins. *Plant Mol Biol* 59:469-484
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Almadanim MC, Alexandre BM, Rosa MTG, Sapeta H, Leitao AE, Ramalho JC, Lam TT, Negrao S, Abreu IA, Oliveira MM (2017) Rice calcium-dependent protein kinase OsCPK17 targets plasma membrane intrinsic protein and sucrose-phosphate synthase and is required for a proper cold stress response. *Plant Cell Environ* 40:1197-1213
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Aroca R, Amedeo G, Fernandez-Illescas S, Herman EM, Chaumont F, Chrispeels MJ (2005) The role of aquaporins and membrane damage in chilling and hydrogen peroxide induced changes in the hydraulic conductance of maize roots. *Plant Physiol* 137:341-353
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Aroca R, Porcel R, Ruiz-Lozano JM (2012) Regulation of root water uptake under abiotic stress conditions. *J Exp Bot* 63:43-57
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Attia Z, Dalal A, Moshelion M (2020) Vascular bundle sheath and mesophyll cells modulate leaf water balance in response to chitin. *Plant J* 101:1368-1377
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Barberon M, Vermeer JE, De Bellis D, Wang P, Naseer S, Andersen TG, Humbel BM, Nawrath C, Takano J, Salt DE, Geldner N (2016) Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell* 164:447-459
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bellati J, Champeyroux C, Hem S, Rofidal V, Krouk G, Maurel C, Santoni V (2016) Novel aquaporin regulatory mechanisms revealed by interactomics. *Mol Cell Proteomics* 15:3473-3487
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Besserer A, Burnotte E, Bienert GP, Chevalier AS, Errachid A, Grefen C, Blatt MR, Chaumont F (2012) Selective regulation of maize plasma membrane aquaporin trafficking and activity by the SNARE SYP121. *Plant Cell* 24:3463-3481
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Boursiac Y, Boudet J, Postaire O, Luu DT, Tournaire-Roux C, Maurel C (2008) Stimulus-induced downregulation of root water transport involves reactive oxygen species-activated cell signalling and plasma membrane intrinsic protein internalization. *Plant J* 56:207-218
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bramley H, Turner NC, Turner DW, Tyerman SD (2010) The contrasting influence of short-term hypoxia on the hydraulic properties of cells and roots of wheat and lupin. *Function Plant Biol* 37:183-193
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Byrt CS, Zhao M, Kourghi M, Bose J, Henderson SW, Qiu J, Gilliam M, Schultz C, Schwarz M, Ramesh SA, Yool A, Tyerman S (2017) Non-selective cation channel activity of aquaporin AtPIP2;1 regulated by Ca²⁺ and pH. *Plant Cell Environ* 40: 802-815
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Caldeira CF, Jeanguenin L, Chaumont F, Tardieu F (2014) Circadian rhythms of hydraulic conductance and growth are enhanced by drought and improve plant performance. *Nat Commun* 5:5365
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ceciliato PHO, Zhang J, Liu Q, Shen X, Hu H, Liu C, Schaffner AR, Schroeder JI (2019) Intact leaf gas exchange provides a robust method for measuring the kinetics of stomatal conductance responses to abscisic acid and other small molecules in *Arabidopsis* and grasses. *Plant Methods* 15:38
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Champeyroux C, Bellati J, Barberon M, Rofidal V, Maurel C, Santoni V (2019) Regulation of a plant aquaporin by a Casparian strip membrane domain protein-like. *Plant Cell Environ* 42:1788-1801
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chaumont F, Barrieu F, Herman EM, Chrispeels MJ (1998) Characterization of a maize tonoplast aquaporin expressed in zones of cell division and elongation. *Plant Physiol* 117:1143-1152
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chaumont F, Tyerman SD (2014) Aquaporins: highly regulated channels controlling plant water relations. *Plant Physiol* 164:1600-1618

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen Q, Liu R, Wu Y, Wei S, Wang Q, Zheng Y, Xia R, Shang X, Yu F, Yang X, Liu L, Huang X, Wang Y, Xie Q (2021) ERAD-related E2 and E3 enzymes modulate the drought response by regulating the stability of PIP2 aquaporins. Plant Cell. doi:10.1093/plcell/koab141

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen W, Yin X, Wang L, Tian J, Yang R, Liu D, Yu Z, Ma N, Gao J (2013) Involvement of rose aquaporin RhPIP1;1 in ethylene-regulated petal expansion through interaction with RhPIP2;1. Plant Mol Biol 83:219-233

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Choudhary MK, Nomura Y, Wang L, Nakagami H, Somers DE (2015) Quantitative circadian phosphoproteomic analysis of Arabidopsis reveals extensive clock control of key components in physiological, metabolic, and signaling pathways. Mol Cell Proteomics 14:2243-2260

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cui Y, Zhao Y, Lu Y, Su X, Chen Y, Shen Y, Lin J, Li X (2021) In vivo single-particle tracking of the aquaporin AtPIP2;1 in stomata reveals cell type-specific dynamics. Plant Physiol 185:1666-1681

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Davies WJ, Bennett MJ (2015) Achieving more crop per drop. Nature Plants 1:15118

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Di Pietro M, Vialaret J, Li G, Hem S, Rossignol M, Maurel C, Santoni V (2013) Coordinated post-translational responses of aquaporins to abiotic and nutritional stimuli in Arabidopsis roots. Mol Cell Proteomics 12:3886-3897

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Doblas VG, Geldner N, Barberon M (2017) The endodermis, a tightly controlled barrier for nutrients. Curr Opin Plant Biol 39:136-143

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dong Q, Bai B, Almutairi BO, Kudla J (2020) Emerging roles of the CBL-CIPK calcium signaling network as key regulatory hub in plant nutrition. J Plant Physiol 257:153335

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dynowski M, Schaaf G, Loque D, Moran O, Ludewig U (2008) Plant plasma membrane water channels conduct the signalling molecule H₂O₂. Biochem J 414:53-61

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gavrin A, Kaiser BN, Geiger D, Tyerman SD, Wen Z, Bisseling T, Fedorova EE (2014) Adjustment of host cells for accommodation of symbiotic bacteria: vacuole defunctionalization, HOPS suppression, and TIP1g retargeting in Medicago. Plant Cell 26:3809-3822

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gloser V, Zwieniecki MA, Orians CM, Holbrook NM (2007) Dynamic changes in root hydraulic properties in response to nitrate availability. J Exp Bot 58:2409-2415

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gorska A, Ye Q, Holbrook NM, Zwieniecki MA (2008) Nitrate control of root hydraulic properties in plants: translating local information to whole plant response. Plant Physiol 148:1159-1167

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grondin A, Rodrigues O, Verdoucq L, Merlot S, Leonhardt N, Maurel C (2015) Aquaporins contribute to ABA-triggered stomatal closure through OST1-mediated phosphorylation. Plant Cell 27:1945-1954

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hachez C, Laloux T, Reinhardt H, Cavez D, Degand H, Grefen C, De Rycke R, Inze D, Blatt MR, Russinova E, Chaumont F (2014a) Arabidopsis SNAREs SYP61 and SYP121 coordinate the trafficking of plasma membrane aquaporin PIP2;7 to modulate the cell membrane water permeability. Plant Cell 26:3132-3147

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hachez C, Veljanovski V, Reinhardt H, Guillaumot D, Vanhee C, Chaumont F, Batoko H (2014b) The Arabidopsis abiotic stress-induced TSPO-related protein reduces cell-surface expression of the aquaporin PIP2;7 through protein-protein interactions and autophagic degradation. Plant Cell 26:4974-4990

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Henzler T, Waterhouse RN, Smyth AJ, Carvajal M, Cooke DT, Schaffner AR, Steudle E, Clarkson DT (1999) Diurnal variations in hydraulic conductivity and root pressure can be correlated with the expression of putative aquaporins in the roots of Lotus japonicus. Planta 210:50-60

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hosy E, Martinieri A, Choquet D, Maurel C, Luu DT (2015) Super-resolved and dynamic imaging of membrane proteins in plant cells reveal contrasting kinetic profiles and multiple confinement mechanisms. Mol Plant 8:339-342

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ishikawa-Sakurai J, Hayashi H, Murai-Hatano M (2014) Nitrogen availability affects hydraulic conductivity of rice roots, possibly through changes in aquaporin gene expression. Plant Soil 379:289-300

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ji R, Zhou L, Liu J, Wang Y, Yang L, Zheng Q, Zhang C, Zhang B, Ge H, Yang Y, Zhao F, Luan S, Lan W (2017) Calcium-dependent protein kinase CPK31 interacts with arsenic transporter AtNIP1;1 and regulates arsenite uptake in *Arabidopsis thaliana*. *PLoS One* 12:e0173681

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Johansson I, Larsson C, Ek B, Kjellbom P (1996) The major integral proteins of spinach leaf plasma membranes are putative aquaporins and are phosphorylated in response to Ca²⁺ and apoplastic water potential. *Plant Cell* 8:1181-1191

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Johnson KD, Chrispeels MJ (1992) Tonoplast-bound protein kinase phosphorylates tonoplast intrinsic protein. *Plant Physiol* 100:1787-1795

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jones AM, Xuan Y, Xu M, Wang RS, Ho CH, Lalonde S, You CH, Sardi MI, Parsa SA, Smith-Valle E, Su T, Frazer KA, Pilot G, Pratelli R, Grossmann G, Acharya BR, Hu HC, Engineer C, Villiers F, Ju C, Takeda K, Su Z, Dong Q, Assmann SM, Chen J, Kwak JM, Schroeder JI, Albert R, Rhee SY, Frommer WB (2014) Border control—a membrane-linked interactome of *Arabidopsis*. *Science* 344:711-716

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jurkiewicz P, Senicourt L, Aye H, Lequin O, Lacapere JJ, Batoko H (2020) A Plant-Specific N-terminal Extension Reveals Evolutionary Functional Divergence within Translocator Proteins. *iScience* 23:100889

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kawa D, Meyer AJ, Dekker HL, Abd-El-Halim AM, Gevaert K, Van De Slijke E, Maszkowska J, Bucholtz M, Dobrowolska G, De Jaeger G, Schuurink RC, Haring MA, Testerink C (2020) SnRK2 Protein Kinases and mRNA Decapping Machinery Control Root Development and Response to Salt. *Plant Physiol* 182:361-377.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kelly G, Sade N, Doron-Faigenboim A, Lerner S, Shatil-Cohen A, Yeselson Y, Egbaria A, Kottapalli J, Schaffer AA, Moshelion M, Granot D (2017) Sugar and hexokinase suppress expression of PIP aquaporins and reduce leaf hydraulics that preserves leaf water potential. *Plant J* 91:325-339.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Laur J, Hacke UG (2013) Transpirational demand affects aquaporin expression in poplar roots. *J Exp Bot* 64:2283-2293

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee HK, Cho SK, Son O, Xu Z, Hwang I, Kim WT (2009) Drought stress-induced Rma1H1, a RING membrane-anchor E3 ubiquitin ligase homolog, regulates aquaporin levels via ubiquitination in transgenic *Arabidopsis* plants. *Plant Cell* 21:622-641

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee SH, Chung GC, Jang JY, Ahn SJ, Zwiazek JJ (2012) Overexpression of PIP2;5 aquaporin alleviates effects of low root temperature on cell hydraulic conductivity and growth in *Arabidopsis*. *Plant Physiol* 159:479-488

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee SH, Zwiazek JJ (2019) Regulation of water transport in *Arabidopsis* by methyl jasmonate. *Plant Physiol Biochem* 139:540-547

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li G, Boudsocq M, Hem S, Vialaret J, Rossignol M, Maurel C, Santoni V (2015a) The calcium-dependent protein kinase CPK7 acts on root hydraulic conductivity. *Plant Cell Environ* 38:1312-1320

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li G, Chen T, Zhang Z, Li B, Tian S (2020a) Roles of Aquaporins in Plant-Pathogen Interaction. *Plants* 9:1134

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li G, Tillard P, Gojon A, Maurel C (2016) Dual regulation of root hydraulic conductivity and plasma membrane aquaporins by plant nitrate accumulation and high-affinity nitrate transporter NRT2.1. *Plant Cell Physiol* 57:733-742

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li L, Wang H, Gago J, Cui H, Qian Z, Kodama N, Ji H, Tian S, Shen D, Chen Y, Sun F, Xia Z, Ye Q, Sun W, Flexas J, Dong H (2015b) Harpin Hpa1 Interacts with Aquaporin PIP1;4 to Promote the Substrate Transport and Photosynthesis in *Arabidopsis*. *Sci Rep* 5:17207

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li P, Zhang L, Mo X, Ji H, Bian H, Hu Y, Majid T, Long J, Pang H, Tao Y, Ma J, Dong H (2019) Rice aquaporin PIP1;3 and harpin Hpa1 of bacterial blight pathogen cooperate in a type III effector translocation. *J Exp Bot* 70:3057-3073

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li X, Liu Q, Feng H, Deng J, Zhang R, Wen J, Dong J, Wang T (2020b) Dehydrin MtCAS31 promotes autophagic degradation under drought stress. *Autophagy* 16:862-877

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li X, Wang X, Yang Y, Li R, He Q, Fang X, Luu DT, Maurel C, Lin J (2011) Single-molecule analysis of PIP2;1 dynamics and partitioning reveals multiple modes of *Arabidopsis* plasma membrane aquaporin regulation. *Plant Cell* 23:3780-3797

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu J, Equiza MA, Navarro-Rodenas A, Lee SH, Zwiazek JJ (2014) Hydraulic adjustments in aspen (*Populus tremuloides*) seedlings following defoliation involve root and leaf aquaporins. *Planta* 240:553–564

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Luu DT, Martinière A, Sorieul M, Runions J, Maurel C (2012) Fluorescence recovery after photobleaching reveals high cycling dynamics of plasma membrane aquaporins in *Arabidopsis* roots under salt stress. *Plant J* 69:894–905

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ma N, Xue J, Li Y, Liu X, Dai F, Jia W, Luo Y, Gao J (2008) Rh-PIP2;1, a rose aquaporin gene, is involved in ethylene-regulated petal expansion. *Plant Physiol* 148:894–907

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ma X, Shatil-Cohen A, Ben-Dor S, Wigoda N, Perera IY, Im YJ, Diminshtein S, Yu L, Boss WF, Moshelion M, Moran N (2015) Do phosphoinositides regulate membrane water permeability of tobacco protoplasts by enhancing the aquaporin pathway? *Planta* 241:741–755

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maathuis FJ, Filatov V, Herzyk P, Krijger GC, Axelsen KB, Chen S, Green BJ, Li Y, Madagan KL, Sanchez-Fernandez R, Forde BG, Palmgren MG, Rea PA, Williams LE, Sanders D, Amtmann A (2003) Transcriptome analysis of root transporters reveals participation of multiple gene families in the response to cation stress. *Plant J* 35:675–692

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Martinez F, Rodrigo G, Aragonés V, Ruiz M, Lodewijk I, Fernandez U, Elena SF, Daros JA (2016) Interaction network of tobacco etch potyvirus NIa protein with the host proteome during infection. *BMC Genomics* 17:87

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Martinière A, Fiche JB, Smokvarska M, Mari S, Alcon C, Dumont X, Hematy K, Jaillais Y, Nollmann M, Maurel C (2019) Osmotic stress activates two reactive oxygen species pathways with distinct effects on protein nanodomains and diffusion. *Plant Physiol* 179:1581–1593

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maurel C, Boursiac Y, Luu DT, Santoni V, Shahzad Z, Verdoucq L (2015) Aquaporins in plants. *Physiol Rev* 95:1321–1358

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maurel C, Nacry P (2020) Root architecture and hydraulics converge for acclimation to changing water availability. *Nat Plants* 6:744–749

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maurel C, Verdoucq L, Rodrigues O (2016) Aquaporins and plant transpiration. *Plant Cell Environ* 39:2580–2587

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McLoughlin F, Arisz SA, Dekker HL, Kramer G, de Koster CG, Haring MA, Munnik T, Testerink C (2013) Identification of novel candidate phosphatidic acid-binding proteins involved in the salt-stress response of *Arabidopsis thaliana* roots. *Biochem J* 450:573–581

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niittylä T, Fuglsang AT, Palmgren MG, Frommer WB, Schulze WX (2007) Temporal analysis of sucrose-induced phosphorylation changes in plasma membrane proteins of *Arabidopsis*. *Mol Cell Proteomics* 6:1711–1726

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Péret B, Li GW, Zhao J, Band LR, Voß U, Postaire O, Luu D-T, Da Ines O, Casimiro I, Lucas M, Wells DM, Lazzerini L, Nacry P, King JR, Jensen OE, Schaffner AR, Maurel C, Bennett MJ (2012) Auxin regulates aquaporin function to facilitate lateral root emergence. *Nature Cell Biol* 14:991–998

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pertl-Obermeyer H, Wu XN, Schrodtt J, Mudsam C, Obermeyer G, Schulze WX (2016) Identification of Cargo for Adaptor Protein (AP) Complexes 3 and 4 by Sucrose Gradient Profiling. *Mol Cell Proteomics* 15:2877–2889

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pou A, Jeanguenin L, Milhiet T, Batoko H, Chaumont F, Hachez C (2016) Salinity-mediated transcriptional and post-translational regulation of the *Arabidopsis* aquaporin PIP2;7. *Plant Mol Biol* 92:731–744

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pou A, Medrano H, Flexas J, Tyerman SD (2013) A putative role for TIP and PIP aquaporins in dynamics of leaf hydraulic and stomatal conductances in grapevine under water stress and re-watering. *Plant Cell Environ* 36:828–8439

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prado K, Boursiac Y, Tournaire-Roux C, Monneuse J-M, Postaire O, Da Ines O, Schaffner AR, Hem S, Santoni V, Maurel C (2013) Regulation of *Arabidopsis* leaf hydraulics involves light-dependent phosphorylation of aquaporins in veins. *Plant Cell* 25:1029–1039

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prado K, Cotellet V, Li G, Bellati J, Tang N, Tournaire-Roux C, Martinière A, Santoni V, Maurel C (2019) Oscillating Aquaporin Phosphorylation and 14-3-3 Proteins Mediate the Circadian Regulation of Leaf Hydraulics. *Plant Cell* 31:417–429

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prak S, Hem S, Boudet J, Viennois G, Sommerer N, Rossignol M, Maurel C, Santoni V (2008) Multiple phosphorylations in the C-terminal tail of plant plasma membrane aquaporins. Role in sub-cellular trafficking of AtPIP2;1 in response to salt stress. Mol Cell Proteomics 7:1019-1030

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Qing D, Yang Z, Li M, Wong WS, Guo G, Liu S, Guo H, Li N (2016) Quantitative and functional phosphoproteomic analysis reveals that ethylene regulates water transport via the C-terminal phosphorylation of aquaporin PIP2;1 in Arabidopsis. Mol Plant 9:158-174

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Qiu J, McGaughey SA, Groszmann M, Tyerman SD, Byrt CS (2020) Phosphorylation influences water and ion channel function of AtPIP2;1. Plant Cell Environ 43:2428-2442

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Quiroga G, Erice G, Ding L, Chaumont F, Aroca R, Ruiz-Lozano JM (2019) The arbuscular mycorrhizal symbiosis regulates aquaporins activity and improves root cell water permeability in maize plants subjected to water stress. Plant Cell Environ 42:2274-2290

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rae L, Lao NT, Kavanagh TA (2011) Regulation of multiple aquaporin genes in Arabidopsis by a pair of recently duplicated DREB transcription factors. Planta 234:429-444

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rahman A, Kawamura Y, Maeshima M, Rahman A, Uemura M (2020) Plasma Membrane Aquaporin Members PIPs Act in Concert to Regulate Cold Acclimation and Freezing Tolerance Responses in Arabidopsis thaliana. Plant Cell Physiol 61:787-802

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ricardi MM, Gonzalez RM, Zhong S, Dominguez PG, Duffy T, Turjanski PG, Salgado Salter JD, Aleva K, Carrari F, Giovannoni JJ, Estevez JM, Iusem ND (2014) Genome-wide data (ChIP-seq) enabled identification of cell wall-related and aquaporin genes as targets of tomato ASR1, a drought stress-responsive transcription factor. BMC Plant Biol 14:29

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodrigues O, Reshetnyak G, Grondin A, Saijo Y, Leonhardt N, Maurel C, Verdoucq L (2017) Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. Proc Natl Acad Sci USA 114:9200-9205

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rosales MA, Maurel C, Nacry P (2019) Absciscic acid coordinates dose-dependent developmental and hydraulic responses of roots to water deficit. Plant Physiol 180:2198-2211

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ruschhaupt M, Mergner J, Mucha S, Papacek M, Doch I, Tischer SV, Hemmler D, Chiasson D, Edel KH, Kudla J, Schmitt-Kopplin P, Kuster B, Grill E (2019) Rebuilding core abscisic acid signaling pathways of Arabidopsis in yeast. Embo J 38:e101859.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sade N, Weng F, Tajima H, Zeron Y, Zhang L, Rubio Wilhelmi MDM, Day G, Peleg Z, Blumwald E (2020) A Cytoplasmic Receptor-like Kinase Contributes to Salinity Tolerance. Plants (Basel) 9:1383

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sanchez-Romera B, Ruiz-Lozano JM, Li G, Luu DT, Martinez-Ballesta Mdel C, Carvajal M, Zamarreno AM, Garcia-Mina JM, Maurel C, Aroca R (2014) Enhancement of root hydraulic conductivity by methyl jasmonate and the role of calcium and abscisic acid in this process. Plant Cell Environ 37:995-1008

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shahzad Z, Canut M, Tournaire-Roux C, Martinière A, Boursiac Y, Loudet O, Maurel C (2016) A potassium-dependent oxygen sensing pathway regulates plant root hydraulics. Cell 167:87-98

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shatil-Cohen A, Attia Z, Moshelion M (2011) Bundle-sheath cell regulation of xylem-mesophyll water transport via aquaporins under drought stress: a target of xylem-borne ABA? Plant J 67:72-80

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smokvarska M, Francis C, Platre MP, Fiche JB, Alcon C, Dumont X, Nacry P, Bayle V, Nollmann M, Maurel C, Jaillais Y, Martinière A (2020) A Plasma Membrane Nanodomain Ensures Signal Specificity during Osmotic Signaling in Plants. Curr Biol 30:4654-4664

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Soma F, Mogami J, Yoshida T, Abekura M, Takahashi F, Kidokoro S, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K (2017) ABA-unresponsive SnRK2 protein kinases regulate mRNA decay under osmotic stress in plants. Nature Plants 3:16204

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stecker KE, Minkoff BB, Sussman MR (2014) Phosphoproteomic Analyses Reveal Early Signaling Events in the Osmotic Stress Response. Plant Physiol 165:1171-1187

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Steudle E (1992) The biophysics of plant water: compartmentation, coupling with metabolic processes, and water flow in plant roots. In:

Somero GN, Osmond CB, Bolis CL (eds) *Water and life: Comparative analysis of water relationships at the organismic, cellular, and molecular levels*. Springer-Verlag, Berlin, pp 273-304

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takase T, Ishikawa H, Murakami H, Kikuchi J, Sato-Nara K, Suzuki H (2011) The circadian clock modulates water dynamics and aquaporin expression in *Arabidopsis* roots. *Plant Cell Physiol* 52:373-383

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tan X, Xu H, Khan S, Equiza MA, Lee SH, Vaziriyeganeh M, Zwiazek JJ (2018) Plant water transport and aquaporins in oxygen-deprived environments. *J Plant Physiol* 227:20-30

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tardieu F, Simonneau T, Parent B (2015) Modelling the coordination of the controls of stomatal aperture, transpiration, leaf growth, and abscisic acid: update and extension of the Tardieu-Davies model. *J Exp Bot* 66:2227-2237

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tian S, Wang X, Li P, Wang H, Ji H, Xie J, Qiu Q, Shen D, Dong H (2016) Plant Aquaporin AtPIP1;4 Links Apoplastic H₂O₂ Induction to Disease Immunity Pathways. *Plant Physiol* 171:1635-1650

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tournaire-Roux C, Sutka M, Javot H, Gout E, Gerbeau P, Luu DT, Bligny R, Maurel C (2003) Cytosolic pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature* 425:393-397

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tran STH, Imran S, Horie T, Qiu J, McGaughey S, Byrt CS, Tyerman SD, Katsuhara M (2020) A Survey of Barley PIP Aquaporin Ionic Conductance Reveals Ca²⁺-Sensitive HvPIP2;8 Na⁺ and K⁺ Conductance. *Int J Mol Sci* 21:7135

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tyerman SD, Bohnert HJ, Maurel C, Steudle E, Smith JA (1999) Plant aquaporins: their molecular biology, biophysics and significance for plant water relations. *J Exp Bot* 50:1055-1071

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tyerman SD, McGaughey SA, Qiu J, Yool AJ, Byrt CS (2021) Adaptable and Multifunctional Ion-Conducting Aquaporins. *Annu Rev Plant Biol*. doi:10.1146/annurev-arplant-081720-013608

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ueda M, Tsutsumi N, Fujimoto M (2016) Salt stress induces internalization of plasma membrane aquaporin into the vacuole in *Arabidopsis thaliana*. *Biochem Biophys Res Commun* 474:742-746

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vandeleur RK, Sullivan W, Athman A, Jordans C, Gilliam M, Kaiser BN, Tyerman SD (2014) Rapid shoot-to-root signalling regulates root hydraulic conductance via aquaporins. *Plant Cell Environ* 37:520-538

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang C, Hu H, Qin X, Zeise B, Xu D, Rappel WJ, Boron WF, Schroeder JI (2016) Reconstitution of CO₂ regulation of SLAC1 anion channel and function of CO₂-permeable PIP2;1 aquaporin as CARBONIC ANHYDRASE4 interactor. *Plant Cell* 28:568-582

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang P, Calvo-Polanco M, Rey G, Barberon M, Champeyroux C, Santoni V, Maurel C, Franke RB, Ljung K, Novak O, Geldner N, Boursiac Y, Salt DE (2019) Surveillance of cell wall diffusion barrier integrity modulates water and solute transport in plants. *Sci Rep* 9:4227

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang R, Wang M, Chen K, Wang S, Mur LAJ, Guo S (2018) Exploring the Roles of Aquaporins in Plant-Microbe Interactions. *Cells* 7:267

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wu F, Sheng P, Tan J, Chen X, Lu G, Ma W, Heng Y, Lin Q, Zhu S, Wang J, Wang J, Guo X, Zhang X, Lei C, Wan J (2015) Plasma membrane receptor-like kinase leaf panicle 2 acts downstream of the DROUGHT AND SALT TOLERANCE transcription factor to regulate drought sensitivity in rice. *J Exp Bot* 66:271-281

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wu X, Sklodowski K, Encke B, Schulze WX (2014) A kinase-phosphatase signaling module with BSK8 and BSL2 involved in regulation of sucrose-phosphate synthase. *J Proteome Res* 13:3397-3409

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wu XN, Chu L, Xi L, Pertl-Obermeyer H, Li Z, Sklodowski K, Sanchez-Rodriguez C, Obermeyer G, Schulze WX (2019) Sucrose-induced Receptor Kinase 1 is Modulated by an Interacting Kinase with Short Extracellular Domain. *Mol Cell Proteomics* 18:1556-1571

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wu XN, Sanchez Rodriguez C, Pertl-Obermeyer H, Obermeyer G, Schulze WX (2013) Sucrose-induced receptor kinase SIK1 regulates a plasma membrane aquaporin in *Arabidopsis*. *Mol Cell Proteomics* 12:2856-2873

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wudick MM, Li X, Valentini V, Geldner N, Chory J, Lin J, Maurel C, Luu DT (2015) Subcellular redistribution of root aquaporins induced by hydrogen peroxide. *Mol Plant* 8:1103-1114

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xin XF, Nomura K, Aung K, Velasquez AC, Yao J, Boutrot F, Chang JH, Zipfel C, He SY (2016) Bacteria establish an aqueous living space in plants crucial for virulence. *Nature* 539:524-529

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yang Z, Guo G, Zhang M, Liu CY, Hu Q, Lam H, Cheng H, Xue Y, Li J, Li N (2013) Stable isotope metabolic labeling-based quantitative phosphoproteomic analysis of *Arabidopsis* mutants reveals ethylene-regulated time-dependent phosphoproteins and putative substrates of constitutive triple response 1 kinase. *Mol Cell Proteomics* 12:3559-3582

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yip Delormel T, Boudsocq M (2019) Properties and functions of calcium-dependent protein kinases and their relatives in *Arabidopsis thaliana*. *New Phytol* 224:585-604

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang L, Chen L, Dong H (2019a) Plant Aquaporins in Infection by and Immunity Against Pathogens - A Critical Review. *Front Plant Sci* 10:632

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang S, Feng M, Chen W, Zhou X, Lu J, Wang Y, Li Y, Jiang CZ, Gan SS, Ma N, Gao J (2019b) In rose, transcription factor PTM balances growth and drought survival via PIP2;1 aquaporin. *Nat Plants* 5:290-299

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhu D, Wu Z, Cao G, Li J, Wei J, Tsuge T, Gu H, Aoyama T, Qu LJ (2014) TRANSLUCENT GREEN, an ERF family transcription factor, controls water balance in *Arabidopsis* by activating the expression of aquaporin genes. *Mol Plant* 7:601-615

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zwiewka M, Nodzynski T, Robert S, Vanneste S, Friml J (2015) Osmotic Stress Modulates the Balance between Exocytosis and Clathrin-Mediated Endocytosis in *Arabidopsis thaliana*. *Mol Plant* 8:1175-1187

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)