



Astrocytic IP3Rs: Beyond IP3R2

Mark W Sherwood, Misa Arizono, Aude Panatier, Katsuhiko Mikoshiba, Stéphane
H R Oliet

► To cite this version:

Mark W Sherwood, Misa Arizono, Aude Panatier, Katsuhiko Mikoshiba, Stéphane H R Oliet. Astrocytic IP3Rs: Beyond IP3R2. *Frontiers in Cellular Neuroscience*, 2021, 15, <10.3389/fncel.2021.695817>. <hal-03447677>

HAL Id: hal-03447677

<https://hal.science/hal-03447677v1>

Submitted on 24 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Astrocytic IP₃Rs: Beyond IP₃R2

Mark W. Sherwood^{1*†}, Misa Arizono^{2*†}, Aude Panatier¹, Katsuhiko Mikoshiba^{3,4,5} and Stéphane H. R. Oliet¹

¹ University of Bordeaux, INSERM, Neurocentre Magendie, U1215, Bordeaux, France, ² University of Bordeaux, CNRS, Interdisciplinary Institute for Neuroscience, IINS, UMR 5297, Bordeaux, France, ³ ShanghaiTech University, Shanghai, China, ⁴ Faculty of Science, Toho University, Funabashi, Japan, ⁵ RIKEN CLST, Kobe, Japan

OPEN ACCESS

Edited by:

Yu-Wei Wu,
Academia Sinica, Taiwan

Reviewed by:

Dmitri A. Rusakov,
University College London,
United Kingdom
Vladimir Grubišić,
Michigan State University,
United States

*Correspondence:

Mark W. Sherwood
MWSherwood@protonmail.com
Misa Arizono
arizono0202@gmail.com

[†]These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Non-Neuronal Cells,
a section of the journal
Frontiers in Cellular Neuroscience

Received: 15 April 2021

Accepted: 30 June 2021

Published: 30 July 2021

Citation:

Sherwood MW, Arizono M,
Panatier A, Mikoshiba K and
Oliet SHR (2021) Astrocytic IP₃Rs:
Beyond IP₃R2.
Front. Cell. Neurosci. 15:695817.
doi: 10.3389/fncel.2021.695817

Astrocytes are sensitive to ongoing neuronal/network activities and, accordingly, regulate neuronal functions (synaptic transmission, synaptic plasticity, behavior, etc.) by the context-dependent release of several gliotransmitters (e.g., glutamate, glycine, D-serine, ATP). To sense diverse input, astrocytes express a plethora of G-protein coupled receptors, which couple, via G_{i/o} and G_q, to the intracellular Ca²⁺ release channel IP₃-receptor (IP₃R). Indeed, manipulating astrocytic IP₃R-Ca²⁺ signaling is highly consequential at the network and behavioral level: Depleting IP₃R subtype 2 (IP₃R2) results in reduced GPCR-Ca²⁺ signaling and impaired synaptic plasticity; enhancing IP₃R-Ca²⁺ signaling affects cognitive functions such as learning and memory, sleep, and mood. However, as a result of discrepancies in the literature, the role of GPCR-IP₃R-Ca²⁺ signaling, especially under physiological conditions, remains inconclusive. One primary reason for this could be that IP₃R2 has been used to represent all astrocytic IP₃Rs, including IP₃R1 and IP₃R3. Indeed, IP₃R1 and IP₃R3 are unique Ca²⁺ channels in their own right; they have unique biophysical properties, often display distinct distribution, and are differentially regulated. As a result, they mediate different physiological roles to IP₃R2. Thus, these additional channels promise to enrich the diversity of spatiotemporal Ca²⁺ dynamics and provide unique opportunities for integrating neuronal input and modulating astrocyte–neuron communication. The current review weighs evidence supporting the existence of multiple astrocytic-IP₃R isoforms, summarizes distinct sub-type specific properties that shape spatiotemporal Ca²⁺ dynamics. We also discuss existing experimental tools and future refinements to better recapitulate the endogenous activities of each IP₃R isoform.

Keywords: astrocyte, inositol triphosphate (IP₃) receptor, IP₃R subtypes, calcium, GPCR, tripartite synapse, gliotransmission

INTRODUCTION

Over the last three decades, Ca²⁺ imaging has revealed new roles for astrocytes. Indeed, astrocytic Ca²⁺ signaling was shown to regulate synaptic transmission, synaptic plasticity, and to influence behavior (Araque et al., 2014; Park and Lee, 2020). Inositol 1,4,5-trisphosphate receptors (IP₃Rs) mediated Ca²⁺ signaling (IP₃R-Ca²⁺) is regarded a primary generator of astrocytic Ca²⁺ signaling. Upon activation of G_q-GPCRs, the main input pathway of astrocytes, phospholipase C breaks down PIP₂ into DAG and IP₃, activating IP₃R predominantly located on the membrane of endoplasmic reticulum (ER) resulting in Ca²⁺ release (Bootman et al., 2001). This IP₃R-mediated Ca²⁺ signaling is considered to trigger the activity-dependent and selective release of chemical transmitters (gliotransmitters) such as glutamate, D-serine, and ATP, which have distinct influences

over neuronal activity. Initially, IP₃R subtype 2 (IP₃R2) was the only recognized Ca²⁺ channel in astrocytes; However, advanced Ca²⁺ imaging techniques have since identified novel Ca²⁺ sources, including mitochondria (Agarwal et al., 2017), transient receptor potential ankyrin 1 (Shigetomi et al., 2012, 2013b), L-type voltage gated Ca²⁺ channels (Letellier et al., 2016), sodium/calcium exchanger (Kirischuk et al., 1997; Boddum et al., 2016; Rose et al., 2020), and transient receptor potential canonical (Shiratori-Hayashi et al., 2020) amongst others, thereby expanding the known Ca²⁺ signaling toolkit of astrocytes. Doubtlessly additional Ca²⁺ channels and sources will emerge in the future.

While the field's focus has moved on from understanding IP₃Rs to identifying new Ca²⁺ sources, understanding IP₃R signaling in astrocytes remains highly relevant. Indeed, IP₃Rs are the primary target for manipulating astrocytic activity, and such manipulations have proven to be very consequential in many studies. Since most of these manipulations indiscriminately influence all IP₃R subtypes, this could reflect the key role played by IP₃R subtypes other than IP₃R2, namely IP₃R1 and IP₃R3, which were mostly overlooked. In this review, we summarize the evidence for different subtypes of IP₃R and discuss how we can better study the role of IP₃R-Ca²⁺ signaling in astrocytes which is one of the core issues in understanding astrocyte physiology.

EVIDENCE FOR THREE SUBTYPES

The Dogma: IP₃R2 the Sole Functional Astrocytic IP₃R

There are three mammalian IP₃R subtypes, i.e., IP₃R1 (Furuichi et al., 1989; Mignery et al., 1989; Yamada et al., 1994), IP₃R2 (Mignery et al., 1990; Südhof et al., 1991; Yamamoto-Hino et al., 1994; Iwai et al., 2005), and IP₃R3 (Blondel et al., 1993; Yamamoto-Hino et al., 1994; Iwai et al., 2005). Among them, IP₃R2 has widely been accepted as the only functional IP₃R subtype present in astrocytes, and consequently, the IP₃R2KO model mouse has been at the center of numerous important studies (Aguilhon et al., 2010; Takata et al., 2011; Chen et al., 2012; Navarrete et al., 2012, 2019; Cao et al., 2013; Perez-Alvarez et al., 2014; Petravic et al., 2014; Gómez-Gonzalo et al., 2015, 2017; Mariotti et al., 2016; Monai et al., 2016; Perea et al., 2016; Martin-Fernandez et al., 2017; Tanaka et al., 2017). Although an important model in the astrocyte field, the belief that knocking out IP₃R type-2 abolishes IP₃ induced Ca²⁺ release (ICR) entirely appears to need remedying. This section considers the historical data from which the dogmatic view of astrocytic-IP₃R2 has flowed and reviews the evidence for other IP₃R subtypes.

Astrocyte Proteome

Several studies explored IP₃Rs using immunohistochemistry which provided a consensus over the expression of IP₃R2 in hippocampal/cortical astrocytes and Bergmann glia (Sharp et al., 1999; Holtzclaw et al., 2002; Hertle and Yeckel, 2007; Takata et al., 2011; Chen et al., 2012). While there are some conflicting reports over the immunoreactivity of IP₃R3 in astrocytes and Bergmann glia (Sugiyama et al., 1994;

Yamamoto-Hino et al., 1995; Hamada et al., 1999; Sharp et al., 1999; Holtzclaw et al., 2002), IP₃R1 immunoreactivity was not initially observed in glia (Nakanishi et al., 1991; Dent et al., 1996; Hamada et al., 1999; Sharp et al., 1999; Holtzclaw et al., 2002; Hertle and Yeckel, 2007). These findings supported the view that IP₃R2 is the predominant astrocytic IP₃R. However, these results may also reflect limitations of the available IP₃R antibodies or the difficulty of accurately assigning proteins located within ultrathin astrocyte processes, which are below the resolution limit of conventional microscopy (Panatier et al., 2014; Arizono et al., 2020) and buried amongst neuronal dendrites.


IP₃R1 immunoreactivity was recently detected in spinal dorsal horn astrocytes (Shiratori-Hayashi et al., 2020) and, albeit with low stringency, in isolated astrocytes from adult mice (Chai et al., 2017). Using a state of the art TurboID construct to biotinylate proteins in the immediate proximity of tripartite synapses, Takano et al. (2020) report enrichment of IP₃R1 protein in the peri-synaptic astrocytic compartment (Takano et al., 2020). This finding, however, should be interpreted with some caution as identified proteins were assigned to astrocytes based on published mRNA datasets (Zhang et al., 2014, 2016). Nevertheless, IP₃R1 protein enrichment in fine astrocytic processes is consistent with Ca²⁺ imaging studies (Sherwood et al., 2017) and could account for the poor detection in various assays which favor detection in large subcellular compartments, i.e., major processes and soma of astrocytes. Notably, IP₃R2, which is reported to be in the soma and main branches (Chen et al., 2012), was not enriched in peri-synaptic astrocytic compartments (Takano et al., 2020), likely reflecting the different subcellular distribution of IP₃R1 and IP₃R2 (Figure 1A and Table 1).

Astrocyte Transcriptome

IP₃R1, IP₃R2, and IP₃R3 are encoded by the respective genes ITPR1, ITPR2, and ITPR3. Notably, mRNA for all three genes have been detected in astrocytes isolated from young and aged mouse brain (Cahoy et al., 2008; Zhang et al., 2014; Chai et al., 2017; Clarke et al., 2018) as well as humans (Zhang et al., 2016). Furthermore, ITPR1 and ITPR2 are actively translated in astrocytes of adult mice (Srinivasan et al., 2016; Chai et al., 2017; Clarke et al., 2018; Yu et al., 2018) indicating that their proteins are produced in astrocytes. In astrocytes the isoform transcript abundance is generally ITPR2 > ITPR1 >>> ITPR3 (ITPR3 mRNA is present in very small amounts and may be negligible). However, these transcripts are developmentally and differentially regulated across brain regions and in some instances ITPR1 mRNA appears to be more abundant than ITPR2 mRNA (Yu et al., 2018; Table 1).

Ca²⁺ Imaging

Over the last decade, studies show that deletion of IP₃R2 does not abolish Ca²⁺ signaling in astrocytes. It is now generally agreed that bulk/somatic cytosolic Ca²⁺ responses are hard to detect in IP₃R2KO astrocytes (Petravic et al., 2008, 2014; Aguilhon et al., 2010, 2013; Takata et al., 2011, 2013; Nizar et al., 2013). However, with a detailed examination, rich Ca²⁺ activity can be observed within fine astrocyte processes (Di Castro et al., 2011; Hausteine et al., 2014; Kanemaru et al., 2014; Srinivasan et al., 2015;



Rungta et al., 2016; Agarwal et al., 2017; Sherwood et al., 2017). Non-IP₃R2 Ca²⁺ activity has largely been interpreted as evidence for non-IP₃R Ca²⁺ stores, nevertheless, these activities could equally arise from IP₃R1 and or IP₃R3 (Tamamushi et al., 2012; Sherwood et al., 2017). Indeed, Ca²⁺ release from the ER was recently reported for IP₃R2KO astrocytes (Okubo et al., 2019). In Bergmann glia, while IP₃R2 was shown to be a dominant subtype, IP₃R1 and IP₃R3 were also shown to contribute to rapid Ca²⁺ events in the processes (Tamamushi et al., 2012). Similarly, using IP₃R2KO and IP₃R2/3 double KO mice, we found that IP₃R2 is involved in global Ca²⁺ release, whereas local Ca²⁺ signals involve IP₃R1 and IP₃R3 (**Figure 1B**) (Sherwood et al., 2017) in hippocampal astrocytes. Knock-down of IP₃R1 in dorsal spinal cord astrocytes of IP₃R2KO mice also unmasked IP₃R1 mediated Ca²⁺ signals (Shiratori-Hayashi et al., 2020). Furthermore, the selective activation of G_q-GPCR/IP₃R/Ca²⁺ signaling in astrocytes, using DREADDs (designer receptor exclusively activated by designer drug, see section “Activation of IP₃ Induced Ca²⁺ Release in Astrocytes”), triggered Ca²⁺ events in the astrocytic processes of IP₃R2KO mice (Wang et al., 2021) indicating the presence of IP₃R1/3.

2-APB was introduced as an antagonist of IP₃Rs (Maruyama et al., 1997) and has been widely used to investigate the contribution of IP₃Rs to cellular Ca²⁺ signaling. 2-APB appears to preferentially block IP₃R1 and IP₃R3, whereas cells predominantly expressing IP₃R2 seem largely insensitive (Kukkonen et al., 2001; Bootman et al., 2002; Saleem et al., 2014). Thus, the fact that 2-APB reduces astrocyte Ca²⁺ amplitude and responsiveness in diverse brain regions (Sul et al., 2004; Young et al., 2010; Tamura et al., 2014; Tang et al., 2015; Arizono et al., 2020) and spinal dorsal horn (Shiratori-Hayashi et al., 2020) may support the presence of functional IP₃R1 and IP₃R3 in astrocytes. The functional role of non-IP₃R2 in astrocytic Ca²⁺ signaling remains to be determined, it is possible that some membrane receptors are functionally coupled solely to IP₃R1 or IP₃R3 and generate local Ca²⁺ events, which may or may not be involved in triggering IP₃R2 dependent Ca²⁺ waves (Petravic et al., 2008).

Alternatively, it is possible that IP₃Rs exist as heterotetramers (Wojcikiewicz and He, 1995; Nucifora et al., 1996).

Phenotypic Comparison

Comparison of WT and total IP₃R2KO mice has revealed some important physiological roles of IP₃R2 signaling in astrocytes, i.e., motor learning (Padmashri et al., 2015) modulating depressive-like behaviors (Cao et al., 2013) but this remains controversial (Petravic et al., 2014). The possible roles of astrocytic IP₃R1 and IP₃R3 may be gleaned by comparing WT and IP₃R2KOs with manipulations that impair all IP₃R subtypes. For example, while IP₃R2KO has no impact on sleep (Cao et al., 2013) overexpressing a transgene for the IP₃ hydrolyzing enzyme, IP₃-5-phosphatase, in astrocytes, suppresses Ca²⁺ release from all IP₃R subtypes and disrupts sleep (Foley et al., 2017). Similarly, although IP₃R2KO had no impact on hippocampal LTP (Agulhon et al., 2010), we showed that loading a single astrocyte with the membrane-impermeable pan-IP₃R antagonist, heparin, blocks long-term synaptic potentiation (Sherwood et al., 2017). These studies, taken together, indicate that non-IP₃R2 IP₃Rs regulate sleep and hippocampal LTP.

Summary

The expression of multiple IP₃R subtypes in astrocytes has a wide-reaching implication in the field. Studies using IP₃R2KO as a model for blocked IICR would need to be re-assessed to include the possibility of IICR mediated by IP₃R1 and IP₃R3.

PROSPECTUS/ADVANTAGE OF MULTIPLE IP₃R ISOFORMS

Different Properties of IP₃R Subtypes

The three IP₃R subtypes share only 65–85% homology accounting for many of the subtype-specific properties leading to particular spatiotemporal features of Ca²⁺ responses. Although

TABLE 1 | Dissecting IP₃R subtypes.

Transcriptome/translatome							
Species	Age (weeks)	Region	ITPR1 (FPKM)	ITPR2 (FPKM)	ITPR3 (FPKM)	Comment	References
Mouse	1	Str	1.4	7.6	0.1	Translatome	Clarke et al., 2018
	4.6		7.8	4.5	0.3		
	9		28.5; 85.0	17.4; 6.4	0.2; 0.1		
	9		27.1	10.1			
	10		10.9	2.6	0.1		
	38		13.1	2.5	0.1		
	96		16.1	4.4	0.0		
	0–2.4	Forebrain	Present	Present	Absent	Transcriptome	Clarke et al., 2018
	1	Ctx	0.4	12.6	0.1		Cahoy et al., 2008
	1		1.5	4.2	0.1	Translatome	Zhang et al., 2014
	Adult	Hc	2.6	7.1			Clarke et al., 2018
	4.6		5.4	6.0	0.6	Translatome	Srinivasan et al., 2016
	10		13.7	5.3	0.1		Clarke et al., 2018
	38		12.3	4.6	0.0		
	96		17.5	4.5	0.0		
	1		0.8	4.8	0.1	Transcriptome	Clarke et al., 2018
	4.6		3.0	5.4	0.3		
	9		6.7	11.7		Translatome	Chai et al., 2017
	10		8.6	4.0	0.1		Clarke et al., 2018
	38		4.4	4.8	0.0		
	96		4.4	6.9	0.1	Transcriptome	Zhang et al., 2016
	Human	8–63 years	Ctx	0.7	13.7		
		17–20 gw	Brain	0.4	2.6	0.2	
Proteome							
Species	Age (weeks)	Region	IP ₃ R1	IP ₃ R2	IP ₃ R3	References	
Rat	Adult	Hc, Cb, Cc, ScNu	No	Yes	No	Holtzclaw et al., 2002	
	>8	Hc	No	Yes	No	Hertle and Yeckel, 2007	
	8	Cb, Ht, Hc	No	No	Yes	Yamamoto-Hino et al., 1995	
		ScNu, Ht, Ctx	No	–	Yes	Hamada et al., 1999	
Mouse		SDH	Yes	–	–	Shiratori-Hayashi et al., 2020	
		Hc, Str	Yes	Yes	No	Chai et al., 2017	
	7	Ctx	Yes	–	–	Takano et al., 2020	
	>6	Ctx	–	Yes	–	Chen et al., 2012	
	8–12	Ctx	–	Yes	–	Takata et al., 2011	
	Human		Ctx	Yes	Yes	Yes	Hur et al., 2010
Available pharmacological/genetic tools							
	Tool	IP ₃ R	References				
Transgenic mice	IP ₃ R1 KO	1	Tamamushi et al., 2012				
	IP ₃ R2 KO	2	Fiacco et al., 2007; Petravicz et al., 2008; Agulhon et al., 2010; Takata et al., 2011, 2013; Navarrete et al., 2012; Tamamushi et al., 2012; Cao et al., 2013; Nizar et al., 2013; Bonder and McCarthy, 2014; Gómez-Gonzalo et al., 2015, 2017; Mariotti et al., 2016; Monai et al., 2016; Perea et al., 2016; Martin-Fernandez et al., 2017; Sherwood et al., 2017; Tanaka et al., 2017; Wang et al., 2021				
	IP ₃ R3 KO	3	Tamamushi et al., 2012				
	IP ₃ R2/3KO	2 and 3	Tamamushi et al., 2012; Sherwood et al., 2017				
	IP ₃ R1 cKO	1	# Sugawara et al., 2013				
	IP ₃ R2 cKO	2	Chen et al., 2012; Petravicz et al., 2014; Padmashri et al., 2015; Wang et al., 2021				
Genetic tools for inhibition	IP ₃ sponge	1, 2, and 3	Xie et al., 2010; Tanaka et al., 2013				
	5ppase	1, 2, and 3	Kanemaru et al., 2007; Foley et al., 2017				
	IP ₃ R1 shRNA	1	Shiratori-Hayashi et al., 2020				

(Continued)

TABLE 1 | Continued

Available pharmacological/genetic tools

	Tool	IP ₃ R	References
Pharmacology	IP ₃ R2 shRNA	2	#
	IP ₃ R3 shRNA	3	#
	IP ₃ R1 Ab	1	## Miyazaki et al., 1992; Inoue et al., 1998; Nishiyama et al., 2000
	IP ₃ R2 Ab	2	## Gerasimenko et al., 2009
	IP ₃ R3 Ab	3	## Gerasimenko et al., 2009
	Heparin	1, 2, and 3	\$ Sherwood et al., 2017
Genetic tools for activation	(1,2,3,4,6)IP ₅ dimer	1; 2/3?	## Konieczny et al., 2016
	MrgA1 (G _i GPCR)	?	Fiocco et al., 2007; Agulhon et al., 2010; Cao et al., 2013
	hM3Dq	2 and 1 or 3	Agulhon et al., 2013; Bonder and McCarthy, 2014; Yang et al., 2015; Martin-Fernandez et al., 2017; Adamsky et al., 2018; Wang et al., 2021
	hM4Di	?	Yang et al., 2015
	Opto-α ₁ AR	?	Figueiredo et al., 2014; Adamsky et al., 2018; Iwai et al., 2021
	Melanopsin	?	Mederos et al., 2019

Untested in astrocytes; \$ requires loading using an astrocytic patch-pipette; ?, unknown; FPKM, fragments per kilobase of transcript per million mapped reads; Ctx, cortex; Str, striatum; Hc, hippocampus; Ht, hypothalamus; Cb, cerebellum; Cc, corpus callosum; ScNu, suprachiasmatic nucleus; SDH, spinal dorsal horn; gw, gestational weeks. Darker shades of blue represents higher FPKM values.

high homology is observed in regions critical for forming the IP₃-gated Ca²⁺ channel, each subtype has a different IP₃ affinity; IP₃R2 > IP₃R1 > IP₃R3 (Zhang et al., 2011). High IP₃ affinity of IP₃R2 has been associated with slower kinetics and more prolonged duration of IP₃R2-mediated Ca²⁺ microdomains, or Ca²⁺ puffs (Mataragka and Taylor, 2018).

Importantly, IP₃R channel activity is not only regulated by IP₃ but also by Ca²⁺ (Finch et al., 1991). The synergy between IP₃ and Ca²⁺ creates repetitive IP₃R activation and inhibition, resulting in Ca²⁺ oscillations. Such oscillations are crucial to protect cells from extended Ca²⁺ elevation that can often be toxic to the cell (Berridge et al., 2000). The oscillatory pattern is considered necessary for many cellular processes, such as fertilization (Miyazaki et al., 1992). IP₃R2 mediates long-lasting, regular Ca²⁺ oscillations, whereas IP₃R1 or IP₃R3 tend to exhibit mono-phasic transients or very rapidly dampened Ca²⁺ oscillations (Miyakawa, 1999).

The difference in IP₃R subtype properties is further characterized by various binding partners, including kinase and phosphatases, which can further fine-tune Ca²⁺ profiles. Interestingly, while there are many interacting partners common to all three IP₃R subtypes, the nature of their regulation can be subtype-specific. For instance, protein kinase C, depending on the IP₃R subtype, can either be stimulatory or inhibitory; this difference likely reflects isoform-specific phosphorylation sites. Further detailed biochemical study (Hamada et al., 2017) is required to investigate how various IP₃R binding molecules specifically regulate each IP₃R isoform (Hamada and Mikoshiba, 2020).

Different Distribution and Role of IP₃R Subtypes Within Various Tissues

One important feature that defines the subtype-specific role of IP₃Rs *in vivo* is the tissue distribution patterns. While IP₃R1 is mostly expressed in the central nervous system, IP₃R2

and IP₃R3 are broadly expressed in various organs such as the heart, pancreas, liver, and salivary glands (Hisatsune and Mikoshiba, 2017). This distribution pattern is tightly linked to the physiological role of IP₃R subtypes. Reflecting its rich expression in Purkinje cells, mice lacking IP₃R1 exhibit severe cerebellar ataxia, a seizure-like posture, impaired cerebellar LTD. (Inoue et al., 1998), and die within 3–4 weeks of birth (Matsumoto et al., 1996). The dysregulation of IP₃R1 is linked with other brain disorders such as Huntington's disease and Alzheimer's disease (Hisatsune and Mikoshiba, 2017). IP₃R2 is associated with sweating (Klar et al., 2014), bone formation (Kuroda et al., 2008), and heart hypertrophy (Nakayama et al., 2010; Drawnel et al., 2012; Vervloessem et al., 2015). IP₃R3 plays a role in taste sensing (Hisatsune et al., 2007) and hair cycle (Sato-Miyaoka et al., 2012). IP₃R2 and IP₃R3 together are associated with the heart's development (Uchida et al., 2010, 2016) and secretion of saliva and tears (Futatsugi, 2005; Inaba et al., 2014).

Different Distribution and Role of IP₃R Subtypes Within a Cell

Some cells express multiple IP₃R subtypes, enabling each subtype to uniquely contribute to Ca²⁺ profiles and cellular functions. For instance, in HeLa cells, knock-down of IP₃R1 terminates Ca²⁺ oscillations, whereas knock-down of IP₃R3 results in more robust and long-lasting Ca²⁺ oscillations (Hattori et al., 2004). The specific contribution of IP₃R subtypes can also depend on their distinct subcellular distribution, as seen in pancreatic acinar cells (Lur et al., 2011), COS cells (Pantazaka and Taylor, 2011), and DT40 cells (Bartok et al., 2019). In Bergmann glia, knocking out IP₃R2 resulted in decreased agonist-induced Ca²⁺ release while knocking out IP₃R1 and IP₃R3 resulted in delaying the peak of agonist-induced Ca²⁺ release specifically in astrocytic processes (Tamamushi et al., 2012), suggesting their subtype-specific distribution. It is likely that astrocytes, which

express multiple IP₃R subtypes, also take advantage of subtype-specific distribution.

In astrocytes, IP₃Rs are predominantly located on thapsigargin sensitive ER Ca²⁺ store. The ER in astrocytes may be found throughout the cell in the soma, major processes (Okubo et al., 2019, 2020), and peri-synaptic astrocytic processes (Bergersen et al., 2012), in close association with adhesion junctions (puncta adherentia) between dendritic spines and astrocytic processes (Spacek and Harris, 1998). Although recent studies indicate that ER and other membrane-bound organelles are absent from peri-synaptic processes (Patrushev et al., 2013) this likely reflects their sensitivity to chemical fixation (Korogod et al., 2015). The ER store in astrocytes is heterogeneous and organized into sub-compartments that can release Ca²⁺ independently (Golovina and Blaustein, 1997, 2000). It would be fascinating to see if IP₃R subtypes are located to specific functional domains, e.g., signaling domains of membrane receptors, ER-mitochondria contacts (Bartok et al., 2019), and ER-plasma membrane junctions (Thillaiappan et al., 2017). In addition to the ER, astrocytic IP₃Rs can also be found on other Ca²⁺ stores, with unique properties, i.e., the large dense-core vesicles (Hur et al., 2010), which may be comparable to the thapsigargin insensitive, Bafilomycin A1 sensitive, acidic Ca²⁺ stores previously described in secretory cells (Gerasimenko et al., 1996, 2009, 2011; Hur et al., 2010). Notably, compared to the ER, acidic Ca²⁺ stores can exhibit enhanced sensitivity to IP₃ (Yoo, 2010). Other potential IP₃-sensitive Ca²⁺ stores include the nuclear envelope (Gerasimenko et al., 1995; Petersen et al., 1998), nucleoplasm (Echevarria et al., 2003), Golgi (Pinton et al., 1998), and plasma membrane (Dellis et al., 2006).

Summary

While IP₃R subtypes are regulated by IP₃ and Ca²⁺ and have many common interacting partners, they differ in how they are affected by these regulators. Such differences enrich the diversity of spatio-temporal Ca²⁺ profiles created by IP₃Rs. The IP₃R subtype expression pattern, *in vivo*, is tissue specific and their subcellular localization is highly variable and dependent on cell types, and this carries important functional implications. Together with recent reports showing the distinct role of IP₃R1 and IP₃R3 in Bergmann glia and astrocytes, these facts support the view that IP₃R isoforms 1 and 3 are unique Ca²⁺ channels that need to be addressed independently of IP₃R2.

TOOLS TO DISSECT THE ROLE OF IP₃R ISOFORMS

Experimental and Analytical Tools

To understand the role of the various Ca²⁺ signals in astrocyte physiology, it will be necessary to make quantitative measurements (Neher, 2008). Progress in this direction has been frustrated by the unique astrocyte morphology and difficulties in interpreting recorded Ca²⁺-dependent fluorescent signals (Rusakov, 2015). To accurately capture Ca²⁺ dynamics

in sub-cellular compartments, there is a need to adopt imaging techniques with improved resolution and to develop tools for efficient analysis in three-dimensional (Bindocci et al., 2017; Romanos et al., 2019). Because of our poor understanding of functional compartments, analysis of astrocytic Ca²⁺ dynamics has been moving toward state-of-the-art event-based analysis (Romanos et al., 2019; Wang et al., 2019; Bjørnstad et al., 2021), nevertheless ROI (region-of-interest) based analysis, informed by cellular anatomy (functional compartments) and molecular architecture, has been critical for understanding neuron physiology (e.g., spines and boutons). To this end, the identification of morphologically distinct subcellular compartments are promising targets for classical ROI based analysis, i.e., “glial microdomains” on Bergmann glial processes (Grosche et al., 1999) and “astrocytic compartments” on major branches (Panatier et al., 2011), both revealed using confocal microscopy, and astrocytic nodes and shafts within the spongiform structure visualized using live STED microscopy (Arizono et al., 2020). The ultimate goal of extracting quantitative Ca²⁺ dynamics from fluorescent data is non-trivial but achievable using realistic biophysical cell models (Rusakov, 2015; Denizot et al., 2019), a task simplified by the recent development of the open-source flexible model builder ASTRO (Savtchenko et al., 2018). For an accurate understanding of Ca²⁺ dynamics, it will be necessary to constrain models further using empirically determined details, e.g., receptor kinetics, expression patterns, endogenous Ca²⁺ buffering, etc.

Pharmacological Tools

It is difficult to disentangle the physiological roles of IP₃R subtypes in cells that typically express complex mixtures of homo- and hetero-tetrameric IP₃Rs. There are no ligands that usefully distinguish among IP₃R subtypes (Saleem et al., 2013a,b) and nor are there effective antagonists that lack serious side effects (Michelangeli et al., 1995). Of the available antagonists, heparin is currently the most useful. Heparin is a membrane impermeant pan-IP₃R inhibitor that may be selectively loaded into astrocytes using a whole-cell patch-pipette (Sherwood et al., 2017). Recent developments report small impermeant competitive antagonists of IP₃R1, which, compared to heparin, are likely to have fewer off-targets (Konieczny et al., 2016). Well-characterized function-blocking monoclonal antibodies are powerful tools to specifically inhibit IP₃R subtypes (Miyazaki et al., 1992; Inoue et al., 1998; Nishiyama et al., 2000; Gerasimenko et al., 2009). This technology has not yet been applied to astrocytes.

Genetic Tools

Inhibition of IP₃ Induced Ca²⁺ Release

IP₃R2KO and conditional-KO (cKO) mice (Petravicz et al., 2014; Padmashri et al., 2015; Wang et al., 2021) are widely used, however, as highlighted above, deletion of IP₃R2 does not abolish IICR. IICR can be suppressed, irrespective of the underlying receptor, using an IP₃-sponge to buffer IP₃ (Xie et al., 2010; Tanaka et al., 2013), or an IP₃-5'-phosphatase transgene to enhance IP₃ metabolism

(Kanemaru et al., 2007; Foley et al., 2017). To study the physiological role of IP₃R subtypes it is necessary to develop inducible cKO for IP₃R1 (Sugawara et al., 2013) and IP₃R3. Specific knock-down of IP₃R subtypes can also be achieved using viruses to introduce short hairpin RNA into astrocytes (Shiratori-Hayashi et al., 2020).

Activation of IP₃ Induced Ca²⁺ Release in Astrocytes Pharmacogenetics

DREADDs (designer receptor exclusively activated by designer drug) enable the selective activation of GPCR-IP₃R-Ca²⁺ signaling in astrocytes. The most used DREADDs are the excitatory G_q or inhibitory G_i-coupled receptors, hM3Dq and hM4Di, respectively (derived from human M3/M4 muscarinic receptor). Both receptors are activated by a pharmacologically inert but bioavailable ligand clozapine-N-oxide (CNO) while being non-responsive to endogenous GPCR ligands (Agulhon et al., 2013). hM3Dq has been used to demonstrate an astrocytic role in behaviors such as food intake (Yang et al., 2015), fear response (Martin-Fernandez et al., 2017), and memory recall (Adamsky et al., 2018). While the DREADDs enables selective activation of astrocytes, they have two major drawbacks: Firstly, the exogenous receptors have not been targeted to specific signaling domains and are likely to be spatially uncoupled from signaling nanodomains critical to IP₃R physiology (Bootman et al., 2001); Secondly, because of the sustained (hour-long) activation by exogenous ligands (Iwai et al., 2021), temporal features of astrocyte signaling are lost. While perhaps mimicking global Ca²⁺ surges, the available DREADDs are unlikely to recapitulate many of the local Ca²⁺ transients typically observed within fine astrocytic processes (Shigetomi et al., 2013a; Arizono et al., 2020).

Optogenetics

To achieve temporal control, an optogenetic approach has been developed for the reliable stimulation of endogenous GPCR-IP₃R-Ca²⁺ signaling cascade using light. Light activation has been achieved by introducing to astrocytes either a mammalian light-sensitive G_q/G_{i/o}-protein-coupled photopigment, Melanopsin (Panda, 2005; Bailes and Lucas, 2013; Mederos et al., 2019), or light-activated chimeric GPCRs, termed OptoXRs. OptoXRs are generated by replacing the intracellular loops of a light-sensitive GPCR, e.g., rhodopsin, with those of a donor GPCR, e.g., G_q-coupled human adrenergic receptor α_{1a} (Airan et al., 2009; Figueiredo et al., 2014; Tang et al., 2014; Adamsky et al., 2018; Iwai et al., 2021).

REFERENCES

- Adamsky, A., Kol, A., Kreisel, T., Doron, A., Ozeri-Engelhard, N., Melcer, T., et al. (2018). Astrocytic activation generates de novo neuronal potentiation and memory enhancement. *Cell* 174, 59–71.e14. doi: 10.1016/j.cell.2018.05.002
- Agarwal, A., Wu, P.-H., Hughes, E. G., Fukaya, M., Tischfield, M. A., Langseth, A. J., et al. (2017). Transient opening of the mitochondrial permeability

Summary – Future Developments

While having great potential for controlling astrocytic activation, a central question is to what extent do the chimeras mimic the signaling of wild-type receptors. GPCRs can have multiple signaling axis, e.g., multiple G-protein axes, β-arrestins, wnt-frizzled, or the hedgehog-smoothened axes (Bailes and Lucas, 2013; Tichy et al., 2019), and the signaling axes bias is often not characterized but can have profound side effects on physiology (Agulhon et al., 2013; Tichy et al., 2019). Indeed, the functional outcome of activating G_q in astrocytes using different exogenous receptor (i.e., hM3Dq and MrgA1) is not reproducible (Agulhon et al., 2010; Adamsky et al., 2018). While multiple signaling axis could confuse the role of IICR, they may be required to obtain an optimal IP₃R response (Konieczny et al., 2017). Another primary concern is that DREADDs and optoXRs likely activate IP₃Rs from cellular compartments distinct from those of the endogenous receptors, limiting their ability to recreate physiologically relevant Ca²⁺ profiles. To address this, next-generation activation tools are being engineered to mimic the subcellular distribution of endogenous receptors (Oh et al., 2010; Masseck et al., 2014; Spoida et al., 2014; Tichy et al., 2019).

In the last decade, substantial progress has been made revealing diverse spatio-temporal Ca²⁺ signaling in astrocytes. Understanding the subtleties of these signals will require detailed knowledge of the astrocytic Ca²⁺ signaling toolbox along with the generation and characterization of more sophisticated tools to control and accurately recapitulate the physiologically relevant Ca²⁺ signals.

AUTHOR CONTRIBUTIONS

MS and MA performed the literature survey, wrote the manuscript, and prepared the figures and tables. MS, MA, AP, KM, and SHRO reviewed, finalized, and approved the final version.

FUNDING

This review was funded by the CNRS, INSERM, LabEx BRAIN, and the Agence Nationale de la Recherche, grant numbers ANR-17-CE16-0002 to SHRO, and ANR-16-CE16-0001-0 to AP. MS was supported by a Takeda Science Foundation fellowship (Japan), and MA was supported by postdoctoral fellowships from RIKEN and JSPS (Japan). KM was supported by Scientific Research S (25221002) of JSPS and ICORP; ICORP-SORST of Japan Science and Technology Agency.

- transition pore induces microdomain calcium transients in astrocyte processes. *Neuron* 93, 587–605.e7. doi: 10.1016/j.neuron.2016.12.034
- Agulhon, C., Boyt, K. M., Xie, A. X., Friocourt, F., Roth, B. L., and McCarthy, K. D. (2013). Modulation of the autonomic nervous system and behaviour by acute glial cell G_q protein-coupled receptor activation in vivo: Glial GPCR signalling in physiology and behaviour in vivo. *J. Physiol.* 591, 5599–5609. doi: 10.1113/jphysiol.2013.261289

- Agulhon, C., Fiacco, T. A., and McCarthy, K. D. (2010). hippocampal short- and long-term plasticity are not modulated by astrocyte Ca²⁺ signaling. *Science* 327, 1250–1254. doi: 10.1126/science.1184821
- Airan, R. D., Thompson, K. R., Fenno, L. E., Bernstein, H., and Deisseroth, K. (2009). Temporally precise in vivo control of intracellular signalling. *Nature* 458, 1025–1029. doi: 10.1038/nature07926
- Araque, A., Carmignoto, G., Haydon, P. G., Oliet, S. H. R., Robitaille, R., and Volterra, A. (2014). Gliotransmitters travel in time and space. *Neuron* 81, 728–739. doi: 10.1016/j.neuron.2014.02.007
- Arizono, M., Inavalli, V. V. G. K., Panatier, A., Pfeiffer, T., Angibaud, J., Levet, F., et al. (2020). Structural basis of astrocytic Ca²⁺ signals at tripartite synapses. *Nat. Commun.* 11:1906. doi: 10.1038/s41467-020-15648-4
- Bailes, H. J., and Lucas, R. J. (2013). Human melanopsin forms a pigment maximally sensitive to blue light ($\lambda_{max} \approx 479$ nm) supporting activation of G q /11 and G i/o signalling cascades. *Proc. R. Soc. B Biol. Sci.* 280:20122987. doi: 10.1098/rspb.2012.2987
- Bartok, A., Weaver, D., Golenár, T., Nichtova, Z., Katona, M., Bánsághi, S., et al. (2019). IP₃ receptor isoforms differently regulate ER-mitochondrial contacts and local calcium transfer. *Nat. Commun.* 10:3726. doi: 10.1038/s41467-019-11646-3
- Bergersen, L. H., Morland, C., Ormel, L., Rinholm, J. E., Larsson, M., Wold, J. F. H., et al. (2012). Immunogold detection of L-glutamate and D-serine in small synaptic-like microvesicles in adult hippocampal astrocytes. *Cereb. Cortex* 22, 1690–1697. doi: 10.1093/cercor/bhr254
- Berridge, M. J., Lipp, P., and Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* 1, 11–21. doi: 10.1038/35036035
- Bindocci, E., Savtchouk, I., Liaudet, N., Becker, D., Carriero, G., and Volterra, A. (2017). Three-dimensional Ca²⁺ imaging advances understanding of astrocyte biology. *Science* 356:eaai8185. doi: 10.1126/science.aai8185
- Bjornstad, D. M., Åbjørnsbråten, K. S., Hennestad, E., Cunen, C., Hermansen, G. H., Bojarskaite, L., et al. (2021). Begonia-A two-photon imaging analysis pipeline for astrocytic Ca²⁺ signals. *Front. Cell. Neurosci.* 15:681066. doi: 10.3389/fncel.2021.681066
- Blondel, O., Takeda, J., Janssen, H., Seino, S., and Bell, G. I. (1993). Sequence and functional characterization of a third inositol trisphosphate receptor subtype, IP3R-3, expressed in pancreatic islets, kidney, gastrointestinal tract, and other tissues. *J. Biol. Chem.* 268, 11356–11363. doi: 10.1016/s0021-9258(18)82132-9
- Boddum, K., Jensen, T. P., Magloire, V., Kristiansen, U., Rusakov, D. A., Pavlov, I., et al. (2016). Astrocytic GABA transporter activity modulates excitatory neurotransmission. *Nat. Commun.* 7:13572. doi: 10.1038/ncomms13572
- Bonder, D. E., and McCarthy, K. D. (2014). Astrocytic Gq-GPCR-linked IP3R-dependent Ca²⁺ signaling does not mediate neurovascular coupling in mouse visual cortex in vivo. *J. Neurosci.* 34, 13139–13150. doi: 10.1523/JNEUROSCI.2591-14.2014
- Bootman, M. D., Collins, T. J., Mackenzie, L., Roderick, H. L., Berridge, M. J., and Peppiatt, C. M. (2002). 2-Aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca²⁺ entry but an inconsistent inhibitor of InsP₃ -induced Ca²⁺ release. *FASEB J.* 16, 1145–1150. doi: 10.1096/fj.02-0037rev
- Bootman, M. D., Lipp, P., and Berridge, M. J. (2001). The organisation and functions of local Ca(2+) signals. *J. Cell Sci.* 114, 2213–2222. doi: 10.1242/jcs.114.12.2213
- Cahoy, J. D., Emery, B., Kaushal, A., Foo, L. C., Zamanian, J. L., Christopherson, K. S., et al. (2008). A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function. *J. Neurosci.* 28, 264–278. doi: 10.1523/JNEUROSCI.4178-07.2008
- Cao, X., Li, L.-P., Wang, Q., Wu, Q., Hu, H.-H., Zhang, M., et al. (2013). Astrocyte-derived ATP modulates depressive-like behaviors. *Nat. Med.* 19, 773–777. doi: 10.1038/nm.3162
- Chai, H., Diaz-Castro, B., Shigetomi, E., Monte, E., Oceau, J. C., Yu, X., et al. (2017). Neural circuit-specialized astrocytes: transcriptomic, proteomic, morphological, and functional evidence. *Neuron* 95, 531–549.e9. doi: 10.1016/j.neuron.2017.06.029
- Chen, N., Sugihara, H., Sharma, J., Perea, G., Petracz, J., Le, C., et al. (2012). Nucleus basalis-enabled stimulus-specific plasticity in the visual cortex is mediated by astrocytes. *Proc. Natl. Acad. Sci. U.S.A.* 109, E2832–E2841. doi: 10.1073/pnas.1206557109
- Clarke, L. E., Liddel, S. A., Chakraborty, C., Münch, A. E., Heiman, M., and Barres, B. A. (2018). Normal aging induces A1-like astrocyte reactivity. *Proc. Natl. Acad. Sci. U.S.A.* 115, E1896–E1905. doi: 10.1073/pnas.1800165115
- Dellis, O., Dedos, S. G., Tovey, S. C., Taufiq-Ur-Rahman, Dubel, S. J., and Taylor, C. W. (2006). Ca²⁺ entry through plasma membrane IP₃ receptors. *Science* 313, 229–233. doi: 10.1126/science.1125203
- Denizot, A., Arizono, M., Nägerl, U. V., Soula, H., and Berry, H. (2019). Simulation of calcium signaling in fine astrocytic processes: effect of spatial properties on spontaneous activity. *PLoS Comput. Biol.* 15:e1006795. doi: 10.1371/journal.pcbi.1006795
- Dent, M. A., Raisman, G., and Lai, F. A. (1996). Expression of type 1 inositol 1,4,5-trisphosphate receptor during axogenesis and synaptic contact in the central and peripheral nervous system of developing rat. *Dev. Camb. Engl.* 122, 1029–1039. doi: 10.1242/dev.122.3.1029
- Di Castro, M. A., Chuquet, J., Liaudet, N., Bhaukaurally, K., Santello, M., Bouvier, D., et al. (2011). Local Ca²⁺ detection and modulation of synaptic release by astrocytes. *Nat. Neurosci.* 14, 1276–1284. doi: 10.1038/nn.2929
- Drawnel, F. M., Wachten, D., Molkentin, J. D., Maillet, M., Aronsen, J. M., Swift, F., et al. (2012). Mutual antagonism between IP3R2 and miRNA-133a regulates calcium signals and cardiac hypertrophy. *J. Cell Biol.* 199, 783–798. doi: 10.1083/jcb.201111095
- Echevarría, W., Leite, M. F., Guerra, M. T., Zipfel, W. R., and Nathanson, M. H. (2003). Regulation of calcium signals in the nucleus by a nucleoplasmic reticulum. *Nat. Cell Biol.* 5, 440–446. doi: 10.1038/ncb980
- Fiacco, T. A., Agulhon, C., Taves, S. R., Petracz, J., Casper, K. B., Dong, X., et al. (2007). Selective stimulation of astrocyte calcium in situ does not affect neuronal excitatory synaptic activity. *Neuron* 54, 611–626. doi: 10.1016/j.neuron.2007.04.032
- Figueiredo, M., Lane, S., Stout, R. F., Liu, B., Pargura, V., Teschemacher, A. G., et al. (2014). Comparative analysis of optogenetic actuators in cultured astrocytes. *Cell Calcium* 56, 208–214. doi: 10.1016/j.ceca.2014.07.007
- Finch, E., Turner, T., and Goldin, S. (1991). Calcium as a coagonist of inositol 1,4,5-trisphosphate-induced calcium release. *Science* 252, 443–446. doi: 10.1126/science.2017683
- Foley, J., Blustein, T., Lee, S., Erneux, C., Halassa, M. M., and Haydon, P. (2017). Astrocytic IP₃/Ca²⁺ signaling modulates theta rhythm and REM sleep. *Front. Neural Circuits* 11:3. doi: 10.3389/fncir.2017.00003
- Furuichi, T., Yoshikawa, S., Miyawaki, A., Wada, K., Maeda, N., and Mikoshiba, K. (1989). Primary structure and functional expression of the inositol 1,4,5-trisphosphate-binding protein P400. *Nature* 342, 32–38. doi: 10.1038/342032a0
- Futatsugi, A. (2005). IP₃ receptor types 2 and 3 mediate exocrine secretion underlying energy metabolism. *Science* 309, 2232–2234. doi: 10.1126/science.1114110
- Gerasimenko, J. V., Lur, G., Ferdek, P., Sherwood, M. W., Ebisui, E., Tepikin, A. V., et al. (2011). Calmodulin protects against alcohol-induced pancreatic trypsinogen activation elicited via Ca²⁺ release through IP₃ receptors. *Proc. Natl. Acad. Sci. U.S.A.* 108, 5873–5878. doi: 10.1073/pnas.1016534108
- Gerasimenko, J. V., Lur, G., Sherwood, M. W., Ebisui, E., Tepikin, A. V., Mikoshiba, K., et al. (2009). Pancreatic protease activation by alcohol metabolite depends on Ca²⁺ release via acid store IP₃ receptors. *Proc. Natl. Acad. Sci. U.S.A.* 106, 10758–10763. doi: 10.1073/pnas.0904818106
- Gerasimenko, O. V., Gerasimenko, J. V., Belan, P. V., and Petersen, O. H. (1996). Inositol trisphosphate and cyclic ADP-ribose-mediated release of Ca²⁺ from single isolated pancreatic zymogen granules. *Cell* 84, 473–480. doi: 10.1016/s0092-8674(00)81292-1
- Gerasimenko, O. V., Gerasimenko, J. V., Tepikin, A. V., and Petersen, O. H. (1995). ATP-dependent accumulation and inositol trisphosphate- or cyclic ADP-ribose-mediated release of Ca²⁺ from the nuclear envelope. *Cell* 80, 439–444. doi: 10.1016/0092-8674(95)90494-8
- Golovina, V. A., and Blaustein, M. P. (1997). Spatially and functionally distinct Ca²⁺ stores in sarcoplasmic and endoplasmic reticulum. *Science* 275, 1643–1648. doi: 10.1126/science.275.5306.1643
- Golovina, V. A., and Blaustein, M. P. (2000). Unloading and refilling of two classes of spatially resolved endoplasmic reticulum Ca(2+) stores in astrocytes. *Glia* 31, 15–28. doi: 10.1002/(sici)1098-1136(200007)31:1<15::aid-glia20>3.0.co;2-h
- Gómez-Gonzalo, M., Martín-Fernández, M., Martínez-Murillo, R., Mederos, S., Hernández-Vivanco, A., Jamison, S., et al. (2017). Neuron-astrocyte signaling

- is preserved in the aging brain: neuron-astrocyte signaling in aging brain. *Glia* 65, 569–580. doi: 10.1002/glia.23112
- Gómez-Gonzalo, M., Navarrete, M., Perea, G., Covelo, A., Martín-Fernández, M., Shigemoto, R., et al. (2015). Endocannabinoids induce lateral long-term potentiation of transmitter release by stimulation of gliotransmission. *Cereb. Cortex* 25, 3699–3712. doi: 10.1093/cercor/bhu231
- Grosche, J., Matyash, V., Möller, T., Verkhratsky, A., Reichenbach, A., and Kettenmann, H. (1999). Microdomains for neuron-glia interaction: parallel fiber signaling to Bergmann glial cells. *Nat. Neurosci.* 2, 139–143. doi: 10.1038/5692
- Hamada, K., and Mikoshiba, K. (2020). IP 3 receptor plasticity underlying diverse functions. *Annu. Rev. Physiol.* 82, 151–176. doi: 10.1146/annurev-physiol-021119-034433
- Hamada, K., Miyatake, H., Terauchi, A., and Mikoshiba, K. (2017). IP 3 -mediated gating mechanism of the IP 3 receptor revealed by mutagenesis and X-ray crystallography. *Proc. Natl. Acad. Sci. U.S.A.* 114, 4661–4666. doi: 10.1073/pnas.1701420114
- Hamada, T., Niki, T., Ziging, P., Sugiyama, T., Watanabe, S., Mikoshiba, K., et al. (1999). Differential expression patterns of inositol trisphosphate receptor types 1 and 3 in the rat suprachiasmatic nucleus. *Brain Res.* 838, 131–135. doi: 10.1016/s0006-8993(99)01719-9
- Hattori, M., Suzuki, A. Z., Higo, T., Miyauchi, H., Michikawa, T., Nakamura, T., et al. (2004). Distinct roles of inositol 1,4,5-Trisphosphate receptor types 1 and 3 in Ca²⁺ signaling. *J. Biol. Chem.* 279, 11967–11975. doi: 10.1074/jbc.M311456200
- Haustein, M. D., Kracun, S., Lu, X.-H., Shih, T., Jackson-Weaver, O., Tong, X., et al. (2014). Conditions and constraints for astrocyte calcium signaling in the hippocampal mossy fiber pathway. *Neuron* 82, 413–429. doi: 10.1016/j.neuron.2014.02.041
- Hertle, D. N., and Yeckel, M. F. (2007). Distribution of inositol-1,4,5-trisphosphate receptor isoforms and ryanodine receptor isoforms during maturation of the rat hippocampus. *Neuroscience* 150, 625–638. doi: 10.1016/j.neuroscience.2007.09.058
- Hisatsune, C., and Mikoshiba, K. (2017). IP 3 receptor mutations and brain diseases in human and rodents. *J. Neurochem.* 141, 790–807. doi: 10.1111/jnc.13991
- Hisatsune, C., Yasumatsu, K., Takahashi-Iwanaga, H., Ogawa, N., Kuroda, Y., Yoshida, R., et al. (2007). Abnormal taste perception in mice lacking the Type 3 Inositol 1,4,5-trisphosphate receptor. *J. Biol. Chem.* 282, 37225–37231. doi: 10.1074/jbc.M705641200
- Holbro, N., Grunditz, A., and Oertner, T. G. (2009). Differential distribution of endoplasmic reticulum controls metabotropic signaling and plasticity at hippocampal synapses. *Proc. Natl. Acad. Sci. U.S.A.* 106, 15055–15060. doi: 10.1073/pnas.0905110106
- Holtzclaw, L. A., Pandhit, S., Bare, D. J., Mignery, G. A., and Russell, J. T. (2002). Astrocytes in adult rat brain express type 2 inositol 1,4,5-trisphosphate receptors. *Glia* 39, 69–84. doi: 10.1002/glia.10085
- Hur, Y. S., Kim, K. D., Paek, S. H., and Yoo, S. H. (2010). Evidence for the existence of secretory granule (dense-core vesicle)-based inositol 1,4,5-trisphosphate-dependent Ca²⁺ signaling system in astrocytes. *PLoS One* 5:e11973. doi: 10.1371/journal.pone.0011973
- Inaba, T., Hisatsune, C., Sasaki, Y., Ogawa, Y., Ebisui, E., Ogawa, N., et al. (2014). Mice lacking inositol 1,4,5-trisphosphate receptors exhibit dry eye. *PLoS One* 9:e99205. doi: 10.1371/journal.pone.0099205
- Inoue, T., Kato, K., Kohda, K., and Mikoshiba, K. (1998). Type 1 inositol 1,4,5-trisphosphate receptor is required for induction of long-term depression in cerebellar purkinje neurons. *J. Neurosci.* 18, 5366–5373. doi: 10.1523/JNEUROSCI.18-14-05366.1998
- Iwai, M., Tateishi, Y., Hattori, M., Mizutani, A., Nakamura, T., Futatsugi, A., et al. (2005). Molecular cloning of mouse type 2 and type 3 inositol 1,4,5-trisphosphate receptors and identification of a novel type 2 receptor splice variant. *J. Biol. Chem.* 280, 10305–10317. doi: 10.1074/jbc.M413824200
- Iwai, Y., Ozawa, K., Yahagi, K., Mishima, T., Akther, S., Vo, C. T., et al. (2021). Transient astrocytic Gq signaling underlies remote memory enhancement. *Front. Neural. Circuits*, 15:658343 doi: 10.3389/fncir.2021.658343
- Kanemaru, K., Okubo, Y., Hirose, K., and Iino, M. (2007). Regulation of neurite growth by spontaneous Ca²⁺ oscillations in astrocytes. *J. Neurosci.* 27, 8957–8966. doi: 10.1523/JNEUROSCI.2276-07.2007
- Kanemaru, K., Sekiya, H., Xu, M., Satoh, K., Kitajima, N., Yoshida, K., et al. (2014). In vivo visualization of subtle, transient, and local activity of astrocytes using an ultrasensitive Ca²⁺ indicator. *Cell Rep.* 8, 311–318. doi: 10.1016/j.celrep.2014.05.056
- Kirischuk, S., Kettenmann, H., and Verkhratsky, A. (1997). Na⁺/Ca²⁺ exchanger modulates kainate-triggered Ca²⁺ signaling in Bergmann glial cells in situ. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 11, 566–572. doi: 10.1096/fasebj.11.7.9212080
- Klar, J., Hisatsune, C., Baig, S. M., Tariq, M., Johansson, A. C. V., Rasool, M., et al. (2014). Abolished InsP3R2 function inhibits sweat secretion in both humans and mice. *J. Clin. Invest.* 124, 4773–4780. doi: 10.1172/JCI70720
- Konieczny, V., Stefanakis, J. G., Sitsanidis, E. D., Ioannidou, N.-A. T., Papadopoulos, N. V., Fylaktakidou, K. C., et al. (2016). Synthesis of inositol phosphate-based competitive antagonists of inositol 1,4,5-trisphosphate receptors. *Org. Biomol. Chem.* 14, 2504–2514. doi: 10.1039/C5OB02623G
- Konieczny, V., Tovey, S. C., Mataragka, S., Prole, D. L., and Taylor, C. W. (2017). Cyclic AMP recruits a discrete intracellular Ca²⁺ store by unmasking hypersensitive IP 3 receptors. *Cell Rep.* 18, 711–722. doi: 10.1016/j.celrep.2016.12.058
- Korogod, N., Petersen, C. C. H., and Knott, G. W. (2015). Ultrastructural analysis of adult mouse neocortex comparing aldehyde perfusion with cryo fixation. *eLife* 4:e05793. doi: 10.7554/eLife.05793
- Kukkonen, J. P., Lund, P.-E., and Åkerman, K. E. O. (2001). 2-aminoethoxydiphenyl borate reveals heterogeneity in receptor-activated Ca²⁺-discharge and store-operated Ca²⁺-influx. *Cell Calcium* 30, 117–129. doi: 10.1054/ceca.2001.0219
- Kuroda, Y., Hisatsune, C., Nakamura, T., Matsuo, K., and Mikoshiba, K. (2008). Osteoblasts induce Ca²⁺ oscillation-independent NFATc1 activation during osteoclastogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8643–8648. doi: 10.1073/pnas.0800642105
- Letellier, M., Park, Y. K., Chater, T. E., Chipman, P. H., Gautam, S. G., Oshima-Takago, T., et al. (2016). Astrocytes regulate heterogeneity of presynaptic strengths in hippocampal networks. *Proc. Natl. Acad. Sci. U.S.A.* 113, E2685–E2694. doi: 10.1073/pnas.1523717113
- Lur, G., Sherwood, M. W., Ebisui, E., Haynes, L., Feske, S., Sutton, R., et al. (2011). InsP3 receptors and Orai channels in pancreatic acinar cells: co-localization and its consequences. *Biochem. J.* 436, 231–239. doi: 10.1042/BJ20110083
- Mariotti, L., Losi, G., Sessolo, M., Marcon, I., and Carmignoto, G. (2016). The inhibitory neurotransmitter GABA evokes long-lasting Ca(2+) oscillations in cortical astrocytes. *Glia* 64, 363–373. doi: 10.1002/glia.22933
- Martin-Fernandez, M., Jamison, S., Robin, L. M., Zhao, Z., Martin, E. D., Aguilar, J., et al. (2017). Synapse-specific astrocyte gating of amygdala-related behavior. *Nat. Neurosci.* 20, 1540–1548. doi: 10.1038/nn.4649
- Maruyama, T., Kanaji, T., Nakade, S., Kanno, T., and Mikoshiba, K. (1997). 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P3-induced Ca²⁺ release. *J. Biochem. (Tokyo)* 122, 498–505. doi: 10.1093/oxfordjournals.jbchem.a021780
- Masseck, O. A., Spoida, K., Dalkara, D., Maejima, T., Rubelowski, J. M., Wallhorn, L., et al. (2014). Vertebrate cone opsins enable sustained and highly sensitive rapid control of G i/o signaling in anxiety circuitry. *Neuron* 81, 1263–1273. doi: 10.1016/j.neuron.2014.01.041
- Mataragka, S., and Taylor, C. W. (2018). All three IP 3 receptor subtypes generate Ca 2+ puffs, the universal building blocks of IP 3 -evoked Ca 2+ signals. *J. Cell Sci.* 131:jcs220848. doi: 10.1242/jcs.220848
- Matsumoto, M., Nakagawa, T., Inoue, T., Nagata, E., Tanaka, K., Takano, H., et al. (1996). Ataxia and epileptic seizures in mice lacking type 1 inositol 1,4,5-trisphosphate receptor. *Nature* 379, 168–171. doi: 10.1038/379168a0
- Mederos, S., Hernández-Vivanco, A., Ramírez-Franco, J., Martín-Fernández, M., Navarrete, M., Yang, A., et al. (2019). Melanopsin for precise optogenetic activation of astrocyte-neuron networks. *Glia* 67, 915–934. doi: 10.1002/glia.23580
- Michelangeli, F., Mezna, M., Tovey, S., and Sayers, L. G. (1995). Pharmacological modulators of the inositol 1,4,5-trisphosphate receptor. *Neuropharmacology* 34, 1111–1122. doi: 10.1016/0028-3908(95)00053-9
- Mignery, G. A., Newton, C. L., Archer, B. T., and Südhof, T. C. (1990). Structure and expression of the rat inositol 1,4,5-trisphosphate receptor. *J. Biol. Chem.* 265, 12679–12685. doi: 10.1016/s0021-9258(19)38397-8

- Mignery, G. A., Südhof, T. C., Takei, K., and De Camilli, P. (1989). Putative receptor for inositol 1,4,5-trisphosphate similar to ryanodine receptor. *Nature* 342, 192–195. doi: 10.1038/342192a0
- Miyakawa, T. (1999). Encoding of Ca²⁺ signals by differential expression of IP₃ receptor subtypes. *EMBO J.* 18, 1303–1308. doi: 10.1093/emboj/18.5.1303
- Miyazaki, S., Yuzaki, M., Nakada, K., Shirakawa, H., Nakanishi, S., Nakade, S., et al. (1992). Block of Ca²⁺ wave and Ca²⁺ oscillation by antibody to the inositol 1,4,5-trisphosphate receptor in fertilized hamster eggs. *Science* 257, 251–255. doi: 10.1126/science.1321497
- Monai, H., Ohkura, M., Tanaka, M., Oe, Y., Konno, A., Hirai, H., et al. (2016). Calcium imaging reveals glial involvement in transcranial direct current stimulation-induced plasticity in mouse brain. *Nat. Commun.* 7:11100. doi: 10.1038/ncomms11100
- Nakanishi, S., Maeda, N., and Mikoshiba, K. (1991). Immunohistochemical localization of an inositol 1,4,5-trisphosphate receptor, P400, in neural tissue: studies in developing and adult mouse brain. *J. Neurosci.* 11, 2075–2086. doi: 10.1523/JNEUROSCI.11-07-02075.1991
- Nakayama, H., Bodi, I., Maillet, M., DeSantiago, J., Domeier, T. L., Mikoshiba, K., et al. (2010). The IP₃ receptor regulates cardiac hypertrophy in response to select stimuli. *Circ. Res.* 107, 659–666. doi: 10.1161/CIRCRESAHA.110.220038
- Navarrete, M., Cuartero, M. I., Palenzuela, R., Draffin, J. E., Konomi, A., Serra, I., et al. (2019). Astrocytic p38 α MAPK drives NMDA receptor-dependent long-term depression and modulates long-term memory. *Nat. Commun.* 10:2968. doi: 10.1038/s41467-019-10830-9
- Navarrete, M., Perea, G., de Sevilla, D. F., Gómez-Gonzalo, M., Núñez, A., Martín, E. D., et al. (2012). Astrocytes mediate in vivo cholinergic-induced synaptic plasticity. *PLoS Biol.* 10:e1001259. doi: 10.1371/journal.pbio.1001259
- Neher, E. (2008). Details of Ca²⁺ dynamics matter. *J. Physiol.* 586:2031. doi: 10.1113/jphysiol.2008.153080
- Nishiyama, M., Hong, K., Mikoshiba, K., Poo, M. M., and Kato, K. (2000). Calcium stores regulate the polarity and input specificity of synaptic modification. *Nature* 408, 584–588. doi: 10.1038/35046067
- Nizar, K., Uhlirva, H., Tian, P., Saisan, P. A., Cheng, Q., Reznichenko, L., et al. (2013). In vivo stimulus-induced vasodilation occurs without IP₃ receptor activation and may precede astrocytic calcium increase. *J. Neurosci.* 33, 8411–8422. doi: 10.1523/JNEUROSCI.3285-12.2013
- Nucifora, F. C., Sharp, A. H., Milgram, S. L., and Ross, C. A. (1996). Inositol 1,4,5-trisphosphate receptors in endocrine cells: localization and association in hetero- and homotetramers. *Mol. Biol. Cell* 7, 949–960. doi: 10.1091/mbc.7.6.949
- Oh, E., Maejima, T., Liu, C., Deneris, E., and Herlitze, S. (2010). Substitution of 5-HT_{1A} Receptor Signaling by a Light-activated G Protein-coupled Receptor. *J. Biol. Chem.* 285, 30825–30836. doi: 10.1074/jbc.M110.147298
- Okubo, Y., Iino, M., and Hirose, K. (2020). Store-operated Ca²⁺ entry-dependent Ca²⁺ refilling in the endoplasmic reticulum in astrocytes. *Biochem. Biophys. Res. Commun.* 522, 1003–1008. doi: 10.1016/j.bbrc.2019.12.006
- Okubo, Y., Kanemaru, K., Suzuki, J., Kobayashi, K., Hirose, K., and Iino, M. (2019). Inositol 1,4,5-trisphosphate receptor type 2-independent Ca²⁺ release from the endoplasmic reticulum in astrocytes. *Glia* 67, 113–124. doi: 10.1002/glia.23531
- Padmashri, R., Suresh, A., Boska, M. D., and Dunaevsky, A. (2015). Motor-Skill learning is dependent on astrocytic activity. *Neural Plast.* 2015:938023. doi: 10.1155/2015/938023
- Panatier, A., Arizono, M., and Nägerl, U. V. (2014). Dissecting tripartite synapses with STED microscopy. *Philos. Trans. R. Soc. B Biol. Sci.* 369:20130597. doi: 10.1098/rstb.2013.0597
- Panatier, A., Vallée, J., Haber, M., Murai, K. K., Lacaille, J.-C., and Robitaille, R. (2011). Astrocytes are endogenous regulators of basal transmission at central synapses. *Cell* 146, 785–798. doi: 10.1016/j.cell.2011.07.022
- Panda, S. (2005). Illumination of the Melanopsin Signaling Pathway. *Science* 307, 600–604. doi: 10.1126/science.1105121
- Pantazaka, E., and Taylor, C. W. (2011). Differential distribution, clustering, and lateral diffusion of subtypes of the inositol 1,4,5-trisphosphate receptor. *J. Biol. Chem.* 286, 23378–23387. doi: 10.1074/jbc.M111.236372
- Park, K., and Lee, S. J. (2020). Deciphering the star codings: astrocyte manipulation alters mouse behavior. *Exp. Mol. Med.* 52, 1028–1038. doi: 10.1038/s12276-020-0468-z
- Patrushev, I., Gavrilov, N., Turlapov, V., and Semyanov, A. (2013). Subcellular location of astrocytic calcium stores favors extrasynaptic neuron-astrocyte communication. *Cell Calcium* 54, 343–349. doi: 10.1016/j.ceca.2013.08.003
- Perea, G., Gómez, R., Mederos, S., Covelo, A., Ballesteros, J. J., Schlosser, L., et al. (2016). Activity-dependent switch of GABAergic inhibition into glutamatergic excitation in astrocyte-neuron networks. *eLife* 5:e20362. doi: 10.7554/eLife.20362
- Perez-Alvarez, A., Navarrete, M., Covelo, A., Martín, E. D., and Araque, A. (2014). Structural and functional plasticity of astrocyte processes and dendritic spine interactions. *J. Neurosci.* 34, 12738–12744. doi: 10.1523/JNEUROSCI.2401-14.2014
- Petersen, O. H., Gerasimenko, O. V., Gerasimenko, J. V., Mogami, H., and Tepikin, A. V. (1998). The calcium store in the nuclear envelope. *Cell Calcium* 23, 87–90. doi: 10.1016/s0143-4160(98)90106-3
- Petravicz, J., Boyt, K. M., and McCarthy, K. D. (2014). Astrocyte IP₃R2-dependent Ca²⁺ signaling is not a major modulator of neuronal pathways governing behavior. *Front. Behav. Neurosci.* 8:384. doi: 10.3389/fnbeh.2014.00384
- Petravicz, J., Fiacco, T. A., and McCarthy, K. D. (2008). Loss of IP₃ receptor-dependent Ca²⁺ increases in hippocampal astrocytes does not affect baseline CA1 pyramidal neuron synaptic activity. *J. Neurosci.* 28, 4967–4973. doi: 10.1523/JNEUROSCI.5572-07.2008
- Pinton, P., Pozzan, T., and Rizzuto, R. (1998). The Golgi apparatus is an inositol 1,4,5-trisphosphate-sensitive Ca²⁺ store, with functional properties distinct from those of the endoplasmic reticulum. *EMBO J.* 17, 5298–5308. doi: 10.1093/emboj/17.18.5298
- Romanos, J., Thieren, L., and Santello, M. (2019). Diving into new depths of astrocyte signaling. *Nat. Neurosci.* 22, 1749–1750. doi: 10.1038/s41593-019-0513-1
- Rose, C. R., Ziemens, D., and Verkhratsky, A. (2020). On the special role of NCX in astrocytes: translating Na⁺-transients into intracellular Ca²⁺ signals. *Cell Calcium* 86:102154. doi: 10.1016/j.ceca.2019.102154
- Rungta, R. L., Bernier, L.-P., Dissing-Olesen, L., Groten, C. J., LeDue, J. M., Ko, R., et al. (2016). Ca²⁺ transients in astrocyte fine processes occur via Ca²⁺ influx in the adult mouse hippocampus: Ca²⁺ Transients in Astrocyte Fine Processes. *Glia* 64, 2093–2103. doi: 10.1002/glia.23042
- Rusakov, D. A. (2015). Disentangling calcium-driven astrocyte physiology. *Nat. Rev. Neurosci.* 16, 226–233. doi: 10.1038/nrn3878
- Saleem, H., Tovey, S. C., Molinski, T. F., and Taylor, C. W. (2014). Interactions of antagonists with subtypes of inositol 1,4,5-trisphosphate (IP₃) receptor. *Br. J. Pharmacol.* 171, 3298–3312. doi: 10.1111/bph.12685
- Saleem, H., Tovey, S. C., Rahman, T., Riley, A. M., Potter, B. V. L., and Taylor, C. W. (2013a). Stimulation of inositol 1,4,5-trisphosphate (IP₃) receptor subtypes by analogues of IP₃. *PLoS One* 8:e54877. doi: 10.1371/journal.pone.0054877
- Saleem, H., Tovey, S. C., Riley, A. M., Potter, B. V. L., and Taylor, C. W. (2013b). Stimulation of inositol 1,4,5-trisphosphate (IP₃) receptor subtypes by adenophostin A and its analogues. *PLoS One* 8:e58027. doi: 10.1371/journal.pone.0058027
- Sato-Miyaoka, M., Hisatsune, C., Ebisui, E., Ogawa, N., Takahashi-Iwanaga, H., and Mikoshiba, K. (2012). Regulation of hair shedding by the Type 3 IP₃ receptor. *J. Invest. Dermatol.* 132, 2137–2147. doi: 10.1038/jid.2012.141
- Savchenko, L. P., Bard, L., Jensen, T. P., Reynolds, J. P., Kraev, I., Medvedev, N., et al. (2018). Disentangling astroglial physiology with a realistic cell model in silico. *Nat. Commun.* 9:3554. doi: 10.1038/s41467-018-05896-w
- Sharp, A. H., Nucifora, F. C., Blondel, O., Sheppard, C. A., Zhang, C., Snyder, S. H., et al. (1999). Differential cellular expression of isoforms of inositol 1,4,5-trisphosphate receptors in neurons and glia in brain. *J. Comp. Neurol.* 406, 207–220. doi: 10.1002/(sici)1096-9861(19990405)406:2<207::aid-cne6>3.0.co;2-7
- Sherwood, M. W., Arizono, M., Hisatsune, C., Bannai, H., Ebisui, E., Sherwood, J. L., et al. (2017). Astrocytic IP₃ Rs: contribution to Ca²⁺ signalling and hippocampal LTP: astrocytic IP₃ Rs: Ca²⁺ Signalling and LTP. *Glia* 65, 502–513. doi: 10.1002/glia.23107
- Shigetomi, E., Bushong, E. A., Hausteiner, M. D., Tong, X., Jackson-Weaver, O., Kracun, S., et al. (2013a). Imaging calcium microdomains within entire astrocyte territories and endfeet with GCaMPs expressed using adeno-associated viruses. *J. Gen. Physiol.* 141, 633–647. doi: 10.1085/jgp.201210949

- Shigetomi, E., Jackson-Weaver, O., Huckstepp, R. T., O'Dell, T. J., and Khakh, B. S. (2013b). TRPA1 channels are regulators of astrocyte basal calcium levels and long-term potentiation via constitutive D-serine release. *J. Neurosci.* 33, 10143–10153. doi: 10.1523/JNEUROSCI.5779-12.2013
- Shigetomi, E., Tong, X., Kwan, K. Y., Corey, D. P., and Khakh, B. S. (2012). TRPA1 channels regulate astrocyte resting calcium and inhibitory synapse efficacy through GAT-3. *Nat. Neurosci.* 15, 70–80. doi: 10.1038/nn.3000
- Shiratori-Hayashi, M., Yamaguchi, C., Eguchi, K., Shiraishi, Y., Kohno, K., Mikoshiba, K., et al. (2020). Astrocytic STAT3 activation and chronic itch require IP3R1/TRPC-dependent Ca2+ signals in mice. *J. Allergy Clin. Immunol.* 147, 1341–1353. doi: 10.1016/j.jaci.2020.06.039
- Spacek, J., and Harris, K. M. (1998). Three-dimensional organization of cell adhesion junctions at synapses and dendritic spines in area CA1 of the rat hippocampus. *J. Comp. Neurol.* 393, 58–68. doi: 10.1002/(sici)1096-9861(19980330)393:1<58::aid-cne6>3.0.co;2-p
- Spoida, K., Massek, O. A., Deneris, E. S., and Herlitze, S. (2014). Gq/5-HT2c receptor signals activate a local GABAergic inhibitory feedback circuit to modulate serotonergic firing and anxiety in mice. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6479–6484. doi: 10.1073/pnas.1321576111
- Srinivasan, R., Huang, B. S., Venugopal, S., Johnston, A. D., Chai, H., Zeng, H., et al. (2015). Ca2+ signaling in astrocytes from Ip3r2-/- mice in brain slices and during startle responses in vivo. *Nat. Neurosci.* 18, 708–717. doi: 10.1038/nn.4001
- Srinivasan, R., Lu, T.-Y., Chai, H., Xu, J., Huang, B. S., Golshani, P., et al. (2016). New transgenic mouse lines for selectively targeting astrocytes and studying calcium signals in astrocyte processes in situ and in vivo. *Neuron* 92, 1181–1195. doi: 10.1016/j.neuron.2016.11.030
- Südhof, T. C., Newton, C. L., Archer, B. T., Ushkaryov, Y. A., and Mignery, G. A. (1991). Structure of a novel InsP3 receptor. *EMBO J.* 10, 3199–3206. doi: 10.1002/j.1460-2075.1991.tb04882.x
- Sugawara, T., Hisatsune, C., Le, T. D., Hashikawa, T., Hirono, M., Hattori, M., et al. (2013). Type 1 inositol trisphosphate receptor regulates cerebellar circuits by maintaining the spine morphology of purkinje cells in adult mice. *J. Neurosci.* 33, 12186–12196. doi: 10.1523/JNEUROSCI.0545-13.2013
- Sugiyama, T., Furuya, A., Monkawa, T., Yamamoto-Hino, M., Satoh, S., Ohmori, K., et al. (1994). Monoclonal antibodies distinctively recognizing the subtypes of inositol 1,4,5-trisphosphate receptor: application to the studies on inflammatory cells. *FEBS Lett.* 354, 149–154. doi: 10.1016/0014-5793(94)01099-4
- Sul, J.-Y., Orosz, G., Givens, R. S., and Haydon, P. G. (2004). Astrocytic connectivity in the hippocampus. *Neuron Glia Biol.* 1, 3–11. doi: 10.1017/S1740925X04000031
- Takano, T., Wallace, J. T., Baldwin, K. T., Purkey, A. M., Uezu, A., Courtland, J. L., et al. (2020). Chemico-genetic discovery of astrocytic control of inhibition in vivo. *Nature* 588, 296–302. doi: 10.1038/s41586-020-2926-0
- Takata, N., Mishima, T., Hisatsune, C., Nagai, T., Ebisui, E., Mikoshiba, K., et al. (2011). Astrocyte Calcium signaling transforms cholinergic modulation to cortical plasticity in vivo. *J. Neurosci.* 31, 18155–18165. doi: 10.1523/JNEUROSCI.5289-11.2011
- Takata, N., Nagai, T., Ozawa, K., Oe, Y., Mikoshiba, K., and Hirase, H. (2013). Cerebral blood flow modulation by Basal forebrain or whisker stimulation can occur independently of large cytosolic Ca2+ signaling in astrocytes. *PLoS One* 8:e66525. doi: 10.1371/journal.pone.0066525
- Tamamushi, S., Nakamura, T., Inoue, T., Ebisui, E., Sugiura, K., Bannai, H., et al. (2012). Type 2 inositol 1,4,5-trisphosphate receptor is predominantly involved in agonist-induced Ca2+ signaling in Bergmann glia. *Neurosci. Res.* 74, 32–41. doi: 10.1016/j.neures.2012.06.005
- Tamura, A., Yamada, N., Yaguchi, Y., Machida, Y., Mori, I., and Osanai, M. (2014). Both neurons and astrocytes exhibited tetrodotoxin-resistant metabotropic glutamate receptor-dependent spontaneous Slow Ca2+ oscillations in striatum. *PLoS One* 9:e85351. doi: 10.1371/journal.pone.0085351
- Tanaka, M., Shih, P.-Y., Gomi, H., Yoshida, T., Nakai, J., Ando, R., et al. (2013). Astrocytic Ca2+ signals are required for the functional integrity of tripartite synapses. *Mol. Brain* 6:6. doi: 10.1186/1756-6606-6-6
- Tanaka, M., Wang, X., Mikoshiba, K., Hirase, H., and Shinohara, Y. (2017). Rearing-environment-dependent hippocampal local field potential differences in wild-type and inositol trisphosphate receptor type 2 knockout mice. *J. Physiol.* 595, 6557–6568. doi: 10.1113/JP274573
- Tang, F., Lane, S., Korsak, A., Paton, J. F. R., Gourine, A. V., Kasparov, S., et al. (2014). Lactate-mediated glia-neuronal signalling in the mammalian brain. *Nat. Commun.* 5:284. doi: 10.1038/ncomms4284
- Tang, W., Szokol, K., Jensen, V., Enger, R., Trivedi, C. A., Hvalby, O., et al. (2015). Stimulation-evoked Ca2+ signals in astrocytic processes at hippocampal CA3-CA1 synapses of adult mice are modulated by glutamate and ATP. *J. Neurosci.* 35, 3016–3021. doi: 10.1523/JNEUROSCI.3319-14.2015
- Thillaiappan, N. B., Chavda, A. P., Tovey, S. C., Prole, D. L., and Taylor, C. W. (2017). Ca2+ signals initiate at immobile IP3 receptors adjacent to ER-plasma membrane junctions. *Nat. Commun.* 8:1505. doi: 10.1038/s41467-017-01644-8
- Tichy, A.-M., Gerrard, E. J., Sexton, P. M., and Janovjak, H. (2019). Light-activated chimeric GPCRs: limitations and opportunities. *Curr. Opin. Struct. Biol.* 57, 196–203. doi: 10.1016/j.sbi.2019.05.006
- Uchida, K., Aramaki, M., Nakazawa, M., Yamagishi, C., Makino, S., Fukuda, K., et al. (2010). Gene knock-outs of inositol 1,4,5-trisphosphate receptors types 1 and 2 result in perturbation of cardiogenesis. *PLoS One* 5:e12500. doi: 10.1371/journal.pone.0012500
- Uchida, K., Nakazawa, M., Yamagishi, C., Mikoshiba, K., and Yamagishi, H. (2016). Type 1 and 3 inositol trisphosphate receptors are required for extra-embryonic vascular development. *Dev. Biol.* 418, 89–97. doi: 10.1016/j.ydbio.2016.08.007
- Vervoesem, T., Yule, D. I., Bultynck, G., and Parys, J. B. (2015). The type 2 inositol 1,4,5-trisphosphate receptor, emerging functions for an intriguing Ca2+-release channel. *Biochim. Biophys. Acta BBA Mol. Cell Res.* 1853, 1992–2005. doi: 10.1016/j.bbamcr.2014.12.006
- Wang, Q., Kong, Y., Wu, D.-Y., Liu, J.-H., Jie, W., You, Q.-L., et al. (2021). Impaired calcium signaling in astrocytes modulates autism spectrum disorder-like behaviors in mice. *Nat. Commun.* 12:3321. doi: 10.1038/s41467-021-23843-0
- Wang, Y., DelRosso, N. V., Vaidyanathan, T. V., Cahill, M. K., Reitman, M. E., Pittolo, S., et al. (2019). Accurate quantification of astrocyte and neurotransmitter fluorescence dynamics for single-cell and population-level physiology. *Nat. Neurosci.* 22, 1936–1944. doi: 10.1038/s41593-019-0492-2
- Wojcikiewicz, R. J., and He, Y. (1995). Type I, II and III inositol 1,4,5-trisphosphate receptor co-immunoprecipitation as evidence for the existence of heterotetrameric receptor complexes. *Biochem. Biophys. Res. Commun.* 213, 334–341. doi: 10.1006/bbrc.1995.2134
- Xie, Y., Wang, T., Sun, G. Y., and Ding, S. (2010). Specific disruption of astrocytic Ca2+ signaling pathway in vivo by adeno-associated viral transduction. *Neuroscience* 170, 992–1003. doi: 10.1016/j.neuroscience.2010.08.034
- Yamada, N., Makino, Y., Clark, R. A., Pearson, D. W., Mattei, M. G., Guénet, J. L., et al. (1994). Human inositol 1,4,5-trisphosphate type-1 receptor, InsP3R1: structure, function, regulation of expression and chromosomal localization. *Biochem. J.* 302, 781–790. doi: 10.1042/bj3020781
- Yamamoto-Hino, M., Miyawaki, A., Kawano, H., Sugiyama, T., Furuichi, T., Hasegawa, M., et al. (1995). Immunohistochemical study of inositol 1,4,5-trisphosphate receptor type 3 in rat central nervous system. *NeuroReport* 6, 273–276. doi: 10.1097/00001756-199501000-00012
- Yamamoto-Hino, M., Sugiyama, T., Hikichi, K., Mattei, M. G., Hasegawa, K., Sekine, S., et al. (1994). Cloning and characterization of human type 2 and type 3 inositol 1,4,5-trisphosphate receptors. *Recept. Channels* 2, 9–22.
- Yang, L., Qi, Y., and Yang, Y. (2015). Astrocytes control food intake by inhibiting AGRP neuron activity via adenosine A1 receptors. *Cell Rep.* 11, 798–807. doi: 10.1016/j.celrep.2015.04.002
- Yoo, S. H. (2010). Secretory granules in inositol 1,4,5-trisphosphate-dependent Ca2+ signaling in the cytoplasm of neuroendocrine cells. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 24, 653–664. doi: 10.1096/fj.09-132456
- Young, S. Z., Platel, J., Nielsen, J. V., Jensen, N., and Bordey, A. (2010). GABAA increases calcium in subventricular zone astrocyte-like cells through L- and T-type voltage-gated calcium channels. *Front. Cell. Neurosci.* 4:8. doi: 10.3389/fncel.2010.00008

- Yu, X., Taylor, A. M. W., Nagai, J., Golshani, P., Evans, C. J., Coppola, G., et al. (2018). Reducing astrocyte calcium signaling in vivo alters striatal microcircuits and causes repetitive behavior. *Neuron* 99, 1170–1187.e9. doi: 10.1016/j.neuron.2018.08.015
- Zhang, S., Fritz, N., Ibarra, C., and Uhlén, P. (2011). Inositol 1,4,5-trisphosphate receptor subtype-specific regulation of calcium oscillations. *Neurochem. Res.* 36, 1175–1185. doi: 10.1007/s11064-011-0457-7
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O’Keeffe, S., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947. doi: 10.1523/JNEUROSCI.1860-14.2014
- Zhang, Y., Sloan, S. A., Clarke, L. E., Caneda, C., Plaza, C. A., Blumenthal, P. D., et al. (2016). Purification and characterization of progenitor and mature human astrocytes reveals transcriptional and functional differences with mouse. *Neuron* 89, 37–53. doi: 10.1016/j.neuron.2015.11.013

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Sherwood, Arizono, Panatier, Mikoshiba and Olié. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.