HAL
open science

Some geometry of irregular connections on curves (slides of talk at Conference on Gauge Theory and Representation Theory, IAS Princeton, 27/11/2007)

Philip P Boalch

▶ **To cite this version:**

Philip P Boalch. Some geometry of irregular connections on curves (slides of talk at Conference on Gauge Theory and Representation Theory, IAS Princeton, 27/11/2007). Doctoral. United States. 2007, pp.27. hal-03447700

HAL Id: hal-03447700

https://hal.science/hal-03447700v1

Submitted on 24 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Some geometry of irregular connections on curves

P. Boalch

Guiding principle for seeking hyperKahler manifolds – (Hitchin, Seminar Bourbaki '91):–

"Look for hyperKahler structures on manifolds with complex symplectic structures"

Representative collection of examples:–

1. Resolutions of rational double point singularities
2. Coadjoint orbits of complex Lie groups
3. Spaces of representations of a surface group in a complex Lie group

⋮

- Here a 'surface group' is the fundamental group $\pi_1(\Sigma)$ of a compact (Riemann) surface Σ
- If G a complex (reductive) group

$$M = \mathrm{Hom}^{irr}(\pi_1(\Sigma), G)/G$$

has a natural complex symplectic structure (Atiyah-Bott, Goldman, ...)

– independent of complex structure of Σ

"symplectic nature of the fundamental group"

Tannakian categories

- Rough idea: axiomatise properties of categories of representations of groups

- Prove:

  any Tannakian category $\cong$ category of representations of some group

Basically: $\pi_1(\Sigma)$ is the Tannaka group of the (Tannakian) category of vector bundles with holom. connections $/\Sigma$

Define 'wild fundamental group' $w\pi_1(\Sigma)$ to be Tannaka group of category of vector bundles with mero. connections $/\Sigma$

– constructed (in some sense) by Martinet-Ramis '91

Picture

$$\mathrm{Hom}(w\pi_1(\Sigma), G)/G$$

much too big

- Has natural Poisson structure & finite dimensional symplectic leaves
- Describe (& study) generic such leaves directly

Main results

① Atiyah-Bott type definition of symplectic str.
- indept. of cx str. of Σ (& "irreg. types")

"symplectic nature of wild fundamental group"

② (w. O. Biquard) HyperKahler metrics & correspondence w. mero. Higgs bundles

(one direction done earlier by Sabbah)
(& simple poles by Simpson, Nakajima, ...)

Alternative viewpoint

"Hyperkahler metrics on total spaces of algebraically completely integrable systems"

- The Hitchin integrable system on moduli spaces of Higgs bundles was extended to meromorphic Higgs bundles by Bottacin, Markman ~ '94

($\mathbb{P}^1$ case by Beauville, Adams-Harnad-Hurtubise, ...)

Plan for rest of talk

1. Simplest examples

2. Quick general description of the spaces

3. More examples
   - surprising link to Poisson Lie groups
   - appearance of G-braid groups

4. Fourier-Laplace transform (simplest case)

5. Generalised Deligne-Simpson problem

Examples

"Quaternionic Curves" $\left\{ \begin{array}{l} \dim_{\mathbb{H}} = 1 \\ \text{real 4 mfds} \end{array} \right.$

– simplest examples have open pieces

$\cong$ Kronheimer's ALE spaces

cx. sympl. not isometrically



cf. Cherkis–Kapustin, ..., Painlevé...

Quick explicit description of the spaces

Σ genus g, m punctures (simple pole case)

Choose m conjugacy classes $\mathcal{C}_i \subset G$

$$M = \mathrm{Hom}_{\mathcal{C}}(\pi_1(\Sigma), G)/G$$

$$\cong \Big\{ \underset{\sim}{A}, \underset{\sim}{B}, \underset{\sim}{C} \;\Big|\; \underbrace{\prod [A_i, B_i] \prod C_j = 1}_{\text{view as multiplicative moment map condition}},\ C_j \in \mathcal{C}_j \Big\} \Big/ G$$

$$=: \underbrace{\mathbb{D} \circledast \cdots \circledast \mathbb{D}}_{g} \circledast \mathcal{C}_1 \circledast \cdots \circledast \mathcal{C}_m /\!/ G$$

where $\mathbb{D} = G \times G$ with map $\mu : \mathbb{D} \to G$; $(A,B) \mapsto ABA^{-1}B^{-1}$

– can construct cx sympl. structure algebraically this way

(quasi-Hamiltonian geometry – Alekseev–Malkin–Meinrenken)

Generic higher order poles :-

(most polar coeff. regular semisimple $\in H_{reg}$)

Replace conjugacy classes $\mathcal{C} \subset G$ by

$$\mathcal{C} = \left((U_+ \times U_-)^{k-1} \times G \right) / H$$

where $k =$ pole order

$H \subset G$ max. complex torus

$U_\pm \subset B_\pm \subset G$ full unipotent subgroups
$(B_+ \cap B_- = H)$

Choose $t \in H$, then 'moment map' is

$$\mu : \mathcal{C} \to G \; ; \quad [\{s_i^\pm\}, c] \longmapsto c s_1^- s_1^+ s_2^- s_2^+ \cdots t c^{-1}$$

($\mathcal{C}$ is a quasi-Hamiltonian G-space)

Have such 'generalised conjugacy class' $\mathcal{C}$ at each pole, then

$$M \cong \underbrace{\mathbb{D} \circledast \cdots \circledast \mathbb{D}}_{g} \circledast \mathcal{C}_1 \circledast \cdots \circledast \mathcal{C}_m /\!/ G$$

- get explicit (algebraic) symplectic structure
- same as that from extended Atiyah-Bott viewpoint


A
A
A
A

Local irregular Riemann-Hilbert correspondence

(generic, order 2 pole)

- $\Delta \subset \mathbb{C}$ unit disk
- $\mathcal{G} = \{$ holomorphic maps $g : \Delta \to G \}$
- $\mathcal{G}_1 = \{ g \in \mathcal{G} \mid g(0) = 1 \}$
- Fix $A_0 \in \mathfrak{h}_{reg}$

Theorem (Balser–Jurkat–Lutz '79, PB '02)

$$\left\{ \left(\frac{A_0}{z^2} + \frac{B}{z} + \Theta \right) dz \;\middle|\; \begin{array}{l} B \in \mathfrak{g} \\ \Theta : \Delta \to \mathfrak{g} \text{ holom.} \end{array} \right\} \Big/ \mathcal{G}_1$$

$$\cong \quad U_+ \times U_- \times H \ni (\underbrace{S_+, S_-}_{\text{Stokes multipliers}}, \underbrace{\Lambda}_{\pi_{\mathfrak{h}}(B)})$$

- still have action of $H \subset \mathcal{G}$ "compatible framing at 0"
- to get $\mathcal{C}$: include arbitrary framing at point $1 \in \partial\Delta$ & quotient by H fixing value of Λ $(t = e^{2\pi i \Lambda})$

— have more intrinsic viewpoint (isom. here depends on choice of sector at 0 $\Rightarrow$ choice of +ve roots)

$$\left.\begin{array}{l}\text{Roots } R \subset \mathfrak{h}^* \\ A_0 \in \mathfrak{h}_{reg}\end{array}\right\} \quad \langle R, A_0 \rangle \subset \mathbb{C}^*$$



Preferred horizontal section on each sector

Φ : sector $\longrightarrow G$ (asymptotics / k-summation)


S_+
t
S_-

$\rightsquigarrow$ $U_+ \times U_- \times H$ inherits complex Poisson str.

<u>Theorem</u> This Poisson manifold is isomorphic to the standard Poisson Lie group G^* (dual to G).

$$G^* = \{ (b_+, b_-, \Lambda) \in B_+ \times B_- \times H \mid \pi_H(b_+) = \pi_H(b_-^{-1}) = e^{\pi i \Lambda} \}$$

- symplectic leaves got by fixing conjugacy class of
$$b_-^{-1} b_+ = s_- s_+ t \in G$$

- (well) approximated by setting $\Theta = 0$, $B \in \mathfrak{g} \cong \mathfrak{g}^*$

map $\quad \nu_{A_0} : \mathfrak{g}^* \longrightarrow G^*$

$$B \longmapsto \left[\frac{A_0}{z^2} + \frac{B}{z} \right]$$

is Poisson for any $A_0 \in \mathfrak{t}_{reg}$

"dual exponential map"

More hyperKahler spaces (examples)

- Typically $\mathcal{O} \overset{\nu}{\hookrightarrow} \mathcal{L} \subset G^*$

$$\mathcal{O} \subset \mathfrak{g}^*$$

Forget framings: $M \cong \mathcal{L} /\!\!/_{\wedge} H$ (hk)

(partial compactification of $\mathcal{O} /\!\!/_{\wedge} H$)

- 2 order 2 poles on $\mathbb{P}^1$:-

with compatible framings $\tilde{M} \cong \Gamma$ "double symplectic groupoid" – Lu–Weinstein

& have $T^*G \hookrightarrow \Gamma$

Forget framings: $M \cong {}_H\!\|\Gamma /\!\!/_H$ (hk)

(partial compactification of ${}_H\!\|T^*G /\!\!/_H$)

Fourier-Laplace transform

$(GL_n,\ \mathbb{P}^1,\ \text{1 order 2 pole} + \text{1 order 1 pole})$

(Balser-Jurkat-Lutz '81)

$A_0 = \mathrm{diag}(a_1, \dots, a_n)$

$$\begin{tikzcd}
 & B \in \mathfrak{g} \cong \mathfrak{g}^* \arrow[dl, "A_i = E_i B"'] \arrow[dr] & \\
\sum_1^n \frac{A_i}{z-a_i}\,dz \arrow[rr, leftrightarrow, "\text{Fourier-Laplace}"'] \arrow[d, squiggly, "\text{Riemann-Hilbert}"] & & \left(\frac{A_0}{z^2} + \frac{B}{z}\right) dz \arrow[d, squiggly, "\text{Stokes}"'] \\
r_1, \dots, r_n \quad r_i = 1 + e_i \otimes \alpha_i \arrow[rr, leftrightarrow, "\text{Killing-Coxeter}", "(\alpha_i(e_j)) = S_+ t - S_-^{-1}"'] \arrow[dr, "r_n \cdots r_1"'] & & S_+, S_-, t \in U_+ \times U_- \times H \arrow[dl, "S_- S_+ t"] \\
 & G &
\end{tikzcd}$$

- Get isomorphisms of moduli spaces
- Led to search for independent general construction
- "Fourier-Laplace is an isometry" now proved by S. Szabo '05
- original interest was via Dubrovin's work on semi-simple Frobenius manifolds (~ '95)

Deligne-Simpson problem

$(GL_n, \mathbb{P}^1, \text{simple poles})$

"When are the spaces non-empty?"

① Additive version: fix orbits $\mathcal{O}_1, \dots, \mathcal{O}_m \subset \mathfrak{g}$

$\exists$? $A_1, \dots, A_m$ s.t. $A_i \in \mathcal{O}_i$, $\sum A_i = 0$, irreducible

$$\mathcal{M}^* = \mathcal{O}_1 \times \dots \times \mathcal{O}_m /\!/ G \ni \left[\sum \frac{A_i}{z-a_i} dz \right]$$

② Multiplicative version: fix classes $\mathcal{C}_1, \dots, \mathcal{C}_m \subset G$

$\exists$? $C_1, \dots, C_m$ s.t. $C_i \in \mathcal{C}_i$, $\prod C_i = 1$, irred.

$$M = \mathcal{C}_1 \circledast \dots \circledast \mathcal{C}_m /\!/ G$$

Lots of progress made by Crawley-Boevey (solved ①, announced ②) using link to quiver varieties (star-shaped quivers)

(Riemann-Hilbert maps ① into ②, if non-resonant)

More general problem

$(GL_n, \mathbb{P}^1, 1$ pole of order $2 + m$ simple poles$)$

now leading coef. $A_0 \in \mathfrak{h}$, not necessarily $\mathfrak{h}_{reg}$

Let $H(\underset{\sim}{n}) = \mathrm{Stab}(A_0) \cong \prod GL_{n_i} \subset GL_n$

$\underset{\sim}{n} = (n_1, \dots, n_k)$, $\sum n_i = n$ (eigenvalue mults)

$\mathfrak{h}(\underset{\sim}{n}) = \mathrm{Lie}(H(\underset{\sim}{n})) = \ker(\mathrm{ad}_{A_0})$

① Fix adjoint orbits $\mathcal{O}_1, \dots, \mathcal{O}_m \subset \mathfrak{g}$ & $\check{\mathcal{O}} \subset \mathfrak{h}(\underset{\sim}{n})$

$(\check{\mathcal{O}} \cong \check{\mathcal{O}}_1 \times \dots \times \check{\mathcal{O}}_k, \quad \check{\mathcal{O}}_i \subset \mathfrak{gl}_{n_i})$

$\exists$? $A_1, \dots, A_m$ s.t $A_i \in \mathcal{O}_i$, $\pi_{\mathfrak{h}(\underset{\sim}{n})}(\sum A_i) \in \check{\mathcal{O}}$, irred.

$$\mathcal{M}^* = \mathcal{O}_1 \times \dots \times \mathcal{O}_m /\!\!/_{\check{\mathcal{O}}} H(\underset{\sim}{n})$$

Key point: These are also quiver varieties

(A_0 scalar $(k=1)$, or (by Fourier-Laplace) $m=1$ get original problem)

② Choose conjugacy classes $e_1, \dots, e_m \subset G$, $\check{e} \subset H(\underset{\sim}{n})$

$(\check{e} \cong \check{e}_1 \times \dots \times \check{e}_k, \quad \check{e}_i \subset GL_{n_i})$

$\exists ?$ $c_1, \dots, c_m$ s.t $c_i \in e_i$, "$\pi_{H(\underset{\sim}{n})}(\vec{\prod} c_i) \in \check{e}$"

+ irreducible

explicitly: need ① $\vec{\prod} c_i \in \underset{\sim}{n}$-big cell

$U_+(\underset{\sim}{n}) H(\underset{\sim}{n}) U_-(\underset{\sim}{n})$

② $H(\underset{\sim}{n})$ component is in $\check{e}$

(affine) Quiver Varieties (... Nakajima)

Quiver = directed finite graph

$E =$ edges

$I =$ vertices

$$\left.\begin{array}{l} E = \text{edges} \\ I = \text{vertices} \end{array}\right\} \quad E \underset{t}{\overset{h}{\rightrightarrows}} I$$

head

tail

Choose $\mathbb{C}$ vector space V_i $(\forall i \in I)$ $d_i = \dim V_i \Rightarrow \underset{\sim}{d}$

Maps in both directions: $\mathbb{V} = T^* \bigoplus_{e \in E} \mathrm{Hom}(V_{t(e)}, V_{h(e)})$

$\ni (g, g')$

$$G = G(\mathbb{V}) = \prod_{i \in I} GL(V_i) \circlearrowright \mathbb{V}$$

Moment map μ, with components $\mu_i : \mathbb{V} \to \mathrm{End}(V_i)$;

$$\mu_i(g, g') = \sum_{e \mid h(e) = i} g_e g'_e - \sum_{e \mid t(e) = i} g'_e g_e$$

Choose scalar at each node $\underset{\sim}{\lambda} = \{ \lambda_i \in \mathbb{C} \mid i \in I \}$

$$QV(\underset{\sim}{\lambda}, \underset{\sim}{d}) = \mathbb{V} /\!\!/_{\underset{\sim}{\lambda}} G(\mathbb{V})$$

e.g.

- ALE spaces, quiver = affine Dynkin diagram (Kronheimer)

- Closures of adjoint orbits $\mathcal{O} \subset \mathfrak{gl}_n$ (Kraft-Procesi, ...)

$d_i = \operatorname{rank}(A-x_1)\cdots(A-x_i)$ where

$A \in \mathcal{O}$, $x_1, \dots, x_w$ roots of min poly of A

(params $\underset{\sim}{\lambda}$ ~ evals)

Equivalently take graph:

or:



$\sum n_i = n$

and just symplectic quotient by the groups at these nodes

Star shaped quivers

$\mathcal{O}_1, \dots, \mathcal{O}_m \Rightarrow m$ legs 

with residual GL_n action.



$QV \cong \mathcal{O}_1 \times \dots \times \mathcal{O}_m /\!/ GL_n$ (if orbits closed)

Supernova shaped quivers

want $QV = \mathcal{O}_1 \times \cdots \times \mathcal{O}_m /\!\!/_{\check{\theta}} H(\underline{n})$

$$\begin{cases} \sum_1^k n_i = n \\ \check{\theta} = \prod_1^k \check{\theta}_i \\ \check{\theta}_i \subset \mathfrak{gl}_{n_i} \end{cases}$$

$m=2,\ k=3$:-



- residual $H(\underline{n})$ action



(e.g $\hat{A}_3$)

Dual viewpoint (cf. Harnad)



$r = r_1 + r_2$

$\Uparrow$



so $QV \cong \theta_1' \times \theta_2' \times \theta_3' /\!/_{\check{\theta}'} H(\underline{r})$ as well

Claim This duality is the Fourier-Laplace transform

(Generalisation of Gelfand-MacPherson duality)