



HAL
open science

Recent progress in our understanding of peptidoglycan assembly in Firmicutes

Adrien Ducret, Christophe Grangeasse

► **To cite this version:**

Adrien Ducret, Christophe Grangeasse. Recent progress in our understanding of peptidoglycan assembly in Firmicutes. *Current Opinion in Microbiology*, 2021, 60, pp.44-50. 10.1016/j.mib.2021.01.011 . hal-03449233

HAL Id: hal-03449233

<https://hal.science/hal-03449233>

Submitted on 13 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Recent progress in our understanding of peptidoglycan assembly in Firmicutes

Adrien Ducret¹ and Christophe Grangeasse^{1*}

¹ Molecular Microbiology and Structural Biochemistry, CNRS UMR 5086, Université de Lyon, 7 passage du Vercors, 69367 Lyon, France

* Correspondence should be addressed to C.G. E-mail: c.grangeasse@ibcp.fr and Tel. +33 (0)4 72 72 26 88)

Keywords: Peptidoglycan / Bacterial cell morphogenesis / Bacterial cell division / Firmicutes

Abstract

Years of intense research have shown that the assembly of peptidoglycan, the extracellular mesh-like polymer surrounding the bacterial cell, is incredibly complex. It requires a suite of reactions catalyzed by dynamic macromolecular protein complexes whose localization and activity should be finely regulated in space and time. In this review, we focus on the main developments reported over the last 5 years for the assembly of peptidoglycan in Firmicutes, a bacterial phylum that comprises monoderm bacteria and that encompasses well studied bacterial models with different cell shapes and lifestyles.

Introduction

The bacterial cell wall consists mainly of a giant biopolymer named peptidoglycan. The latter surrounds the whole cell and is commonly described as a mesh-like structure consisting of glycan strands that are cross-linked by short peptides [1] (Figure 1). The assembly of this cage-like structure starts at the inner face of the cytoplasmic membrane by a series of reactions involving a set of well conserved proteins generating a lipid-linked disaccharide-pentapeptide precursor named lipid II. This precursor is then flipped across the membrane, polymerized and inserted into the existing peptidoglycan mesh. Polymerization involves both transglycosylation reactions to elongate the glycan stands and transpeptidation reactions that cross-link the pentapeptides of two adjacent glycan strands [1]. These reactions are performed by either the penicillin-binding proteins (PBPs), the L,D-transpeptidases (Ldt), and the SEDS (Shape, Elongation, Division and Sporulation) proteins. The peptidoglycan polymer is also continuously remodeled by a variety of hydrolases, cleaving either the glycan strand, the link between the sugar moieties and the peptide or inside the peptide itself, to promote the insertion of new precursors or to simply mediate cell separation after division [2] (Figure 2).

This general view of peptidoglycan synthesis derives mostly from studies on model organisms such as *Escherichia coli* or *Bacillus subtilis*, but does not reflect the great diversity encountered in bacteria with different morphology or physiology. The most prominent sources of variability are the composition of the peptide, the mode of pentapeptides cross-linking or the sites of peptidoglycan synthesis (polar, septal or lateral) [3] (Figure 1). In this review, we highlight some of the recent breakthroughs that have improved, and even changed, our understanding of some key aspects of peptidoglycan assembly in monoderm bacteria of the Firmicute phylum. Firmicutes are a well-studied group of Gram-positive bacteria and a number of important new discoveries regarding peptidoglycan synthesis have been published

during the last five years. To get more information about other bacterial models and the assembly of other compounds of the cell wall, we suggest the following reviews [4-6]. Furthermore, recent developments allowing the high through peptidoglycan analysis as well as advances in microscopy techniques allowing the investigation of spatial and temporal changes in peptidoglycan synthesis will be covered in this issue in a review by Sara B. Hernández and Felipe Cava [7].

Lipid II transport, polymerization and cross-linking

Once assembled at the inner face of the cytoplasmic membrane by the Mur proteins [1], the lipid II precursor must be translocated across the membrane in order to be incorporated in the peptidoglycan mesh. For many years, it was thought that the translocation step was carried out by the two SEDS proteins, FtsW and RodA, whereas polymerization and cross-linking were performed by two classes of PBPs : the bi-functional class A PBPs that display both a transpeptidase and a transglycosylase activity (aPBPs) and the mono-functional class B PBPs that display only a transpeptidase activity (bPBPs) [1]. This model has however been considerably modified based on the results of several seminal studies, revealing that MurJ mediates the translocation of lipid II [8] and that the two SEDS proteins, FtsW and RodA, possess an unexpected glycosyltransferase activity [9] [10] (Figure 1). Subsequently, it has been shown that the interaction of SEDS proteins with bPBPs was required to stimulate their respective activity [9]. Although initially demonstrated for *B. subtilis*, a series of evidence support that the same model is true for other Firmicutes such as *Staphylococcus aureus* in which the RodA/bPBP3 and FtsW/bPBP1 pairs mediate the sidewall and septal assembly of the peptidoglycan framework, respectively [11], *Streptococcus pneumoniae* in which FtsW works with bPBP2x in septal peptidoglycan assembly [12] or *Listeria monocytogenes* in which RodA and FtsW might work with the bPBP B1 and B2, respectively

[13]. Interestingly, *L. monocytogenes* possesses two FtsW and three RodA homologs that can compensate each other, at least partially, and whose expression levels are modulated upon treatment with antibiotics [13]. In addition to MurJ, a second unrelated protein, Amj, was shown to also flip the lipid II across the membrane in *B. subtilis* [14]. Amj forms a synthetic lethal pair with MurJ and its expression can be induced by cell wall stress. Together, these observations suggest that multiple SEDS and translocases might have redundant and/or specific role to support peptidoglycan synthesis during different environmental conditions/stresses.

As bPBPs work in concert with SEDS proteins to build the peptidoglycan framework, the function of aPBPs had to be revised. A first study has proposed that aPBPs could work outside and independently of the SEDS/bPBP complexes [15]. This view was recently strengthened in *S. pneumoniae* in which evidence was provided that aPBPs are required for the repair and/or maturation of the primary peptidoglycan framework independently of the SEDS/bPBP pairs FtsW/bPBP2x and RodA/bPBP2b [16]. A similar function is also proposed for aPBPs in diderm bacteria like *E. coli* [17]. Altogether, those recent findings support a new model with a division of labor between the SEDS/bPBPs and the aPBPs: while the SEDS/bPBPs complexes are essential for the synthesis of the nascent peptidoglycan, aPBPs would have complementary functions in the maintenance and maturation of the existing framework (Figure 1). The wide diversity in the number of aPBPs with an apparent redundancy among the Firmicutes suggests the existence of a whole repertoire of ancillary functions for aPBPs that have yet to be determined. Understanding how aPBPs operate and how their activity is regulated will be discussed in more detail in the same issue in a review by Manuel Pazos and Waldemar Vollmer [18], and will also be an exciting area of research for the coming years.

Modes of peptidoglycan synthesis

To maintain cell-wall homeostasis, the insertion of new material into the existing peptidoglycan mesh must be regulated in space and time. Different modes of peptidoglycan assembly (polar, lateral, septal) have been identified and characterized by the association between a scaffolding protein and a multiprotein complex responsible for the synthesis and the insertion of new material. In *B. subtilis*, a multiprotein complex called the elongasome which is scaffolded by the actin-like MreB protein, allows lateral insertion of peptidoglycan for cell elongation whereas another complex called the divisome which is scaffolded by the tubulin-like protein FtsZ, assembles the septal peptidoglycan [19]. The contribution of the scaffolding proteins MreB and FtsZ in the peptidoglycan synthesis is still actively debated, but over the last few years numerous insights have considerably improved our understanding of their dynamic and their function during the lateral and septal modes of peptidoglycan synthesis.

Initially thought to form a helical filament distributed along the length of cell, it is now clear that MreB forms small assemblies that move around the rod circumference and promote peptidoglycan synthesis in the radial direction in *B. subtilis* and *E. coli*. Thanks to the emergence of new labelling methods for nascent peptidoglycan and the use of super-resolution imaging, it has also become clear that this circumferential rotation is powered by the active peptidoglycan assembly (reviewed in [20]). While the molecular details regarding the organization of these MreB assemblies are still debated [21], it has become evident that the density of rotating MreB assemblies relies on RodA/bPBP2a but not on the activity of aPBPs [22]. In addition, there is growing evidence that the orientation and/or the motion of MreB filaments depend on the membrane curvature and fluidity, and very interestingly, on the nutrient availability and the growth rate suggesting an intimate link between the cell elongation and the central metabolism [23] [24] [25]. Resource sharing between central

metabolism and cell envelope synthesis will be covered in more detail in this issue in a review by Ankita J. Sachla and John D. Helmann [26].

FtsZ was initially thought to form a continuous structure named the Z-ring at the division septa. However, the advent of super-resolution imaging has also drastically changed our conception of the organization of the Z-ring and revealed that FtsZ forms discontinuous patches/filaments that also move circumferentially around the cell division site [27, 28]. Unlike MreB, the apparent circumferential movement of FtsZ was shown to be independent of cell wall synthesis and rather due to a treadmilling mechanism in which one end of the polymer grows and the other shrinks. While FtsZ treadmilling could guide the progressive insertion of nascent peptidoglycan during cell division, the functional link between FtsZ and peptidoglycan assembly remains more enigmatic and seems to be strain-specific. For instance, it has been shown that the treadmilling of FtsZ filaments drive the motion of bPBP2b in *B. subtilis* whereas the rotation of the bPBP2x:FtsW complex depends on peptidoglycan synthesis and not on FtsZ treadmilling in *S. pneumoniae* [12, 27]. In *S. aureus*, FtsZ treadmilling is proposed to serve for the initial invagination of the membrane while peptidoglycan synthesis occurs at the cell periphery. Upon recruitment of the protein complex DivIB/DivIC/FtsL and the protein MurJ, peptidoglycan assembly is reoriented at the division site and would provide the constriction force independently of FtsZ treadmilling [29]. Further studies will likely tell us whether these observations reflect the specificity of each species in regards to their cell cycle and/or in response to the growth conditions.

Regulation of peptidoglycan hydrolases

To promote the insertion of new precursors into the existing peptidoglycan or to simply mediate cell separation after division, the peptidoglycan polymer needs to be continuously remodeled by a variety of proteins with hydrolytic functions (Figure 2). These

hydrolases also serve other functions, like cell wall recycling or signaling in the host, that are detailed in [30, 31]. For cell division and morphogenesis, these hydrolases include glycosidases that cleave the glycosidic bonds in the glycan strand or amidases and peptidases that cleave the amide bond either between the glycan strand and the first amino acid of the stem peptide or within the stem peptide itself, respectively [2] (Figure 2). As observed for the synthesis machineries, the action of hydrolases must be regulated to maintain cell-wall homeostasis. While many of them are biochemically and structurally well characterized, their regulation remains largely unknown.

Over the last few years, the conserved transmembrane FtsEX complex has emerged as a major regulator of different hydrolases, notably in *S. pneumoniae* and *B. subtilis*. In *S. pneumoniae*, the cell division endopeptidase PcsB is autoinhibited and activated by an ATP-driven conformational change of the FtsEX complex [32] [33]. In *B. subtilis*, a similar observation was made for the endopeptidase CwlO that also requires FtsEX for activation [34]. CwlO however degrades peptidoglycan at the lateral cell wall allowing cell elongation and two new co-factors, SweC and SweD, are required for the FtsEX-CwlO complex to function outside the divisome [35]. In *E. coli*, FtsEX is able to interact with several cell division proteins including FtsZ and FtsA [36]. Based on these findings, it has been proposed that FtsEX could act as a mechano-transmitter coupling cell division with the activation of the endopeptidases required for cell division and morphogenesis. Recently, a new amidase, LytH, contributing to the coordination between cell expansion and cell division was discovered in *S. aureus* [37]. Although independent of FtsEX, LytH is also controlled by a new co-factor, named ActH, to be fully active. Last, the glucosaminidase SagB was recently shown to work in complex with the integral membrane protein SpdC to release the growing glycan strand from the membrane. Indeed, SpdC would bind the lipid carrier and would orient the SagB active site for efficient clivage [38].

These few examples illustrate that we likely only know the tip of the iceberg regarding the regulation of the peptidoglycan hydrolase activities. Interesting directions for future research will be the identification of additional co-factors of cell-wall hydrolases and to investigate the mechanisms by which these hydrolytic enzymes are regulated.

New factors and regulatory mechanisms

An impressive array of studies has led to the characterization of new factors involved in peptidoglycan assembly. Notable examples are the *S. pneumoniae* CozE protein, which directs the activity of the aPBP PBP1a to specific cell wall areas [39]. It has also been proposed that CozE also interacts with MreC and MreD to promote cell elongation. Interestingly, a CozE homolog, named CozEb, was then identified both in *S. pneumoniae* and *S. aureus* [40] [41]. The two proteins interact together and harbor some overlapping functions required for the maintenance of the cell shape. Another factor that is crucial for pneumococcal cell morphogenesis is the RNA-binding protein EloR (also named KhpA) [42] [43]. Inactivation of EloR suppresses the requirement of the essential bPBP PBP2b and SEDS RodA and generates shorter cells indicating that EloR is involved in cell elongation. Interestingly, EloR is phosphorylated by the serine-threonine kinase StkP [42], the key regulator of the pneumococcal cell elongation and division [44]. In absence of EloR phosphorylation, cells length is reduced as for $\Delta eloR$ cells suggesting that EloR is likely inactive. By contrast, the increase in EloR phosphorylation is tolerated only in the presence of suppressor mutations that reduce the functionality of the cell elongation proteins MreC or RodZ [42]. Altogether, it is thus proposed that EloR phosphorylation would stimulate the elongosome activity. EloR phosphorylation echoes another study that has identified a second regulator, specific of the aPBP PBP2a in *S. pneumoniae* [45]. The phosphorylation of this protein, named MacP, by StkP is also required for promoting PBP2a function.

The contribution of serine-threonine-kinases in peptidoglycan assembly is not restricted to *S. pneumoniae* and it has been detected in many Firmicutes, including *Clostridium difficile* [46], *B. subtilis* [47], *Enterococcus faecalis* [48], *L. monocytogenes* [49] or *S. aureus* [50]. Quantitative phosphoproteomic studies also suggest that many more proteins involved in peptidoglycan assembly are likely regulated by phosphorylation and protein-kinases [51]. In line with this claim, the recently identified transglycosylase MltG was found to be phosphorylated in *S. pneumoniae* [51]. MltG was proposed to contribute to pneumococcal cell elongation as its mutation suppresses the requirement of PBP2b, RodA and other proteins from the elongasome [52]. Similarly, the glucosaminidase LytB, that is required for final separation of daughter cell, is controlled by the serine-threonine-kinase StkP [53]. Last, a protein of unknown function was found phosphorylated in both *E. faecalis* (IreB) and *L. monocytogenes* (ReoM). While the function IreB remains unclear [54], ReoM was shown to activate the ClpCP-dependent proteolytic degradation of MurA that is essential for the synthesis of the lipid II [55].

Conclusion and perspectives

The key observations described in this review have changed our view about the biosynthesis of peptidoglycan. However, many other findings, like the knowledge gained about the cell division protein GpsB (reviewed in [56]) have also provided interesting observations and clarifications regarding some aspects of the peptidoglycan assembly. In addition, recent observations provide new directions of research. For instance, peptidoglycan homeostasis is also tightly interconnected with the cell metabolism ([57,58]), regulated by other protein post-translational modification like acetylation ([59]) and the production of second messengers like c-di-AMP ([60,61]) or still dependent on the biosynthesis of other cell wall compounds like teichoic acids ([62]). Together with the recent advances in

peptidoglycan imaging [63,64], all these observations promise further discoveries as exciting as those presented in this review for the next years.

Acknowledgement

We apologize to colleagues whose original work could not be included due to space limitations. This work was supported by the Centre National de la Recherche Scientifique, The Université de Lyon, the Agence National de la Recherche (ANR-18-CE11-0017-02, ANR-19-CE15-0011-01) and the Bettencourt-Schueller Foundation.

References and recommended reading

Papers of particular interest, published within the last two years, have been highlighted as:

* **Special interest**

** **Outstanding interest**

1. Egan AJ, Cleverley RM, Peters K, Lewis RJ, Vollmer W: **Regulation of bacterial cell wall growth.** *FEBS J* 2017, **284**:851-867.
2. Alcorlo M, Martinez-Caballero S, Molina R, Hermoso JA: **Carbohydrate recognition and lysis by bacterial peptidoglycan hydrolases.** *Curr Opin Struct Biol* 2017, **44**:87-100.
3. Shaku M, Ealand C, Matlhabe O, Lala R, Kana BD: **Peptidoglycan biosynthesis and remodeling revisited.** *Adv Appl Microbiol* 2020, **112**:67-103.
4. Randich AM, Brun YV: **Molecular mechanisms for the evolution of bacterial morphologies and growth modes.** *Front Microbiol* 2015, **6**:580.
5. Mueller EA, Levin PA: **Bacterial Cell Wall Quality Control during Environmental Stress.** *mBio* 2020, **11**.
6. van Teeseling MCF, de Pedro MA, Cava F: **Determinants of Bacterial Morphology: From Fundamentals to Possibilities for Antimicrobial Targeting.** *Front Microbiol* 2017, **8**:1264.
7. Hernández SB, Cava F: **New approaches and techniques for bacterial cell wall analysis.** *Curr Opin Microbiol* 2021.
8. Sham LT, Butler EK, Lebar MD, Kahne D, Bernhardt TG, Ruiz N: **Bacterial cell wall. MurJ is the flippase of lipid-linked precursors for peptidoglycan biogenesis.** *Science* 2014, **345**:220-222.
9. Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, Walker S, Kruse AC, Bernhardt TG, Rudner DZ: **SEDS proteins are a widespread family of bacterial cell wall polymerases.** *Nature* 2016, **537**:634-638.
10. Taguchi A, Welsh MA, Marmont LS, Lee W, Sjodt M, Kruse AC, Kahne D, Bernhardt TG, Walker S: **FtsW is a peptidoglycan polymerase that is functional only in complex with its cognate penicillin-binding protein.** *Nat Microbiol* 2019.
- ** This study demonstrates that FtsW polymerizes lipid II and requires its partner bPBP to produce peptidoglycan.
11. Reichmann NT, Tavares AC, Saraiva BM, Jousselin A, Reed P, Pereira AR, Monteiro JM, Sobral RG, VanNieuwenhze MS, Fernandes F, et al.: **SEDS-bPBP pairs direct lateral**

- and septal peptidoglycan synthesis in *Staphylococcus aureus*.** *Nat Microbiol* 2019, **4**:1368-1377.
- ** This study shows that RodA and FtsW together with their cognate bPBPs PBP3 and PBP1 mediate sidewall and septal peptidoglycan incorporation, respectively.
12. Perez AJ, Cesbron Y, Shaw SL, Bazan Villicana J, Tsui HT, Boersma MJ, Ye ZA, Tovpeko Y, Dekker C, Holden S, et al.: **Movement dynamics of divisome proteins and PBP2x:FtsW in cells of *Streptococcus pneumoniae*.** *Proc Natl Acad Sci U S A* 2019, **116**:3211-3220.
- * The authors analyzed FtsZ treadmilling in *S. pneumoniae* and showed that movement of the FtsW/bPBP2x complex depends on peptidoglycan synthesis rather than FtsZ treadmilling.
13. Rismondo J, Halbedel S, Grundling A: **Cell Shape and Antibiotic Resistance Are Maintained by the Activity of Multiple FtsW and RodA Enzymes in *Listeria monocytogenes*.** *mBio* 2019, **10**.
14. Meeske AJ, Sham LT, Kimsey H, Koo BM, Gross CA, Bernhardt TG, Rudner DZ: **MurJ and a novel lipid II flippase are required for cell wall biogenesis in *Bacillus subtilis*.** *Proc Natl Acad Sci U S A* 2015, **112**:6437-6442.
15. Cho H, Wivagg CN, Kapoor M, Barry Z, Rohs PD, Suh H, Marto JA, Garner EC, Bernhardt TG: **Bacterial cell wall biogenesis is mediated by SEDS and PBP polymerase families functioning semi-autonomously.** *Nat Microbiol* 2016:16172.
16. Straume D, Piechowiak KW, Olsen S, Stamsas GA, Berg KH, Kjos M, Heggenhougen MV, Alcorlo M, Hermoso JA, Havarstein LS: **Class A PBPs have a distinct and unique role in the construction of the pneumococcal cell wall.** *Proc Natl Acad Sci U S A* 2020, **117**:6129-6138.
- ** This study demonstrates that aPBPs work independently of the divisome and elongasome activities and suggests that aPBPs mature the peptidoglycan framework.
17. Vigouroux A, Cordier B, Aristov A, Alvarez L, Ozbaykal G, Chaze T, Oldewurtel ER, Matondo M, Cava F, Bikard D, et al.: **Class-A penicillin binding proteins do not contribute to cell shape but repair cell-wall defects.** *Elife* 2020, **9**.
18. Pazos M, Vollmer W: **Regulation and function of class A Penicillin-binding proteins.** *Curr Opin Microbiol* 2021.
19. Egan AJF, Errington J, Vollmer W: **Regulation of peptidoglycan synthesis and remodelling.** *Nat Rev Microbiol* 2020, **18**:446-460.
20. Errington J: **Bacterial morphogenesis and the enigmatic MreB helix.** *Nat Rev Microbiol* 2015, **13**:241-248.

21. Billaudeau C, Yao Z, Cornilleau C, Carballido-Lopez R, Chastanet A: **MreB Forms Subdiffraction Nanofilaments during Active Growth in Bacillus subtilis.** *mBio* 2019, **10**.
22. Dion MF, Kapoor M, Sun Y, Wilson S, Ryan J, Vigouroux A, van Teeffelen S, Oldenbourg R, Garner EC: **Bacillus subtilis cell diameter is determined by the opposing actions of two distinct cell wall synthetic systems.** *Nat Microbiol* 2019, **4**:1294-1305.
23. Billaudeau C, Chastanet A, Yao Z, Cornilleau C, Mirouze N, Fromion V, Carballido-Lopez R: **Contrasting mechanisms of growth in two model rod-shaped bacteria.** *Nat Commun* 2017, **8**:15370.
24. Hussain S, Wivagg CN, Szwedziak P, Wong F, Schaefer K, Izore T, Renner LD, Holmes MJ, Sun Y, Bisson-Filho AW, et al.: **MreB filaments align along greatest principal membrane curvature to orient cell wall synthesis.** *Elife* 2018, **7**.
25. Zielinska A, Savietto A, de Sousa Borges A, Martinez D, Berbon M, Roelofsen JR, Hartman AM, de Boer R, Van der Klei IJ, Hirsch AK, et al.: **Flotillin-mediated membrane fluidity controls peptidoglycan synthesis and MreB movement.** *Elife* 2020, **9**.
- * This study shows that flotillins controls membrane fluidity and that the latter modulates MreB dynamics.
26. Sachla AJ, Helmann JD: **Resource sharing between central metabolism and cell envelope synthesis.** *Curr Opin Microbiol* 2021.
27. Bisson-Filho AW, Hsu YP, Squyres GR, Kuru E, Wu F, Jukes C, Sun Y, Dekker C, Holden S, VanNieuwenhze MS, et al.: **Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division.** *Science* 2017, **355**:739-743.
28. Yang X, Lyu Z, Miguel A, McQuillen R, Huang KC, Xiao J: **GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis.** *Science* 2017, **355**:744-747.
29. Monteiro JM, Pereira AR, Reichmann NT, Saraiva BM, Fernandes PB, Veiga H, Tavares AC, Santos M, Ferreira MT, Macario V, et al.: **Peptidoglycan synthesis drives an FtsZ-treadmilling-independent step of cytokinesis.** *Nature* 2018, **554**:528-532.
30. Mayer C, Kluj RM, Muhleck M, Walter A, Unsleber S, Hottmann I, Borisova M: **Bacteria's different ways to recycle their own cell wall.** *Int J Med Microbiol* 2019, **309**:151326.
31. Bastos PAD, Wheeler R, Boneca IG: **Uptake, recognition and responses to peptidoglycan in the mammalian host.** *FEMS Microbiol Rev* 2020.
32. Rued BE, Alcorlo M, Edmonds KA, Martinez-Caballero S, Straume D, Fu Y, Bruce KE, Wu H, Havarstein LS, Hermoso JA, et al.: **Structure of the Large Extracellular Loop of FtsX and Its Interaction with the Essential Peptidoglycan Hydrolase PcsB in Streptococcus pneumoniae.** *mBio* 2019, **10**.

33. Alcorlo M, Straume D, Lutkenhaus J, Havarstein LS, Hermoso JA: **Structural Characterization of the Essential Cell Division Protein FtsE and Its Interaction with FtsX in *Streptococcus pneumoniae***. *mBio* 2020, **11**.
34. Meisner J, Montero Llopis P, Sham LT, Garner E, Bernhardt TG, Rudner DZ: **FtsEX is required for CwIO peptidoglycan hydrolase activity during cell wall elongation in *Bacillus subtilis***. *Mol Microbiol* 2013, **89**:1069-1083.
35. Brunet YR, Wang X, Rudner DZ: **SweC and SweD are essential co-factors of the FtsEX-CwIO cell wall hydrolase complex in *Bacillus subtilis***. *PLoS Genet* 2019, **15**:e1008296.
- * The authors have identified two co-factors of FtsEX that are required for the function of the peptidoglycan hydrolase CwIO outside the divisome.
36. Pichoff S, Du S, Lutkenhaus J: **Roles of FtsEX in cell division**. *Res Microbiol* 2019, **170**:374-380.
37. Do T, Schaefer K, Santiago AG, Coe KA, Fernandes PB, Kahne D, Pinho MG, Walker S: ***Staphylococcus aureus* cell growth and division are regulated by an amidase that trims peptides from uncrosslinked peptidoglycan**. *Nat Microbiol* 2020, **5**:291-303.
- * Characterization of a new amidase together with its activator. Most recent study illustrating the importance of peptidoglycan hydrolase activities and regulation in peptidoglycan assembly.
38. Schaefer K, Owens TW, Page JE, Santiago M, Kahne D, Walker S: **Structure and reconstitution of a hydrolase complex that may release peptidoglycan from the membrane after polymerization**. *Nat Microbiol* 2021, **6**:34-43.
- * The authors propose that the SagB-spdC complex cleave the nascent peptidoglycan from the membrane to allow its integration into the cell wall.
39. Fenton AK, Mortaji LE, Lau DT, Rudner DZ, Bernhardt TG: **CozE is a member of the MreCD complex that directs cell elongation in *Streptococcus pneumoniae***. *Nat Microbiol* 2016, **2**:16237.
40. Stamsas GA, Myrbraten IS, Straume D, Salehian Z, Veening JW, Havarstein LS, Kjos M: **CozEa and CozEb play overlapping and essential roles in controlling cell division in *Staphylococcus aureus***. *Mol Microbiol* 2018, **109**:615-632.
41. Stamsas GA, Restelli M, Ducret A, Freton C, Garcia PS, Havarstein LS, Straume D, Grangeasse C, Kjos M: **A CozE homolog contributes to cell size homeostasis of *Streptococcus pneumoniae***. *MBio* 2020, in press.
42. Stamsas GA, Straume D, Ruud Winther A, Kjos M, Frantzen CA, Havarstein LS: **Identification of EloR (Spr1851) as a regulator of cell elongation in *Streptococcus pneumoniae***. *Mol Microbiol* 2017.

43. Zheng JJ, Perez AJ, Tsui HT, Massidda O, Winkler ME: **Absence of the KhpA and KhpB (JAG/EloR) RNA-Binding Proteins Suppresses the Requirement for PBP2b by Overproduction of FtsA in Streptococcus pneumoniae D39.** *Mol Microbiol* 2017.
44. Grangeasse C: **Rewiring the Pneumococcal Cell Cycle with Serine/Threonine- and Tyrosine-kinases.** *Trends Microbiol* 2016, **24**:713-724.
45. Fenton AK, Manuse S, Flores-Kim J, Garcia PS, Mercy C, Grangeasse C, Bernhardt TG, Rudner DZ: **Phosphorylation-dependent activation of the cell wall synthase PBP2a in Streptococcus pneumoniae by MacP.** *Proc Natl Acad Sci U S A* 2018, **115**:2812-2817.
46. Cuenot E, Garcia-Garcia T, Douche T, Gorgette O, Courtin P, Denis-Quanquin S, Hoys S, Tremblay YDN, Matondo M, Chapot-Chartier MP, et al.: **The Ser/Thr Kinase PrkC Participates in Cell Wall Homeostasis and Antimicrobial Resistance in Clostridium difficile.** *Infect Immun* 2019, **87**.
47. Pompeo F, Byrne D, Mengin-Lecreulx D, Galinier A: **Dual regulation of activity and intracellular localization of the PASTA kinase PrkC during Bacillus subtilis growth.** *Sci Rep* 2018, **8**:1660.
48. Labbe BD, Hall CL, Kellogg SL, Chen Y, Koehn O, Pickrum AM, Mirza SP, Kristich CJ: **Reciprocal Regulation of PASTA Kinase Signaling by Differential Modification.** *J Bacteriol* 2019, **201**.
49. Pensinger DA, Boldon KM, Chen GY, Vincent WJ, Sherman K, Xiong M, Schaezner AJ, Forster ER, Coers J, Striker R, et al.: **The Listeria monocytogenes PASTA Kinase PrkA and Its Substrate Yvck Are Required for Cell Wall Homeostasis, Metabolism, and Virulence.** *PLoS Pathog* 2016, **12**:e1006001.
50. Hardt P, Engels I, Rausch M, Gajdiss M, Ulm H, Sass P, Ohlsen K, Sahl HG, Bierbaum G, Schneider T, et al.: **The cell wall precursor lipid II acts as a molecular signal for the Ser/Thr kinase PknB of Staphylococcus aureus.** *Int J Med Microbiol* 2017, **307**:1-10.
51. Hirschfeld C, Gomez-Mejia A, Bartel J, Hentschker C, Rohde M, Maass S, Hammerschmidt S, Becher D: **Proteomic Investigation Uncovers Potential Targets and Target Sites of Pneumococcal Serine-Threonine Kinase StkP and Phosphatase PhpP.** *Front Microbiol* 2019, **10**:3101.
52. Tsui HC, Zheng JJ, Magallon AN, Ryan JD, Yunck R, Rued BE, Bernhardt TG, Winkler ME: **Suppression of a deletion mutation in the gene encoding essential PBP2b reveals a new lytic transglycosylase involved in peripheral peptidoglycan synthesis in Streptococcus pneumoniae D39.** *Mol Microbiol* 2016, **100**:1039-1065.
53. Zucchini L, Mercy C, Garcia PS, Cluzel C, Gueguen-Chaignon V, Galisson F, Fretton C, Guiral S, Brochier-Armanet C, Gouet P, et al.: **PASTA repeats of the protein kinase StkP**

- interconnect cell constriction and separation of *Streptococcus pneumoniae*.** *Nat Microbiol* 2018, **3**:197-209.
54. Labbe BD, Kristich CJ: **Growth- and Stress-Induced PASTA Kinase Phosphorylation in *Enterococcus faecalis*.** *J Bacteriol* 2017, **199**.
55. Wamp S, Rutter ZJ, Rismondo J, Jennings CE, Moller L, Lewis RJ, Halbedel S: **PrkA controls peptidoglycan biosynthesis through the essential phosphorylation of ReoM.** *Elife* 2020, **9**.
- * Most recent study illustrating the regulatory function of protein phosphorylation in peptidoglycan assembly. Phosphorylated ReoM controls the degradation of MurA that is required for the first step of peptidoglycan synthesis.
56. Hammond LR, White ML, Eswara PJ: **inverted exclamation markVIVA la DivIVA!** *J Bacteriol* 2019, **201**.
57. Sperber AM, Herman JK: **Metabolism Shapes the Cell.** *J Bacteriol* 2017, **199**.
58. Patel V, Wu Q, Chandrangsu P, Helmann JD: **A metabolic checkpoint protein GlmR is important for diverting carbon into peptidoglycan biosynthesis in *Bacillus subtilis*.** *PLoS Genet* 2018, **14**:e1007689.
59. Carabetta VJ, Greco TM, Tanner AW, Cristea IM, Dubnau D: **Temporal Regulation of the *Bacillus subtilis* Acetylome and Evidence for a Role of MreB Acetylation in Cell Wall Growth.** *mSystems* 2016, **1**.
60. Rismondo J, Gibhardt J, Rosenberg J, Kaefer V, Halbedel S, Commichau FM: **Phenotypes Associated with the Essential Diadenylate Cyclase CdaA and Its Potential Regulator CdaR in the Human Pathogen *Listeria monocytogenes*.** *J Bacteriol* 2016, **198**:416-426.
61. Massa SM, Sharma AD, Siletti C, Tu Z, Godfrey JJ, Gutheil WG, Huynh TN: **C-di-AMP accumulation impairs muropeptide synthesis in *Listeria monocytogenes*.** *J Bacteriol* 2020.
62. Caveney NA, Li FK, Strynadka NC: **Enzyme structures of the bacterial peptidoglycan and wall teichoic acid biogenesis pathways.** *Curr Opin Struct Biol* 2018, **53**:45-58.
63. Radkov AD, Hsu YP, Booher G, VanNieuwenhze MS: **Imaging Bacterial Cell Wall Biosynthesis.** *Annu Rev Biochem* 2018, **87**:991-1014.
64. Pasquina-Lemonche L, Burns J, Turner RD, Kumar S, Tank R, Mullin N, Wilson JS, Chakrabarti B, Bullough PA, Foster SJ, et al.: **The architecture of the Gram-positive bacterial cell wall.** *Nature* 2020, **582**:294-297.
- ** The authors applied atomic force microscopy to show that the peptidoglycan layer could be composed of two different zones: an outer-surface composed of loose and disordered thick glycan strands and a dense inner-surface with thinner glycan strands.

Figure Legends

Figure 1: Composition and diversity of the peptidoglycan polymer

Glycan strands are made of alternating β -1,4-linked N-acetylmuramic acid (MurNac) and N-acetyl-glucosamine (GlcNac) cross-linked by peptide links. The assembly of this structure starts at the inner face of the cytoplasmic membrane by a series of reactions involving a set of well conserved proteins (MurABCDEFGL and MraY) generating a lipid-linked disaccharide-pentapeptide precursor named lipid II. The composition of the stem peptide could vary between species. For instance, the third amino acid could be a meso-DAP (meso-diaminopimelic acid) in proteobacteria, bacilli and mycobacteria or a L-Lys in many Firmicutes. The fifth amino acid could be a D-Ala in most of Firmicutes or a D-Lac in *Lactobacillus plantarum* (a). The lipid II is anchored to the membrane with a undecaprenol phosphate (C55-P). Lipid II is then flipped across the cytoplasmic membrane by MurJ and is used as the substrate for the polymerization of the nascent peptidoglycan chains by glycosyltransferases (GT: SEDS protein & class A PBPs) and transpeptidases (TP: class A & B PBPs). Transpeptidation reactions occur between the third amino acid (meso-DAP or L-Lys) of the acceptor stem and either the fourth amino acid (b and c) (DD-transpeptidation: 3-4 link) or the third amino acid (g) (LD-transpeptidation: 3-3 link) of the donor stem to form a dimer. Cross-linking can occur with or between two dimers producing cross-linked trimers (c) and tetramers. In some species the transpeptidation reactions can occur between the third amino acid (meso-DAP or L-Lys) of the acceptor stem and a bridge structure which is attached to the L-Lys of the third position of the stem. The composition and the length of the bridge structure could vary between species. For instance, the bridge may contain one amino acid in *Enterococcus faecium* (d) (D-Asp or D-Asn), one or two amino acid(s) in *S.*

pneumoniae (e) (L-Ala and L-Ala or L-Ser) or up to 5 amino acids in *S. aureus* (f) (Gly or Ser). The MurNac and the GlcNac can be O-acetylated and N-deacetylated, respectively.

Figure 2: Hydrolases required for peptidoglycan remodeling

The peptidoglycan structure is continuously remodeled by a variety of proteins with hydrolytic functions. For instance, peptides are trimmed by either DD-carboxypeptidases or LD-carboxypeptidases which cleave the bond respectively between the fourth D-Ala and the fifth D-Ala (or D-Lac) amino acid and between the third meso-DAP (or L-Lys) and the fourth D-Ala amino acid of the stem peptide. Crosslinks are cleaved by DD-endopeptidases or LD-endopeptidases while the peptide is removed from the glycan chain by amidases. Glycosydases cleave the backbone of glycan and consist of N-acetylglucosaminidases which hydrolyses N-acetyl-D-glucosamine (GlcNAc) residues from adjacent sugar residues and N-acetylmuramidases which cleaves the β 1-4 glycosidic bond between N-Acetylmuramic acid (MurNAc) and GlcNAc.





