



HAL
open science

Moment polytopes in real symplectic geometry II: applications to singular value inequalities

Paul-Emile Paradan

► **To cite this version:**

Paul-Emile Paradan. Moment polytopes in real symplectic geometry II: applications to singular value inequalities. 2021. hal-03450349

HAL Id: hal-03450349

<https://hal.science/hal-03450349>

Preprint submitted on 25 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Moment polytopes in real symplectic geometry II : applications to singular value inequalities

Paul-Emile Paradan*

November 25, 2021

Abstract

In this work, we study some convex cones associated to isotropic representations of symmetric spaces. We explain the inequalities that describe them by means of cohomological conditions. In particular, we study the singular Horn cone which is the counterpart of the classical Horn cone, where the eigenvalues of Hermitian square matrices are replaced by the singular values of rectangular matrices.

Contents

1	Introduction	1
1.1	Kähler-Hamiltonian manifold with involution	3
1.2	Description of the main result	6
1.3	Bruhat decompositions	8
1.4	Comparison with a result of Kapovich-Leeb-Millson	9
1.5	Levi-movability	10
1.6	Singular Horn cone	13
2	Parametrization of facets of real Kirwan polytopes	16
2.1	Admissible elements	17
2.2	Balanced Ressayre's pair	17
2.3	Parametrization of the facets	19
3	Proof of the main theorem	19
3.1	Admissible elements	20
3.2	Balanced Ressayre's pair : infinitesimal conditions	22
3.3	Cohomological conditions	23
3.4	Conclusion	24

*IMAG, Univ Montpellier, CNRS, email : paul-emile.paradan@umontpellier.fr

4	Singular value inequalities for matrix sums	25
4.1	Multiplicative formulas in $H^*(\mathbb{F}(r, n - r; n), \mathbb{Z})$	26
4.2	Inequalities of Weyl, Lidskii and Thompson	27
4.3	Horn conditions	30
4.4	Examples : Singular($p, 1$), Singular($p, 2$) and Singular($3, 3$)	32
4.5	The convex cone Singular(∞, q)	35
4.6	Proof of Theorem ??	38

1 Introduction

This paper is concerned with convexity properties associated to isotropy representations of symmetric spaces. Let G/K be a Riemannian symmetric space of the non-compact type and let $\mathfrak{p} := \mathbf{T}_e G/K$ be the isotropy representation of the compact Lie group K (that we suppose connected). The K -orbits in \mathfrak{p} are parametrized by a closed cone \mathfrak{a}_+ contained in a maximal abelian subspace $\mathfrak{a} \subset \mathfrak{p}$.

Suppose that $G/K \hookrightarrow \tilde{G}/\tilde{K}$ is an embedding of Riemannian symmetric spaces of the non-compact type. Thus K is a closed subgroup of \tilde{K} and we have an equivariant orthogonal projection $\pi : \tilde{\mathfrak{p}} \rightarrow \mathfrak{p}$. The main purpose of this article is the description of the following convex cone

$$\Pi_{\mathfrak{p}}(\tilde{K}, K) := \left\{ (\tilde{\xi}, \xi) \in \tilde{\mathfrak{a}}_+ \times \mathfrak{a}_+; K\xi \subset \pi(\tilde{K}\tilde{\xi}) \right\}.$$

An efficient way to show that $\Pi_{\mathfrak{p}}(\tilde{K}, K)$ is convex is to use the classical duality between the symmetric spaces of the non-compact type with those of the compact type. In our study, we suppose that $G \subset \tilde{G}$ are linear. In this context, there exists complex reductive groups $U_{\mathbb{C}} \subset \tilde{U}_{\mathbb{C}}$ equipped with an anti-linear involution σ such that the connected component of the fixed point subgroups $(U_{\mathbb{C}})^{\sigma} \subset (\tilde{U}_{\mathbb{C}})^{\sigma}$ and $U^{\sigma} \subset \tilde{U}^{\sigma}$ are respectively $G \subset \tilde{G}$ and $K \subset \tilde{K}$ (see §1.1).

We consider maximal torus $T \subset U$ and $\tilde{T} \subset \tilde{U}$, that are invariant under σ , and such that the subspaces $\mathfrak{t}^{-\sigma}$ and $\tilde{\mathfrak{t}}^{-\sigma}$ are of maximal dimension. Then we can choose Weyl chambers \mathfrak{t}_+ (resp. $\tilde{\mathfrak{t}}_+$) such that $\mathfrak{t}^{-\sigma} \cap \mathfrak{t}_+$ (resp. $\tilde{\mathfrak{t}}^{-\sigma} \cap \tilde{\mathfrak{t}}_+$) parametrizes the K -orbits in $\mathfrak{u}^{-\sigma}$ (resp. the \tilde{K} -orbits in $\tilde{\mathfrak{u}}^{-\sigma}$).

The convex cone

$$\Pi(\tilde{U}, U) := \left\{ (\tilde{\xi}, \xi) \in \tilde{\mathfrak{t}}_+ \times \mathfrak{t}_+; U\xi \subset \pi(\tilde{U}\tilde{\xi}) \right\}.$$

has been the subject of numerous studies in recent decades. Let us mention the works of Horn [11], Klyachko [14], Belkale [1] and Knutson-Tao [15], when $U = U(n)$ and $\tilde{U} = (U(n))^s$. The case $\tilde{U} = (U)^s$ was studied by Belkale-Kumar [2] and Kapovich-Leeb-Millson [12], and the general setting $U \subset \tilde{U}$ was considered by Berenstein-Sjamaar [3] and Ressayre [21, 23].

In particular, Ressayre obtained in [23] a complete description of the general¹ faces of $\Pi(\tilde{U}, U)$. When $\tilde{U} = U^s$, he shows that the general faces of $\Pi(U^s, U)$ are parameterized by the set of $(\mathbb{P}, \mathfrak{X}_{w_1}^{\mathbb{P}}, \dots, \mathfrak{X}_{w_{s+1}}^{\mathbb{P}})$ where \mathbb{P} is a standard parabolic subgroup of $U_{\mathbb{C}}$ and the $\mathfrak{X}_{w_k}^{\mathbb{P}}$'s are Schubert varieties of $U_{\mathbb{C}}/\mathbb{P}$ such that $[\mathfrak{X}_{w_1}^{\mathbb{P}}] \odot_0 \cdots \odot_0 [\mathfrak{X}_{w_{s+1}}^{\mathbb{P}}] = [pt]$. Here \odot_0 is the Belkale-Kumar's product on the cohomology groups $H^*(U_{\mathbb{C}}/\mathbb{P}, \mathbb{Z})$.

The link between $\Pi(\tilde{U}, U)$ and $\Pi_{\mathfrak{p}}(\tilde{K}, K)$ is given by a theorem of O'Shea and Sjamaar [19] that we recall in the next Section: it says that we have a natural identification

$$\Pi_{\mathfrak{p}}(\tilde{K}, K) \simeq \Pi(\tilde{U}, U) \cap \tilde{\mathfrak{t}}^{-\sigma} \times \mathfrak{t}^{-\sigma}.$$

From the above isomorphism, the convex cone $\Pi_{\mathfrak{p}}(\tilde{K}, K)$ is completely determined since the equations of the cone $\Pi(\tilde{U}, U)$ are known. But by doing so, the inequalities describing $\Pi_{\mathfrak{p}}(\tilde{K}, K)$ are in general highly overdetermined.

Our main contribution is a description of a smaller list of inequalities describing $\Pi_{\mathfrak{p}}(\tilde{K}, K)$. Let us give an overview of our result when $\tilde{G} = G^s$ and $\tilde{K} = K^s$ with $s \geq 2$. A more precise version is explained in Sections 1.2 and 1.5.

Theorem 1.1 *The convex cone $\Pi_{\mathfrak{p}}(K^s, K)$ can be described by a system of inequalities parameterized by the set of $(\mathbb{P}, \mathfrak{X}_{w_1}^{\mathbb{P}}, \dots, \mathfrak{X}_{w_{s+1}}^{\mathbb{P}})$ where*

- \mathbb{P} is a σ -invariant parabolic subgroup of $U_{\mathbb{C}}$,
- \mathbb{P} is maximal among the σ -invariant parabolic subgroups of $U_{\mathbb{C}}$,
- the $\mathfrak{X}_{w_k}^{\mathbb{P}} = \overline{\mathbb{B}[w_k]}$ are Schubert varieties such that $\mathbb{B}[w_k] \cap (U_{\mathbb{C}}/\mathbb{P})^{\sigma} \neq \emptyset$,
- the relation $[\mathfrak{X}_{w_1}^{\mathbb{P}}] \odot_0 \cdots \odot_0 [\mathfrak{X}_{w_{s+1}}^{\mathbb{P}}] = [pt]$ holds in $H^*(U_{\mathbb{C}}/\mathbb{P}, \mathbb{Z})$.

A description of $\Pi_{\mathfrak{p}}(K^s, K)$ was also obtained by Kapovich-Leeb-Millson in [12]. We explain in §1.4 why our result is more accurate than theirs.

When working with the reductive group $G = U(p, q)$ and its maximal compact subgroup $K = U(p) \times U(q)$, the corresponding convex cone $\Pi_{\mathfrak{p}}(K^2, K)$ is the singular Horn cone which we denote by $\text{Singular}(p, q)$. In §1.6, we describe $\text{Singular}(p, q)$ by means of the inequalities of Theorem 1.1.

Acknowledgements

I would like to thank Michèle Vergne and Velleda Baldoni for their precious help in the calculation of some Littlewood-Richardson coefficients.

1.1 Kähler-Hamiltonian manifold with involution

The purpose of this section is to explain the theorem of O'Shea and Sjamaar [19] that we mentioned above.

¹A face of $\Pi(\tilde{U}, U)$ is called *general* when it intersects the interior of the cone $\tilde{\mathfrak{t}}_+ \times \mathfrak{t}_+$.

Duality

Let $\tilde{G} \subset \mathrm{GL}(n, \mathbb{R})$ be a connected linear real reductive subgroup with maximal compact subgroup $\tilde{K} \subset \mathrm{SO}(n, \mathbb{R})$. Let $\tilde{\mathfrak{p}}$ be the intersection of the Lie algebra $\tilde{\mathfrak{g}}$ of \tilde{G} with the subspace of $n \times n$ symmetric matrices. Let $\tilde{U} \subset \mathrm{GL}(n, \mathbb{C}) \subset \mathrm{GL}(2n, \mathbb{R})$ be the compact connected Lie group with Lie algebra $\tilde{\mathfrak{u}} = \tilde{\mathfrak{k}} \oplus i\tilde{\mathfrak{p}}$. Let us consider the involution $\sigma(A) = JAJ^{-1}$ of $\mathrm{GL}(2n, \mathbb{R})$ defined by $J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$. We see that σ commutes with the Cartan involution, leaves stable the subgroups \tilde{U} , and that the morphism $j : \mathrm{GL}(n, \mathbb{R}) \rightarrow \mathrm{GL}(2n, \mathbb{R})$, $j(g) = \mathrm{Diag}(g, g)$ defines an isomorphism between \tilde{K} and the connected component of \tilde{U}^σ .

The compact Lie groups $\tilde{U} \subset U(n) \subset \mathrm{SO}(2n, \mathbb{R})$ admit a complexification $\tilde{U}_{\mathbb{C}} \subset \mathrm{GL}(2n, \mathbb{C})$ (see §III.8 in [7]). We extend the involution σ on $\mathrm{GL}(2n, \mathbb{R})$ to the conjugate linear involution $\sigma(g) = J\bar{g}J^{-1}$ on $\mathrm{GL}(2n, \mathbb{C})$. We see then that the morphism $j : \mathrm{GL}(n, \mathbb{R}) \rightarrow \mathrm{GL}(2n, \mathbb{C})$ defines an isomorphism between \tilde{G} and the connected component of the subgroup $\tilde{U}_{\mathbb{C}}^\sigma$ fixed by the involution.

Let $\tilde{\mathfrak{t}} \subset \tilde{\mathfrak{u}}$ be the Lie algebra of a maximal torus $\tilde{T} \subset \tilde{U}$ stable under the involution and such that $\tilde{\mathfrak{a}} := \frac{1}{i}\tilde{\mathfrak{t}}^{-\sigma} \subset \tilde{\mathfrak{p}}$ is of maximal dimension. One can choose the Weyl chamber $\tilde{\mathfrak{t}}_+ \subset \tilde{\mathfrak{t}}$ appropriately so that the restricted Weyl chamber $\tilde{\mathfrak{a}}_+ = \frac{1}{i}\tilde{\mathfrak{t}}_+ \cap \tilde{\mathfrak{t}}^{-\sigma}$ parameters the \tilde{K} -orbits on $\tilde{\mathfrak{p}} = \frac{1}{i}\tilde{\mathfrak{u}}^{-\sigma}$ (see the Appendix in [19]).

If we work with a connected linear real reductive subgroup $G \subset \tilde{G}$ the same construction holds. We have a compact connected subgroup $U \subset \tilde{U}$ with Lie algebra $\mathfrak{u} = \mathfrak{k} \oplus i\mathfrak{p}$. Its complexification $U_{\mathbb{C}} \subset \tilde{U}_{\mathbb{C}}$ is stable under the conjugate linear involution σ and the real reductive group G admits a natural identification with the connected component of the subgroup $U_{\mathbb{C}}^\sigma$. The maximal torus $T \subset U$ is taken invariant under σ , contained in \tilde{T} , and such that $\mathfrak{a} = \frac{1}{i}\mathfrak{t}^{-\sigma} \subset \tilde{\mathfrak{a}}$ is of maximal dimension. The Lie algebras \mathfrak{u} and $\tilde{\mathfrak{u}}$ belong to the vector space $\mathfrak{gl}(2n, \mathbb{R})$ that is equipped with the canonical Euclidean norm $\|X\|^2 = \sqrt{\mathrm{Tr}(XX^t)}$. We denote by $\pi : \tilde{\mathfrak{u}} \rightarrow \mathfrak{u}$ the orthogonal projection.

O'Shea-Sjamaar's Theorem

We consider the following geometrical setting : the group $\tilde{U}_{\mathbb{C}}$ viewed as a complex manifold equipped with the following action of $\tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}} : (\tilde{g}, g) \cdot x = \tilde{g}xg^{-1}$.

We identify the complex reductive group $\tilde{U}_{\mathbb{C}}$ with the tangent bundle $\mathbf{T}\tilde{U}$ through the isomorphism $\varphi_2 : \mathbf{T}\tilde{U} \rightarrow \tilde{U}_{\mathbb{C}}, \varphi_2(\tilde{a}, \tilde{X}) = \tilde{a}e^{i\tilde{X}}$ given by the Cartan decomposition. If we use φ_2 to transport the complex structure of $\tilde{U}_{\mathbb{C}}$ we get a complex structure \mathbb{J} on $\mathbf{T}\tilde{U}$.

Since $\mathbf{T}^*\tilde{U} \simeq \tilde{U} \times \tilde{\mathfrak{u}}^*$ and $\mathbf{T}\tilde{U} \simeq \tilde{U} \times \tilde{\mathfrak{u}}$ through left translations, the isomorphism $\varphi_1 : \mathbf{T}\tilde{U} \simeq \mathbf{T}^*\tilde{U}$ is obtain by means of the identification $\tilde{\mathfrak{u}} \simeq \tilde{\mathfrak{u}}^*$ given by the invariant scalar product. If we use φ_1 to transport the canonical symplectic form of $\mathbf{T}^*\tilde{U}$, the resulting symplectic structure Ω on $\mathbf{T}\tilde{U}$ is invariant relatively to the $\tilde{U} \times U$ -action, and is compatible with the complex structure \mathbb{J} (see [10], §3). The moment map relative to the

$\tilde{U} \times U$ -action on $(\mathbf{T}\tilde{U}, \Omega)$ is a proper map $\Phi = \Phi_{\tilde{U}} \oplus \Phi_U : \mathbf{T}\tilde{U} \rightarrow \tilde{\mathfrak{u}} \times \mathfrak{u}$ defined by

$$(1) \quad \Phi_{\tilde{U}}(\tilde{a}, \tilde{X}) = -\tilde{a}\tilde{X}, \quad \Phi_U(\tilde{a}, \tilde{X}) = \pi(\tilde{X}),$$

The conjugate linear involution σ on $\tilde{U}_{\mathbb{C}}$ defines (through φ_2) an involution $\tau : \mathbf{T}\tilde{U} \rightarrow \mathbf{T}\tilde{U}$ defined by $\tau(\tilde{a}, \tilde{X}) = (\sigma(\tilde{a}), -\sigma(\tilde{X}))$.

The $\tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}}$ -manifold $\mathbf{T}\tilde{U}$ equipped with the data $(\Omega, \mathbb{J}, \Phi, \tau, \sigma)$ is a Kähler-Hamiltonian manifold with involution :

- $(\mathbf{T}\tilde{U}, \Omega, \mathbb{J})$ is a Kähler manifold,
- the $\tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}}$ -action preserves the complex structure \mathbb{J} ,
- the $\tilde{U} \times U$ -action is Hamiltonian with proper moment map Φ ,
- $\tau^*(\Omega) = -\Omega$ and $\tau^*(\mathbb{J}) = -\mathbb{J}$,
- $\tau(g \cdot x) = \sigma(g) \cdot \tau(x)$, and $\Phi(\tau(x)) = -\sigma(\Phi(x))$, $\forall (g, x) \in \tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}} \times \mathbf{T}\tilde{U}$.

We finish the section by recalling the O'Shea-Sjamaar's Theorem [19]. Let $\Delta_{\tilde{U} \times U}(\mathbf{T}\tilde{U}) \subset \tilde{\mathfrak{t}}_+ \times \mathfrak{t}_+$ be the Kirwan polytope associated to Φ . Equations (1) show that

$$\Delta_{\tilde{U} \times U}(\mathbf{T}\tilde{U}) = \left\{ (\tilde{X}, X) \in \tilde{\mathfrak{t}}_+ \times \mathfrak{t}_+ \mid -UX \subset \pi(\tilde{U}\tilde{X}) \right\}.$$

We consider now the submanifold $(\mathbf{T}\tilde{U})^\tau$ fixed by τ . Let \mathcal{Z} be the connected component of $(\mathbf{T}\tilde{U})^\tau$ containing $(e, 0)$: through the diffeomorphism $\varphi_2 : \mathbf{T}\tilde{U} \simeq \tilde{U}_{\mathbb{C}}$, we see that $\mathcal{Z} \simeq \tilde{G}$. The restriction of the moment Φ to \mathcal{Z} takes value in $\tilde{\mathfrak{u}}^{-\sigma} \times \mathfrak{u}^{-\sigma}$. If we use the canonical isomorphism $(\tilde{X}, X) \in \tilde{\mathfrak{p}} \times \mathfrak{p} \rightarrow i(\tilde{X}, X) \in \tilde{\mathfrak{u}}^{-\sigma} \times \mathfrak{u}^{-\sigma}$, we obtain a gradient map

$$\begin{aligned} \Psi : \tilde{G} \simeq \tilde{K} \times \tilde{\mathfrak{p}} &\longrightarrow \tilde{\mathfrak{p}} \times \mathfrak{p} \\ (\tilde{k}, \tilde{X}) &\longmapsto (-\tilde{k}\tilde{X}, \pi(\tilde{X})). \end{aligned}$$

The corresponding Kirwan polytope is $\Delta_{\tilde{K} \times K}(\tilde{G}) = \{(\tilde{\xi}, \xi) \in \tilde{\mathfrak{a}}_+ \times \mathfrak{a}_+; -K\xi \subset \pi(\tilde{K}\tilde{\xi})\}$.

Theorem 1.2 (O'Shea-Sjamaar) *Through the isomorphism $(\tilde{X}, X) \mapsto i(\tilde{X}, X)$, $\tilde{\mathfrak{a}}_+ \times \mathfrak{a}_+ \rightarrow \tilde{\mathfrak{t}}_+ \times \mathfrak{t}_+ \cap \tilde{\mathfrak{t}}^{-\sigma} \times \mathfrak{t}^{-\sigma}$ we have an identification*

$$\Delta_{\tilde{K} \times K}(\tilde{G}) \simeq \Delta_{\tilde{U} \times U}(\mathbf{T}\tilde{U}) \cap \tilde{\mathfrak{t}}^{-\sigma} \times \mathfrak{t}^{-\sigma}.$$

If w_0 denotes the longest element of the Weyl group $W = N_U(T)/T$, we see that the cones $\Delta_{\tilde{U} \times U}(\mathbf{T}\tilde{U})$ and $\Pi(\tilde{U}, U)$ are related as follows: $(\tilde{X}, X) \in \Delta_{\tilde{U} \times U}(\mathbf{T}\tilde{U}) \iff (\tilde{X}, -w_0X) \in \Pi(\tilde{U}, U)$. The element w_0 is σ -invariant and satisfies $w_0(\mathfrak{a}_+) = -\mathfrak{a}_+$ (see [19], Lemma 5.4). Hence we obtain the same relations : $(\tilde{X}, X) \in \Delta_{\tilde{K} \times K}(\tilde{G}) \iff (\tilde{X}, -w_0X) \in \Pi_{\mathfrak{p}}(\tilde{K}, K)$.

Finally, the O'Shea-Sjamaar's Theorem tells us that $\Pi_{\mathfrak{p}}(\tilde{K}, K) \simeq \Pi(\tilde{U}, U) \cap \tilde{\mathfrak{t}}^{-\sigma} \times \mathfrak{t}^{-\sigma}$. Nevertheless, this identity does not give an effective description of $\Pi_{\mathfrak{p}}(\tilde{K}, K)$ because most of the inequalities of the cone $\Pi(\tilde{U}, U)$ become redundant when we restrict them to $\tilde{\mathfrak{t}}^{-\sigma} \times \mathfrak{t}^{-\sigma}$. In the next section, we propose a smaller list of inequalities describing $\Pi_{\mathfrak{p}}(\tilde{K}, K)$.

1.2 Description of the main result

Let $G \subset \tilde{G} \subset \mathrm{GL}(n, \mathbb{R})$ be two connected linear real reductive subgroups with maximal compact subgroups $K \subset \tilde{K} \subset \mathrm{SO}(n, \mathbb{R})$. Let $\mathfrak{a} \subset \tilde{\mathfrak{a}}$ be maximal abelian subspaces of $\mathfrak{p} \subset \tilde{\mathfrak{p}}$.

In this article, we work under the following assumption:

Assumption 1.3 *If \mathfrak{b} is an ideal of $\tilde{\mathfrak{g}}$ contained in \mathfrak{g} , then $\mathfrak{b} \subset Z(\tilde{\mathfrak{g}}) \cap \mathfrak{g}$.*

1.2.1 Admissible elements and polarized trace

An element $\gamma \in \mathfrak{a}$ is called *rational* if the eigenvalues of the symmetric endomorphism $\mathrm{ad}(\gamma) : \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}}$ are rational. We let $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \subset \mathfrak{a}^*$ denote the set of non-zero weights relative to the \mathfrak{a} -action on $\tilde{\mathfrak{g}}/\mathfrak{g}$.

Lemma 1.4 *Under Assumption 1.3, we see that $\mathrm{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}))^\perp = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$.*

Proof : The set $\mathfrak{b} := \{X \in \mathfrak{g}, [X, \tilde{\mathfrak{g}}] \subset \mathfrak{g}\}$ is an ideal of $\tilde{\mathfrak{g}}$ contained in \mathfrak{g} . Thanks to Assumption 1.3, we see that $\mathfrak{b} = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{g}$. Then the vector space $\mathrm{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}))^\perp = \{X \in \mathfrak{a}, [X, \tilde{\mathfrak{g}}] \subset \mathfrak{g}\}$ coincides with $Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{a}$. \square

If $\gamma \in \mathfrak{a}$, we denote by $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp$ the subset of weights vanishing against γ .

Definition 1.5 *An element $\gamma \in \mathfrak{a}$ is admissible if it is rational and if*

$$(2) \quad \mathrm{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \mathrm{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp.$$

If $\ell : E \rightarrow E$ is a symmetric endomorphism of an Euclidean space, we define $\mathrm{Tr}_\ell(E^{\ell>0}) = \sum_{a>0} a \dim(E^{\ell=a})$, where $E^{\ell=a} = \{v \in E, \ell(v) = av\}$.

If $\gamma \in \mathfrak{a}$, its adjoint action $\mathrm{ad}(\gamma) : \mathfrak{g} \rightarrow \mathfrak{g}$ defines a symmetric endomorphism, hence for any subspace $\mathfrak{m} \subset \mathfrak{g}$ stable under $\mathrm{ad}(\gamma)$ one can define the number $\mathrm{Tr}_{\mathrm{ad}(\gamma)}(\mathfrak{m}^{\mathrm{ad}(\gamma)>0})$ that we denote simply by $\mathrm{Tr}_\gamma(\mathfrak{m}^{\gamma>0})$.

1.2.2 Parabolic subgroups

The maximal torus $T \subset U$ is stable under the involution σ , so the Weyl group $W = N_U(T)/T$ is equipped also with an involution still denoted σ , and we denote by W^σ the subgroup fixed by it. Let \mathfrak{R} be the set of roots relative to the action of T on $\mathfrak{u}_\mathbb{C}$.

The maximal compact subgroup $K \subset G$ correspond to the connected component of the subgroup U^σ . We consider the restricted Weyl group $W_{\mathfrak{a}} = N_K(\mathfrak{a})/Z_K(\mathfrak{a})$ relative to the maximal abelian subspace $\mathfrak{a} = \frac{1}{i}\mathfrak{t}^{-\sigma} \subset \frac{1}{i}\mathfrak{u}^{-\sigma} = \mathfrak{p}$. Let Σ be the set of (restricted) roots relative to the action of \mathfrak{a} on \mathfrak{g} .

Let us consider $\gamma \in \mathfrak{a}$ and the subgroups $W^\gamma \subset W$ and $W_{\mathfrak{a}}^\gamma \subset W_{\mathfrak{a}}$ fixing γ . The following fact is explained in §1.3.

Lemma 1.6 *We have canonical identifications $(W/W^\gamma)^\sigma \simeq W^\sigma/W^\sigma \cap W^\gamma \simeq W_{\mathfrak{a}}/W_{\mathfrak{a}}^\gamma$.*

We choose a Weyl chamber $\mathfrak{t}_+ \subset \mathfrak{t}$ so that the restricted Weyl chamber $\mathfrak{a}_+ = \frac{1}{i}\mathfrak{t}_+ \cap \mathfrak{t}^{-\sigma}$ parameters the K -orbits on $\tilde{\mathfrak{p}}$. The choice of \mathfrak{t}_+ determines a system of positive roots \mathfrak{R}^+ , and we denote by $\mathbb{B} \subset U_{\mathbb{C}}$ the Borel subgroup with Lie algebra $\mathfrak{t}_{\mathbb{C}} \oplus \sum_{\alpha \in \mathfrak{R}^+} (\mathfrak{u}_{\mathbb{C}})_{\alpha}$.

To any element $\gamma \in \mathfrak{a}$, we associate the parabolic subgroup

$$(3) \quad \mathbb{P}_\gamma = \{g \in U_{\mathbb{C}}, \lim_{t \rightarrow \infty} e^{t\gamma} g e^{-t\gamma} \text{ exists}\}.$$

Notice that \mathbb{P}_γ is invariant under the involution σ and that $\mathbb{B} \subset \mathbb{P}_\gamma$ when γ is anti-dominant, i.e. $\gamma \in -\mathfrak{a}_+$.

1.2.3 Schubert Calculus

Here we work with two complex reductive groups $U_{\mathbb{C}} \subset \tilde{U}_{\mathbb{C}}$ equipped with an antilinear involution σ . We choose maximal tori $T \subset \tilde{T}$ invariant by σ , such that the corresponding subspaces $\mathfrak{a} := \frac{1}{i}\mathfrak{t}^{-\sigma} \subset \tilde{\mathfrak{a}} := \frac{1}{i}\tilde{\mathfrak{t}}^{-\sigma}$ are maximal abelian in $\mathfrak{p} \subset \tilde{\mathfrak{p}}$.

To any $\gamma \in \mathfrak{a}$, we associate via (3) the parabolic subgroups $\mathbb{P}_\gamma \subset \tilde{\mathbb{P}}_\gamma$ and we define the flag varieties $\mathcal{F}_\gamma = U_{\mathbb{C}}/\mathbb{P}_\gamma$ and $\tilde{\mathcal{F}}_\gamma = \tilde{U}_{\mathbb{C}}/\tilde{\mathbb{P}}_\gamma$. Notice that the involution σ defines an antiholomorphic involution on the flag varieties \mathcal{F}_γ and $\tilde{\mathcal{F}}_\gamma$ that we still denoted by σ . As $\mathbb{P}_\gamma = U_{\mathbb{C}} \cap \tilde{\mathbb{P}}_\gamma$, we have a canonical embedding $\iota : \mathcal{F}_\gamma \hookrightarrow \tilde{\mathcal{F}}_\gamma$. We denote $\iota^* : H^*(\tilde{\mathcal{F}}_\gamma, \mathbb{Z}) \rightarrow H^*(\mathcal{F}_\gamma, \mathbb{Z})$ the pullback in cohomology.

Thanks to the Bruhat decomposition, $\mathcal{F}_\gamma = \bigcup_{w \in W/W^\gamma} \mathbb{B}[w]$, we know that the \mathbb{B} -orbits on \mathcal{F}_γ are parametrized by W/W^γ . We associate to any $w \in W/W^\gamma$, the Schubert cell $\mathfrak{X}_{w,\gamma}^o = \mathbb{B}[w]$, the Schubert variety $\mathfrak{X}_{w,\gamma} := \overline{\mathfrak{X}_{w,\gamma}^o}$ and its cycle class in cohomology $[\mathfrak{X}_{w,\gamma}] \in H^*(\mathcal{F}_\gamma, \mathbb{Z})$.

Thanks to Lemma 1.6, we know that we can attach a Schubert cell $\mathfrak{X}_{w,\gamma}^o$ to any element $w \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^\gamma \simeq (W/W^\gamma)^\sigma$. We will see in §1.3 that an element $w \in W/W^\gamma$ is σ -invariant if and only if $\mathfrak{X}_{w,\gamma}^o \cap (\mathcal{F}_\gamma)^\sigma \neq \emptyset$.

1.2.4 Main result

Let $G \subset \tilde{G}$ be two linear real reductive groups satisfying Assumption 1.3. Let $\pi : \tilde{\mathfrak{p}} \rightarrow \mathfrak{p}$ be the orthogonal projection.

Let $w_0 \in W_{\mathfrak{a}}$ be the unique element such that $w_0(\mathfrak{a}_+) = -\mathfrak{a}_+$. The choice of Weyl chamber \mathfrak{a}_+ defines a system of positive roots Σ^+ , and we denote by $\mathfrak{n} = \sum_{\beta \in \Sigma^+} \mathfrak{g}_\beta$ the corresponding real nilpotent subalgebra of \mathfrak{g} .

Recall that the cohomology class $[pt]$ associated to a singleton $\{pt\} \subset \mathcal{F}_\gamma$ is a basis of $H^{max}(\mathcal{F}_\gamma, \mathbb{Z})$.

Theorem 1.7 *Let $(\tilde{\xi}, \xi) \in \tilde{\mathfrak{a}}_+ \times \mathfrak{a}_+$. We have $K\xi \subset \pi(\tilde{K}\tilde{\xi})$ if and only if*

$$(\tilde{\xi}, \tilde{w}\gamma) \geq (\xi, w_0w\gamma)$$

for any rational antidominant element $\gamma \in \mathfrak{a}$ and any $(w, \tilde{w}) \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^\gamma \times W_{\tilde{\mathfrak{a}}}/W_{\tilde{\mathfrak{a}}}^\gamma$ satisfying the following properties:

- a) $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$.
- b) $[\mathfrak{X}_{w,\gamma}] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma}]) = [pt]$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.
- c) $\text{Tr}_{w\gamma}(\mathfrak{n}^{w\gamma>0}) + \text{Tr}_{\tilde{w}\gamma}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma>0}) = \text{Tr}_\gamma(\tilde{\mathfrak{g}}^{\gamma>0})$.

The result still holds if we replace condition b) by the weaker condition

$$b') \quad [\mathfrak{X}_{w,\gamma}] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma}]) = \ell[pt] \text{ with } \ell \geq 1, \text{ in } H^*(\mathcal{F}_\gamma, \mathbb{Z}).$$

1.3 Bruhat decompositions

The complex parabolic subgroup \mathbb{P}_γ is stable under the involution σ . The intersection $P_\gamma := \mathbb{P}_\gamma \cap G \subset (\mathbb{P}_\gamma)^\sigma$ defines a real parabolic subgroup of G (that is not necessarily connected). Let us explain why the fixed point submanifold $(\mathcal{F}_\gamma)^\sigma$ corresponds to the real flag variety $\mathcal{F}_\gamma^{\mathbb{R}} = G/P_\gamma$. As the flag variety $\mathcal{F}_\gamma = U_{\mathbb{C}}/\mathbb{P}_\gamma \simeq U/U^\gamma$, equipped with the involution σ , admits an identification with the adjoint orbit $U\gamma$, we have

$$(\mathcal{F}_\gamma)^\sigma \simeq U\gamma \cap i\mathfrak{u}^{-\sigma} = K\gamma \simeq K/K^\gamma \simeq G/P_\gamma = \mathcal{F}_\gamma^{\mathbb{R}}.$$

The crucial point here is the equality $U\gamma \cap i\mathfrak{u}^{-\sigma} = K\gamma$ (see [19], Example 2.9).

We fix an element γ_o in the interior of the Weyl chamber \mathfrak{a}_+ and we consider the parabolic subgroup $\mathbb{P} := \mathbb{P}_{-\gamma_o} \subset U_{\mathbb{C}}$. Then $P := \mathbb{P} \cap G$ is the minimal parabolic subgroup of G with Lie algebra $Z_{\mathfrak{k}}(\mathfrak{a}) \oplus \mathfrak{a} \oplus \mathfrak{n}$.

Let us consider the following Bruhat decompositions (see [8, 24]) :

1. $\mathcal{F}_\gamma = \bigcup_{w \in W/W^\gamma} \mathbb{B}[w]$ relative to the \mathbb{B} -action on \mathcal{F}_γ ,
2. $\mathcal{F}_\gamma = \bigcup_{u \in W^{\gamma_o} \backslash W/W^\gamma} \mathbb{P}[u]$ relative to the \mathbb{P} -action on \mathcal{F}_γ ,
3. $\mathcal{F}_\gamma^{\mathbb{R}} = \bigcup_{v \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^\gamma} P[v]$ relative to the P -action on $\mathcal{F}_\gamma^{\mathbb{R}}$.

Now we give a proof of Lemma 1.6. First we recall a standard result relating the Weyl groups $W_{\mathfrak{a}}$ and W with the involution σ (see the Appendix B in [19]). The subgroup W^σ fixed by σ is equal to the normalizer subgroup $N_W(\mathfrak{a})$. Since γ_o is a regular element of \mathfrak{a} ,

the subgroup W^{γ_0} coincides with the centralizer subgroup $Z_W(\mathfrak{a})$, and the restricted Weyl group $W_{\mathfrak{a}}$ admits a canonical identification with $N_W(\mathfrak{a})/Z_W(\mathfrak{a})$.

Thus, we have two exact sequences

$$0 \rightarrow W^{\gamma_0} \rightarrow W^{\sigma} \rightarrow W_{\mathfrak{a}} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow W^{\gamma_0} \rightarrow W^{\sigma} \cap W^{\gamma} \rightarrow W_{\mathfrak{a}}^{\gamma} \rightarrow 0,$$

that induce an isomorphism $W^{\sigma}/W^{\sigma} \cap W^{\gamma} \simeq W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$. The other isomorphism $(W/W^{\gamma})^{\sigma} \simeq W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$ follows from the fact that $W/W^{\gamma} \simeq W^{\gamma} = U^{\gamma} \cap it$: it gives

$$(W/W^{\gamma})^{\sigma} \simeq (W^{\gamma})^{\gamma} = U^{\gamma} \cap it^{-\sigma} = K^{\gamma} \cap it^{-\sigma} = W_{\mathfrak{a}}^{\gamma} \simeq W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}.$$

Notice also that we have an inclusion $W^{\sigma}/W^{\sigma} \cap W^{\gamma} \hookrightarrow W^{\gamma_0} \backslash W/W^{\gamma}$: it is due to the fact that W^{γ_0} is a distinguished subgroup of W^{σ} contained in W^{γ} . Finally we have proven $W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$ can be seen as a subset of $W^{\gamma_0} \backslash W/W^{\gamma}$. The next Lemma characterizes this subset in terms of the intersections $(\mathbb{B}[w])^{\sigma} = \mathbb{B}[w] \cap \mathcal{F}_{\gamma}^{\mathbb{R}}$.

Lemma 1.8 *Let $w \in W$. The following statements are equivalent*

1. $(\mathbb{B}[w])^{\sigma} \neq \emptyset$.
2. $(\mathbb{P}[w])^{\sigma} \neq \emptyset$.
3. *the class of w in $W^{\gamma_0} \backslash W/W^{\gamma}$ is contained in $W^{\sigma}/W^{\sigma} \cap W^{\gamma} \simeq W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$.*

Moreover, if one of the statement is true, then $\mathbb{B}[w] = \mathbb{P}[w]$ and $(\mathbb{B}[w])^{\sigma} = \mathbb{P}[w]$.

Proof : The implication 1. \implies 2. is obvious since $\mathbb{B}[w] \subset \mathbb{P}[w]$. For the implication 2. \implies 3. we consider the Bruhat decomposition $\mathcal{F}_{\gamma}^{\mathbb{R}} = \bigcup_{v \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}} P[v]$. As we have already explained, each class in $W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$ can be represented by an element of W^{σ} . If $(\mathbb{P}[w])^{\sigma} \neq \emptyset$, there exists $v \in W^{\sigma}$ such that $P[v] \subset (\mathbb{P}[w])^{\sigma}$. Hence $\mathbb{P}[v] = \mathbb{P}[w]$, and then $w = v$ in $W^{\gamma_0} \backslash W/W^{\gamma}$. The implication 2. \implies 3. is settled and the last one 3. \implies 1. is immediate.

Consider now $u \in W^{\sigma}$. Since $\mathbb{B} \subset \mathbb{P}$, the Bruhat decomposition 1. tells us that $\mathbb{P}[u] = \bigcup_w \mathbb{B}[w]$ where the union runs over the $w \in W/W^{\gamma}$ such that $w = u$ in $W^{\gamma_0} \backslash W/W^{\gamma}$. As $\sigma(u) = u$, the last relation implies that $w = u$ in W/W^{γ} . Thus we have proven that $\mathbb{P}[u] = \mathbb{B}[u]$. Now the Bruhat decomposition 3. shows that $(\mathbb{B}[u])^{\sigma} = \mathbb{P}[u] \cap \mathcal{F}_{\gamma}^{\mathbb{R}} = \bigcup_v P[v]$ where the union runs over the $v \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma} \simeq W^{\sigma}/W^{\sigma} \cap W^{\gamma}$ such that $P[v] \subset \mathbb{P}[u]$. The last inclusion implies that $v = u$ in $W^{\gamma_0} \backslash W/W^{\gamma}$, and then $v = u$ in $W^{\sigma}/W^{\sigma} \cap W^{\gamma}$. We have proven that $(\mathbb{B}[u])^{\sigma} = \mathbb{P}[u]$. \square

1.4 Comparison with a result of Kapovich-Leeb-Millson

Suppose that $\tilde{G} = G^s$ for $s \geq 2$: Assumption 1.3 holds automatically. Here $\tilde{\mathfrak{p}} = \mathfrak{p}^s$, $\pi(X_1, \dots, X_s) = \sum_{j=1}^s X_j$, and the set $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})$ corresponds to the set Σ of (non-zero) weights relative to the \mathfrak{a} -action on \mathfrak{g} . The restricted Weyl group $W_{\tilde{\mathfrak{a}}}$ is equal to $W_{\mathfrak{a}}^s$.

Recall that we can associate to any $w \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$, the Schubert cell $\mathfrak{X}_{w,\gamma}^o := \mathbb{B}[w] \subset \mathcal{F}_{\gamma}$ and the Schubert variety $\mathfrak{X}_{w,\gamma} := \overline{\mathfrak{X}_{w,\gamma}^o}$.

In this setting, Theorem 1.7 becomes

Theorem 1.9 *Let $(\xi_0, \xi_1, \dots, \xi_s) \in (\mathfrak{a}_+)^{s+1}$. We have $K\xi_0 \subset \sum_{j=1}^s K\xi_j$ if and only if*

$$\sum_{j=1}^s (\xi_j, w_j \gamma) \geq (\xi_0, w_0 w \gamma)$$

for any rational antidominant element $\gamma \in \mathfrak{a}$ and any $(w, w_1, \dots, w_s) \in (W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma})^{s+1}$ satisfying the following properties:

- a) $\text{Vect}(\Sigma \cap \gamma^{\perp}) = \text{Vect}(\Sigma) \cap \gamma^{\perp}$.
- b) $[\mathfrak{X}_{w,\gamma}] \cdot [\mathfrak{X}_{w_1,\gamma}] \cdot \dots \cdot [\mathfrak{X}_{w_s,\gamma}] = [pt]$ in $H^*(\mathcal{F}_{\gamma}, \mathbb{Z})$.
- c) $\text{Tr}_{w\gamma}(\mathfrak{n}^{w\gamma > 0}) + \sum_{j=1}^s \text{Tr}_{w_j\gamma}(\mathfrak{n}^{w_j\gamma > 0}) = s \text{Tr}_{\gamma}(\mathfrak{g}^{\gamma > 0})$.

The submanifold of \mathcal{F}_{γ} fixed by the conjugate linear involution is naturally diffeomorphic to the real flag $\mathcal{F}_{\gamma}^{\mathbb{R}} := G/P_{\lambda}$. Thanks to the Bruhat decomposition $\mathcal{F}_{\gamma}^{\mathbb{R}} = \bigcup_{w \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}} P[w]$, we know that the cycle classes $[\mathfrak{X}_{w,\gamma}^{\mathbb{R}}], w \in W_{\mathfrak{a}}/W_{\mathfrak{a}}^{\gamma}$ associated to the real Bruhat-Schubert varieties $\mathfrak{X}_{w,\gamma}^{\mathbb{R}} := \overline{P[w]}$ defines a basis of the cohomology $H^*(\mathcal{F}_{\gamma}^{\mathbb{R}}, \mathbb{Z}_2)$ with \mathbb{Z}_2 -coefficients [26].

We know from Lemma 1.8 that the real Bruhat-Schubert varieties $\mathfrak{X}_{w,\gamma}^{\mathbb{R}}$ corresponds to the real part of the complex Bruhat-Schubert varieties $\mathfrak{X}_{w,\gamma}$. In this context, a theorem of Borel and Haefliger [5] tells us that

$$b) \quad [\mathfrak{X}_{w,\gamma}] \cdot [\mathfrak{X}_{w_1,\gamma}] \cdot \dots \cdot [\mathfrak{X}_{w_s,\gamma}] = [pt] \quad \text{in} \quad H^*(\mathcal{F}_{\gamma}, \mathbb{Z}).$$

implies the following

$$b^{\mathbb{R}}) \quad [\mathfrak{X}_{w,\gamma}^{\mathbb{R}}] \cdot [\mathfrak{X}_{w_1,\gamma}^{\mathbb{R}}] \cdot \dots \cdot [\mathfrak{X}_{w_s,\gamma}^{\mathbb{R}}] = [pt] \quad \text{in} \quad H^*(\mathcal{F}_{\gamma}^{\mathbb{R}}, \mathbb{Z}_2).$$

The description of $\Pi_{\mathbb{P}}(K^s, K)$ obtained by Kapovich-Leeb-Millson [12] was in terms of elements $(\gamma, w, w_1, \dots, w_s) \in \mathfrak{a} \times W_{\mathfrak{a}}^{s+1}$ verifying conditions a) and b^ℝ). Our description is therefore more accurate, first by adding condition c) and then by taking the refined condition b).

1.5 Levi-movability

In this section, we recall Belkale-Kumar's notion of Levi-movability [2] and explain its connection with conditions b) and c) of Theorem 1.9.

Recall that we work with a reductive Lie group $U_{\mathbb{C}}$ equipped with an anti-linear involution σ . The Lie algebra $\mathfrak{g} := (\mathfrak{u}_{\mathbb{C}})^{\sigma}$ is a real form of the complex Lie algebra $\mathfrak{u}_{\mathbb{C}}$. We

have chosen a maximal torus $T \subset U$ with Lie algebra \mathfrak{t} , that is invariant under σ , and such that $\mathfrak{t}^{-\sigma}$ is of maximal dimension. Let $\mathfrak{a} = \frac{1}{i}\mathfrak{t}^{-\sigma} \subset (\mathfrak{u}_{\mathbb{C}})^{\sigma}$.

We choose a Weyl chamber $\mathfrak{t}_+ \subset \mathfrak{t}$ such that $\mathfrak{a}_+ = \frac{1}{i}(\mathfrak{t}^{-\sigma} \cap \mathfrak{t}_+)$ is a Weyl chamber relatively to the restricted root system Σ . Let $\mathbb{B} \subset U_{\mathbb{C}}$ be the Borel subgroup with Lie algebra $\mathfrak{t}_{\mathbb{C}} \oplus \sum_{\alpha \in \mathfrak{R}^+} (\mathfrak{u}_{\mathbb{C}})_{\alpha}$. To any element $\gamma \in \mathfrak{a}$, we associate the parabolic subgroup $\mathbb{P}_{\gamma} = \{g \in U_{\mathbb{C}}, \lim_{t \rightarrow \infty} e^{t\gamma} g e^{-t\gamma} \text{ exists}\}$ that is invariant under the involution σ . Notice that $\mathbb{B} \subset \mathbb{P}_{\gamma}$ when γ is anti-dominant, i.e. $\gamma \in -\mathfrak{a}_+$.

The antilinear involution σ defines an anti-holomorphic involution on the flag variety $\mathcal{F}_{\gamma} = U_{\mathbb{C}}/\mathbb{P}_{\gamma}$. We consider the cellular decomposition $\mathcal{F}_{\gamma} = \coprod_{w \in W/W^{\gamma}} \mathbb{B}[w]$ parameterized by the Weyl group $W = N_U(\mathfrak{t})/T$ and the subgroup W^{γ} fixing $\gamma \in \mathfrak{a}$. For any $w \in W/W^{\gamma}$, we denote by $\mathfrak{X}_{w,\gamma}^{\circ} = \mathbb{B}[w] \subset \mathcal{F}_{\gamma}$ the Schubert cell.

The cycle class of the Schubert variety $\mathfrak{X}_{w,\gamma} = \overline{\mathbb{B}[w]}$ in $H^*(\mathcal{F}_{\gamma}, \mathbb{Z})$ is denoted by $[\mathfrak{X}_{w,\gamma}]$ and it is called a Schubert class. Let $\mathfrak{X}_{w_i,\gamma}$ be d Schubert classes parameterized by $w_i \in W/W^{\gamma}$, $i \in [d]$. If there exists an integer k such that $[\mathfrak{X}_{w_1,\gamma}] \cdots [\mathfrak{X}_{w_d,\gamma}] = k[pt]$, then we set $c_{\gamma}(w_1, \dots, w_d) = k$; we set $c_{\gamma}(w_1, \dots, w_d) = 0$ otherwise.

Using the transversality theorem of Kleiman, Belkale and Kumar showed in [2], Proposition 2, the following important lemma.

Lemma 1.10 *Suppose that γ is anti-dominant. The coefficient $c_{\gamma}(w_1, \dots, w_d)$ is nonzero if and only if for generic $(p_1, \dots, p_d) \in (\mathbb{P}_{\gamma})^d$, the intersection $p_1 w_1^{-1} \mathfrak{X}_{w_1,\gamma}^{\circ} \cap \cdots \cap p_d w_d^{-1} \mathfrak{X}_{w_d,\gamma}^{\circ}$ is transverse at e .*

Then Belkale-Kumar defined Levi-movability.

Definition 1.11 *Suppose that γ is anti-dominant. A d -uplet $(w_1, \dots, w_d) \in W^d$ is Levi-movable if for generic $(l_1, \dots, l_d) \in (U_{\mathbb{C}}^{\gamma})^d$, the intersection $l_1 w_1^{-1} \mathfrak{X}_{w_1,\gamma}^{\circ} \cap \cdots \cap l_d w_d^{-1} \mathfrak{X}_{w_d,\gamma}^{\circ}$ is transverse at e .*

Belkale-Kumar have defined in [2], a new product \odot_0 on $H^*(\mathcal{F}_{\gamma}, \mathbb{Z})$ that is closely related to the notion of Levi-movability. By definition of this product, the identity

$$[\mathfrak{X}_{w_1,\gamma}] \odot_0 \cdots \odot_0 [\mathfrak{X}_{w_d,\gamma}] = \ell[pt]$$

holds for some $\ell \geq 1$ if and only if $[\mathfrak{X}_{w_1,\gamma}] \cdots [\mathfrak{X}_{w_d,\gamma}] = \ell[pt]$ and the d -uplet (w_1, \dots, w_d) is Levi-movable.

In the next proposition we concentrate ourself to d -uplets (w_1, \dots, w_d) of σ -invariant elements. Recall that $\mathfrak{n} \subset \mathfrak{g}$ is the nilpotent Lie subalgebra defined as the subspace $\mathfrak{g}^{\gamma_{\circ} > 0}$ for any regular dominant element $\gamma_{\circ} \in \mathfrak{a}_+$.

Proposition 1.12 *Let $\gamma \in \mathfrak{a}$ be a rational anti-dominant element. A d -uplet $(w_1, \dots, w_d) \in (W^{\sigma})^d$ satisfies the following conditions*

b) $[\mathfrak{X}_{w_1,\gamma}] \cdots [\mathfrak{X}_{w_d,\gamma}] = [pt]$ in $H^*(\mathcal{F}_{\gamma}, \mathbb{Z})$.

$$c) \sum_{j=1}^d \mathrm{Tr}_{w_j \gamma}(\mathfrak{n}^{w_j \gamma > 0}) = (d-1) \mathrm{Tr}_\gamma(\mathfrak{g}^{\gamma > 0}).$$

if and only if $[\mathfrak{X}_{w_1, \gamma}] \odot_0 \dots \odot_0 [\mathfrak{X}_{w_d, \gamma}] = [pt]$.

Proof: Since \mathfrak{g} is the real part of the complex Lie algebra $\mathfrak{u}_\mathbb{C}$, we have the following identity of \mathfrak{a} -modules : $\mathfrak{u}_\mathbb{C} \simeq \mathfrak{g}^2$. Thus $\mathrm{Tr}_\gamma(\mathfrak{g}^{\gamma > 0}) = \frac{1}{2} \mathrm{Tr}_\gamma(\mathfrak{u}_\mathbb{C}^{\gamma > 0})$. The nilpotent radical of the Lie algebra of \mathbb{B} is $\mathfrak{n}_{\mathfrak{u}_\mathbb{C}} = E \oplus \mathfrak{n}_\mathbb{C}$, where $E = \mathfrak{n}_{\mathfrak{u}_\mathbb{C}} \cap Z_{\mathfrak{u}_\mathbb{C}}(\mathfrak{a})$. Thus for any $w \in W_{\mathfrak{a}}$, the vector space $\mathfrak{n}_{\mathfrak{u}_\mathbb{C}}^{w \gamma > 0}$ is the complexification of $\mathfrak{n}^{w \gamma > 0}$. It follows that $\mathrm{Tr}_{w \gamma}(\mathfrak{n}^{w \gamma > 0}) = \frac{1}{2} \mathrm{Tr}_{w \gamma}(\mathfrak{n}_{\mathfrak{u}_\mathbb{C}}^{w \gamma > 0})$. We have proven that $c)$ is equivalent to $\sum_{j=1}^d \mathrm{Tr}_{w_j \gamma}(\mathfrak{n}_{\mathfrak{u}_\mathbb{C}}^{w_j \gamma > 0}) = (d-1) \mathrm{Tr}_\gamma(\mathfrak{u}_\mathbb{C}^{\gamma > 0})$. Let ρ_u be half the sum of the positive roots and let $\gamma' = i\gamma \in \mathfrak{t}^{-\sigma} \cap -\mathfrak{t}_+$. Standard computations give that $\mathrm{Tr}_{w \gamma}(\mathfrak{n}_{\mathfrak{u}_\mathbb{C}}^{w \gamma > 0}) = \langle \rho_u, w \gamma' \rangle - \langle \rho_u, \gamma' \rangle$ and $\mathrm{Tr}_\gamma(\mathfrak{u}_\mathbb{C}^{\gamma > 0}) = -2 \langle \rho_u, \gamma' \rangle$. Finally we see that $c)$ is equivalent to

$$c') \quad \sum_{j=1}^d \langle \rho_u, w_j \gamma' \rangle = (2-d) \langle \rho_u, \gamma' \rangle.$$

Consider the complex $U_\mathbb{C}$ -manifold $M = (U_\mathbb{C}/\mathbb{B})^d$. The set $C = \prod_{k=1}^d U_\mathbb{C}^{\gamma'} [w_k^{-1}]$ is a connected component of the submanifold $M^{\gamma'}$. The Bialynicki-Birula's complex submanifold $C^- := \{m \in M, \lim_{t \rightarrow \infty} \exp(-it\gamma')m \in C\}$ is equal to $C = \prod_{k=1}^d \mathbb{P}' [w_k^{-1}]$ where \mathbb{P}' is the parabolic subgroup with Lie algebra $\mathfrak{t}_\mathbb{C} \oplus \sum_{\langle \alpha, \gamma' \rangle \leq 0} (\mathfrak{u}_\mathbb{C})_\alpha$. Notice that the Borel subgroup \mathbb{B} is contained in \mathbb{P}' as $\gamma' \in -\mathfrak{t}_+$.

Now we consider the map $\pi : U_\mathbb{C} \times_{\mathbb{P}'} C^- \rightarrow M$ that sends $[g, x]$ to gx . Standard computations shows the following statement are equivalent

1. Conditions $b)$ and $c')$ holds.
2. π is a birational map with an exceptional set that does not contains C .
3. $[\mathfrak{X}_{w_1, \gamma}] \odot_0 \dots \odot_0 [\mathfrak{X}_{w_d, \gamma}] = [pt]$.

See §5 in [6]. \square

Proposition 1.12 permits us to describe the convex cone $\Pi_{\mathbb{p}}(K^s, K)$ in terms of the Belkale-Kumar's product \odot_0 .

Theorem 1.13 *Let $(\xi_0, \xi_1, \dots, \xi_s) \in (\mathfrak{a}_+)^{s+1}$. We have $K\xi_0 \subset \sum_{j=1}^s K\xi_j$ if and only if $\sum_{j=1}^s (\xi_j, w_j \gamma) \geq (\xi_0, w_0 w \gamma)$ for any rational antidominant element $\gamma \in \mathfrak{a}$ and any $(w, w_1, \dots, w_s) \in (W_{\mathfrak{a}}/W_{\mathfrak{a}}^\gamma)^{s+1}$ satisfying the following properties:*

1. $\mathrm{Vect}(\Sigma \cap \gamma^\perp) = \mathrm{Vect}(\Sigma) \cap \gamma^\perp$.
2. $[\mathfrak{X}_{w, \gamma}] \odot_0 [\mathfrak{X}_{w_1, \gamma}] \odot_0 \dots \odot_0 [\mathfrak{X}_{w_s, \gamma}] = [pt]$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.

1.6 Singular Horn cone

Let $n \geq 1$. If A is an Hermitian $n \times n$ matrix, we denote by $s(A) = (s_1(A) \geq \dots \geq s_n(A))$ its spectrum. The Horn cone $\text{Horn}(n)$ is defined as the set of triplets $(s(A), s(B), s(C))$ where A, B, C are Hermitian $n \times n$ matrices satisfying $A + B + C = 0$.

Denote the set of cardinality r subsets $I = \{i_1 < i_2 < \dots < i_r\}$ of $[n] = \{1, \dots, n\}$ by \mathcal{P}_r^n . To each $I \in \mathcal{P}_r^n$ we associate

- a weakly decreasing sequence of non-negative integers $\lambda(I) = (\lambda_1 \geq \dots \geq \lambda_r)$ where $\lambda_a = n - r + a - i_a$ for $a \in [r]$.
- a Schubert class $[\mathfrak{X}_I] \in H^*(\mathbb{G}(r, n), \mathbb{Z})$ where $\mathbb{G}(r, n)$ denotes the Grassmann variety of r -dimensional subspaces of \mathbb{C}^n .

The following Horn's conjecture [11] was settled in the affirmative by combining the work of A. Klyachko [14] with the work of A. Knutson and T. Tao [15] on the "saturation" problem. We refer the reader to survey articles [9, 6] for details.

If $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $I \subset [n]$, we define $|x|_I = \sum_{i \in I} x_i$ and $|x| = \sum_{i=1}^n x_i$. Let $1^r = (1, \dots, 1) \in \mathbb{R}^r$.

Theorem 1.14 (Horn's conjecture) *An element $(x, y, z) \in (\mathbb{R}_+^n)^3$ belongs to $\text{Horn}(n)$ if and only if the following conditions holds : $|x| + |y| + |z| = 0$ and*

$$|x|_I + |y|_J + |z|_K \leq 0$$

for any $r \in [n - 1]$, for any $I, J, K \in \mathcal{P}_r^n$ such that

$$(4) \quad (\lambda(I), \lambda(J), \lambda(K) - (n - r)1^r) \in \text{Horn}(r).$$

Thanks to the saturation Theorem of Knutson-Tao, we know that (4) is equivalent to the following relation in $H^*(\mathbb{G}(r, n), \mathbb{Z})$

$$(5) \quad [\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = \ell[pt] \quad \ell \geq 1,$$

where $[pt] \in H^{\max}(\mathbb{G}(r, n), \mathbb{Z})$ denote the Poincaré dual class of the point. Belkale proved in [1] that Theorem 1.14 holds if we replace condition (5) by

$$(6) \quad [\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = [pt].$$

Finally, Knutson-Tao-Woodward proved in [16] that the list of inequalities parameterized by the (r, I, J, K) satisfying (6) is optimal.

We turn now our attention to the singular Horn cone. Suppose that $n = p + q$ with $p \geq q \geq 1$. If A is a complex $p \times q$ matrix, we denote by $\tau(A) = (\tau_1(A) \geq \dots \geq \tau_q(A) \geq 0)$ its singular spectrum : $\tau_k(A) = \sqrt{s_k(A^*A)}$ for any $k \in [q]$.

Definition 1.15 *The singular Horn cone, denoted $\text{Singular}(p, q)$, is defined as the set of triplets $(\tau(A), \tau(B), \tau(C))$ where A, B, C are complex $p \times q$ matrices satisfying $A + B + C = 0$.*

We have a natural connection between $\text{Singular}(p, q)$ and $\text{Horn}(n)$ because, for any complex $p \times q$ matrix A , the spectrum of the $n \times n$ matrix

$$\begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix}$$

is $(\tau_1(A) \geq \dots \geq \tau_q(A) \geq 0 \geq \dots \geq 0 \geq -\tau_q(A) \geq \dots \geq -\tau_1(A))$. To any $x = (x_1, \dots, x_q) \in \mathbb{R}^q$ we associate the vector $\tilde{x} = (x_1, \dots, x_q, 0, \dots, 0, -x_q, \dots, -x_1) \in \mathbb{R}^n$. From the previous remark, we see that

$$(x, y, z) \in \text{Singular}(p, q) \implies (\tilde{x}, \tilde{y}, \tilde{z}) \in \text{Horn}(n).$$

The O'Shea-Sjamaar theorem tells us that the implication above is in fact an equivalence. Hence we obtain a first description of $\text{Singular}(p, q)$.

Proposition 1.16 *Let $\mathbb{R}_{++}^q = \{x = (x_1 \geq \dots \geq x_q \geq 0)\}$. An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if*

$$(7) \quad |\tilde{x}|_I + |\tilde{y}|_J + |\tilde{z}|_K \leq 0$$

for any $r \in [n-1]$, for any $I, J, K \in \mathcal{P}_r^n$ such that $[\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = [pt]$ in $H^*(\mathbb{G}(r, n), \mathbb{Z})$.

The main issue with this first description is that most of the inequalities (7) are redundant.

Now we explain the more precise description of $\text{Singular}(p, q)$ that we obtain by applying Theorem 1.13. For any $r \in [q]$, let $\mathcal{P}_r^{p,q} \subset \mathcal{P}_r^n$ be the collection of subsets I satisfying² $I \cap I^o = \emptyset$ and $I \cap \{q+1, \dots, p\} = \emptyset$. We see that any $I \in \mathcal{P}_r^{p,q}$ is equal to the union $I_+ \coprod I_-$ where I_+, I_- are disjoint subsets of $[q]$.

Notice that, when $I, J, K \in \mathcal{P}_r^{p,q}$, the inequality (7) becomes

$$(\star)_{I_{\pm}, J_{\pm}, K_{\pm}} \quad |x|_{I_+} + |y|_{J_+} + |z|_{K_+} \leq |x|_{I_-} + |y|_{J_-} + |z|_{K_-}.$$

The first version of our result is the following theorem that answers (partially) a conjecture of A. S. Buch (see [9], §5).

Theorem 1.17 *An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ holds for any $r \in [q]$, for any $I, J, K \in \mathcal{P}_r^{p,q}$ such that $[\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = [pt]$ in $H^*(\mathbb{G}(r, n), \mathbb{Z})$.*

² $I^o = \{n+1-k, k \in I\}$

The above description can be refined as follows. For any $r \in [q]$ such that $r < \frac{n}{2}$, we denote by $\mathbb{F}(r, n-r; n)$ the two-steps flag variety parameterizing nested sequences of linear subspaces $E \subset F^\perp \subset \mathbb{C}^n$ where $\dim E = \dim F = r$. Here the orthogonal is taken relatively to the bilinear product $(x, y) = \sum_{i=1}^n x_i y_i$. In the case where $p = q$ and $r = q = \frac{n}{2}$, we denote by $\mathbb{F}(q, q; n)$ (or by $\mathbb{G}(q, n)$) the Grassmannian parameterizing linear subspaces $E \subset \mathbb{C}^n$ of dimension q .

Definition 1.18 *Let $A \subset [n]$ be a subset of cardinal $2r$. We say that A is polarized if it admits a decomposition $A = A_+ \amalg A_-$ into disjoint subsets. If A_+ and A_- are both of cardinal r , we say that A is balanced. We denote by Pol_{2r}^n the collection of balanced polarized subsets of $[n]$ of cardinal $2r$.*

The orbits of the Borel subgroup $B_n \subset GL_n(\mathbb{C})$ of upper-triangular matrices on $\mathbb{F}(r, n-r; n)$ are parametrized by the balanced polarized subsets of $[n]$ of cardinal $2r$. To any balanced polarized subset $A = A_+ \amalg A_- \in \text{Pol}_{2r}^n$, we associate

- the flag $\mathbb{C}^{A_+} \subset (\mathbb{C}^{A_-})^\perp$, where $\mathbb{C}^{A_\pm} = \text{Vect}(e_i, i \in A_\pm) \subset \mathbb{C}^n$. Notice that in the case where $p = q$ and $r = q = \frac{n}{2}$, we have $\mathbb{C}^{A_+} = (\mathbb{C}^{A_-})^\perp$.
- the Schubert variety $\mathfrak{X}_A = \overline{B_n \cdot (\mathbb{C}^{A_+} \subset (\mathbb{C}^{A_-})^\perp)} \subset \mathbb{F}(r, n-r; n)$, and its cycle class $[\mathfrak{X}_A] \in H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$ in cohomology.

To any subset $I \in \mathcal{P}_r^{p,q}$, we associate the balanced polarized subset $\widehat{I} = I \amalg I^o \in \text{Pol}_{2r}^n$ and the cohomology class $[\mathfrak{X}_{\widehat{I}}] \in H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$. The final description of $\text{Singular}(p, q)$ is given in the following Theorem (whose proof is postponed at §4.6).

Theorem 1.19 *An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if $(\star)_{I_\pm, J_\pm, K_\pm}$ holds for any $r \in [q]$, for any $I, J, K \in \mathcal{P}_r^{p,q}$ such that*

$$(8) \quad [\mathfrak{X}_{\widehat{I}}] \odot_0 [\mathfrak{X}_{\widehat{J}}] \odot_0 [\mathfrak{X}_{\widehat{K}}] = [pt] \quad \text{in} \quad H^*(\mathbb{F}(r, n-r; n), \mathbb{Z}).$$

In Section §4.1, we will explain why condition (8) implies condition (6).

Let us end this section with the example of $\text{Singular}(3, 3)$. The first description obtained from the O'Shea-Sjamaar theorem says that $\text{Singular}(3, 3)$ admits a natural identification with the intersection $\text{Horn}(6) \cap V^3$ where $V = \{(a, b, c, -c, -b, -a)\} \subset \mathbb{R}^6$. Thanks to computer calculation we know that $\text{Horn}(6) \subset (\mathbb{R}^6)^3$ is described with a minimal list of 536 inequalities. Theorem 1.19 allows to describe $\text{Singular}(3, 3)$ in terms of $87 + 9$ inequalities (see §4.4).

2 Parametrization of facets of real Kirwan polytopes

Let $L_{\mathbb{C}}$ be a connected complex reductive group with maximal compact subgroup L . Let σ be a conjugate linear involution on $L_{\mathbb{C}}$ commuting with the Cartan involution. We work with a σ -invariant scalar product $(-, -)$ on \mathfrak{l} , allowing an identification $\mathfrak{l} \simeq \mathfrak{l}^*$.

Let us consider a proper Kähler-Hamiltonian L -manifold with involution $(M, \Omega, \mathbb{J}, \Phi_{\mathfrak{l}}, \tau, \sigma)$ (see §1.1). By that we mean :

- $L_{\mathbb{C}}$ acts holomorphically on the complex manifold (M, \mathbb{J}) and the Kähler form Ω is L -invariant,
- There is a L -equivariant *proper* moment mapping $\Phi_{\mathfrak{l}} : M \rightarrow \mathfrak{l}$ satisfying the relation³ $d(\Phi_{\mathfrak{l}}, X) = -\Omega(X_M, -), \forall X \in \mathfrak{l}$.
- The anti-holomorphic involution $\tau : M \rightarrow M$ satisfies the following relations : $\tau^*(\Omega) = -\Omega$, and $\tau(g \cdot m) = \sigma(g) \cdot \tau(m)$, $\Phi_{\mathfrak{l}}(\tau(m)) = -\sigma(\Phi_{\mathfrak{l}}(m))$, for all $(g, m) \in L_{\mathbb{C}} \times M$.

Let us denote by G_L the connected component of the subgroup $(L_{\mathbb{C}})^{\sigma}$. The Lie algebra of G_L admits the Cartan decomposition $\mathfrak{g}_L = \mathfrak{k}_L \oplus \mathfrak{p}_L$ where $\mathfrak{p}_L = i\mathfrak{l}^{-\sigma}$ and \mathfrak{k}_L is the Lie algebra of $K_L := (L^{\sigma})_o$.

The submanifold $Z = \{m \in M, \tau(m) = m\}$, that we suppose *non-empty*, is a Lagrangian submanifold of (M, Ω) , not necessarily connected, and stable under the action of the real reductive group G_L . Let \mathcal{Z} be a connected component of Z . We define the K_L -equivariant map

$$\Psi : \mathcal{Z} \longrightarrow \mathfrak{p}_L$$

by the relation $\Psi(z) = \frac{1}{i}\Phi_{\mathfrak{l}}(z)$. One sees that, for any $\beta \in \mathfrak{p}_L$, $\beta_{\mathcal{Z}}$ is the gradient vector field of the function (Ψ, β) .

The O'Shea-Sjamaar Theorem [19] tells us that the image $\Psi(\mathcal{Z})$ is equal to $\Phi_{\mathfrak{l}}(M) \cap \mathfrak{l}^{-\sigma}$ through the identification $\mathfrak{p}_L \simeq \mathfrak{l}^{-\sigma}$ (for another proof see [20]). Let $\mathfrak{a}_L \subset \mathfrak{p}_L$ be a maximal abelian subspace. The K_L -orbits of \mathfrak{p}_L are parametrized by a Weyl chamber $(\mathfrak{a}_L)_+ \subset \mathfrak{a}_L$. Hence the K_L -orbits of $\Psi(\mathcal{Z})$ are parametrized by the real Kirwan polytope

$$\Delta(\mathcal{Z}) := \Psi(\mathcal{Z}) \cap (\mathfrak{a}_L)_+.$$

Thanks to the O'Shea-Sjamaar Theorem, we know that $\Delta(\mathcal{Z})$ is a convex polytope. In the next sections we explain the parametrization of the facets of $\Delta(\mathcal{Z})$ in terms of balanced Ressayre's pairs that we obtained in [20].

³Here $X_M(m) = -X \cdot m = \frac{d}{dt}|_{t=0} e^{-tX} m$ is the vector field generated by $X \in \mathfrak{l}$.

2.1 Admissible elements

The stabilizer subgroups of $m \in M$ relatively to the L and G_L actions are denoted respectively by L^m and $(G_L)^m$: their Lie algebras are denoted by \mathfrak{l}^m and $(\mathfrak{g}_L)^m$. We will take a particular attention to the subspace $(\mathfrak{p}_L)^m = \{X \in \mathfrak{p}_L, X \cdot m = 0\} \subset \mathfrak{g}_m$. If $\gamma \in \mathfrak{a}_L$, we denote by \mathcal{Z}^γ the submanifold where the vector field $z \mapsto \gamma \cdot z$ vanishes.

Definition 2.1 *Let us define*

$$\dim_{\mathfrak{p}_L}(\mathcal{X}) := \min_{z \in \mathfrak{X}} \dim((\mathfrak{p}_L)^z)$$

for any subset $\mathcal{X} \subset \mathcal{Z}$. A non-zero element $\gamma \in \mathfrak{a}$ is called *admissible relatively to the G -manifold \mathcal{Z}* if γ is rational, and if $\dim_{\mathfrak{p}_L}(\mathcal{Z}^\gamma) - \dim_{\mathfrak{p}_L}(\mathcal{Z}) \in \{0, 1\}$.

The next results, proven in [20], are useful to determine $\dim_{\mathfrak{p}_L}(\mathcal{Z})$.

Lemma 2.2 *There exists a subspace $\mathfrak{h} \subset \mathfrak{p}_L$ such that*

1. $\forall x \in \mathcal{Z}, \exists k \in K_L$ such that $\text{Ad}(k)(\mathfrak{h}) \subset (\mathfrak{p}_L)^x$,
2. $\dim(\mathfrak{h}) = \dim((\mathfrak{p}_L)^x)$ on a dense open subset of \mathcal{Z} .

Therefore $\dim_{\mathfrak{p}_L}(\mathcal{Z}) = \dim(\mathfrak{h})$.

The subspace \mathfrak{h} is called *the generic \mathfrak{p}_L -stabilizer of \mathcal{Z}* .

Corollary 2.3 *Let $\mathfrak{h}_o \subset \mathfrak{p}_L$ be a subspace such that the set $K\mathcal{Z}^{\mathfrak{h}_o}$ has a non-empty interior in \mathcal{Z} . Then $\dim_{\mathfrak{p}_L}(\mathcal{Z}) \geq \dim(\mathfrak{h}_o)$.*

2.2 Balanced Ressayre's pair

Let $\gamma \in \mathfrak{a}_L$ be a non-zero element, and let $C \subset M^\gamma$ be a connected component intersecting \mathcal{Z} . The Kähler submanifold C is stable under the involution τ and under the action of the stabilizer subgroup $L_C^\gamma = \{g \in L_C, g\gamma = \gamma\}$. The real submanifold $\mathcal{C} = C \cap \mathcal{Z} \subset \mathcal{Z}^\gamma$ is non-empty, not necessarily connected, and stable under the action of the group $G_L^\gamma = L_C^\gamma \cap G_L$. The function (Ψ, γ) is locally constant on M^γ and we denote by $(\Psi(\mathcal{C}), \gamma)$ its value on \mathcal{C} .

The Bialynicki-Birula's complex submanifold

$$(9) \quad C^- := \{m \in M, \lim_{t \rightarrow \infty} \exp(t\gamma)m \in C\}.$$

is invariant under the involution of M . The infinitesimal action of γ defines a symmetric endomorphism of the tangent bundle $\mathbf{TM}|_C$, and one sees that $\mathbf{TC}^-|_C = (\mathbf{TM}|_C)^{\gamma \leq 0}$.

Consider now the parabolic subgroups defined by

$$(10) \quad \mathbb{P}_\gamma = \{g \in L_C, \lim_{t \rightarrow \infty} \exp(t\gamma)g \exp(-t\gamma) \text{ exists}\} \quad \text{and} \quad P_\gamma = G_L \cap \mathbb{P}_\gamma.$$

Here the Lie algebra of \mathbb{P}_γ and P_γ are respectively $(\mathfrak{t}_\mathbb{C})^{\gamma \leq 0}$ and $(\mathfrak{g}_L)^{\gamma \leq 0}$. We denote by $\mathbb{P} \subset L_\mathbb{C}$ and $P \subset G_L$ the parabolic subgroups $\mathbb{P}_{-\gamma_o}$ and $P_{-\gamma_o}$, where γ_o is an element of the interior of the Weyl chamber $(\mathfrak{a}_L)_+$. The nilradical of the Lie algebra of P is $\mathfrak{n}_L := \mathfrak{g}^{\gamma_o > 0}$.

The submanifold C^- is invariant under the action of \mathbb{P}_γ , hence we can consider the complex manifold $\mathbb{P} \times_{\mathbb{P} \cap \mathbb{P}_\gamma} C^-$ and the holomorphic map

$$q_\gamma^{\mathbb{C}} : \mathbb{P} \times_{\mathbb{P} \cap \mathbb{P}_\gamma} C^- \rightarrow M$$

that sends $[p, m]$ to pm . The complex manifold $\mathbb{P} \times_{\mathbb{P} \cap \mathbb{P}_\gamma} C^-$ is equipped with a natural anti-holomorphic involution $\tau_\gamma : [p, m] \mapsto [\sigma(p), \tau(m)]$ such that $\tau \circ q_\gamma^{\mathbb{C}} = q_\gamma^{\mathbb{C}} \circ \tau_\gamma$. The submanifold fixed by the involution is $P \times_{P \cap P_\gamma} C^-$ where $C^- = C^- \cap \mathcal{Z}$ is a submanifold of \mathcal{Z} . Notice that for any $z \in \mathcal{C}$, we have $\mathbf{T}_z C^- = (\mathbf{T}_z \mathcal{Z})^{\gamma \leq 0}$.

We can now define the *real* counterpart of the map $q_\gamma^{\mathbb{C}}$: the map

$$q_\gamma : P \times_{P \cap P_\gamma} C^- \rightarrow \mathcal{Z}.$$

The tangent bundle at the identity of $P/P \cap P_\gamma$ admits a canonical identification with $(\mathfrak{n}_L)^{\gamma > 0}$, so that for any $z \in \mathcal{C}$, the tangent map $\mathbf{T}q_\gamma|_{[e, z]} : (\mathfrak{n}_L)^{\gamma > 0} \times (\mathbf{T}_z \mathcal{Z})^{\gamma \leq 0} \mapsto \mathbf{T}_z \mathcal{Z}$ is defined by the relation $\mathbf{T}q_\gamma|_{[e, z]}(X, v) = X \cdot z \oplus v$. Hence for any $z \in \mathcal{C}$ the following statements are equivalent :

- $\mathbf{T}q_\gamma^{\mathbb{C}}|_{[e, z]}$ is an isomorphism
- $\mathbf{T}q_\gamma|_{[e, z]}$ is an isomorphism
- $\rho_z^\gamma : (\mathfrak{n}_L)^{\gamma > 0} \mapsto (\mathbf{T}_z \mathcal{Z})^{\gamma > 0}$, $X \mapsto X \cdot z$ is an isomorphism.

Definition 2.4 *The data (γ, \mathcal{C}) is called a **balanced Ressayre's pair** of \mathcal{Z} if there exists τ -invariant Zariski open subsets $\mathcal{V}^{\mathbb{C}} \subset M$ and $\mathcal{U}^{\mathbb{C}} \subset C^-$ such that : $\mathcal{V}^{\mathbb{C}}$ is \mathbb{P} -invariant, $\mathcal{U}^{\mathbb{C}}$ is $\mathbb{P} \cap \mathbb{P}_\gamma$ -invariant and intersects C , and the map $q_\gamma^{\mathbb{C}}$ defines a diffeomorphism $\mathbb{P} \times_{\mathbb{P} \cap \mathbb{P}_\gamma} \mathcal{U}^{\mathbb{C}} \simeq \mathcal{V}^{\mathbb{C}}$.*

*If furthermore we have $\dim_{\mathfrak{p}}(\mathcal{C}) - \dim_{\mathfrak{p}}(\mathcal{Z}) \in \{0, 1\}$, and γ is rational, we call (γ, \mathcal{C}) a **regular balanced Ressayre's pair**.*

If $\mathcal{U}^{\mathbb{C}}$ is a τ -invariant Zariski open subset of C^- that intersects the complex submanifold C , then $\mathcal{U}^{\mathbb{C}} \cap C$ is a τ -invariant Zariski open subset of C . It implies that $\mathcal{U}^{\mathbb{C}} \cap C^\tau$ is dense in C^τ , and in particular that $\mathcal{U}^{\mathbb{C}} \cap \mathcal{C}$ is non-empty (see Appendix B in [20]). When the map $q_\gamma^{\mathbb{C}}$ defines a diffeomorphism $\mathbb{P} \times_{\mathbb{P} \cap \mathbb{P}_\gamma} \mathcal{U}^{\mathbb{C}} \simeq \mathcal{V}^{\mathbb{C}}$, we see then that the infinitesimal map $\rho_z^\gamma : (\mathfrak{n}_L)^{\gamma > 0} \mapsto (\mathbf{T}_z \mathcal{Z})^{\gamma > 0}$ is an isomorphism for any $z \in \mathcal{U}^{\mathbb{C}} \cap \mathcal{C}$.

We can weaken the condition on the balanced Ressayre's pair.

Definition 2.5 *The data (γ, \mathcal{C}) is called a **infinitesimal Ressayre's pair** of \mathcal{Z} if there exists $z \in \mathcal{C}$ such that $\rho_z^\gamma : (\mathfrak{n}_L)^{\gamma > 0} \mapsto (\mathbf{T}_z \mathcal{Z})^{\gamma > 0}$ is an isomorphism. If furthermore we have $\dim_{\mathfrak{p}}(\mathcal{C}) - \dim_{\mathfrak{p}}(\mathcal{Z}) \in \{0, 1\}$, and γ is rational, we call (γ, \mathcal{C}) a **regular infinitesimal Ressayre's pair** of \mathcal{Z} .*

When (γ, \mathcal{C}) is an infinitesimal real Ressayre's pair of \mathcal{Z} , there exists $z \in \mathcal{C}$ so that $\rho_z^\gamma : (\mathfrak{n}_L)^{\gamma > 0} \longrightarrow (\mathbf{T}_x \mathcal{Z})^{\gamma > 0}$ is an isomorphism commuting with the infinitesimal action of γ . It gives us two identities

$$\begin{aligned} (C_1) \quad & \dim(\mathfrak{n}_L)^{\gamma > 0} = \text{rank}(\mathbf{T}\mathcal{Z}|_{\mathcal{C}})^{\gamma > 0}, \\ (C_2) \quad & \text{Tr}_\gamma((\mathfrak{n}_L)^{\gamma > 0}) = \text{Tr}_\gamma((\mathbf{T}\mathcal{Z}|_{\mathcal{C}})^{\gamma > 0}). \end{aligned}$$

If (γ, \mathcal{C}) is a balanced Ressayre's pair then

$$(C_3) \quad \left\{ x \in M, \#\{(q_\gamma^{\mathbb{C}})^{-1}(x)\} = 1 \right\} \text{ contains a Zariski open subset of } M.$$

Condition (C_3) can be made more flexible by considering

$$(C'_3) \quad \left\{ x \in M, (q_\gamma^{\mathbb{C}})^{-1}(x) \neq \emptyset \right\} \text{ contains a Zariski open subset of } M.$$

2.3 Parametrization of the facets

The following result will be the main tool used in the next sections.

Theorem 2.6 ([20]) *For $\xi \in (\mathfrak{a}_L)_+$, the following statements are equivalent:*

- $\xi \in \Delta_{\mathfrak{p}}(\mathcal{Z})$.
- $(\xi, \gamma) \geq (\Psi(\mathcal{C}), \gamma)$ holds for any regular infinitesimal real Ressayre's pair (γ, \mathcal{C}) of \mathcal{Z} .
- $(\xi, \gamma) \geq (\Psi(\mathcal{C}), \gamma)$ holds for any regular balanced Ressayre's pair (γ, \mathcal{C}) of \mathcal{Z} .
- $(\xi, \gamma) \geq (\Psi(\mathcal{C}), \gamma)$ holds for any (γ, \mathcal{C}) such that
 - a) γ is rational,
 - b) $\dim_{\mathfrak{p}}(\mathcal{C}) - \dim_{\mathfrak{p}}(\mathcal{Z}) \in \{0, 1\}$,
 - c) (C_1) , (C_2) and (C_3) hold.

The result is still true if we replace condition c) by

$$c') \quad (C_1), (C_2) \text{ and } (C'_3) \text{ hold.}$$

3 Proof of the main theorem

We come back to the setting of §1.1. We will explain how Theorem 1.7 is a consequence of Theorem 2.6 applied to the Kähler-Hamiltonian manifold $M = \mathbf{T}\tilde{U}$ equipped with the action of the reductive group $L_{\mathbb{C}} = \tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}}$.

The real reductive group $G \subset \tilde{G}$ are the connected components of the subgroups $U_{\mathbb{C}}^{\sigma} \subset \tilde{U}_{\mathbb{C}}^{\sigma}$. We have a Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where $\mathfrak{k} = \mathfrak{u}^{\sigma}$ and $\mathfrak{p} = i\mathfrak{u}^{-\sigma}$ (idem for $\tilde{\mathfrak{g}}$). Let $\mathfrak{a} \subset \tilde{\mathfrak{a}}$ be maximal abelian subspaces of $\mathfrak{p} \subset \tilde{\mathfrak{p}}$.

3.1 Admissible elements

We see \tilde{G} as a submanifold of $\tilde{U}_\mathbb{C}$ equipped with the following $\tilde{G} \times G$ -action : $(\tilde{g}, g) \cdot \tilde{x} = \tilde{g}\tilde{x}g^{-1}$. Let (\tilde{X}, X) be an element of $\tilde{\mathfrak{a}} \times \mathfrak{a}$. In the following lemma, we describe the manifold $\tilde{G}^{(\tilde{X}, X)}$ of zeroes of the vector field (\tilde{X}, X) on \tilde{G} . We denote by $\tilde{G}^X \subset \tilde{G}$ the subgroup that fixes X .

Lemma 3.1 • *If $\tilde{X} \notin \tilde{K}X$, then $\tilde{G}^{(\tilde{X}, X)} = \emptyset$*
 • *If $\tilde{X} = \tilde{k}X$ with $\tilde{k} \in \tilde{K}$, then $\tilde{G}^{(\tilde{X}, X)} = \tilde{k} \cdot \tilde{G}^X$.*

Proof : The set $\tilde{G}^{(\tilde{X}, X)}$ is non-empty if and only if \tilde{X} belongs to the adjoint orbit $\tilde{G}X$. The Lemma follows then from the fact that the intersection $\tilde{G}X \cap \tilde{\mathfrak{p}}$ is equal to $\tilde{K}X$. \square

Let $W_{\tilde{\mathfrak{a}}} = N_{\tilde{K}}(\tilde{\mathfrak{a}})/Z_{\tilde{K}}(\tilde{\mathfrak{a}})$ be the restricted Weyl group. Since both elements \tilde{X} and X belongs to $\tilde{\mathfrak{a}}$, we notice that if $\tilde{X} = \tilde{k}X$ for some $\tilde{k} \in \tilde{K}$, then there exists $\tilde{w} \in W_{\tilde{\mathfrak{a}}}$ such that $\tilde{X} = \tilde{w}X$. So, the admissible elements relative to the action of $\tilde{G} \times G$ on \tilde{G} are of the form $(\tilde{w}\gamma, \gamma)$ where $\gamma \in \mathfrak{a}$ is a rational element and $\tilde{w} \in W_{\tilde{\mathfrak{a}}}$.

We let $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \subset \mathfrak{a}^*$ denote the set of weights relative to the \mathfrak{a} -action on $\tilde{\mathfrak{g}}/\mathfrak{g}$. If $\gamma \in \mathfrak{a}$, we denote by $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp$ the subset of weights vanishing against γ . The main purpose of this section is the proof of the following result.

Proposition 3.2 *Let $G \subset \tilde{G}$ be two linear real reductive groups satisfying Assumption 1.3. An element $(\tilde{w}\gamma, \gamma)$ is admissible relatively to the $\tilde{G} \times G$ -action on \tilde{G} if and only if γ is rational and if $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$.*

The proof of Proposition 3.2 is a consequence of the next Lemma.

Lemma 3.3 *Let $\gamma \in \mathfrak{a}$ and let $\tilde{X}_o \in \tilde{\mathfrak{a}}$ be a regular element.*

1. *We have $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G} = \dim_{\mathfrak{p}} \tilde{K}\tilde{X}_o$ and $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G}^{(\tilde{w}\gamma, \gamma)} = \dim_{\mathfrak{p}} \tilde{K}^\gamma \tilde{X}_o$.*
2. *Under Assumption 1.3, we have*
 - $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G} = \dim Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$.
 - $\dim_{\mathfrak{p}} \tilde{K}^\gamma \tilde{X}_o - \dim_{\mathfrak{p}} \tilde{K}\tilde{X}_o \in \{0, 1\}$ *if and only if* $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp)$ *is equal to* $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$.

Proof : For any $x \in \tilde{G}$, the stabilizer $(\tilde{\mathfrak{p}} \times \mathfrak{p})^x$ admits a canonical identification with $\text{Ad}(x)(\mathfrak{p}) \cap \tilde{\mathfrak{p}}$. If we write $x = \tilde{k}'e^{\tilde{Y}}$, with $\tilde{k}' \in \tilde{K}$ and $\tilde{Y} \in \tilde{\mathfrak{p}}$, we see that $\text{Ad}(x)(\mathfrak{p}) \cap \tilde{\mathfrak{p}} \simeq \text{Ad}(e^{\tilde{Y}})(\mathfrak{p}) \cap \tilde{\mathfrak{p}} = \mathfrak{p} \cap \tilde{\mathfrak{p}}^{\tilde{Y}}$. Thus $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G} = \min_{\tilde{Y} \in \tilde{\mathfrak{p}}} \dim(\mathfrak{p} \cap \tilde{\mathfrak{p}}^{\tilde{Y}})$. If we write $\tilde{Y} = \tilde{k} \cdot \tilde{X}$, with $\tilde{X} \in \tilde{\mathfrak{a}}$, we have $\text{Ad}(\tilde{k})(\tilde{\mathfrak{a}}) \subset \tilde{k} \cdot \tilde{\mathfrak{p}}^{\tilde{X}} = \tilde{\mathfrak{p}}^{\tilde{Y}}$, and the equality $\text{Ad}(\tilde{k})(\tilde{\mathfrak{a}}) = \tilde{\mathfrak{p}}^{\tilde{Y}}$ holds when \tilde{Y} is regular. So $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G} = \min_{\tilde{k} \in \tilde{K}} \dim(\mathfrak{p} \cap \text{Ad}(\tilde{k})(\tilde{\mathfrak{a}}))$. On the other hand, when

$\tilde{X}_o \in \tilde{\mathfrak{a}}$ is a regular element it is immediate that $\dim_{\mathfrak{p}} \tilde{K} \tilde{X}_o = \min_{\tilde{k} \in \tilde{K}} \dim(\mathfrak{p} \cap \text{Ad}(\tilde{k})(\tilde{\mathfrak{a}}))$. We obtain finally that $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G} = \dim_{\mathfrak{p}} \tilde{K} \tilde{X}_o$. Since $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G}^{(\tilde{w}\gamma, \gamma)} = \dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G}^\gamma$, the identity $\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G}^{(\tilde{w}\gamma, \gamma)} = \dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o$ admits the same proof than before. First one sees that

$$\dim_{\tilde{\mathfrak{p}} \times \mathfrak{p}} \tilde{G}^\gamma = \min_{\tilde{k} \in \tilde{K}^\gamma} \dim(\mathfrak{p} \cap \text{Ad}(\tilde{k})(\tilde{\mathfrak{a}})) = \min_{\tilde{k} \in \tilde{K}^\gamma} \dim(\mathfrak{p}^\gamma \cap \text{Ad}(\tilde{k})(\tilde{\mathfrak{a}}))$$

and then we use that $\min_{\tilde{k} \in \tilde{K}^\gamma} \dim(\mathfrak{p}^\gamma \cap \text{Ad}(\tilde{k})) = \dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o$. The first point is settled.

$\tilde{K} \tilde{X}_o$ is the real part of the coadjoint orbit $\tilde{U} \tilde{X}_o$. Thanks to Lemma 2.2, we know that exists a subspace $\mathfrak{h} \subset \mathfrak{p}$ such that $\forall x \in \tilde{K} \tilde{X}_o, \exists k \in K$ such that $\text{Ad}(k)(\mathfrak{h}) \subset \mathfrak{p}^x$, and $\dim(\mathfrak{h}) = \dim(\mathfrak{p}^x)$ on a dense open subset of $\tilde{K} \tilde{X}_o$: \mathfrak{h} is the generic \mathfrak{p} -stabilizer on $\tilde{K} \tilde{X}_o$ and $\dim_{\mathfrak{p}}(\tilde{K} \tilde{X}_o) = \dim(\mathfrak{h})$.

Let us check that under Assumption 1.3, we have $\mathfrak{h} = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$. If $\beta' \in \mathfrak{h}$, then $K(\tilde{K} \tilde{X}_o)^{\beta'} = \tilde{K} \tilde{X}_o$. Let us write $\beta' = \text{Ad}(k')(\beta)$ with $\beta \in \mathfrak{a}$ and $k' \in K$. Since $(\tilde{K} \tilde{X}_o)^\beta = \cup_{\tilde{w} \in W_{\tilde{\mathfrak{a}}}} \tilde{K}^\beta \tilde{w} \tilde{X}_o$, the relation $K(\tilde{K} \tilde{X}_o)^{\beta'} = \tilde{K} \tilde{X}_o$ implies that a subset $K \tilde{K}^\beta \tilde{w} \tilde{X}_o \subset \tilde{K} \tilde{X}_o$ has a non empty interior. Let $g = k \tilde{k} \tilde{w} \in K \tilde{K}^\beta \tilde{w}$ such that $g \tilde{X}_o$ belongs to the interior of $K \tilde{K}^\beta \tilde{w} \tilde{X}_o$: this implies the identity

$$(11) \quad \mathfrak{k} + \text{Ad}(k)(\tilde{\mathfrak{k}}^\beta) + \text{Ad}(g)(\tilde{\mathfrak{k}}^{\tilde{X}_o}) = \tilde{\mathfrak{k}}.$$

But $\text{Ad}(g)(\tilde{\mathfrak{k}}^{\tilde{X}_o}) = \text{Ad}(k \tilde{k})(\tilde{\mathfrak{k}}^{\tilde{X}_o}) \subset \text{Ad}(k)(\tilde{\mathfrak{k}}^\beta)$, hence (11) is equivalent to $\mathfrak{k} + \tilde{\mathfrak{k}}^\beta = \tilde{\mathfrak{k}}$. Now standard computation shows that

$$\mathfrak{k} + \tilde{\mathfrak{k}}^\beta = \tilde{\mathfrak{k}} \iff [\beta, \tilde{\mathfrak{k}}] \subset \mathfrak{p} \iff [\beta, \tilde{\mathfrak{p}}] \subset \mathfrak{k} \iff [\beta, \tilde{\mathfrak{g}}] \subset \mathfrak{g} \iff [\beta', \tilde{\mathfrak{g}}] \subset \mathfrak{g}.$$

We have proved that any element $\beta' \in \mathfrak{h} \subset \mathfrak{p}$ belongs to the ideal $\mathfrak{b} := \{X \in \mathfrak{g}, [X, \tilde{\mathfrak{g}}] \subset \mathfrak{g}\}$. Thanks to Assumption 1.3, we know that $\mathfrak{b} = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{g}$. We have proved that $\mathfrak{h} \subset Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$. On the other hand, it is immediate that $Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \subset \mathfrak{p}^x$ for any $x \in \tilde{K} \tilde{X}_o$. We have proved that the equality $Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} = \mathfrak{p}^x$ holds on a dense open subset of $\tilde{K} \tilde{X}_o$. The first part of the second point is settled.

Let us concentrated on the last part of the second point.

If $\gamma \in Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$ then $\dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o = \dim_{\mathfrak{p}} \tilde{K} \tilde{X}_o$ and $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}))$.

Suppose now that $\gamma \notin Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p}$. We see that $Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma \subset (\mathfrak{p}^\gamma)^x$ for any $x \in \tilde{K}^\gamma \tilde{X}_o$, hence $\dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o - \dim_{\mathfrak{p}} \tilde{K} \tilde{X}_o = 1$ if and only if $Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma$ is the generic \mathfrak{p}^γ -stabilizer on $\tilde{K}^\gamma \tilde{X}_o$.

Suppose that $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$. For any $\beta \in \mathfrak{a}$, we have the equivalences

$$\begin{aligned} [\beta, \tilde{\mathfrak{g}}^\gamma] \subset \mathfrak{g}^\gamma &\iff \langle \alpha, \beta \rangle = 0, \forall \alpha \in \Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp &\iff \beta \in \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp)^\perp \\ &&\iff \beta \in \left(\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp \right)^\perp \\ &&\iff \beta \in \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}))^\perp + \mathbb{R}\gamma \\ &&\iff \beta \in Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma. \end{aligned}$$

Let $\mathfrak{h}_\gamma \subset \mathfrak{p}^\gamma$ be the generic \mathfrak{p}^γ -stabilizer on $\tilde{K}^\gamma \tilde{X}_o$. Let $\beta' \in \mathfrak{h}_\gamma$ that we write $\beta' = k\beta$ with $\beta \in \mathfrak{a}$ and $k \in K^\gamma$. Since $K^\gamma(\tilde{K}^\gamma \tilde{X}_o)^{\beta'} = K^\gamma(\tilde{K}^\gamma \tilde{X}_o)^\beta$ is equal to $\tilde{K}^\gamma \tilde{X}_o$, the same arguments than those used before show that $[\beta, \tilde{\mathfrak{g}}^\gamma] \subset \mathfrak{g}^\gamma$ and then $\beta' = \beta \in Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma$. We have proved that $\mathfrak{h}_\gamma = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma$ is the generic \mathfrak{p}^γ -stabilizer on $\tilde{K}^\gamma \tilde{X}_o$.

Suppose now that $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) \neq \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$. It means that there exists $\beta \notin Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}))^\perp + \mathbb{R}\gamma$ such that $\langle \alpha, \beta \rangle = 0, \forall \alpha \in \Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp$, i.e. $[\beta, \tilde{\mathfrak{g}}^\gamma] \subset \mathfrak{g}^\gamma$. Let consider the subspace $\mathfrak{h}_o = Z(\tilde{\mathfrak{g}}) \cap \mathfrak{p} \oplus \mathbb{R}\gamma \oplus \mathbb{R}\beta$. As we have shown before, the inclusion $[\beta, \tilde{\mathfrak{g}}^\gamma] \subset \mathfrak{g}^\gamma$ shows that the set $K^\gamma(\tilde{K}^\gamma \tilde{X}_o)^{\mathfrak{h}_o} = K^\gamma(\tilde{K}^\gamma \tilde{X}_o)^\beta$ has a non-empty interior. Thanks to Lemma 2.3, we can conclude that $\dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o \geq \dim(\mathfrak{h}_o)$ and then $\dim_{\mathfrak{p}^\gamma} \tilde{K}^\gamma \tilde{X}_o - \dim_{\mathfrak{p}} \tilde{K}^\gamma \tilde{X}_o \geq 2$. The last part of the second point is proved. \square

3.2 Balanced Ressayre's pair : infinitesimal conditions

Let $(\tilde{w}, \gamma) \in W_{\tilde{\mathfrak{a}}} \times \mathfrak{a}$ such that $\gamma_{\tilde{w}} := (\tilde{w}\gamma, \gamma)$ is an admissible element relative to the action of $\tilde{G} \times G$ on $\tilde{G} \subset \tilde{U}_{\mathbb{C}}$.

The complex submanifold of $\tilde{U}_{\mathbb{C}}$ fixed by $\gamma_{\tilde{w}}$ is $C_{\tilde{w}, \gamma} := \tilde{w}\tilde{U}_{\mathbb{C}}^\gamma$ and the real part contained in \tilde{G} is $\mathcal{C}_{\tilde{w}, \gamma} := C_{\tilde{w}, \gamma} \cap \tilde{G} = \tilde{w}\tilde{G}^\gamma$. The Bialynicki-Birula's complex submanifold associated to $C_{\tilde{w}, \gamma}$ is $C_{\tilde{w}, \gamma}^- := \tilde{w}\tilde{\mathbb{P}}_\gamma$ where $\tilde{\mathbb{P}}_\gamma$ is the complex parabolic subgroup of $\tilde{U}_{\mathbb{C}}$ associated to γ (see (10)). The parabolic subgroup of $\tilde{U}_{\mathbb{C}} \times U_{\mathbb{C}}$ associated to the weight $\gamma_{\tilde{w}}$ is

$$\mathbb{P}_{\tilde{w}, \gamma} := \text{Ad}(\tilde{w})(\tilde{\mathbb{P}}_\gamma) \times \mathbb{P}_\gamma.$$

Let $\mathbb{P} \subset U_{\mathbb{C}}$ (resp. $\tilde{\mathbb{P}} \subset \tilde{U}_{\mathbb{C}}$) be the parabolic subgroup associated to the choice of the Weyl chamber \mathfrak{a}_+ (resp. $\tilde{\mathfrak{a}}_+$). We can now consider the holomorphic map

$$q_{\tilde{w}, \gamma}^{\mathbb{C}} : (\tilde{\mathbb{P}} \times \mathbb{P}) \times_{(\tilde{\mathbb{P}} \times \mathbb{P}) \cap \mathbb{P}_{\tilde{w}, \gamma}} \tilde{w}\tilde{\mathbb{P}}_\gamma \longrightarrow \tilde{U}_{\mathbb{C}}$$

that sends $[\tilde{p}, p; \tilde{w}\tilde{x}]$ to $\tilde{p}\tilde{w}\tilde{x}p^{-1}$. Let us recall the definition of regular balanced Ressayre's pair in this setting.

Definition 3.4 *The data $(\gamma_{\tilde{w}}, \mathcal{C}_{\tilde{w}, \gamma})$ is a regular balanced Ressayre's pair of \tilde{G} if there exists τ -invariant Zariski open subsets $\mathcal{V}^{\mathbb{C}} \subset \tilde{U}_{\mathbb{C}}$ and $\mathcal{U}^{\mathbb{C}} \subset \tilde{w}\tilde{\mathbb{P}}_\gamma$ such that, $\mathcal{V}^{\mathbb{C}}$ is $\tilde{\mathbb{P}} \times \mathbb{P}$ -invariant, $\mathcal{U}^{\mathbb{C}}$ is $(\tilde{\mathbb{P}} \times \mathbb{P}) \cap \mathbb{P}_{\tilde{w}, \gamma}$ -invariant and intersects $\tilde{w}\tilde{U}_{\mathbb{C}}^\gamma$, and the map $q_{\tilde{w}, \gamma}^{\mathbb{C}}$ defines a diffeomorphism*

$$(12) \quad (\tilde{\mathbb{P}} \times \mathbb{P}) \times_{(\tilde{\mathbb{P}} \times \mathbb{P}) \cap \mathbb{P}_{\tilde{w}, \gamma}} \mathcal{U}^{\mathbb{C}} \simeq \mathcal{V}^{\mathbb{C}}.$$

As we have explain in §2.2, the set $\mathcal{U}^{\mathbb{C}} \cap \tilde{w}\tilde{G}^\gamma$ is nonempty and the diffeomorphism (12) insures that the infinitesimal map $\rho_z^{\gamma_{\tilde{w}}} : \tilde{\mathfrak{n}}^{\tilde{w}\gamma > 0} \times \mathfrak{n}^{\gamma > 0} \longrightarrow (\mathbf{T}_z \tilde{G})^{\gamma_{\tilde{w}} > 0}$ is an isomorphism for any $z \in \mathcal{U}^{\mathbb{C}} \cap \tilde{w}\tilde{G}^\gamma$.

Lemma 3.5 *Let $(\gamma_{\tilde{w}}, \mathcal{C}_{\tilde{w}, \gamma})$ be a regular balanced Ressayre's pair of \tilde{G} . Then we have*

$$(C_1) \quad \dim(\tilde{\mathfrak{n}}^{\tilde{w}\gamma > 0}) + \dim(\mathfrak{n}^{\gamma > 0}) = \dim(\tilde{\mathfrak{g}}^{\gamma > 0}),$$

$$(C_2) \quad \text{Tr}_\gamma(\mathfrak{n}^{\gamma > 0}) + \text{Tr}_{\tilde{w}\gamma}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma > 0}) = \text{Tr}_\gamma(\tilde{\mathfrak{g}}^{\gamma > 0}).$$

Proof : Let $z \in \tilde{w}\tilde{G}^\gamma$ such that $\rho_z^{\gamma\tilde{w}} : \tilde{\mathfrak{n}}^{\tilde{w}\gamma>0} \times \mathfrak{n}^{\gamma>0} \longrightarrow (\mathbf{T}_z\tilde{G})^{\gamma\tilde{w}>0}$ is an isomorphism. Since $\rho_z^{\gamma\tilde{w}}$ commutes with the infinitesimal action of $\gamma\tilde{w}$, it implies that $\dim(\tilde{\mathfrak{n}}^{\tilde{w}\gamma>0}) + \dim(\mathfrak{n}^{\gamma>0}) = \dim((\mathbf{T}_z\tilde{G})^{\gamma\tilde{w}>0})$ and $\mathrm{Tr}_{\gamma\tilde{w}}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma>0} \times \mathfrak{n}^{\gamma>0}) = \mathrm{Tr}_{\gamma\tilde{w}}((\mathbf{T}_z\tilde{G})^{\gamma\tilde{w}>0})$.

Since $(\mathbf{T}_z\tilde{G})^{\gamma\tilde{w}>0} \simeq \tilde{\mathfrak{g}}^{\gamma>0}$, relations (C₁) and (C₂) follows. \square

3.3 Cohomological conditions

To any $\tilde{w} \in W_{\tilde{a}}$, we associate the Schubert cell $\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o = \tilde{\mathbb{P}}[\tilde{w}] = \tilde{\mathbb{B}}[\tilde{w}] \subset \tilde{\mathcal{F}}_\gamma$ and the Schubert variety $\tilde{\mathfrak{X}}_{\tilde{w},\gamma} = \overline{\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o}$ (see §1.3). Let us denote $\mathfrak{X}_\gamma^o := \mathbb{B}[e] \subset \mathcal{F}_\gamma$ and the corresponding Schubert variety $\mathfrak{X}_\gamma = \overline{\mathfrak{X}_\gamma^o}$.

Let $[\tilde{\mathfrak{X}}_{\tilde{w},\gamma}] \in H^{2n_{\tilde{w},\gamma}}(\tilde{\mathcal{F}}_\gamma, \mathbb{Z})$ and $[\mathfrak{X}_\gamma] \in H^{2n_\gamma}(\mathcal{F}_\gamma, \mathbb{Z})$ be the cycle classes defined by these algebraic subvarieties. Here $n_{\tilde{w},\gamma} = \mathrm{codim}_{\mathbb{C}}(\tilde{\mathfrak{X}}_{\tilde{w},\gamma})$ and $n_\gamma = \mathrm{codim}_{\mathbb{C}}(\mathfrak{X}_\gamma)$.

A standard computation shows that condition (C₁) of Lemma 3.5 is equivalent to $\mathrm{codim}_{\mathbb{C}}(\tilde{\mathfrak{X}}_{\tilde{w},\gamma}) + \mathrm{codim}_{\mathbb{C}}(\mathfrak{X}_\gamma) = \dim_{\mathbb{C}}(\mathcal{F}_\gamma)$.

If $(\gamma\tilde{w}, \mathcal{C}_{\tilde{w},\gamma})$ is a balanced Ressayre's pair of \tilde{G} , then the map $q_{\tilde{w},\gamma}^{\mathbb{C}}$ satisfies condition (C₃) : $\{\tilde{g} \in \tilde{U}_{\mathbb{C}}, \#\{(q_{\tilde{w},\gamma}^{\mathbb{C}})^{-1}(\tilde{g}^{-1})\} = 1\}$ contains a Zariski open subset of $\tilde{U}_{\mathbb{C}}$.

Let us consider the map

$$\pi_{\tilde{w},\gamma} : (\tilde{\mathbb{P}} \times \mathbb{P}) \times_{(\tilde{\mathbb{P}} \times \mathbb{P}) \cap \mathbb{P}_{\tilde{w},\gamma}} \tilde{w}\tilde{\mathbb{P}}_\gamma \longrightarrow \tilde{\mathcal{F}}_\gamma \times \mathcal{F}_\gamma$$

that sends $[\tilde{p}, p; \tilde{w}x]$ to $([\tilde{p}\tilde{w}], [p])$.

The following facts are standard (see Proposition 4.3 in [6] or Section 6.1 in [20]). For any $\tilde{g} \in \tilde{U}_{\mathbb{C}}$, the map $\pi_{\tilde{w},\gamma}$ is injective on the fiber $(q_{\tilde{w},\gamma}^{\mathbb{C}})^{-1}(\tilde{g}^{-1})$, and the set $\pi_{\tilde{w},\gamma}((q_{\tilde{w},\gamma}^{\mathbb{C}})^{-1}(\tilde{g}^{-1}))$ is equal to

$$\mathfrak{X}_\gamma^o \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o) \simeq \{(x, y) \in \tilde{\mathbb{P}}[\tilde{w}] \times \mathbb{P}[e], \tilde{g}x = \iota(y)\} \subset \tilde{\mathcal{F}}_\gamma \times \mathcal{F}_\gamma.$$

Here $\iota : \mathcal{F}_\gamma \rightarrow \tilde{\mathcal{F}}_\gamma$ is the canonical embedding. Thus conditions (C₃) and (C'₃) becomes

$$(C_3) \quad \left\{ \tilde{g} \in \tilde{U}_{\mathbb{C}}, \#\{\mathfrak{X}_\gamma^o \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o)\} = 1 \right\} \text{ contains a Zariski open subset of } \tilde{U}_{\mathbb{C}}.$$

$$(C'_3) \quad \left\{ \tilde{g} \in \tilde{U}_{\mathbb{C}}, \mathfrak{X}_\gamma^o \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o) \neq \emptyset \right\} \text{ contains a Zariski open subset of } \tilde{U}_{\mathbb{C}}.$$

According to a transversality theorem due to Kleiman, the intersection $\mathfrak{X}_\gamma \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma})$ is transverse for any \tilde{g} in a nonempty Zariski open subset of $\tilde{U}_{\mathbb{C}}$; moreover, $\mathfrak{X}_\gamma^o \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma}^o)$ is dense in this intersection. We deduce that

1. (C₃) is equivalent to asking that the cohomology class of $\mathfrak{X}_\gamma \cap \iota^{-1}(\tilde{g}\tilde{\mathfrak{X}}_{\tilde{w},\gamma})$ is equal to the cohomology class of a point : $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma}]) = [pt]$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.
2. (C₁) and (C'₃) are equivalent to asking that $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma}]) = \ell[pt]$ with $\ell \geq 1$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.

Let us notice that $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma}]) = [pt]$ implies that $\mathrm{codim}_{\mathbb{C}}(\tilde{\mathfrak{X}}_{\tilde{w},\gamma}) + \mathrm{codim}_{\mathbb{C}}(\mathfrak{X}_\gamma) = \dim_{\mathbb{C}}(\mathcal{F}_\gamma)$, hence (C₁) is a consequence of (C₃).

3.4 Conclusion

Let us consider the gradient map $\Psi : \tilde{G} \simeq \tilde{K} \times \tilde{\mathfrak{p}} \rightarrow \tilde{\mathfrak{p}} \times \mathfrak{p}, (\tilde{k}, \tilde{X}) \mapsto (-\tilde{k}\tilde{X}, \pi(\tilde{X}))$.

Let $(\tilde{w}, \gamma) \in W_{\tilde{\mathfrak{a}}} \times \mathfrak{a}$ and $\mathcal{C}_{\tilde{w}, \gamma} = \tilde{w}\tilde{G}^\gamma$. In the previous sections, we have proved the following

- $\gamma_{\tilde{w}} := (\tilde{w}\gamma, \gamma)$ is admissible if and only if $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$.
- Condition (C₂) for $(\gamma_{\tilde{w}}, \mathcal{C}_{\tilde{w}, \gamma})$ is equivalent to $\text{Tr}_\gamma(\mathfrak{n}^{\gamma>0}) + \text{Tr}_{\tilde{w}\gamma}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma>0}) = \text{Tr}_\gamma(\tilde{\mathfrak{g}}^{\gamma>0})$.
- Condition (C₃) for $(\gamma_{\tilde{w}}, \mathcal{C}_{\tilde{w}, \gamma})$ is equivalent to $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w}, \gamma}]) = [pt]$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.
- (C₁) is a consequence of (C₃).
- Conditions (C₁) and (C'₃) for $(\gamma_{\tilde{w}}, \mathcal{C}_{\tilde{w}, \gamma})$ are equivalent to $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w}, \gamma}]) = \ell[pt]$ with $\ell \geq 1$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.

We notice that $(\Psi(\mathcal{C}_{\tilde{w}, \gamma}), \gamma_{\tilde{w}}) = 0$. Thus Theorem 2.6 gives us the following

Theorem 3.6 *Let $(\tilde{\xi}, \xi) \in \tilde{\mathfrak{a}}_+ \times \mathfrak{a}_+$. We have $-K\xi \subset \pi(\tilde{K}\tilde{\xi})$ if and only if*

$$(13) \quad (\tilde{\xi}, \tilde{w}\gamma) + (\xi, \gamma) \geq 0$$

for any $(\gamma, \tilde{w}) \in \mathfrak{a} \times W_{\tilde{\mathfrak{a}}}$ satisfying the following properties:

- a) γ is rational and $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$.
- b) $[\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w}, \gamma}]) = [pt]$ in $H^*(\mathcal{F}_\gamma, \mathbb{Z})$.
- c) $\text{Tr}_\gamma(\mathfrak{n}^{\gamma>0}) + \text{Tr}_{\tilde{w}\gamma}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma>0}) = \text{Tr}_\gamma(\tilde{\mathfrak{g}}^{\gamma>0})$.

The result still holds if we replace condition b) with

$$b') \quad [\mathfrak{X}_\gamma] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w}, \gamma}]) = \ell[pt] \text{ with } \ell \geq 1 \text{ in } H^*(\mathcal{F}_\gamma, \mathbb{Z}).$$

We have to make some small modifications in the previous theorem in order to obtain Theorem 1.7. The longest element $w_0 \in W$, which is invariant by σ , can be seen as the unique element of $W_{\mathfrak{a}}$ such that $w_0(\mathfrak{a}_+) = -\mathfrak{a}_+$. Any elements $\xi \in \mathfrak{a}_+$ and $\gamma \in \mathfrak{a}$ can be written $\xi = -w_0(\xi')$ and $\gamma = w\gamma'$ where $\xi' \in \mathfrak{a}_+$, $\gamma' \in -\mathfrak{a}_+$ and $w \in W_{\mathfrak{a}}$. Since $W_{\mathfrak{a}}\gamma = K\gamma \cap \mathfrak{a}$ is contained in $W_{\tilde{\mathfrak{a}}}\gamma = \tilde{K}\gamma \cap \tilde{\mathfrak{a}}$, for any $(w, \tilde{w}) \in W_{\mathfrak{a}} \times W_{\tilde{\mathfrak{a}}}$, there exists $\tilde{w}' \in W_{\tilde{\mathfrak{a}}}$ such that $\tilde{w}w\gamma' = \tilde{w}'\gamma'$.

We leaves the reader verify the following

- $-K\xi \subset \pi(\tilde{K}\tilde{\xi})$ is equivalent to $K\xi' \subset \pi(\tilde{K}\tilde{\xi})$
- (13) is equivalent to $(\tilde{\xi}, \tilde{w}'\gamma') \geq (\xi', w_0w\gamma')$

- $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap \gamma^\perp$ is equivalent to $\text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) \cap (\gamma')^\perp) = \text{Vect}(\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g})) \cap (\gamma')^\perp$.
- Condition *b*) is equivalent to $[\mathfrak{X}_{w,\gamma'}] \cdot \iota^*([\tilde{\mathfrak{X}}_{\tilde{w},\gamma'}]) = [pt]$ in $H^*(\mathcal{F}_{\gamma'}, \mathbb{Z})$.
- Condition *c*) is equivalent to $\text{Tr}_{w\gamma'}(\mathfrak{n}^{w\gamma'>0}) + \text{Tr}_{\tilde{w}\gamma'}(\tilde{\mathfrak{n}}^{\tilde{w}\gamma'>0}) = \text{Tr}_{\gamma'}(\tilde{\mathfrak{g}}^{\gamma'>0})$.

We see then that Theorem 1.7 is a reformulation of Theorem 3.6.

4 Singular value inequalities for matrix sums

Let $p \geq q \geq 1$ and $n = p + q$. We consider the canonical action of the group $K := U(p) \times U(q)$ on the vector space $M_{p,q}(\mathbb{C})$ of complex matrices of size $p \times q$. Let $\mathfrak{a} \subset M_{p,q}(\mathbb{C})$ be the real subspace consisting of all matrices of the form

$$Y(x) = \begin{pmatrix} 0 & \cdots & x_1 \\ \vdots & \ddots & \vdots \\ x_q & \cdots & 0 \\ 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{pmatrix}, x \in \mathbb{R}^q.$$

The chamber $\mathfrak{a}_+ := \{Y(x), x_1 \geq x_2 \geq \cdots \geq x_q \geq 0\}$ parametrizes the K -orbits on $M_{p,q}(\mathbb{C})$: in other words a matrix X belongs to the orbit $K \cdot Y(x)$ if and only x is equal to the singular spectrum of X (see §1.6). Thus we can consider

$$\text{Singular}(p, q) := \{(a, b, c) \in (\mathbb{R}_{++}^q)^3, K \cdot Y(c) \subset K \cdot Y(a) + K \cdot Y(b)\},$$

where $\mathbb{R}_{++}^q = \{x = (x_1 \geq x_2 \geq \cdots \geq x_q \geq 0)\}$. Notice that $K \cdot Y(x) = -K \cdot Y(x)$ for any $x \in \mathbb{R}_{++}^q$, hence $\text{Singular}(p, q)$ can be defined with the symmetric condition $0 \in K \cdot Y(a) + K \cdot Y(b) + K \cdot Y(c)$.

We need to recall some notations introduced in §1.6. For any $r \in [q]$, let $\mathcal{P}_r^{p,q}$ be the collection of subsets $I \subset [n]$ of cardinal r , satisfying $I \cap I^o = \emptyset$ and $I \cap \{q+1, \dots, p\} = \emptyset$, where $I^o = \{n+1-k, k \in I\}$. We see that any $I \in \mathcal{P}_r^{p,q}$ is equal to the disjoint union $I_+ \amalg I_-$ where I_+, I_- are disjoint subsets of $[q]$. A subset $A \subset [n]$ is polarized if it admits a decomposition $A = A_+ \amalg A_-$ into disjoint subsets. If A_+ and A_- are both of cardinal r , we say that A is balanced. We denote by Pol_{2r}^n the collection of balanced polarized subsets of $[n]$ of cardinal $2r$.

To any subset $I \in \mathcal{P}_r^{p,q}$, we associate the balanced polarized subset $\hat{I} = I \amalg I^o \in \text{Pol}_{2r}^n$ and the cohomology class $[\mathfrak{X}_{\hat{I}}] \in H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$. The proof of the following result is given in §4.6.

Theorem 4.1 *An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if*

$$(\star)_{I_{\pm}, J_{\pm}, K_{\pm}} \quad |x|_{I_+} + |y|_{J_+} + |z|_{K_+} \leq |x|_{I_-} + |y|_{J_-} + |z|_{K_-}$$

holds for any $r \in [q]$, for any $I, J, K \in \mathcal{P}_r^{p,q}$ such that

$$(14) \quad [\mathfrak{X}_{\hat{I}}] \odot_0 [\mathfrak{X}_{\hat{J}}] \odot_0 [\mathfrak{X}_{\hat{K}}] = [pt] \quad \text{in} \quad H^*(\mathbb{F}(r, n-r; n), \mathbb{Z}).$$

The result still holds if we replace (14) by the weaker condition $[\mathfrak{X}_{\hat{I}}] \odot_0 [\mathfrak{X}_{\hat{J}}] \odot_0 [\mathfrak{X}_{\hat{K}}] = \ell[pt]$ with $\ell \geq 1$.

The following examples are detailed in §4.4.

Example 4.2 *Let us consider the case $p \geq q = 1$. Thus $(a, b, c) \in (\mathbb{R}_+)^3$ belongs to $\text{Singular}(p, 1)$ if and only if the Weyl inequalities holds : $a + b \geq c$, $a + c \geq b$, and $b + c \geq a$.*

Example 4.3 *Let us consider the case $p \geq q = 2$. Thus $(a, b, c) \in (\mathbb{R}_{++}^2)^3$ belongs to $\text{Singular}(p, 2)$ if and only if the following 18 inequalities holds*

1. *the Weyl inequalities*

- $a_1 + b_1 \geq c_1$ (and 2 permutations),
- $a_1 + b_2 \geq c_2$ (and 5 permutations),

2. *the Lidskii inequalities*

- $a_1 + a_2 + b_1 + b_2 \geq c_1 + c_2$ (and 2 permutations),

3. *the signed Lidskii inequalities*

- $a_1 + a_2 + b_1 - b_2 \geq c_1 - c_2$ (and 5 permutations).

4.1 Multiplicative formulas in $H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$

In this section, we use some multiplicative formula obtained by Ressayre [22] and Richmond [25] for the structure constants in the cohomology of flag varieties to simplify condition (14). We are interested by their multiplicative formula in the case of two steps flag varieties $\mathbb{F}(r, n-r; n)$. The Schubert varieties of $\mathbb{F}(r, n-r; n)$ are parametrized by the set Pol_{2r}^n of balanced polarized subsets of $[n]$ of cardinal $2r$. If $A = A_+ \amalg A_- \in \text{Pol}_{2r}^n$, the subset $(A_+)^c \subset [n]$ is of cardinal $n-r$: $(A_+)^c = \{u_1 < \dots < u_{n-r}\}$. Since A_- is contained in $(A_+)^c$ we can define the following subsets of cardinal r : $A' = \{i \in [n-r], u_i \in A_-\}$ and $A'' = \{k \in [n-r], n-r+1-k \in A'\}$. Therefore, we can associate the cycles classes $[\mathfrak{X}_{A_+}] \in H^*(\mathbb{G}(r, n), \mathbb{Z})$ and $[\mathfrak{X}_{A''}] \in H^*(\mathbb{G}(r, n-r), \mathbb{Z})$ to any $A \in \text{Pol}_{2r}^n$.

The following result is a particular case of a general multiplicative formula obtained by Ressayre [22] and Richmond [25].

Proposition 4.4 *Let $A, B, C \in \text{Pol}_{2r}^n$. The relation $[\mathfrak{X}_A] \odot_0 [\mathfrak{X}_B] \odot_0 [\mathfrak{X}_C] = \ell[pt]$, with $\ell > 0$, holds in $H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$ if and only if we have*

1. $[\mathfrak{X}_{A_+}] \cdot [\mathfrak{X}_{B_+}] \cdot [\mathfrak{X}_{C_+}] = \ell'[pt]$ with $\ell' > 0$ in $H^*(\mathbb{G}(r, n), \mathbb{Z})$,
2. $[\mathfrak{X}_{A''}] \cdot [\mathfrak{X}_{B''}] \cdot [\mathfrak{X}_{C''}] = \ell''[pt]$ with $\ell'' > 0$ in $H^*(\mathbb{G}(r, n-r), \mathbb{Z})$,
3. $|A_+| + |B_+| + |C_+| = |A_-| + |B_-| + |C_-| + r(n-r)$.

Moreover we have $\ell = \ell' \ell''$.

When we work with $I, J, K \in \mathcal{P}_r^{p,q}$ and the associated balanced polarized subsets $\widehat{I} = I \amalg I^o$, $\widehat{J} = J \amalg J^o$, and $\widehat{K} = K \amalg K^o$, we see that

$$\begin{aligned} |\widehat{I}_+| + |\widehat{J}_+| + |\widehat{K}_+| &= |\widehat{I}_-| + |\widehat{J}_-| + |\widehat{K}_-| + r(n-r) \iff \\ |I| + |J| + |K| &= 2r(n-r) + \frac{3}{2}r(r+1) \iff \\ \text{codim}(\mathfrak{X}_I) + \text{codim}(\mathfrak{X}_J) + \text{codim}(\mathfrak{X}_K) &= r(n-r). \end{aligned}$$

Therefore condition 3. is a consequence of condition 1. in Proposition 4.4 when working with $\widehat{I}, \widehat{J}, \widehat{K} \in \text{Pol}_{2r}^n$.

If $X \in \mathcal{P}_r^{p,q}$, we define $\widetilde{X} \in \mathcal{P}_r^{n-r}$ as equal to $(\widehat{X})''$. Thanks to Proposition 4.4, we can state another version of Theorem 1.19.

Theorem 4.5 *An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ holds for any $r \in [q]$, for any $I, J, K \in \mathcal{P}_r^{p,q}$ satisfying both conditions :*

$$(15) \quad [\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = [pt] \quad \text{in} \quad H^*(\mathbb{G}(r, n), \mathbb{Z}),$$

$$(16) \quad [\mathfrak{X}_{\widetilde{I}}] \cdot [\mathfrak{X}_{\widetilde{J}}] \cdot [\mathfrak{X}_{\widetilde{K}}] = [pt] \quad \text{in} \quad H^*(\mathbb{G}(r, n-r), \mathbb{Z}).$$

The result is still true if we work with the weaker conditions : $[\mathfrak{X}_I] \cdot [\mathfrak{X}_J] \cdot [\mathfrak{X}_K] = \ell'[pt]$ and $[\mathfrak{X}_{\widetilde{I}}] \cdot [\mathfrak{X}_{\widetilde{J}}] \cdot [\mathfrak{X}_{\widetilde{K}}] = \ell''[pt]$ with $\ell', \ell'' \geq 1$.

4.2 Inequalities of Weyl, Lidskii and Thompson

The purpose of this section is to give some examples of triplets (I, J, K) satisfying (15) and (16) which will allow us to recover the classical inequalities of Weyl, Lidskii and Thompson.

We need to recall a standard feature of Littlewood-Richardson coefficients. Let $m \geq 2$ and $r \in [m]$. For any subset $X = \{x_1 < \dots < x_r\} \subset [q]$ of cardinal r , we associate several objects : the partition $\lambda(X) = (\lambda_1 \geq \dots \geq \lambda_r)$ where $\lambda_a = m - r + a - x_a$; the polynomial irreducible representation $V_{\lambda(X)}$ of $U(r)$ and the cycle class $[\mathfrak{X}_X] \in H^{2n(X)}(\mathbb{G}(r, m-r), \mathbb{Z})$,

where $n(X) = |\lambda(X)|$. A classical result of Lesieur [17] tell us that for any subsets $A, B, C \subset [m]$ of cardinal r , if $|\lambda(A)| + |\lambda(B)| + |\lambda(C)| = r(m - r)$ we have

$$[\mathfrak{X}_A] \cdot [\mathfrak{X}_B] \cdot [\mathfrak{X}_C] = c(A, B, C)[pt]$$

where $c(A, B, C) = \dim(V_{\lambda(A)} \otimes V_{\lambda(B)} \otimes V_{\lambda(C)} \otimes \det^{r-m})^{U(r)}$. Since $|\lambda(A)| + |\lambda(B)| + |\lambda(C)| = r(m - r)$ when $c(A, B, C) \neq 0$, we obtain the following equivalence for any $\ell \geq 1$:

$$[\mathfrak{X}_A] \cdot [\mathfrak{X}_B] \cdot [\mathfrak{X}_C] = \ell[pt] \iff \dim(V_{\lambda(A)} \otimes V_{\lambda(B)} \otimes V_{\lambda(C)} \otimes \det^{r-m})^{U(r)} = \ell.$$

Weyl inequalities

Let $m \geq 2$. To any element $a \in [m]$ we associate the Schubert variety $\mathfrak{X}_a = \overline{B_m \cdot (\mathbb{C}e_a)} \subset \mathbb{G}(1, m)$ of codimension $m - a$. Lesieur's result leads to the following simple criterion.

Lemma 4.6 *Let $i, j, k \in [m]$. Then $[\mathfrak{X}_i] \cdot [\mathfrak{X}_j] \cdot [\mathfrak{X}_k] = [pt]$ in $H^*(\mathbb{G}(1, m), \mathbb{Z})$ if and only if $i + j + k = 2m + 1$.*

Let $i, j \in [q]$ such that $i + j \leq q + 1$. Consider the following triplet of $\mathcal{P}_1^{p,q}$: $I = \emptyset \coprod \{n + 1 - i\}$, $J = \emptyset \coprod \{n + 1 - j\}$ and $K = \{i + j - 1\} \coprod \emptyset$. On one side, the cycles classes in $H^*(\mathbb{G}(1, n), \mathbb{Z})$ are $[\mathfrak{X}_I] = [\mathfrak{X}_{n+1-i}]$, $[\mathfrak{X}_J] = [\mathfrak{X}_{n+1-j}]$ and $[\mathfrak{X}_K] = [\mathfrak{X}_{i+j-1}]$. On the other side, the cycles classes in $H^*(\mathbb{G}(1, n - 1), \mathbb{Z})$ are $[\mathfrak{X}_{\tilde{I}}] = [\mathfrak{X}_{n-i}]$, $[\mathfrak{X}_{\tilde{J}}] = [\mathfrak{X}_{n-j}]$ and $[\mathfrak{X}_{\tilde{K}}] = [\mathfrak{X}_{i+j-1}]$. Thanks to the previous Lemma, we see that Conditions (15) and (16) are satisfied. The corresponding inequalities are the *Weyl inequalities* [28] :

$$\tau_{i+j-1}(A + B) \leq \tau_i(A) + \tau_j(B), \quad \forall A, B \in M_{p,q}(\mathbb{C}).$$

whenever $i + j \leq q + 1$.

Lidskii inequalities

Let us consider the element $X_r^n := \emptyset \coprod \{n - r + 1, \dots, n\} \in \mathcal{P}_r^{p,q}$: here $\tilde{X}_r^n = \{n - 2r + 1, \dots, n - r\} \in \mathcal{P}_r^{n-r}$. The Schubert varieties $\mathfrak{X}_{X_r^n}$ and $\mathfrak{X}_{\tilde{X}_r^n}$ are respectively equal to the whole manifolds $\mathbb{G}(r, n)$ and $\mathbb{G}(r, n - r)$, hence $[\mathfrak{X}_{X_r^n}] = 1$ and $[\mathfrak{X}_{\tilde{X}_r^n}] = 1$. Recall that the identity $[\mathfrak{X}_I] \cdot [\mathfrak{X}_J] = [pt]$ holds in $H^*(\mathbb{G}(r, n), \mathbb{Z})$ if and only if $J = I^o$.

Thus we obtain the following

Lemma 4.7 *Let $I \in \mathcal{P}_r^{p,q}$. The triplet (X_r^n, I, I^o) satisfies (15) and (16) if and only if $\tilde{I}^o = (\tilde{I})^o$.*

We leave it to the reader to verify that the identity $\tilde{I}^o = (\tilde{I})^o$ is satisfied in the following two cases:

1. $I = \{i_1 < \dots < i_r\} \coprod \emptyset$. The corresponding inequalities are the *Lidskii inequalities* [18]:

$$\sum_{k=1}^r \tau_{i_k}(A + B) \leq \sum_{k=1}^r \tau_{i_k}(B) + \sum_{k=1}^r \tau_k(A), \quad \forall A, B \in M_{p,q}(\mathbb{C}).$$

2. $I = \{i_1 < \dots < \widehat{i_s} < \dots < i_r\} \coprod \{n+1-i_s\}$. The corresponding inequalities will be referred to as the *signed Lidskii inequalities* :

$$\sum_{k \neq s} \tau_{i_k}(A+B) - \tau_{i_s}(A+B) \leq \sum_{k \neq s} \tau_{i_k}(B) - \tau_{i_s}(B) + \sum_{k=1}^r \tau_k(A), \quad \forall A, B \in M_{p,q}(\mathbb{C}).$$

Thompson inequalities

If $X = \{x_1 < \dots < x_r\} \subset [q]$ is a subset of cardinal $r \geq 1$, we define the partition $\mu_X = (x_r - r \geq \dots \geq x_1 - 1)$ and the irreducible representation V_{μ_X} of $U(r)$ with highest μ_X . Recall that $V_{\mu_X}^* = V_{\mu_X^*}$ where $\mu_X^* = (1 - x_1 \geq \dots \geq r - x_r)$.

Let $r \in [q]$. Let us consider three subsets of $[q]$ of cardinal r : $I_- := \{i_1 < \dots < i_r\}$, $J_- := \{j_1 < \dots < j_r\}$ and $K_+ = \{k_1 < \dots < k_r\}$. For any $p \geq q$, we define three elements of $\mathcal{P}_r^{p,q}$ as follows : $I^p = \emptyset \coprod \{p+q+1-i, i \in I_-\}$, $J^p = \emptyset \coprod \{p+q+1-j, j \in J_-\}$ and $K^p = K_+ \coprod \emptyset$.

Lemma 4.8 *The triplet (I^p, J^p, K^p) defined above satisfies (15) and (16) if and only if $\dim(V_{\mu_{I_-}} \otimes V_{\mu_{J_-}} \otimes V_{\mu_{K_+}}^*)^{U(r)} = 1$. If $(V_{\mu_{I_-}} \otimes V_{\mu_{J_-}} \otimes V_{\mu_{K_+}}^*)^{U(r)} \neq \{0\}$, then the weak versions of (15) and (16) are true.*

Proof : A direct computation gives : $\lambda(I^p) = \lambda(\widetilde{I^p}) = \mu_{I_-}$, $\lambda(J^p) = \lambda(\widetilde{J^p}) = \mu_{J_-}$ and $\lambda(K^p) = (n-r)1^r + \mu_{K_+}^*$, $\lambda(\widetilde{K^p}) = (n-2r)1^r + \mu_{K_+}^*$. Using Lesieur's result one sees that, for any $\ell \geq 1$, the following statements are equivalent :

- $[\mathfrak{X}_{I^p}] \cdot [\mathfrak{X}_{J^p}] \cdot [\mathfrak{X}_{K^p}] = \ell[pt]$ in $H^*(\mathbb{G}(r, n), \mathbb{Z})$,
- $[\mathfrak{X}_{\widetilde{I^p}}] \cdot [\mathfrak{X}_{\widetilde{J^p}}] \cdot [\mathfrak{X}_{\widetilde{K^p}}] = \ell[pt]$ in $H^*(\mathbb{G}(r, n-r), \mathbb{Z})$.
- $\dim(V_{\mu_{I_-}} \otimes V_{\mu_{J_-}} \otimes V_{\mu_{K_+}}^*)^{U(r)} = \ell$.

□

Corollary 4.9 *We have*

$$\sum_{k \in K_+} \tau_k(A+B) \leq \sum_{j \in J_-} \tau_j(B) + \sum_{i \in I_-} \tau_i(A), \quad \forall A, B \in M_{p,q}(\mathbb{C})$$

if $(V_{\mu_{I_-}} \otimes V_{\mu_{J_-}} \otimes V_{\mu_{K_+}}^*)^{U(r)} \neq \{0\}$.

Let us consider the situation where $k_\alpha = i_\alpha + j_\alpha - \alpha$, $\forall \alpha \in [r]$. These relations imply that $\mu_{K_+} = \mu_{I_-} + \mu_{J_-}$, and a standard fact of representation theory tells us that $\dim(V_{\mu_{I_-}} \otimes V_{\mu_{J_-}} \otimes V_{\mu_{I_-} + \mu_{J_-}}^*)^{U(r)} = 1$. We obtain here Thompson's inequalities [27]

$$\sum_{\alpha=1}^r \tau_{i_\alpha + j_\alpha - \alpha}(A+B) \leq \sum_{\alpha=1}^r \tau_{j_\alpha}(B) + \sum_{\alpha=1}^r \tau_{i_\alpha}(A), \quad \forall A, B \in M_{p,q}(\mathbb{C}),$$

which are valid for any subsets $\{i_1 < \dots < i_r\}$ and $\{j_1 < \dots < j_r\}$ satisfying $i_r + j_r - r \leq q$.

4.3 Horn conditions

In the previous sections we parameterized the inequality $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ in terms of subsets I, J, K leaving in $\mathcal{P}_r^{p,q}$. To simplify the presentation, we will choose a parameterization using directly the $I_{\pm}, J_{\pm}, K_{\pm}$.

The cardinal of a finite set Z is denoted by $\sharp Z$. Let start from a polarized subset $X_{\bullet} = X_+ \amalg X_-$ of $[q]$. To any $p \geq q$, we associate the subset $X_{\bullet}^p \subset [p+q]$ of cardinal $r = \sharp X_{\bullet}$ defined as $X_{\bullet}^p = X_+ \amalg \{p+q+1-x, x \in X_-\}$.

The complement of X_{\bullet}^p , $(X_{\bullet}^p)^c = \{u_1 < \dots < u_{p+q-r}\}$, contains $(X_{\bullet}^p)^o = \{p+q+1-x, x \in X_{\bullet}^p\}$. Hence we can define the following subsets of cardinal r : $(X_{\bullet}^p)' = \{i \in [p+q-r], u_i \in (X_{\bullet}^p)^o\}$ and

$$\tilde{X}_{\bullet}^p = \{k \in [p+q-r], p+q-r+1-k \in (X_{\bullet}^p)'\}.$$

Finally we associate the cycles classes $[\mathfrak{X}_{X_{\bullet}^p}] \in H^*(\mathbb{G}(r, p+q), \mathbb{Z})$ and $[\mathfrak{X}_{\tilde{X}_{\bullet}^p}] \in H^*(\mathbb{G}(r, p+q-r), \mathbb{Z})$ to any polarized subset $X_{\bullet} \subset [q]$ of cardinal $r \geq 1$, and to any $p \geq q$.

Recall that $\lambda(X_{\bullet}^p)$ (resp. $\lambda(\tilde{X}_{\bullet}^p)$) is the partition associated to the subset $X_{\bullet}^p \subset [p+q]$ (resp. $\tilde{X}_{\bullet}^p \subset [p+q-r]$). When $I_{\bullet}, J_{\bullet}, K_{\bullet}$ are three polarized subsets of $[q]$ of same cardinal r , the relation $[\mathfrak{X}_{I_{\bullet}^p}] \cdot [\mathfrak{X}_{J_{\bullet}^p}] \cdot [\mathfrak{X}_{K_{\bullet}^p}] = \ell[p\ell] \quad \ell \geq 1$, in $H^*(\mathbb{G}(r, p+q), \mathbb{Z})$ is equivalent to $(\lambda(I_{\bullet}^p), \lambda(J_{\bullet}^p), \lambda(K_{\bullet}^p) - (p+q-r)1^r) \in \text{Horn}(r)$. In the same way, the relation $[\mathfrak{X}_{\tilde{I}_{\bullet}^p}] \cdot [\mathfrak{X}_{\tilde{J}_{\bullet}^p}] \cdot [\mathfrak{X}_{\tilde{K}_{\bullet}^p}] = \ell[p\ell] \quad \ell \geq 1$, in $H^*(\mathbb{G}(r, p+q-r), \mathbb{Z})$ is equivalent to $(\lambda(\tilde{I}_{\bullet}^p), \lambda(\tilde{J}_{\bullet}^p), \lambda(\tilde{K}_{\bullet}^p) - (p+q-2r)1^r) \in \text{Horn}(r)$.

Let us rewrite a weak version of Theorem 4.5 with these new notations.

Theorem 4.10 *An element $(x, y, z) \in (\mathbb{R}_{++}^q)^3$ belongs to $\text{Singular}(p, q)$ if and only if $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ holds for any triplet $I_{\bullet}, J_{\bullet}, K_{\bullet}$ of polarized subsets of $[q]$ satisfying the following conditions :*

1. $\sharp I_{\bullet} = \sharp J_{\bullet} = \sharp K_{\bullet} = r \in [q]$,
2. $(\lambda(I_{\bullet}^p), \lambda(J_{\bullet}^p), \lambda(K_{\bullet}^p) - (p+q-r)1^r) \in \text{Horn}(r)$,
3. $(\lambda(\tilde{I}_{\bullet}^p), \lambda(\tilde{J}_{\bullet}^p), \lambda(\tilde{K}_{\bullet}^p) - (p+q-2r)1^r) \in \text{Horn}(r)$.

Conditions 2. and 3. implies respectively that $|\lambda(I_{\bullet}^p)| + |\lambda(J_{\bullet}^p)| + |\lambda(K_{\bullet}^p)| = (p+q-r)r$ and $|\lambda(\tilde{I}_{\bullet}^p)| + |\lambda(\tilde{J}_{\bullet}^p)| + |\lambda(\tilde{K}_{\bullet}^p)| = (p+q-2r)r$. We end this section by calculating precisely these two relations.

Definition 4.11 *For a polarized subset $X_{\bullet} \subset [q]$, we denote by $\delta_{X_{\bullet}}$ the cardinal of the set $\{(a, b) \in X_+ \times X_-, a < b\}$.*

Let $r \in [q]$. For any $0 \leq \alpha \leq r$, we denote by $1_{\alpha}^r = (1, \dots, 1, 0, \dots, 0) \in \mathbb{R}^r$ the vector where 1 appears α times and 0 appears $r - \alpha$ times.

Lemma 4.12 *Let $X_{\bullet} \subset [q]$ be a polarized subset of cardinal $r \geq 1$.*

1. For any $p \geq q$, we have

$$(17) \quad |\lambda(X_{\bullet}^p)| = |X_-| - |X_+| + (p+q+1)\sharp X_+ - \frac{r(r+1)}{2},$$

$$(18) \quad |\lambda(\tilde{X}_{\bullet}^p)| = |X_-| - |X_+| + (p+q+1)\sharp X_+ - \frac{r(r+1)}{2} - (\sharp X_+)^2 - 2\delta_{X_{\bullet}}.$$

$$\text{Hence } |\lambda(X_{\bullet}^p)| - |\lambda(\tilde{X}_{\bullet}^p)| = (\sharp X_+)^2 + 2\delta_{X_{\bullet}}.$$

2. For any $p' \geq p \geq q$, we have

$$\lambda(X_{\bullet}^{p'}) - \lambda(X_{\bullet}^p) = (p' - p)1_{\sharp X_+}^r \quad \text{and} \quad \lambda(\tilde{X}_{\bullet}^{p'}) - \lambda(\tilde{X}_{\bullet}^p) = (p' - p)1_{\sharp X_+}^r.$$

Proof : Let us prove the first point. A direct computation gives $|X_{\bullet}^p| = |X_+| - |X_-| + (p+q+1)(r - \sharp X_+)$ and then, as $|\lambda(X_{\bullet}^p)| = (p+q-r)r + \frac{r(r+1)}{2} - |X_{\bullet}^p|$, we obtain (17).

Recall that $(X_{\bullet}^p)' \subset [p+q-r]$ is defined as the collection of the ranks of the elements of $\{p+q+1-x, x \in X_{\bullet}^p\}$ in $[n] - X_{\bullet}^p$. A short computation gives that

$$(19) \quad |(X_{\bullet}^p)'| = |X_-| - |X_+| - 2\delta_{X_{\bullet}} + (p+q+1 - \sharp X_+)\sharp X_+.$$

and $|\tilde{X}_{\bullet}^p| = (p+q-r+1)r - |(X_{\bullet}^p)'|$. As $|\lambda(\tilde{X}_{\bullet}^p)| = (p+q-2r)r + \frac{r(r+1)}{2} - |\tilde{X}_{\bullet}^p|$, we obtain

$$\begin{aligned} |\lambda(\tilde{X}_{\bullet}^p)| &= (p+q-2r)r + \frac{r(r+1)}{2} - (p+q-r+1)r + |(X_{\bullet}^p)'| \\ &= |(X_{\bullet}^p)'| - \frac{r(r+1)}{2} \\ &= |X_-| - |X_+| + (p+q+1)\sharp X_+ - \frac{r(r+1)}{2} - (\sharp X_+)^2 - 2\delta_{X_{\bullet}}. \end{aligned}$$

The last equality is a consequence of (19).

The second point, which can be proved by direct calculation, is left to the reader. \square

Corollary 4.13 *The inequality $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ appears in the description of the convex cone $\text{Singular}(p, q)$ only if the triplet $I_{\bullet}, J_{\bullet}, K_{\bullet}$ of polarized subsets of $[q]$ satisfies the relations :*

$$(C1) \quad \sharp I_{\bullet} = \sharp J_{\bullet} = \sharp K_{\bullet} = r \in [q],$$

$$(C2) \quad |I_+| + |J_+| + |K_+| - (|I_-| + |J_-| + |K_-|) + \frac{r(r+1)}{2} = (p+q+1)(\sharp I_+ + \sharp J_+ + \sharp K_+ - r),$$

$$(C3) \quad (\sharp I_+)^2 + (\sharp J_+)^2 + (\sharp K_+)^2 + 2(\delta_{I_{\bullet}} + \delta_{J_{\bullet}} + \delta_{K_{\bullet}}) = r^2.$$

4.4 Examples : Singular($p, 1$), Singular($p, 2$) and Singular($3, 3$)

In the next examples, the notation $X_\bullet = \{i_1^\epsilon, \dots, i_r^\epsilon\}$ means that $X_+ = \{i_k^\epsilon, \epsilon = +\}$ and $X_- = \{i_k^\epsilon, \epsilon = -\}$.

Example 4.14 Let $p \geq q = 1$. Thus $(a, b, c) \in (\mathbb{R}_+)^3$ belongs to Singular($p, 1$) if and only if the Weyl inequalities holds : $a + b \geq c$, $a + c \geq b$, and $b + c \geq a$.

Proof : Let $I_\bullet, J_\bullet, K_\bullet \subset [1]$ satisfying the relations of Theorem 4.10. If we use relation (C3) of Corollary 4.13, we obtain that $(\#I_+)^2 + (\#J_+)^2 + (\#K_+)^2 = 1$. Up to a permutation, we must have $I_\bullet = J_\bullet = \{1^-\}$ and $K_\bullet = \{1^+\}$. This case corresponds to the Weyl inequality $a + b \geq c$. \square

Example 4.15 Let $p \geq q = 2$. The element $(a, b, c) \in (\mathbb{R}_{++}^2)^3$ belongs to Singular($p, 2$) if and only if the following 18 inequalities holds

1. the Weyl inequalities

- $a_1 + b_1 \geq c_1$ (and 2 permutations),
- $a_1 + b_2 \geq c_2$ (and 5 permutations),

2. the Lidskii inequalities

- $a_1 + a_2 + b_1 + b_2 \geq c_1 + c_2$ (and 2 permutations),

3. the signed Lidskii inequalities

- $a_1 + a_2 + b_1 - b_2 \geq c_1 - c_2$ (and 5 permutations).

Proof : Let $I_\bullet, J_\bullet, K_\bullet \subset [2]$ satisfying the relations of Theorem 4.10. Let $r := \#I_\bullet = \#J_\bullet = \#K_\bullet$.

First case : $r = 1$. Relation (C3) of Corollary 4.13 gives $(\#I_+)^2 + (\#J_+)^2 + (\#K_+)^2 = 1$. Up to a permutation, we have $I_\bullet = \{i^-\}$, $J_\bullet = \{j^-\}$, $K_\bullet = \{k^+\}$ with $i, j, k \in \{1, 2\}$. Relation (C2) of Corollary 4.13 imposes that $k = i + j - 1$, thus the triplet (i, j, k) belongs to $(1, 1, 1)$, $(1, 2, 2)$ and $(2, 1, 2)$. All this cases corresponds to the Weyl inequalities : $a_1 + b_1 \geq c_1$, $a_1 + b_2 \geq c_2$, and $a_2 + b_1 \geq c_2$.

Second case : $r = 2$. Here relation (C3) becomes

$$(\#I_+)^2 + (\#J_+)^2 + (\#K_+)^2 + 2(\delta_{I_\pm} + \delta_{J_\pm} + \delta_{K_\pm}) = 4.$$

Up to a permutation, we can suppose that $\#I_+ \leq \#J_+ \leq \#K_+$. There are two possibilities :

- $(\#I_+, \#J_+, \#K_+) = (0, 0, 2)$. Thus $I_\bullet = J_\bullet = \{1^-, 2^-\}$ and $K_\bullet = \{1^+, 2^+\}$. This case corresponds to the Lidskii inequality $a_1 + a_2 + b_1 + b_2 \geq c_1 + c_2$.

- $(\#I_+, \#J_+, \#K_+) = (0, 1, 1)$. Up to a permutation, we have $\delta_{K_\bullet} = 1$ and $\delta_{J_\bullet} = 0$. Here $I_\bullet = \{1^-, 2^-\}$, $J_\bullet = \{1^-, 2^+\}$ and $K_\bullet = \{1^+, 2^-\}$. This situation corresponds to the signed Lidskii inequality $a_1 + a_2 + b_1 - b_2 \geq c_1 - c_2$. \square

Example 4.16 *An element $(a, b, c) \in (\mathbb{R}_{++}^3)^3$ belongs to $\text{Singular}(3, 3)$ if and only if the following 87 inequalities holds :*

1. *the Weyl inequalities*

- $a_1 + b_1 \geq c_1$ (and 2 permutations),
- $a_1 + b_2 \geq c_2$ (and 5 permutations),
- $a_2 + b_2 \geq c_3$ (and 2 permutations),
- $a_1 + b_3 \geq c_3$ (and 5 permutations).

2. *the Lidskii inequalities*

- $a_1 + a_2 + b_1 + b_2 \geq c_1 + c_2$ (and 2 permutations),
- $a_1 + a_2 + b_1 + b_3 \geq c_1 + c_3$ (and 5 permutations),
- $a_1 + a_2 + b_2 + b_3 \geq c_2 + c_3$ (and 5 permutations),
- $a_1 + a_2 + a_3 + b_1 + b_2 + b_3 \geq c_1 + c_2 + c_3$ (and 2 permutations).

3. *the signed Lidskii inequalities*

- $a_1 + a_2 + b_1 - b_2 \geq c_1 - c_2$ (and 5 permutations),
- $a_1 + a_2 + b_1 - b_3 \geq c_1 - c_3$ (and 5 permutations),
- $a_1 + a_2 + b_2 - b_3 \geq c_2 - c_3$ (and 5 permutations),
- $a_1 + a_2 + a_3 + b_1 + b_2 - b_3 \geq c_1 + c_2 - c_3$ (and 5 permutations),
- $a_1 + a_2 + a_3 + b_1 - b_2 + b_3 \geq c_1 - c_2 + c_3$ (and 5 permutations),
- $a_1 + a_2 + a_3 - b_1 + b_2 + b_3 \geq -c_1 + c_2 + c_3$ (and 5 permutations).

4. *others inequalities*

- $a_1 + a_3 + b_1 + b_3 \geq c_2 + c_3$ (and 2 permutations),
- $a_1 + a_3 + b_1 - b_3 \geq c_2 - c_3$ (and 5 permutations),
- $(a_1 + a_2 - a_3) + (b_1 - b_2 + b_3) + (-c_1 + c_2 + c_3) \geq 0$ (and 5 permutations).

Proof : Let $I_\bullet, J_\bullet, K_\bullet \subset [3]$ satisfying the relations of Theorem 4.10. Let $r := \#J_\bullet = \#K_\bullet = \#K_\bullet$.

First case : $r = 1$.

We obtain here the Weyl inequalities $a_i + b_j \geq c_{i+j-1}$ (up to permutations).

Second case : $r = 2$.

Let us use the relation $(\#I_+)^2 + (\#J_+)^2 + (\#K_+)^2 + 2(\delta_{I_\pm} + \delta_{J_\pm} + \delta_{K_\pm}) = 4$. Up to a permutation, we can suppose that $\#I_+ \leq \#J_+ \leq \#K_+$. There are two possibilities :

- $(\#I_+, \#J_+, \#K_+) = (0, 0, 2)$. Here we have $I_+ = J_+ = K_- = \emptyset$. Relation (C3) of Corollary 4.13 gives $|I_-| + |J_-| = |K_+| + 3$. Knowing that $|I_-|, |J_-|, |K_+| \in \{3, 4, 5\}$, we obtain few cases related to Lidskii inequalities:

1. $(|I_-|, |J_-|, |K_+|) = (3, 3, 3) : a_1 + a_2 + b_1 + b_2 \geq c_1 + c_2$,
2. $(|I_-|, |J_-|, |K_+|) = (3, 4, 4)$ or $(4, 3, 4) : a_1 + a_2 + b_1 + b_3 \geq c_1 + c_3$ or $a_1 + a_3 + b_1 + b_2 \geq c_1 + c_3$,
3. $(|I_-|, |J_-|, |K_+|) = (3, 5, 5)$ or $(5, 3, 5) : a_1 + a_2 + b_2 + b_3 \geq c_2 + c_3$ or $a_2 + a_3 + b_1 + b_2 \geq c_2 + c_3$.

The only case left is $(|I_-|, |J_-|, |K_+|) = (4, 4, 5)$. Here $I_\bullet = J_\bullet = \{1^-, 3^-\}$ and $K_\bullet = \{2^+, 3^+\}$. A small computation shows that the Horn conditions are satisfied in this case. We obtain the inequality $a_1 + a_3 + b_1 + b_3 \geq c_2 + c_3$.

- $(\#I_+, \#J_+, \#K_+) = (0, 1, 1)$. Up to a permutation, we can suppose that $\delta_{K_\bullet} = 1$ and $\delta_{J_\bullet} = 0$. Thus $I_\bullet = \{i_1^- < i_2^-\}$, $J_\bullet = \{j_1^- < j_2^+\}$ and $K_\bullet = \{k_1^+ < k_2^-\}$. Relation (C2) of Corollary 4.13 gives $\underbrace{i_1^- + i_2^-}_{\alpha} + \underbrace{j_1^- + k_1^-}_{\beta} = \underbrace{j_2^+ + k_1^+}_{\gamma} + 3$.

Knowing that $\alpha, \beta, \gamma \in \{3, 4, 5\}$, we obtain

1. $(\alpha, \beta, \gamma) = (3, 3, 3), (3, 4, 4)$, or $(3, 5, 5)$. It gives us the signed Lidskii inequalities.
2. $(\alpha, \beta, \gamma) = (4, 3, 4)$, or $(5, 3, 5)$. No solutions.
3. $(\alpha, \beta, \gamma) = (4, 4, 5)$. Here $I_\bullet = \{1^-, 3^-\}$, $J_\bullet = \{1^-, 3^+\}$ and $K_\bullet = \{2^+, 3^-\}$. A small computation shows that the Horn conditions are satisfied in this case. It corresponds to the inequality $a_1 + a_3 + b_1 - b_3 \geq c_2 - c_3$.

Last case : $r = 3$.

Let start with the relation (C3) : $(\#I_+)^2 + (\#J_+)^2 + (\#K_+)^2 + 2(\delta_{I_\pm} + \delta_{J_\pm} + \delta_{K_\pm}) = 9$. Up to a permutation, we can suppose that $\#I_+ \leq \#J_+ \leq \#K_+$. First, let us list the triples $(\#I_+ \leq \#J_+ \leq \#K_+)$ that cannot satisfy (C3) : $(0, 0, 0)$; $(0, 0, 1)$; $(0, 1, 1)$; $(0, 0, 2)$; $(1, 1, 2)$; $(0, 2, 2)$; $(2, 2, 2)$. There are four cases left :

1. $(\#I_+, \#J_+, \#K_+) = (0, 0, 3)$. Here $I_\bullet = J_\bullet = \{1^-, 2^-, 3^-\}$ and $K_\bullet = \{1^+, 2^+, 3^+\}$. We obtain the Lidskii inequality $a_1 + a_2 + a_3 + b_1 + b_2 + b_3 \geq c_1 + c_2 + c_3$.
2. $(\#I_+, \#J_+, \#K_+) = (0, 1, 2)$. We must have $\delta_{J_\bullet} + \delta_{K_\bullet} = 2$, so we have three possibilities
 - a) $(\delta_{J_\bullet}, \delta_{K_\bullet}) = (0, 2) : \text{ here } I_\bullet = \{1^-, 2^-, 3^-\}$, $J_\bullet = \{1^-, 2^-, 3^+\}$ and $K_\bullet = \{1^+, 2^+, 3^-\}$. We obtain the signed Lidskii inequality $a_1 + a_2 + a_3 + b_1 + b_2 - b_3 \geq c_1 + c_2 - c_3$.

- b) $(\delta_{J_\bullet}, \delta_{K_\bullet}) = (1, 1)$: here $I_\bullet = \{1^-, 2^-, 3^-\}$, $J_\bullet = \{1^-, 2^+, 3^-\}$ and $K_\bullet = \{1^+, 2^-, 3^+\}$. We obtain the signed Lidskii inequality $a_1 + a_2 + a_3 + b_1 - b_2 + b_3 \geq c_1 - c_2 + c_3$.
- c) $(\delta_{J_\bullet}, \delta_{K_\bullet}) = (2, 0)$: here $I_\bullet = \{1^-, 2^-, 3^-\}$, $J_\bullet = \{1^+, 2^-, 3^-\}$ and $K_\bullet = \{1^-, 2^+, 3^+\}$. We obtain the signed Lidskii inequality $a_1 + a_2 + a_3 - b_1 + b_2 + b_3 \geq -c_1 + c_2 + c_3$.
3. $(\#I_+, \#J_+, \#K_+) = (1, 2, 2)$. We must have $\delta_{I_\bullet} = \delta_{J_\bullet} = \delta_{K_\bullet} = 0$, thus $I_\bullet = \{1^-, 2^-, 3^+\}$, $J_\bullet = \{1^-, 2^+, 3^+\}$ and $K_\bullet = \{1^-, 2^+, 3^+\}$. The associated partitions are $\lambda(I_\bullet^3) = (1, 0, 0)$ and $\lambda(J_\bullet^3) = \lambda(K_\bullet^3) = (2, 2, 0)$. As $((1, 0, 0), (2, 2, 0), (-1, -1, -3))$ does not belongs to $\text{Horn}(3)$, this triplet $(I_\bullet, J_\bullet, K_\bullet)$ does not satisfies the conditions of Theorem 4.10.
4. $(\#I_+, \#J_+, \#K_+) = (1, 1, 1)$. We must have $\delta_{I_\bullet} + \delta_{J_\bullet} + \delta_{K_\bullet} = 3$. Up to a permutation, two situations arise
- a) $\delta_{I_\bullet} = \delta_{J_\bullet} = \delta_{K_\bullet} = 1$: here $I_\bullet = J_\bullet = K_\bullet = \{1^-, 2^+, 3^-\}$. The corresponding partition is $\mu = (2, 1, 0)$, and one checks that $(\mu, \mu, \mu - 3 \cdot 1^3) \in \text{Horn}(3)$. Thus this triplet satisfies the conditions of Theorem 4.10. The corresponding inequality,
- $$a_1 + a_3 + b_1 + b_3 + c_1 + c_3 \geq a_2 + b_2 + c_2,$$
- is nevertheless redundant since it is a consequence of the inequality $x_1 \geq x_2$ satisfied by a, b, c .
- b) $\delta_{I_\bullet} = 0, \delta_{J_\bullet} = 1$ and $\delta_{K_\bullet} = 2$: here $I_\bullet = \{1^-, 2^-, 3^+\}$, $J_\bullet = \{1^-, 2^+, 3^-\}$ and $K_\bullet = \{1^+, 2^-, 3^-\}$. The corresponding partition are $\lambda = \lambda(I_\bullet^3) = (1, 0, 0)$, $\mu = \lambda(J_\bullet^3) = (2, 1, 0)$ and $\nu = \lambda(K_\bullet^3) = (3, 1, 1)$. One checks that $(\lambda, \mu, \nu - 3 \cdot 1^3) \in \text{Horn}(3)$, thus the triplet $(I_\bullet, J_\bullet, K_\bullet)$ satisfies the conditions of Theorem 4.10. The corresponding inequality is $a_1 + a_2 + b_1 + b_3 + c_2 + c_3 \geq a_3 + b_2 + c_1$.

Remark 4.17 *The case where $I_\bullet, J_\bullet, K_\bullet$ are all equal to $A_\bullet = \{1^-, 2^+, 3^-\}$ yields the example of a triplet satisfying the Horn conditions but giving a redundant inequality. This redundancy phenomenon can be explained as the triplet $(A_\bullet, A_\bullet, A_\bullet)$ does not satisfy the cohomological conditions of Theorem 4.5 : one has $[\mathfrak{X}_{A_\bullet^3}] \cdot [\mathfrak{X}_{A_\bullet^3}] \cdot [\mathfrak{X}_{A_\bullet^3}] = 2[pt]$ in $H^*(\mathbb{G}(3, 6), \mathbb{Z})$ (see Example 4.2.1 in [4]).*

4.5 The convex cone $\text{Singular}(\infty, q)$

Let $p \geq q$. To any matrix $X \in M_{p,q}(\mathbb{C})$, we associate

$$X^\vee = \begin{pmatrix} X \\ 0 \dots 0 \end{pmatrix} \in M_{p+1,q}(\mathbb{C})$$

It is immediate to see that X and X^\vee have the same singular values : $\tau(X) = \tau(X^\vee)$ for any $X \in M_{p,q}(\mathbb{C})$. From this basic consideration, we see that if $(\tau(A), \tau(B), \tau(A +$

$B)) \in \text{Singular}(p, q)$ then $(\tau(A), \tau(B), \tau(A + B)) = (\tau(A^\vee), \tau(B^\vee), \tau(A^\vee + B^\vee))$ belongs to $\text{Singular}(p + 1, q)$. Hence, $\text{Singular}(p, q) \subset \text{Singular}(p + 1, q)$ for any $p \geq q$.

This leads us to define the following convex cone

$$\text{Singular}(\infty, q) = \bigcup_{p \geq q} \text{Singular}(p, q).$$

Let us denote by \mathcal{E}_q the set formed by all inequalities

$$(\star)_{I_\pm, J_\pm, K_\pm} : \quad |x|_{I_+} + |y|_{J_+} + |z|_{K_+} \leq |x|_{I_-} + |y|_{J_-} + |z|_{K_-}, \quad x, y, z \in \mathbb{R}^q.$$

associated to triplet $(I_\bullet, J_\bullet, K_\bullet)$ of polarized subsets of $[q]$ of same cardinal.

Definition 4.18 *Let us denote by $\mathcal{E}_{p,q} \subset \mathcal{E}_q$ the set of inequalities associated to triplet $(I_\bullet, J_\bullet, K_\bullet)$ satisfying the Horn conditions of Theorem 4.10. Let $\mathcal{E}_{p,q}^{\min} \subset \mathcal{E}_{p,q}$ be the minimal system of inequalities describing the convex cone $\text{Singular}(p, q)$.*

An inequality $(\star)_{I_\pm, J_\pm, K_\pm}$ is called regular if there are twice as many terms on the right side as on the left side, e.g. $\#I_- + \#J_- + \#K_- = 2(\#I_+ + \#J_+ + \#K_+)$.

We start with a basic observation.

Lemma 4.19 *There exists $p_o \geq q$, such that $\text{Singular}(\infty, q) = \text{Singular}(p_o, q)$. In other words, $\text{Singular}(p, q) = \text{Singular}(p + 1, q)$ for any $p \geq p_o$.*

Proof : Thanks to Theorem 4.10, we know that for any $p \geq q$ the convex cone $\text{Singular}(p, q)$ is determined by a finite system of inequalities $\mathcal{E}_{p,q} \subset \mathcal{E}_q$. As \mathcal{E}_q is finite, there exists $\mathcal{R} \subset \mathcal{E}_q$ such that $\mathbb{N}(\mathcal{R}) := \{p \geq q, \mathcal{E}_{p,q} = \mathcal{R}\}$ is infinite. Take $p_o = \inf \mathbb{N}(\mathcal{R})$. Let us verify that $\text{Singular}(\infty, q) \subset \text{Singular}(p_o, q)$: as the other inclusion $\text{Singular}(p_o, q) \subset \text{Singular}(\infty, q)$ is obvious, the proof will be completed. Let $x \in \text{Singular}(\infty, q)$: there exists $p \geq q$ such that $x \in \text{Singular}(p, q)$. As $\mathbb{N}(\mathcal{R})$ is infinite, there exists $p' \in \mathbb{N}(\mathcal{R})$ such that $p' \geq p$. It follows that $x \in \text{Singular}(p, q) \subset \text{Singular}(p', q) = \text{Singular}(p_o, q)$. \square

We have now a nice characterization.

Proposition 4.20 *The identity $\text{Singular}(\infty, q) = \text{Singular}(p, q)$ holds if and only if all the inequalities of $\mathcal{E}_{p,q}^{\min}$ are regular.*

Proof : Let $p \geq q$ such that $\text{Singular}(\infty, q) = \text{Singular}(p, q)$. It means that $\mathcal{E}_{p',q}^{\min} = \mathcal{E}_{p,q}^{\min}$ for any $p' > p$. Let $(I_\bullet, J_\bullet, K_\bullet)$ such that $(\star)_{I_\pm, J_\pm, K_\pm} \in \mathcal{E}_{p,q}^{\min}$. Corollary 4.13 tells us that

$$(C2) \quad |I_+| + |J_+| + |K_+| - (|I_-| + |J_-| + |K_-|) + \frac{r(r+1)}{2} = (p+q+1)(\#I_+ + \#J_+ + \#K_+ - r).$$

Notice that $(\star)_{I_\pm, J_\pm, K_\pm}$ is a *regular* if and only if $\#I_+ + \#J_+ + \#K_+ - r = 0$. Hence, if $(\star)_{I_\pm, J_\pm, K_\pm}$ is not regular, the identity (C2) does not hold anymore when we change p in

$p' > p$. It implies that $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}} \notin \mathcal{E}_{p', q}$ for any $p' > p$. It is in contradiction with the fact that $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}} \in \mathcal{E}_{p', q}^{\min}$.

Suppose now that all the inequalities of $\mathcal{E}_{p, q}^{\min}$ are regular. In order to prove that $\text{Singular}(\infty, q)$ coincides with $\text{Singular}(p, q)$, it is sufficient to show that $\mathcal{E}_{p, q}^{\min} \subset \mathcal{E}_{p', q}$ for any $p' \geq p$. Let $(I_{\bullet}, J_{\bullet}, K_{\bullet})$ such that $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}} \in \mathcal{E}_{p, q}^{\min}$. By definition, we have

1. $\#I_{\bullet} = \#J_{\bullet} = \#K_{\bullet} = r \in [q]$,
2. $\#I_{+} + \#J_{+} + \#K_{+} = r$,
3. $\Lambda^p := (\lambda(I_{\bullet}^p), \lambda(J_{\bullet}^p), \lambda(K_{\bullet}^p)) - (p + q - r)1^r \in \text{Horn}(r)$,
4. $\tilde{\Lambda}^p := (\lambda(\tilde{I}_{\bullet}^p), \lambda(\tilde{J}_{\bullet}^p), \lambda(\tilde{K}_{\bullet}^p)) - (p + q - 2r)1^r \in \text{Horn}(r)$.

Identity 2. is due to our assumption that $(\star)_{I_{\pm}, J_{\pm}, K_{\pm}}$ is a regular inequality.

Our goal is achieved if we show that $\Lambda^{p'}$ and $\tilde{\Lambda}^{p'}$ belong to $\text{Horn}(r)$ for any $p' > p$. If $0 \leq \alpha \leq r$, we denote by $1_{\alpha}^r = (1, \dots, 1, 0, \dots, 0) \in \mathbb{R}^r$ the element where 1 appears α -times. We have seen in Lemma 4.12 that for any polarized subset $X_{\bullet} \subset [q]$ of cardinal $r \geq 1$ we have $\lambda(X_{\bullet}^{p'}) - \lambda(X_{\bullet}^p) = \lambda(\tilde{X}_{\bullet}^{p'}) - \lambda(\tilde{X}_{\bullet}^p) = (p' - p)1_{\#X_{+}}^r$, $\forall p' \geq p$. This leads to the relations

$$\Lambda^{p'} - \Lambda^p = \tilde{\Lambda}^{p'} - \tilde{\Lambda}^p = (p' - p) \left(1_{\#I_{+}}^r, 1_{\#J_{+}}^r, 1_{\#K_{+}}^r - 1^r \right), \quad \forall p' \geq p.$$

Now it is an easy matter to check that $(1_{\#I_{+}}^r, 1_{\#J_{+}}^r, 1_{\#K_{+}}^r - 1^r) \in \text{Horn}(r)$ since $\#I_{+} + \#J_{+} + \#K_{+} = r$. Using the fact that $\text{Horn}(r)$ is a convex cone, we can conclude that $\Lambda^{p'} = \Lambda^p + (p' - p)(1_{\#I_{+}}^r, 1_{\#J_{+}}^r, 1_{\#K_{+}}^r - 1^r)$ and $\tilde{\Lambda}^{p'} = \tilde{\Lambda}^p + (p' - p)(1_{\#I_{+}}^r, 1_{\#J_{+}}^r, 1_{\#K_{+}}^r - 1^r)$ belongs to $\text{Horn}(r)$ for any $p' \geq p$. \square

Corollary 4.21 *We have the following identities :*

- $\text{Singular}(\infty, 1) = \text{Singular}(1, 1)$,
- $\text{Singular}(\infty, 2) = \text{Singular}(2, 2)$,
- $\text{Singular}(\infty, 3) = \text{Singular}(3, 3)$.

Proof : The first two relations follow from the computations done in Examples 4.14 and 4.15. In Example 4.16, we have proved that $\text{Singular}(3, 3)$ is determined by a system of *regular* inequalities. The relation $\text{Singular}(\infty, 3) = \text{Singular}(3, 3)$ is then a consequence of the previous proposition. \square

4.6 Proof of Theorem 1.19

We work with the conjugate linear involution $\sigma_{p,q}(g) = I_{p,q}(g^*)^{-1}I_{p,q}$ on $GL_n(\mathbb{C})$, where $I_{p,q}$ is the diagonal matrix $\text{Diag}(I_p, -I_q)$. Let $U_{\mathbb{C}} = GL_n(\mathbb{C})$ embedded diagonally in $\tilde{U}_{\mathbb{C}} = GL_n(\mathbb{C}) \times GL_n(\mathbb{C})$. We define on $\tilde{U}_{\mathbb{C}}$ the conjugate linear involution $\sigma_{p,q}(g_1, g_2) = (\sigma_{p,q}(g_1), \sigma_{p,q}(g_2))$.

We see then that $G = U(p, q)$ and $\tilde{G} = G \times G$ are the subgroups fixed by $\sigma_{p,q}$. The Cartan decomposition of the Lie algebra of G is $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ where $\mathfrak{k} = \mathfrak{u}(p) \times \mathfrak{u}(q)$ and $\mathfrak{p} \simeq M_{p,q}(\mathbb{C})$ as a $U(p) \times U(q)$ -module.

Let $\mathfrak{t} \subset \mathfrak{u}(n)$ be the maximal torus composed by all the matrices of the form

$$i \begin{pmatrix} D(a) & 0 & M(c) \\ 0 & D(b) & 0 \\ M(\tilde{c}) & 0 & D(\tilde{a}) \end{pmatrix}, \quad (a, b, c) \in \mathbb{R}^q \times \mathbb{R}^{p-q} \times \mathbb{R}^q.$$

Here $M(c) = \begin{pmatrix} 0 & \cdots & c_1 \\ \vdots & \ddots & \vdots \\ c_q & \cdots & 0 \end{pmatrix}$ and $D(a) = \begin{pmatrix} a_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & a_q \end{pmatrix}$. We denoted $\tilde{x} = (x_q, \dots, x_1)$ for any $x = (x_1, \dots, x_q) \in \mathbb{R}^q$.

We notice that \mathfrak{t} is invariant under the involution $\sigma_{p,q}$, and that $\mathfrak{a} = i\mathfrak{t}^{-\sigma_{p,q}}$ is the maximal abelian subspace of \mathfrak{p} consisting of all matrices of the form

$$(20) \quad X(c) := \begin{pmatrix} 0 & 0 & M(c) \\ 0 & 0 & 0 \\ M(\tilde{c}) & 0 & 0 \end{pmatrix}, \quad c \in \mathbb{R}^q.$$

Let f_i be the member of \mathfrak{a}^* whose value on the matrix $X(a)$ is a_i . The set $\Sigma(\mathfrak{g})$ of restricted roots includes all linear functionals $\pm f_i \pm f_j$ with $i \neq j$ and $\pm 2f_i$ for all i . Also the $\pm f_i$ are restricted roots if $p \neq q$ (see Chapter 6 in [13]). Thus we can choose the following Weyl chamber

$$\mathfrak{a}_+ := \{X(a), a_1 \geq a_2 \geq \cdots \geq a_q \geq 0\}.$$

In this case the restricted Weyl group $W_{\mathfrak{a}}$ is the semi-direct product $\mathfrak{S}_q \rtimes \{1, -1\}^q$ where \mathfrak{S}_q acts on \mathbb{R}^q by permuting the indices and $\{1, -1\}^q = \{\epsilon = (\epsilon_1, \dots, \epsilon_q) \mid \epsilon_k = \pm 1\}$ acts by sign changes: $\epsilon \cdot a = (\epsilon_1 a_1, \dots, \epsilon_q a_q)$.

We start by determining the admissible elements.

Lemma 4.22 *Let $G = U(p, q)$ and $\tilde{G} = G \times G$. The admissible elements relatively to the $\tilde{G} \times G$ -action on \tilde{G} are⁴ of the form $(w_1\gamma_r, w_2\gamma_r, \gamma_3\gamma_r)$ where $w_i \in W_{\mathfrak{a}}, \forall i$ and*

$$\gamma_r := \begin{pmatrix} 0 & 0 & M(c_r) \\ 0 & 0 & 0 \\ M(\tilde{c}_r) & 0 & 0 \end{pmatrix},$$

⁴Up to multiplication by a positive rational.

where $c_r = (\underbrace{-1, \dots, -1}_{r \text{ times}}, 0, \dots, 0)$ for some $r \in \{1, \dots, q\}$.

Proof : First we note that $\tilde{\mathfrak{g}}/\mathfrak{g} \simeq \mathfrak{g}$ as a \mathfrak{g} -module, thus $\Sigma(\tilde{\mathfrak{g}}/\mathfrak{g}) = \Sigma(\mathfrak{g})$. We look at a rational vector γ such that $(\star) \text{Vect}(\Sigma(\mathfrak{g}) \cap \gamma^\perp) = \text{Vect}(\Sigma(\mathfrak{g})) \cap \gamma^\perp$. Up to the action of the Weyl group $W_{\mathfrak{a}}$, we can suppose that $\gamma = (\gamma_1 \geq \dots \geq \gamma_q \geq 0)$ is dominant. Since the set $\Sigma(\mathfrak{g})$ is generated by the simple roots $S(\mathfrak{g}) = \{f_i - f_{i+1}, 1 \leq i \leq q-1\} \cup \{f_q\}$, we see that for γ dominant, the condition (\star) is equivalent to asking that all linear forms of $S(\mathfrak{g})$ except one must vanish against γ . This last condition implies that γ is proportional to a certain γ_r for $r \in \{1, \dots, q\}$. \square

Now we have to detailed condition b) of Theorem 1.13 for an admissible element $(w_1\gamma_r, w_2\gamma_r, w_3\gamma_r)$. To do so, we use the conjugation by the element $\theta \in O(n)$ defined by

$$\theta = \frac{1}{\sqrt{2}} \begin{pmatrix} I_q & 0 & J_q \\ 0 & \sqrt{2}I_{p-q} & 0 \\ J_q & 0 & -I_q \end{pmatrix}.$$

Here $J_q = \begin{pmatrix} 0 & \dots & 1 \\ 0 & \dots & 0 \\ 1 & \dots & 0 \end{pmatrix}$ is a $q \times q$ matrix. We see that $\theta^2 = I_n$ and that $\tilde{\mathfrak{t}} := \text{Ad}(\theta)(\mathfrak{t}) \subset \mathfrak{u}(n)$ is the (usual) maximal torus of diagonal matrices. More precisely, the image of the matrix $\begin{pmatrix} D(a) & 0 & M(c) \\ 0 & D(b) & 0 \\ M(\tilde{c}) & 0 & D(\tilde{a}) \end{pmatrix}$ by $\text{Ad}(\theta)$ is equal to $\begin{pmatrix} D(a+c) & 0 & 0 \\ 0 & D(b) & 0 \\ 0 & 0 & D(\tilde{a}-\tilde{c}) \end{pmatrix}$.

Let $E_\theta = \theta(\mathbb{C}^r) \subset F_\theta = \theta(\mathbb{C}^{n-r})$ be the image of the flag $\mathbb{C}^r \subset \mathbb{C}^{n-r}$ by θ . The map $\varphi([g]) = g(E_\theta \subset F_\theta)$ defines a diffeomorphism between the flag manifold $\mathcal{F}_{\gamma_r} := GL_n(\mathbb{C})/\mathbb{P}_{\gamma_r}$ and the two-steps flag variety $\mathbb{F}(r, n-r; n)$. Moreover, for any $w \in W_{\mathfrak{a}}$ the image of the Schubert variety $\mathfrak{X}_{w, \gamma_r}$ through the diffeomorphism φ is equal to some $\theta(\mathfrak{X}_{\tilde{X}}) \subset \mathbb{F}(r, n-r; n)$ where $X_+ \cup X_- = w(\{1, \dots, r\})$. To be more precise the polarized subset $X_+ \cup X_-$ is equal to $\{|w(i)|, 1 \leq i \leq r\}$ and $k \in X_+$ if and only if there exists $i \in [r]$ with $k = w(i) = |w(i)|$.

Through the diffeomorphism φ , we see that the condition

$$b) \quad [\mathfrak{X}_{w_1, \gamma_r}] \odot_0 [\mathfrak{X}_{w_2, \gamma_r}] \odot_0 [\mathfrak{X}_{w_3, \gamma_r}] = [pt] \quad \text{in} \quad H^*(\mathcal{F}_{\gamma_r}, \mathbb{Z})$$

becomes $[\mathfrak{X}_{\tilde{I}}] \odot_0 [\mathfrak{X}_{\tilde{J}}] \odot_0 [\mathfrak{X}_{\tilde{K}}] = [pt]$ in $H^*(\mathbb{F}(r, n-r; n), \mathbb{Z})$, where $I_+ \cup I_- = w_1(\{1, \dots, r\})$, $J_+ \cup J_- = w_2(\{1, \dots, r\})$ and $K_+ \cup K_- = w_3(\{1, \dots, r\})$ are polarized subsets of $[q]$.

References

- [1] P. Belkale : Geometric proofs of Horn and saturation conjectures. J. Algebraic Geom. **15**, 133–173 (2006).

- [2] P. Belkale and S. Kumar : Eigenvalue problem and a new product in cohomology of flag varieties. *Invent. Math.* **166**, 185–228 (2006).
- [3] A. Berenstein and R. Sjamaar : Coadjoint orbits, moment polytopes, and the Hilbert-Mumford criterion, *Journal of the A.M.S.* **13**, 433–466 (2000).
- [4] N. Berline, M. Vergne and M. Walter : The Horn inequalities from a geometric point of view. *L’Enseignement Mathématique* **63**, 403–470 (2018).
- [5] A. Borel and A. Haefliger : La classe d’homologie fondamentale d’un espace analytique. *Bulletin de la Société Mathématique de France* **89**, 461–513 (1961).
- [6] M. Brion : Restriction de représentations et projections d’orbites coadjointes. *Séminaire Bourbaki*, (2012).
- [7] T. Bröcker and T. Tom Dieck. *Representations of compact Lie groups* (Vol. 98). Springer Science & Business Media (2013).
- [8] J. J. Duistermaat, J. A. C. Kolk and V. S. Varadarajan : Functions, flows and oscillatory integrals on flag manifolds and conjugacy classes in real semisimple Lie groups. *Compositio Mathematica* **49**, 309–398 (1983).
- [9] W. Fulton : Eigenvalues, invariant factors, highest weights, and Schubert calculus. *Bulletin of the American Mathematical Society* **37**, 209–249 (2000).
- [10] B. Hall : Phase space bounds for quantum mechanics on a compact Lie group. *Commun. Math. Phys.* **184**, 233–250 (1997).
- [11] A. Horn : Eigenvalues of sums of Hermitian matrices. *Pacific J. Math.* **12**, 225–241 (1962).
- [12] M. Kapovich B. Leeb and J. Millson : The generalized triangle inequalities in symmetric spaces and buildings with applications to algebra, *Mem. Amer. Math. Soc.* **192** , no. 896, (2008).
- [13] A. W. Knap : *Lie groups beyond an introduction* (Vol. 140). Boston: Birkhäuser (1996).
- [14] A. Klyachko : Stable bundles, representation theory and Hermitian operators, *Selecta Mathematica* **4**, 419–445 (1998).
- [15] A. Knutson and T. Tao : The honeycomb model of $GL_n(\mathbb{C})$ tensor products I : Proof of the saturation conjecture. *Journal of the A.M.S.* **12**, 1055–1090 (1999).
- [16] A. Knutson, T. Tao and C. Woodward : The honeycomb model of $GL_n(\mathbb{C})$ tensor products II: Puzzles determine facets of the Littlewood-Richardson cone. *Journal of the A.M.S.* **17**, 19–48 (2004).
- [17] L. Lesieur : Les problèmes d’intersection sur une variété de Grassmann, *C. R. Acad. Sci.* **225**, 916–917 (1947).
- [18] B. V. Lidskii : The proper values of the sum and product of symmetric matrices, *Dokl. Akad. Nauk SSSR* **74**, 769–772 (1950).
- [19] L. O’Shea and R. Sjamaar : Moment maps and Riemannian symmetric pairs. *Math. Annalen* **317**, 415–457 (2000).
- [20] P.-E. Paradan : Moment polytopes in real symplectic geometry I. arXiv preprint arXiv:2012.08837.

- [21] N. Ressayre : Geometric invariant theory and the generalized eigenvalue problem. *Inventiones mathematicae* **180**, 389–441 (2010).
- [22] N. Ressayre : Multiplicative formulas in Schubert calculus and quiver representation. *Indagationes Mathematicae* **22**, 87–102 (2011).
- [23] N. Ressayre : Geometric invariant theory and generalized eigenvalue problem II. *Annales de L’institut Fourier* **61**, 1467–1491 (2011).
- [24] R. W. Richardson : Intersections of double cosets in algebraic groups. *Indagationes Mathematicae* **3**, 69–77 (1992).
- [25] E. Richmond : A multiplicative formula for structure constants in the cohomology of flag varieties. *Michigan Mathematical Journal* **61**, 3–17 (2012).
- [26] M. Takeuchi: Cell decompositions and Morse equalities on certain symmetric spaces, *J. Fac. Sci. Univ. Tokyo*, **12** , 81–192 (1965).
- [27] R. C. Thompson : Singular value inequalities for matrix sums and minors, *Linear Algebra Appl.* **11**, 251–269 (1975).
- [28] H. Weyl : Das asymptotische Verteilungsgesetz der Eigenwerte lineare partieller Differentialgleichungen, *Math. Ann.* **71** , 441–479 (1912).