



HAL
open science

The canonical complex of the weak order

Doriann Albertin, Vincent Pilaud

► **To cite this version:**

Doriann Albertin, Vincent Pilaud. The canonical complex of the weak order. *Order. A Journal on the Theory of Ordered Sets and its Applications*, 2023, 40, pp.349-370. 10.1007/s11083-022-09605-7. hal-03451490

HAL Id: hal-03451490

<https://hal.science/hal-03451490>

Submitted on 26 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE CANONICAL COMPLEX OF THE WEAK ORDER

DORIANN ALBERTIN AND VINCENT PILAUD

ABSTRACT. We define and study the canonical complex of a finite semidistributive lattice L . It is the simplicial complex on the join or meet irreducible elements of L which encodes each interval of L by recording the canonical join representation of its bottom element and the canonical meet representation of its top element. This complex behaves properly with respect to lattice quotients of L , in the sense that the canonical complex of a quotient of L is the subcomplex of the canonical complex of L induced by the join or meet irreducibles of L uncontracted in the quotient. We then describe combinatorially the canonical complex of the weak order on permutations in terms of semi-crossing arc bidiagrams, formed by the superimposition of two non-crossing arc diagrams of N . Reading. We provide explicit direct bijections between the semi-crossing arc bidiagrams and the weak order interval posets of G. Châtel, V. Pilaud and V. Pons. Finally, we provide an algorithm to describe the Kreweras maps in any lattice quotient of the weak order in terms of semi-crossing arc bidiagrams.




FIGURE 1. The canonical complex of the weak order on \mathfrak{S}_4 labeled by arcs.

CONTENTS

1. Introduction	2
2. The canonical complex of a finite semidistributive lattice	3
2.1. Recollection on lattices	3
2.1.1. Join representations and semidistributive lattices	3
2.1.2. Kreweras maps	5
2.1.3. Lattice congruences	5
2.2. The canonical complex	7
3. Semi-crossing arc bidiagrams	9
3.1. Non-crossing arc diagrams	9
3.2. Weak order on arcs	12
3.3. Semi-crossing arc bidiagrams	12
3.4. Weak order interval posets	15
3.5. Kreweras maps in quotients of the weak order	16
acknowledgements	17
References	17

1. INTRODUCTION

A finite lattice L is join semidistributive when any element admits a canonical join representation (see *e.g.* [FN95] for a classical reference on lattices). This enables us to define the canonical join complex of L [Rea15, Bar19], whose vertices are the join irreducible elements of L and whose simplices are the canonical join representations in L . When L is both join and meet semidistributive, it thus admits both a canonical join complex and a canonical meet complex which are actually isomorphic flag simplicial complexes [Bar19].

In the first part of this paper, we define the canonical complex of a finite semidistributive lattice L , a larger flag simplicial complex where the canonical join complex and the canonical meet complex naturally live and interact. More precisely, its vertex set is the disjoint union of the set of join irreducible elements of L with the set of meet irreducible elements of L , and its simplices are the disjoint unions $J \sqcup M$ of a canonical join representation J in L with a canonical meet representation M in L such that $\bigvee J \leq \bigwedge M$. In other words, each interval $[x, y]$ in L contributes to a simplex of the canonical complex given by the disjoint union of the canonical join representation of x with the canonical meet representation of y . This provides a model for the intervals of L which is compatible with lattice quotients. Namely, the canonical complex of a quotient L/\equiv is the subcomplex of the canonical complex of L induced by the join and meet irreducibles of L uncontracted by the congruence \equiv .

In the second part of this paper, we study the combinatorics of the canonical complex of the weak order. N. Reading showed in [Rea15] that join irreducible permutations correspond to certain arcs wiggling around the horizontal axis, and that canonical join representations of permutations correspond to non-crossing arc diagrams. We show that the elements of the canonical complex can be interpreted as semi-crossing arc bidiagrams, defined as pairs $\delta_\vee \sqcup \delta_\wedge$ of non-crossing arc diagrams where only certain types of crossings are allowed between an arc of δ_\vee and an arc of δ_\wedge . It thus follows that the canonical complex of any quotient of the weak order is isomorphic to a subcomplex of the semi-crossing complex induced by arcs contained in an upper ideal of the subarc order. We then provide explicit direct bijections between the semi-crossing arc bidiagrams and the weak order interval posets of G. Châtel, V. Pilaud and V. Pons [CPP19], which are both in bijection with the intervals of the weak order. Finally, we provide an algorithm to describe the Kreweras maps in any lattice quotient of the weak order in terms of semi-crossing arc bidiagrams, generalizing the classical Kreweras complement on non-crossing partitions.

2. THE CANONICAL COMPLEX OF A FINITE SEMIDISTRIBUTIVE LATTICE

This section deals with canonical meet and join representations in a finite semidistributive lattice and its quotients. We start with a recollection on join semidistributive lattices, their canonical join representations, their canonical join complexes, their Kreweras maps, and their lattice congruences (Section 2.1). We then define the canonical complex of a semidistributive lattice L which encodes the intervals of L and contains both the canonical join complex and the canonical meet complex of L (Section 2.2).

2.1. Recollection on lattices. We start by a quick recollection on semidistributive lattices, canonical representations, canonical complexes, Kreweras maps and lattice congruences. All the material covered here is classical, we refer for instance to [FN95, Rea16, Rea15, Bar19]. Following [Bar19, Exm. 10], we illustrate this section with the case of distributive lattices.

2.1.1. Join representations and semidistributive lattices. Consider a finite lattice (L, \leq, \vee, \wedge) where \vee is the join and \wedge is the meet. We see \vee and \wedge as internal binary operators on L and try to factorize the elements of L in some canonical way. It is first important to understand the irreducible elements for \vee and \wedge .

Definition 1. An element $x \in L$ is called *join* (resp. *meet*) *irreducible* if it covers (resp. is covered by) a unique element denoted x_* (resp. x^*). We denote by $\mathcal{JI}(L)$ (resp. $\mathcal{MI}(L)$) the subposet of L induced by the set of join (resp. meet) irreducible elements of L .

Definition 2. A *join representation* of $x \in L$ is a subset $J \subseteq L$ such that $x = \bigvee J$. Such a representation is *irredundant* if $x \neq \bigvee J'$ for any strict subset $J' \subsetneq J$. The irredundant join representations in L are antichains of L , and are ordered by containment of the lower sets of their elements (*i.e.* $J \leq J'$ if and only if for any $y \in J$ there exists $y' \in J'$ such that $y \leq y'$ in L). The *canonical join representation* of x , denoted $\mathbf{cjr}(x)$, is the minimal irredundant join representation of x for this order, when it exists.

Note that when it exists, $\mathbf{cjr}(x)$ is an antichain of $\mathcal{JI}(L)$. The following statement characterizes the lattices where canonical join representations exist.

Proposition 3 ([FN95, Thm. 2.24 & Thm. 2.56]). *A finite lattice L is join semidistributive when the following equivalent conditions hold:*

- (i) $x \vee y = x \vee z$ implies $x \vee (y \wedge z) = x \vee y$ for any $x, y, z \in L$,
- (ii) for any cover relation $x < y$ in L , the set

$$K_{\vee}(x, y) := \{z \in L \mid z \not\leq x \text{ but } z \leq y\} = \{z \in L \mid x \vee z = y\}$$

- has a unique minimal element $k_{\vee}(x, y)$ (which is then automatically join irreducible),
- (iii) any element of L admits a canonical join representation.

Moreover, the canonical join representation of $y \in L$ is $\mathbf{cjr}(y) = \{k_{\vee}(x, y) \mid x < y\}$.

Note that in a finite join semidistributive lattice L , we can associate to any meet irreducible element m of L a join irreducible element $\kappa_{\vee}(m) := k_{\vee}(m, m^*)$ of L . Moreover, the existence of canonical join representations enable us to consider the following complex, illustrated in Figure 2. It was initially defined in [Rea15], where a combinatorial model was provided for the weak order (see Section 3.1), and studied in [Bar19] for arbitrary finite semidistributive lattices.

Definition 4. The *canonical join complex* $\mathcal{CJC}(L)$ of a finite join semidistributive lattice L is the simplicial complex on $\mathcal{JI}(L)$ whose faces are the canonical join representations of the elements of L .

The *meet semidistributivity*, the maps K_{\wedge} , k_{\wedge} and κ_{\wedge} , the *canonical meet representation* $\mathbf{cmr}(x)$ and the *canonical meet complex* $\mathcal{CMC}(L)$ are all defined dually. A lattice L is *semidistributive* if it is both meet and join semidistributive. In this case, the maps κ_{\vee} and κ_{\wedge} define inverse bijections between $\mathcal{MI}(L)$ and $\mathcal{JI}(L)$, and the complexes $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ behave particularly nicely.




FIGURE 2. Some semidistributive lattices and their canonical join (red) and meet (blue) complexes. The letters label all join or meet irreducible elements, and we denote by x_V (resp. x_\wedge) the element x when it is considered as a join (resp. meet) irreducible. Note that we always consistently color joinands in red and meetands in blue. The bottom two lattices are distributive while the top two are only semidistributive.

Proposition 5 ([Bar19, Thm. 2 & Coro. 5]). *If L is a finite semidistributive lattice, then*

- (i) $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ are flag simplicial complexes (i.e. their minimal non-faces are edges, or equivalently they are the clique complexes of their graphs),
- (ii) the maps κ_V and κ_\wedge induce inverse isomorphisms between $\mathcal{CMC}(L)$ and $\mathcal{CJC}(L)$.

In fact, it was proved in [Bar19, Thm. 2] that $\mathcal{CJC}(L)$ is flag if and only if L is semidistributive. We will not use the “only if” direction in this paper.

Example 6 (Distributive lattices). The name semidistributivity actually comes from the well understood class of distributive lattices. A lattice L is *distributive* if $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ for any $x, y, z \in L$. Note that the dual condition $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ for any $x, y, z \in L$ is actually equivalent to the primal one. The fundamental theorem for distributive lattices affirms that L is distributive if and only if it is isomorphic to the lattice of lower sets of its join irreducible poset P . In other words, any antichain of join irreducible elements in P forms a canonical join representation in L . To be more precise, consider, for an antichain A of P , the two lower sets

$$j_A := \{x \in P \mid x \leq y \text{ for some } y \in A\} \quad \text{and} \quad m^A := \{x \in P \mid x \not\leq y \text{ for all } y \in A\}.$$

Said differently, A is the set of maximal elements of j_A and the set of minimal elements of $P \setminus m^A$. For $y \in P$, we abbreviate $j_{\{y\}}$ into j_y and $m^{\{y\}}$ into m^y . Then

- the join (resp. meet) irreducibles of L are precisely the lower sets j_y (resp. m^y) for $y \in P$,
- the map κ_\vee (resp. κ_\wedge) is given by $\kappa_\vee(m^y) = j_y$ (resp. $\kappa_\wedge(j_y) = m^y$),
- the canonical join representation of j_A and the canonical meet representation of m^A are

$$\mathbf{cjr}(j_A) = \{j_y \mid y \in A\} \quad \text{and} \quad \mathbf{cmr}(m^A) = \{m^y \mid y \in A\}.$$

- the canonical join and meet complexes $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ are both (isomorphic to) the clique complex on the incomparability graph of P .

See [Bar19, Exm. 10].

2.1.2. *Kreweras maps.* In a semidistributive lattice L , each element has both a canonical join representation and a canonical meet representation. It is natural to consider the maps that exchange the canonical join representation with the canonical meet representation of the same element.

Definition 7. The *Kreweras maps* $\eta_\vee : \mathcal{CMC}(L) \rightarrow \mathcal{CJC}(L)$ and $\eta_\wedge : \mathcal{CJC}(L) \rightarrow \mathcal{CMC}(L)$ are defined by

$$\eta_\vee(M) := \mathbf{cjr}\left(\bigwedge M\right) \quad \text{and} \quad \eta_\wedge(J) := \mathbf{cmr}\left(\bigvee J\right).$$

Note that some authors call Kreweras maps the compositions $\eta_\vee \circ \kappa_\wedge : \mathcal{CJC}(L) \rightarrow \mathcal{CJC}(L)$ and $\eta_\wedge \circ \kappa_\vee : \mathcal{CMC}(L) \rightarrow \mathcal{CMC}(L)$, see for instance [Bar19]. As will be discussed in Example 50, the Kreweras maps for the Tamari lattice are closely related to the classical Kreweras complement on non-crossing partitions. For the moment, we recall that the Kreweras maps for the distributive lattices are related to rowmotion.

Example 8 (Distributive lattices). With the notations of Example 6, for an antichain A in P , we denote by $\text{row}_\vee(A)$ the set of maximal elements of m^A and by $\text{row}_\wedge(A)$ the set of minimal elements of $P \setminus j_A$. In other words, we have $m^A = j_{\text{row}_\vee(A)}$ and $j_A = m_{\text{row}_\wedge(A)}$. Hence, by Example 6, the Kreweras maps η_\vee and η_\wedge are given by

$$\eta_\vee(\{m^y \mid y \in A\}) = \{j_y \mid y \in \text{row}_\vee(A)\} \quad \text{and} \quad \eta_\wedge(\{j_y \mid y \in A\}) = \{m^y \mid y \in \text{row}_\wedge(A)\}.$$

See [Bar19, Rem. 32].

2.1.3. *Lattice congruences.* We now discuss quotients of the lattice L , considered as an algebraic structure with two internal binary operators \vee and \wedge . We thus need equivalence relations on L that respects \vee and \wedge .

Definition 9. A *congruence* \equiv on L is an equivalence relation on L such that $x \equiv x'$ and $y \equiv y'$ implies $x \vee y \equiv x' \vee y'$ and $x \wedge y \equiv x' \wedge y'$. Equivalently, the equivalence classes are intervals, and the maps π_\downarrow^\equiv and π_\uparrow^\equiv sending an element to the minimum and maximum elements in its congruence class are order preserving.

Definition 10. The *lattice quotient* L/\equiv is the lattice structure on the congruence classes, where for any two congruence classes X and Y ,

- the order is given by $X \leq Y$ if and only if $x \leq y$ for some representatives $x \in X$ and $y \in Y$,
- the join $X \vee Y$ (resp. meet $X \wedge Y$) is the congruence class of $x \vee y$ (resp. $x \wedge y$) for any representatives $x \in X$ and $y \in Y$.

Note that the lattice quotient L/\equiv is isomorphic to the subposet of L induced by the minimal (or maximal) elements in their congruence classes. This subposet is a join (resp. meet) subsemilattice of L but may fail to be a sublattice of L . We now consider all congruences of L .

Definition 11. The *congruence lattice* $\text{con}(L)$ is the set of all congruences of L ordered by refinement.

The congruence lattice $\text{con}(L)$ is a distributive lattice where the meet is the intersection of relations and the join is the transitive closure of union of relations. For any join irreducible element $j \in \mathcal{JI}(L)$, we denote by $\text{con}(j)$ the unique minimal congruence of L that *contracts* j , that is with $j_* \equiv j$. It turns out that $\text{con}(j)$ is join irreducible in $\text{con}(L)$ and that all join irreducible congruences in $\text{con}(L)$ are of this form. Hence, any congruence of L is completely determined by the set of join irreducible elements of L that it contracts. We denote by $\mathcal{UJI}(\equiv)$ the set of join

irreducible elements of L uncontracted by \equiv . Not all subsets of join irreducible elements of L are of the form $\mathcal{UJI}(\equiv)$ for some congruence \equiv of L . The possible subsets are governed by the following relation.

Definition 12. For $j, j' \in \mathcal{JI}(L)$, we say that j *forces* j' , and write $j \succcurlyeq j'$, if $\text{con}(j) \geq \text{con}(j')$, that is if any congruence contracting j also contracts j' .

The forcing relation is a preorder \preccurlyeq (i.e. a transitive and reflexive, but not necessarily antisymmetric, relation) on $\mathcal{JI}(L)$, whose upper sets correspond to the congruences of L .

Proposition 13 ([Rea16, Prop. 9-5.16]). *The following conditions are equivalent for $J \subseteq \mathcal{JI}(L)$:*

- J is an upper set of the forcing preorder (i.e. $j \succcurlyeq j'$ and $j \in J$ implies $j' \in J$).
- $J = \mathcal{UJI}(\equiv)$ for some congruence \equiv of L .

As already mentioned, the set $\mathcal{UJI}(\equiv)$ characterizes \equiv . It moreover enables to understand the elements of L which are minimal in their congruence classes and their canonical representations as follows.

Proposition 14 ([Rea16, Prop. 9-5.29]). *Let \equiv be a congruence of a finite join semidistributive lattice L . Then*

- an element $x \in L$ is minimal in its congruence class if and only if $\mathbf{cjr}(x) \subseteq \mathcal{UJI}(\equiv)$,
- the quotient L/\equiv is join semidistributive and the canonical joinands of a congruence class X in L/\equiv are the congruence classes of the canonical joinands of the minimal element in X .

Proposition 14 translates as follows to the canonical join complex $\mathcal{CJC}(L)$.

Proposition 15. *Let \equiv be a congruence on a finite join semidistributive lattice L . Then the canonical join complex $\mathcal{CJC}(L/\equiv)$ of the quotient L/\equiv is isomorphic to the subcomplex $\mathcal{CJC}(\equiv)$ of the canonical join complex $\mathcal{CJC}(L)$ of L induced by $\mathcal{UJI}(\equiv)$.*

We will need the following statement relating the canonical join representation of an element x of L with the canonical join representation of the minimal element $\pi_{\downarrow}^{\equiv}(x)$ in its equivalence class. It ensures that we can look for $\mathbf{cjr}(\pi_{\downarrow}^{\equiv}(x))$ among the antichains of join irreducible elements of L uncontracted by \equiv and below a join irreducible element of $\mathbf{cjr}(x)$. Unfortunately, it remains difficult in general to describe $\mathbf{cjr}(\pi_{\downarrow}^{\equiv}(x))$ because not all such antichains define a canonical representation of L . We provide a proof here as we have not found this statement explicitly in the literature.

Proposition 16. *Let \equiv be a congruence of a finite join semidistributive lattice L . For any $x \in L$, the lower ideal of L generated by $\mathbf{cjr}(x)$ contains $\mathbf{cjr}(\pi_{\downarrow}^{\equiv}(x))$.*

Proof. The proof works by induction on the size of the interval $[\pi_{\downarrow}^{\equiv}(x), x]$. The statement is immediate if $\pi_{\downarrow}^{\equiv}(x) = x$. Otherwise, there exists $j \in \mathbf{cjr}(x) \setminus \mathcal{UJI}(\equiv)$. Let $y := \bigvee (\mathbf{cjr}(x) \Delta \{j, j_{\star}\})$, where Δ denotes the symmetric difference. Since $j \equiv j_{\star}$, we have $x \equiv y$ and thus $\pi_{\downarrow}^{\equiv}(x) = \pi_{\downarrow}^{\equiv}(y)$. Since y has a join representation strictly contained in the lower ideal of L generated by $\mathbf{cjr}(x)$, we have $y < x$ so that $[\pi_{\downarrow}^{\equiv}(y), y]$ is strictly contained in $[\pi_{\downarrow}^{\equiv}(x), x]$. By induction hypothesis, we thus obtain that $\mathbf{cjr}(\pi_{\downarrow}^{\equiv}(y))$ is contained in the lower ideal of L generated by $\mathbf{cjr}(y)$. Moreover, observe that $J := (\mathbf{cjr}(x) \setminus \{j\}) \cup \mathbf{cjr}(j_{\star})$ is a join representation for y , so that $\mathbf{cjr}(y)$ is contained in the lower ideal of L generated by J , which is itself contained in the lower ideal of L generated by $\mathbf{cjr}(x)$ (since any element of $\mathbf{cjr}(j_{\star})$ is lower than j_{\star} and thus than j). We conclude that $\mathbf{cjr}(\pi_{\downarrow}^{\equiv}(x))$ is indeed in the lower ideal of L generated by $\mathbf{cjr}(x)$. \square

Dual statements hold using meets instead of joins, and we denote by $\mathcal{UMI}(\equiv)$ the meet irreducible elements of L uncontracted by \equiv , and by $\mathcal{CMC}(\equiv)$ the subcomplex of $\mathcal{CMC}(L)$ induced by $\mathcal{UMI}(\equiv)$ for a congruence \equiv on L . Due to Proposition 15, we will always work with the subcomplexes $\mathcal{CJC}(\equiv)$ and $\mathcal{CMC}(\equiv)$ rather than with the complexes $\mathcal{CJC}(L/\equiv)$ and $\mathcal{CMC}(L/\equiv)$.

When the lattice L is semidistributive, the two sets $\mathcal{UJI}(\equiv)$ and $\mathcal{UMI}(\equiv)$ and the two subcomplexes $\mathcal{CJC}(\equiv)$ and $\mathcal{CMC}(\equiv)$ are connected by the maps κ_{\vee} and κ_{\wedge} . We provide a proof here as we have not found this statement explicitly in the literature.

Proposition 17. *Let \equiv be a congruence on a finite semidistributive lattice L . Then we have $\mathcal{UJI}(\equiv) = \kappa_{\vee}(\mathcal{UMI}(\equiv))$ and $\mathcal{UMI}(\equiv) = \kappa_{\wedge}(\mathcal{UJI}(\equiv))$. Hence, the maps κ_{\vee} and κ_{\wedge} induce inverse isomorphisms between the subcomplexes $\mathcal{CMC}(\equiv)$ and $\mathcal{CJC}(\equiv)$.*

Proof. Let $m \in \mathcal{MI}(L)$ and $j = \kappa_{\vee}(m)$. By definition, we have $m \vee j = m^*$ and $m \vee j_{\star} = m$. Hence, $j \equiv j_{\star}$ implies that $m = m \vee j_{\star} \equiv m \vee j = m^*$. In other words, $\kappa_{\vee}(\mathcal{UMI}(\equiv)) \subseteq \mathcal{UJI}(\equiv)$. By symmetry, we have $\kappa_{\wedge}(\mathcal{UJI}(\equiv)) \subseteq \mathcal{UMI}(\equiv)$. Since κ_{\vee} and κ_{\wedge} are reversed bijections, this yields equalities. The last sentence of the statement thus follows from Proposition 5 (ii). \square

Example 18 (Distributive lattices). In a distributive lattice L , there is no forcing at all. Hence, any subset of join irreducible elements of L defines a congruence of L . In other words, with the notations of Example 6, any subset Y of P defines a congruence \equiv_Y with

$$\mathcal{UJI}(\equiv_Y) = \{j_y \mid y \in Y\} \quad \text{and} \quad \mathcal{UMI}(\equiv_Y) = \{m^y \mid y \in Y\}.$$

The lattice quotient L/\equiv is again distributive and isomorphic to the lattice of lower ideals of the restriction of the poset P to Y .

2.2. The canonical complex. We now define another complex that connects the canonical join complex $\mathcal{CJC}(L)$ to the canonical meet complex $\mathcal{CMC}(L)$ using intervals of L . This complex is illustrated in Figure 3.

Definition 19. The *canonical complex* $\mathcal{CC}(L)$ of a finite semidistributive lattice L is the simplicial complex whose

- ground set is the **disjoint** union $\mathcal{JI}(L) \sqcup \mathcal{MI}(L)$ of the sets of join irreducible and of meet irreducible elements of L , and
- faces are the **disjoint** unions $J \sqcup M$ where $J \in \mathcal{CJC}(L)$ is a canonical join representation, $M \in \mathcal{CMC}(L)$ is a canonical meet representation, and $\bigvee J \leq \bigwedge M$.

To avoid any confusion, let us insist here that when an element $x \in L$ is both join irreducible and meet irreducible, then it appears twice in $\mathcal{CC}(L)$ and may appear twice in some faces of $\mathcal{CC}(L)$, once as a join irreducible element and once as a meet irreducible element. We will thus always write the faces of $\mathcal{CC}(L)$ explicitly as disjoint unions. In our pictures, we denote by x_{\vee} (resp. x_{\wedge}) and color red (resp. blue) the element $x \in L$ considered as a join (resp. meet) irreducible.

Our first observation is that $\mathcal{CC}(L)$ contains both $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ as induced subcomplexes.

Proposition 20. *For a finite semidistributive lattice L , the canonical join (resp. meet) complex $\mathcal{CJC}(L)$ (resp. $\mathcal{CMC}(L)$) is the subcomplex of the canonical complex $\mathcal{CC}(L)$ induced by the join (resp. meet) irreducible elements $\mathcal{JI}(L)$ (resp. $\mathcal{MI}(L)$).*

Proof. We have $J \in \mathcal{CJC}(L) \iff J \sqcup \emptyset \in \mathcal{CC}(L)$ and $M \in \mathcal{CMC}(L) \iff \emptyset \sqcup M \in \mathcal{CC}(L)$. \square

We next observe that, while the canonical join and meet complexes $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ encode the individual elements of L , the canonical complex $\mathcal{CC}(L)$ encodes the intervals of L .

Proposition 21. *For a finite semidistributive lattice L , the faces of the canonical complex $\mathcal{CC}(L)$ are in bijection with the intervals of L .*

Proof. Send a face $J \sqcup M$ of $\mathcal{CC}(L)$ to the interval $[\bigvee J, \bigwedge M]$ of L , and conversely an interval $[x, y]$ to the face $\mathbf{cjr}(x) \sqcup \mathbf{cmr}(y)$ of $\mathcal{CC}(L)$. \square

Our next statement is the analogue of Proposition 5 (i).

Proposition 22. *For a finite semidistributive lattice L , the canonical complex $\mathcal{CC}(L)$ is a flag simplicial complex.*

Proof. Consider $J \subseteq J'$ and $M \subseteq M'$ such that $J' \sqcup M' \in \mathcal{CC}(L)$. Then $J \in \mathcal{CJC}(L)$ since $J' \in \mathcal{CJC}(L)$ and $M \in \mathcal{CMC}(L)$ since $M' \in \mathcal{CMC}(L)$ (since $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ are simplicial complexes), and $\bigvee J \leq \bigvee J' \leq \bigwedge M' \leq \bigwedge M$. Hence $J \sqcup M \in \mathcal{CC}(L)$ so that $\mathcal{CC}(L)$ is indeed a simplicial complex.




FIGURE 3. The canonical complexes of the semidistributive lattices of Figure 2. The corresponding join (resp. meet) canonical complexes of Figure 2 are highlighted in red (resp. blue). Since the canonical complexes are flag by Propositions 5 and 22, it is sufficient to represent their graphs. The letters label all join or meet irreducible elements, and we denote by x_v (resp. x_\wedge) the element x when it is considered as a join (resp. meet) irreducible. The vertices of the canonical complex are positioned so that the map κ of Remark 24 acts by central symmetry.

Consider $J \subseteq \mathcal{JI}(L)$ and $M \subseteq \mathcal{MI}(L)$ such that any two elements of $J \sqcup M$ form a face of $\mathcal{CC}(L)$. Then $J \in \mathcal{CJC}(L)$ and $M \in \mathcal{CMC}(L)$ (since $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ are flag by Proposition 5 (i)), and $\bigvee J \leq \bigwedge M$ since $j \leq m$ for any $j \in J$ and $m \in M$. \square

Our next statement says that $\mathcal{CC}(L)$ is not only a simplicial complex, it is actually naturally embedded on the boundary of a cross-polytope.

Proposition 23. *For any $j \in \mathcal{JI}(L)$, the pair $\{j, \kappa_\wedge(j)\}$ is not in $\mathcal{CC}(L)$. Hence, for any labeling $\lambda : \mathcal{JI}(L) \rightarrow \|\mathcal{JI}(L)\|$, the map sending j to $e_{\lambda(j)}$ and $\kappa_\wedge(j)$ to $-e_{\lambda(j)}$ defines an embedding of $\mathcal{CC}(L)$ to the boundary of the $|\mathcal{JI}(L)|$ -dimensional cross-polytope.*

Proof. By definition, $j \not\leq \kappa_\wedge(j)$ so that $\{j, \kappa_\wedge(j)\}$ is not in $\mathcal{CC}(L)$. The second sentence thus follows from Proposition 22. \square

Remark 24. Denote by κ the map on $\mathcal{JI}(L) \sqcup \mathcal{MI}(L)$ defined by $\kappa(m) := \kappa_v(m)$ for $m \in \mathcal{MI}(L)$ and $\kappa(j) := \kappa_\wedge(j)$ for $j \in \mathcal{JI}(L)$. It corresponds to the central symmetry on the corresponding cross-polytope. By Proposition 5 (ii), the two subcomplexes $\mathcal{CJC}(L)$ and $\mathcal{CMC}(L)$ are symmetric under the action of κ . However, the full canonical complex $\mathcal{CC}(L)$ is not invariant under the action of κ . See Figure 3 for examples.




FIGURE 4. Hasse diagrams of the right weak orders of size 2, 3, and 4.

Example 25. The canonical complex of the boolean lattice on $[n]$ is isomorphic to the boundary of the n -dimensional cross-polytope.

Finally, analogously to Proposition 15, the canonical complex is compatible with lattice congruences of L .

Proposition 26. For any congruence \equiv of a finite semidistributive lattice L , the canonical complex $\mathcal{CC}(L/\equiv)$ of the quotient L/\equiv is isomorphic to the subcomplex $\mathcal{CC}(\equiv)$ of the canonical complex $\mathcal{CC}(L)$ of L induced by the disjoint union $\mathcal{UJI}(\equiv) \sqcup \mathcal{UMI}(\equiv)$ of the join and meet irreducible elements of L uncontracted by \equiv .

Proof. This immediately follows from Proposition 15 and its dual version. □

Example 27 (Distributive lattices). With the notations of Example 6, we have $j_y \subseteq m^z \iff y \not\leq z$. Hence, the canonical complex $\mathcal{CC}(L)$ is the clique complex of the graph whose vertex set is made of two copies P_\vee and P_\wedge of P and whose edge set is the union of two copies I_\vee and I_\wedge of the incomparability graph of P with the edges $\{y_\vee, z_\wedge\}$ for $y \not\leq z$ in P .

3. SEMI-CROSSING ARC BIDIAGRAMS

In this section, we apply the lattice theoretic results presented in Section 2 to the classical weak order on permutations. We first recall that the canonical join and meet representations of the weak order can be encoded as non-crossing arc diagrams as defined by N. Reading in [Rea15] (Section 3.1). We then briefly study the restriction of the weak order on join or meet irreducibles in terms of arcs (Section 3.2). We then describe the intervals and the canonical complex of the weak order in terms of semi-crossing arc diagrams (Section 3.3). We then provide direct bijections between the semi-crossing arc bidiagrams and the weak order interval posets of [CPP19] (Section 3.4). Finally, we provide an algorithm to compute the Kreweras maps in any lattice quotient of the weak order, generalizing the classical Kreweras complement on non-crossing partitions (Section 3.5).

3.1. Non-crossing arc diagrams. We consider the following classical order on the set \mathfrak{S}_n of permutations of $[n] := \{1, \dots, n\}$, illustrated in Figure 4.

Definition 28. An *inversion* of a permutation $\sigma \in \mathfrak{S}_n$ is a pair (u, v) with $1 \leq u < v \leq n$ and $\sigma^{-1}(u) > \sigma^{-1}(v)$ (in other words, u is smaller than v but u appears after v in σ). The (right) *weak order* of size n is the order on permutations of \mathfrak{S}_n defined by inclusion of their inversion sets.




FIGURE 5. NCADs and SCABs of the permutations 2537146, 2531746, 2513746, and 2513476. The first line represents the table $(\sigma(i), i)$ of a permutation σ with ascents in blue and descents in red, the second line is the join diagram $\delta_{\vee}(\sigma)$, the third line is the meet diagram $\delta_{\wedge}(\sigma)$, and the fourth line is the superimposition $\delta_{\vee}(\sigma) \sqcup \delta_{\wedge}(\sigma)$.

Note that a cover relation in the weak order corresponds to the swap of two values σ_i and σ_{i+1} at consecutive positions. The swap is increasing in the weak order if i is an *ascent* *i.e.* $\sigma_i < \sigma_{i+1}$, and decreasing if i is a *descent* *i.e.* $\sigma_i > \sigma_{i+1}$.

It is classical that the weak order is a semidistributive lattice. We now describe its join (resp. meet) irreducible elements and its canonical join (resp. meet) representations in terms of the arcs and non-crossing arc diagrams introduced by N. Reading in [Rea15].

Definition 29 ([Rea15]). An *arc* is a quadruple (a, b, A, B) where $1 \leq a < b \leq n$ and $A \sqcup B =]a, b[$ forms a partition of $]a, b[:= \{a+1, \dots, b-1\}$. Two arcs $\alpha := (a, b, A, B)$ and $\alpha' := (a', b', A', B')$ *cross* if there exist $u \neq v$ such that $u \in (A' \cup \{a', b'\}) \cap (B \cup \{a, b\})$ and $v \in (A \cup \{a, b\}) \cap (B' \cup \{a', b'\})$. A *non-crossing arc diagram* (or *NCAD* for short) is a collection of pairwise non-crossing arcs. The *non-crossing complex* is the clique complex of the non-crossing relation on all arcs.

Remark 30. Visually, an arc (a, b, A, B) is represented by an x -monotone curve wiggling around the horizontal axis, starting at a and ending at b , and passing above points of A and below points of B . Two arcs cross if they cross in their interior or start at the same point or end at the same point (but they do not cross if one ends where the other starts). See Figure 5 for illustrations of arcs and of non-crossing arc diagrams.

Observe that the join (resp. meet) irreducible elements of the weak order are precisely the permutations with exactly one descent (resp. ascent). Hence, we associate to an arc $\alpha := (a, b, A, B)$ with $A := \{a_1 < \dots < a_k\}$ and $B := \{b_1 < \dots < b_\ell\}$

- a join irreducible permutation $\sigma_{\vee}(\alpha) := [1, \dots, (a-1), a_1, \dots, a_k, b, a, b_1, \dots, b_\ell, (b+1), \dots, n]$,
- a meet irreducible permutation $\sigma_{\wedge}(\alpha) := [n, \dots, (b+1), a_k, \dots, a_1, a, b, b_\ell, \dots, b_1, (a-1), \dots, 1]$,

where we use the one-line notation of permutations $\sigma = [\sigma_1, \dots, \sigma_n]$.

Conversely, consider a permutation $\sigma \in \mathfrak{S}_n$ represented by its permutation table formed by dots at coordinates (σ_i, i) for $i \in [n]$. Draw segments between consecutive dots (σ_i, i) and $(\sigma_{i+1}, i+1)$,

colored red for a descent $\sigma_i > \sigma_{i+1}$ and blue for an ascent $\sigma_i < \sigma_{i+1}$. Finally, flatten the picture vertically to the horizontal line, allowing segments to bend but not to pass points. The resulting picture is the superimposition of a set $\delta_\vee(\sigma)$ of red arcs and a set $\delta_\wedge(\sigma)$ of blue arcs. See Figure 5. More formally, $\delta_\vee(\sigma) := \{\alpha_\vee(\sigma, i) \mid \sigma_i < \sigma_{i+1}\}$ and $\delta_\wedge(\sigma) := \{\alpha_\wedge(\sigma, i) \mid \sigma_i > \sigma_{i+1}\}$ where

$$\alpha_\vee(\sigma, i) := (\sigma_i, \sigma_{i+1}, \{\sigma_j \mid j < i \text{ and } \sigma_i < \sigma_j < \sigma_{i+1}\}, \{\sigma_j \mid j > i + 1 \text{ and } \sigma_i < \sigma_j < \sigma_{i+1}\}),$$

and $\alpha_\wedge(\sigma, i) := (\sigma_{i+1}, \sigma_i, \{\sigma_j \mid j < i \text{ and } \sigma_i > \sigma_j > \sigma_{i+1}\}, \{\sigma_j \mid j > i + 1 \text{ and } \sigma_i > \sigma_j > \sigma_{i+1}\}).$

Proposition 31 ([Rea15]). *The map δ_\vee (resp. δ_\wedge) is a bijection between the set of permutations of \mathfrak{S}_n and the set of non-crossing arc diagrams of size n . Moreover, the canonical join (resp. meet) representation of a permutation $\sigma \in \mathfrak{S}_n$ is given by $\mathbf{cjr}(\sigma) = \{\sigma_\vee(\alpha_\vee) \mid \alpha_\vee \in \delta_\vee(\sigma)\}$ (resp. $\mathbf{cmr}(\sigma) = \{\sigma_\wedge(\alpha_\wedge) \mid \alpha_\wedge \in \delta_\wedge(\sigma)\}$). Hence, the canonical join (resp. meet) complex of the weak order is isomorphic to the non-crossing complex.*

By construction, the non-crossing arc diagrams are adapted to the maps κ_\vee and κ_\wedge and to quotients of the weak order. First, our next statement says that the maps κ_\vee and κ_\wedge only change the colors of the arcs. We provide a proof as we have not found it explicitly in [Rea15].

Proposition 32. $\kappa_\vee(\sigma_\wedge(\alpha)) = \sigma_\vee(\alpha)$ and $\kappa_\wedge(\sigma_\vee(\alpha)) = \sigma_\wedge(\alpha)$ for any arc α .

Proof. Let $\alpha := (a, b, A, B)$, let $\sigma := \sigma_\wedge(\alpha) = [n, \dots, (b+1), a_k, \dots, a_1, a, b, b_\ell, \dots, b_1, (a-1), \dots, 1]$ and let $\sigma^* = [n, \dots, (b+1), a_k, \dots, a_1, b, a, b_\ell, \dots, b_1, (a-1), \dots, 1]$ denote the only element covering σ . Any permutation τ with $\tau \not\leq \sigma$ but $\tau \leq \sigma^*$ must have all elements of A before b before a before all elements of B . Therefore, the minimal such permutation is clearly $\sigma_\vee(\alpha) = [1, \dots, (a-1), a_1, \dots, a_k, b, a, b_1, \dots, b_\ell, (b+1), \dots, n]$. \square

Proposition 33 ([Rea15]). *For any arcs $\alpha := (a, b, A, B)$ and $\alpha' := (a', b', A', B')$, the join irreducible $\sigma_\vee(\alpha)$ forces the join irreducible $\sigma_\vee(\alpha')$ if and only if α is a **subarc** of α' , meaning that $a' \leq a < b \leq b'$ and $A \subseteq A'$ while $B \subseteq B'$. Hence, to each upper ideal I of the subarc order corresponds a lattice congruence \equiv_I of the weak order, and the canonical join (resp. meet) complex of the quotient of the weak order by \equiv_I is isomorphic to the non-crossing complex on I .*




FIGURE 6. The subarc order on arcs of sizes 3 (top left), 4 (top right), and 5 (bottom).

Remark 34. Visually, α is a subarc of α' if the endpoints of α are weakly in between the endpoints of α' , and α follows α' between its endpoints. The subarc order on arcs of size 3 to 5 is represented in Figure 6.

Example 35. The prototypical congruence of the weak order is the *sylvestre congruence* \equiv_{syly} [LR98, HNT05], which can be defined equivalently as

- the fiber of the binary search tree insertion (inserting a permutation from right to left),
- the congruence where each class is the set of linear extensions of a binary tree (labeled in inorder and oriented toward its root),
- the transitive closure of the rewriting rule $UacVbW \equiv UcaVbW$ for $1 \leq a < b < c \leq n$,
- the congruence corresponding to the upper ideal of the subarc order given by all up arcs $(a, b,]a, b[, \emptyset)$ (or equivalently, generated by the long up arc $(1, n,]1, n[, \emptyset)$.

Hence, a permutation is minimal (resp. maximal) in its class if and only if it avoids the pattern 312 (resp. 132). The quotient of the weak order by the sylvestre congruence is (isomorphic to) the classical *Tamari lattice* [Tam51, HT72], whose elements are the binary trees on n nodes and whose cover relations are rotations in binary trees. The canonical representations in the Tamari lattice correspond to non-crossing sets of up arcs, also known as *non-crossing partitions*. The sylvestre congruence was extended in [Rea06] to Cambrian congruences and in [PP18] to permutree congruences.

3.2. Weak order on arcs. We now briefly compare join or meet irreducible elements in the weak order in terms of arcs. For this, we first observe that the inversions of $\sigma_{\vee}(\alpha)$ and $\sigma_{\wedge}(\alpha)$ are easily read on the arc α .

Lemma 36. *For any arc $\alpha := (a, b, A, B)$ and any $u < v$, the pair (u, v) is an inversion of $\sigma_{\vee}(\alpha)$ (resp. of $\sigma_{\wedge}(\alpha)$) if and only if $u \in B \cup \{a\}$ and $v \in A \cup \{b\}$ (resp. if $u \notin A \cup \{a\}$ or $v \notin B \cup \{b\}$).*

Proof. Immediate from the definition $\sigma_{\vee}(\alpha) := [1, \dots, (a-1), a_1, \dots, a_k, b, a, b_1, \dots, b_\ell, (b+1), \dots, n]$ and $\sigma_{\wedge}(\alpha) := [n, \dots, (b+1), a_k, \dots, a_1, a, b, b_\ell, \dots, b_1, (a-1), \dots, 1]$. \square

Corollary 37. *For any two arcs $\alpha := (a, b, A, B)$ and $\alpha' := (a', b', A', B')$, we have*

- (i) $\sigma_{\vee}(\alpha) \leq \sigma_{\vee}(\alpha')$ if and only if $a \in B' \cup \{a'\}$ while $b \in A' \cup \{b'\}$ and $A \subseteq A'$ while $B \subseteq B'$,
- (ii) $\sigma_{\wedge}(\alpha) \leq \sigma_{\wedge}(\alpha')$ if and only if $a' \in B \cup \{a\}$ while $b' \in A \cup \{b\}$ and $A' \subseteq A$ while $B' \subseteq B$,
- (iii) $\sigma_{\vee}(\alpha) \leq \sigma_{\wedge}(\alpha')$ if and only if there is no $u < v$ such that $u \in (A' \cup \{a'\}) \cap (B \cup \{a\})$ and $v \in (A \cup \{b\}) \cap (B' \cup \{b'\})$.

Remark 38. Figure 7 shows the weak order on arcs defined by $\alpha \leq \alpha'$ if $\sigma_{\vee}(\alpha) \leq \sigma_{\vee}(\alpha')$. Visually, $\alpha \leq \alpha'$ if α is a subarc of α' which starts weakly below α' and ends weakly above α' . Note that $\alpha := (a, b, A, B)$ covers at most two arcs, namely $(\min B, b, A \cap]\min b, b[, B \setminus \min B)$ and $(a, \max A, A \setminus \max A, B \cap]a, \max A])$ when they are defined. Similar remarks hold for the order defined by $\sigma_{\wedge}(\alpha)$ instead of $\sigma_{\vee}(\alpha)$.

Remark 39. As illustrated in Figure 7, the weak order on join irreducible of \mathfrak{S}_n has interesting enumerative properties. Let us just mention here that it has

- $2^n - n - 1$ elements (permutations with a single descent, or arcs) [OEI10, A000295],
- $2^{n+1} - n^2 - n - 2$ cover relations (in bijection with arcs of size $n+1$ crossing the horizontal axis, or with subsets of $[n+1]$ crossing their complement) [OEI10, A324172],
- $n(n+1)2^{n-2}$ intervals (including the singletons) [OEI10, A001788].

3.3. Semi-crossing arc bidiagrams. We now describe the canonical complex of the weak order as defined in Section 2.2 in terms of the following combinatorial objects, illustrated in Figure 8.

Definition 40. A *semi-crossing arc bidiagram* (or *SCAB* for short) is a pair $(\delta_{\vee}, \delta_{\wedge})$ of non-crossing arc diagrams such that for any arcs $\alpha_{\vee} := (a_{\vee}, b_{\vee}, A_{\vee}, B_{\vee}) \in \delta_{\vee}$ and $\alpha_{\wedge} := (a_{\wedge}, b_{\wedge}, A_{\wedge}, B_{\wedge}) \in \delta_{\wedge}$, there is no $u < v$ with $u \in (A_{\wedge} \cup \{a_{\wedge}\}) \cap (B_{\vee} \cup \{a_{\vee}\})$ and $v \in (A_{\vee} \cup \{b_{\vee}\}) \cap (B_{\wedge} \cup \{b_{\wedge}\})$. The *semi-crossing complex* is the simplicial complex whose ground set contains two copies α_{\vee} and α_{\wedge} of each arc α and whose simplices are all semi-crossing arc bidiagrams.




FIGURE 7. The weak orders of size 3 (top left), 4 (top right), and 5 (bottom) restricted to their join irreducibles represented by the corresponding arcs.




FIGURE 8. SCABs of the intervals $[2531746, 2531746]$, $[2531746, 2537146]$, $[2513476, 2537146]$ and $[5264137, 6574231]$.

Remark 41. Visually, a semi-crossing arc bidiagram $\delta_\vee \sqcup \delta_\wedge$ is a collection of arcs such that

- no two arcs of δ_\vee (resp. of δ_\wedge) cross in their interior, or start or end at the same points,
- no two arcs $\alpha_\vee \in \delta_\vee$ and $\alpha_\wedge \in \delta_\wedge$ cross in their interiors with α_\vee going up and α_\wedge going down at the crossing, or start at the same point with α_\vee leaving above α_\wedge , or end at the same point with α_\vee arriving below α_\wedge at this point.

Remark 42. Before going further, we report in Table 1 on the number of semi-crossing arc bidiagrams $\delta_\vee \sqcup \delta_\wedge$ according to the cardinalities $|\delta_\vee|$ and $|\delta_\wedge|$ for $n = 2$ to 6.

In these tables, observe that

- the first row (resp. column) corresponds to the intervals $[\sigma, w_\circ]$ (resp. $[e, \sigma]$) for the permutations σ of $[n]$ with k ascents (resp. descents) and are thus counted by the Eulerian numbers [OEI10, A008292],
- the last row (resp. column) corresponds to the single interval $[w_\circ, w_\circ]$ (resp. $[e, e]$).

We now connect the semi-crossing arc bidiagrams with the canonical complex of the weak order using Corollary 37.

Proposition 43. *The map $[\sigma, \tau] \mapsto \delta_\vee(\sigma) \sqcup \delta_\wedge(\tau)$ is a bijection between the intervals of the weak order on \mathfrak{S}_n and the semi-crossing arc bidiagrams. Hence, the canonical complex of the weak order is isomorphic to the semi-crossing complex.*

Proof. By Proposition 31, $\delta_\vee(\sigma)$ and $\delta_\wedge(\tau)$ are both non-crossing arc diagrams, and correspond to the canonical join representation of σ and the canonical meet representation of τ respectively.

0	1	1	0	1	2	0	1	2	3	3	
0	1	1	0	1	4	1	0	1	11	11	1
1	1	0	1	4	6	0	1	11	54	24	0
			2	1	0	0	2	11	24	2	0
							3	1	0	0	0

0	1	26	66	26	1	0	1	57	302	302	57	1
1	26	300	420	80	0	1	57	1340	4145	2505	240	0
2	66	420	320	20	0	2	302	4145	8270	3035	120	0
3	26	80	20	0	0	3	302	2505	3035	562	5	0
4	1	0	0	0	0	4	57	240	120	5	0	0
						5	1	0	0	0	0	0

TABLE 1. The number of semi-crossing arc bidiagrams $\delta_\vee \sqcup \delta_\wedge$ according to the cardinalities $|\delta_\vee|$ and $|\delta_\wedge|$ for $n = 2$ to 6 .




FIGURE 9. The weak order on \mathfrak{S}_3 with permutations labeled by semi-crossing arc bidiagrams, and the canonical complex of \mathfrak{S}_3 with join and meet irreducible permutations labeled by arcs.

By Corollary 37(iii), $\sigma \leq \tau$ implies that $\delta_\vee(\sigma) \sqcup \delta_\wedge(\tau)$ is a semi-crossing arc bidiagram, and corresponds to the canonical representation of $[\sigma, \tau]$. \square

For instance, the canonical complexes of the weak orders on \mathfrak{S}_3 and \mathfrak{S}_4 are illustrated in Figures 1 and 9. As usual, since the canonical complex is flag by Proposition 22, we only represent its graph. The central symmetry corresponds to the map κ of Remark 24, which just corresponds to the exchange of color of the arcs by Proposition 32.

Let us insist again here that this combinatorial model for the intervals of the weak order is adapted to the study of its quotients. The next statement follows from Propositions 32 and 33.

Proposition 44. *For any lower ideal I of the subarc order, the canonical complex of the quotient of the weak order by \equiv_I is isomorphic to the subcomplex of the semi-crossing complex induced by $\{\alpha_\vee \mid \alpha \in I\} \sqcup \{\alpha_\wedge \mid \alpha \in I\}$.*

Remark 45. Observe that the semi-crossing arc bidiagram corresponding to a singleton $[\sigma, \sigma]$ is just a path alternating between increasing blue steps and decreasing red steps corresponding to the ascents and descents of σ . See Figure 5. In general, we conjecture that the semi-crossing arc bidiagram corresponding to an inclusion minimal interval in a lattice quotient of the weak order does not contain crossing in the interior of an arc.

3.4. Weak order interval posets. Following a classical result of A. Björner and M. Wachs [BW91, Thm. 6.8], G. Châtel, V. Pilaud and V. Pons already studied in [CPP19] a family of posets in bijection with the intervals of the weak order.

Proposition 46 ([BW91, Thm. 6.8] & [CPP19, Prop. 26]). *The following conditions are equivalent for a poset \triangleleft on $[n]$:*

- the linear extensions of \triangleleft form an interval $[\sigma, \tau]$ of the weak order,
- there are $\sigma \leq \tau$ in the weak order such that the increasing relations of \triangleleft are the non-inversions of τ and the decreasing relations are the inversions of σ ,
- $a < c$ implies $a \triangleleft b$ or $b \triangleleft c$, and $a \triangleright c$ implies $a \triangleright b$ or $b \triangleright c$ for all $1 \leq a < b < c \leq n$.

Such a poset is called a **weak order interval poset** (or **WOIP** for short).

Although they are specific to the weak order and do not behave well with respect to its quotients, the WOIPs are combinatorial objects in bijection with intervals of the weak order, and thus with SCABs. It is thus relevant to provide direct explicit bijections between SCABs and WOIPs, which are illustrated in Figure 10. We need the following definitions, illustrated in Figure 10.

From SCABs to WOIPs. For any arc $\alpha := (a, b, A, B)$, we denote by \triangleleft_α the set of increasing relations $u \triangleleft_\alpha v$ (by \triangleright_α the set of decreasing relations $u \triangleright_\alpha v$) where $u < v$ with $u \in A \cup \{a\}$ and $v \in B \cup \{b\}$. For a NCAD δ , we denote by \triangleleft_δ (resp. \triangleright_δ) the transitive closure of the union of the decreasing relations $\{\triangleright_{\alpha_\vee} \mid \alpha_\vee \in \delta_\vee\}$ (resp. of the increasing relations $\{\triangleleft_{\alpha_\wedge} \mid \alpha_\wedge \in \delta_\wedge\}$).

From WOIPs to SCABs. Fix a WOIP $\triangleright \sqcup \triangleleft$ where \triangleright denote the decreasing relations and \triangleleft denote the increasing relations. We say that an increasing (resp. decreasing) cover relation $a \triangleleft b$ (resp. $a \triangleright b$) is **maximal** if there is no cover relation $a' \triangleleft b$ (resp. $a' \triangleright b$) with $a' < a$ or $a \triangleleft b'$ (resp. $a \triangleright b'$) with $b < b'$. To a maximal decreasing (resp. increasing) cover relation $a \triangleright b$ (resp. $a \triangleleft b$), we associate the arc $\alpha(a \triangleright b) := (a, b,]a, b[\cap \langle a \rangle_\downarrow,]a, b[\cap \langle b \rangle^\uparrow)$ (resp. $\alpha(a \triangleleft b) := (a, b,]a, b[\cap \langle b \rangle_\downarrow,]a, b[\cap \langle a \rangle^\uparrow)$), where $\langle x \rangle_\downarrow$ and $\langle x \rangle^\uparrow$ denote the lower and upper ideals of \triangleleft generated by x . Finally, we denote by $\delta(\triangleright)$ (resp. $\delta(\triangleleft)$) the set of arcs $\alpha(a \triangleright b)$ (resp. $\alpha(a \triangleleft b)$) for all maximal decreasing (resp. increasing) cover relations of \triangleleft .

Proposition 47. *The maps $\delta_\vee \sqcup \delta_\wedge \mapsto \triangleright_{\delta_\vee} \sqcup \triangleleft_{\delta_\wedge}$ and $\triangleright \sqcup \triangleleft \mapsto \delta(\triangleright) \sqcup \delta(\triangleleft)$ are inverse bijections between SCABs and WOIPs.*

Proof. Consider an interval $[\sigma, \tau]$ of the weak order corresponding to a SCAB $\delta_\vee \sqcup \delta_\wedge$. For any arc α_\vee , the decreasing relations of $\triangleright_{\alpha_\vee}$ are precisely the inversions of $\sigma_\vee(\alpha_\vee)$ by Lemma 36. Hence, the decreasing relations of $\triangleright_{\delta_\vee}$ are precisely the inversions of $\sigma = \bigvee \{\sigma_\vee(\alpha_\vee) \mid \alpha_\vee \in \delta_\vee\}$ (since the inversion set of a join is the transitive closure of the union of the inversion sets of the joinands). Similarly, the increasing relations of $\triangleleft_{\delta_\wedge}$ are precisely the non-inversions of $\tau = \bigwedge \{\sigma_\wedge(\alpha_\wedge) \mid \alpha_\wedge \in \delta_\wedge\}$. Hence $\triangleright_{\delta_\vee} \sqcup \triangleleft_{\delta_\wedge}$ is indeed the WOIP of the interval $[\sigma, \tau]$ by Proposition 46 (ii). Finally, to see that the two maps are inverse to each other, we just need to observe that the relations $a \triangleleft_{\delta_\wedge} b$ created out of the extremities of the arcs (a, b, A, B) of δ_\wedge are precisely the maximal cover relations of $\triangleleft_{\delta_\wedge}$ (and similarly for $\triangleright_{\delta_\vee}$). \square




FIGURE 10. The bijection between SCABs (top) and WOIPs (bottom). We represent a WOIP \triangleleft with arcs joining $a < b$ from above (resp. below) in blue (resp. red) when $a < b$ (resp. $a > b$).

3.5. Kreweras maps in quotients of the weak order. We finally describe the Kreweras maps defined in Section 2.1.2 in all quotients of the weak order in terms of semi-crossing arc bidiagrams. For this, we first connect the canonical join representation of a permutation to the canonical join representation of the minimal element in its class for a given congruence, as illustrated in Figure 12. In the following proposition, we call weak order on arcs the order $\alpha \leq \alpha'$ if $\sigma_\vee(\alpha) \leq \sigma_\vee(\alpha')$ (see Section 3.2 and Figure 7).

Proposition 48. *Consider an upper ideal I of the subarc order and a permutation σ . Let X be the intersection of I with the lower ideal generated by the non-crossing arc diagram $\delta_\vee(\sigma)$ in the weak order on arcs. Let Y be the set of arcs (a, b, A, B) of X such that there is $a < p < b$ such that both arcs $(a, p, A \cap]a, p[$, $B \cap]a, p[$) and $(p, b, A \cap]p, b[$, $B \cap]p, b[$) belong to X . Then the non-crossing arc diagram $\delta_\vee(\pi_{\downarrow}^{\equiv I}(\sigma))$ is the set of maximal elements of $X \setminus Y$ in the weak order on arcs.*

Proof. By Propositions 14 and 16, $\delta_\vee(\pi_{\downarrow}^{\equiv I}(\sigma))$ is a non-crossing arc diagram contained in X . Among all options, we need to choose the non-crossing arc diagram with maximal join. This implies that the arcs of Y cannot appear in $\delta_\vee(\pi_{\downarrow}^{\equiv I}(\sigma))$ since

$$\sigma_\vee(a, b, A, B) \leq \sigma_\vee(a, p, A \cap]a, p[$$
, $B \cap]a, p[$) \vee $\sigma_\vee(p, b, A \cap]p, b[$, $B \cap]p, b[$)

for any arc (a, b, A, B) and any $a < p < b$ by Lemma 36. We finally claim that any two incomparable elements in $X \setminus Y$ cannot cross. Hence, the maximal elements of $X \setminus Y$ form a non-crossing arc diagram, which must therefore be $\delta_\vee(\pi_{\downarrow}^{\equiv I}(\sigma))$. To prove the claim, consider two arcs $\alpha := (a, b, A, B)$ and $\alpha' := (a', b', A', B')$ in X which are incomparable and crossing. Assume for instance that there are $u < v$ such that $u \in (A \cup \{a\}) \cap (B' \cup \{a'\})$ and $v \in (A' \cup \{b'\}) \cap (B \cup \{b\})$. We can moreover assume that α and α' agree on $]u, v[$, meaning that $A \cap]u, v[= A' \cap]u, v[$ and $B \cap]u, v[= B' \cap]u, v[$. Since α and α' are incomparable, this implies that $u \neq a$ or $v \neq b$. Hence, at least two of the three arcs

$$(a, u, A \cap]a, u[$$
, $B \cap]a, u[$), $(u, v, A' \cap]u, v[$, $B' \cap]u, v[$) and $(u, b, A \cap]u, b[$, $B \cap]u, b[$)

are non-trivial (not reduced to a single point). Moreover, they also belong to X , which implies that α is in Y . \square

This enables us to compute the Kreweras maps in quotients of the weak order directly on non-crossing arc diagrams. For this, let us extend the notations of Section 2.1.2 to quotients and transport them to non-crossing arc diagrams. For an upper ideal I of the subarc order, each equivalence class of \equiv_I is an interval $[x, y]$ of the weak order and thus corresponds to two non-crossing arc diagrams $\delta_\vee := \delta_\vee(x)$ and $\delta_\wedge := \delta_\wedge(y)$. We denote by η_\vee^I and η_\wedge^I the two opposite maps defined by $\eta_\vee^I(\delta_\wedge) = \delta_\vee$ and $\eta_\wedge^I(\delta_\vee) = \delta_\wedge$. We just write η_\vee and η_\wedge when I is the set of all arcs. Note that $\eta_\vee = \delta_\vee \circ \delta_\wedge^{-1}$ and $\eta_\wedge = \delta_\wedge \circ \delta_\vee^{-1}$ are easily computed from the descriptions of the maps δ_\vee and δ_\wedge (see Section 3.1) and of their inverses (see the explicit description in [Rea15]). Proposition 48 enables to compute η_\vee^I and η_\wedge^I in general.

Corollary 49. *Consider an upper ideal I of the subarc order and a non-crossing arc diagram δ_\wedge with all arcs in I . Then the non-crossing arc diagram $\eta_\vee^I(\delta_\wedge)$ is obtained from $\eta_\vee(\delta_\wedge)$ by applying the algorithm of Proposition 48.*

Example 50. When I is the upper ideal of up arcs corresponding to the sylvester congruence, the description of Corollary 49 can be translated to the classical description of the Kreweras complement of a non-crossing partition. Namely, the Kreweras complement of a non-crossing partition is obtained by shifting the points and connecting the points in the same connected component. See Figure 11.




FIGURE 11. Classical Kreweras complement on non-crossing partitions.




FIGURE 12. Illustrations of computations of $\delta_V(\pi_{\downarrow}^{\equiv I}(\sigma))$ and $\eta_V^I(\delta_{\wedge})$ in the trivial congruence, the descent congruence, the sylvester congruence, and a generic congruence.

ACKNOWLEDGEMENTS

We are grateful to J.-C. Novelli for many discussions and suggestions on the content and presentation of this paper, and to S. Giraudo for a careful proofreading of a preliminary version.

REFERENCES

- [Bar19] Emily Barnard. The canonical join complex. *Electron. J. Combin.*, 26(1):Paper No. 1.24, 25, 2019.
- [BW91] Anders Björner and Michelle L. Wachs. Permutation statistics and linear extensions of posets. *J. Combin. Theory Ser. A*, 58(1):85–114, 1991.
- [CPP19] Grégory Chatel, Vincent Pilaud, and Viviane Pons. The weak order on integer posets. *Algebraic Combinatorics*, 2(1):1–48, 2019.
- [FN95] Ralph Freese and James B. Nation. Free lattices. *Mathematical Surveys and Monographs*, 75:93–106, 1995.
- [HNT05] Florent Hivert, Jean-Christophe Novelli, and Jean-Yves Thibon. The algebra of binary search trees. *Theoret. Comput. Sci.*, 339(1):129–165, 2005.
- [HT72] Samuel Huang and Dov Tamari. Problems of associativity: A simple proof for the lattice property of systems ordered by a semi-associative law. *J. Combin. Theory Ser. A*, 13:7–13, 1972.
- [LR98] Jean-Louis Loday and María O. Ronco. Hopf algebra of the planar binary trees. *Adv. Math.*, 139(2):293–309, 1998.
- [OEI10] The On-Line Encyclopedia of Integer Sequences. Published electronically at <http://oeis.org>, 2010.
- [PP18] Vincent Pilaud and Viviane Pons. Permutrees. *Algebraic Combinatorics*, 1(2):173–224, 2018.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea15] Nathan Reading. Noncrossing arc diagrams and canonical join representations. *SIAM J. Discrete Math.*, 29(2):736–750, 2015.
- [Rea16] Nathan Reading. Lattice theory of the poset of regions. In *Lattice theory: special topics and applications. Vol. 2*, pages 399–487. Birkhäuser/Springer, Cham, 2016.
- [Tam51] Dov Tamari. *Monoides préordonnés et chaînes de Malcev*. PhD thesis, Université Paris Sorbonne, 1951.

(DA) LIGM, UNIVERSITÉ GUSTAVE EIFFEL, CNRS, ESIEE PARIS, F-77454 MARNE-LA-VALLÉE, FRANCE

Email address: `doriann.albertin@u-pem.fr`

URL: <https://doriann-albertin.github.io/site/>

(VP) CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU

Email address: `vincent.pilaud@lix.polytechnique.fr`

URL: <http://www.lix.polytechnique.fr/~pilaud/>