



HAL
open science

A remarkable new *Penia* species (Coleoptera: Elateridae: Dimini) from Guizhou, China.

Lu Qiu

► **To cite this version:**

Lu Qiu. A remarkable new *Penia* species (Coleoptera: Elateridae: Dimini) from Guizhou, China..
Faunitaxys, 2021, 9 (37), pp.1-4. hal-03452937

HAL Id: hal-03452937

<https://hal.science/hal-03452937>

Submitted on 27 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*



Volume 9
Numéro 37

Novembre 2021

ISSN : 2269 - 6016
Dépôt légal : Novembre 2021

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*

ZooBank : <http://zoobank.org/79A36B2E-F645-4F9A-AE2B-ED32CE6771CC>

Directeur de la publication, rédacteur, conception graphique et PAO:

Lionel Delaunay

Cette revue ne peut pas être vendue

Elle est distribuée par échange aux institutions (version papier)

et sur simple demande aux particuliers (format PDF)

à l'adresse suivante:

AFCFF (Association française de Cartographie de la Faune et de la Flore)

28, rue Voltaire, F- 42100 Saint Etienne

E-mail: lionel.delaunay@free.fr

Elle est disponible librement au téléchargement à partir du site:

<http://faunitaxys.fr/>

La parution de *Faunitaxys* est apériodique

Faunitaxys est indexé dans / *Faunitaxys* is indexed in:

- **Zoological Record**

Articles and nomenclatural novelties are referenced by:

- **ZooBank** (<http://zoobank.org>)

Online Archives:

- **HAL (archive ouverte)** (<http://hal.archives-ouvertes.fr>)

Imprimée sur les presses de SPEED COPIE

6, rue Tréfilerie, F- 42100 Saint-Etienne

Imprimé le 26 novembre 2021

A remarkable new *Penia* species (Coleoptera: Elateridae: Dimini) from Guizhou, China

LU QIU

Engineering Research Center for Forest and Grassland Disaster Prevention and Reduction, Mianyang Normal University, Mianxing West Road, 621000, Mianyang, China. - e-mail: 123church@163.com - <https://orcid.org/0000-0002-0946-1634>
- ZooBank: <http://zoobank.org/DFDB0DF3-222C-4AEF-A7AD-5EC1E61BBF55>

Mots-clés :

Coleoptera; morphology;
Elateroidea; taxonomy;
Dendrometrinae; new species;
Penia; Guizhou;
click-beetle; China.

Abstract. – *Penia jiangi* sp. nov. is described and illustrated from South Guizhou, China. Habitus and diagnostic characters of the new species are illustrated.

Qiu L., 2021. – A remarkable new *Penia* species (Coleoptera: Elateridae: Dimini) from Guizhou, China. *Faunitaxys*, 9(37) : 1 – 4.

ZooBank: <http://zoobank.org/DCD27E43-5A3A-4474-87DB-13A434F652C7>

Introduction

Genus *Penia* Laporte, 1838 (Elateridae: Dendrometrinae) is the most species-rich genus (109 species) in the tribe Dimini (Kundrata et al. 2018; Kundrata and Németh 2019). This genus is mainly distributed from Himalaya region to East and South East Asia. Currently, 27 species are recorded from China, and 24 of which were described by R. Schimmel (Schimmel 1993; 1996a, b; 1998; 1999; 2001; 2003; 2006a, b; 2015a, b). Most species of *Penia* are not well studied. The original descriptions of these species are brief, and most of the illustrations are insufficient for identification. Although Kundrata et al. (2018) made an essential catalogue for *Penia*, more comprehensive works are needed to be done in this genus.

After years' accumulation of Dimini specimens, I have collected most of the Chinese *Penia* species. Comprehensive revisional works are preparing. Considering they will be time-consuming works, thus, I decided to separately publish some remarkable new species to reduce the workload. In this paper, an easily distinguished *Penia* species is described from Guizhou, China.

Material and methods

The specimen was softened by hot water, then the genital segments were cut down and dissected after treatment in 80°C 10% KOH for 10 minutes. Habitus and diagnostic characters were photographed using a Canon® EOS RP digital camera + Mount Adapter EF-EOS R plus a Laowa® 25 mm F2.8 2.5-5× Ultra Macro Lens and a Laowa 100 mm F2.8 CA-Dreamer Macro 2× lens (all for Canon EF). All figures were modified in Adobe Photoshop® CC 2019. Body length was measured from the anterior margin of head to the apex of elytra, pronotal length was measured at mid-line, pronotal width was measured at hind angles, and width of body was measured at the widest place (on elytra). The type specimens of the new species are stored in the Invertebrate Collection of Mianyang Normal University, Mianyang, Sichuan, China (MYNU). The collecting data is quoted verbatim (in Chinese) in quotation marks. English translation of the data, as well as additional information, is given in square brackets.

Systematics

Family **Elateridae**

Subfamily **Dendrometrinae**

Genus ***Penia*** Laporte, 1838

***Penia jiangi* sp. nov.**

(Fig. 1-4)

ZooBank: <http://zoobank.org/826E231F-DCA0-4320-8897-648AF09694CE>

Holotype, ♂, “2017.IV.27 贵州荔波县茂兰 必左景区 478-507m 姜日新” [China: Guizhou Province, Libo County, Maolan N.R., Bizuo Scenic Spot, 25°17'16"N, 108°04'18"E, 478-507 m, 27.IV.2017, Ri-Xin Jiang leg.] (MYNU).

Paratypes

- 1 ♂, “2019.IV.13 贵州荔波县茂兰 必左景区 487 m 李泊言” [China: Guizhou Province, Libo County, Maolan N.R., Bizuo, 487 m, 13.IV.2019, Bo-Yan Li leg.] (MYNU);
- 1 ♀, same data as holotype (MYNU).

Diagnosis. – This species is easily distinguished from other *Penia* species by the reddish head, pronotum and scutellar shield, as well as the four large whitish spots on elytra.

Description (based on male holotype)

Dimensions (mm).

- Body length 9.1; width 3.4;
- Antenna length 7.8;
- Pronotum length 1.4, width 2.3;
- Elytron length 7.0, width 1.7.

Shape and coloration. – Tricolor and median-sized species with slender antennae and legs (Fig. 1A-B). Body elongate, flat, surface covered with erect, setaceous, and moderate long pubescence. Head yellowish red (including labrum and mandibles), apex and outline of mandibles blackish, maxillary palpi brown, antennae blackish brown. Pronotum, prothorax, and scutellar shield yellowish red, margins of pronotum and scutellar shield blackish. Prothorax dully brown, mesoventrite yellowish. Leg blackish brown, coxae, trochanters and tarsi palely yellowish brown. Elytra blackish brown, striae of basal half slightly reddish brown, each elytron with two large whitish spots.

Abdomen dark brown, with lateral portions slightly brownish. Pubescence yellowish brown to dark brown.

Head. – Head including eyes 0.6 times as wide as pronotum. Head surface sparsely with shallow punctures; punctures small, intervals between punctures mostly equal 1–3 puncture diameter. Eyes large, hemispheric. Edge of frons smooth apically, not carinated, not overhanging the base of labrum; surface above antennal insertions elevated and forming carinae. Maxillary palpus with palpomere III greatly longer than wide. Antenna very long, almost reaching the apical portion of elytron, consists of 11 antennomeres, length ratio of antennomeres II–XI = 1.0 : 1.9 : 3.3 : 3.8 : 3.8 : 3.8 : 3.8 : 3.8 : 4.6; scape robust, pedicel the shortest; antennomeres from III on almost elongate column-shaped, antennomere III distinctly longer than pedicel; antennomeres V–X similar in length, slightly longer than antennomere IV; ultimate antennomere the longest, narrowed apically, apex blunt.

Pronotum. – Very narrow when compared with its congeners, shape somewhat quadrated, widest at hind angles. Wider than long when length measured between mid-line and width between hind angles. In lateral view, pronotum only slightly arched medially. Sides slightly convex; posterior angles long, straight, not curved, divergent as ca. 45-degree angle, apices truncated. Disc sparsely punctate; punctures shallow, intervals between punctures on average subequal to two to five diameters of one puncture; interstices smooth.

Hypomeron and prosternum. – Hypomeron with much more densely punctate than pronotum, intervals between punctures on average subequal to one to two diameters of one puncture. Prosternal sutures excavated anteriorly, sinuate at basal half. Prosternum with punctation slightly sparser than on hypomeron, intervals between punctures on average subequal to two to three diameters of one puncture. Prosternal process small, arched dorsad, curved ventrad, apex blunt.

Scutellar shield. – Typically shield-shaped.

Pterothorax. – Mesoventrite with short medial protrusion of frontal margin, apically only slightly emarginate medially; procoxal rests distinct; sides of mesoventral cavity not elevated above mesocoxae, hind margin with an elongate extension between mesocoxae. Mesanepisternum with distinct large lateral extensions of procoxal rests. Metacoxal plate narrowly elongate.

Elytra. – Elongate, together about 2.1 times as long as wide, widest at ca. 2/3 of their length from base. Elytral apices meeting together. Striae of the elytra formed by lines of large and deep punctures; intervals between punctures on average subequal to two to three diameters of one puncture. Interstriae smooth, with four to six small punctures between the nearest four punctures on the striae. Two large,

round, whitish spots situated near the middle of elytra, while the other two large, semicircle, whitish spots situated apically.

Abdomen. – Densely pubescent. Tergite VIII (Fig. 2A) near trapezoid-shaped, 1.5 times as long as wide, apically covered with sparse setae and dense micro pubescence. Sternite VIII (Fig. 2A) with two large, dark colored lobes, margins with long setae. Tergite IX (Fig. 2B) subtriangular, 1.3 times as long as wide, median deeply and roundly emarginate, two lobes stout, rounded apically, lateral sides with long setae; tergite X (Fig. 2B) membranous, emarginate apically. Sternite IX (Fig. 2C) elongate, about 2.6 times as long as wide, apically rounded and setose.

Aedeagus (Fig. 3A–C). – Robust median lobe notably surpassing parameres, gradually narrowed, apical portion sharply narrowing; apex bluntly rounded, the lobe bearing sclerite inside, which with a triangular hook apically. Paramere elongate, consisting of two portions, the lateral portion well sclerotized, the inner portion transparent. Sides of the sclerotized portion paralleled, apical portion slightly sinuate, apex curved, spinous, pointing ventrad. The transparent portion subulate-shaped, apex blunt, notably surpassing the apex of the sclerotized portion, with four to six long pubescence at apical-lateral portion. Phallobase narrowed, U-shaped.

Male paratype. – Body length 7.2 mm, distinctly smaller and narrower than holotype, but other characters are almost the same as the holotype.

Female paratype. – Body length 10.2 mm, distinctly stouter than the male, sides of pronotum and elytra more convex than male (Fig. 1C). Tergite VIII longer than wide, 0.8 times as long as wide (Fig. 4A); sternite VIII almost as wide as long, with middle part and apex transparent, apical portion rounded, spiculum ventrale 2.3 times of sternite VIII length (Fig. 4B). Ovipositor (Fig. 4C) with robust styli; bursa copulatrix simple, without spines.

Immature stages. – Unknown.

Distribution. – China: Guizhou (Libo).

Etymology. – The specific epithet is dedicated to Dr. Ri-Xin Jiang (Guizhou University, China), the collector of the holotype.

Natural history. – This species was collected by beating the branches of the shrubs by umbrella (Ri-Xin Jiang and Bo-Yan Li, personal communications).



Fig. 1. *Penia jiangi* sp. nov., habitus. A & B) Holotype, ♂. C) Paratype, ♀. Scale bar = 10 mm.




Fig. 2. *Penia jiangi* sp. nov., holotype, ♂ (Scale bar = 1 mm).

A) Abdominal sternite VIII and tergite VIII, ventral view. B) Abdominal tergites IX–X, dorsal view. C) Abdominal sternite IX, dorsal view.



Fig. 3. *Penia jiangi* sp. nov., holotype, ♂, genitalia (Scale bar = 1 mm).




Fig. 4. *Penia jiangi* sp. nov., paratype, ♀ (Scale bar = 1 mm).

A) Abdominal tergite VIII, dorsal view. B) Abdominal sternite VIII, ventral view. C) Ovipositor.

Remarks. – This new species gave an example to show the general outline of pronotum may be varied among one species in *Penia*, especially among the male and female.

Acknowledgements

I thank Dr. Ri-Xin Jiang and Mr. Bo-Yan Li (Guizhou University, China) for collecting and providing the precious specimens of the new species. I am grateful to Dr. Alexander S. Prosvirov (Moscow State University, Russia), Dr. Yong-Ying Ruan (Shenzhen Polytechnic, China), and the late Rainer Schimmel (Germany) for various help. This study was partially supported by the Happy Snail Collecting Fund [No. 01231991].

References

- Kundrata R., Musalkova M. & Kubackzova M., 2018. – Annotated catalogue of the click-beetle tribe Dimini (Coleoptera: Elateridae: Dendrometrinae). *Zootaxa*, 4412 (1): 1-75.
- Kundrata R. & Németh R., 2019. – Description of *Penia mantillerii* sp. nov. (Coleoptera: Elateridae: Dimini), with a key to *Penia* species from Vietnam and nearby areas. *Zootaxa*, 4612 (2): 275-281.
- Schimmel R., 1993. – Neue Arten sowie eine neue Gattung der Unterfamilie Diminae Candèze, 1863 aus Ostasien und dem Balkan (Coleoptera: Elateridae). *Koleopterologische Rundschau*, 63: 245-259.
- Schimmel R., 1996a. – Das Monophylum Diminae Candèze, 1863 (Insecta: Coleoptera: Elateridae). *Pollichia-Buch*, 33, 1-370.
- Schimmel R., 1996b. – Neue und wenig bekannte Elateriden aus Griechenland und Ostasien (Coleoptera: Elateridae). *Koleopterologische Rundschau*, 66: 161-177.
- Schimmel R., 1998. – Neue Elateriden aus Ostasien und dem Iran (Coleoptera: Elateridae). *Koleopterologische Rundschau*, 68: 217-226.
- Schimmel R., 1999. – Neue und wenig bekannte Elateriden sowie eine neue Gattung, *Acumenator* n. gen., aus Südostasien (Insecta: Coleoptera, Elateridae). *Mitteilungen der Pollichia*, 85: 231-259.
- Schimmel R., 2001. – Neue und wenig bekannte Diminae-Arten aus China, Indien, Laos, Myanmar, Nepal und Vietnam (Insecta, Coleoptera: Elateridae). *Mitteilungen der Pollichia*, 88: 205-231.
- Schimmel R., 2003. – Neue Ampedini-, Physorhinini-, Pectocerini-, Elaterini- und Diminae-Arten aus Südostasien (Insecta, Coleoptera, Elateridae). *Mitteilungen der Pollichia*, 90: 265–292.
- Schimmel R., 2006a. – Neue Megapenthini-, Physorhinini-, Diminae- und Senodoniina-Arten aus Südostasien (Insecta: Coleoptera, Elateridae). *Mitteilungen der Pollichia*, 92: 107-130.
- Schimmel R., 2006b. Neue Ampedini-, Physorhinini-, Denticollini- und Diminae-Arten sowie eine neue Gattung, *Poggiellus* n. gen., aus Südostasien (Insecta: Coleoptera, Elateridae). *Mitteilungen der Pollichia*, 92: 131-158.
- Schimmel R., 2015a. – Neue Arten aus der Gattung *Penia* Castelnau, 1838, aus China. In: Diehl P., Imhoff A. & Möller L. (Eds.), Wissensgesellschaft Pfalz – 90 Jahre Pfälzische Gesellschaft zur Förderung der Wissenschaften. Pfälzische Gesellschaft zur Förderung der Wissenschaften Schriftenreihe 1927-2015, Bd. 116, pp. 367-375.
- Schimmel R., 2015b. – New species of the genera *Csikia*, *Limoniscus*, *Neopsephus* and *Penia* from China (Insecta: Coleoptera: Elateridae). *Vernate*, 34: 285-298.

Résumé

Qiu L., 2021. – Une remarquable espèce nouvelle du genre *Penia* (Coleoptera: Elateridae: Dimini) décrite de Guizhou, Chine. *Faunitaxys*, 9(37) : 1 – 4.

Penia jiangi n. sp. est décrite de la province de Guizhou, Chine. L'habitus et les caractères définissant cette nouvelle espèce sont illustrés.

Mots-clés. – Coleoptera, Elateroidea, Dendrometrinae, *Penia*, taupin, morphologie, taxonomie, nouvelle espèce, Guizhou, Chine.

Derniers articles publiés

- Fleck G., 2020. – *Onychogomphus* (*Siriusonychogomphus*) *louissiriusi*, a new species and new subgenus from Thailand (Odonata: Anisoptera: Gomphidae). *Faunitaxys*, 8(7) : 1 – 9.
- Huchet J-B., Romé D. & Touroult J., 2020. – *Hybosorus illigeri* Reiche, 1853, première mention pour les Petites Antilles (Coleoptera, Scarabaeoidea, Hybosoridae). *Faunitaxys*, 8(8) : 1 – 7.
- Huchet J-B., 2020. – *Chiron elegans*, nouvelle espèce d'Afrique subsaharienne (Coleoptera : Scarabaeoidea : Chironidae). *Faunitaxys*, 8(9) : 1 – 3.
- Keith D., 2020. – Description d'une troisième espèce dans le genre *Pantolasius* Lansberge, 1887 (Coleoptera : Scarabaeoidea, Hybosoridae). *Faunitaxys*, 8(10) : 1 – 2.
- Coache A. & Rainon B., 2020. – Contribution à la connaissance des Cassidinae du Bénin (Coleoptera, Chrysomelidae). *Faunitaxys*, 8(11) : 1 – 53.
- Gerstmeier R., 2020. – *Trogodendron bartolozzii*, a new species of Cleridae from Australia (Coleoptera: Cleridae: Clerinae). *Faunitaxys*, 8(12) : 1 – 2.
- Keith D., 2020. – Description d'espèces nouvelles du genre *Miridiba* Reitter, 1902 (Coleoptera: Scarabaeidae, Melolonthinae, Rhizotrogini). *Faunitaxys*, 8(13) : 1 – 5.
- Vives E., 2020. – Descripción de dos nuevos Lepturini del sudeste asiático (Coleoptera Cerambycidae). Notes on Lepturinae (20). *Faunitaxys*, 8(14) : 1 – 3.
- Théry T. & Sokolov A. V., 2020. – *Eucurtiopsis davaoensis* n. sp., a new Chlamydopsinae from Philippines (Coleoptera, Histeridae). *Faunitaxys*, 8(15) : 1 – 5.
- Bezark L. G. & Santos-Silva A., 2020. – Three new genera and three new species of American Cerambycidae (Coleoptera). *Faunitaxys*, 8(16) : 1 – 11.
- Devesa S. & Santos-Silva A., 2020. – A new species of *Pseudosparna* Mermudes & Monné, 2009 from Costa Rica (Coleoptera, Cerambycidae, Lamiinae). *Faunitaxys*, 8(17) : 1 – 5.
- Gomy Y., 2020. – Description de deux nouvelles espèces de *Cylistosoma* Lewis, 1905 de Madagascar (Coleoptera, Histeridae) (Septième contribution à la connaissance des Histeridae de Madagascar). *Faunitaxys*, 8(18) : 1 – 7.
- Porion T. & Audibert C., 2020. – Sur deux nouvelles espèces de Fulgoridae des Philippines (Hemiptera : Fulgoromorpha). *Faunitaxys*, 8(19) : 1 – 5.
- Bezark L. G., Santos-Silva A. & Devesa S., 2020. – New species of *Amphicnaeia* Bates, 1866, and key to species of the genus (Coleoptera, Cerambycidae, Lamiinae, Apomecynini). *Faunitaxys*, 8(20) : 1 – 13.
- Limoges R. & Le Tirant S., 2020. – Description d'une nouvelle espèce du genre *Eupholus* de Papouasie occidentale, Indonésie (Coleoptera, Curculionidae, Entiminae). *Faunitaxys*, 8(21) : 1 – 5.
- Gomy Y. & Tishechkin A., 2020. – Contribution à la connaissance des Histeridae de l'archipel du Vanuatu (Coleoptera). 3. *Faunitaxys*, 8(22) : 1 – 20.
- Huchet J-B., 2020. – Un nouveau *Phoberus* MacLeay, 1819, aptère du KwaZulu-Natal (Coleoptera : Scarabaeoidea : Trogidae). *Faunitaxys*, 8(23) : 1 – 5.
- Devesa S. & Santos-Silva A., 2021. – Description of two new species of Hemilophini (Coleoptera, Cerambycidae, Lamiinae). *Faunitaxys*, 9(1) : 1 – 6.
- Lin J.-Z., 2021. – Description of *Lucanus yulaoensis* sp. nov., a new species stag beetle from northern Taiwan (Coleoptera, Lucanidae). *Faunitaxys*, 9(2) : 1 – 5.
- Wappes J. E. & Santos-Silva A., 2021. – Descriptions, transference, notes and designation of lectotype in Rhinotragini (Coleoptera, Cerambycidae, Cerambycinae). *Faunitaxys*, 9(3) : 1 – 12.
- Gao H. R. & Liang L., 2021. – A new subspecies of *Trachythorax* Redtenbacher, 1908 (Phasmatodea: Necrosiinae) from Yunnan, China. *Faunitaxys*, 9(4) : 1 – 5.
- Ythier E. & Dupré G., 2021. – Description of a new species of *Hottentotta* Birula, 1908, from the Democratic Republic of the Congo (Scorpiones, Buthidae). *Faunitaxys*, 9(5) : 1 – 5.
- Delahaye N., Komiya Z., Drumont A. & Shapovalov A., 2021. – A new species of the genus *Psalidosphryon* Komiya, 2001 from West Papua, Indonesia (Coleoptera, Cerambycidae, Prioninae). *Faunitaxys*, 9(6) : 1 – 7.
- Lin J.-Z. & Chou W.-I., 2021. – Description of a new species of the genus *Neolucanus* Thomson, 1862 from Taiwan, with new localities record of *N. taiwanus* (Coleoptera, Lucanidae). *Faunitaxys*, 9(7) : 1 – 9.
- Wang Y., Ehrmann R. & Borer M., 2021. – A new species in the praying mantis genus *Rhombomantis* Ehrmann & Borer (Mantodea: Mantidae) from Indochina. *Faunitaxys*, 9(8) : 1 – 23.
- Devesa S., Lingafelter S. W. & Santos-Silva A., 2021. – New species of *Anelaphus* and *Poecilomallus* (Coleoptera, Cerambycidae, Cerambycinae, Elaphidiini) from Nicaragua. *Faunitaxys*, 9(9) : 1 – 6.
- Oremans P., Pyrcz T. & Zúbrík M., 2021. – Contribution à l'étude des *Euphaedra* de la République Centre Africaine et description d'une nouvelle espèce (Lepidoptera Nymphalidae). *Faunitaxys*, 9(10) : 1 – 4.
- Oremans P., 2021. – Une forme inédite de *Papilio Menestheus* de République de Côte d'Ivoire (Lepidoptera Papilionidae). *Faunitaxys*, 9(10) : 5 – 6.
- Ythier E., 2021. – Two new species of *Hadruiroides* Pocock, 1893 from Peru and Ecuador (Scorpiones, Caraboctonidae). *Faunitaxys*, 9(11) : 1 – 8.
- Vives E., 2021. – *Hesperoleptura* nuevo subgénero de Lepturini de las Islas Canarias (Coleoptera, Cerambycidae, Lepturinae). Notes on Lepturinae (21). *Faunitaxys*, 9(12) : 1 – 3.

- Lourenço W. R., 2021. – Une nouvelle espèce appartenant au genre *Buthus* Leach, 1815 (Scorpiones : Buthidae) collectée dans le Parc Naturel de la ‘Serra da Estrela’ au Centre du Portugal. *Faunitaxys*, 9(13) : 1 – 7.
- Lourenço W. R. & Velten J., 2021. – One more new genus and species of scorpion from Early Cretaceous Burmese amber (Scorpiones: Protoischnuridae). *Faunitaxys*, 9(14): 1 – 5.
- Théry T., 2021. – Description of *Eucurtiopsis marysae* n. sp., a singular species of Chlamydopsinae from the Philippines (Coleoptera, Histeridae). *Faunitaxys*, 9(15): 1 – 5.
- Cumming R. T. & Le Tirant S., 2021. – Review of the Cretaceous †Archaeatropidae and †Empheriidae and description of a new genus and species from Burmese amber (Psocoptera). *Faunitaxys*, 9(16): 1 – 11.
- Lourenço W. R., 2021. – Further comments on the elements of the family Palaeoburmesebuthidae Lourenço, 2015 with description of a new species of *Spinoburmesebuthus* Lourenço, 2017 from Early Cretaceous Burmite amber (Scorpiones). *Faunitaxys*, 9(17): 1 – 6.
- Zhao M. -Z., 2021. – On the genus *Cyphochilus* Waterhouse, 1867 from Hainan Island, China (Coleoptera: Scarabaeidae: Melolonthinae). *Faunitaxys*, 9(18): 1 – 8.
- Cliquennois N., 2021. – Description d’*Achrioptera hugeli*, nouvelle espèce de phasme de la Grande Comore (Phasmatodea, Achriopteridae). *Faunitaxys*, 9(19): 1 – 7.
- Lassalle B. & Roux P., 2021. – Contribution à la connaissance des *Pheropsophus* africains (Coleoptera, Caraboidea, Brachinidae). *Faunitaxys*, 9(20) : 1 – 12.
- Ythier E., Sadine S. E., Haddadi M. H. & Lourenço W. R., 2021. – A new species of *Buthus* Leach, 1815 from Algeria (Scorpiones: Buthidae) and an interesting new case of vicariance. *Faunitaxys*, 9(21): 1 – 9.
- Zhang Y. & Barclay M. V. L., 2021. – A remarkable new species of Prioninae (Coleoptera: Cerambycidae) from Guadalcanal, Solomon Islands. *Faunitaxys*, 9(22): 1 – 5.
- Qi Z.-H., 2021. – *Lucanus moae* sp. nov., a new species from Sichuan, China (Coleoptera: Lucanidae: Lucaninae). *Faunitaxys*, 9(23): 1 – 7.
- Chen Z.-T., 2021. – Checklist and key to the earwig genera from China (Dermaptera). *Faunitaxys*, 9(24): 1 – 8.
- Ythier E., 2021. – The southwesternmost scorpion species in Europe: *Buthus gabani* sp. n. from Cape St. Vincent, Algarve, Portugal (Scorpiones: Buthidae). *Faunitaxys*, 9(25): 1 – 6.
- Gomy Y., 2021. – Une anthropoducie accidentelle possible évitée à Madagascar (Coleoptera : Histeridae) (Huitième contribution à la connaissance des Histeridae de Madagascar). *Faunitaxys*, 9(26) : 1 – 3.
- Bian C.-Z. & Zhan Z., 2021. – A new species of *Lucanus* Scopoli, 1763 (Coleoptera, Lucanidae) from Tibet, China. *Faunitaxys*, 9(27): 1 – 4.
- Santos-Silva A. & Lingafelter S. W., 2021. – A new species of *Mephritus* Pascoe from Ecuador (Coleoptera, Cerambycidae, Cerambycinae, Elaphidiini). *Faunitaxys*, 9(28): 1 – 4.
- Wang J.-S., 2021. – A new species of the genus *Sinopanorpa* Cai & Hua, 2008 from Hubei Province, China (Mecoptera: Panorpidae). *Faunitaxys*, 9(29): 1 – 6.
- Lourenço W. R. & Velten J., 2021. – Early Cretaceous Burmite fossils of the genus *Chaerilobuthus* Lourenço & Beigel, 2011 (Scorpiones: Chaerilobuthidae) and description of a particular new species. *Faunitaxys*, 9(30): 1 – 5.
- Ythier E., 2021. – A new species of *Androctonus* Ehrenberg, 1828 from the Sahelian wooded steppes of Burkina Faso (Scorpiones: Buthidae). *Faunitaxys*, 9(31): 1 – 7.
- Gereys B., Coache A. & Filippi G., 2021. – Présence en France métropolitaine d’un frelon allochtone : *Vespa orientalis* Linnaeus, 1771 (Le Frelon oriental) (Hymenoptera, Vespidae, Vespinae). *Faunitaxys*, 9(32) : 1 – 5.
- Yu K., Zhang S., Zhang F., Li Z. & Yang Z., 2021. – Taxonomic studies on five species of Selenocosmiinae from China (Araneae, Theraphosidae). *Faunitaxys*, 9(33): 1 – 13.
- Vlasak J. & Santos-Silva A., 2021. – A new species of *Elaphidion* Audinet-Serville from the Antilles (Coleoptera, Cerambycidae, Cerambycinae, Elaphidiini). *Faunitaxys*, 9(34): 1 – 6.
- Écharoux D. & Roux P., 2021. – *Pheropsophus lisaee*, espèce nouvelle d’Éthiopie (Caraboidea, Brachinidae). *Faunitaxys*, 9(35) : 1 – 4.
- Santos-Silva A., 2021. – A new species of *Euderces* LeConte from Guatemala (Coleoptera, Cerambycidae, Cerambycinae). *Faunitaxys*, 9(36): 1 – 3.

Faunitaxys est échangée avec les revues suivantes (« print versions ») :

- Annali del Museo Civico di Storia Naturale G. Doria (Italie)
- Boletín de la Asociación española de Entomología (Espagne)
- Boletín de la Sociedad Andaluza de Entomología (Espagne)
- Bollettino del Museo di Storia Naturale di Venezia (Italie)
- Bulletin de la Société linnéenne de Lyon (France)
- Bulletin of Insectology (Italie)
- Heteropterus Rev. Entomol. (Espagne)
- Israel Journal of Entomology (Israël)
- Klapalekiana (République Tchèque)
- Koleopterologische Rundschau (Allemagne)
- Memorie del Museo Civico di Storia Naturale di Verona (Italie)
- Nova Supplementa Entomologica (Allemagne)
- Proceedings of the Entomological Society of Washington (USA)
- Revue suisse de Zoologie (Suisse)
- Spixiana (Allemagne)
- Stuttgarter Beiträge zur Naturkunde A, Biologie (Allemagne)
- Zoosystematica Rossica (Russie)

Faunitaxys

Volume 9, Numéro 37, Novembre 2021

SOMMAIRE

Une remarquable espèce nouvelle du genre *Penia* (Coleoptera: Elateridae: Dimini) décrite de Guizhou, Chine.

Lu Qiu 1 – 4

CONTENTS

A remarkable new *Penia* species (Coleoptera: Elateridae: Dimini) from Guizhou, China.

Lu Qiu 1 – 4

Illustration de la couverture: Maolan Nature Reserve, the type locality of *Penia jiangi* sp. nov.

Crédits:

© **Lu Qiu**: Fig. 1-4 & couverture.