



HAL
open science

Separate the wheat from the chaff: genomic scan for local adaptation in the red coral *Corallium rubrum*

Marine Pratlong, Anne Haguenaer, K. Brener, G. Mitta, E. Toulza, J. Garrabou, N. Bensoussan, P. Pontarotti, Didier Aurelle

► To cite this version:

Marine Pratlong, Anne Haguenaer, K. Brener, G. Mitta, E. Toulza, et al.. Separate the wheat from the chaff: genomic scan for local adaptation in the red coral *Corallium rubrum*. Peer Community Journal, 2021, 1, pp.e31. 10.24072/pcjournal.12 . hal-03454660v2

HAL Id: hal-03454660

<https://hal.science/hal-03454660v2>

Submitted on 29 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Peer Community Journal

Section: Evolutionary Biology

RESEARCH ARTICLE

Published
2021-11-26

Cite as

M. Pratlong, A. Haguenaier, K. Brener, G. Mitta, E. Toulza, J. Garrabou, N. Bensoussan, P. Pontarotti and D. Aurelle (2021) *Separate the wheat from the chaff: genomic scan for local adaptation in the red coral *Corallium rubrum**, Peer Community Journal, 1: e31.

Correspondence

didier.aurelle@univ-amu.fr

Peer-review

Peer reviewed and recommended by PCI Evolutionary Biology, <https://doi.org/10.24072/pci.evolbiol.100061>



This article is licensed under the Creative Commons Attribution 4.0 License.

Separate the wheat from the chaff: genomic scan for local adaptation in the red coral *Corallium rubrum*

M. Pratlong^{1,2}, A. Haguenaier^{1,3}, K. Brener⁴, G. Mitta⁴, E. Toulza⁴, J. Garrabou^{5,6}, N. Bensoussan⁷, P. Pontarotti^{#,2,8}, and D. Aurelle^{#,1,5,9}

Volume 1 (2021), article e31

<https://doi.org/10.24072/pcjournal.12>

Abstract

Genomic data allow an in-depth and renewed study of local adaptation. The red coral (*Corallium rubrum*, Cnidaria) is a highly genetically structured species and a promising model for the study of adaptive processes along an environmental gradient. Here, we used RAD-Sequencing in order to study the vertical genetic structure of this species and to search for signals of local adaptation to depth and thermal regime in the red coral. Previous studies have shown different thermotolerance levels according to depth in this species which could correspond to genetic or environmental differences. We designed a sampling scheme with six pairs of shallow vs deep populations distributed in three geographical regions as replicates. Our results showed significant differentiation among locations and among sites separated by around 20 m depth. The tests of association between genetics and environment allowed the identification of candidate loci under selection but with a potentially high rate of false positive. We discuss the methodological obstacles and biases encountered for the detection of selected loci in such a strongly genetically structured species. On this basis, we also discuss the significance of the candidate loci for local adaptation detected in each geographical region and the evolution of red coral populations along environmental gradients.

¹Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE, Marseille, France, ²Aix Marseille Univ, CNRS, Centrale Marseille, I2M, Equipe Evolution Biologique et Modélisation, Marseille, France, ³CRIOBE - Perpignan, USR3278 CRIOBE EPHE-CNRS-UPVD, France, ⁴IHPE, Univ Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Perpignan, France, ⁵Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille, France, ⁶Institute of Marine Sciences (ICM-CSIC), Barcelona, 08003, Spain, ⁷IPSO FACTO, SCOPArI, Pole Océanologie, Marseille, 13001, France, ⁸Aix Marseille Univ, IRD, APHM, Microbe, Evolution, Phylogénie, Infection, IHU Méditerranée Infection, Marseille, France, ⁹Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, 57 rue Cuvier, 75005 Paris, France, #Equal contribution

Peer Community Journal is a member of the
Centre Mersenne for Open Scientific Publishing
<http://www.centre-mersenne.org/>



CENTRE
MERSENNE

Introduction

The study of the mechanisms of adaptation by species to their local environment is of great interest in evolutionary biology. The interaction between environmental conditions, biological traits and evolutionary factors (selection, drift, migration and mutation) will shape the relative importance of genetic and plastic responses for each species facing heterogeneous environmental conditions. If selection is predominant, and if the environmental gradient is persistent for an extended period of time, each local population exposed to local selection could become genetically adapted to the corresponding local environment (Kawecki & Ebert, 2004; Gagnaire & Gaggiotti, 2016). An organism can also cope with local environmental conditions via plasticity or acclimatization, whereby a given genotype develops during its lifetime morphological or physiological responses (DeWitt et al., 1998; Pigliucci, 2001). Although particular situations favoring local adaptation or acclimatization are documented, it is often difficult to disentangle the effects of these two mechanisms and establish their relative contributions to adaptability (Palumbi et al., 2014). In addition, understanding these mechanisms has a fundamental interest in the current context of climate change for improving predictive models and proposing management strategies (Mumby et al., 2011; Gagnaire & Gaggiotti, 2016).

The evolution of adaptive processes depends on the interaction between different processes, including local selection and gene flow. Gene flow can hinder local adaptation through the input in a population of potentially maladapted individuals (migration load; Lenormand, 2002). Conversely, several theoretical studies have shown that gene flow can counteract the effects of genetic drift and promote local adaptation (Hastings and Rohlf, 1974; Felsenstein, 1975; Slatkin & Maruyama, 1975; Nagylaki, 1978; Alleaume-Benharira et al., 2006). The use of high throughput sequencing renewed the study of local adaptation. Various bottom – up approaches are now available to study local adaptation through the identification signals of selection along the genome (Barrett & Hoekstra, 2011). In the marine realm, such studies have been conducted at very large scale on highly dispersive teleost species (Bradbury et al., 2010; Limborg et al., 2012; Wang et al., 2013; Milano et al., 2014; Bernardi et al., 2016; Guo et al., 2016), and on benthic invertebrates with a highly dispersive, planctonic larvae stage (Chu et al., 2014; Bay & Palumbi, 2014; Araneda et al., 2016; Benestan et al., 2016). Marine species with high genetic structure are less frequent than more dispersive ones, and genomic studies of local adaptation in such species are still scarce (see Bongaerts et al., 2017 for a recent example). The study of local adaptation in a context of high genetic structure may also be difficult from a methodological point of view: high average F_{ST} values can lead to a high number of false positives in outlier tests for the detection of selection by the corresponding increase in the variance of F_{ST} values (Bierne et al., 2013; Hoban et al., 2016). Furthermore, in a context of high average genomic differentiation, it could be difficult to identify selected loci with a higher differentiation than expected under the neutral model. Finally, if genetic drift is strong, it can generate outlier loci with apparent correlation with an environmental variable outside any selective effect (Kawecki & Ebert, 2004; Hofer et al., 2009; Coop et al., 2010). Therefore the empirical study of local adaptation in such situation remains often challenging and with few empirical data in the marine realm.

Marine coastal environments offer particularly interesting conditions for studies of local adaptation, because of the gradual changes in environmental conditions along coastline at small scale, the more or less gradual vertical changes from shallow to deep water and the patchy distribution of contrasted habitats at different scales (Sanford & Kelly, 2011; Lundgren et al., 2013; Wrange et al., 2014). This interest promoted studies of local adaptation in coastal ecosystems (Ayre, 1995; Ulstrup & Van Oppen, 2003; Smith et al., 2007; Sherman & Ayre, 2008; Barshis et al., 2010; Bongaerts et al., 2011; Barshis et al., 2013; Lundgren et al., 2013; Kersting et al., 2013; Haguenaer et al., 2013; Ziegler et al., 2014; Palumbi et al., 2014; Bay & Palumbi, 2014; Ledoux et al., 2015; Pivotto et al., 2015; Jin et al., 2016; Bongaerts et al., 2017). Studying the genetic basis of local adaptation and the connectivity between habitats, could also give some information on the response to climate change (e.g. Bongaerts et al., 2017). In this context genome scans are powerful approaches to explore adaptive processes in natural populations (Manel et al., 2016).

The red coral (*Corallium rubrum*) is an asymbiotic (without *Symbiodinium*) temperate octocoral distributed from 5 to 1016 m depth in the Mediterranean sea and the near Atlantic (Boavida et al., 2016; Knittweis et al., 2016). It is a sessile and long-living species (more than 100 years), with low growth and

recruitment rates (Marschal et al., 2004; Santangelo et al., 2012). This species is included in two international conventions (annex III of the Bern Convention and annex V of the European Union Habitats directive), and its harvesting is regulated by national legislations (Ledoux et al., 2013). The study of a few microsatellite loci has demonstrated a strong genetic structure in this species (Ledoux et al., 2010a; Ledoux, et al., 2010b). The shallowest populations, above the seasonal thermocline, are exposed to high maximum temperatures and to frequent and intense thermal fluctuations in summer (Haguenaer et al., 2013). The intensity and frequency of extreme thermal events decrease with depth, and the deepest populations are exposed to stable thermal regimes. Since the observation of mass mortality events affecting this species during thermal anomalies in 1999 and 2003, the thermotolerance of the red coral has been intensively studied in the region of Marseille (France; Garrabou et al., 2001, 2009). Common garden experiments highlighted differences in polyp activities, calcification rate, necrosis rate and expression of HSP70 between shallow and deep individuals (10 or 20 m compared to 40 m depth) facing thermal stress (Torrents et al., 2008 ; Ledoux et al. 2015; Haguenaer et al., 2013). Transcriptomes of individuals from 5 and 40 m were compared and several genes were detected as differentially expressed without the application of any stress (Pratlong et al., 2015). These results suggested the possibility of local adaptation to depth in this species, but the possibility of environmental effects could not be excluded.

Together, these studies highlighted phenotypic differences in thermotolerance levels between individuals from different depths in Marseille, with shallower individuals more tolerant than deeper ones. Nevertheless we still do not know if these differences are the result of local adaptation or of individual acclimatization, or both. Previous works on this species enabled us to have a precise idea of the geographic scale at which local adaptation may occur, and were useful to optimize our sampling design. Because populations from different regions may have evolved similar responses to thermal stress, through similar or different genetic basis, it is interesting to investigate local adaptation in pairs of 'shallow vs deep' populations exposed to contrasted thermal regimes in distinct geographical regions (Jones et al., 2012; Hoban et al., 2016). Finally, the study of the genetic structure of this species would be useful to better understand the potential role of deeper populations in reseeding shallower ones following disturbances (Bongaerts et al., 2017).

Here we applied Restriction site Associated DNA sequencing (RAD-Seq) to individuals from pairs of 'shallow vs. deep' populations in three geographical regions of the Mediterranean Sea. The goal of this study was to characterize the neutral and adaptive genomic variation in this species and to test the possibility of local adaptation to depth through a genome scan approach. Our results enable us to discuss the neutral genetic structure of the red coral. Then we highlight the methodological obstacles expected in the detection of local adaptation in this context. Finally, we discuss the robustness of the candidates of local adaptation detected in each geographical region.

1. Material and Methods

Sampling and DN extraction

Corallium rubrum colonies were collected by scuba diving at two depths of two sites in three geographical regions (Marseille, Banyuls, Corsica) between February and August 2013 (Fig. S1, Table 1). Red coral populations from these three regions correspond to different genetic clusters according to microsatellites (Ledoux et al., 2010b) and RAD-Seq (see results). The two depths of each site presented contrasted thermal regimes with higher mean, maximum and standard deviation of temperature at shallower depths (surveys from March 2012 to October 2014; Table 2). Samples from the two depths at each site will be referred as shallow and deep. The three geographical regions presented different annual variations of temperatures between the two studied depths: a difference of 3.8 °C between the maximum observed at the two depths in Marseille, 1.7 °C in Corsica and 0.5 °C in Banyuls (Table 2). Thirty individuals per site and depth were collected (total 360 individuals), preserved in 95 % ethanol and stored at -20 °C until DNA extraction. Total genomic DNA was extracted according to the protocol of Sambrook et al. (1989), followed by a purification using Qiagen DNeasy blood and tissue spin columns (Qiagen). Genomic DNA concentration was quantified using a Qubit 2.0 Fluorometer (Life Technologies).

RAD-Sequencing

Twelve RAD libraries were prepared according to the protocol described in Etter et al. (2011), with small modifications. Briefly, 1 µg of genomic DNA for each sample was digested using high-fidelity PstI during 60 min at 37 °C. P1 adapters, with 4-6 bp individual barcodes were then ligated to each sample using 0.5 µL of T4 DNA ligase (NEB), 0.5 µL of rATP 100 mM (Promega), 1 µL of DTT 500 mM (Promega), 1 µL of 10X T4 ligase buffer (NEB) and incubated during 60 min at 22 °C, 10 min at 65 °C and 1 min at 64 °C. Individual samples were pooled by 32 (generally by location), sheared, size selected and P2-barcoded. Final PCR for RAD-tags enrichment were performed with 16 cycles and primers dimers were removed during a final AMPure Beads Purification (Agencourt). Libraries were sequenced on an Illumina HiSeq2000 using 100 bp single-end reads, at the Biology Institute of Lille (IBL, UMR 8199 CNRS) and at the MGX sequencing platform in Montpellier (France).

The STACKS pipeline (Catchen et al., 2011, 2013) was used for the loci de novo assembly and genotyping. Quality filtering and demultiplexing were performed with the process_radtags module with default parameters which enables to remove any read with uncalled base and to perform a phred-33 quality filtering of raw reads. Exact-matching RAD loci (putative orthologous tags) were individually assembled using ustacks with a minimum depth of coverage of five reads per allele ($m = 5$) and a maximum of five nucleotide mismatches between allele ($M = 5$). These parameters were optimized during preliminary runs. Cstacks was used to build a catalog of consensus loci from all individuals, with five mismatches allowed between individuals at the same locus ($n = 5$). Matches of individual RAD loci to the catalog of loci were searched using sstacks. Finally, the population module was used to obtain the loci that were successfully genotyped in at least 75 % of individuals from all populations. We observed an increase in the number of SNPs from position 86 bp to 91 bp and we removed these positions from the analysis which were due to sequencing problems. In order to filter for poor-quality SNPs and artifacts due paralogous sequences, we used VCFtools (Danecek et al., 2011) to remove SNPs that were not at the Hardy-Weinberg equilibrium within at least one of the 12 populations with a p-value threshold of 0.01. SNPs with a minor allele frequencies below 0.01 were removed using VCFtools. Individuals with more than 30 % of missing genotypes were discarded. Finally, only the first SNP of each RAD locus was kept for further analysis. The whole dataset has been previously used for the study of sex determinism in *C. rubrum* (Pratlong et al., 2017); we develop here the study of genetic structure and local adaptation.

Diversity and neutral genetic structure

Global F_{IS} over alleles and expected heterozygosity were estimated using GENEPOP and ARLEQUIN v.3.5 (Rousset, 2008; Excoffier and Lischer, 2010). The *C. rubrum* genetic structure was first analyzed by principal component analysis (PCA) using the package adegenet in R (Jombart, 2008; R Core Team, 2016). This analysis was performed on the total dataset (12 populations) and inside each of the three studied geographical regions (four populations in each). The dataset was centered and missing data were replaced by the mean allele frequency for each locus (<http://adegenet.r-forge.r-project.org/files/tutorial-basics.pdf>). In a second step, we performed a Bayesian population clustering implemented in the program STRUCTURE v.2.3.4 (Pritchard et al., 2000; Falush et al., 2003, 2007; Hubisz et al., 2009). We performed ten independent replicates from $K = 1$ to 10 with a burn-in of 50 000 and a number of MCMC iterations after burn-in of 100 000, with the model allowing for admixture and correlated allele frequencies. We calculated the ΔK statistic of Evanno et al. (2005) to help in the choice of the most appropriate number of genetic clusters but we also considered different K values. We used CLUMPAK to summarize the STRUCTURE results from the ten independent runs (Kopelman et al., 2015). The global and pairwise populations F_{ST} and exact tests for population differentiation were computed with GENEPOP 4.0.10 (Rousset, 2008). The correlation between the spatial distance between the two depths of the same site and the corresponding population pairwise F_{ST} was tested with the correlation test of Spearman implemented in R (R Core Team, 2016). Finally, we conducted an analysis of molecular variance (AMOVA) in ARLEQUIN v.3.5 (Excoffier and Lischer, 2010) with 10 000 permutations. The hierarchy for this analysis was chosen to follow the three geographical regions of our samples (Marseille, Corsica and Banyuls). This choice was justified by the PCA on the overall dataset. Finally, we performed the PCA and F_{ST} calculation using a dataset comprising only putatively neutral SNPs (without the SNPs detected as outliers by F_{ST} outlier methods, see below).

Detection of local adaptation

In order to search for loci potentially involved in local adaptation, we first used BayeScEnv (Villemerieul & Gaggiotti, 2015). This method identifies F_{ST} outlier loci that show a relationship between genetic differentiation and environmental differentiation. Runs were performed using default parameters, except the number of pilot runs that was set at 40. The maximal temperature recorded in each site was used as environmental variable (Table 2). We tested other descriptors of the thermal regime and we got similar results (data not shown). The convergence of runs was checked with the Gelman and Rubin's diagnostic using the R package coda (Plummer et al., 2006).

Second, we searched for F_{ST} outliers among red coral populations using ARLEQUIN v.3.5 (Hofer et al., 2009; Excoffier & Lischer, 2010). Because hierarchical genetic structures are known to lead to a high number of false positives in the search of outlier loci (Hofer et al., 2009), we performed this analysis independently in the three geographical regions in order to down a level in the structure. With this method, a distribution of F_{ST} across loci as a function of heterozygosity between populations is obtained by performing simulations under a hierarchical island model (two depths in one site and two sites in one geographical region). Outliers were identified as loci being in the tails of the generated distribution ($p < 0.01$). Outliers detected by ARLEQUIN could be false positives or the result of a selective pressure independent of depth. Therefore, we selected among these candidate loci, those linked with depth differentiation by searching, inside each geographical regions, loci with significant differences in genotypic frequencies between depths according to a χ^2 test. Significant differences were identified by using a false discovery rate of 0.05 (Benjamini & Hochberg, 1995).

Finally, we used the R package pcadapt to search for outliers loci by taking into account population structure and individual admixture (Luu et al., 2017). This method is recommended in cases of hierarchical genetic structure for a better control of the false positive rate. By identifying outliers loci linked with a particular principal component, pcadapt enabled us to focus on candidates linked with our biological question. From the pcadapt analyses, we selected outliers candidates linked with the relevant principal components with a q-value cutoff of 0.01.

Functional annotation and enrichment tests

The RAD tags were aligned on the red coral transcriptome (Pratlong et al., 2015) using the Burrows-Wheeler Alignment Tool (BWA; Li & Durbin, 2009). Blast2GO was used for the annotation of resulting contigs and functional enrichment analysis (Conesa et al., 2005). First, a blastp was first performed on the NCBI nr database with an e-value threshold of 10^{-10} (Altschul et al., 1990). Then, Blast2GO retrieved Gene Ontology (GO) terms associated with the obtained BLAST hits. Finally, in order to identify function potentially over-represented in outliers, we performed an enrichment analysis using a Fisher's exact test corrected using a false discovery rate of 0.05 (Benjamini & Hochberg, 1995).

Table 1. Characteristics of red coral sampling sites.

Population	Geographic region	Site	Depth (m)	GPS	GPS
FIG8	Marseille	MarseilleSouth	8	43° 12.330'N	5° 26.790'E
MOR40	Marseille	MarseilleSouth	40	43° 12.060'N	5° 27.100'E
ELV12	Marseille	MarseilleNorth	12	43° 19.780'N	5° 14.210'E
MEJ40	Marseille	MarseilleNorth	40	43° 19.700'N	5° 13.480'E
BANN20	Banyuls	BanyulsNorth	25	42° 26.890'N	3° 10.330'E
BANN40	Banyuls	BanyulsNorth	35	42° 26.890'N	3° 10.330'E
BANS20	Banyuls	BanyulsSouth	26	42° 26.390'N	3° 10.790'E
BANS40	Banyuls	BanyulsSouth	36	42° 26.390'N	3° 10.790'E
POR20	Corsica	Porto	21	42° 16.292'N	8° 41.255'E
POR40	Corsica	Porto	33	42° 16.292'N	8° 41.255'E
GAL20	Corsica	Galeria	26	42° 28.210'N	8° 38.950'E
GAL40	Corsica	Galeria	36	42° 28.210'N	8° 38.950'E

Table 2. Temperatures (in °C) characteristics of the sampling sites from March 2012 to October 2014. The corresponding regions are indicated by a letter: M – Marseille, B – Banyuls, C - Corsica

	Depth (m)	Minimum	Maximum	Mean	GPS
FIG8 (M)	8	12.63	26.92	17.03	3.52
MOR40 (M)	40	12.73	23.06	15.40	2.11
ELV12 (M)	12	11.81	26.70	16.60	3.24
MEJ40 (M)	40	11.86	22.87	15.29	2.18
BANN20 (B)	25	12.22	24.29	17.20	2.63
BANN40 (B)	35	9.41	23.83	14.49	2.45
BANS20 (B)	26	12.22	24.29	17.20	2.63
BANS40 (B)	36	9.41	23.83	14.49	2.45
POR20 (C)	21	12.51	25.91	17.51	3.41
POR40 (C)	33	12.56	23.83	16.26	2.45
GAL20 (C)	26	12.46	25.09	17.13	3.13
GAL40 (C)	36	12.56	23.83	16.26	2.45

Results

RAD-Sequencing and genotyping

An average of 191 ± 21 millions of reads by library was obtained after sequencing. After the demultiplexing and cleaning processes of the STACKS's process_radtags module, an average of 180 ± 22 millions of reads by library was obtained. From these reads, we were able to assemble 138 810 unique consensus RAD-tags present in at least 75 % of our 360 individuals. After all quality filter steps (Table 3), 27 461 SNPs were available. The mean proportion of missing data per locus before filtering for MAF 0.01 was 4.22 % (standard error : 2.78%). Finally, we removed six individuals presenting more than 30 % of missing data (one individual from the MEJ40 population, two from the BANN40 population, two from the GAL20 population and one from the GAL40 population). Our final dataset used for further analysis consisted in 359 individuals genotyped on 27 461 SNPs.

Table 3. Counts of SNP loci after each filtering step.

Step	Number of SNPs	Software
After assembly raw data	138 810	Stacks (Catchen et al., 2011, 2013)
Excluding loci not in within population HWE	86 520	VCftools (Danecek et al., 2011)
MAF 1 %	56 844	VCftools (Danecek et al., 2011)
One SNP per RAD-tag	27 461	

Genetic diversity

Multilocus values of the F_{IS} ranged between 0.005 (ELV12) and 0.065 (BANS40) (Table 4). Expected heterozygosity varied from 0.12 (GAL20) to 0.18 (all populations of Marseille) (Table 4). Populations of Marseille had higher values of expected heterozygosity than populations from Corsica and Banyuls ($p = 0.02$, Wilcoxon–Mann–Whitney test).

Population structure analysis

The positioning of individuals with respect to the first two principal components reflected the geographical, and, partly, the depth origin of the individuals (Fig. 1A). Individuals from Marseille and from Banyuls formed two clear and homogeneous groups while individuals from the two sites of Corsica formed two different groups with an important distance between them on the second axis. The first PCA axis explained 7.28 % of the total genotypic variance and separated individuals from Marseille from individuals from Banyuls and Corsica. The second axis explained 4.4 % of the total genetic variance and separated individuals from the Porto site in Corsica from other individuals. The fifth axis of the PCA separated all individuals according to their sex, independently from their geographical origin, as observed in a previous study (Pratlong et al., 2017). This axis can be explained by loci diagnostic of sex, heterozygous in females and homozygous in males (see Pratlong et al., 2017). Concerning PCA inside geographical regions, individuals from the two sites of Corsica and Marseille (north and south) were

separated along the first axis (13.41 % and 6.74 % of the total genetic variance respectively; Fig. 1B and 1C). The second axis (2.77 % of the total genetic variance) separated populations from the two depths of the two sites of Marseille. In Corsica and Banyuls, no PCA axis showed clear association with depth. Individuals from the two depths of the Galeria (GAL) site of Corsica were separated along the second axis (4.02 % of the total genetic variance) but this was not the case for individuals from the two depths of the Porto site (POR). Individuals from Banyuls showed much less structure than individuals from Marseille and Corsica (Fig. 1D). The first axis (2.99 % of the total genetic variance) separated individuals from the two sites (north and south). The second axis (2.46 % of the total genetic variance) separated individuals according to their sex (Pratlong et al., 2017). The PCA on the overall dataset and inside each geographical region gave similar results when only putatively neutral SNPs were considered (Fig. S2).

The delta(K) criterion (Evanno et al., 2005) indicated $K = 2$ as the most informative number of clusters for the STRUCTURE analysis. We present here the results for $K = 2$ to $K = 4$ which captured the main information (Fig. 2). In all cases, all clusters corresponded to the main geographical boundaries and the two depths of each site always clustered together. For $K = 2$, a clear separation between the Marseille regions and the Corsica / Banyuls regions was observed, confirming the separation of populations along the first PCA axis (Fig. 1). The clustering at $K = 3$ separated the three geographical regions in 7/10 replicates, and the remaining replicates grouped either one or the other Corsican sites with Banyuls populations (Fig. S3). Finally, $K = 4$ separated the two Corsican sites.

Table 4. Summary statistics of genetic diversity. The nucleotide diversities and numbers of private alleles are based on the dataset with all SNPs for each RAD-loci, excluding those with the lowest quality, and the removal of loci with minor allele frequencies below 0.01 (total 56 844 SNPs). Measures of F_{IS} and expected heterozygosities were based on the final dataset comprising 27 461 SNPs (with one SNP / locus). The corresponding regions are indicated by a letter: M – Marseille, B – Banyuls, C – Corsica.

Population	Samplesize	Nucleotide diversity	Average number of private alleles	F_{IS}	Expected heterozygosity
BANN20 (B)	30	0.172	0	0.018	0.15
BANN40 (B)	28	0.174	0	0.012	0.15
BANS20 (B)	30	0.173	0	0.019	0.15
BANS40 (B)	30	0.172	0	0.065	0.13
ELV12 (M)	30	0.192	0.001	0.005	0.17
MEJ40 (M)	29	0.195	0	0.053	0.18
FIG8 (M)	30	0.192	0.001	0.005	0.18
MOR40 (M)	30	0.198	0.001	0.036	0.18
GAL20 (C)	28	0.124	0.002	0.013	0.09
GAL40 (C)	29	0.149	0.001	0.019	0.13
POR20 (C)	30	0.165	0	0.023	0.13
POR40 (C)	30	0.168	0.006	0.009	0.15

The overall multilocus F_{ST} of the total dataset was 0.13. Pairwise F_{ST} values ranged from 0.01 (BANS20 vs BANS40, BANS20 vs BANN40 and BANN40 vs BANS40) to 0.24 (ELV12 vs GAL20 and FIG8 vs GAL20; Table 5). The exact test of genetic differentiation was highly significant for all pairwise comparisons ($p < 0.001$), even for the Banyuls and Galeria populations separated by 10 m depth ($F_{ST} = 0.022$ for BANN20 vs BANN40, $F_{ST} = 0.012$ for BANS20 vs BANS40 and $F_{ST} = 0.10$ for GAL20 vs GAL40). Considering the F_{ST} between depths, high F_{ST} values can be observed for different loci and different samples comparisons (Fig. S4). The average F_{ST} between the two depths of the same site was 0.04 in Marseille, 0.08 in Corsica and 0.02 in Banyuls (0.04 for the total dataset). Considering only putatively neutral loci (see below for outliers loci), the overall F_{ST} of the total dataset was 0.12 and pairwise F_{ST} values ranged from 0.01 (BANS20 vs BANS40, BANS20 vs BANN40 and BANN40 vs BANS40) to 0.23 (ELV12 vs GAL20) (Table S1). There was no correlation between the distance between two depths of the same site and the corresponding population pairwise F_{ST} ($p = 0.75$). We obtained a similar result ($p = 1$) if we removed the four populations of Marseille whose sampling sites for the two considered depths were not exactly the same (653 m horizontal distance between FIG8 and MOR40 and 995 m between ELV12 and MEJ40). Finally, the F_{ST} between the two shallow sites inside a geographical region was in all three cases higher than those between the two deep sites of the same region (0.10 vs 0.058 in Marseille, 0.20 vs 0.14 in Corsica and 0.025 vs 0.014 in Banyuls. This difference was significant in each of the three regions

($p < 1.10^{-16}$ with a t-test in all cases). In a similar way, the F_{ST} between two shallow sites of two different geographical regions were in all cases higher than those between the two corresponding deep sites, except for the comparisons between the Porto sites and the Banyuls sites ($p = 0.38$ and $p = 0.02$ for the POR/BANN and POR/BANS comparisons respectively; $p < 1.10^{-16}$ for the other comparisons).

The AMOVA indicated a similar percentage of the molecular variance explicated by differences among group and within groups (7.8 and 7.07 % respectively) and approximately 85 % of variance explicated by differences within populations (Table 6). There was significant genetic differentiation at the three studied levels ($F_{ST} = 0.15$, $F_{SC} = 0.08$, $F_{CT} = 0.08$; $p < 0.001$ in the three cases).




Figure 1. Principal component analysis (Axes 1 and 2) of A) the 12 red coral populations ($n = 354$ individuals, 27 461 SNPs), B) the four red coral populations from Marseille ($n = 119$ individuals, 27 461 SNPs), C) the four red coral populations from Corsica ($n = 117$ individuals, 27 461 SNPs), D) the four red coral populations from Banyuls ($n = 118$ individuals, 27 461 SNPs).




Figure 2. Results from Bayesian individual clustering with STRUCTURE for $K = 2$ to $K = 4$. For $K = 2$ and $K = 4$, all ten replicates produced the same structure. For $K = 3$, the major mode presented here was the result of 7/10 replicates. Minor modes are presented in Fig. S3.

Table 5. Pairwise F_{ST} estimates. All comparisons were highly significant. Intra-region comparisons are highlighted.

Population	BAN 20	BANN 40	BANS 20	BANS 40	ELV 12	MEJ 40	FIG 8	MOR 40	GAL 20	GAL 40	POR 20	POR 40
BANN20	-											
BANN40	0.02	-										
BANS20	0.03	0.01	-									
BANS40	0.03	0.03	0.01	-								
ELV12	0.13	0.13	0.13	0.13	-							
MEJ40	0.11	0.11	0.11	0.11	0.03	-						
FIG8	0.14	0.13	0.13	0.13	0.10	0.08	-					
MOR40	0.11	0.10	0.10	0.10	0.08	0.06	0.05	-				
GAL20	0.18	0.17	0.17	0.17	0.24	0.22	0.24	0.21	-			
GAL40	0.11	0.10	0.10	0.10	0.19	0.16	0.18	0.15	0.10	-		
POR20	0.14	0.13	0.13	0.13	0.17	0.16	0.17	0.14	0.20	0.13	-	
POR40	0.14	0.13	0.13	0.13	0.17	0.15	0.17	0.14	0.21	0.14	0.05	-

Table 6. Percent of the variation explained by grouping populations according to their geographical region on the analysis of molecular variance (performed with ARLEQUIN).

Source of variation	d.f.	Percentage of variation
Among groups	2	7.80
Among populations within groups	9	7.07
Within populations	696	85.13

Outlier SNPs

We identified 82 outliers with BayeScEnv. However, we noticed that all these outliers seemed to be driven by the divergence between particular populations, with one allele being always fixed in one or several populations without logical association with depth. ARLEQUIN detected 563 loci potentially under selection in Marseille, 869 in Corsica and 397 in Banyuls. Among these SNPs, all corresponded to a signal of divergent selection in Marseille and Banyuls, 207 of the 869 candidate loci corresponded to a signal of balanced selection in Corsica and the remaining 662 loci corresponded to a signal of divergent selection. Considering only these outliers, the overall F_{ST} of the total dataset was 0.25 and pairwise F_{ST} values ranged from 0.02 (BANS20 vs BANS40) to 0.42 (GAL20 vs POR40) (Table S2). The 207 loci potentially under balanced selection in Corsica were linked with sex differentiation and were not further analyzed here (Pratlong et al. 2017). Eight outlier SNPs were detected both in Marseille and Banyuls, 12 both in Marseille and Corsica and 12 both in Corsica and Banyuls. No SNP was detected as potentially under divergent selection and common in the three regions. The complementary χ^2 test of homogeneity of

genotypic frequencies between depths inside each region detected 162 candidate loci in Marseille, 1 371 in Corsica and 3 in Banyuls. Among these loci, 35, 248 and 2 were also respectively detected with the ARLEQUIN analysis. The numbers of outlier loci were correlated with the variance and the average of F_{ST} values inside each geographical regions (correlation coefficient of 0.97). The second axis of the PCA using the Marseille individuals showed apparent association with depth and appeared to be influenced by the variation of the candidates for local adaptation to depth detected by ARLEQUIN: 51 % of these loci were in the top 1 % of the contributions to this second axis, and 86 % were in the top 5 %.

Because the Marseille region is the only one presenting a principal component linked with depth (Fig. 1C), the *pcadapt* results obtained for the Corsica et Banyuls regions have poor biological relevance for our biological question. We chose thus to present only the *pcadapt* results obtained for the Marseille region. *Pcadapt* detected 58 outlier loci linked with the second PCA axis, the one which was linked to depth differentiation. All these candidates were detected by the ARLEQUIN analyses and 20 were also common with the χ^2 test presented above.

Functional annotation

Among the 27 461 analyzed RAD-tags, 6 376 had hits on the red coral transcriptome (23.2 %). Concerning SNPs detected as outliers by ARLEQUIN and contributing to the depth divergence, 8 on the 35 detected in Marseille had hits on the transcriptome, 46 on the 248 detected in Corsica and 2 on the 2 detected in Banyuls (Table S3). We did not observe any GO term enriched in coding regions among candidate SNPs, nor any functional enrichment in these outliers.

Discussion

Genetic diversity and structure

Our results confirm at a genomic scale the high genetic structure of the red coral, which was observed with a small number of microsatellite loci (Costantini et al., 2007; Ledoux et al., 2010b). Here the differentiation was significant for all comparisons, even with low average F_{ST} in some cases (0.01 or 0.02 between depths in Banyuls). Such low but significant F_{ST} can be explained by the high number of loci used here, and by the high variance in F_{ST} values. Indeed for all comparisons some important F_{ST} were observed in the dataset (see below). This significant differentiation at all distances could be the consequence of reduced mean larval dispersal distance, despite a quite long pelagic larval duration estimated in aquarium (from 16 to 42 days; Martínez-Quintana et al., 2015). Genetic incompatibilities could also contribute to the observed differentiation at least for some loci (Kulmuni & Westram, 2017). Our analysis of the genetic structure of the red coral revealed several clusters mainly corresponding to the geographical distributions of this species. A high differentiation was observed here with PCA and STRUCTURE between the two sites of Corsica separated by around 22 km. This pattern of genetic structure could be explained by a putative barrier to gene flow (through currents or lack of suitable habitats) between the two Corsican sites, or it could also correspond to an historical separation of these populations: two lineages could then be present in Corsica, with one being related to Banyuls populations.

We reported here a significant vertical genetic structure between the two depths of the same site (populations separated by less than 20 m). This differentiation was also observed when outlier loci were removed indicating that it is also shaped by neutral processes (migration / drift) as suggested previously with microsatellites (e.g. Ledoux et al., 2010a). In a study of the vertical genetic structure of red coral in two western Mediterranean sites (Cap de Creus, Spain and Portofino, Italia), Costantini et al. (2011) observed a drop in connectivity around 40 – 50 m depth, with genetic diversity declining with depth, but our sampling scheme did not allow us to test this hypothesis. The study of the vertical genetic structure in corals is important in the context of climate change. As deeper populations may be less affected by climate change, they could possibly reseed shallower populations (Bongaerts et al., 2017). Nevertheless this possibility of reseeding depends on the connectivity or potential barriers between depths. A lack of connectivity could erroneously be inferred in cases of cryptic species (Pante et al., 2015). Contrary to Prada et al. (2008) who showed the existence of two cryptic lineages at two different depths in a tropical octocoral, the populations of red coral from the two studied depths clearly correspond here to the same species, and the differentiation between depth was lower than the differentiation between sites and

regions. This vertical structure may be the result of both inherent life history traits and environmental variables. Weinberg (1979) has described a negative geotropism for the planulae of *C. rubrum*, and Martínez-Quintana et al. (2015) demonstrated that this was an active behavior. Depending on the orientation of the substrate, this can limit the connectivity between depths. The seasonal stratification during larval emission (which can occur from June to September; Haguenaer A., pers. comm.) could also limit dispersal. According to these hypotheses the larval behavior and oceanographic factors should lead to the genome-wide neutral differentiation between depths. This differentiation is probably also shaped by drift induced by the small effective size of red coral populations (Ledoux et al., 2010a).

The horizontal genetic differentiation between the two shallow sites was higher than those between the two corresponding deep sites, inside and between geographical region. This suggests a higher connectivity or lower rate of genetic drift for deep populations compared to shallow ones. The repeated colonization of shallow depths from deeper ones or the higher harvesting pressure on shallow populations compared to deep one could enhance genetic drift as well (Rossi et al., 2008; Cannas et al., 2016). We did not observe here a reduction in expected heterozygosity for shallow populations, connectivity differences then seem to be more probable in explaining the observed differences of genetic differentiation. Interestingly, Rossi et al. (2008) observed a higher frequency of patches of red coral below 50 m compared to above 50 m: if such pattern is present in the area and depths considered here, then it could increase gene flow through stepping stones migration. The observed vertical and horizontal genetic structure could indicate reduced recolonization abilities following disturbances such as mortality event induced by heat waves. Nevertheless the observed genetic structure could also be shaped by colonization history and monopolization effect (Orsini et al., 2013). In this case, a disturbance leading to free habitats would facilitate recolonization from other populations.

Potential biases in the search of outlier loci

We observed high F_{ST} values for different loci and different sample comparisons, and not only between depths (Fig. S4). The methods used here to identify selected loci will most likely detect loci with strong effects (Pritchard & Di Rienzo, 2010; Gagnaire & Gaggiotti, 2016), and it is highly improbable to observe such a high number of selected loci. Both hierarchical genetic structure and high levels of differentiation are known to lead to a high number of false positives in genomic studies of local adaptation, as observed here (Bierne et al., 2013; Hoban et al. 2016).

The markers density obtained with RAD-Seq may be also insufficient to detect a RAD-tag in linkage disequilibrium with a selected locus (Lowry et al., 2017; McKinney et al., 2017; Catchen et al., 2017). With a genome size of about 500 Mb for the red coral (Ganot et al., 2016), we expected a SNP sampling of 1 for 18 kb after the SNPs filtering steps (55 SNPs per Mb). Nevertheless, a potential high linkage disequilibrium may be present here because of drift, and we thus expected that our RAD-tags to at least detect a signal of genetic adaptation, which remains to be confirmed. Approaches dedicated to the study of polygenic adaptation (Daub et al., 2013) or to the genomic distribution of F_{ST} or nucleotide diversity (Hohenlohe et al., 2010) could be interesting here, but a reference genome is still lacking for the red coral. On a more theoretical point of view, outlier loci could be linked to intrinsic genetic incompatibilities whose allelic frequencies coupled with environmental barriers (Bierne et al., 2011). The frequency of genetic incompatibilities in marine populations is largely unknown but probably under-estimated (Plough et al., 2016). Even if not directly linked to local adaptation, such loci are important factors in the evolution of red coral populations.

Local adaptation to depth in the red coral

We focused on candidate loci meeting the following criteria: i) detection with ARLEQUIN and pcadapt, ii) significant differentiation between depth, iii) function relevant to the adaptation to thermal regime. These loci are the most relevant as factors of local adaptation. The absence of candidate SNPs common to the three geographical regions could indicate that the adaptation to comparable shallow environmental pressures in these independent regions are based on different genetic pathways, or on non-genetic mechanisms (Putnam and Gates, 2015). However, most candidate loci should be in linkage disequilibrium with selected loci, and such association can easily be lost between distant locations through recombination. Differences in the strength of selective pressure in the three regions could also explain the differences in the detected loci. In Marseille we evidenced a clear signal of differentiation

between depths according to multivariate and outlier loci analyses. This detection of a signal of local adaptation in the Marseille region is consistent with the observations from studies of thermotolerance differences in this region (Torrents et al., 2008; Haguenaer et al., 2013; Ledoux et al., 2015; Pratlong et al., 2015). In the case of the Marseille region there are then strong evidences of the existence of adaptive differentiation at a scale of few tens of meters only. Concerning Corsican populations, Ledoux et al. (2015) reported no phenotypic signal of local adaptation after reciprocal transplant experiment. Here, the most promising candidate for the adaptation to thermal regime, was an homologous to an allene oxide synthase-lipoxygenase which is known to be involved in the response to thermal stress in octocorals (Löhelaid et al., 2015). This indirect argument would support the presence of local adaptation in this area as well, but more experimental analyzes will be necessary to confirm the involvement of this function in adaptation to thermal stress in this species. Finally, the detection of a reduced number of candidate loci (two) in the Banyuls region would be consistent with the weaker selective pressure here (see above, Table 2).

Previously, we have sequenced the transcriptome of individuals from the two depths of the Marseille site studied here in Marseille (Pratlong et al., 2015). Several genes were differentially expressed between individuals from the two depths outside thermal stress conditions. Some of these genes, such as those from the Tumor Necrosis Factors Receptor Associated Factors (TRAF) family, have been identified as involved in the response to thermal stress in the hexacoral *Acropora hyacinthus* (Barshis et al., 2013). However, none of these differentially expressed genes were identified in our RAD-Seq study. Additionally the differences of expression may result from acclimatization, and are not necessarily adaptive. The use of a reference genome would be useful here as well to study the potential link between candidate SNPs and genes location and function (Manel et al., 2016).

Conclusion

To our knowledge, the red coral presents among the highest levels of differentiation among studies of local adaptation through genome scans approaches in marine environment (Bradbury et al., 2010; Limborg et al., 2012; Wang et al., 2013; Chu et al., 2014; Milano et al., 2014; Bay and Palumbi, 2014; Bernardi et al., 2016; Araneda et al., 2016; Guo et al., 2016; Benestan et al., 2016; Bongaerts et al., 2017). This study enabled us to empirically emphasize the limitations in the detection and the interpretation of signals of local adaptation using usual statistical methods in this strongly structured species. Both neutral and adaptive divergence highlighted here demonstrate the genetic singularity of shallow populations of the red coral, especially in the Marseille region where the shallowest populations of this species are found. Together, the strong genetic structure we observed between shallow populations, the low dispersal abilities of the red coral and the local adaptation of these individuals to the highly variable thermal conditions they experiment, raise strong concerns about the evolution of shallow populations and the possibility of loss of adaptive variations in case of mortality events. Extending the genomic study initiated here would be useful to study the evolution of this species in heterogeneous and changing environments. Whatever their origin (genetic or environmental), the different thermotolerance levels observed between depths and populations in the red coral should also be taken into account in future studies of adaptive evolution in this species. These results would also be useful for the management of this harvested species which is present in two international conservation conventions.

Acknowledgements

We thank Nicolas Fernandez and Béatrice Loriod from the Marseille TGML platform for their invaluable help and advice with the preparation of the RAD libraries; the team of the MGX platform for the sequencing of the RAD libraries. We thank the molecular biology service of the IMBE, the informatic service of the Pytheas Institute, and Frédéric Zuberer of the Pytheas Institute for his support in sampling. We thank the Scandola Natural Reserve, especially Jean-Marie Dominici. We also thank Manuela Carenzi, Pierre-Alexandre Gagnaire, François Bonhomme and Nicolas Bierne for stimulating discussions. Three reviewers, Guillaume Achaz, Lucas Gonçalves da Silva, and an anonymous one, greatly helped us to

improve this article. A previous version of this article has been reviewed and recommended by Peer Community In Evolutionary Biology (<https://doi.org/10.24072/pci.evolbiol.100061>).

Funding

This work is a contribution to the Labex OT-Med (n° ANR-11-LABX-0061) funded by the French Government “Investissements d’Avenir” program of the French National Research Agency (ANR) through the A*MIDEX project (n° ANR-11-IDEX-0001-02). This project has been funded by the ADACNI program of the French National Research Agency (ANR) (project n°ANR-12- ADAP-0016; <http://adacni.imbe.fr>). We thank the ECCOREV Research Federation (FR 3098) for the financial support of part of this study. The project leading to this publication has received funding from European FEDER Fund under project 1166-39417. The authors thank the UMR 8199 LIGAN-PM Genomics platform (Lille, France, especially Véronique Dhennin) which belongs to the 'Federation de Recherche' 3508 Labex EGID (European Genomics Institute for Diabetes; ANR-10-LABX-46) and was supported by the ANR Equipex 2010 session (ANR-10-EQPX-07- 01; 'LIGAN-PM'). The LIGAN-PM Genomics platform (Lille, France) is also supported by the FEDER and the Region Nord-Pas-de-Calais-Picardie. This study is set within the framework of the « Laboratoire d’Excellence (LabEx) » TULIP (ANR-10-LABX-41).

Conflict of interest disclosure

The authors declare that they have no financial conflict of interest with the content of this article.

Data, script and code availability

The raw DNA sequences are available in the Short Read Archive (SRA) database under accession no. SRR5186771-SRR5187129. The filtered SNP file is available from the Dryad Digital Repository: <http://doi.org/10.5061/dryad.rs7bm>.

References

- Alleaume-Benharira M, Pen I, Ronce O (2006) Geographical patterns of adaptation within a species’ range: interactions between drift and gene flow. *Journal of evolutionary biology*, **19**, 203–215. <https://doi.org/10.1111/j.1420-9101.2005.00976.x>
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of molecular biology*, **215**, 403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Araneda C, Larraín MA, Hecht B, Narum S (2016) Adaptive genetic variation distinguishes Chilean blue mussels (*Mytilus chilensis*) from different marine environments. *Ecology and Evolution*, **6**, 3632–3644. <https://doi.org/10.1002/ece3.2110>
- Ayre DJ (1995) Localized adaptation of sea anemone clones: evidence from transplantation over two spatial scales. *Journal of animal ecology*, 186–196. <https://doi.org/10.2307/5754>
- Barrett RD, Hoekstra HE (2011) Molecular spandrels: tests of adaptation at the genetic level. *Nature Reviews Genetics*, **12**, 767–780. <https://doi.org/10.1038/nrg3015>
- Barshis DJ, Ladner JT, Oliver TA, Seneca FO, Traylor-Knowles N, Palumbi SR (2013) Genomic basis for coral resilience to climate change. *Proceedings of the National Academy of Sciences*, **110**, 1387–1392. <https://doi.org/10.1073/pnas.1210224110>
- Barshis DJ, Stillman JH, Gates RD, Toonen RJ, Smith LW, Birkeland C (2010) Protein expression and genetic structure of the coral *Porites lobata* in an environmentally extreme Samoan back reef: does host genotype limit phenotypic plasticity? *Molecular Ecology*, **19**, 1705–1720. <https://doi.org/10.1111/j.1365-294X.2010.04574.x>
- Bay RA, Palumbi SR (2014) Multilocus Adaptation Associated with Heat Resistance in Reef-Building Corals. *Current Biology*, **24**, 2952–2956. <https://doi.org/10.1016/j.cub.2014.10.044>
- Benestan L, Quinn BK, Maaroufi H, Laporte M, Clark FK, Greenwood SJ, Rochette R, Bernatchez L (2016) Seascape genomics provides evidence for thermal adaptation and current-mediated population

- structure in American lobster (*Homarus americanus*). *Molecular Ecology*, **25**, 5073–5092. <https://doi.org/10.1111/mec.13811>
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, **57**, 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Bernardi G, Azzurro E, Golani D, Miller MR (2016) Genomic signatures of rapid adaptive evolution in the bluespotted cornetfish, a Mediterranean Lessepsian invader. *Molecular ecology*, **25**, 3384–3396.
- Bierne N, Roze D, Welch JJ (2013) Pervasive selection or is it...? why are FST outliers sometimes so frequent? *Molecular Ecology*, **22**, 2061–2064. <https://doi.org/10.1111/mec.12241>
- Bierne N, Welch, John J, Loire E, Bonhomme F, David P (2011) The coupling hypothesis: why genome scans may fail to map local adaptation genes. *Molecular Ecology*, **20**, 2044–2072. <https://doi.org/10.1111/j.1365-294X.2011.05080.x>
- Boavida J, Paulo D, Aurelle D, Arnaud-Haond S, Marschal C, Reed J, Goncalves JM, Serrao EA (2016) A well-kept treasure at depth: precious red coral rediscovered in Atlantic deep coral gardens (SW Portugal) after 300 years. *PloS one*, **11**, e0147228. <https://doi.org/10.1371/journal.pone.0147228>
- Bongaerts P, Riginos C, Brunner R, Englebert N, Smith SR, Hoegh-Guldberg O (2017) Deep reefs are not universal refuges: Reseeding potential varies among coral species. *Science Advances*, **3**, e1602373. <https://doi.org/10.1126/sciadv.1602373>
- Bongaerts P, Riginos C, Hay KB, van Oppen MJ, Hoegh-Guldberg O, Dove S (2011) Adaptive divergence in a scleractinian coral: physiological adaptation of *Seriatopora hystrix* to shallow and deep reef habitats. *BMC evolutionary biology*, **11**, 1–15. <https://doi.org/10.1186/1471-2148-11-303>
- Bradbury IR, Hubert S, Higgins B, Borza T, Bowman S, Paterson IG, Snelgrove PV, Morris CJ, Gregory RS, Hardie DC (2010) Parallel adaptive evolution of Atlantic cod on both sides of the Atlantic Ocean in response to temperature. *Proceedings of the Royal Society B: Biological Sciences*, **277**, 3725–3734. <https://doi.org/10.1098/rspb.2010.0985>
- Cannas R, Sacco F, Cau A, Cuccu D, Follesa MC, Cau A (2016) Genetic monitoring of deep-water exploited banks of the precious Sardinia coral *Corallium rubrum* (L., 1758): useful data for a sustainable management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **26**, 236–250. <https://doi.org/10.1002/aqc.2522>
- Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH (2011) Stacks: building and genotyping loci de novo from short-read sequences. *G3: Genes, genomes, genetics*, **1**, 171–182. <https://doi.org/10.1534/g3.111.000240>
- Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Molecular Ecology*, **22**, 3124–3140. <https://doi.org/10.1111/mec.12354>
- Catchen JM, Hohenlohe PA, Bernatchez L, Funk WC, Andrews KR, Allendorf FW (2017) Unbroken: RADseq remains a powerful tool for understanding the genetics of adaptation in natural populations. *Molecular Ecology Resources*, **17**, 362–365. <https://doi.org/10.1111/1755-0998.12669>
- Chu ND, Kaluziak ST, Trussell GC, Vollmer SV (2014) Phylogenomic analyses reveal latitudinal population structure and polymorphisms in heat stress genes in the North Atlantic snail *Nucella lapillus*. *Molecular Ecology*, **23**, 1863–1873. <https://doi.org/10.1111/mec.12681>
- Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*, **21**, 3674–3676.
- Coop G, Witonsky D, Di Rienzo A, Pritchard JK (2010) Using environmental correlations to identify loci underlying local adaptation. *Genetics*, **185**, 1411–1423. <https://doi.org/10.1534/genetics.110.114819>
- Costantini F, Fauvelot C, Abbiati M (2007) Genetic structuring of the temperate gorgonian coral (*Corallium rubrum*) across the western Mediterranean Sea revealed by microsatellites and nuclear sequences. *Molecular Ecology*, **16**, 5168–5182. <https://doi.org/10.1111/j.1365-294X.2007.03579.x>
- Costantini F, Rossi S, Pintus E, Cerrano C, Gili J-M, Abbiati M (2011) Low connectivity and declining genetic variability along a depth gradient in *Corallium rubrum* populations. *Coral Reefs*, **30**, 991–1003. <https://doi.org/10.1007/s00338-011-0771-1>
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>

- Daub JT, Hofer T, Cutivet E, Dupanloup I, Quintana-Murci L, Robinson-Rechavi M, Excoffier L (2013) Evidence for polygenic adaptation to pathogens in the human genome. *Molecular Biology and Evolution*, **30**, 1544–1558. <https://doi.org/10.1093/molbev/mst080>
- De Villemereuil P, Gaggiotti OE (2015) A new FST-based method to uncover local adaptation using environmental variables. *Methods in Ecology and Evolution*, **6**, 1248–1258. <https://doi.org/10.1111/2041-210X.12418>
- DeWitt TJ, Sih A, Wilson DS (1998) Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, **13**, 77–81. [https://doi.org/10.1016/S0169-5347\(97\)01274-3](https://doi.org/10.1016/S0169-5347(97)01274-3)
- Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F, English CA, Galindo HM, Grebmeier JM, Hollowed AB, Knowlton N, Polovina J, Rabalais NN, Sydeman WJ, Talley LD (2011) Climate Change Impacts on Marine Ecosystems. *Annual Review of Marine Science*, **4**, 11–37. <https://doi.org/10.1146/annurev-marine-041911-111611>
- Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA (2011) SNP discovery and genotyping for evolutionary genetics using RAD sequencing. *Methods in Molecular Biology* (Clifton, N.J.), **772**, 157–178. https://doi.org/10.1007/978-1-61779-228-1_9
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, **14**, 2611–20. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Excoffier L, Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, **10**, 564–567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Falush D, Stephens M, Pritchard JK (2003) Inference of Population Structure Using Multilocus Genotype Data: Linked Loci and Correlated Allele Frequencies. *Genetics*, **164**, 1567–1587. <https://doi.org/10.1093/genetics/164.4.1567>
- Falush D, Stephens M, Pritchard JK (2007) Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Molecular Ecology Notes*, **7**, 574–578. <https://doi.org/10.1111/j.1471-8286.2007.01758.x>
- Felsenstein J (1975) Genetic drift in clines which are maintained by migration and natural selection. *Genetics*, **81**, 191–207. <https://doi.org/10.1093/genetics/81.1.191>
- Gagnaire P-A, Gaggiotti OE (2016) Detecting polygenic selection in marine populations by combining population genomics and quantitative genetics approaches. *Current Zoology*, **62**, 603–616. <https://doi.org/10.1093/cz/zow088>
- Ganot P, Fritz M, Rausch T, Aurelle D, Haguenaer A, Romans P, Zoccola D, Tambutté E, Aranda M, Benes V, Allemand D, Tambutté S (2016) Calcification in *Corallium rubrum*: dissecting the process using genomics and transcriptomics together with scalpels and scissors. In *13th International Coral Reef Symposium, Honolulu, Hawai'i*.
- Garrabou J, Perez T, Sartoretto S, Harmelin J (2001) Mass mortality event in red coral *Corallium rubrum* populations in the Provence region (France, NW Mediterranean). *Marine Ecology Progress Series*, **217**, 263–272. <https://doi.org/10.3354/meps217263>
- Guo B, Li Z, Merilä J (2016) Population genomic evidence for adaptive differentiation in the Baltic Sea herring. *Molecular Ecology*, **25**, 2833–2852. <https://doi.org/10.1111/mec.13657>
- Haguenaer A, Zuberer F, Ledoux J-B, Aurelle D (2013) Adaptive abilities of the Mediterranean red coral *Corallium rubrum* in a heterogeneous and changing environment: from population to functional genetics. *Journal of Experimental Marine Biology and Ecology*, **449**, 349–357. <https://doi.org/10.1016/j.jembe.2013.10.010>
- Hastings A, Rohlfs FJ (1974) Gene flow: effect in stochastic models of differentiation. *The American Naturalist*, **108**, 701–705. <https://doi.org/10.1086/282947>
- Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, Poss ML, Reed LK, Storfer A, Whitlock MC (2016) Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions. *The American Naturalist*, **188**, 379–397. <https://doi.org/10.1086/688018>
- Hofer T, Ray N, Wegmann D, Excoffier L (2009) Large allele frequency differences between human continental groups are more likely to have occurred by drift during range expansions than by selection. *Annals of Human Genetics*, **73**, 95–108. <https://doi.org/10.1111/j.1469-1809.2008.00489.x>

- Hohenlohe PA, Bassham S, Etter PD, Stiffler N, Johnson EA, Cresko WA (2010) Population genomics of parallel adaptation in threespine stickleback using sequenced RAD tags. *PLoS genetics*, **6**, e1000862.
- Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with the assistance of sample group information. *Molecular Ecology Resources*, **9**, 1322–1332. <https://doi.org/10.1371/journal.pgen.1000862>
- Jin YK, Lundgren P, Lutz A, Raina J-B, Howells EJ, Paley AS, Willis BL, Van Oppen MJ (2016) Genetic markers for antioxidant capacity in a reef-building coral. *Science Advances*, **2**, e1500842. <https://doi.org/10.1126/sciadv.1500842>
- Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*, **24**, 1403–1405. <https://doi.org/10.1093/bioinformatics/btn129>
- Jones FC, Grabherr MG, Chan YF, Russell P, Mauceci E, Johnson J, Swofford R, Pirun M, Zody MC, White S (2012) The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, **484**, 55–61. <https://doi.org/10.1038/nature10944>
- Kawecki TJ, Ebert D (2004) Conceptual issues in local adaptation. *Ecology Letters*, **7**, 1225–1241. <https://doi.org/10.1111/j.1461-0248.2004.00684.x>
- Kersting DK, Bensoussan N, Linares C (2013) Long-term responses of the endemic reef-builder *Cladocora caespitosa* to Mediterranean warming. *PLoS One*, **8**, e70820. <https://doi.org/10.1371/journal.pone.0070820>
- Knittweis L, Aguilar R, Alvarez H, Borg JA, Evans J, Garcia S, Schembri PJ (2016) New depth record of the precious red coral *Corallium rubrum* for the Mediterranean. *Rapport du Congrès de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, **41**. <https://www.um.edu.mt/library/oar/handle/123456789/18340>
- Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, **15**, 1179–1191. <https://doi.org/10.1111/1755-0998.12387>
- Kulmuni J, Westram AM (2017) Intrinsic incompatibilities evolving as a by-product of divergent ecological selection: Considering them in empirical studies on divergence with gene flow. *Molecular Ecology*, **26**, 3093–3103. <https://doi.org/10.1111/mec.14147>
- Ledoux J-B, Aurelle D, Bensoussan N, Marschal C, Féral J-P, Garrabou J (2014) Potential for adaptive evolution at species range margins: contrasting interactions between red coral populations and their environment in a changing ocean. *Ecology and Evolution*, **5**(6), 1178–1192. <https://doi.org/10.1002/ece3.1324>
- Ledoux J-B, Aurelle D, Féral J-P, Garrabou J (2013) Molecular forensics in the precious Mediterranean red coral, *Corallium rubrum*: testing DNA extraction and microsatellite genotyping using dried colonies. *Conservation Genetics Resources*, **5**, 327–330. <https://doi.org/10.1007/s12686-012-9795-2>
- Ledoux J, Garrabou J, Bianchimani O, Drap P, Féral J-P, Aurelle D (2010) Fine-scale genetic structure and inferences on population biology in the threatened Mediterranean red coral, *Corallium rubrum*. *Molecular Ecology*, **19**, 4204–4216. <https://doi.org/10.1111/j.1365-294X.2010.04814.x>
- Ledoux J-B, Mokhtar-Jamaï K, Roby C, Féral J-P, Garrabou J, Aurelle D (2010) Genetic survey of shallow populations of the Mediterranean red coral [*Corallium rubrum* (Linnaeus, 1758)]: new insights into evolutionary processes shaping nuclear diversity and implications for conservation. *Molecular Ecology*, **19**, 675–690. <https://doi.org/10.1111/j.1365-294X.2009.04516.x>
- Lenormand T (2002) Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, **17**, 183–189. [https://doi.org/10.1016/S0169-5347\(02\)02497-7](https://doi.org/10.1016/S0169-5347(02)02497-7)
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, **25**, 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
- Limborg MT, Helyar SJ, De Bruyn M, Taylor MI, Nielsen EE, Ogden R, Carvalho GR, FPT Consortium, Bekkevold D (2012) Environmental selection on transcriptome-derived SNPs in a high gene flow marine fish, the Atlantic herring (*Clupea harengus*). *Molecular Ecology*, **21**, 3686–3703. <https://doi.org/10.1111/j.1365-294X.2012.05639.x>
- Löhelaid H, Teder T, Samel N (2015) Lipoxygenase-allene oxide synthase pathway in octocoral thermal stress response. *Coral Reefs*, **34**, 143–154. <https://doi.org/10.1007/s00338-014-1238-y>

- Lowry DB, Hoban S, Kelley JL, Lotterhos KE, Reed LK, Antolin MF, Storfer A (2017) Breaking RAD: an evaluation of the utility of restriction site-associated DNA sequencing for genome scans of adaptation. *Molecular Ecology Resources*, **17**, 142–152. <https://doi.org/10.1111/1755-0998.12635>
- Lundgren P, Vera J, Peplow L, Manel S, van Oppen M (2013) Genotype - environment correlations in corals from the Great Barrier Reef. *BMC Genetics*, **14**, 9. <https://doi.org/10.1186/1471-2156-14-9>
- Luu K, Bazin E, Blum MG (2017) pcadapt: an R package to perform genome scans for selection based on principal component analysis. *Molecular Ecology Resources*, **17**, 67–77. <https://doi.org/10.1111/1755-0998.12592>
- Manel S, Perrier C, Pratlong M, Abi-Rached L, Paganini J, Pontarotti P, Aurelle D (2016) Genomic resources and their influence on the detection of the signal of positive selection in genome scans. *Molecular Ecology*, **25**, 170–184. <https://doi.org/10.1111/mec.13468>
- Marschal C, Garrabou J, Harmelin J, Pichon M (2004) A new method for measuring growth and age in the precious red coral *Corallium rubrum* (L.). *Coral Reefs*, **23**, 423–432. <https://doi.org/10.1007/s00338-004-0398-6>
- Martínez-Quintana A, Bramanti L, Viladrich N, Rossi S, Guizien K (2015) Quantification of larval traits driving connectivity: the case of *Corallium rubrum* (L. 1758). *Marine Biology*, **162**, 309–318. <https://doi.org/10.1007/s00227-014-2599-z>
- McKinney GJ, Larson WA, Seeb LW, Seeb JE (2017) RADseq provides unprecedented insights into molecular ecology and evolutionary genetics: comment on Breaking RAD by Lowry et al.(2016). *Molecular Ecology Resources*, **17**, 356–361. <https://doi.org/10.1111/1755-0998.12649>
- Milano I, Babbucci M, Cariani A, Atanassova M, Bekkevold D, Carvalho GR, Espiñeira M, Fiorentino F, Garofalo G, Geffen AJ (2014) Outlier SNP markers reveal fine-scale genetic structuring across E uropean hake populations (*Merluccius merluccius*). *Molecular Ecology*, **23**, 118–135. <https://doi.org/10.1111/mec.12568>
- Mumby PJ, Elliott IA, Eakin CM, Skirving W, Paris CB, Edwards HJ, Enríquez S, Iglesias-Prieto R, Cherubin LM, Stevens JR (2011) Reserve design for uncertain responses of coral reefs to climate change. *Ecology Letters*, **14**, 132–140. <https://doi.org/10.1111/j.1461-0248.2010.01562.x>
- Nagylaki T (1978) Random genetic drift in a cline. *Proceedings of the National Academy of Sciences*, **75**, 423–426. <https://doi.org/10.1073/pnas.75.1.423>
- Orsini L, Vanoverbeke J, Swillen I, Mergeay J, Meester L (2013) Drivers of population genetic differentiation in the wild: isolation by dispersal limitation, isolation by adaptation and isolation by colonization. *Molecular Ecology*, **22**, 5983–5999. <https://doi.org/10.1111/mec.12561>
- Palumbi SR, Barshis DJ, Traylor-Knowles N, Bay RA (2014) Mechanisms of reef coral resistance to future climate change. *Science*, **344**, 895–898. <https://doi.org/10.1126/science.1251336>
- Pante E, Puillandre N, Viricel A, Arnaud-Haond S, Aurelle D, Castelin M, Chenuil A, Destombe C, Forcioli D, Valero M (2015) Species are hypotheses: avoid connectivity assessments based on pillars of sand. *Molecular Ecology*, **24**, 525–544. <https://doi.org/10.1111/mec.13048>
- Pigliucci M (2001) Phenotypic plasticity: beyond nature and nurture. JHU Press.
- Pivotto I, Nerini D, Masmoudi M, Kara H, Chaoui L, Aurelle D (2015) Highly contrasted responses of Mediterranean octocorals to climate change along a depth gradient. *Royal Society Open Science*, **2**, 140493. <https://doi.org/10.1098/rsos.140493>
- Plough L, Shin G, Hedgecock D (2016) Genetic inviability is a major driver of type III survivorship in experimental families of a highly fecund marine bivalve. *Molecular Ecology*, **25**, 895–910.
- Plummer M, Best N, Cowles K, Vines K (2006) CODA: Convergence diagnosis and output analysis for MCMC. *R news*, **6**, 7–11. <https://doi.org/10.1111/mec.13524>
- Prada C, Schizas NV, Yoshioka PM (2008) Phenotypic plasticity or speciation? A case from a clonal marine organism. *BMC Evolutionary Biology*, **8**, 1–19. <https://doi.org/10.1186/1471-2148-8-47>
- Pratlong M, Haguenaer A, Chabrol O, Klopp C, Pontarotti P, Aurelle D (2015) The red coral (*Corallium rubrum*) transcriptome: a new resource for population genetics and local adaptation studies. *Molecular Ecology Resources*, **15**, 1205–1215. <https://doi.org/10.1111/1755-0998.12383>
- Pratlong M, Haguenaer A, Chenesseau S, Brener K, Mitta G, Toulza E, Bonabaud M, Rialle S, Aurelle D, Pontarotti P (2017) Evidence for a genetic sex determination in Cnidaria, the Mediterranean red coral (*Corallium rubrum*). *Royal Society Open Science*, **4**. <https://doi.org/10.1098/rsos.160880>

- Pritchard JK, Di Rienzo A (2010) Adaptation—not by sweeps alone. *Nature Reviews Genetics*, **11**, 665–667. <https://doi.org/10.1038/nrg2880>
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of Population Structure Using Multilocus Genotype Data. *Genetics*, **155**, 945–959. <https://doi.org/http://dx.doi.org/10.1093/genetics/155.2.945>
- Putnam HM, Gates RD (2015) Preconditioning in the reef-building coral *Pocillopora damicornis* and the potential for trans-generational acclimatization in coral larvae under future climate change conditions. *The Journal of Experimental Biology*, **218**, 2365–2372. <https://doi.org/10.1242/jeb.123018>
- R Core Team (2015) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.
- Rossi S, Tsounis G, Orejas C, Padrón T, Gili J-M, Bramanti L, Teixidó N, Gutt J (2008) Survey of deep-dwelling red coral (*Corallium rubrum*) populations at Cap de Creus (NW Mediterranean). *Marine Biology*, **154**, 533–545. <https://doi.org/10.1007/s00227-008-0947-6>
- Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources*, **8**, 103–106. <https://doi.org/10.1111/j.1471-8286.2007.01931.x>
- Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning: a laboratory manual. Cold spring harbor laboratory press.
- Sanford E, Kelly MW (2010) Local Adaptation in Marine Invertebrates. *Annual Review of Marine Science*, **3**, 509–535. <https://doi.org/10.1146/annurev-marine-120709-142756>
- Santangelo G, Bramanti L, Rossi S, Tsounis G, Vielmini I, Lott C, Gili JM (2012) Patterns of variation in recruitment and post-recruitment processes of the Mediterranean precious gorgonian coral *Corallium rubrum*. *Journal of Experimental Marine Biology and Ecology*, **411**, 7–13. <https://doi.org/10.1016/j.jembe.2011.10.030>
- Sherman CD, Ayre DJ (2008) Fine-scale adaptation in a clonal sea anemone. *Evolution: International Journal of Organic Evolution*, **62**, 1373–1380. <https://doi.org/10.1111/j.1558-5646.2008.00375.x>
- Slatkin M, Maruyama T (1975) Genetic drift in a cline. *Genetics*, **81**, 209–222. <https://doi.org/10.1093/genetics/81.1.209>
- Smith L, Barshis D, Birkeland C (2007) Phenotypic plasticity for skeletal growth, density and calcification of *Porites lobata* in response to habitat type. *Coral Reefs*, **26**, 559–567. <https://doi.org/10.1007/s00338-007-0216-z>
- Szulkin M, Gagnaire P, Bierne N, Charmantier A (2016) Population genomic footprints of fine-scale differentiation between habitats in Mediterranean blue tits. *Molecular Ecology*, **25**, 542–558. <https://doi.org/10.1111/mec.13486>
- Torrents O, Tambutté E, Caminiti N, Garrabou J (2008) Upper thermal thresholds of shallow vs. deep populations of the precious Mediterranean red coral *Corallium rubrum* (L.): Assessing the potential effects of warming in the NW Mediterranean. *Journal of Experimental Marine Biology and Ecology*, **357**, 7–19. <https://doi.org/10.1016/j.jembe.2007.12.006>
- Ulstrup K, Van Oppen M (2003) Geographic and habitat partitioning of genetically distinct zooxanthellae (*Symbiodinium*) in *Acropora* corals on the Great Barrier Reef. *Molecular Ecology*, **12**, 3477–3484. <https://doi.org/10.1046/j.1365-294X.2003.01988.x>
- Wang L, Liu S, Zhuang Z, Guo L, Meng Z, Lin H (2013) Population genetic studies revealed local adaptation in a high gene-flow marine fish, the small yellow croaker (*Larimichthys polyactis*). *PloS one*, **8**, e83493. <https://doi.org/10.1371/journal.pone.0083493>
- Weinberg S (1979) The light-dependent behaviour of planula larvae of *Eunicella singularis* and *Corallium rubrum* and its implication for octocorallian ecology. *Bijdragen tot de Dierkunde*, **49**, 16–30. <https://doi.org/10.1163/26660644-04901002>
- Wrange A-L, André C, Lundh T, Lind U, Blomberg A, Jonsson PJ, Havenhand JN (2014) Importance of plasticity and local adaptation for coping with changing salinity in coastal areas: a test case with barnacles in the Baltic Sea. *BMC Evolutionary Biology*, **14**, 1–15. <https://doi.org/10.1186/1471-2148-14-156>
- Ziegler M, Roder CM, Büchel C, Voolstra CR (2014) Limits to physiological plasticity of the coral *Pocillopora verrucosa* from the central Red Sea. *Coral Reefs*, **33**, 1115–1129. <https://doi.org/10.1007/s00338-014-1192-8>

Supplementary information




Figure S1. Location of the sampling sites of the red coral among the three studied geographical regions.




Figure S2. Principal component analysis (Axes 1 and 2), using only putative neutral SNPs, of the A) 12 red coral populations ($n = 354$ individuals, 25 669 SNPs), B) four red coral populations from Marseille ($n = 119$ individuals, 26 898 SNPs), C) four red coral populations from Corsica ($n = 117$ individuals, 26 592 SNPs), D) four red coral populations from Banyuls ($n = 118$ individuals, 27 069 SNPs).




Figure S3. Results from Bayesian individual clustering with STRUCTURE for $K = 3$. The three figures correspond to major and minor modes detected.




Figure S4. Joint distribution of between-depths F_{ST} in the three geographical regions.

Table S1. Pairwise F_{ST} estimates using only putatively neutral SNPs. All comparisons were highly significant. Intra-region comparisons are highlighted.

Population	BAN 20	BANN 40	BANS 20	BANS 40	ELV 12	MEJ 40	FIG 8	MOR 40	GAL 20	GAL 40	POR 20	POR 40
BANN20	-											
BANN40	0.02	-										
BANS20	0.02	0.01	-									
BANS40	0.02	0.01	0.01	-								
ELV12	0.13	0.12	0.12	0.12	-							
MEJ40	0.11	0.10	0.10	0.10	0.03	-						
FIG8	0.13	0.12	0.12	0.12	0.09	0.07	-					
MOR40	0.10	0.10	0.09	0.09	0.07	0.05	0.04	-				
GAL20	0.17	0.16	0.16	0.16	0.23	0.21	0.22	0.20	-			
GAL40	0.11	0.09	0.09	0.09	0.18	0.16	0.18	0.15	0.10	-		
POR20	0.13	0.12	0.12	0.12	0.17	0.15	0.16	0.14	0.17	0.11	-	
POR40	0.13	0.12	0.12	0.12	0.16	0.15	0.16	0.14	0.18	0.12	0.04	-

Table S2. Pairwise F_{ST} estimates using only outlier SNPs from the ARLEQUIN analysis. All comparisons were highly significant. Intra-region comparisons are highlighted.

Population	BAN 20	BANN 40	BANS 20	BANS 40	ELV 12	MEJ 40	FIG 8	MOR 40	GAL 20	GAL 40	POR 20	POR 40
BANN20	-											
BANN40	0.04	-										
BANS20	0.06	0.03	-									
BANS40	0.06	0.03	0.02	-								
ELV12	0.19	0.18	0.19	0.18	-							
MEJ40	0.16	0.14	0.15	0.15	0.04	-						
FIG8	0.19	0.17	0.18	0.18	0.23	0.19	-					
MOR40	0.14	0.12	0.13	0.13	0.19	0.14	0.08	-				
GAL20	0.27	0.27	0.28	0.27	0.37	0.35	0.36	0.32	-			
GAL40	0.15	0.13	0.15	0.13	0.26	0.24	0.26	0.22	0.17	-		
POR20	0.23	0.21	0.21	0.21	0.26	0.22	0.22	0.18	0.41	0.30	-	
POR40	0.24	0.23	0.24	0.23	0.26	0.23	0.23	0.19	0.42	0.31	0.07	-

Table S3. Results of the annotation analysis of candidates for local adaptation in the three geographical regions.

	value	Description	GO	Region
Contig_16793	1.1E-117	LON peptidase N-terminal domain and RINGfinger partial	ubiquitin-protein transferase activity zinc ion binding metal ion binding ATP-dependent peptidase activity protein ubiquitination	
Contig_20016	0	Chromodomain-helicase-DNA-binding 1-like		
Contig_23068	2.8E-117	RNA-directed DNA polymerase from mobileelement jockey-like	nucleic acid phosphodiester bondhydrolysis RNA-directed DNA polymerase activity endonuclease activity RNA-dependent DNA biosyntheticprocess	Marseille
Contig_38936	4.1E-23	E3 ubiquitin-ligase DZIP	nucleic acid binding zinc ion binding	
Contig_44372	3.1E-61	PREDICTED : uncharacterized proteinLOC107346707	binding	
Contig_47492	7.8E-98	No description	metal ion binding oxidoreductase activity metabolic process oxidation-reduction process	
Contig_7346	2.3E-150	FAM46C-like		
Contig_10570	4.8E-37	PREDICTED : uncharacterized protein		
Contig_10731	2.8E-68	allene oxide synthase-lipoxygenase	oxidoreductase activity, acting on singledonors with incorporation of molecular oxygen, incorporation of two atoms of oxygen metal ion binding fatty acid metabolic process	
Contig_11268	2.3E-90	neuronal acetylcholine receptor subunit alpha-9	integral component of membraneplasma membrane part extracellular ligand-gated ion channelactivity ion transport synaptic transmission response to stimulus biological regulation	
Contig_11731	0	E3 ubiquitin-ligase RNF213	binding	
Contig_12920	5.5E-149	Stonin-2 isoform X2	cytoplasmic part vesicle-mediated transport single-organism process intracellular transport	Corsica
Contig_13771	4.0E-151	A-kinase anchor mitochondrial	Hemopoiesis cell differentiation	

Contig_16202	6.5E-65	PIN2 TERF1-interacting telomerase inhibitor 1	Nucleus chromosome intracellular organelle part nucleic acid binding regulation of cellular process
Contig_16843	0	Succinate-semialdehyde mitochondrial	aldehyde dehydrogenase (NAD) activity succinate-semialdehyde dehydrogenase (NAD+) activity succinate-semialdehyde dehydrogenase [NAD(P)+] activity glycosylceramide metabolic process multicellular organism development glutamine family amino acid metabolic process gamma-aminobutyric acid catabolic process dicarboxylic acid metabolic process oxidation-reduction process
Contig_16868	4.3E-24	glioma tumor suppressor candidate region gene 1-like	Membrane
Contig_17255	7.0E-53	Nanos 1	integral component of membrane
Contig_18282	1.3E-44	centromere K	zinc ion binding RNA binding
Contig_19099	0	epithelial growth factor receptor substrate 15-like partial	nucleus
Contig_19611	2.0E-134	DDB1- and CUL4-associated factor 5	Calcium ion binding
Contig_24102	1.6E-62	nuclease HARB1	mitochondrion CUL4-RING E3 ubiquitin ligase complex
Contig_24221	0	tRNA (guanine(26)-N(2))-diethyltransferase	Nucleus mitochondrion tRNA (guanine-N2)-methyltransferase activity binding tRNA N2-guanine methylation binding
Contig_31623	0	No description	Nucleus
Contig_32690	1.4E-163	ubiquitin carboxyl-terminal hydrolase isozyme L5	cytoplasm thio-dependent ubiquitin-specific protease activity binding ubiquitin-dependent protein catabolic process protein deubiquitination regulation of proteolysis forebrain development regulation of cellular protein metabolic process
Contig_33144	6.6E-56	Dok-7	protein kinase binding positive regulation of protein tyrosine

Contig_35407	3.8E-11	hypothetical protein AC249_AIPGENE14243	kinase activity insulin receptor-binding
Contig_36059	5.9E-47	cytosolic non-specific dipeptidase	exopeptidase activity metabolic process
Contig_36102	2.8E-135	nucleolar complex 4 homolog	Nuclear part Membrane cell part
Contig_37478	8.9E-28	fibroblast growth factor receptor 3	protein kinase activity phosphorylation positive regulation of phosphorylation regulation of primary metabolic process
Contig_38721	2.3E-41	L-seryl-tRNA(Sec) kinase	Phosphorylation kinase activity
Contig_38739	2.0E-56	Dr1	Ada2/Gcn5/Ada3 transcription activator complex DNA binding transcription corepressor activity TBP-class protein binding protein heterodimerization activity negative regulation of transcription from RNA polymerase II promoter transcription DNA-templated histone H3 acetylation
Contig_39033	7.1E-155	homeodomain transcription factor 1 isoform X1	intracellular membrane-bounded organelle nucleic acid phosphodiester bond hydrolysis
Contig_40626	1.0E-51	RNA-directed DNA polymerase from mobile element jockey-like	RNA-directed DNA polymerase activity RNA binding endonuclease activity RNA-dependent DNA biosynthetic process
Contig_41259	9.9E-45	serine protease 23-like	proteolysis serine-type endopeptidase activity serine-type peptidase activity peptidase activity
Contig_41360	8.9E-134	tonsoku	nuclear lumen protein complex DNA metabolic process cellular response to DNA damage stimulus single-organism metabolic process single-organism cellular process
Contig_41417	3.3E-27	eukaryotic translation initiator factor 4E-binding 1	Cytosol eukaryotic initiation factor 4E binding translational initiation signal transduction negative regulation of translational initiation
Contig_41500	1.8E-128	Cytoplasmic 1-like	Cytoskeleton plasma membrane focal adhesion

Contig_42135	0	No description	dense body membrane exopeptidase activity
Contig_42435	3.8E-84	Tensin-partial	focal adhesionactin binding cell-substrate junction assembly fibroblast migration
Contig_43083	2.0E-47	No description	early endosome
Contig_45623	0	kinesin KIF16B isoform X2	cytosol kinesin complex phosphatidylinositol-3,4,5-trisphosphate binding ATP-dependent microtubule motor activity, plus-end-directed phosphatidylinositol-3-phosphate binding phosphatidylinositol-3,4-bisphosphate binding phosphatidylinositol-3,5-bisphosphate binding formation of primary germ layer regulation of receptor recycling Golgi to endosome transport microtubule-based movement epidermal growth factor receptor signaling pathway endoderm developmentfibroblast growth factor receptor signaling pathway cytoskeleton-dependent intracellular transport receptor catabolic process early endosome to late endosome transport fibroblast growth factor receptor signaling pathway cytoskeleton-dependent intracellular transport receptor catabolic process early endosome to late endosome transport
Contig_45771	8.3E-94	No description	membrane dopamine neurotransmitter receptoractivity G-protein coupled receptor signalingpathway regulation of Rho protein signaltransduction metabolic process Rho guanyl-nucleotide exchange factoractivity transferase activity positive regulation of GTPase activity
Contig_47280	2.0E-168	No description	oxidoreductase activity, acting on singledonors with incorporation of molecular oxygen, incorporation of two atoms of oxygen

Contig_47344	0	allene oxide synthase-lipoxygenase	metal ion binding fatty acid metabolic process	
Contig_5993	1.5E-95	GA-binding subunit beta-partial	Nucleus protein homodimerization activity transcription regulatory region DNA binding protein heterodimerization activity transcription, DNA-templated positive regulation of transcription from RNA polymerase II promoter	
Contig_7742	0	Indole-3-acetaldehyde oxidase-like	oxidoreductase activity ion binding single-organism metabolic process	
Contig_8936	5.2E-105	drebrin isoform X1	actin filament binding neuron projection morphogenesis receptor-mediated endocytosis intracellular	
Contig_8963	3.4E-169	tubuline delta chain	actin binding	
Contig_26377	6.7E-138	ribokinase isoform X1	Intracellular part ribokinase activity D-ribose metabolic process carbohydrate phosphorylation	Banyuls