

## STABILITY OF EQUIVARIANT LOGARITHMIC TANGENT SHEAVES ON TORIC VARIETIES OF PICARD RANK TWO

Achim Napame

### ▶ To cite this version:

Achim Napame. STABILITY OF EQUIVARIANT LOGARITHMIC TANGENT SHEAVES ON TORIC VARIETIES OF PICARD RANK TWO. 2021. hal-03457163v1

## HAL Id: hal-03457163 https://hal.science/hal-03457163v1

Preprint submitted on 30 Nov 2021 (v1), last revised 27 Jul 2023 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# STABILITY OF EQUIVARIANT LOGARITHMIC TANGENT SHEAVES ON TORIC VARIETIES OF PICARD RANK TWO

#### ACHIM NAPAME

ABSTRACT. For an equivariant log pair (X, D) where X is a normal toric variety and D a reduced Weil divisor, we study slope-stability of the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$ . We give a complete description of divisors D and polarizations L such that  $\mathscr{T}_X(-\log D)$  is (semi)stable with respect to L when X has a Picard rank one or two.

#### 1. INTRODUCTION

The notion of slope-stability for vector bundles was first introduced by Mumford [Mum62] in his attempt to construct the moduli spaces of vector bundles over a curve. This notion was generalized in higher dimension by Takemoto [Tak72]. The study of stability of coherent sheaves over polarized varieties is a difficult problem. This problem is simplified when additional structures are added on the sheaf.

A quasi-coherent sheaf  $\mathscr{E}$  on a toric variety X with torus T is said to be an *equivariant sheaf* if it admits a lift of the T-action on X, which is linear on the stalks of  $\mathscr{E}$ . Klyachko in [Kly90] gave a complete classification of equivariant vector bundles over toric varieties in terms of a family of filtrations of vector spaces. This classification was generalized in the case of quasi-coherent sheaves on toric varieties by Perling in [Per04].

By using the equivariant structure of the tangent bundle, Hering-Nill-Süss in [HNS19] and Dasgupta-Dey-Khan in [DDK20] studied slope-stability of the tangent bundle TX of a smooth projective toric variety X of Picard rank one or two. Inspired by *Iitaka's philosophy*, in this paper, we extend the result of [HNS19] and [DDK20] to the case of log pairs (X, D). More precisely, if X is a normal toric variety and D a reduced Weil divisor such that the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$  is equivariant, we are interested by the set of polarizations L on X such that  $\mathscr{T}_X(-\log D)$  is (semi)stable with respect to L.

Assume that X is a Kähler manifold and  $D = \sum_i a_i D_i$  a simple normal crossing divisor with  $a_i \in [0; 1] \cap \mathbb{Q}$ . If  $K_X + D$  is ample, there exists a Kähler-Einstein metric  $\omega$  on the pair (X, D) by [BG16, Theorem C]; moreover, if D is reduced, then the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$  is semi-stable with respect to  $K_X + D$ by [Gue16, Theorem A]. If the pair (X, D) is Fano, by [Ber16, Section 4.3], the pair (X, D) admits a Kähler-Einstein metric  $\omega$  if for all  $i, 0 < a_i < 1$ . In this case, (X, D) is K-polysatble. According to [Li20, Theorem 1.4], the orbifold tangent sheaf  $\mathscr{T}_X(-\log D)$  is slope semi-stable with respect to  $-(K_X + D)$ .

In this paper, we are interested in the case where  $a_i \in \{0, 1\}$ . According to [BB13, Section 1.2.1 and Section 3], when (X, D) is Fano, we don't have a natural notion of Kähler-Einstein metric on (X, D). We will use another method to study the slope-stability of  $\mathscr{T}_X(-\log D)$ .

In Section 2, we recall some notions about toric varieties and equivariant sheaves. Let X be a normal toric variety over  $\mathbb{C}$  and T be its torus. We denote by  $\Sigma$  the fan of X and  $\Sigma(1)$  the set of rays of  $\Sigma$ . In Proposition 2.10, we show that : for a reduced Weil divisor D, the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$  is equivariant if and only if

$$D = \sum_{\rho \in \Delta} D_{\rho}$$

where  $\Delta$  is a subset of  $\Sigma(1)$  and  $D_{\rho}$  the Zariski closure of the orbit  $O(\rho)$  corresponding to the ray  $\rho$ . Using the family of multifiltrations defined in [Per04, Section 5], we give in Section 3.1 the family of multifiltrations corresponding to the equivariant sheaf  $\mathscr{T}_X(-\log D)$ . In Section 3.2, we give some conditions on  $\Sigma$  and  $\Delta$ , such that  $\mathscr{T}_X(-\log D)$  is decomposable. In Section 4, we recall some notions about slope-stability and we show that :

**Proposition 1.1.** Let X be a toric variety without torus factor and p be the rank of its class group Cl(X). If

$$+ p \le \operatorname{card}(\Delta) \le \dim(X) + p - 1$$

then  $\mathscr{T}_X(-\log D)$  is not semi-stable with respect to any polarizations.

1

By Corollary 3.6, if  $\Delta = \emptyset$ , then  $\mathscr{T}_X(-\log D)$  is the tangent sheaf  $\mathscr{T}_X$  and when  $\Delta = \Sigma(1), \ \mathscr{T}_X(-\log D)$  is isomorphic to the trivial vector bundle of rank dim X. If X has no torus factor, we have  $\operatorname{card}(\Sigma(1)) = \dim(X) + p$  by [CLS11, Theorem 4.1.3]. According to Proposition 1.1, it is therefore sufficient to study the stability of  $\mathscr{T}_X(-\log D)$  when  $\operatorname{card}(\Delta) \leq p$ . In this paper we will be interested in the case where  $p \in \{1, 2\}$ . As the case  $\operatorname{card}(\Delta) = 0$  corresponds to the tangent sheaf, we will study the slope-stability of  $\mathscr{T}_X(-\log D)$  when  $1 \leq \operatorname{card}(\Delta) \leq p$ .

In Section 4.4, we study slope-stability of  $\mathscr{T}_X(-\log D)$  when X is a toric orbifold of Picard rank one. If  $X = \mathbb{P}^n$  and  $\Delta \subset \Sigma(1)$ , we show that :

**Proposition 1.2.** If  $\operatorname{card}(\Delta) = 1$ , then  $\mathscr{T}_X(-\log D)$  is poly-stable but not stable with respect to any polarizations.

In the last sections (Sections 5 and 6) we study slope-stability of  $\mathscr{T}_X(-\log D)$  when X is a smooth toric variety of Picard rank 2. These parts form the core of this paper.

1.1. Smooth toric varieties of Picard rank two. Let X be a smooth toric variety of dimension n with fan  $\Sigma$  in  $\mathbb{R}^n$  such that  $\operatorname{rk}\operatorname{Pic}(X) = 2$ . By [CLS11, Theorem 7.3.7] due to Kleinschmidt (see [Kle88]), there exists  $r, s \in \mathbb{N}^*$  with r + s = n and  $a_1, \ldots, a_r \in \mathbb{N}$  with  $a_1 \leq a_2 \leq \ldots \leq a_r$  such that

(1) 
$$X = \mathbb{P}\left(\mathscr{O}_{\mathbb{P}^s} \oplus \bigoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^s}(a_i)\right)$$

We denote by  $\pi: X \longrightarrow \mathbb{P}^s$  the projection to the base  $\mathbb{P}^s$ . By [CLS11, Section 7.3] and [DDK20, Section 4.2], the rays of  $\Sigma$  are given by the half-lines generated by  $w_0, w_1, \ldots, w_s, v_0, v_1, \ldots, v_r$  where  $(w_1, \ldots, w_s)$  is the standard basis of  $\mathbb{Z}^s \times 0_{\mathbb{Z}^r}$ ,  $(v_1, \ldots, v_r)$  the standard basis of  $0_{\mathbb{Z}^s} \times \mathbb{Z}^r$ ,

 $v_0 = -(v_1 + \ldots + v_r)$  and  $w_0 = a_1v_1 + \ldots + a_rv_r - (w_1 + \ldots + w_s)$ .

We denote by  $D_{v_i}$  the divisor corresponding to the ray  $\text{Cone}(v_i)$  and  $D_{w_j}$  the divisor corresponding to the ray  $\text{Cone}(w_j)$ .

**Notation 1.3.** Let  $\operatorname{Amp}(X) \subset N^1(X) \otimes_{\mathbb{Z}} \mathbb{R}$  be the *ample cone* of X. We define :

$$\operatorname{Stab}(\mathscr{T}_X(-\log D)) = \left\{ \begin{array}{l} L \in \operatorname{Amp}(X) : \mathscr{T}_X(-\log(D)) \\ \text{is stable with respect to } L \end{array} \right\}$$

and

$$\operatorname{sStab}(\mathscr{T}_X(-\log D)) = \left\{ \begin{array}{l} L \in \operatorname{Amp}(X) : \mathscr{T}_X(-\log(D)) \\ \text{is semi-stable with respect to } L \end{array} \right\}.$$

1.2. Main results. Let X be a smooth toric variety of Picard rank two and  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\mu) \otimes \mathscr{O}_X(\lambda)$  be a polarization on X. We set  $\nu = \frac{\mu}{\lambda}$ . In Section 5, more precisely in Proposition 5.10, we give a criterion that allows to verify the stability of  $\mathscr{T}_X(-\log D)$ . For all divisors  $D = \sum_{\rho \in \Delta} D_\rho$  with  $1 \leq \operatorname{card}(\Delta) \leq 2$ , we give in tables 1, 2, 3 and 4 the values of  $\nu$  for which  $\mathscr{T}_X(-\log D)$  is (semi)stable with respect to L. We give here some results.

If  $a_1 = \ldots = a_r = 0$ , the variety X given in (1) is  $\mathbb{P}^s \times \mathbb{P}^r$ . Let  $\mathscr{D}_1 = \{D_{v_i} : 0 \le i \le r\} \cup \{D_{w_j} : 0 \le j \le s\} \cup \{D_{v_i} + D_{w_j} : 0 \le i \le r, 0 \le j \le s\}$  and

$$\mathscr{D}_2 = \{ D_{v_i} + D_{v_j} : 0 \le i < j \le r \} \cup \{ D_{w_i} + D_{w_j} : 0 \le i < j \le s \} .$$

We have :

**Theorem 1.4.** If  $D \in \mathscr{D}_1 \cup \mathscr{D}_2$ , then  $\operatorname{Stab}(\mathscr{T}_X(-\log D)) = \varnothing$ . But :

- (1) if  $D \in \mathscr{D}_2$ ,  $\operatorname{sStab}(\mathscr{T}_X(-\log D)) = \varnothing$ ;
- (2) if  $D \in \mathscr{D}_1$ ,  $\mathscr{T}_X(-\log D)$  is poly-stable with respect to L if and only if L is a power of the polarization corresponding to  $-(K_X + D)$ .

We now consider the general case, that is X given by (1) with  $a_r \ge 1$ .

**Theorem 1.5** (Stability of  $\mathscr{T}_X(-\log D_{v_r})$ ). We have  $\operatorname{Stab}(\mathscr{T}_X(-\log D_{v_r})) \neq \varnothing$ if and only if  $\operatorname{SStab}(\mathscr{T}_X(-\log D_{v_r})) \neq \varnothing$  if and only if  $a_r = 1$  and  $a_{r-1} = 0$ . If  $a_r = 1$  and  $a_{r-1} = 0$ , then the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D_{v_r})$  is stable (resp. semi-stable) with respect to  $\pi^*\mathscr{O}_{\mathbb{P}^s}(\mu) \otimes \mathscr{O}_X(\lambda)$  if and only if  $0 < \frac{\mu}{\lambda} < \nu_0$ (resp.  $0 < \frac{\mu}{\lambda} \leq \nu_0$ ) where  $\nu_0$  is the unique positive root of

$$\mathsf{P}_{0}(x) = \sum_{k=0}^{s-1} \binom{s+r-1}{k} x^{k} - s \binom{s+r-1}{s} x^{s} \; .$$

If  $a_r \geq 1$ , we see that this theorem is similar to [HNS19, Theorem 1.4]. If we fix r and s, there is only one smooth toric variety with Picard rank two such that  $\operatorname{Stab}(\mathscr{T}_X(-\log D_{v_r})) \neq \emptyset$  or  $\operatorname{Stab}(\mathscr{T}_X) \neq \emptyset$ ; this variety is given by (1) with  $a_1 = \ldots = a_{r-1} = 0$  and  $a_r = 1$ . If  $a_r \geq 1$  and  $(a_1, \ldots, a_r) \neq$  $(0, \ldots, 0, 1)$ , we have  $\operatorname{sStab}(\mathscr{T}_X) = \emptyset$  by [HNS19, Theorem 1.4]. In the case of log pairs, with some conditions on the  $a_i$ , there exists another divisor D such that  $\operatorname{Stab}(\mathscr{T}_X(-\log D)) \neq \emptyset$ .

**Theorem 1.6** (Stability of  $\mathscr{T}_X(-\log D_{v_0})$ ). We assume that  $a_r \geq 1$ .

- (1) If r = 1, then  $\operatorname{Stab}(\mathscr{T}_X(-\log D_{v_0})) \neq \varnothing$ .
- (2) If  $r \ge 2$ , then  $\operatorname{Stab}(\mathscr{T}_X(-\log D_{v_0})) \ne \varnothing$  if and only if  $a_1 = \ldots = a_r$ and  $a_r < \frac{s+1}{r-1}$ .

If  $D \notin \{D_{v_0}, D_{v_r}\}$ , then  $\mathscr{T}_X(-\log D)$  is not stable with respect to any polarizations.

Acknowledgments. I would like to thank my advisor Carl TIPLER for our discussions on this subject and also Henri GUENANCIA for some references.

#### A. NAPAME

2. Toric varieties and equivariant logarithmic tangent sheaves

2.1. Toric varieties. We give here some notions about toric varieties. We refer to [CLS11, Chapter 2, 3] and [CT20, Section 2.1].

**Definition 2.1** ([CLS11, Definition 3.1.1]). A toric variety is an irreducible variety X containing a torus  $T \simeq (\mathbb{C}^*)^n$  as a Zariski open subset such that the action of T on itself extends to an algebraic action of T on X.

Notation 2.2. Let T be a torus. We denote by M the lattice of characters of T and N the lattice of one-parameter subgroups of T; we have  $T \cong N \otimes_{\mathbb{Z}} \mathbb{C}^*$  (it is a group isomorphism). We set  $N_{\mathbb{K}} = N \otimes_{\mathbb{Z}} \mathbb{K}$  and  $M_{\mathbb{K}} = M \otimes_{\mathbb{Z}} \mathbb{K}$  for  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . For  $m \in M$ , we associate the character  $\chi^m : T \longrightarrow \mathbb{C}^*$  and for  $u \in N$ , we associate the oneparameter subgroup  $\lambda^u : \mathbb{C}^* \longrightarrow T$ . There is a natural pairing  $\langle \cdot, \cdot \rangle : M \times N \longrightarrow \mathbb{Z}$ between M and N.

**Definition 2.3** ([CLS11, Definition 3.1.2]). A fan  $\Sigma$  in  $N_{\mathbb{R}}$  is a finite collection of comes  $\sigma \subseteq N_{\mathbb{R}}$  such that :

- (1) Every  $\sigma \in \Sigma$  is a strongly convex rational polyhedral cone.
- (2) For all  $\sigma \in \Sigma$ , each face of  $\sigma$  is also in  $\Sigma$ .
- (3) For all  $\sigma_1, \sigma_2 \in \Sigma$ , the intersection  $\sigma_1 \cap \sigma_2$  is a face of each.

We will write  $\tau \preceq \sigma$  if  $\tau$  is a face of  $\sigma$ . If  $\Sigma$  is fan, the support of  $\Sigma$  is  $|\Sigma| = \bigcup_{\sigma \in \Sigma} \sigma$ .

We say that  $\Sigma$  is *complete*, if  $|\Sigma| = N_{\mathbb{R}}$ . We denote by  $\Sigma(r)$  the set of r-dimensional cones of  $\Sigma$ . We call  $\Sigma(1)$  the set of rays of  $\Sigma$ . If  $\rho \in \Sigma(1)$ , we denote by  $u_{\rho} \in N$  its minimal generator.

Let  $X_{\Sigma}$  be a toric variety associated to a fan  $\Sigma$ . The variety  $X_{\Sigma}$  is obtained by gluing affine charts  $(U_{\sigma})_{\sigma \in \Sigma}$ , with  $U_{\sigma} = \operatorname{Spec}(\mathbb{C}[S_{\sigma}])$  and  $\mathbb{C}[S_{\sigma}]$  is the semi-group algebra of

$$S_{\sigma} = \sigma^{\vee} \cap M = \{ m \in M : \langle m, u \rangle \ge 0 \text{ for all } u \in \sigma \} .$$

By [CLS11, Theorem 3.1.5],  $X_{\Sigma}$  is a normal separated toric variety.

**Theorem 2.4** ([CLS11, Corollary 3.1.8]). Let X be a normal separated toric variety with torus T. Then, there exists a fan  $\Sigma$  in  $N_{\mathbb{R}}$  such that X is isomorphic to  $X_{\Sigma}$ .

From now on, a normal toric variety X will be defined by a fan  $\Sigma$ .

**Definition 2.5** ([CLS11, Proposition 3.3.9]). A normal toric variety X has a *torus* factor if and only if the  $u_{\rho}, \rho \in \Sigma(1)$  do not span  $N_{\mathbb{R}}$ .

A cone  $\sigma \in \Sigma$  gives the torus orbit  $O(\sigma) \cong \operatorname{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}^*)$  where

$$\sigma^{\perp} \cap M = \{ m \in M : \langle m, u \rangle = 0 \text{ for all } u \in \sigma \} .$$

We have :

Theorem 2.6 (Orbit-Cone Correspondence, [CLS11, Theorem 3.2.6]).

(1) There is a bijective correspondence

$$\begin{array}{rcl} \{Cone \ \sigma \ in \ \Sigma\} & \longleftrightarrow & \{T - orbits \ in \ X\} \\ \sigma & \longleftrightarrow & O(\sigma) \end{array}$$

and  $\dim O(\sigma) = \dim N_{\mathbb{R}} - \dim \sigma$ .

- (2) The affine open subset  $U_{\sigma}$  is the union of orbits  $U_{\sigma} = \bigcup_{\tau \preceq \sigma} O(\tau)$ .
- (3)  $\tau \preceq \sigma$  if and only if  $O(\sigma) \subset \overline{O(\tau)}$ , and  $\overline{O(\tau)} = \bigcup \overset{\tau \supseteq \sigma}{O(\sigma)}$  where  $\overline{O(\tau)}$ denotes the closure in both the classical and Zariski topologies.

For  $\rho \in \Sigma(1)$ , we define the *T*-invariant Weil divisor  $D_{\rho}$  as the closure in the Zariski topology of  $O(\rho)$ .

**Lemma 2.7** ([CLS11, Section 4.1]). A Weil divisor D on X is invariant under the action of T if and only if

$$D = \sum_{\rho \in \Sigma(1)} a_{\rho} D_{\rho} \quad with \quad a_{\rho} \in \mathbb{Z} \;.$$

**Order Relation.** Let  $\Sigma$  be a fan in  $N_{\mathbb{R}}$  and  $\sigma \in \Sigma$ . We define an order relation  $\preceq_{\sigma}$  on M by setting  $m \preceq_{\sigma} m'$  if and only if  $m' - m \in S_{\sigma}$ . We write  $m \prec_{\sigma} m'$  if we have  $m \preceq_{\sigma} m'$  but not  $m' \preceq_{\sigma} m$ .

**Group action.** Let G be an algebraic group acting on the affine toric variety  $Y = \operatorname{Spec}(R)$ . For any  $g \in G$ , let  $\phi_g : Y \longrightarrow Y$  the map defined by  $\phi_g(x) = g \cdot x$ . We define an action of G on R by setting : for  $g \in G$  and  $\varphi \in R$ ,

$$g \cdot \varphi = \left(\phi_{g^{-1}}\right)^* \varphi$$

that is, for any  $y \in Y$ ,  $(g \cdot \varphi)(y) = \varphi(g^{-1} \cdot y)$ .

2.2. Equivariant sheaves. We refer to [Per03, Section 2.2.2] for properties about equivariant coherent sheaf. Let T be the torus of X.

Let  $\sigma: T \times X \longrightarrow X$  be an action of algebraic group T on X,  $\mu: T \times T \longrightarrow T$  the group multiplication,  $p_2: T \times X \longrightarrow X$  the projection onto the second factor and  $p_{23}: T \times T \times X \longrightarrow T \times X$  the projection onto the second and the third factor.

We call a sheaf  $\mathscr{E}$  on X equivariant or T-linearized if there exists an isomorphism  $\Phi: \sigma^* \mathscr{E} \xrightarrow{\cong} p_2^* \mathscr{E}$  such that

(2) 
$$(\mu \times \mathrm{Id}_X)^* \Phi = p_{23}^* \Phi \circ (\mathrm{Id}_T \times \sigma)^* \Phi$$

For  $t \in T$ , let  $\alpha_t : X \longrightarrow T \times X$  and  $\phi_t : X \longrightarrow X$  be the morphisms defined by

$$\alpha_t(x) = (t, x)$$
 and  $\phi_t(x) = \sigma(t, x)$ .

We get an isomorphism  $\Phi_t := \alpha_t^* \Phi : \phi_t^* \mathscr{E} \xrightarrow{\cong} \mathscr{E}$ . For any  $t, t' \in T$ , the cocyle condition (2) factors as follows :

(3) 
$$(\phi_{t'\cdot t})^* \mathscr{E} \xrightarrow{\Phi_{t'\cdot t}} \mathscr{E}$$
$$(\phi_{t'\cdot t})^* \mathscr{E} \xrightarrow{\phi_{t'} \Phi_{t'}} \mathscr{E}$$

2.3. Logarithmic tangent sheaves. We recall here the definition of the logarithmic tangent sheaf of a log pair (X, D) where X is a normal projective variety and D a reduced Weil divisor on X. We refer to [Gue16, Section 3.1].

**Definition 2.8.** We say that a pair (X, D) is log-smooth if X is smooth and D is reduced snc (simple normal crossing) divisor.

We denote by  $(X, D)_{reg}$  the snc locus of the pair (X, D), that is, the locus of points  $x \in X$  where (X, D) is log-smooth in a neighborhood of x.

If (X, D) is log-smooth, we define the logarithmic tangent bundle  $T_X(-\log D)$  as the dual of the bundle of logarithmic differential form  $\Omega^1_X(\log D)$  where  $\Omega^1_X(\log D)$ is defined in [Ita76, §1]. By [Kaw78, Definition 4] and [Sai75, §1], we can see the space of sections of  $T_X(-\log D)$  as the set of vector fields on X which vanish along D. If D is locally given by  $(z_1 \cdots z_k = 0)$ , then  $T_X(-\log D)$  as a sheaf is the locally free  $\mathscr{O}_X$ -module generated by

$$z_1 \frac{\partial}{\partial z_1}, \ldots, z_k \frac{\partial}{\partial z_k}, \frac{\partial}{\partial z_{k+1}}, \ldots, \frac{\partial}{\partial z_n}$$
.

**Definition 2.9** ([Gue16, Definition 3.4]). Let (X, D) be a log pair and  $X_0 = (X, D)_{\text{reg.}}$ . The logarithmic tangent sheaf of (X, D), denoted by  $\mathscr{T}_X(-\log D)$ , is defined as  $j_*T_{X_0}(-\log D_{|X_0})$  where  $j: X_0 \longrightarrow X$  is the open immersion.

The sheaf  $\mathscr{T}_X(-\log D)$  (as well as its dual) is coherent ; by [Har80, Proposition 1.6], this sheaf is reflexive.

We now consider the case where X is a toric variety. We will give a condition on D such that  $\mathscr{T}_X(-\log D)$  is equivariant. Let  $\Sigma$  be the fan of X and  $X_0$  the toric variety corresponding to the fan  $\Sigma^1 = \Sigma(0) \cup \Sigma(1)$ . We denote by  $j: X_0 \longrightarrow X$  the open immersion.

**Proposition 2.10.** Let D be a reduced Weil divisor on X. The sheaf  $\Omega^1_X(\log D)$  is equivariant if and only if D is an invariant divisor under the torus action.

Proof. We assume that D is an invariant divisor under the torus action. Let  $D_0$  be the restriction of D on  $X_0$ . For  $t \in T$ , let  $\phi_t : X \longrightarrow X$  the map defined by  $\phi_t(x) = \sigma(t, x)$  and  $\Phi_t$  the map defined by  $\Phi_t = (d\phi_t)^{-1}$  where  $d\phi_t$  is the differential of  $\phi_t$ . If  $\mathscr{E} = TX_0$ , we have an isomorphism  $\Phi_t : \phi_t^* \mathscr{E} \longrightarrow \mathscr{E}$  and the diagram (3) is verified. Now if we replace  $\mathscr{E}$  by  $T_{X_0}(-\log D_0)$ , the diagram (3) remains true ; so  $T_{X_0}(-\log D_0)$  is equivariant. Hence  $\Omega_{X_0}^1(\log D_0)$  is equivariant. Like

(4) 
$$\Omega^1_X(\log D) \cong j_*\Omega^1_{X_0}(\log D_0) ,$$

we deduce that  $\Omega^1_X(\log D)$  is equivariant.

exact sequences

We now assume that  $\Omega^1_X(\log D)$  is equivariant. We write  $D = \sum_{j=1}^s D_j$ .

*First case.* We assume that X is smooth. By [EV92, Section 2.3] we have an exact sequence

$$0 \longrightarrow \Omega^1_X \longrightarrow \Omega^1_X \left( \log D \right) \longrightarrow \bigoplus_{j=1}^s \mathscr{O}_{D_j} \longrightarrow 0$$

where  $\mathscr{O}_{D_j}$  is viewing as a sheaf on X via extension by zero. The first part of the proof is to show that : for any  $t \in T$ ,  $t \cdot Z = Z$  where  $Z = X \setminus D$ . Let  $x \in Z$  and assume that there is  $t \in T$  such that  $y = \sigma(t, x) \in D$ . We have two

$$0 \longrightarrow \Omega^{1}_{X, x} \longrightarrow \Omega^{1}_{X} (\log D)_{x} \longrightarrow \bigoplus_{j=1}^{s} \mathscr{O}_{D_{j}, x} \longrightarrow 0$$
$$0 \longrightarrow \Omega^{1}_{X, y} \longrightarrow \Omega^{1}_{X} (\log D)_{y} \longrightarrow \bigoplus_{j=1}^{s} \mathscr{O}_{D_{j}, y} \longrightarrow 0$$

Like  $\Omega^1_X$  and  $\Omega^1_X(\log D)$  are equivariant, we have an isomorphism

$$\bigoplus_{j=1}^{s} \mathscr{O}_{D_{j}, x} \cong \bigoplus_{j=1}^{s} \mathscr{O}_{D_{j}, y} ;$$

this is absurd. Therefore, for any  $t \in T$ , we have  $t \cdot Z \subset Z$ , that is  $t \cdot Z = Z$ . As  $\Omega^1_X(\log D)$  is equivariant, by using the fact that  $D = X \setminus Z$ , for any  $t \in T$ , we have  $t \cdot D = D$ ; thus, D is a T-invariant divisor.

Second case. We assume that X is a normal variety. By (4), as  $\Omega^1_X(\log D)$  is equivariant, we also have the same property for  $\Omega^1_{X_0}(\log D_0)$ . By the first case,  $D_0$  is an invariant divisor under the action of T on  $X_0$ . Like  $\operatorname{codim}(X \setminus X_0) \ge 2$ , we deduce that D is the Zariski closure of  $D_0$  on X. Thus, D is an invariant divisor under the action of T on X.

#### 3. Multifiltrations of logarithmic tangent sheaf

3.1. Family of multifiltrations. We give here the family of multifiltrations of equivariant logarithmic tangent sheaf on toric varieties. Let X be a toric variety of dimension n with fan  $\Sigma$ . We first give some results.

**Proposition 3.1** ([DDK20, Corollary 2.2.17]). The family  $(\mathcal{T}, {\mathcal{T}^{\rho}(i)}_{\rho \in \Sigma(1), i \in \mathbb{Z}})$  of multifiltrations associated to the tangent sheaf  $\mathcal{T}_X$  are given by

$$\mathscr{T}^{\rho}(i) = \begin{cases} 0 & \text{if } i \leq -2\\ \operatorname{Span}(u_{\rho}) & \text{if } i = -1\\ N \otimes_{\mathbb{Z}} \mathbb{C} & \text{if } i \geq 0 \end{cases}$$

Remark 3.2. If X is smooth, the tangent sheaf  $\mathscr{T}_X$  if the tangent bundle TX.

**Lemma 3.3.** For every ray  $\rho \in \Sigma(1)$ , we have  $(U_{\rho} \cap D_{\rho}) \cap T = \emptyset$ .

*Proof.* Let  $\rho \in \Sigma(1)$ . We assume that  $\rho = \text{Cone}(u_1)$  where  $u_1 \in N$ . Let  $(u_1, \ldots, u_n)$  be a basis of N and  $(e_1, \ldots, e_n)$  its dual basis. We set  $x_j = \chi^{e_j}$ , we have

 $U_{\rho} = \operatorname{Spec}(\mathbb{C}[x_1, x_2^{\pm 1}, \dots, x_n^{\pm 1}]) \cong \mathbb{C} \times (\mathbb{C}^*)^{n-1}$ 

and

$$T = \operatorname{Spec}(\mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]) \cong (\mathbb{C}^*)^n .$$

As on  $U_{\rho}$  the divisor  $D_{\rho}$  is defined by  $x_1 = 0$ , we deduce that  $(U_{\rho} \cap D_{\rho}) \cap T = \emptyset$ .  $\Box$  **Lemma 3.4.** Let  $\rho_1$  and  $\rho_2$  be two distinct rays of  $\Sigma$ . If  $i \neq j$ , then  $U_{\rho_i} \cap D_{\rho_j} = \emptyset$ . *Proof.* By the orbit-cone correspondence (Theorem 2.6), we have  $U_{\rho_i} = O(\rho_i) \cup T$ and  $O(\rho_i) \cap D_{\rho_j} = \emptyset$  if  $i \neq j$ . As

$$U_{\rho_i} \cap D_{\rho_j} = \left( O(\rho_i) \cap D_{\rho_j} \right) \cup \left( T \cap D_{\rho_j} \right)$$

by Lemma 3.3, we conclude that  $U_{\rho_i} \cap D_{\rho_i} = \emptyset$ .

Let  $\Delta \subset \Sigma(1)$  and D be the T-invariant divisor of X defined by

$$D = \sum_{\rho \in \Delta} D_{\rho} \; .$$

For  $\rho \in \Sigma(1)$ , we set  $E^{\rho} = \Gamma(U_{\rho}, \mathscr{T}_X(-\log D))$ .

**Theorem 3.5.** The family of multifiltrations  $(E, \{E^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  corresponding to the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$  are given by

$$E^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -1 \\ N_{\mathbb{C}} & \text{if } j \geq 0 \end{cases} \quad \text{if } \rho \in \Delta$$

and by

$$E^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -2\\ \text{Span}(u_{\rho}) & \text{if } j = -1\\ N_{\mathbb{C}} & \text{if } j \geq 0 \end{cases} \quad \text{if } \rho \notin \Delta .$$

*Proof.* According to Lemma 3.4, if  $\rho \in \Delta$ , we have  $U_{\rho} \cap D = D_{\rho}$  and for  $\rho \notin \Delta$ ,  $U_{\rho} \cap D = \emptyset$ . We can reduce the problem to the case where  $\Delta = \{\rho_1\}$  and  $D = D_{\rho_1}$ . For the rest of the proof, we assume that  $\Delta = \{\rho_1\}$ .

First case : We assume that  $\rho = \rho_1$ . Let  $(u_1, \ldots, u_n)$  be a basis of N such that  $u_1 = u_{\rho}$ . We denote by  $(e_1, \ldots, e_n)$  the dual basis of  $(u_1, \ldots, u_n)$  and  $x_i = \chi^{e_i}$ . We have  $\mathbb{C}[S_{\rho}] = \mathbb{C}[x_1, x_2^{\pm 1}, \ldots, x_n^{\pm 1}]$  and  $U_{\rho} = \operatorname{Spec}(\mathbb{C}[S_{\rho}])$ . Like on  $U_{\rho}$  the divisor D is defined by the equation  $x_1 = 0$ , we have

$$E^{\rho} = \left( \mathbb{C}[S_{\rho}] \cdot x_1 \frac{\partial}{\partial x_1} \right) \oplus \left( \bigoplus_{i=2}^n \mathbb{C}[S_{\rho}] \cdot \frac{\partial}{\partial x_i} \right)$$

that is

$$E^{\rho} = \bigoplus_{i=2}^{n} \left( \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^{m} \frac{\partial}{\partial x_{i}} \right) \oplus \left( \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^{m+e_{1}} \frac{\partial}{\partial x_{1}} \right) \,.$$

We set

$$L_1^{\rho} = \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^{m+e_1} \frac{\partial}{\partial x_1}$$

and for  $i \in \{2, \ldots, n\}$ , we set

$$L_i^{\rho} = \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^m \, \frac{\partial}{\partial x_i}$$

For  $t \in T$  and  $m \in M$ , we have  $t \cdot \chi^m = \chi^{-m}(t)\chi^m$ , so  $t \cdot dx_i = \chi^{-e_i}(t) dx_i$ . Thus, we have  $t \cdot \frac{\partial}{\partial x_i} = \chi^{e_i}(t) \frac{\partial}{\partial x_i}$ . For  $i \in \{1, \ldots, n\}$ , we write

$$L_i^{\rho} = \bigoplus_{m \in M} (L_i^{\rho})_m \quad \text{where} \quad (L_i^{\rho})_m = \{ f \in L_i^{\rho} : t \cdot f = \chi^{-m}(t) f \} .$$

We have

$$(L_1^{\rho})_m = \begin{cases} \mathbb{C} \cdot \chi^{m+e_1} \frac{\partial}{\partial x_1} & \text{if } 0 \preceq_{\rho} m \\ 0 & \text{otherwise} \end{cases}$$

and for  $i \in \{2, \ldots, n\}$ , we have

$$\left(L_{i}^{\rho}\right)_{m} = \begin{cases} \mathbb{C} \cdot \chi^{m+e_{i}} \frac{\partial}{\partial x_{i}} & \text{if } -e_{i} \preceq_{\rho} m \\ 0 & \text{otherwise} \end{cases}$$

For  $i \in \{1, \ldots, n\}$ , we set  $\mathbb{L}_i^{\rho} = \mathbb{C} \cdot \frac{\partial}{\partial x_i}$ . If  $i \in \{2, \ldots, n\}$ , we have  $(L_i^{\rho})_{-e_i} = \mathbb{L}_i^{\rho}$ . For all  $m \in M$ :

- if  $i \geq 2$  and  $-e_i \preceq_{\rho} m$ , we identify  $(L_i^{\rho})_m$  with  $\mathbb{L}_i^{\rho}$  through the multiplication by the character  $\chi^{-m-e_i}$ .
- If i = 1 and  $0 \leq_{\rho} m$ , we identify  $(L_i^{\rho})_m$  with  $\mathbb{L}_1^{\rho}$  through the multiplication by  $\chi^{-m-e_1}$ .

For  $m \in M$ , we set  $j = \langle m, u_1 \rangle$ .

- If j ≤ −1, for all i ∈ {1,..., n}, we have (L<sup>ρ</sup><sub>i</sub>)<sub>m</sub> = 0.
  If j ≥ 0, for all i ∈ {1,..., n}, (L<sup>ρ</sup><sub>i</sub>)<sub>m</sub> ≃ L<sup>ρ</sup><sub>i</sub>.

The torus T is a Lie group. The tangent space of T at the neutral element is isomorphic to  $N_{\mathbb{C}}$ . Like the tangent space of T at the neutral element is generated by  $\left(\frac{\partial}{\partial x_i}\right)_{1 \le i \le n}$ , for all  $i \in \{1, \ldots, n\}$ , we can identify  $\frac{\partial}{\partial x_i}$  with  $u_i$ . Like

$$E^{\rho} = \bigoplus_{m \in M} E^{\rho}(\langle m, u_1 \rangle) \quad \text{where} \quad E^{\rho}(\langle m, u_1 \rangle) = E^{\rho}_m = \bigoplus_{i=1}^n (L^{\rho}_i)_m + E^{\rho}_m = E$$

we get  $E^{\rho}(j) \cong \begin{cases} 0 & \text{if } j \leq -1 \\ N_{\mathbb{C}} & \text{if } j \geq 0 \end{cases}$ 

Second case : We assume that  $\rho \in \Sigma(1) \setminus \Delta$ . We start as in the first case. Like  $U_{\rho} \cap D = \emptyset$ , we have

$$E^{\rho} = \bigoplus_{i=1}^{n} \mathbb{C}[S_{\rho}] \cdot \frac{\partial}{\partial x_{i}} = \bigoplus_{i=1}^{n} \left( \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^{m} \frac{\partial}{\partial x_{i}} \right)$$

For all  $i \in \{1, \ldots, n\}$ , we set

$$L_i^{\rho} = \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^m \, \frac{\partial}{\partial x_i}$$

We have

$$L_i^{\rho} = \bigoplus_{m \in M} \left( L_i^{\rho} \right)_m \quad \text{where} \quad \left( L_i^{\rho} \right)_m = \begin{cases} \mathbb{C} \cdot \chi^{m+e_i} \frac{\partial}{\partial x_i} & \text{if } -e_j \preceq_{\rho} m \\ 0 & \text{oherwise} \end{cases}$$

As in the first case, with a small modification, for all  $i \in \{1, \ldots, n\}$ , we identify  $(L_i^{\rho})_m$  with  $\mathbb{L}_i^{\rho}$  if  $m \in M$  satisfies  $-e_i \preceq_{\rho} m$ . By setting  $j = \langle m, u_{\rho} \rangle$ , we get

$$E^{\rho}(j) \cong \begin{cases} 0 & \text{if } j \leq -2\\ \operatorname{Span}(u_{\rho}) & \text{if } j = -1\\ N \otimes_{\mathbb{Z}} \mathbb{C} & \text{if } j \geq 0 \end{cases}$$

Hence we get the theorem.

The sheaf of regular section of the trivial vector bundle  $X \times \mathbb{C} \longrightarrow X$  of rank 1 is  $\mathscr{O}_X$ . For  $\rho \in \Sigma(1)$ , we set  $F^{\rho} = \mathscr{O}_X(U_{\rho})$ . Like  $F^{\rho} = \mathbb{C}[S_{\rho}]$ , we have

$$F^{\rho} = \bigoplus_{m \in S_{\rho}} \mathbb{C} \cdot \chi^{m} = \bigoplus_{m \in M} F^{\rho}_{m} \quad \text{where} \quad F^{\rho}_{m} = \{f \in F^{\rho} : t \cdot f = \chi^{-m}(t) f\} \;.$$

Thus  $F_m^{\rho} = 0$  if  $m \notin S_{\rho}$  and  $F_m^{\rho} = \mathbb{C} \cdot \chi^m$  if  $m \in S_{\rho}$  (i.e  $0 \leq_{\rho} m$ ). For all  $m \in S_{\rho}$ , we identify  $F_m^{\rho}$  with  $F_0^{\rho} = \mathbb{C}$ . By setting  $j = \langle m, u_{\rho} \rangle$ , we get

$$F^{\rho}(j) \cong \begin{cases} 0 & \text{if } j \le -1 \\ \mathbb{C} & \text{if } j \ge 0 \end{cases}$$

**Corollary 3.6.** Let  $\Delta \subset \Sigma(1)$  and  $D = \sum_{\rho \in \Delta} D_{\rho}$ .

- 1. If  $\Delta = \emptyset$ , then  $\mathscr{T}_X(-\log D)$  is the tangent sheaf  $\mathscr{T}_X$ .
- 2. If  $\Delta = \Sigma(1)$ , then  $\mathscr{T}_X(-\log D)$  is isomorphic to the trivial vector bundle of rank n.

*Proof.* If  $\Delta = \emptyset$ , the family of multifiltrations of  $\mathscr{T}_X(-\log D)$  is identical to the family of multifiltrations given in Proposition 3.1. If  $\Delta = \Sigma(1)$ , for all  $\rho \in \Sigma(1)$ , we have

$$E^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -1 \\ N \otimes_{\mathbb{Z}} \mathbb{C} & \text{if } j \geq 0 \end{cases}$$

Hence,  $\mathscr{T}_X(-\log D)$  is isomorphic to the trivial vector bundle of rank n.

From now on, we will assume that  $\emptyset \subsetneq \Delta \subsetneq \Sigma(1)$  and D the T-invariant Weil divisor defined by

$$D = \sum_{\rho \in \Delta} D_{\rho} \; .$$

**Notation 3.7.** Let G be a sub-vector space of  $N_{\mathbb{C}}$ . We denote by  $\mathscr{E}_G$  the sub-sheaf of  $\mathscr{E} = \mathscr{T}_X(-\log D)$  defined by the family of multifiltrations  $(E_G, \{G^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  where  $E_G = G$  and  $G^{\rho}(j) = E^{\rho}(j) \cap G$ . If  $\rho \in \Delta$  or  $u_{\rho} \notin G$ , then

$$G^{
ho}(j) = \left\{ egin{array}{cc} 0 & ext{if } j \leq -1 \ G & ext{if } j \geq 0 \end{array} 
ight.$$

If  $\rho \notin \Delta$  and  $u_{\rho} \in G$ , then

$$G^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -2\\ \text{Span}(u_{\rho}) & \text{if } j = -1\\ G & \text{if } j \geq 0 \end{cases}$$

#### A. NAPAME

3.2. Decomposition of equivariant logarithmic tangent sheaf. In this part, we give some conditions on  $\Sigma$  and  $\Delta$  which ensure that the logarithmic tangent sheaf is decomposable. We first recall the family of multifiltration of a direct sum of equivariant reflexive sheaves.

**Proposition 3.8** ([DDK20, Remark 2.2.15]). Let  $\mathscr{F}$  and  $\mathscr{G}$  be two equivariant reflexive sheaves with  $(F, \{F^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  and  $(G, \{G^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  for family of filtrations. The family of multifiltrations of  $\mathscr{F} \oplus \mathscr{G}$  is given by

(5)  $(F \oplus G, \{(F \oplus G)^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  where  $(F \oplus G)^{\rho}(j) = F^{\rho}(j) \oplus G^{\rho}(j)$ .

We assume that X is toric variety without torus factor. We denote by p the rank of the class group Cl(X) of X. By [CLS11, Theorem 4.1.3], we have an exact sequence

(6) 
$$0 \longrightarrow M \longrightarrow \bigoplus_{\rho \in \Sigma(1)} \mathbb{Z} \cdot D_{\rho} \longrightarrow \operatorname{Cl}(X) \longrightarrow 0$$

and  $\operatorname{card}(\Sigma(1)) = n + p$ .

**Theorem 3.9.** We assume that  $\operatorname{card}(\Delta) = p$ . We set  $\Sigma(1) \setminus \Delta = \{\rho_1, \ldots, \rho_n\}$  where  $\rho_k = \operatorname{Cone}(u_k)$  and  $u_k \in N$ . If  $N_{\mathbb{R}} = \operatorname{Span}(u_1, \ldots, u_n)$ , then  $\mathscr{E} = \mathscr{T}_X(-\log D)$  is decomposable and

$$\mathscr{E} = \bigoplus_{k=1}^n \mathscr{E}_{F_k}$$

where  $\mathscr{E}_{F_k}$  is the sub-sheaf of  $\mathscr{E}$  corresponding to the vector space  $F_k = \text{Span}(u_k)$ .

*Proof.* For all  $k \in \{1, ..., n\}$ , the family of multifiltration  $(F_k, \{F_k^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  of  $\mathscr{E}_{F_k}$  are given by

$$F_k^{\rho}(j) = \begin{cases} 0 & \text{if } j \le -1 \\ F_k & \text{if } j \ge 0 \end{cases} \quad \text{if } \rho \neq \text{Cone}(u_k)$$

and

$$F_k^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -2\\ \text{Span}(u_{\rho}) & \text{if } j = -1\\ F_k & \text{if } j \geq 0 \end{cases} \text{ for } \rho = \text{Cone}(u_k) \ .$$

For all  $\rho \in \Sigma(1)$  and  $j \in \mathbb{Z}$ , we have

$$\bigoplus_{k=1}^{n} F_{k}^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -1 \\ N_{\mathbb{C}} & \text{if } j \geq 0 \end{cases} \quad \text{if } \rho \in \Delta$$

and

$$\bigoplus_{k=1}^{n} F_{k}^{\rho}(j) = \begin{cases} 0 & \text{if } j \leq -2 \\ \text{Span}(u_{\rho}) & \text{if } j = -1 \\ N_{\mathbb{C}} & \text{if } j \geq 0 \end{cases} \quad \text{if } \rho \notin \Delta \ .$$

Hence, by (5) and Theorem 3.5 we have  $\mathscr{E} = \bigoplus_{k=1}^{n} \mathscr{E}_{F_k}$ .

**Proposition 3.10.** We assume that  $\Delta$  satisfies  $1 + p \leq \operatorname{card}(\Delta) \leq n + p - 1$ . The sheaf  $\mathscr{E} = \mathscr{T}_X(-\log D)$  is decomposable and  $\mathscr{E} = \mathscr{E}_G \oplus \mathscr{E}_F$  where  $G = \operatorname{Span}(u_\rho : \rho \in \Sigma(1) \setminus \Delta)$  and F a sub-space of  $N_{\mathbb{C}}$  such that  $N_{\mathbb{C}} = G \oplus F$ .

*Proof.* It suffices to work with the family of multifiltration as in the proof of Theorem 3.9.

4.1. Some stability notions. We recall now the notions of stability that we will consider in this paper. For this part, we refer to [Gue16, Section 3.1] and [CT20, Section 4.1].

**Definition 4.1.** Let  $\mathscr{E}$  be a torsion-free coherent sheaf on X. The *degree* of  $\mathscr{E}$  with respect to an ample class  $L \in Amp(X)$  is the real number obtained by intersection:

$$\deg_L(\mathscr{E}) = c_1(\mathscr{E}) \cdot L^{n-1}$$

and its *slope* with respect to L is given by

$$\mu_L(\mathscr{E}) = \frac{\deg_L(\mathscr{E})}{\operatorname{rk}(\mathscr{E})}.$$

**Definition 4.2.** A torsion-free coherent sheaf  $\mathscr{E}$  is said to be  $\mu$ -semi-stable with respect to  $L \in \operatorname{Amp}(X)$  if for any proper coherent sub-sheaf of lower rank  $\mathscr{F}$  of  $\mathscr{E}$  with  $0 < \operatorname{rk} \mathscr{F} < \operatorname{rk} \mathscr{E}$ , one has

$$\mu_L(\mathscr{F}) \le \mu_L(\mathscr{E}) \; .$$

When strict inequality always holds, we say that  $\mathscr{E}$  is  $\mu$ -stable. Finally,  $\mathscr{E}$  is said to be  $\mu$ -polystable if it is the direct sum of  $\mu$ -stable sub-sheaves of the same slope.

**Lemma 4.3** ([CT20, Lemma 4.3]). Let  $\mathscr{E}$  be an equivariant reflexive sheaf with family of multifiltrations  $(E, \{E^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$ . We have

$$\mu_L(\mathscr{E}) = \frac{1}{\operatorname{rk}(\mathscr{E})} \sum_{\rho \in \Sigma(1)} \sum_{i \in \mathbb{Z}} i \, e^{\rho}(i) \, \deg_L(D_{\rho})$$

where  $e^{\rho}(i) = \dim E^{\rho}(i-1) - \dim E^{\rho}(i)$ .

Following [Koo11, Proposition 4.13] and [HNS19, Proposition 2.3], we have :

**Proposition 4.4** ([CT20, Proposition 4.2]). Let  $\mathscr{E}$  be a *T*-equivariant reflexive sheaf on *X* with family of multifiltrations  $(E, \{E^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$ . Then  $\mathscr{E}$  is  $\mu$ semi-stable (resp.  $\mu$ -stable) with respect to *L* if and only if for all proper vector sub-spaces  $W \subset E$ ,  $\mu_L(\mathscr{E}_W) \leq \mu_L(\mathscr{E})$  (resp.  $\mu_L(\mathscr{E}_W) < \mu_L(\mathscr{E})$ ), where  $\mathscr{E}_W$  is defined in notation 3.7.

4.2. **Polystability and semi-stability.** We give here a link between poly-stability and semi-stability.

**Proposition 4.5** ([Koo11, Claim 2 of Proposition 4.13]). A reflexive  $\mu$ -polystable sheaf on X is a  $\mu$ -semi-stable sheaf on X isomorphic to a (finite, nontrivial) direct sum of reflexive  $\mu$ -stable sheaves. Let  $\mathscr{E}$  be a  $\mu$ -semi-stable reflexive sheaf on X. Then  $\mathscr{E}$  contains a unique maximal reflexive  $\mu$ -polystable sub-sheaf of the same slope as  $\mathscr{E}$ .

**Corollary 4.6.** Let  $\mathcal{E}$  be an equivariant reflexive sheaf such that

$$\mathscr{E} = \bigoplus_{i=1}^r \mathscr{E}_i$$

with  $r \geq 2$ . Let  $L \in Amp(X)$  such that for all  $i \in \{1, \ldots, r\}$ ,  $\mathscr{E}_i$  is stable with respect to L. Then,  $\mathscr{E}$  is poly-stable with respect to L if and only if  $\mathscr{E}$  is semi-stable with respect to L.

A. NAPAME

*Proof.* If  $\mathscr{E}$  is poly-stable with respect to L, then  $\mathscr{E}$  is semi-stable with respect to L. We now assume that  $\mathscr{E}$  is semi-stable with respect to L; by using the identity

$$\operatorname{rk}(\mathscr{E})\,\mu_L(\mathscr{E}) = \sum_{i=1}^r \operatorname{rk}(\mathscr{E}_i)\,\mu_L(\mathscr{E}_i) \;,$$

we get  $\mu_L(\mathscr{E}_i) = \mu_L(\mathscr{E})$  for all  $i \in \{1, \ldots, r\}$ . Hence,  $\mathscr{E}$  is poly-stable.  $\Box$ 

4.3. An instability condition for the logarithmic tangent sheaves. Let  $\Delta \subset \Sigma(1)$  and

$$D = \sum_{\rho \in \Delta} D_{\rho}$$

a *T*-invariant Weil divisor on *X*. Let *G* be a sub-vector space of  $N_{\mathbb{C}}$  of dimension l with  $1 \leq l < n$ . Let  $(E, \{E^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  be the family of multifiltrations corresponding to  $\mathscr{E} = \mathscr{T}_X(-\log D)$  and  $(E_G, \{G^{\rho}(j)\}_{\rho \in \Sigma(1), j \in \mathbb{Z}})$  be the family of multifiltrations corresponding to  $\mathscr{E}_G$ . By Lemma 4.3, we have

(7) 
$$\mu_L(\mathscr{E}) = \frac{1}{n} \sum_{\rho \notin \Delta} \deg_L(D_\rho)$$

and

(8) 
$$\mu_L(\mathscr{E}_G) = \frac{1}{l} \sum_{\rho \notin \Delta \text{ and } u_\rho \in G} \deg_L(D_\rho) .$$

Therefore, we have (9)

$$\mu_L(\mathscr{E}) - \mu_L(\mathscr{E}_G) = \left(\frac{1}{n} - \frac{1}{l}\right) \left(\sum_{\rho \notin \Delta, \ u_\rho \in G} \deg_L(D_\rho)\right) + \frac{1}{n} \left(\sum_{\rho \notin \Delta, \ u_\rho \notin G} \deg_L(D_\rho)\right).$$

To study the stability of  $\mathscr{E}$  with respect to  $L \in \operatorname{Amp}(X)$ , it suffices to compare  $\mu_L(\mathscr{E}_G)$  with  $\mu_L(\mathscr{E}_G)$  where  $G \subset \operatorname{Span}(u_{\rho} : \rho \notin \Delta)$  and  $1 \leq \dim G \leq n-1$ .

**Theorem 4.7.** If  $1 \leq \operatorname{card}(\Sigma(1) \setminus \Delta) \leq n-1$ , then for any  $L \in \operatorname{Amp}(X)$ , the logarithmic tangent sheaf  $\mathscr{E} = \mathscr{T}_X(-\log D)$  is not semi-stable with respect to L.

*Proof.* We assume that  $\Sigma(1) \setminus \Delta = \{\rho_1, \ldots, \rho_k\}$  where  $1 \le k \le n-1$  and we denote by  $D_j$  the divisor corresponding to  $\rho_j = \text{Cone}(u_j)$ . Let  $G = \text{Span}(u_1, \ldots, u_k)$  and  $l = \dim G$ . If  $L \in \text{Amp}(X)$ , we have

$$\mu_L(\mathscr{E}) - \mu_L(\mathscr{E}_G) = \left(\frac{1}{n} - \frac{1}{l}\right) \sum_{j=1}^k \deg_L(D_j) < 0$$

because the numbers  $\deg_L(D_j)$  are positive and different to zero. Thus,  $\mathscr{E}$  is not semi-stable with respect to L.

We assume that X has no torus factor. We have seen that  $\operatorname{card}(\Sigma(1)) = n + \operatorname{rk} \operatorname{Cl}(X)$ , so the Theorem 4.7 becomes :

**Corollary 4.8.** We set  $p = \operatorname{rk} \operatorname{Cl}(X)$ . If  $1 + p \leq \operatorname{card}(\Delta) \leq n + p - 1$ , then for any  $L \in \operatorname{Amp}(X)$ , the logarithmic tangent sheaf  $\mathscr{T}_X(-\log D)$  is not semi-stable with respect to L.

*Proof.* If  $1 + p \leq \operatorname{card}(\Delta) \leq n + p - 1$ , by using

$$\operatorname{card}(\Sigma(1)) = n + p = \operatorname{card}(\Delta) + \operatorname{card}(\Sigma(1) \setminus \Delta) ,$$

we get  $1 \leq \operatorname{card}(\Sigma(1) \setminus \Delta) \leq n-1$ ; we can conclude with Theorem 4.7.

Remark 4.9. By Corollary 3.6, if  $\operatorname{card}(\Delta) = n + p$ ,  $\mathscr{T}_X(-\log D)$  is semi-stable.

From now on, when we study the stability or semi-stability of  $\mathscr{T}_X(-\log D)$ , we will consider only the case where  $1 \leq \operatorname{card}(\Delta) \leq p = \operatorname{rk} \operatorname{Cl}(X)$  and  $p \in \{1, 2\}$ .

4.4. **Stability on toric orbifolds of Picard rank one.** In this section, we will study the semi-stability of logarithmic tangent sheaf on toric orbifolds of Picard rank one.

Let  $q_0, q_1, \ldots, q_n \in \mathbb{N}^*$  such that  $gcd(q_0, \ldots, q_n) = 1$ . We set

$$N = \mathbb{Z}^{n+1} / \mathbb{Z}(q_0, \dots, q_n)$$

The dual lattice of N is  $M = \{(a_0, \ldots, a_n) \in \mathbb{Z}^{n+1} : a_0 q_0 + \ldots + a_n q_n = 0\}$ . Let  $\{u_i : 0 \le i \le n\}$  be the images in N of the standard basis vectors in  $\mathbb{Z}^{n+1}$ , so the relation

$$q_0 u_0 + q_1 u_1 + \ldots + q_n u_n = 0$$

holds in N. Let  $\Sigma$  be the fan in  $N_{\mathbb{R}}$  defined by

$$\Sigma = \{ \operatorname{Cone}(A) : A \subsetneq \{u_0, \dots, u_n\} \} .$$

We define X to be the toric variety of the fan  $\Sigma$ . Like  $\Sigma$  is simplicial, by [CLS11, Proposition 4.2.7], Pic(X) has finite index in Cl(X). By (6), we deduce that rk Pic(X) = 1. We denote by  $D_i$  the Weil divisor corresponding to the ray Cone $(u_i)$ .

In the Corollary 4.8, we see that if  $2 \leq \operatorname{card}(\Delta) \leq n$ , then  $\mathscr{T}_X(-\log D)$  is not semistable with respect to any  $L \in \operatorname{Amp}(X)$ . We assume that  $\operatorname{card}(\Delta) = 1$ . Let  $i \in \{0, \ldots, n\}$  and  $A_i = \{0, \ldots, n\} \setminus \{i\}$ . We set  $\mathscr{E} = \mathscr{T}_X(-\log D_i)$ .

**Lemma 4.10.** For all  $j \in A_i$ , the divisor  $q_iD_j$  is linearly equivalent to the divisor  $q_jD_i$ .

*Proof.* We fix  $j \in A_i$ . Let  $m = (a_0, \ldots, a_n) \in M$  defined by  $a_i = q_j$ ,  $a_j = -q_i$  and  $a_k = 0$  if  $k \in \{0, \ldots, n\} \setminus \{i, j\}$ . Like  $\operatorname{div}(\chi^m) = q_j D_i - q_i D_j$ , we deduce that  $q_i D_j$  is linearly equivalent to  $q_j D_i$ .

**Theorem 4.11.** Let  $L \in Amp(X)$ . The sheaf  $\mathscr{E}$  is poly-stable with respect to L if and only if there is  $q \in \mathbb{N}^*$  such that for all  $j \in A_i$ ,  $q_j = q$ .

*Proof.* The assumptions of Theorem 3.9 are verified. Hence,

$$\mathscr{E} = \bigoplus_{j \in A_i} \mathscr{E}_{F_j}$$

where  $F_j = \text{Span}(u_j)$ . By (8),  $\mu_L(\mathscr{E}_{F_j}) = \deg_L(D_j)$  for all  $j \in A_i$ . By Lemma 4.10, for all  $j \in A_i$ ,

$$\mu_L(\mathscr{E}_{F_j}) = \frac{q_j}{q_i} \deg_L(D_i) \; .$$

If  $\mathscr{E}$  is poly-stable with respect to L, there is  $r \in \mathbb{Q}$  such that for all  $j \in A_i$ ,  $q_j = r q_i$ . Hence, we have the existence of  $q \in \mathbb{N}^*$  such that for all  $j \in A_i$ ,  $q_j = q$ . For the converse, if for all  $j \in A_i$ , we have  $q_j = q$ , then  $\mathscr{E}$  is poly-stable.

According to Theorem 4.11 and Corollary 4.6, we have :

**Corollary 4.12.** For all  $i \in \{0, ..., n\}$ ,  $\operatorname{sStab}(\mathscr{T}_X(-\log D_i)) \neq \emptyset$  if and only if there exists  $q \in \mathbb{N}^*$  such that for all  $j \in A_i$ ,  $q_j = q$ . Moreover, if for all  $j \in A_i$ , we have  $q_j = q$ , then

$$\operatorname{sStab}(\mathscr{T}_X(-\log D_i)) = \operatorname{Amp}(X)$$
.

**Corollary 4.13.** For all  $i \in \{0, \ldots, n\}$ , we have  $\operatorname{Stab}(\mathscr{T}_X(-\log D_i)) = \varnothing$ .

**Corollary 4.14.** If  $q_0 = \ldots = q_n = 1$ , that is  $X = \mathbb{P}^n$ , then for all  $i \in \{0, \ldots, n\}$ , we have

$$\emptyset = \operatorname{Stab}(T_X(-\log D_i)) \subsetneq \operatorname{sStab}(T_X(-\log D_i)) = \operatorname{Amp}(X)$$

#### 5. Description of smooth projective toric varieties of Picard Rank two

Let X be a smooth projective toric variety of dimension n such that  $\operatorname{rk}\operatorname{Pic}(X) = 2$ . There exists  $r, s \in \mathbb{N}^*$  with r + s = n and  $a_1, \ldots, a_r \in \mathbb{N}$  with  $a_1 \leq a_2 \leq \ldots \leq a_r$  such that

(10) 
$$X = \mathbb{P}\left(\mathscr{O}_{\mathbb{P}^s} \oplus \bigoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^s}(a_i)\right) \ .$$

We keep the notation of Section 1.1. We denote by  $\pi : X \longrightarrow \mathbb{P}^s$  the projection onto the basis  $\mathbb{P}^s$ . We have the following linear equivalence,

(11)  $D_{v_i} \sim_{\text{lin}} D_{v_0} - a_i D_{w_0}$  for  $i \in \{1, \ldots, r\}$  and  $D_{w_j} \sim_{\text{lin}} D_{w_0}$  for  $j \in \{1, \ldots, s\}$ .

By (11), we deduce that Pic(X) is generated by  $D_{v_0}$  and  $D_{w_0}$ .

**Proposition 5.1** ([DDK20, Proposition 4.2.1]). Let  $D = \mu D_{w_0} + \lambda D_{v_0}$  be a *T*-invariant divisor of X with  $\mu, \lambda \in \mathbb{Z}$ . The divisor D is ample if and only if  $\mu > 0$  and  $\lambda > 0$ .

Remark 5.2. Let  $\mu$ ,  $\lambda \in \mathbb{N}^*$  and  $L = \mathscr{O}_X(\mu D_{w_0} + \lambda D_{v_0})$ . We have an isomorphism  $L \cong \pi^* \mathscr{O}_{\mathbb{P}^s}(\mu) \otimes \mathscr{O}_X(\lambda)$ .

By [CLS11, Theorem 8.1.2], the anti-canonical divisor of X is given by

(12) 
$$-K_X = \sum_{i=0}^r D_{v_i} + \sum_{j=0}^s D_{w_j} \sim_{\text{lin}} (s+1-a_1-\ldots-a_r)D_{w_0} + (r+1)D_{v_0} .$$

Hence, X is Fano if and only if  $a_1 + \ldots + a_r \leq s$ .

Let D be a reduced T-invariant Weil divisor on X. To study the stability of  $\mathscr{E} = T_X(-\log D)$  with respect to  $L = \mathscr{O}_X(Z)$  where  $Z = \mu D_{w_0} + \lambda D_{v_0} (\mu, \lambda \in \mathbb{N}^*)$ , we will study the stability of  $\mathscr{E}$  with respect to the ample  $\mathbb{Q}$ -divisor  $Z' = \nu D_{w_0} + D_{v_0}$  where  $\nu = \frac{\mu}{\lambda}$ .

5.1. Computation of the degree. Let  $P \subset M_{\mathbb{R}}$  be a polytope such that the fan  $\Sigma_P$  of P is the fan  $\Sigma$  of X. For each  $\rho \in \Sigma(1)$  we denote by  $P^{\rho}$  the facet of P corresponding to a ray  $\rho$ .

We recall that a lattice M defines a measure  $\nu$  on  $M_{\mathbb{R}}$  as the pull-back of Haar measure on  $M_{\mathbb{R}}/M$ . It is determined by the properties

- i.  $\nu$  is translation invariant,
- ii.  $\nu(M_{\mathbb{R}}/M) = 1.$

For all  $\rho \in \Sigma(1)$ , we denote by  $\operatorname{vol}(P^{\rho})$  the volume of  $P^{\rho}$  with respect to the measure determined by the lattice  $\operatorname{Span}(P^{\rho}) \cap M$ .

**Proposition 5.3** ([Dan78, Section 11]). Let (X, L) be a polarized toric variety corresponding to a lattice polytope P. For all ray  $\rho$ , we have  $\deg_L(D_{\rho}) = \operatorname{vol}(P^{\rho})$ .

5.2. Polytope corresponding to a  $\mathbb{Q}$ -ample divisor. Here, we describe the polytope corresponding to the  $\mathbb{Q}$ -polarized variety (X, L) where

$$L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \times \mathscr{O}_X(1)$$

We refer to [HNS19, Section 4].

Let  $\nu \in \mathbb{Q}^+$  such that  $\nu \neq 0$  and  $P_s = \text{Conv}(0, w_1, w_2, \dots, w_s)$ . Then,

$$P = \operatorname{Conv}\left(\nu P_s \times \{0\} \cup (a_1 + \nu) P_s \times \{v_1\} \cup \ldots \cup (a_r + \nu) P_s \times \{v_r\}\right)$$

is the polytope corresponding to the  $\mathbb{Q}$ -polarized variety (X, L). We denote by  $P^{v_i}$  (resp.  $P^{w_j}$ ) the facet of P corresponding to the ray  $\operatorname{Cone}(v_i)$  (resp.  $\operatorname{Cone}(w_j)$ ). The facet  $P^{v_i}$  is the convex hull of

 $\nu P_s \times \{0\} \cup \ldots \cup (a_{i-1}+\nu)P_s \times \{v_{i-1}\} \cup (a_{i+1}+\nu)P_s \times \{v_{i+1}\} \cup \ldots \cup (a_r+\nu)P_s \times \{v_r\}$ and  $P^{w_i}$  is isomorphic to

$$\nu P'_s \times \{0\} \cup (a_1 + \nu) P'_s \times \{v_1\} \cup \ldots \cup (a_r + \nu) P'_s \times \{v_r\}$$

where  $P'_{s} = \text{Conv}(0, w_{1}, \dots, w_{s-1})$ .

Notation 5.4. For all  $i \in \{0, 1, \ldots, r\}$ , we set  $V_i = \operatorname{vol}(P^{v_i})$ . Like for all  $j \in \{1, \ldots, s\}$ ,  $\operatorname{vol}(P^{w_j}) = \operatorname{vol}(P^{w_0})$ , we set  $\mathsf{W} = \operatorname{vol}(P^{w_0})$ .

**Proposition 5.5** ([HNS19, Proposition 4.3]). Let  $c_0, c_1, \ldots, c_r \in \mathbb{N}, \nu > 0$  and  $P_s = \text{Conv}(0, w_1, w_2, \ldots, w_s)$ . The volume of the polytope

 $P = \text{Conv}\left((c_0 + \nu)P_s \times \{0\} \cup (c_1 + \nu)P_s \times \{v_1\} \cup \ldots \cup (c_r + \nu)P_s \times \{v_r\}\right)$ 

is given by

$$\sum_{k=0}^{s} \binom{s+r}{k} \left( \sum_{d_0+\ldots+d_r=s-k} c_0^{d_0} \cdots c_r^{d_r} \right) \nu^k \; .$$

By Proposition 5.5, we have

$$W = \sum_{k=0}^{s-1} {s+r-1 \choose k} \left( \sum_{d_1+\ldots+d_r=s-k-1} a_1^{d_1} \cdots a_r^{d_r} \right) \nu^k$$
$$V_0 = \sum_{k=0}^{s} {s+r-1 \choose k} \left( \sum_{d_1+\ldots+d_r=s-k} a_1^{d_1} \cdots a_r^{d_r} \right) \nu^k$$

and for  $i \in \{1, ..., r\}$ ,

$$\mathsf{V}_{i} = \sum_{k=0}^{s} \binom{s+r-1}{k} \left( \sum_{\substack{d_{1}+\ldots+d_{i-1}\\+d_{i+1}+\ldots+d_{r}=s-k}} a_{1}^{d_{1}} \cdots a_{i-1}^{d_{i-1}} a_{i+1}^{d_{i+1}} \cdots a_{r}^{d_{r}} \right) \nu^{k} \; .$$

For  $\nu$  fixed, we give some relations between W and the numbers  $V_i$ . If  $a_1 = \ldots = a_r = 0$ , then

(13) 
$$W = \binom{s+r-1}{s-1}\nu^{s-1} \text{ and } V_i = \binom{s+r-1}{s}\nu^s$$

We now assume that  $(a_1, ..., a_r) \neq (0, ..., 0)$  with  $a_1 \leq a_2 \leq ... \leq a_r$ . Let  $z \in \{0, 1, ..., r-1\}$  such that  $a_z = 0$  and  $a_{z+1} > 0$  (we set  $a_0 = 0$ ). For  $k \in \{0, ..., s\}$  and  $i \in \{z + 1, ..., r\}$ , we set

$$W_{k} = \sum_{\substack{d_{z+1}+\dots+d_{r}=s-1-k}} a_{z+1}^{d_{z+1}} \cdots a_{r}^{d_{r}}$$
$$V_{0k} = \sum_{\substack{d_{z+1}+\dots+d_{r}=s-k}} a_{z+1}^{d_{z+1}} \cdots a_{r}^{d_{r}}$$
$$V_{ik} = \sum_{\substack{d_{z+1}+\dots+d_{i-1}\\ +d_{i+1}+\dots+d_{r}=s-k}} a_{z+1}^{d_{z+1}} \cdots a_{i-1}^{d_{i-1}} a_{i+1}^{d_{i+1}} \cdots a_{r}^{d_{r}}$$

where  $W_s = 0$ . If  $i \in \{1, ..., z\}$ , we set  $V_{ik} = V_{0k}$ . We have  $W_{s-1} = 1$  and for  $i \in \{0, ..., r\}, V_{is} = 1$ .

*Remark* 5.6. If r = 1, we set  $V_{1s} = 1$  and for  $k \in \{0, \dots, s-1\}, V_{1k} = 0$ .

**Lemma 5.7.** For  $k \in \{0, ..., s - 1\}$  and  $i \in \{1, ..., r\}$ ,  $a_i W_k + V_{ik} = V_{0k}$ .

*Proof.* If  $i \in \{1, ..., z\}$ , the equality is true because  $a_i = 0$ . We assume that  $i \in \{z + 1, ..., r\}$ , we have

$$V_{0k} = \sum_{\substack{d_{z+1}+\dots+d_r=s-k \\ a_{z+1}=\dots+a_r=s-k}} a_{z+1}^{d_{z+1}} \cdots a_r^{d_r}$$
$$= \sum_{\substack{d_{z+1}+\dots+d_r=s-k \\ a_i=0}} a_{z+1}^{d_{z+1}} \cdots a_r^{d_r} + \sum_{\substack{d_{z+1}+\dots+d_r=s-k \\ a_i\geq 1}} a_{z+1}^{d_{z+1}} \cdots a_r^{d_r}$$

The first term of the second line corresponds to the number  $V_{ik}$  and the second to  $a_i W_k$  (it suffices to replace  $d_i$  by  $d'_i + 1$ ). Hence,  $V_{0k} = V_{ik} + a_i W_k$ .

**Corollary 5.8.** *For all*  $i \in \{1, ..., r\}$ *,*  $V_0 = a_i W + V_i$ *.* 

5.3. A necessary condition for stability. In this part, we adapt some results of [HNS19, Section 4] for the study of the stability of  $T_X(-\log D)$ .

The following lemma will be useful in the proof of Proposition 5.10 which is the main result of this part. Let  $z \in \{0, \ldots, r-1\}$  such that  $a_z = 0$  and  $a_{z+1} > 0$ . We have the following lemma.

Lemma 5.9 ([HNS19, Lemma 4.2]).

Let  $I' \subset \{0, 1, \ldots, r\}$  and  $G = \text{Span}(v_i : i \in I')$ . The vector  $a_1v_1 + \ldots + a_rv_r$ belongs to G if and only if

i.  $\{z + 1, ..., r\} \subset I'$  or ii.  $\{0, ..., z\} \subset I'$ ,  $card(\{z + 1, ..., r\} \setminus I') \geq 1$  and  $a_i = a_j$  for all  $i, j \in \{z + 1, ..., r\} \setminus I'$ .

Let 
$$\Delta \subset \Sigma(1)$$
 and  $D = \sum_{\rho \in \Delta} D_{\rho}$ .  
Let  $I_{\Sigma} = \{\operatorname{Cone}(v_0), \dots, \operatorname{Cone}(v_r)\}$  and  $J_{\Sigma} = \{\operatorname{Cone}(w_0), \dots, \operatorname{Cone}(w_s)\}$ . We set  
 $I = \{i \in \{0, 1, \dots, r\} : \operatorname{Cone}(v_i) \in I_{\Sigma} \setminus (I_{\Sigma} \cap \Delta)\}$  and  
 $J = \{j \in \{0, 1, \dots, s\} : \operatorname{Cone}(w_j) \in J_{\Sigma} \setminus (J_{\Sigma} \cap \Delta)\}$ .

Let  $L \in \operatorname{Amp}(X)$ . To study the stability of  $\mathscr{E} = T_X(-\log D)$ , it suffices to compare  $\mu_L(\mathscr{E})$  and  $\mu_L(\mathscr{E}_G)$  where  $G = \operatorname{Span}(v_i, w_j : i \in I', j \in J')$  with  $I' \subset I, J' \subset J$  and  $1 \leq \dim G < (r+s)$ . By Proposition 5.3, and by using of (7), (8) we have

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} \left( \sum_{i \in I} \mathsf{V}_i + \operatorname{card}(J) \cdot \mathsf{W} \right)$$

and

$$\mu_L(\mathscr{E}_G) = \frac{1}{\dim G} \left( \sum_{i \in I'} \mathsf{V}_i + \operatorname{card}(J') \cdot \mathsf{W} \right) \;.$$

Here is a variant of [HNS19, Proposition 4.1] which will be useful for us in the study of stability of  $T_X(-\log D)$ .

**Proposition 5.10.** The logarithmic tangent bundle  $\mathscr{E} = T_X(-\log D)$  is stable (resp. semi-stable) with respect to L if and only if

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} \left( \sum_{i \in I} \mathsf{V}_i + \operatorname{card}(J) \cdot \mathsf{W} \right)$$

is greater than (resp. greater than or equal to) the maximum of

- (1)  $V_{i_0}$  where  $i_0 = \min I$  if  $I \neq \emptyset$ ;
- (2)  $\frac{1}{r'}\sum_{i \in T} \mathsf{V}_i$ , if  $r' = \dim \operatorname{Span}(v_i : i \in I) \neq 0$ ; (3)  $\frac{\operatorname{card}(J) \cdot \mathsf{W}}{s'}$ , if  $s' = \dim \operatorname{Span}(w_j : j \in J) \neq 0$  and s' < r + s;
- (4)  $\frac{1}{s+k} \left( \sum_{i \in I'} \mathsf{V}_i + (s+1) \mathsf{W} \right), \text{ if } \operatorname{card}(J') = s+1, \ k = \operatorname{card}(I') < r \text{ and} \\ \{z+1, \dots, r\} \subset I' \subset I;$ (5)  $\frac{1}{s+k} \left( \sum_{i \in I'} \mathsf{V}_i + (s+1) \mathsf{W} \right)$ , if  $\operatorname{card}(J') = s+1$ ,  $k = \operatorname{card}(I') < r$  and
  - $I' \subset I$  such that the condition ii. of Lemma 5.9 is verified.

*Proof.* Let  $G = \text{Span}(v_i, w_j : i \in I', j \in J')$  where  $I' \subset I$  and  $J' \subset J$ . The expressions (1) of Proposition 5.10 corresponds to  $G = \text{Span}(v_{i_0})$ , (2) corresponds to  $G = \text{Span}(v_i : i \in I)$  and (3) corresponds to  $G = \text{Span}(w_j : j \in J)$ .

• If  $\operatorname{card}(J') = 0$ , then for  $\varnothing \subsetneq I' \subseteq I$ , we have dim  $G \leq r$  and

$$\mu_L(\mathscr{E}_G) = \frac{1}{\dim G} \sum_{i \in I'} \mathsf{V}_i \; ;$$

this number is less than or equal to the maximum of the numbers given in (1) and (2).

• If  $\operatorname{card}(I') = 0$ , then for  $\emptyset \subsetneq J' \subseteq J$  such that  $\dim G < r + s$ , we have

$$\mu_L(\mathscr{E}_G) = \frac{\operatorname{card}(J') \cdot W}{\dim G} ;$$

this number is less than or equal to that given in (3).

• If  $\operatorname{card}(I') = r + 1$ , then  $\dim G < r + s$  if and only if  $s' := \operatorname{card}(J') < s$ . If  $1 \leq s' < s$ , then

$$\mu_L(\mathscr{E}_G) = \frac{1}{r+s'} \left( \sum_{i \in I'} \mathsf{V}_i + s' \mathsf{W} \right) \le \max\left( \frac{1}{r} \sum_{i \in I'} \mathsf{V}_i, \mathsf{W} \right) ;$$

this number is less than or equal to the maximum of numbers given in (2) and (3). • If  $1 \leq \operatorname{card}(I') \leq r$ ,  $1 \leq \operatorname{card}(J') \leq s$  and  $\dim G < r + s$ , then  $\mu_L(\mathscr{E}_G)$  is less than or equal to the maximum of numbers given in (1), (2) and (3).

• It remains to study the case where  $\operatorname{card}(J') = s + 1$  and  $1 \leq \operatorname{card}(I') < r$  (because if  $\operatorname{card}(I') \ge r$ , then  $\dim G = r + s$ ). We will treat it in two cases.

First case :  $a_r = 0$ . For all  $i \in \{1, \ldots, r\}$ ,  $V_i = V_0$ . If  $r' = \operatorname{card}(I')$  and  $1 \le r' < r$ , then

$$\mu_L(\mathscr{E}_G) = \frac{1}{r'+s} \left( \sum_{i \in I'} \mathsf{V}_i + (s+1)\mathsf{W} \right) \le \max\left(\mathsf{V}_0, \frac{(s+1)\mathsf{W}}{s}\right)$$

Second case :  $a_r > 0$ . We set  $r' = \operatorname{card}(I')$ . If I' satisfies the first (resp. second) condition of Lemma 5.9, we get (4) (resp. (5)).

If I' doesn't satisfy the conditions of Lemma 5.9, then dim G = r' + (s + 1). Moreover, if r' + (s+1) < r+s, the number  $\mu_L(\mathscr{E}_G)$  is less than or equal to the maximum of the numbers given in (1) and (3).

#### A. NAPAME

Remark 5.11. If  $a_1 = \ldots = a_r = 0$ , to check the stability of  $\mathscr{E}$  with respect to L, it is enough to compare  $\mu_L(\mathscr{E})$  with the numbers given by the points 1, 2 and 3 of Proposition 5.10.

To check that  $\mathscr{E} = T_X(-\log D)$  is not semi-stable, we will need the lemma below. The purpose of the lemma is to avoid comparing  $\mu_L(\mathscr{E})$  with all the numbers given in the Proposition 5.10.

#### Lemma 5.12.

- (1) If  $a_r \geq 2$ , then  $s \mathsf{V}_0 (s+1)\mathsf{W} \geq s \mathsf{V}_r$ .
- (2) Let  $r \ge 2$  and  $i \in \{1, \ldots, r-1\}$ . If  $a_r > a_i$ , then  $V_i W \ge V_r$ .

*Proof.* If  $a_r \ge 2$ , then  $\left(s - \frac{s+1}{a_r}\right) = \frac{a_r s - (s+1)}{a_r} \ge \frac{2s - (s+1)}{a_r} \ge 0$  because  $s \ge 1$ . Hence,

$$s \mathsf{V}_0 - (s+1)\mathsf{W} = s \mathsf{V}_0 - \frac{s+1}{a_r} (\mathsf{V}_0 - \mathsf{V}_r) \quad \text{because } a_r \mathsf{W} = \mathsf{V}_0 - \mathsf{V}_r$$
$$= \left(s - \frac{s+1}{a_r}\right) \mathsf{V}_0 + \frac{s+1}{a_r} \mathsf{V}_r$$
$$\ge \left(s - \frac{s+1}{a_r}\right) \mathsf{V}_r + \frac{s+1}{a_r} \mathsf{V}_r = s \mathsf{V}_r \ .$$

Since  $V_0 = a_i W + V_i = a_r W + V_r$ , we have  $V_i = (a_r - a_i)W + V_r$ . If  $a_r > a_i$ , then  $a_r - a_i \ge 1$ ; therefore  $V_i \ge W + V_r$ , i.e  $V_i - W \ge V_r$ .

5.4. Stability of logarithmic tangent bundle on  $X = \mathbb{P}^r \times \mathbb{P}^s$ . In this part, we assume that  $a_1 = a_2 = \ldots = a_r = 0$ . We have  $X \cong \mathbb{P}^s \times \mathbb{P}^r$ . We denote by  $\pi_1 : X \longrightarrow \mathbb{P}^s$  and  $\pi_2 : X \longrightarrow \mathbb{P}^r$  the projection maps.

We first show that  $T_X(-\log D)$  is decomposable when  $\operatorname{card}(\Delta) \in \{1, 2\}$ .

**Lemma 5.13.** For  $i \in \{0, \ldots, r\}$ , the vector bundle  $\mathscr{E} = T_X(-\log D_{v_i})$  satisfies

$$\mathscr{E} = \left( \bigoplus_{k=0, \, k \neq i}^{r} \mathscr{E}_{F_k} \right) \oplus \mathscr{E}_G$$

where  $F_k = \text{Span}(v_k)$  and  $G = \text{Span}(w_0, \dots, w_s)$ .

**Lemma 5.14.** For  $j \in \{0, \ldots, s\}$ , the vector bundle  $\mathscr{E} = T_X(-\log D_{w_j})$  satisfies

$$\mathscr{E} = \left( \bigoplus_{k=0, \, k \neq j}^{s} \mathscr{E}_{G_k} \right) \oplus \mathscr{E}_F$$

where  $G_k = \operatorname{Span}(w_k)$  and  $F = \operatorname{Span}(v_0, \ldots, v_r)$ .

**Lemma 5.15.** Let  $i, i' \in \{0, \ldots, r\}$  distinct and  $j, j' \in \{0, \ldots, s\}$  distinct. We have

$$T_X(-\log(D_{v_i} + D_{v_{i'}})) \cong T_{\mathbb{P}^s} \oplus T_{\mathbb{P}^r}(-\log(\pi_2(D_{v_i}) + \pi_2(D_{v_{i'}})))$$
$$T_X(-\log(D_{w_i} + D_{w_{i'}})) \cong T_{\mathbb{P}^s}(-\log(\pi_1(D_{w_i}) + \pi_1(D_{w_{i'}}))) \oplus T_{\mathbb{P}^r}$$

For the proofs of these three lemmas it suffices to work with the family of multifiltrations as in the proof of Theorem 3.9. If  $i \in \{0, ..., r\}$  and  $j \in \{0, ..., s\}$ , by Theorem 3.9 the vector bundle  $T_X(-\log(D_{v_i} + D_{w_j}))$  is decomposable.

We now study (semi)stability. In this table, we give the values of  $\nu$  for which  $\mathscr{E} = T_X(-\log D)$  is (semi)stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ .

Divisor D $\operatorname{Stab}(\mathscr{E})$  $sStab(\mathcal{E})$ References  $\nu = \frac{s+1}{r}$  $\nu = \frac{r+1}{s}$ Theorem 5.16 Ø  $D_{v_i}$ ,  $0 \le i \le r$ Theorem 5.16Ø  $D_{w_i}$ ,  $0 \le j \le s$  $\nu = \frac{s}{r}$ Theorem 5.19 Ø  $D_{v_i} + D_{w_j}$  $\begin{array}{ll} D_{v_i} + D_{v_j} \ , & 0 \leq i < j \leq r \\ D_{w_i} + D_{w_j} \ , & 0 \leq i < j \leq s \end{array}$ Theorem 5.18 Ø Ø Theorem 5.18 Ø Ø

TABLE 1. Stability of  $T_X(-\log D)$  when  $a_1 = \ldots = a_r = 0$ 

By (13), we have  $V = \frac{r\nu}{2}W$ .

**Theorem 5.16.** Let  $i \in \{0, 1, ..., r\}$  and  $j \in \{0, 1, ..., s\}$ . For any  $L \in Amp(X)$ , the logarithmic tangent bundles  $T_X(-\log D_{v_i})$  and  $T_X(-\log D_{w_i})$  are not stable with respect to L. We have :

- (1)  $T_X(-\log D_{v_i})$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \frac{s+1}{r}$ ;
- (2)  $T_X(-\log^r D_{w_j})$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \frac{r+1}{s}$ .

*Proof.* We start with  $\mathscr{E} = T_X(-\log D_{v_i})$ . We have :

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} (r \mathsf{V} + (s+1)\mathsf{W}) = \frac{r^2 \nu + s^2 + s}{s(r+s)} \mathsf{W} = \frac{r^2 \nu + s^2 + s}{r \nu(r+s)} \mathsf{V}$$

By Proposition 5.10, to have stability or semi-stability, it is enough to compare  $\mu_L(\mathscr{E})$  with  $\max\left(\frac{(s+1)W}{s}, \mathsf{V}\right)$ . If  $\mu_L(\mathscr{E}) \ge \mathsf{V}$ , then  $\frac{r^2\nu + s^2 + s}{r\nu(r+s)} \ge 1$ , i.e.  $(r^2\nu + s^2 + s) \ge (r^2\nu + rs\nu)$ ; hence,  $\nu \leq \frac{s+1}{r} \text{. If } \mu_L(\mathscr{E}) \geq \frac{(s+1) \mathbb{W}}{s} \text{, then } \frac{r^2 \nu + s^2 + s}{s(r+s)} \geq \frac{s+1}{s} \text{, i.e } \nu \geq \frac{s+1}{r} \text{.}$ 

If  $\nu \neq \frac{s+1}{r}$ , the numbers  $\mu_L(\mathscr{E}) - \mathsf{V}$  and  $\mu_L(\mathscr{E}) - \frac{s+1}{s}\mathsf{W}$  are non-zero and have opposite signs. Therefore,  $T_X(-\log D_{v_i})$  is not semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1).$ 

If  $\nu = \frac{s+1}{r}$ , then  $\mu_L(\mathscr{E}) = \mathsf{V} = \frac{s+1}{s}\mathsf{W}$ . Thus,  $T_X(-\log D_{v_i})$  is semi-stable but not stable with respect to  $\pi^*\mathscr{O}_{\mathbb{P}^s}(\nu) \overset{\otimes}{\otimes} \mathscr{O}_X(1)$ .

If we regard the case where  $\mathscr{E} = T_X(-\log D_{w_j})$ , it is enough to compare  $\mu_L(\mathscr{E})$ with  $\max\left(\frac{(r+1)\mathsf{V}}{r},\mathsf{W}\right)$ . We get the result by exchanging the roles of r and sand also the roles of  $\mathsf{V}$  and  $\mathsf{W}$  in the computation above. 

Corollary 5.17. With the decomposition given in Lemma 5.13 (resp. Lemma 5.14), the vector bundle  $T_X(-\log D_{v_i})$  (resp.  $T_X(-\log D_{w_i})$ ) is poly-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \frac{s+1}{r}$  (resp.  $\nu = \frac{r+1}{s}$ ). **Theorem 5.18.** Let  $i, i' \in \{0, ..., r\}$  distinct and  $j, j' \in \{0, ..., s\}$  distinct. For any  $L \in \operatorname{Amp}(X)$ , the logarithmic tangent bundles  $T_X(-\log(D_{v_i} + D_{v_{i'}}))$  and  $T_X(-\log(D_{w_j} + D_{w_{j'}}))$  are not semi-stable with respect to L.

*Proof.* Let  $\mathscr{E} = T_X(-\log(D_{v_i} + D_{v_{i'}}))$ . We have :

$$\mu_L(\mathscr{E}) = \frac{1}{r+s}((r-1)\mathsf{V} + (s+1)\mathsf{W}) = \frac{r(r-1)\nu + s^2 + s}{s(r+s)}\,\mathsf{W} = \frac{r(r-1)\nu + s^2 + s}{r\nu(r+s)}\,\mathsf{V}\,.$$

We assume that r = 1. We have  $\mu_L(\mathscr{E}) = W$ . If we set  $F = \text{Span}(w_0, \ldots, w_s)$ , we have  $\mu_L(\mathscr{E}_F) = \frac{(s+1)W}{s}$ ; thus  $T_X(-\log(D_{v_i} + D_{v_{i'}}))$  is not semi-stable with respect to L.

We now assume that  $r \ge 2$ . To check the stability or semi-stability, it is enough to compare  $\mu_L(\mathscr{E})$  with  $\max\left(\frac{(s+1)W}{s}, \mathsf{V}\right)$ . If  $\mu_L(\mathscr{E}) \ge \mathsf{V}$ , then  $\frac{r(r-1)\nu + s^2 + s}{r\nu(r+s)} \ge 1$ ; i.e  $\nu \le \frac{s}{r}$ . If  $\mu_L(\mathscr{E}) \ge \frac{(s+1)W}{s}$ , then  $\frac{r(r-1)\nu + s^2 + s}{s(r+s)} \ge \frac{s+1}{s}$ ; i.e  $\nu \ge \frac{s+1}{r-1} > \frac{s}{r}$ .

Like  $\nu$  cannot satisfy this two conditions, we deduce that  $T_X(-\log(D_{v_i} + D_{v_{i'}}))$  is not semi-stable with respect to L.

**Theorem 5.19.** Let  $i \in \{0, ..., r\}$ ,  $j \in \{0, ..., s\}$  and  $D = D_{v_i} + D_{w_j}$ . We have :

- (1)  $T_X(-\log D)$  is not stable with respect to any  $L \in \operatorname{Amp}(X)$ .
  - (2)  $T_X(-\log D)$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \frac{s}{r}$ .

Proof. We have

$$\mu_L(\mathscr{E}) = \frac{1}{r+s}(r\mathsf{V} + s\mathsf{W}) = \frac{r^2\nu + s^2}{s(r+s)}\mathsf{W} = \frac{r^2\nu + s^2}{r\nu(r+s)}\mathsf{V}$$

To check stability or semi-stability, it is enough to compare  $\mu_L(\mathscr{E})$  with  $\max(\mathsf{V}, \mathsf{W})$ . If  $\mu_L(E) \ge \mathsf{V}$ , then  $\frac{r^2\nu + s^2}{r\nu(r+s)} \ge 1$ ; i.e  $\nu \le \frac{s}{r}$ . If  $\mu_L(E) \ge \mathsf{W}$ , then  $\frac{r^2\nu + s^2}{s(r+s)} \ge 1$ ; i.e  $\nu \ge \frac{s}{r}$ . Hence,  $T_X(-\log D)$  is semi-stable with respect to  $\pi^*\mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only

if  $\nu = \frac{s}{r}$ .

**Corollary 5.20.** With the decomposition given in theorem 3.9,  $T_X(-\log(D_{v_i} + D_{w_j}))$  is poly-stable with respect to  $\pi^* \mathcal{O}_{\mathbb{P}^s}(\nu) \otimes \mathcal{O}_X(1)$  if and only if  $\nu = \frac{s}{r}$ .

According to (11) and (12), when  $a_1 = \ldots = a_r = 0$ , we have :

 $D_{v_i} \sim_{\text{lin}} D_{v_0}$ ,  $D_{w_j} \sim_{\text{lin}} D_{w_0}$  and  $-K_X \sim_{\text{lin}} (s+1)D_{w_0} + (r+1)D_{v_0}$ .

By the above study when  $\operatorname{sStab}(T_X(-\log D)) \neq \emptyset$ , we see that  $T_X(-\log D)$  is semi-stable with respect to L if and only if  $L \cong \mathcal{O}_X(-\alpha (K_X + D))$  with  $\alpha \in \mathbb{N}^*$ .

#### 6. Stability of $T_X(-\log D)$ on smooth toric varieties of Picard Rank two when $a_r \ge 1$

In this part, we will study the stability of  $T_X(-\log D)$  when X is given by (10) with  $a_r \ge 1$ . If  $D = \sum_{\rho \in \Delta} D_\rho$  with  $\Delta \subset \Sigma(1)$ , by Corollary 4.8, we will only study the case where card  $\Delta \in \{1, 2\}$ . The case card  $\Delta = 0$  was treated by HERING-NILL-

the case where card  $\Delta \in \{1, 2\}$ . The case card  $\Delta = 0$  was treated by HERING-NILL-Süss in [HNS19].

Let  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  be an element of  $\operatorname{Amp}(X) \subset N^1(X) \otimes_{\mathbb{Z}} \mathbb{R}$ . We recall that the numbers  $\mathsf{V}_0, \ldots, \mathsf{V}_r$  defined on Section 5.2 are polynomials of  $\nu$  of degree s and  $\mathsf{W}$  is polynomial of degree s - 1. If  $\mathscr{E} = T_X(-\log D)$ , the number  $\mu_L(\mathscr{E})$  is polynomial of degree at most s. Let  $\mathsf{P}_0, \mathsf{P}_1, \mathsf{P}_2$  and  $\mathsf{Q}$  be the polynomials of  $\nu$  defined by

$$\begin{split} \mathsf{P}_0 &= \mu_L(\mathscr{E}) - \mathsf{V}_0 \;, \quad \mathsf{P}_1 = \mu_L(\mathscr{E}) - \mathsf{V}_1 \;, \quad \mathsf{P}_2 = \mu_L(\mathscr{E}) - \mathsf{V}_2 \quad \text{and} \quad \mathsf{Q} = \mu_L(\mathscr{E}) - \mathsf{W} \;. \end{split}$$
 We first recall the sign changes rule of Descartes. We refer to [Mig89, Chapter 5, Section 4.3].

**Theorem 6.1** (Descartes). Let  $P = c_n X^n + c_{n-1} X^{n-1} + \ldots + c_0$  be a polynomial with real coefficients where  $c_n c_0 \neq 0$ . Let p the number of sign changes in the sequence  $(c_0, \ldots, c_n)$  of its coefficients and q the numbers of positive real roots, counted with their order of multiplicity. Then, there is  $m \in \mathbb{N}$  such that q = p - 2m.

Under certain conditions on  $a_i$ , r and s, these polynomials ( $\mathsf{P}_0, \mathsf{P}_1, \mathsf{P}_2$  and  $\mathsf{Q}$ ) have respectively one or no positive root. If the positive root exists, we denote by

- $\nu_i$  the unique positive root of  $\mathsf{P}_i$  where  $i \in \{0, 1, 2\}$
- $\nu_3$  the unique positive root of Q.

In these tables, we give the values of  $\nu$  for which  $\mathscr{E} = T_X(-\log D)$  is stable or semi-stable with respect to  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ .

Divisor D	Condition on $r$ and $a_i$	Condition on $s$	$\mathrm{Stab}(\mathscr{E})$	$\mathrm{sStab}(\mathscr{E})$
$ \begin{array}{c} D_{w_j} \\ 0 \le j \le s \\ \text{Theorem 6.2} \end{array} $	$r \ge 1$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$ \begin{array}{c} D_{v_i} \\ 1 \le i \le r-1 \\ \text{Prop. 6.4} \end{array} $	$r \ge 2$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$D_{v_r}$	$r \ge 1, a_r = 1$ and $a_{r-1} = 0$	$s \ge 1$	$0 < \nu < \nu_0$	$0 < \nu \leq \nu_0$
Theorem 6.6	$r \ge 1$ and $(a_r \ge 2$ or $a_{r-1} \ne 0)$	$s \ge 1$	Ø	Ø
	r = 1	$s \ge 1$	$0 < \nu < \nu_1$	$0 < \nu \le \nu_1$
$D_{v_0}$	$r \ge 2$ and $a_1 < a_r$	$s \ge 1$	Ø	Ø
Theorem 6.9	$r \ge 2$ and	$a \ge \frac{s+1}{r-1}$	Ø	Ø
Lemma 6.8	$a_1 = a_r = a$	$\frac{s}{r} \le a < \frac{s+1}{r-1}$	$0 < \nu < \nu_1$	$0 < \nu \le \nu_1$
Theorem 6.11		a r < s	$\nu_3 < \nu < \nu_1$	$\nu_3 \le \nu \le \nu_1$

TABLE 2. Stability of  $T_X(-\log D)$  when  $a_r \ge 1$ 

#### A. NAPAME

Divisor $D$	Condition on $r$ and $a_i$	Condition on $s$	$\mathrm{Stab}(\mathscr{E})$	$\mathrm{sStab}(\mathscr{E})$
$D_{w_i} + D_{w_j}$ $0 \le i < j \le s$ Theorem 6.2	$r \ge 1$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$D_{v_i} + D_{v_j}$ $1 \le i < j \le r$ Corollary 6.5	$r \ge 2$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$D_{v_i} + D_{w_j} , j \ge 0$ and $1 \le i \le r - 1$ Proposition 6.4	$r \ge 2$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$D_{v_r} + D_{w_j} , \ j \ge 0$ Corollary 6.7	$r \ge 1$ and $a_r \ge 1$	$s \ge 1$	Ø	Ø
$D_{v_0} + D_{w_j}, 0 \le j \le s$	r = 1	$s \ge 1$	Ø	$\nu = \nu_3$
Theorem $6.9$	$r \ge 2$ and $a_1 < a_r$	$s \ge 1$	Ø	Ø
Lemma 6.8	$r \ge 2$ and	$s \le a(r-1)$	Ø	Ø
Theorem 6.13	$a_1 = a_r = a$	s > a(r-1)	Ø	$\nu = \nu_3$
$D_{v_0} + D_{v_i} , \ 2 \le i \le r$	$r \ge 2$ and $a_1 < a_r$	$s \ge 1$	Ø	Ø
Lemma 6.8	$r \ge 2$ and	$s \le a(r-1)$	Ø	Ø
Theorem 6.13	$a_1 = a_r = a$	s > a(r-1)	Ø	$\nu = \nu_3$

TABLE 3. Stability of  $\mathscr{T}_X(-\log D)$  when  $a_r \geq 1$ 

TABLE 4. Stability of  $T_X(-\log D)$  when  $a_r \ge 1$ 

Divisor $D$	Condition on $r$ and $a_i$	Condition on $s$	$\mathrm{Stab}(\mathscr{E})$	$\operatorname{sStab}(\mathscr{E})$
	r = 1	$s \ge 1$	Ø	$\nu > 0$
$D_{v_0} + D_{v_1}$	$r \ge 2$ and $0 = a_1 < a_r$	$s \ge 1$	Ø	Ø
	$r \geq 2$ and	$s \le a(r-1)$	Ø	Ø
Theorem $6.9$	$a_1 = a_r = a$	s > a(r-1)	Ø	$\nu = \nu_3$
Prop. 6.14	r=2 and	$s \le \delta_2$	Ø	Ø
Theorem 6.13	$0 < a_1 < a_2$	$s > \delta_2$	Ø	$\nu = \nu_3$
Prop. 6.17	$r \ge 3$ and $a_2 < a_r$	$s \ge 1$	Ø	Ø
Lemma $6.15$	$r \geq 3$ and	$s \leq \delta_r$	Ø	Ø
Prop. 6.19	$0 < a_1 < a_2 = \ldots = a_r$	$s > \delta_r$	Ø	$\nu = \nu_3$

6.1. The case  $D = D_{w_j}$  and  $D = D_{w_j} + D_{w_i}$ . Let  $i, j \in \{0, \ldots, s\}$  distinct and  $\mathscr{E} = T_X(-\log D_{w_i}), \ \mathscr{F} = T_X(-\log(D_{w_i} + D_{w_j}))$ . If  $a_r \geq 1$ , the Lemma 5.14 is again true. The vector bundles  $\mathscr{E}$  and  $\mathscr{F}$  are decom-

If  $a_r \ge 1$ , the Lemma 5.14 is again true. The vector bundles  $\delta$  and  $\mathcal{F}$  are decomposable and we have

$$\mathscr{E} = \left( \bigoplus_{k=0, \, k \neq i}^{s} \mathscr{E}_{G_{k}} \right) \oplus \mathscr{E}_{F} \quad \text{and} \quad \mathscr{F} = \left( \bigoplus_{k=0, \, k \neq i}^{s} \mathscr{F}_{G_{k}} \right) \oplus \mathscr{F}_{F}$$

where  $G_k = \text{Span}(w_k)$  and  $F = \text{Span}(v_0, \dots, v_r)$ .

**Theorem 6.2.** Let  $(a_1, \ldots, a_r) \neq (0, \ldots, 0)$ . For any  $L \in Amp(X)$ , the vector bundles  ${\mathscr E}$  and  ${\mathscr F}$  are not semi-stable with respect to L .

*Proof.* Let  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ , we have

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} \left( s \mathsf{W} + \left( \mathsf{V}_0 + \ldots + \mathsf{V}_r \right) \right)$$

and

$$\mu_L(\mathscr{F}) = \frac{1}{r+s} \left( \left( s-1 \right) \mathsf{W} + \left( \mathsf{V}_0 + \ldots + \mathsf{V}_r \right) \right) ;$$

thus,  $\mu_L(\mathscr{F}) < \mu_L(\mathscr{E})$ . The points 4 and 5 of Proposition 5.10 are not verifying. To check that  $\mathscr{E}$  (resp.  $\mathscr{F}$ ) is not semi-stable with respect to L it is enough to compare  $\mu_L(\mathscr{E})$  (resp.  $\mu_L(\mathscr{F})$ ) with

$$\max\left(\mathsf{V}_0, \frac{1}{r}(\mathsf{V}_0 + \mathsf{V}_1 + \ldots + \mathsf{V}_r)\right)$$

By Corollary 5.8, we have  $V_0 = a_r W + V_r$ . Like  $a_r \ge 1$ , we have  $V_0 \ge W + V_r$ , i.e  $V_0 - W \ge V_r$ . Thus,

$$\begin{aligned} (r+s)\left(\mathsf{V}_0-\mu_L(\mathscr{E})\right) &= s(\mathsf{V}_0-\mathsf{W}) + (r\,\mathsf{V}_0-(\mathsf{V}_0+\ldots+\mathsf{V}_{r-1})) - \mathsf{V}_r\\ &\geq s(\mathsf{V}_0-\mathsf{W}) - \mathsf{V}_r \quad \text{because } \mathsf{V}_i \leq \mathsf{V}_0\\ &\geq (s-1)\mathsf{V}_r \ . \end{aligned}$$

If  $s \geq 2$ , we have  $\mathsf{V}_0 - \mu_L(\mathscr{E}) > 0$  and  $\mathsf{V}_0 - \mu_L(\mathscr{F}) > 0$ . Thus,  $\mathscr{E}$  and  $\mathscr{F}$  are not semi-stables with respect to L.

We now assume that s = 1. By using the formula defining  $V_i$  and W given in Section 5.2, we have

W = 1,  $V_0 = (a_1 + \ldots + a_r) + r\nu$  and  $V_i = V_0 - a_i$  for  $i \in \{1, \ldots, r\}$ . We have

$$\mu_L(\mathscr{E}) = \frac{1}{r+1} \left( 1 + (r+1)\mathsf{V}_0 - (a_1 + \ldots + a_r) \right)$$

and

$$\mathbf{V}(\mathbf{V}_0 + \ldots + \mathbf{V}_r) = \frac{1}{r} \left( (r+1)\mathbf{V}_0 - (a_1 + \ldots + a_r) \right)$$

Therefore,  $\frac{\mathsf{V}_0 + \ldots + \mathsf{V}_r}{r} - \mu_L(\mathscr{E}) = \frac{(r+1)\mathsf{V}_0 - (a_1 + \ldots + a_r) - r}{r(r+1)}$ . We have  $(r+1)V_0 - (a_1 + \ldots + a_r) - r = (r+1)(a_1 + \ldots + a_r) + (r+1)r\nu$ 

$$-(a_1 + \ldots + a_r) - r$$
  
=  $r(a_1 + \ldots + a_r - 1) + (r+1)r\nu > 0$ 

because  $(a_1 + \ldots + a_r - 1) \ge 0$  and  $\nu > 0$ . Thus,  $\mathscr{E}$  and  $\mathscr{F}$  are not semi-stables with respect to L.

6.2. The case where  $\operatorname{Cone}(v_0) \notin \Delta$ . If  $i \in \{1, \ldots, r\}$  and  $j \in \{0, \ldots, s\}$ , by Theorem 3.9, the logarithmic tangent bundle  $T_X(-\log(D_{v_i} + D_{w_j}))$  is decomposable. Let  $r \ge 2$  and  $D \in \{D_{v_i} + D_{v_j} : 1 \le i < j \le r\}$ ; we set  $\mathscr{E} = T_X(-\log D)$ , we have the following proposition :

**Lemma 6.3.** The logarithmic tangent bundle  $\mathcal{E}$  is decomposable. If  $a_i < a_j$ , then

$$\mathscr{E} = \left( \bigoplus_{l=0}^{s} \mathscr{E}_{G_{l}} \right) \oplus \left( \bigoplus_{k=0, \, k \notin \{i, \, j\}}^{r} \mathscr{E}_{F_{k}} \right)$$

where  $G_l = \text{Span}(w_l)$  and  $F_k = \text{Span}(v_k)$ . If  $a_i = a_j$ , then

$$\mathscr{E} = \mathscr{E}_G \oplus \mathscr{E}_F$$

where  $G = \operatorname{Span}(w_l, v_k : l \in \{0, \dots, s\}, k \in \{0, \dots, r\} \setminus \{i, j\})$  and  $F = \operatorname{Span}(v_i)$ or  $F = \operatorname{Span}(v_j)$ .

*Proof.* For the proof we use the families of multifiltrations. We use the Lemma 5.9 to explain the choice of this decomposition.

If  $D \in \{D_{v_i} : 1 \leq i \leq r\}$ , we will not search to know if  $\mathscr{E} = T_X(-\log D)$  is decomposable because it depends on the numbers r, s and  $a_1, \ldots, a_r$ . In particular, if we assume that r = 2, s = 1,  $a_1 = 0$  and  $a_2 = 1$ , then  $\mathscr{E} = T_X(-\log D_{v_1})$  is decomposable and  $\mathscr{E} = \mathscr{E}_F \oplus \mathscr{E}_G$  where  $F = \operatorname{Span}(v_2, w_1)$  and  $G = \operatorname{Span}(v_0)$ . But  $\mathscr{F} = T_X(-\log D_{v_2})$  is not decomposable.

We first study the stability of  $T_X(-\log D)$  when  $r \ge 2$  and

$$D \in \{D_{v_i} : 1 \le i \le r - 1\} \cup \{D_{v_i} + D_{w_j} : 1 \le i \le r - 1 \text{ and } 0 \le j \le s\}$$
$$\cup \{D_{v_i} + D_{v_j} : 1 \le i < j \le r\}.$$

**Proposition 6.4.** Let  $r \ge 2$ ,  $(a_1, \ldots, a_r) \ne (0, \ldots, 0)$ ,  $i \in \{1, \ldots, r-1\}$  and  $j \in \{0, \ldots, s\}$ . For any  $L \in \operatorname{Amp}(X)$ , the logarithmic tangent bundles  $T_X(-\log D_{v_i})$  and  $T_X(-\log(D_{v_i} + D_{w_j}))$  are not semi-stables with respect to L.

*Proof.* We set  $\mathscr{E} = T_X(-\log D_{v_i})$  and  $\mathscr{F} = T_X(-\log(D_{v_i} + D_{w_j}))$ . Let  $L \in \operatorname{Amp}(X)$ , we have

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} \left( (s+1)W + (V_0 + \ldots + V_{i-1} + V_{i+1} + \ldots + V_r) \right)$$

and  $\mu_L(\mathscr{F}) < \mu_L(\mathscr{E})$ . We will show that  $\mathsf{V}_0 > \mu_L(\mathscr{E})$ . We have

$$(r+s)(\mathsf{V}_0 - \mu_L(\mathscr{E})) = (s+1)(\mathsf{V}_0 - \mathsf{W}) - \mathsf{V}_r + ((r-1)\mathsf{V}_0 - (\mathsf{V}_0 + \ldots + \mathsf{V}_{i-1} + \mathsf{V}_{i+1} + \ldots + \mathsf{V}_{r-1})) \geq (s+1)(\mathsf{V}_0 - \mathsf{W}) - \mathsf{V}_r \geq (s+1)\mathsf{V}_r - \mathsf{V}_r = s \mathsf{V}_r$$

By Proposition 5.10,  $\mathscr{E}$  and  $\mathscr{F}$  are not semi-stables with respect to L.

**Corollary 6.5.** Let  $r \ge 2$  and  $(a_1, \ldots, a_r) \ne (0, \ldots, 0)$ . Let  $L \in Amp(X)$ ; for all  $i, j \in \{1, \ldots, r\}$  distinct, the logarithmic tangent bundle  $T_X(-\log(D_{v_i} + D_{v_j}))$  is not semi-stable with respect to L.

*Proof.* If we set  $\mathscr{G} = T_X(-\log(D_{v_i} + D_{v_j}))$ , by using the proof of Proposition 6.4, we have  $\mu_L(\mathscr{G}) < \mu_L(\mathscr{E}) < \mathsf{V}_0$ . Thus,  $\mathscr{G}$  is not semi-stable with respect to L.  $\Box$ 

We now study the stability of  $T_X(-\log D)$  when

 $D \in \{D_{v_r}\} \cup \{D_{v_r} + D_{w_i} : 0 \le j \le s\} .$ 

**Theorem 6.6.** Let  $r \ge 1$  and  $a_r \ge 1$ . We have  $\operatorname{Stab}(T_X(-\log D_{v_r})) \ne \emptyset$  if and only if  $\operatorname{sStab}(T_X(-\log D_{v_r})) \ne \emptyset$  if and only if  $a_r = 1$  and  $a_{r-1} = 0$ .

If  $a_r = 1$  and  $a_{r-1} = 0$ , then the logarithmic tangent bundle  $T_X(-\log D_{v_r})$  is stable (resp. semi-stable) with respect to  $\pi^* \mathcal{O}_{\mathbb{P}^s}(\nu) \otimes \mathcal{O}_X(1)$  if and only if  $0 < \nu < \nu_0$ (resp.  $0 < \nu \leq \nu_0$ ) where  $\nu_0$  is the unique positive root of

$$\mathsf{P}_{0}(x) = \sum_{k=0}^{s-1} \binom{s+r-1}{k} x^{k} - s \binom{s+r-1}{s} x^{s} .$$

*Proof.* Let  $\mathscr{E} = T_X(-\log D_{v_r})$  and  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ . We have

$$(r+s)\mu_L(\mathscr{E}) = (s+1)W + V_0 + V_1 + \ldots + V_{r-1}$$
.

• If  $a_r \ge 2$ , by using the first point of Lemma 5.12 and the fact that  $V_i \le V_0$ , we get :

$$(r+s)[\mathsf{V}_0 - \mu_L(\mathscr{E})] = (s\mathsf{V}_0 - (s+1)\mathsf{W}) + r\mathsf{V}_0 - (\mathsf{V}_0 + \ldots + \mathsf{V}_{r-1}) \ge s\mathsf{V}_r$$
.

By Proposition 5.10,  $T_X(-\log D_{v_r})$  is not semi-stable with respect to L.

• We assume that  $r \geq 2$  and  $a_{r-1} = a_r = 1$ . Like  $V_{r-1} = V_r$ , we have

$$(r+s)[\mathsf{V}_0 - \mu_L(\mathscr{E})] = (s+1)[\mathsf{V}_0 - \mathsf{W}] - \mathsf{V}_{r-1} + [(r-1)\mathsf{V}_0 - (\mathsf{V}_0 + \ldots + \mathsf{V}_{r-2})]$$
  

$$\geq (s+1)\mathsf{V}_r - \mathsf{V}_{r-1} \quad \text{because } \mathsf{V}_0 - \mathsf{W} \geq \mathsf{V}_r$$
  

$$\geq s\mathsf{V}_r$$

By Proposition 5.10,  $T_X(-\log D_{v_r})$  is not semi-stable with respect to L.

• Let  $r \ge 1$ . We now assume that  $a_{r-1} = 0$  and  $a_r = 1$ . By using the expressions of Section 5.2, we have  $V_0 = \ldots = V_{r-1} = V$  where

$$\mathsf{V} = \sum_{k=0}^{s} \binom{s+r-1}{k} \nu^{k} \quad \text{and} \quad \mathsf{W} = \sum_{k=0}^{s-1} \binom{s+r-1}{k} \nu^{k} .$$

The points 4 and 5 of Proposition 5.10 are not verified in this case. To check the stability of  $\mathscr E$  it is enough to compare

$$\mu_L(\mathscr{E}) = \frac{1}{r+s} (r\mathsf{V} + (s+1)\mathsf{W})$$

with  $\max(V, W)$ . We have

$$(r+s)[\mu_L(\mathscr{E}) - \mathsf{W}] = r\mathsf{V} - (r-1)\mathsf{W} > 0$$
 because  $\mathsf{W} < \mathsf{V}$ 

and

$$\begin{split} (r+s)[\mu_L(\mathcal{E})-\mathsf{V}] =& (s+1)\mathsf{W}-s\mathsf{V} \\ =& \sum_{k=0}^{s-1} \binom{s+r-1}{k} \nu^k - s\binom{s+r-1}{s} \nu^s = \mathsf{P}_0(\nu) \ . \end{split}$$

By the sign rule of Descartes (see Theorem 6.1), the polynomial  $\mathsf{P}_0$  have a unique positive root  $\nu_0$ . If  $\nu > 0$ , then  $\mathsf{P}_0(\nu) > 0$  (resp.  $\mathsf{P}_0(\nu) \ge 0$ ) if and only if  $\nu < \nu_0$  (resp.  $\nu \le \nu_0$ ). Thus,  $T_X(-\log D_{\nu_r})$  is stable (resp. semi-stable) with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $0 < \nu < \nu_0$  (resp.  $0 < \nu \le \nu_0$ ).  $\Box$ 

**Corollary 6.7.** We assume that  $r \ge 1$  and  $(a_1, \ldots, a_r) \ne (0, \ldots, 0)$ . Let  $j \in \{0, \ldots, s\}$  and  $D = D_{v_r} + D_{w_j}$ . For any  $L \in \text{Amp}(X)$ , the logarithmic tangent bundle  $T_X(-\log D)$  is not semi-stable with respect to L.

*Proof.* If  $\mathscr{E} = T_X(-\log D)$ , we have  $V_0 > \mu_L(\mathscr{E})$ . By Proposition 5.10,  $T_X(-\log D)$  is not semi-stable with respect to L.

6.3. The case where  $\operatorname{Cone}(v_0) \in \Delta$ . In this part we study the stability of the logarithmic tangent bundle  $T_X(-\log D)$  when  $r \geq 2$  and

 $D \in \{D_{v_0}\} \cup \{D_{v_0} + D_{w_j} : 0 \le j \le s\} \cup \{D_{v_0} + D_{v_i} : 2 \le i \le r\}.$ 

The last case  $D = D_{v_0} + D_{v_1}$  will be study in the Section 6.4. We recall that  $a_0 = 0$ . By Lemma 6.3 and Theorem 3.9, for all

$$D \in \{D_{v_0} + D_{w_i} : 0 \le j \le s\} \cup \{D_{v_0} + D_{v_i} : 2 \le i \le r\}$$

 $T_X(-\log D)$  is decomposable. If  $D = D_{v_0}, r \ge 2$  and  $a_1 = 0$ , then  $\mathscr{E} = T_X(-\log D)$  is decomposable and  $\mathscr{E} = \mathscr{E}_G \oplus \mathscr{E}_F$  where  $G = \operatorname{Span}(w_0, w_1, \ldots, w_s, v_2, \ldots, v_r)$  and  $F = \operatorname{Span}(v_1)$ . By Lemma 5.9, if  $a_1 > 0$ , then  $T_X(-\log D_{v_0})$  cannot be written as a sum of two equivariant vector bundles.

A. NAPAME

We now study the semi-stability of these bundles.

**Lemma 6.8.** Let  $r \geq 2$ ,  $(a_1, \ldots, a_r) \neq (0, \ldots, 0)$  such that  $a_1 < a_r$ ,  $i \in \{2, \ldots, r\}$  and  $j \in \{0, \ldots, s\}$ . We set  $\mathscr{E} = T_X(-\log D_{v_0})$ ,  $\mathscr{F} = T_X(-\log(D_{v_0} + D_{v_i}))$  and  $\mathscr{G} = T_X(-\log(D_{v_0} + D_{w_j}))$ . For any  $L \in \operatorname{Amp}(X)$ , the vector bundles  $\mathscr{E}$ ,  $\mathscr{F}$  and  $\mathscr{G}$  are not semi-stables with respect to L.

*Proof.* We have  $\mu_L(\mathscr{E}) > \mu_L(\mathscr{F})$  and  $\mu_L(\mathscr{E}) > \mu_L(\mathscr{G})$ . We will show that  $V_1 > \mu_L(\mathscr{E})$ . By Lemma 5.12, we have  $V_1 - W \ge V_r$ . Therefore

$$(r+s)(V_1 - \mu_L(\mathscr{E})) = (r+s)V_1 - (V_1 + \ldots + V_r) - (s+1)W$$
  
=  $(s+1)[V_1 - W] - V_r + [(r-1)V_1 - (V_1 + \ldots + V_{r-1})]$   
 $\ge (s+1)[V_1 - W] - V_r$   
 $\ge sV_r$ 

By Proposition 5.10,  $\mathscr{E}$ ,  $\mathscr{F}$  and  $\mathscr{G}$  are not semi-stables with respect to L.

Let  $a \in \mathbb{N}^*$ . We now study what happen in Lemma 6.8 when  $a_1 = \ldots = a_r = a$ . 6.3.1. We assume that r = 1.

 $\square$ 

**Theorem 6.9.** We assume that  $X = \mathbb{P}(\mathscr{O}_{\mathbb{P}^s} \oplus \mathscr{O}_{\mathbb{P}^s}(a))$ . Let  $\mathsf{P}_1$  and  $\mathsf{Q}$  the polynomials defined by

$$\mathsf{P}_{1}(x) = (s+1) \sum_{k=0}^{s-1} \binom{s}{k} a^{s-k-1} x^{k} - s \, x^{s} \quad and \quad \mathsf{Q}(x) = x^{s} - \sum_{k=0}^{s-1} \binom{s}{k} a^{s-k-1} x^{k} \, .$$

We have :

- (1)  $T_X(-\log D_{\nu_0})$  is stable (resp. semi-stable) with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $0 < \nu < \nu_1$  (resp.  $0 < \nu \le \nu_1$ ) where  $\nu_1$  is the unique positive root of  $\mathsf{P}_1$ .
- (2) If  $j \in \{0, ..., s\}$ , then  $\operatorname{Stab}(T_X(-\log(D_{v_0} + D_{w_j}))) = \emptyset$ ;  $T_X(-\log(D_{v_0} + D_{w_j}))$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \nu_3$  where  $\nu_3$  is the unique positive root of  $\mathbb{Q}$ .
- (3)  $\varnothing = \operatorname{Stab}(T_X(-\log(D_{v_0} + D_{v_1}))) \subseteq \operatorname{sStab}(T_X(-\log(D_{v_0} + D_{v_1}))) = \operatorname{Amp}(X).$

*Proof.* By the sign rule of Descartes (cf. Theorem 6.1),  $\mathsf{P}_1$  and  $\mathsf{Q}$  have respectively one positive root. Let  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ , by using the expressions of Section 5.2, we have :

$$\mathsf{V}_1 = \nu^s$$
 and  $\mathsf{W} = \sum_{k=0}^{s-1} \binom{s}{k} a^{s-k-1} \nu^k$ .

Let  $\mathscr{E} = T_X(-\log D_{v_0})$ . By Proposition 5.10, to check the stability of  $\mathscr{E}$ , it is enough to compare

$$\mu_L(\mathscr{E}) = \frac{\mathsf{V}_1 + (1+s)\mathsf{W}}{1+s}$$

with  $\max(V_1, W)$ . We don't use the points 4 and 5 because the hypothesis are not verifying. We have

$$\mu_L(\mathscr{E}) > \mathsf{W}$$
 and  $(1+s)(\mu_L(\mathscr{E}) - \mathsf{V}_1) = \mathsf{P}_1(\nu)$ .

Thus,  $\mathscr{E}$  is stable (resp. semi-stable) with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $0 < \nu < \nu_1$  (resp.  $0 < \nu \leq \nu_1$ ).

Let  $\mathscr{F} = T_X(-\log(D_{v_0} + D_{w_i}))$ . By Proposition 5.10, we compare

$$\mu_L(\mathscr{F}) = \frac{\mathsf{V}_1 + s\mathsf{W}}{1+s}$$

with  $\max(V_1, W)$ . We have

$$(1+s)(\mu_L(\mathscr{F}) - \mathsf{W}) = \mathsf{Q}(\nu) \text{ and } (1+s)(\mu_L(\mathscr{F}) - \mathsf{V}_1) = -s \,\mathsf{Q}(\nu)$$

Thus, for any  $\nu > 0$ ,  $\mathscr{F}$  is not stable with respect to L. The vector bundle  $\mathscr{F}$  is semi-stable with respect to L if and only if  $\nu = \nu_3$ .

Let  $\mathscr{G} = T_X(-\log(D_{v_0} + D_{v_1}))$ . We have  $\mu_L(\mathscr{G}) = \mathsf{W}$ . By Proposition 5.10,  $\mathscr{G}$  is semi-stable not stable with respect to L.

6.3.2. We assume that  $r \geq 2$ . We assume that  $a_1 = \ldots = a_r = a$  with  $a \in \mathbb{N}^*$ .

Lemma 6.10. We have

$$\operatorname{card}\{(\alpha_1,\ldots,\alpha_p)\in\mathbb{N}^p:\alpha_1+\ldots+\alpha_p=m\}=\binom{m+p-1}{m}$$

By Lemma 6.10, for all  $k \in \{0, ..., s - 1\}$ , we have

$$W_k = \sum_{d_1 + \dots + d_r = s - k - 1} a_1^{d_1} \cdots a_r^{d_r} = \binom{s - k + r - 2}{s - k - 1} a^{s - k - 1}$$

and

$$\mathsf{V}_{1k} = \sum_{d_2 + \dots + d_r = s-k} a_2^{d_2} \cdots a_r^{d_r} = \binom{s-k+r-2}{s-k} a^{s-k}$$

We recall that  $V_{1s} = 1$ . By using the equality  $\binom{n}{p-1} = \frac{p}{n-p+1} \binom{n}{p}$ , for all  $k \in \{0, \dots, s-1\}$ ,

$$\mathsf{W}_{k} = \binom{s-k+r-2}{s-k-1} a^{s-k-1} = \frac{s-k}{r-1} \binom{s-k+r-2}{s-k} a^{s-k-1} = \frac{s-k}{a(r-1)} \mathsf{V}_{1k} \; .$$

Let  $\mathsf{P}_1$  and  $\mathsf{Q}$  the polynomials defined by :

$$\mathsf{P}_{1}(x) = \sum_{k=0}^{s-1} \left[ \left( -s + \frac{(s-k)(s+1)}{a(r-1)} \right) \binom{s+r-1}{k} \mathsf{V}_{1k} \right] x^{k} - s\binom{s+r-1}{s} x^{s}$$

$$\mathsf{Q}(x) = \sum_{k=0}^{s-1} \left[ \left( r - \frac{s-k}{a} \right) \binom{s+r-1}{k} \mathsf{V}_{1k} \right] x^{k} + r\binom{s+r-1}{s} x^{s}$$

**Theorem 6.11.** Let  $r \geq 2$  and  $X = \mathbb{P}(\mathscr{O}_{\mathbb{P}^s} \oplus \mathscr{O}_{\mathbb{P}^s}(a_1) \oplus \ldots \oplus \mathscr{O}_{\mathbb{P}^s}(a_r))$  with  $a_1 = \ldots = a_r = a$  where  $a \in \mathbb{N}^*$ . We set  $\mathscr{E} = T_X(-\log D_{v_0})$ .

- (1) If  $a < \frac{s}{r}$ , then  $\mathscr{E}$  is stable (resp. semi-stable) with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu_3 < \nu < \nu_1$  (resp.  $\nu_3 \le \nu \le \nu_1$ ) where  $\nu_1$  and  $\nu_3$  are respectively the positive roots of  $\mathsf{P}_1$  and  $\mathsf{Q}$ .
- respectively the positive roots of  $\mathsf{P}_1$  and  $\mathsf{Q}$ . (2) If  $\frac{s}{r} \leq a < \frac{s+1}{r-1}$ , then  $\mathscr{E}$  is stable (resp. semi-stable) with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $0 < \nu < \nu_1$  (resp.  $0 < \nu \leq \nu_1$ ) where  $\nu_1$  is the unique positive root of  $\mathsf{P}_1$ .
- is the unique positive root of  $\mathsf{P}_1$ . (3) If  $a \ge \frac{s+1}{r-1}$ , then for any  $L \in \operatorname{Amp}(X)$ ,  $\mathscr{E}$  is not semi-stable with respect to L.

Before giving the proof of this theorem, we will explain the condition which ensure the existence of positive roots on  $P_1$  and Q. We will use the Descartes sign rule.

We write 
$$\mathsf{P}_1$$
 as  $\mathsf{P}_1(x) = \sum_{k=0}^{s} \alpha_k x^k$ . We have  $\alpha_s < 0$ . If  $-s + \frac{(s-k)(s+1)}{a(r-1)} > 0$ ,

A. NAPAME

then  $s-k > \frac{a s(r-1)}{s+1}$ , i.e.  $k < \left(1 - \frac{a(r-1)}{s+1}\right) s$ . If  $\frac{a(r-1)}{s+1} \ge 1$ , then for any  $x \ge 0$ ,  $\mathsf{P}_1(x) < 0$ . If  $a < \frac{s+1}{r-1}$ , then for all  $k < \left(1 - \frac{a(r-1)}{s+1}\right) s$ , we have  $\alpha_k > 0$  and for all  $k \ge \left(1 - \frac{a(r-1)}{s+1}\right) s$ ,  $\alpha_k \le 0$ . Hence  $\mathsf{P}_1$  has only one positive root  $\nu_1$ . We write  $\mathsf{Q}(x) = \sum_{k=0}^{s} \beta_k x^k$ . We have  $\beta_s > 0$ . If  $r - \frac{s-k}{a} < 0$ , then r a - (s-k) < 0, i.e. k < s - r a. If  $a \ge \frac{s}{r}$ , then for any  $x \ge 0$ ,  $\mathsf{Q}(x) > 0$ . If  $a < \frac{s}{r}$ , then for all k < s - r a, we have  $\beta_k < 0$  and for all  $k \ge s - r a$ ,  $\beta_k \ge 0$ . Thus,  $\mathsf{Q}$  has only one positive root  $\nu_3$ .

**Lemma 6.12.** If  $a < \frac{s}{r}$ , then  $\nu_3 < \nu_1$ .

*Proof.* If  $a < \frac{s}{r}$ , then  $a < \frac{s+1}{r-1}$ . By the above analysis,  $\nu_1$  and  $\nu_3$  exist. For  $x \ge 0$ , we have  $\mathsf{P}_1(x) > 0$  if and only if  $0 \le x < \nu_1$ . Like

$$\frac{\mathsf{P}_1(x)}{-s} - \frac{\mathsf{Q}(x)}{r} = \sum_{k=0}^{s-1} \left[ \left( 1 - \frac{(s-k)(s+1)}{a\,s(r-1)} - 1 + \frac{s-k}{r\,a} \right) \binom{s+r-1}{k} \mathsf{V}_{1k} \right] x^k \\ = \frac{-(r+s)}{a\,s\,r(r-1)} \sum_{k=0}^{s-1} (s-k) \binom{s+r-1}{k} \mathsf{V}_{1k} x^k = \mathsf{P}(x)$$

we have  $\frac{\mathsf{P}_1(\nu_3)}{-s} - \frac{\mathsf{Q}(\nu_3)}{r} = \mathsf{P}(\nu_3) < 0$ . Hence,  $\mathsf{P}_1(\nu_3) > 0$ , i.e  $\nu_3 < \nu_1$ .

Now, we can prove Theorem 6.11.

**Proof of Theorem 6.11.** Like  $a_1 = \ldots = a_r$ , we have  $V_1 = \ldots = V_r$ . Therefore

$$\mu_L(\mathscr{E}) = \frac{r \,\mathsf{V}_1 + (s+1)\mathsf{W}}{r+s}$$

In this case the point 4 and 5 are not verified. To check the stability of  $\mathscr{E} = T_X(-\log D_{v_0})$ , it is enough to compare  $\mu_L(\mathscr{E})$  with  $\max(\mathsf{V}_1,\mathsf{W})$ . We have

$$\begin{split} (r+s)(\mu_L(\mathscr{E}) - \mathsf{V}_1) &= -s\,\mathsf{V}_1 + (s+1)\mathsf{W} \\ &= -s\sum_{k=0}^s \binom{s+r-1}{k}\mathsf{V}_{1k}\,\nu^k + (s+1)\sum_{k=0}^{s-1} \binom{s+r-1}{k}\mathsf{W}_k\,\nu^k \\ &= \mathsf{P}_1(\nu) \end{split}$$

and

$$\begin{split} (r+s)(\mu_L(\mathscr{E}) - \mathsf{W}) &= r \,\mathsf{V}_1 - (r-1)\mathsf{W} \\ &= r \sum_{k=0}^s \binom{s+r-1}{k} \mathsf{V}_{1k} \,\nu^k - (r-1) \sum_{k=0}^{s-1} \binom{s+r-1}{k} \mathsf{W}_k \,\nu^k \\ &= \mathsf{Q}(\nu) \end{split}$$

We have :

i. If 
$$a \ge \frac{s+1}{r-1}$$
, then for any  $\nu > 0$ , we have  $\mathsf{P}_1(\nu) < 0$ .

- ii. If  $a < \frac{s+1}{r-1}$ , then  $\mathsf{P}_1(\nu) > 0$  (resp.  $\mathsf{P}_1(\nu) \ge 0$ ) if and only if  $0 < \nu < \nu_1$
- (resp.  $0 < \nu \leq \nu_1$ ). iii. If  $a \geq \frac{s}{r}$ , then for any  $\nu > 0$ , we have  $Q(\nu) > 0$ . iv. If  $a < \frac{s}{r}$ , then  $Q(\nu) > 0$  (resp.  $Q(\nu) \geq 0$ ) if and only if  $\nu > \nu_3$  (resp.  $\dots > \dots > \dots$ )

The point *i*. shows the third point of the theorem. By using the points *ii*. , *iv*. and the Lemma 6.12, we get the first point of theorem. The points *ii.* and *iii.* give the second point of the theorem. 

Let  $i \in \{1, \ldots, r\}$  and  $j \in \{0, \ldots, s\}$ . We now study the stability of  $\mathscr{F}_j =$  $T_X(-\log(D_{v_0} + D_{w_i}))$  and  $\mathscr{G}_i = T_X(-\log(D_{v_0} + D_{v_i}))$ . Let Q the polynomial defined by

$$\mathsf{Q}(x) = \sum_{k=0}^{s-1} \left[ \left( 1 - \frac{s-k}{a(r-1)} \right) \binom{s+r-1}{k} \mathsf{V}_{1k} \right] x^k + \binom{s+r-1}{s} x^s .$$

**Theorem 6.13.** We assume that  $r \geq 2$  and  $X = \mathbb{P}(\mathcal{O}_{\mathbb{P}^s} \oplus \mathcal{O}_{\mathbb{P}^s}(a_1) \oplus \ldots \oplus \mathcal{O}_{\mathbb{P}^s}(a_r))$ with  $a_1 = \ldots = a_r = a$  where  $a \in \mathbb{N}^*$ . Let  $i \in \{1, \ldots, r\}$  and  $j \in \{0, \ldots, s\}$ .

- (1) For any  $L \in \operatorname{Amp}(X)$ ,  $\mathscr{F}_j$  and  $\mathscr{G}_i$  are not stables with respect to L. (2) If  $a \geq \frac{s}{r-1}$ , then for any  $L \in \operatorname{Amp}(X)$ ,  $\mathscr{F}_j$  and  $\mathscr{G}_i$  are not semi-stables with respect to L.
- (3) If  $a < \frac{s}{r-1}$ , then  $\mathscr{F}_j$  and  $\mathscr{G}_i$  are semi-stables with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes$  $\mathscr{O}_X(1)$  if and only if  $\nu = \nu_3$  where  $\nu_3$  is the unique root of Q.

*Proof.* We first study the polynomial Q. We write  $Q(x) = \sum_{k=0}^{s} \alpha_k x^k$ . We have  $\alpha_s > 0 \, . \ \text{If} \ \ 1 - \frac{s-k}{a(r-1)} < 0 \ , \ \text{then} \ \ a(r-1) - (s-k) < 0 \ , \ \text{i.e} \ \ k < s - a(r-1) \, .$ If  $a \ge \frac{s}{r-1}$ , then for any  $x \ge 0$ , Q(x) > 0. If  $a < \frac{s}{r-1}$ , then for all k < s-a(r-1), we have  $\alpha_k < 0$  and for  $k \ge s-a(r-1)$ ,  $\alpha_k \geq 0$ . Thus, **Q** has a unique positive root  $\nu_3$ .

Like  $a_1 = \ldots = a_r$ , we have  $V_1 = \ldots = V_r$ . Thus,

$$\mu_L(\mathscr{F}_j) = \frac{r \operatorname{V}_1 + s \operatorname{W}}{r+s}$$
 and  $\mu_L(\mathscr{G}_i) = \frac{(r-1)\operatorname{V}_1 + (s+1)\operatorname{W}}{r+s}$ 

By using Proposition 5.10, to check the stability of  $\mathscr{F}_i$  (resp.  $\mathscr{G}_i$ ), it is enough to compare  $\mu_L(\mathscr{F}_j)$  (resp.  $\mu_L(\mathscr{G}_i)$ ) with  $\max(V_1, W)$ . We have

$$\begin{cases} (r+s)(\mu_L(\mathscr{F}_j) - \mathsf{V}_1) = s(\mathsf{W} - \mathsf{V}_1) = -s \, \mathsf{Q}(\nu) \\ (r+s)(\mu_L(\mathscr{F}_j) - \mathsf{W}) = r(\mathsf{V}_1 - \mathsf{W}) = r \, \mathsf{Q}(\nu) \end{cases}$$

and

$$\begin{cases} (r+s)(\mu_L(\mathscr{G}_i) - \mathsf{V}_1) = (s+1)(\mathsf{W} - \mathsf{V}_1) = -(s+1)\,\mathsf{Q}(\nu) \\ (r+s)(\mu_L(\mathscr{G}_i) - \mathsf{W}) = (r-1)(\mathsf{V}_1 - \mathsf{W}) = (r-1)\,\mathsf{Q}(\nu) \end{cases}$$

If  $a \geq \frac{s}{r-1}$ , then for any  $\nu > 0$ ,  $Q(\nu) > 0$ ; thus,  $\mu_L(\mathscr{F}_j) < \mathsf{V}_1$  and  $\mu_L(\mathscr{G}_i) < \mathsf{V}_1$ . Hence, for any  $\nu > 0$ ,  $\mathscr{F}_j$  and  $\mathscr{G}_i$  are not semi-stables with respect to L. We now assume that  $a < \frac{s}{r-1}$ . Let  $\nu_3$  the positive root of Q.

If  $\nu \neq \nu_3$ , by the above equalities,  $\mathscr{F}_i$  and  $\mathscr{G}_i$  are not semi-stables with respect to

L. If  $\nu = \nu_3$ , the logarithmic tangent bundles  $\mathscr{F}_j$  and  $\mathscr{G}_i$  are semi-stables but not stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$ .

6.4. The case  $D = D_{v_0} + D_{v_1}$ . In this part, we assume that  $D = D_{v_0} + D_{v_1}$  and  $\mathscr{E} = T_X(-\log D)$ . In the beginning of Section 6.3, we see that  $\mathscr{E}$  is decomposable. In this part, we will study the stability of  $\mathscr{E}$  when  $r \ge 2$  and  $a_1 < a_r$ . The stability of  $\mathscr{E}$  when r = 1 was treated in Theorem 6.9. When  $r \ge 2$ , in Theorem 6.13, we study the stability of  $\mathscr{E}$  when  $a_1 = \ldots = a_r$  with  $a_1 \in \mathbb{N}^*$ .

**Proposition 6.14.** If  $a_1 = 0$ , then for any  $L \in Amp(X)$ ,  $\mathscr{E}$  is not semi-stable with respect to L.

*Proof.* We have  $\mu_L(\mathscr{E}) = \frac{1}{r+s} \left( (s+1)\mathsf{W} + \mathsf{V}_2 + \ldots + \mathsf{V}_r \right)$ . Like  $\operatorname{card}\{2, \ldots, r\} = r-1$ , by using the point 4 of Proposition 5.10 with  $I' = \{2, \ldots, r\}$ , we have

$$\frac{1}{r+s-1}\left(\sum_{i\in I'} \mathsf{V}_i + (s+1)\mathsf{W}\right) = \frac{1}{r+s-1}\left(\mathsf{V}_2 + \ldots + \mathsf{V}_r + (s+1)\mathsf{W}\right) \ .$$

Thus,  $T_X(-\log(D_{v_0} + D_{v_1}))$  is not semi-stable with respect to L.

We also have :

**Lemma 6.15.** Let  $r \ge 3$ ,  $(a_1, \ldots, a_r) \ne (0, \ldots, 0)$  such that  $a_2 < a_r$ . For any  $L \in \operatorname{Amp}(X)$ , the vector bundle  $\mathscr{E}$  is not semi-stable with respect to L.

*Proof.* We have  $\mu_L(\mathscr{E}) = \frac{1}{r+s}(\mathsf{V}_2 + \ldots + \mathsf{V}_r + (s+1)\mathsf{W})$ . We will show that  $\mathsf{V}_2 > \mu_L(\mathscr{E})$ . By Lemma 5.12, we have  $\mathsf{V}_2 - \mathsf{W} \ge \mathsf{V}_r$ , thus

$$(r+s)(V_2 - \mu_L(\mathscr{E}) = (r+s)V_2 - (V_2 + \ldots + V_r) - (s+1)W$$
  
= (s+1)[V\_2 - W] + [(r-1)V\_2 - (V\_2 + \ldots + V\_r)]  
\ge (s+1)[V\_2 - W]  
\ge (s+1)V\_r

Hence, by Proposition 5.10,  $\mathscr{E}$  is not semi-stable with respect to L.

We now assume that  $0 < a_1 < a_2 = \ldots = a_r$ . For the stability of  $\mathscr{E}$ , we see that: the point 4 and 5 of Proposition 5.10 are not verified here. To check the stability of  $\mathscr{E}$ , it is enough to compare  $\mu_L(\mathscr{E})$  with  $\max(V_2, W)$ . We have

$$\mu_L(\mathscr{E}) = \frac{(r-1)\mathsf{V}_2 + (s+1)\mathsf{W}}{r+s}$$

and

(14) 
$$\begin{cases} (r+s)[\mu_L(\mathscr{E}) - \mathsf{V}_2] = (s+1)(\mathsf{W} - \mathsf{V}_2) \\ (r+s)[\mu_L(\mathscr{E}) - \mathsf{W}] = -(r-1)(\mathsf{W} - \mathsf{V}_2) \end{cases}$$

We have :

**Proposition 6.16.** Let  $r \geq 2$ . If  $0 < a_1 < a_2 = \ldots = a_r$ , then  $\operatorname{Stab}(\mathscr{E}) = \varnothing$ .

The vector bundle  $\mathscr{E} = T_X(-\log(D_{v_0} + D_{v_1}))$  is semi-stable with respect to  $L = \pi^* \mathscr{O}_{\mathbb{P}^s}(\nu_3) \otimes \mathscr{O}_X(1)$  if and only if  $\nu_3$  is a positive root of the polynomial  $\mathsf{Q}(\nu) = \mathsf{W} - \mathsf{V}_2$  (W and  $\mathsf{V}_2$  depend on  $\nu$ ).

6.4.1. We assume that r = 2 and  $0 < a_1 < a_2$ .

**Proposition 6.17.** Let r = 2 and  $0 < a_1 < a_2$ . We define  $\delta = \frac{\ln(1 + a_2 - a_1)}{\ln(a_2) - \ln(a_1)}$ and Q the polynomial

$$\mathsf{Q}(x) = \sum_{k=0}^{s-1} \left[ \frac{a_1^{s-k}}{a_2 - a_1} \left( \left( \frac{a_2}{a_1} \right)^{s-k} - 1 - a_2 + a_1 \right) \binom{s+1}{k} \right] x^k - (s+1)x^s \; .$$

We have :

- (1) If  $s \leq \delta$ , then  $\operatorname{sStab}(T_X(-\log(D_{v_0} + D_{v_1}))) = \emptyset$ ;
- (2) If  $s > \delta$ , then  $T_X(-\log(D_{v_0}+D_{v_1}))$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \nu_3$  where  $\nu_3$  is the positive root of  $\mathbb{Q}$ .

Proof. We have

$$V_{2} = \sum_{k=0}^{s} {\binom{s+1}{k}} a_{1}^{s-k} \nu^{k}$$
$$W = \sum_{k=0}^{s-1} {\binom{s+1}{k}} \left(\sum_{d_{1}+d_{2}=s-k-1} a_{1}^{d_{1}} a_{2}^{d_{2}}\right) \nu^{k} = \sum_{k=0}^{s-1} \frac{a_{2}^{s-k} - a_{1}^{s-k}}{a_{2} - a_{1}} {\binom{s+1}{k}} \nu^{k}$$

hence,

$$\mathsf{W}-\mathsf{V}_{2} = \sum_{k=0}^{s-1} \left[ \frac{a_{1}^{s-k}}{a_{2}-a_{1}} \left( \left( \frac{a_{2}}{a_{1}} \right)^{s-k} - 1 - a_{2} + a_{1} \right) \binom{s+1}{k} \right] \nu^{k} - (s+1)\nu^{s} = \mathsf{Q}(\nu) \; .$$

We write  $Q(x) = \sum_{k=0} \alpha_k x^k$ . We have  $\alpha_s < 0$ . The inequality

$$\left(\frac{a_2}{a_1}\right)^{s-k} - 1 - a_2 + a_1 > 0$$

gives  $k < s - \frac{\ln(1 + a_2 - a_1)}{\ln(a_2) - \ln(a_1)} = s - \delta$ . If  $s \le \delta$ , then for any  $x \ge 0$ , Q(x) < 0. If  $s > \delta$ , then by the Descartes rule, Q has a unique positive root  $\nu_3$ .

Hence, if  $s > \delta$ , then  $\mathscr{E}$  is semi-stable with respect to  $\pi^*\mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \nu_3$ .

6.4.2. We assume that  $r \ge 3$  and  $a_1 < a_r$ . Let  $a, b \in \mathbb{N}^*$  such that a < b. We assume that  $a_1 = a$  and  $a_2 = \ldots = a_r = b$ . By Lemma 6.10, for all  $k \in \{0, \ldots, s-1\}$ , we have :

$$W_{k} = \sum_{\substack{d_{1}+\ldots+d_{r}\\ = s-k-1}} a_{1}^{d_{1}} \cdots a_{r}^{d_{r}} = \sum_{j=0}^{s-k-1} a^{s-k-1-j} \left(\sum_{\substack{d_{2}+\ldots+d_{r}=j\\ j}} b^{j}\right)$$
$$= \sum_{j=0}^{s-k-1} {j + r - 2 \choose j} b^{j} a^{s-k-1-j}$$

and

$$V_{2k} = V_{rk} = \sum_{\substack{d_1 + \dots + d_{r-1} \\ = s-k}} a_1^{d_1} \cdots a_{r-1}^{d_{r-1}} = \sum_{j=0}^{s-k} a^{s-k-j} \left( \sum_{\substack{d_2 + \dots + d_{r-1} = j \\ d_2 + \dots + d_{r-1} = j}} b^j \right)$$
$$= \sum_{j=0}^{s-k} {j + r - 3 \choose j} b^j a^{s-k-j} .$$

For  $k \in \{0, ..., s-1\}$ , we have  $(s-k) \in \{1, ..., s\}$ . For  $p \in \{1, ..., s\}$ , we set  $\alpha_p = W_{s-p} - V_{2, s-p}$ ; we have

$$\alpha_p = \sum_{j=0}^{p-1} \binom{j+r-2}{j} b^j a^{p-1-j} - \sum_{j=0}^p \binom{j+r-3}{j} b^j a^{p-j} .$$

Let  $Q_s$  be the polynomial defined by

$$Q_{s}(x) = \sum_{k=0}^{s-1} {\binom{s+r-1}{k}} \alpha_{s-k} x^{k} - {\binom{s+r-1}{s}} x^{s}$$

We have  $W - V_2 = Q_s(\nu)$ . We now search a condition on s which ensure the existence of positive root on Q. By using the identity  $\binom{n}{p-1} = \frac{p}{n-p+1}\binom{n}{p}$ , we have

$$\alpha_p = \sum_{j=0}^{p-1} \frac{j+1}{r-2} \binom{j+r-2}{j+1} b^j a^{p-1-j} - \sum_{j=1}^p \binom{j+r-3}{j} b^j a^{p-j} - a^p$$
$$= \sum_{j=0}^{p-1} \left[ \left( \frac{j+1}{r-2} - b \right) \binom{j+r-2}{j+1} b^j a^{p-1-j} \right] - a^p .$$

If  $1 \le p \le b(r-2)$ , then for all  $j \in \{0, \ldots, p-1\}$ , we have

$$\frac{j+1}{r-2} - b \le \frac{p}{r-2} - b = \frac{p - b(r-2)}{r-2} \le 0 ;$$

thus  $\alpha_p < 0$ . Hence, if  $\alpha_p > 0$ , then we must have p > b(r-2). If there is p > b(r-2) such that  $\alpha_p > 0$ , then for any  $q \ge p$ , we have  $\alpha_q > 0$ ; this follows from these equalities.

$$\alpha_{p+1} = \sum_{j=0}^{p-1} \left[ \left( \frac{j+1}{r-2} - b \right) \left( \frac{j+r-2}{j+1} \right) b^j a^{p-j} \right] - a^{p+1} + \left( \frac{p+1}{r-2} - b \right) \left( \frac{p+r-2}{p+1} \right) b^p$$
$$= a \,\alpha_p + \left( \frac{p+1}{r-2} - b \right) \left( \frac{p+r-2}{p+1} \right) b^p$$

We denote by |x| the floor of  $x \in \mathbb{R}$ .

**Lemma 6.18.** Let m = b(r-2). There is a unique integer  $\delta_r \in [m+1; \lfloor 3.2 m \rfloor + 1]$  such that : if  $p \leq \delta_r$ , then  $\alpha_p \leq 0$  and if  $p > \delta_r$ , then  $\alpha_p > 0$ .

Let  $\delta_r$  be the integer given in Lemma 6.18. If  $s \leq \delta_r$ , then all coefficients of  $Q_s$  are negative; thus, for any x > 0,  $Q_s(x) < 0$ . If  $s > \delta_r$ , then by the Descartes rule (see Theorem 6.1),  $Q_s$  has a only one positive root  $\nu_3$ . We deduce :

**Proposition 6.19.** Let  $r \geq 3$  and  $a, b \in \mathbb{N}^*$  such that a < b and  $a_1 = a$ ,  $a_2 = \ldots = a_r = b$ .

- (1) If  $s \leq \delta_r$ , then  $\operatorname{sStab}(T_X(-\log(D_{v_0} + D_{v_1}))) = \emptyset$ ;
- (2) If  $s > \delta_r$ , then  $T_X(-\log(D_{v_0}+D_{v_1}))$  is semi-stable with respect to  $\pi^* \mathscr{O}_{\mathbb{P}^s}(\nu) \otimes \mathscr{O}_X(1)$  if and only if  $\nu = \nu_3$ .

We now give the proof of Lemma 6.18.

Proof. We have 
$$\alpha_p = \sum_{j=0}^{m-1} \left[ \left( \frac{j+1}{r-2} - b \right) \binom{j+r-2}{j+1} b^j a^{p-1-j} \right] - a^p + \beta_p$$
 where  

$$\beta_p = \sum_{l=m}^{p-1} \left[ \left( \frac{l+1-(r-2)b}{r-2} \right) \binom{l+r-2}{l+1} b^l a^{p-1-l} \right]$$

$$= \sum_{l=0}^{p-1-m} \left[ \frac{l+1}{r-2} \binom{l+m+r-2}{l+m+1} b^{l+m} a^{p-1-(l+m)} \right]$$

The goal of this proof is to find an integer p such that  $\alpha_p > 0$ . We will search an integer p such that

(15) 
$$\beta_p \ge a^p + \sum_{j=0}^{m-1} b \binom{j+r-2}{j+1} b^j a^{p-1-j} .$$

We have  $\binom{l+m+r-2}{l+m+1} = \binom{l+m+r-2}{r-3} = \frac{r-2}{l+m+1} \binom{l+m+r-2}{r-2}$ . From the equality

$$\sum_{j=n}^{r} \binom{j}{n} = \binom{r+1}{n+1} \text{ we have } \binom{r+1}{n+1} = 1 + \sum_{j=0}^{r-n-1} \binom{j+n+1}{n};$$

Hence,

$$\binom{l+m+r-2}{r-2} = 1 + \sum_{j=0}^{l+m-1} \binom{j+r-2}{r-3} = 1 + \sum_{j=0}^{l+m-1} \binom{j+r-2}{j+1} \\ \ge 1 + \sum_{j=0}^{m-1} \binom{j+r-2}{j+1} .$$

Thus,

$$\beta_p \ge \left(1 + \sum_{j=0}^{m-1} \binom{j+r-2}{j+1}\right) \sum_{l=0}^{p-1-m} \frac{l+1}{l+1+m} b^{l+m} a^{p-1-(l+m)}$$
$$\ge b^m a^{p-1-m} \left(1 + \sum_{j=0}^{m-1} \binom{j+r-2}{j+1}\right) \sum_{l=0}^{p-1-m} \frac{l+1}{l+1+m} \left(\frac{b}{a}\right)^l.$$

We have

$$\sum_{l=0}^{p-1-m} \frac{l+1}{l+1+m} \ge \sum_{l=0}^{p-1-m} \int_{l}^{l+1} \frac{x \, dx}{x+m} = \int_{0}^{p-m} \frac{x}{x+m} dx$$
$$= \left[\frac{x}{1} - m\ln(x+m)\right]_{0}^{p-m}$$
$$= p - m - m\ln\left(\frac{p}{m}\right)$$

We set  $k = \frac{p}{m}$ . If  $k \ge \frac{17}{5} = 3.2$ , then  $(k - 1 - \ln(k)) > 1$ . If we set  $p = \lfloor 3.2 m \rfloor + 1$ , we get

$$\sum_{l=0}^{p-1-m} \frac{l+1}{l+1+m} \left(\frac{b}{a}\right)^l \ge \sum_{l=0}^{p-1-m} \frac{l+1}{l+1+m} \ge m \ge b \; .$$

For  $j \in \{0, ..., m-1\}$ , we have  $b^m a^{p-1-m} \ge b^j a^{p-1-j}$ . If  $p = \lfloor 3.2 m \rfloor + 1$ , we get

$$\begin{split} \beta_p &\geq b \, b^m \, a^{p-1-m} \left( 1 + \sum_{j=0}^{m-1} \binom{j+r-2}{j+1} \right) \\ &\geq b^{m+1} \, a^{p-1-m} + \sum_{j=0}^{m-1} b \binom{j+r-2}{j+1} b^j \, a^{p-1-j} \; ; \end{split}$$

this proves the inequality (15). Thus,  $\alpha_p > 0$  for  $p = \lfloor 3.2 \, m \rfloor + 1$ . Hence, we deduce the existence of the integer  $\delta_r$  in the interval  $[m+1; \lfloor 3.2 \, m \rfloor + 1]$ .

#### References

[BB13]	Robert J. BERMAN, BO BERNDTSSON, Real Monge-Ampère equations and Kähler-Ricci solitons on toric log Fano varieties. Annales de la faculté des sciences de Toulouse
	Mathématiques 22.4 (2013): 649-711. http://afst.cedram.org/item?id=AFST_2013_ 6.22.4.649_0
[Ber16]	Robert J. BERMAN, K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics, Invent. Math. 203 (2016), no. 3, p. 973–1025
[BG16]	Robert J. BERMAN, Henri GUENANCIA, Kähler-Einstein metrics on stable varieties and log canonical pairs. ArXiv preprint 1304.2087v2, https://arxiv.org/abs/1304. 2087v2, 2016.
[CLS11]	D. COX, J. LITTLE, SCHENCK, <i>Toric varieties</i> . AMS Graduate Studies in Mathematics volume : 124 ; 2011.
[CT20]	A. CLARKE, C. TIPLER, Equivariant stable sheaves and toric GIT. ArXiv preprint 2001.01500, https://arxiv.org/abs/2001.01500v1, 2020.
[Dan78]	V. I. DANILOV, The geometry of toric varieties. Uspekhi Mat. Nauk, 33(2(200)):85–134, 247, 1978.
[DDK20]	T. DASGUPTA, A. DEY, B. KHAN, Stabiliy of equivariant vector bundle over toric varieties. Documenta Mathematica 25 : 1787–1833, 2020.
[EV92]	H. Esnault, E. Viehweg, <i>Lectures on Vanishing Theorems</i> . DMV Seminar, vol. 20, Birkhäuser Verlag, Basel, 1992.
[Ful93]	William FULTON, <i>Introduction to toric varieties</i> . Princeton University Press, Princeton New Jersey, 1993.
[GH93]	P. GRIFFITHS, J. HARRIS, <i>Principles of algebraic geometry</i> . Wiley Classics Library, 1994.
[GS01]	GE. DETHLOFF, S. SHIN-YI LU, Logarithmic jet bundles and applications. Osaka J. Math. 38 (2001), 185-237.
[GT21]	Henri GUENANCIA, Behrouz TAJI, Orbifold Stability and Miyaoka-Yau Inequality for minimal pairs. ArXiv preprint 1611.05981v2, https://arxiv.org/abs/1611.05981v2, 2021.
[Gue16]	Henri GUENANCIA, Semistability of the tangent sheaf of singular varieties. Alebraic Geometry 3 (5): 508 - 542, 2016.
[Har80] [HNS19]	Robin HARTSHORNE, Stable reflexive sheaves, Math. Ann. 254 (1980), 121–176. M. HERING, B. NILL, H. SÜSS, Stability of tangent bundles on smooth toric Picard- rank-2 varieties and surfaces. ArXiv preprint 1910.08848, https://arxiv.org/abs/ 1910.08848v1, 2019.
[Ita76]	Shigeru IITAKA, Logarithmic forms of algebraic varieties. J. Fac. Sci., Univ. Tokyo, Sec. 1A, 23, 525 - 544 (1976).
[Kaw78]	Yujiro KAWAMATA, On deformations of compactifiable complex manifolds. Math. Ann. 235, 247–265, Springer 1978.
[Kle88]	Peter KLEINSCHMIDT, A classification of toric varieties with few generators. Aequationes Math. 35 (1988), 254–266.
[Kly90]	Alexander A. KLYACHKO, <i>Equivariant Bundles on Toral Varieties</i> . Math. USSR Izvestiya, 35(2): 337–375, 1990.
[Kob13]	Shoshichi KOBAYASHI, Differential geometry of complex vector bundle. Mathematical Society of Japan, 2013.

[Koo11] Martijn KOOL, Fixed point loci of moduli spaces of sheaves on toric varieties. Adv. Math., 227(4): 1700–1755, 2011.

[Li20]	Chi Li, On the stability of extensions of tangent sheaves on Kähler-Einstein
	Fano/Calabi-Yau pairs. Mathematische Annalen, 2020.
[Mig89]	Maurice MIGNOTTE, Mathématiques pour le calcul formel. Presses Universitaires de
	France, 1989.
[Mum62]	David MUMFORD, Projective invariants of projective structures and applications. Proc.
	Internat. Congr. Mathematicians (Stockholm, 1962), 526-530.
[Per03]	Markus PERLING, Resolutions and moduli for equivariant sheaves over toric varieties.
	PhD Dessertion University of Kaiserslautern, 2003.
[Per04]	Markus PERLING, Graded rings and equivariant sheaves on toric varieties. Math.
	Nachr. 263/264:181-197. 2004.

- [Sai75] Kyoji SAITO, On the uniformization of complements of discriminant loci. A.M.S. Summer Institute 1975.
- [Tak72] Fumio TAKEMOTO, Stable vector bundles on algebraic surfaces. Nagoya Math. J. Vol. 47 (1972), 29-48.

UNIV BREST, UMR CNRS 6205, LABORATOIRE DE MATHÉMATIQUES DE BRETAGNE ATLAN-TIQUE, FRANCE

Email address: achim.napame@univ-brest.fr