



**HAL**  
open science

## Genomic studies across the lifespan point to early mechanisms determining subcortical volumes

Quentin Le Grand, Claudia L. Satizabal, Muralidharan Sargurupremraj, Aniket Mishra, Aicha Soumare, Alexandre Laurent, Fabrice Crivello, Ami Tsuchida, Jean Shin, Melissa Macalli, et al.

► **To cite this version:**

Quentin Le Grand, Claudia L. Satizabal, Muralidharan Sargurupremraj, Aniket Mishra, Aicha Soumare, et al.. Genomic studies across the lifespan point to early mechanisms determining subcortical volumes. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 2021, 10.1016/j.bpsc.2021.10.011 . hal-03466940

**HAL Id: hal-03466940**

**<https://hal.science/hal-03466940>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Genomic studies across the lifespan point to early mechanisms determining subcortical volumes

Quentin Le Grand<sup>1</sup>, Claudia L Satizabal<sup>2,3,4,5</sup>, Muralidharan Sargurupremraj<sup>1,2</sup>, Aniket Mishra<sup>1</sup>, Aicha Soumaré<sup>1</sup>, Alexandre Laurent<sup>6,7,8</sup>, Fabrice Crivello<sup>6,7,8</sup>, Ami Tsuchida<sup>6,7,8</sup>, Jean Shin<sup>9</sup>, Mélissa Macalli<sup>1</sup>, Baljeet Singh<sup>10</sup>, Alexa S Beiser<sup>4,5,11</sup>, Charles DeCarli<sup>10</sup>, Evan Fletcher<sup>10</sup>, Tomas Paus<sup>12,13</sup>, Mark Lathrop<sup>14</sup>, Hieab H. H. Adams<sup>15,16</sup>, Joshua C. Bis<sup>17</sup>, Sudha Seshadri<sup>2,3,4,5</sup>, Christophe Tzourio<sup>1,18</sup>, Bernard Mazoyer<sup>6,7,8,19</sup>, Stéphanie Debette<sup>1,20</sup>

## Author affiliations:

1 University of Bordeaux, INSERM, Bordeaux Population Health Center, UMR1219, F-33000 Bordeaux, France

2 Glenn Biggs Institute for Alzheimer's and Neurodegenerative Diseases, UT Health San Antonio, San Antonio, TX, USA.

3 Department of Population Health Sciences, UT Health San Antonio, San Antonio, TX, USA.

4 The Framingham Heart Study, Framingham, MA, USA.

5 Department of Neurology, Boston University School of Medicine, Boston, MA, USA.

6 University of Bordeaux, Institute of Neurodegenerative Diseases, UMR5293, Neurofunctional imaging group, F-33000 Bordeaux, France

7 CNRS, Institute of Neurodegenerative Diseases, UMR5293, Neurofunctional imaging group, F-33000 Bordeaux, France

8 CEA, Institute of Neurodegenerative Diseases, UMR5293, Neurofunctional imaging group, F-33000 Bordeaux, France

9 The Hospital for Sick Children, and Departments of Physiology and Nutritional Sciences, University of Toronto, Toronto, ON, Canada

10 Imaging of Dementia and Aging (IDeA) Laboratory, Department of Neurology, University of California Davis, Davis, California, USA.

11 Department of Biostatistics, Boston University School of Public Health, Boston, MA.

12 Department of Psychiatry, Faculty of Medicine and Centre Hospitalier Universitaire Sainte-Justine, University of Montreal, Montreal, Quebec, H3T 1C5, Canada

13 Departments of Psychiatry and Psychology, University of Toronto, Toronto, Ontario M5T 1R8, Canada.

14 University of McGill Genome Center, Montreal, Quebec H3A 0G1, Canada

15 Department of Radiology and Nuclear Medicine, Erasmus MC, Rotterdam, the Netherlands.

16 Department of Clinical Genetics, Erasmus MC, Rotterdam, the Netherlands.

17 Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, USA.

18 Bordeaux University Hospital, Department of Medical Informatics, F-33000 Bordeaux, France

19 Bordeaux University Hospital, Department of Neuroradiology, F-33000 Bordeaux, France

20 Bordeaux University Hospital, Department of Neurology, Institute of Neurodegenerative Diseases, F-33000 Bordeaux, France

**Correspondence to:** Prof. Stéphanie Debette

University of Bordeaux, INSERM, Bordeaux Population Health Center, UM1219, Team VINTAGE, 146, rue Léo Saignat, F-33000 Bordeaux, France

E-mail: stephanie.debette@u-bordeaux.fr

**Keywords:** Genomics; Epidemiology; Lifecourse approach; Subcortical volumes; Dementia

**Running title:** Lifespan subcortical volumes genetics

**Abbreviations:** AD = Alzheimer's disease; eQTL = expression Quantitative Trait Loci; GWAS = Genome-Wide Association Study; LD = Linkage Disequilibrium; LDSR = LD Score Regression; GRS = Genetic Risk Score; MRI = Magnetic Resonance Imaging; PD = Parkinson's disease; SNP = Single Nucleotide Polymorphism; TWAS = Transcriptome-Wide Association Studies

## Abstract

**Background:** Subcortical brain structures play a key role in pathological processes of age-related neurodegenerative disorders. Mounting evidence also suggests that early-life factors may have an impact on the development of common late-life neurological diseases, including genetic factors that can influence both brain maturation and neurodegeneration.

**Methods:** Using large population-based brain imaging datasets across the lifespan ( $N \leq 40,628$ ) we aimed to: (i) estimate the heritability of subcortical volumes in young (18-35), middle (35-65), and older age (65+), and their genetic correlation across age groups; (ii) identify whether genetic loci associated with subcortical volumes in older persons also show associations in early adulthood, and explore underlying genes using transcriptome-wide association studies; (iii) explore their association with neurological phenotypes.

**Results:** Heritability of subcortical volumes consistently decreased with increasing age. Genetic risk scores for smaller caudate nucleus, putamen and hippocampus volume in older adults were associated with smaller volumes in young adults. Individually, ten loci associated with subcortical volumes in older adults also showed associations in young adults. Within these loci, transcriptome-wide association studies showed that expression of several genes in brain tissues (especially *MYLK2* and *TUFM*) was associated with subcortical volumes in both age-groups. One risk variant for smaller caudate nucleus volume (*TUFM* locus) was associated with lower cognitive performance. Genetically-predicted Alzheimer's disease was associated with smaller subcortical volumes in middle and older age.

**Conclusions:** Our findings provide novel insights into the genetic determinants of subcortical volumes across the lifespan. More studies are needed to decipher the underlying biology and clinical impact.

## Introduction

Subcortical structures (hippocampus, caudate nucleus, putamen, pallidum, accumbens, amygdala and thalamus) are involved in many neural processes, from autonomic and sensorimotor functions to memory, decision making and processing of reward and threat signals (1–5). These structures also play a key role in pathological processes of age-related neurodegenerative disorders, including Parkinson disease (PD) and Alzheimer’s disease (AD). Volumes of subcortical structures can be measured quantitatively in large population-based samples using brain Magnetic Resonance Imaging (MRI) and were shown to decrease with ageing and neurodegenerative diseases (6–15).

Mounting evidence suggests that factors already present at an early age may have an impact on the development of late-life neurological diseases, likely due to a complex combination of genetic and environmental factors that influence both brain maturation and neurodegeneration (16,17). Identifying early-life determinants of neurological diseases occurring in older age could therefore be crucial for developing more efficient prevention strategies. MRI-based measures of subcortical brain structures represent a powerful intermediate phenotype to decipher these early determinants, as they show alterations many years before the onset of clinical symptoms (18).

Recently, genome-wide association studies (GWAS) have enabled substantial progress in deciphering genes underlying variations in volumes of brain structures, identifying 38 independent genetic risk loci for subcortical volumes (15,19–21). To date, most of these studies have focused on middle-aged and older adults. A recent study found that AD genetic risk variants were associated with smaller hippocampal volume in young adults, suggesting that genetic determinants of late-life neurodegenerative diseases may already be associated with subtle variations in brain structure early in life (22). However, genetic determinants of MRI-

based measures may also change over time, either because of different underlying physiological or pathological processes between young and older persons or because of time-varying exposures to environmental risk factors across the lifespan (16,23,24).

Exploring the shared genetic variation underpinning early and late structural brain variations, specifically of subcortical brain volumes, could provide important novel insight into the time-course of structural brain changes throughout the adult lifespan and into the mechanisms underlying brain aging and their connection with factors modulating brain maturation. This could contribute to exploring early biological processes driving late-onset neurodegenerative diseases and open new avenues for preventive approaches.

Using large population-based brain imaging datasets across the adult lifespan we aimed to: (i) estimate the heritability of subcortical volumes from young adulthood to older age, and assess the amount of shared genetic variation across age groups; (ii) identify whether genetic loci associated with subcortical volumes in older age also show association with subcortical volumes in early adulthood, and explore underlying genes using transcriptome-wide association studies; (iii) explore the clinical significance of these results in relation with cognition, AD, and PD.

# **Methods and Materials**

## **Study Population**

Analyses were conducted in three age groups: young (18-35), middle-aged (35-65), and older adults (65+). The 35-years cut-off (also corresponding to the upper-age limit of i-Share participants) was based on previous neuroimaging studies indicating that some brain structures, including subcortical structures, reach a peak volume around this age (6,25). The 65-years cut-off (also the lower-age limit for the 3C-Dijon participants) corresponds to an age after which aging features on brain imaging tend to accelerate more steeply (6,25).

## **Heritability analyses**

Heritability analyses were conducted on individual level data from three cohorts:

### **Population-based cohort studies of unrelated individuals**

The Internet-based Students Health Research Enterprise (i-Share) study is an ongoing prospective population-based cohort study of French-speaking students (26). 1,777 participants aged 18-35 years had both brain MRI and genome-wide genotypes (**Supplementary Methods**).

The Three-City Dijon (3C-Dijon) Study is a population-based cohort study (27). 1,440 participants aged 65 years and older had both brain MRI and genome-wide genotypes, excluding participants with dementia, stroke history, or brain tumors at baseline (**Supplementary Methods**).

### **Family-based cohort studies**

The Framingham Heart Study (FHS) is a community-based cohort study comprising three generations of participants. 1,999 participants aged 36-64 years and 1,828 participants aged 65 years and older had both brain MRI and genome-wide genotypes, excluding participants with



stroke history or other neurologic disorders confounding the assessment of brain volumes at time of MRI (**Supplementary Methods**).

### **Analyses of shared genetic variation across the lifespan**

To analyze genetic associations with subcortical volumes in young adults we used the aforementioned 1,777 i-Share participants with high quality brain MRI and genome-wide genotype data. To derive genome-wide significant associations with subcortical structures in middle-aged to older adults we used summary statistics of the largest published meta-analyses of subcortical volumes GWAS (detailed in **Supplementary Methods**) for optimal power (15,20). In secondary analyses we also used smaller meta-analyses without a subset of cohorts comprising younger age groups leading to a sample size of 19,555 participants (detailed in **Supplementary Methods**). For hippocampal volume we used a previously published GWAS meta-analysis (21). Participants with prevalent dementia, stroke or other neurological pathologies potentially influencing brain measurements at the time of MRI were excluded in all but one meta-analyses (20).

This study was approved by the ethics committees of participating studies, and written informed consent was obtained from all participants.

### **MRI Acquisition and Phenotyping**

MRI acquisition parameters and phenotyping methods in individual cohorts are presented in the **Supplementary Methods** and have been described in detail (15,20,21,28).

We defined subcortical volumes by the total (left+right) grey matter volumes (cm<sup>3</sup>) of the accumbens, amygdala, caudate nucleus, hippocampus, pallidum, putamen, and thalamus. The accumbens volume was not available in the 3C-Dijon study.

## **Genotyping, quality control, and imputation**

Genome-wide genotyping platforms, quality control and imputation procedures are described in the **Supplementary Methods**.

## **Statistical analyses**

Analyses performed in this study are summarized in **Figure 1**.

### **Heritability analyses**

Heritability analyses were adjusted for age, sex, total intracranial volume, and the first four principal components of population stratification.

### **Population-based cohort studies of unrelated individuals**

To estimate heritability for each subcortical volume in i-Share and 3C-Dijon, we used GCTA (v1.26.0) to estimate the proportion of phenotype variance explained by genome-wide single nucleotide polymorphisms (SNPs) (**Figure 1, Supplementary Methods**) (29).

### **Family-based cohort studies**

To estimate heritability for subcortical volumes in FHS, we used the variance component model implemented in SOLAR accounting for familial relationships to determine the ratio of the genetic variance to the phenotypic variance (30). Mixed models were fit including fixed effects for covariates and additive effects for additive polygenetic and residual error terms.

### **Genetic correlation analyses**

We used LD Score Regression (LDSR) to estimate the genetic correlation between subcortical volumes in young and older adults (**Figure 1**) (31,32). For the older population, we first used the latest, largest published GWAS meta-analysis of subcortical brain volumes (15,20). To confirm that the effects were not driven by the youngest individuals in the GWAS meta-

analyses, we conducted secondary analyses using subcortical brain volumes GWAS meta-analyses without cohorts or consortia comprising younger participants (as described for analyses of shared genetic variation across the lifespan). For young adults, we performed GWAS of subcortical volumes in i-Share, adjusting for age, sex, total intracranial volume, and the first four principal components of population stratification. We used a linear mixed model implemented in GCTA-MLMA-LOCO (33). We used a method implemented in R (matSPDlite) to correct for multiple testing (34). Among the seven subcortical volumes studied, the number of independent tests was estimated at 4.0 based on the correlation matrix between all phenotypes, leading to a Bonferroni corrected significance threshold of  $p < 1.25 \times 10^{-2}$ .

### **Single variant analyses and Genetic Risk Score approaches**

First, we selected published genome-wide significant associations in middle-aged to older adults (15,20) and looked up associations with the corresponding phenotype in young adults (i-Share), for the lead SNP and nearby variants ( $\pm 250$  kb) in moderate to high linkage disequilibrium (LD) ( $LD-r^2 > 0.5$ ). To define significance thresholds we corrected for the four independent phenotypes in i-Share and the number of independent loci tested for each subcortical volume: accumbens:  $p < 3.13 \times 10^{-3}$ ; amygdala:  $p < 1.25 \times 10^{-2}$ ; caudate:  $p < 1.25 \times 10^{-3}$ ; hippocampus:  $p < 2.08 \times 10^{-3}$ ; pallidum:  $p < 2.08 \times 10^{-3}$ ; putamen:  $p < 1.39 \times 10^{-3}$ ; thalamus:  $p < 6.25 \times 10^{-3}$ .

Second, we generated genetic risk scores (GRS) for lower subcortical volumes in young adults (i-Share) by summing the number of published genome-wide significant independent risk alleles identified from subcortical volume GWAS in middle-aged to older adults, weighting each risk allele by the regression coefficient for the corresponding SNP in the published GWAS (**Supplementary Methods**). Associations were tested using linear regression models in R v3.6.1 and adjusted for age, sex, total intracranial volume, and the first four principal

components of population stratification. To account for multiple testing, we corrected for four independent phenotypes ( $p < 1.25 \times 10^{-2}$ ). As a sensitivity analysis, GRS analyses were repeated using summary statistics of the subcortical volumes GWAS conducted after removing cohorts with young participants as described above (**Figure 1**).

### **Transcriptome-wide association study**

To explore genes underlying genetic associations with subcortical volumes across the lifespan we performed transcriptome-wide association studies (TWAS) using TWAS-Fusion (**Figure 1, Supplementary Methods**) (35). We used summary statistics from the aforementioned published GWAS meta-analyses of subcortical volumes (15,20) and 17 publicly available gene expression quantitative trait loci (eQTL) reference panels from blood, brain and peripheral nerve tissues (**Supplementary Methods**). We used TWAS-Fusion to estimate the TWAS association statistics between predicted expression and each subcortical volume (35). Transcriptome-wide significant genes were determined in each tissue expression panel after correcting for the average number of features (3793.5 genes) and 4 independent phenotypes ( $p < 3.30 \times 10^{-6}$ ). Transcriptome-wide significant genes were then tested in conditional analysis (TWAS-Fusion) (35). Next, to ensure that observed associations did not reflect random correlation between gene expression and non-causal variants associated with subcortical volumes, we performed a colocalization analysis (COLOC) on the conditionally significant genes ( $p < 0.05$ ) to estimate the posterior probability of a shared causal variant between the gene expression and trait association (PP4) (36). Genes presenting a  $PP4 \geq 0.75$ , for which eQTLs did not reach genome-wide significance in association with subcortical volumes, and were not in LD ( $r^2 < 0.01$ ) with a lead SNP of genome-wide significant risk loci for subcortical volumes, were considered as novel.

In order to test colocalized associations in a younger population, we used the individual-level prediction of the gene expression option implemented in the Fusion software to generate expression-weights in i-Share (**Supplementary Methods**). The significance threshold accounted for the number of genes colocalized in  $\geq 1$  tissue for each phenotype (accumbens:  $p < 0.05$ ; amygdala:  $p < 0.05$ ; caudate:  $p < 2.17 \times 10^{-3}$ ; hippocampus:  $p < 1.67 \times 10^{-2}$ ; pallidum:  $p < 2.94 \times 10^{-3}$ ; putamen:  $p < 3.85 \times 10^{-3}$ ; thalamus:  $p < 0.05$ ).

### **Lifetime brain gene expression profile**

We examined the spatio-temporal expression pattern of genes in loci reaching genome-wide significance in the subcortical volumes GWAS, TWAS-COLOC significance and at least nominal significance in the i-Share TWAS. We used a public database (<https://hbatlas.org/>) comprising genome-wide exon-level transcriptome data from 1,340 tissue samples from 16 brain regions of 57 postmortem human brains, from embryonic development to late adulthood (37).

### **Clinical correlates**

First, we explored the relation of loci associated with subcortical volumes in young and older populations or at least nominally significant in TWAS in i-Share (**Table 1**) with AD, PD, and general cognitive function using summary statistics of the latest, largest published GWAS (38–40). Accounting for 10 independent loci ( $r^2 < 0.5$ ) and three traits, the significance threshold was  $p < 1.67 \times 10^{-3}$ .

Second, we tested whether genetically predicted AD or PD have an impact on subcortical volumes in the general population in young, middle-aged and older adults using the generalised summary data-based Mendelian randomization (GSMR) tool implemented in GCTA (**Supplementary Methods**) (29,41). The significance threshold accounted for four independent subcortical volumes and two diseases ( $p < 6.25 \times 10^{-3}$ ).

## **Data availability**

Data supporting these results can be made available upon reasonable request from the corresponding author.

## Results

### Heritability and genetic correlation of subcortical volumes across the lifespan

SNP-heritability estimates for all subcortical volumes in young, middle-aged, and older adults are presented in **Figure 2** and **Table S1**. We observed a decreasing trend of the average SNP-heritability estimates of subcortical volumes in cohorts of increasing age (**Figure 2 – Panel A**), and a similar decreasing trend for most individual subcortical volumes (**Figure 2 – Panel B**), both in unrelated and family-based population-based studies.

Most subcortical volumes were genetically correlated with each other among middle-aged to older adults, while in young adults genetic correlations reached significance after correction for multiple testing between hippocampus and amygdala, pallidum and caudate nucleus, and putamen and pallidum. Genetic correlations between young and middle-aged to older adults for the same individual subcortical volumes were significant for the caudate nucleus and the pallidum (**Figure 3**). These results were similar when using the published GWAS meta-analyses of subcortical volumes in middle-aged to older adults (**Figure 3 – Panel A**), and the secondary GWAS meta-analyses in older adults (**Figure 3 – Panel B**).

### Genetic associations with subcortical volumes across the lifespan

Out of the 38 genome-wide significant loci for subcortical volumes in middle-aged to older adults, 10 were significantly associated with the same volume, in the same direction, in young adults (**Table 1, Table S2**): four in the caudate nucleus, two in the hippocampus and the putamen respectively, and one in the amygdala and the pallidum. The most significant association in young adults was for rs8017172 ( $p=1.43\times 10^{-7}$ ), near *KTNI*, with putamen volume.

GRSs for smaller subcortical volumes derived from GWAS in middle-aged to older adults were significantly associated with smaller volumes of the same structure in young adults for putamen, caudate nucleus, hippocampus, amygdala, and pallidum (**Table 2**). The most significant association was observed for the putamen ( $p=5.04\times 10^{-11}$ ). When deriving the GRS from the secondary meta-analyses of GWAS in older adults exclusively, associations remained significant for the putamen, caudate nucleus, and hippocampus.

We then sought to explore putative causal genes underlying genome-wide significant associations with subcortical volumes using TWAS, initially based on published GWAS meta-analyses in middle-aged and older adults (**Figure 1**). Among all genes showing transcriptome-wide significant associations, 54 presented a high colocalization posterior probability of sharing a causal variant between the gene expression and trait association ( $\text{COLOC PP4}\geq 0.75$ ) with at least one subcortical volume, mostly in brain tissues (**Figure S1, Table S3**): 23 with caudate nucleus, 17 with pallidum, 13 with putamen, three with hippocampus, and one respectively with nucleus accumbens, amygdala, and thalamus. Among these 54 genes, 30 were in loci that did not reach genome-wide significance in the subcortical volumes GWAS and can be considered as novel (**Table S3**). Although we lacked power to detect transcriptome-wide associations in i-Share, we detected significant signals after multiple testing correction for 4 of the 54 colocalized genes described above, all at the genome-wide significant subcortical volume GWAS locus chr20q11.21 (**Figure 4, Figure S1, Table S3**): lower expression of *MYLK2* in putamen was associated with smaller caudate nucleus and putamen volume, higher expression of *FRG1B*, *MLLT10P1* and *RP4-610C12.4* in basal ganglia and cerebellum with smaller pallidum volume. With an exploratory purpose, we also considered 13 additional genes showing nominally significant associations in the young adult TWAS (**Figure 4**), including: 7 genes/transcripts for caudate nucleus (at chr16p11.2-12.1: *CCDC101*, *NPIPB7*, *NPIPB9*, *TUFM*, *EIF3C*, *RP11-1348G14.4*, *RP11-22P6.2*), 2 for pallidum (at chr20p11.21: *ENTPD6*,



*PYGB*), and 4 for putamen (at chr5q14.3: *CTC-498M16.4*, *TMEM161B*, *TMEM161B-AS1*; and chr20q11.21: *MLLT10P1*). Of note, the chr16p11.2-12.1 locus also showed significant association with the caudate nucleus in young adults (**Table 1** and **Table S2**). Most TWAS-COLOC genes from genome-wide significant risk loci for subcortical volumes that also showed nominally significant TWAS association in young adults were described to have constant expression levels in subcortical regions throughout the life course, including in the prenatal period (**Figure S2**).

## **Clinical correlates**

When exploring the association of the 14 genetic variants (in 10 independent loci) associated with subcortical volumes in both young and older adults (**Table 1**) with AD, PD, and general cognitive function we observed a genome-wide significant association of the lead SNP for smaller caudate nucleus volume at chr16p11.2-12.1 with lower general cognitive function (**Table 3**). No association reached significance with AD and PD after multiple testing correction.

Using GSMR, genetically predicted AD was significantly associated with smaller volumes of most subcortical structures except pallidum in dementia-free middle-aged and older adults from the general population, and with larger putamen volume in young adults (**Figure 5, Table S4**). Genetically predicted PD was significantly associated with larger thalamus volume in young adults (**Figure 5, Table S4**).

## Discussion

In large population-based cohort studies across the adult lifespan, we identified a consistent trend towards decreasing heritability of subcortical volumes with increasing age. We observed significant genetic correlation of caudate nucleus and pallidum volumes between young and older adults. GRSs for smaller caudate nucleus, putamen or hippocampus volumes in older adults were significantly associated with smaller volumes of the same structures in young adults. Individually, ten of 38 independent loci associated with subcortical volumes in older adults also showed significant associations with the corresponding volumes in young adults. Using TWAS with colocalization analyses, we found evidence for expression levels of 16 genes to be significantly associated with caudate nucleus, putamen, or pallidum volume both in older and young adults, pointing to biological pathways underlying structural brain changes across the adult lifespan. One genome-wide significant caudate nucleus locus in older and young adults (at *TUFM*) was associated with lower general cognitive function at genome-wide significance. We also observed an association of genetically determined AD with smaller volumes of most subcortical structures in middle-aged and older dementia-free adults.

The observed heritability trends of subcortical volumes across the adult lifespan, both in an unrelated and family-based population-based setting, support and expand on a previous meta-analysis of brain volume heritability estimates in twin and family studies. The latter showed increasing heritability from childhood to early adulthood and decreasing heritability from young adulthood to old age (42). Potential explanations include an increase of the environmental contribution to variation in subcortical volumes with increasing age, thus leading to a relative decrease of the genetic contribution, differential timing of gene expression during life, or age of onset of some disorders (43).

Genetic variants associated with smaller subcortical volumes in older persons were associated with smaller volumes of the same structure already in early adulthood, both individually and aggregated in a GRS. These results suggest that biological pathways influencing subcortical volumes in older age already have an impact on the latter in young adulthood. Although our analyses only focused on adult age, one could speculate that at least some of the susceptibility variants for smaller subcortical structures already showing significant associations at age 20 may be involved in developmental processes. This is supported by a recent study showing that polygenic risk scores for subcortical volumes in middle-aged to older adults were already associated with these volumes during infancy and early childhood (44). Several genes in loci associated with subcortical volumes in older persons were reported to be involved in neurodevelopmental processes in experimental work and are implicated in Mendelian disorders (15). *FAT3* at the chr11q14.3 locus that also showed association with caudate nucleus volume in i-Share, was for instance shown to be involved in neuronal morphogenesis and cell migration (15).

Ten individual loci associated with smaller subcortical volumes in older adults were already associated with smaller volumes of the same structures in young adults. We also showed association of subcortical volumes with up- or down-regulation of three genes in loci associated with caudate nucleus, putamen and pallidum volumes both in older and young adults: *MYLK2*, *FRG1B* and *MLLT10P1*. Among these, upregulation of *MYLK2* expression in brain tissues was significantly associated with larger caudate nucleus and putamen volume, in both older and young adults (**Figure 4, Figure S3**). In brain single-cell RNA sequencing analyses in mice, *mylk2* appears to be expressed in arterial and arteriolar smooth muscle cells and in pericytes (45,46). *MYLK2* encodes a myosin light chain kinase, a calcium/calmodulin dependent enzyme, harboring rare mutations causing hypertrophic cardiomyopathy (47).

Expression levels of four genes (*CCDC101*, *NPIP7*, *NPIP9* and *TUFM*) in a caudate nucleus GWAS locus that also shows significant association in young adults (chr16p11.2-12.1, **Table 1**) were associated with caudate nucleus volume at transcriptome-wide significance in older adults and nominal significance in young adults. Among these, increased *TUFM* expression was associated with smaller caudate nucleus volume, with evidence for colocalization in several tissues (**Figure 4**, **Figure S3**). *TUFM* is involved in combined oxidative phosphorylation deficiency 4 (COXPD4), a disease causing developmental regression, microcephaly and basal ganglia atrophy (48,49). Interestingly, the lead variant associated with smaller caudate nucleus at chr16p11.2-12.1 was associated with lower general cognitive function at genome-wide significance. Additional studies are required to confirm these findings and explore the role of *TUFM*.

The most significant association with subcortical volumes in young adults was observed for an intergenic variant at chr14q22.3, with *KTN1* as the closest gene, reaching near to genome-wide significance in association with putamen volume despite a limited sample size. Genetic variants in this region had previously been shown to be associated with putamen shape in healthy adolescents (19), our results thus strengthen the evidence for an important role of this locus in modulating putamen structure across the lifespan.

Noteworthy, genetic correlations between young and older adults were significant for striatal and pallidal volumes only and most significant associations of genetic risk scores and individual variants across age-groups, as well as transcriptome-wide associations were observed for these structures, and to a lesser extent the hippocampus. This could be at least partly explained by the recent observation that striatal and pallidal volumes peak in childhood and decline steadily thereafter, while volumes of the thalamus, amygdala, and hippocampus peak later and start declining from the sixth decade onwards (6). Moreover, interindividual variability of thalamus, amygdala, and hippocampus were found to increase with age,

suggesting that these structures may be more susceptible to environmental factors or late-acting genes (6).

Using Mendelian randomization (GSMR), we identified significant associations of genetically determined AD, but not PD, with all subcortical volumes except pallidum in older dementia-free adults, consistent with prior observations from studies using polygenic risk scores of AD on hippocampus and amygdala (50–54). These associations were not observed in young adults, in contrast with other studies, which identified an association between polygenic risk scores of AD and hippocampal volume in adolescent and young adults (22,55). These had used polygenic risk scores composed of SNPs selected at less stringent p-value thresholds (22,55). In young adults we found that genetically determined AD was associated with larger putamen volume and genetically determined PD with larger thalamus volume. If confirmed these results could reflect complex mechanisms whereby biological pathways contributing to larger maximal volumes in certain brain regions early in life could also enhance late-life neurodegenerative processes, akin to the observation that rates of age-related maturation are significantly correlated with rates of decline of white matter tracts (56).

To our knowledge, this is one of the first studies exploring associations of genetic variants with subcortical volumes across the adult lifespan. Our analyses were based on high quality MRI measurements and genome-wide genotype data in several cohorts, including a unique cohort of young students and leverage large scale meta-analyses conducted within the CHARGE consortium. We acknowledge limitations. While we describe trends in heritability estimates, we could not formally compare whether these differed significantly between age groups, therefore decreasing heritability estimates with increasing age need to be interpreted with caution. The sample size of our young adult cohort was limited, particularly compared with the large-scale meta-analysis of GWAS in older adults, and included a majority of women. Our study was restricted to the adult lifespan and should be complemented by cohorts of children

and adolescents to capture the full spectrum of genetic determinants across the entire life course. Whereas a longitudinal design would be most appropriate to explore changes in heritability estimates and genetic determinants across the lifespan in the same individuals, repeated MRIs and genetic analyses in large population-based samples are limited and have been available for the past twenty years only. While we show compelling association of genetic risk variants for AD with smaller subcortical volumes in middle and older dementia-free community persons, we did not observe such an association in young adults. Moreover, while one of the loci associated with lower caudate nucleus volume showed genome-wide significant association with lower general cognitive function, most of the loci associated with subcortical volumes across the adult lifespan were not associated with neurodegenerative diseases after multiple testing correction. This may suggest that these loci are not necessarily reflecting early neurodegenerative processes, but may point to developmental or non-pathological processes related to healthy aging. The fact that genetic loci associated with subcortical volumes in middle-aged to older adults (aggregated in polygenic risk scores) were recently found to be associated with these volumes already during infancy and early childhood could be a potential argument for a stronger role of developmental processes (44). We may have been underpowered to measure the modifying effects (deleterious or protective) of loci modulating subcortical volumes on the occurrence of neurodegenerative diseases. Finally, we cannot exclude selective survival bias for some variants when exploring their relation with late-life neurodegenerative diseases (if variants associated with subcortical volumes also influence survival).

In conclusion, our findings provide novel insight into the genetic determinants of subcortical volumes across the adult lifespan, with some evidence suggesting that specific genes such as *MYLK2* and *TUFM* may have a causal role in determining subcortical volumes already in young adulthood. Further research is warranted to decipher the underlying biological

mechanisms and inform prevention strategies for common late-life neurodegenerative diseases, for which pathological processes are known to start long before their clinical diagnosis.

## **Acknowledgements**

We thank Dr. Mathieu Bourgey, Dr. Hans Markus Munter, Alexandre Belisle, and Rui Li at the McGill Genome Center for processing the i-Share genome-wide genotype data. We thank Pr. Philippe Amouyel for supporting the funding of the 3C-Dijon study genome-wide genotyping.

Part of the computations were performed at the Bordeaux Bioinformatics Center (CBiB), University of Bordeaux and at the CREDIM (Centre de Ressource et Développement en Informatique Médicale) at University of Bordeaux, on a server infrastructure supported by the Fondation Claude Pompidou.

Quentin Le Grand benefited from the Digital Public Health Graduate Program (DPH), a PhD program supported by the French Investment for the Future Programme (17-EURE-0019).

Claudia L Satizabal receives support from R01 AG059727.

Dr. DeCarli received funding from P30 AG0101029.

## **Study funding**

The i-Share study is conducted by the Universities of Bordeaux and Versailles Saint-Quentin-en-Yvelines (France). The i-Share study has received funding by the French National Agency (Agence Nationale de la Recherche, ANR), via the ‘Investissements d’Avenir’ programme (grand number ANR-10-COHO-05) and from the University of Bordeaux Initiative of Excellence (IdEX). This project has also received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No 640643.



The 3C-Study is conducted under a partnership agreement among the Institut National de la Santé et de la Recherche Médicale (INSERM), the University of Bordeaux, and Sanofi-Aventis. The Fondation pour la Recherche Médicale funded the preparation and initiation of the study. The 3C-Study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, Mutuelle Générale de l'Éducation Nationale (MGEN), Institut de la Longévité, Conseils Régionaux of Aquitaine and Bourgogne, Fondation de France, and Ministry of Research–INSERM Programme “Cohortes et collections de données biologiques.” Christophe Tzourio and Stéphanie Debette have received investigator-initiated research funding from the French National Research Agency (ANR) and from the Fondation Leducq. This work was supported by the National Foundation for Alzheimer’s disease and related disorders, the Institut Pasteur de Lille, the labex DISTALZ and the Centre National de Génotypage.

This project is an EU Joint Programme -Neurodegenerative Disease Research (JPND) project. The project is supported through the following funding organisations under the aegis of JPND-[www.jpnd.eu](http://www.jpnd.eu): Australia, National Health and Medical Research Council, Austria, Federal Ministry of Science, Research and Economy; Canada, Canadian Institutes of Health Research; France, French National Research Agency; Germany, Federal Ministry of Education and Research; Netherlands, The Netherlands Organisation for Health Research and Development; United Kingdom, Medical Research Council.

This project has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 643417.

This project has also received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement

No 640643 and from the European Union's Horizon 2020 research and innovation programme under grant agreements No 667375 and 754517.

This work was also supported by a grant overseen by the French National Research Agency (ANR) as part of the “Investment for the Future Programme” ANR-18-RHUS-0002.

The FHS was supported by the National Heart, Lung and Blood Institute’s Framingham Heart Study (Contract No. N01-HC-25195, No. HHSN268201500001I and No. 75N92019D00031).

This study was also supported by grants from the National Institute of Aging (R01s AG031287, AG054076, AG049607, AG059421, AG059725, U01s AG049505, AG052409), the National Institute of Neurological Disorders and Stroke (R01 NS017950). The generation of FreeSurfer data was supported by the Alzheimer’s Association (AARGD-16-443384). Funding for SHARe Affymetrix genotyping was provided by NHLBI Contract N02-HL64278. The computational work reported in this paper was performed on the Shared Computing Cluster which is administered by Boston University’s Research Computing Services. We also thank all the FHS study participants.

Infrastructure for the CHARGE Consortium is supported in part by the National Heart, Lung, and Blood Institute grant R01HL105756 and NeuroCHARGE is supported in part by the National Institute on Aging grant R01AG033193.

Alexandre Laurent and Ami Tsuchida benefited a grant from Fondation pour la Recherche Médicale (DIC202161236446).

## **Disclosures**

The authors report no biomedical financial interests or potential conflicts of interest.

## References

1. Yin HH, Knowlton BJ (2006): The role of the basal ganglia in habit formation. *Nat Rev Neurosci* 7: 464–476.
2. McDonald AJ, Mott DD (2017): Functional neuroanatomy of amygdalohippocampal interconnections and their role in learning and memory: Amygdalohippocampal Interconnections. *J Neurosci Res* 95: 797–820.
3. Hikosaka O, Kim HF, Yasuda M, Yamamoto S (2014): Basal Ganglia Circuits for Reward Value–Guided Behavior. *Annu Rev Neurosci* 37: 289–306.
4. Salzman CD, Fusi S (2010): Emotion, Cognition, and Mental State Representation in Amygdala and Prefrontal Cortex. *Annu Rev Neurosci* 33: 173–202.
5. Floresco SB (2015): The Nucleus Accumbens: An Interface Between Cognition, Emotion, and Action. *Annu Rev Psychol* 66: 25–52.
6. Dima D, Modabbernia A, Papachristou E, Doucet GE, Agartz I, Aghajani M, *et al.* (2021): Subcortical volumes across the lifespan: Data from 18,605 healthy individuals aged 3-90 years. *Hum Brain Mapp*. <https://doi.org/10.1002/hbm.25320>
7. Coupé P, Manjón JV, Lanuza E, Catheline G (2019): Lifespan Changes of the Human Brain In Alzheimer’s Disease. *Sci Rep* 9: 3998.
8. de Jong LW, van der Hiele K, Veer IM, Houwing JJ, Westendorp RGJ, Bollen ELEM, *et al.* (2008): Strongly reduced volumes of putamen and thalamus in Alzheimer’s disease: an MRI study. *Brain* 131: 3277–3285.
9. Mak E, Su L, Williams GB, Firbank MJ, Lawson RA, Yarnall AJ, *et al.* (2015): Baseline and longitudinal grey matter changes in newly diagnosed Parkinson’s disease: ICICLE-PD study. *Brain* 138: 2974–2986.

10. Carriere N, Besson P, Dujardin K, Duhamel A, Defebvre L, Delmaire C, Devos D (2014): Apathy in Parkinson's disease is associated with nucleus accumbens atrophy: A magnetic resonance imaging shape analysis. *Mov Disord* 29: 897–903.
11. Möller C, Dieleman N, van der Flier WM, Versteeg A, Pijnenburg Y, Scheltens P, *et al.* (2015): More Atrophy of Deep Gray Matter Structures in Frontotemporal Dementia Compared to Alzheimer's Disease. *J Alzheimers Dis* 44: 635–647.
12. Graff-Radford J, Williams L, Jones DT, Benarroch EE (2017): Caudate nucleus as a component of networks controlling behavior. *Neurology* 89: 2192–2197.
13. Pasquini J, Durcan R, Wiblin L, Gersel Stockholm M, Rochester L, Brooks DJ, *et al.* (2019): Clinical implications of early caudate dysfunction in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 90: 1098–1104.
14. Luo X, Mao Q, Shi J, Wang X, Li C-SR (2019): Putamen gray matter volumes in neuropsychiatric and neurodegenerative disorders. *World J Psychiatry Ment Health Res* 3.
15. Satizabal CL, Adams HHH, Hibar DP, White CC, Knol MJ, Stein JL, *et al.* (2019): Genetic architecture of subcortical brain structures in 38,851 individuals. *Nat Genet* 51: 1624–1636.
16. Glahn DC, Kent JW, Sprooten E, Diego VP, Winkler AM, Curran JE, *et al.* (2013): Genetic basis of neurocognitive decline and reduced white-matter integrity in normal human brain aging. *Proc Natl Acad Sci* 110: 19006–19011.
17. Hunter DJ (2005): Gene–environment interactions in human diseases. *Nat Rev Genet* 6: 287–298.
18. Jack CR, Holtzman DM (2013): Biomarker Modeling of Alzheimer's Disease. *Neuron* 80: 1347–1358.

19. Hibar DP, Stein JL, Renteria ME, Arias-Vasquez A, Desrivières S, Jahanshad N, *et al.* (2015): Common genetic variants influence human subcortical brain structures. *Nature* 520: 224–229.
20. Hibar DP, Adams HHH, Jahanshad N, Chauhan G, Stein JL, Hofer E, *et al.* (2017): Novel genetic loci associated with hippocampal volume. *Nat Commun* 8: 13624.
21. Bis JC, DeCarli C, Smith AV, van der Lijn F, Crivello F, Fornage M, *et al.* (2012): Common variants at 12q14 and 12q24 are associated with hippocampal volume. *Nat Genet* 44: 545–551.
22. Foley SF, Tansey KE, Caseras X, Lancaster T, Bracht T, Parker G, *et al.* (2017): Multimodal Brain Imaging Reveals Structural Differences in Alzheimer’s Disease Polygenic Risk Carriers: A Study in Healthy Young Adults. *Biol Psychiatry* 81: 154–161.
23. Brouwer RM, Panizzon MS, Glahn DC, Hibar DP, Hua X, Jahanshad N, *et al.* (2017): Genetic influences on individual differences in longitudinal changes in global and subcortical brain volumes: Results of the ENIGMA plasticity working group: Heritability Estimates of Brain Changes. *Hum Brain Mapp* 38: 4444–4458.
24. Chouinard-Decorte F, McKay DR, Reid A, Khundrakpam B, Zhao L, Karama S, *et al.* (2014): Heritable changes in regional cortical thickness with age. *Brain Imaging Behav* 8: 208–216.
25. Pomponio R, Erus G, Habes M, Doshi J, Srinivasan D, Mamourian E, *et al.* (2020): Harmonization of large MRI datasets for the analysis of brain imaging patterns throughout the lifespan. *NeuroImage* 208: 116450.
26. Montagni I, Guichard E, Kurth T (2016): Association of screen time with self-perceived attention problems and hyperactivity levels in French students: a cross-sectional study. *BMJ Open* 6: e009089.

27. 3C Study Group (2003): Vascular Factors and Risk of Dementia: Design of the Three-City Study and Baseline Characteristics of the Study Population. *Neuroepidemiology* 22: 316–325.
28. Maillard P, Delcroix N, Crivello F, Dufouil C, Gicquel S, Joliot M, *et al.* (2008): An automated procedure for the assessment of white matter hyperintensities by multispectral (T1, T2, PD) MRI and an evaluation of its between-centre reproducibility based on two large community databases. *Neuroradiology* 50: 31–42.
29. Yang J, Lee SH, Goddard ME, Visscher PM (2011): GCTA: A Tool for Genome-wide Complex Trait Analysis. *Am J Hum Genet* 88: 76–82.
30. Almasy L, Blangero J (1998): Multipoint Quantitative-Trait Linkage Analysis in General Pedigrees. *Am J Hum Genet* 62: 1198–1211.
31. Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Patterson N, *et al.* (2015): LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 47: 291–295.
32. Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh P-R, *et al.* (2015): An atlas of genetic correlations across human diseases and traits. *Nat Genet* 47: 1236–1241.
33. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL (2014): Advantages and pitfalls in the application of mixed-model association methods. *Nat Genet* 46: 100–106.
34. Li J, Ji L (2005): Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix. *Heredity* 95: 221–227.
35. Gusev A, Ko A, Shi H, Bhatia G, Chung W, Penninx BWJH, *et al.* (2016): Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet* 48: 245–252.

36. Giambartolomei C, Vukcevic D, Schadt EE, Franke L, Hingorani AD, Wallace C, Plagnol V (2014): Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. *PLoS Genet* 10: e1004383.
37. Kang HJ, Kawasawa YI, Cheng F, Zhu Y, Xu X, Li M, *et al.* (2011): Spatio-temporal transcriptome of the human brain. *Nature* 478: 483–489.
38. Schwartzenuber J, Cooper S, Liu JZ, Barrio-Hernandez I, Bello E, Kumasaka N, *et al.* (2021): Genome-wide meta-analysis, fine-mapping and integrative prioritization implicate new Alzheimer’s disease risk genes. *Nat Genet* 53: 392–402.
39. Nalls MA, Blauwendraat C, Vallerga CL, Heilbron K, Bandres-Ciga S, Chang D, *et al.* (2019): Identification of novel risk loci, causal insights, and heritable risk for Parkinson’s disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 18: 1091–1102.
40. Davies G, Lam M, Harris SE, Trampush JW, Luciano M, Hill WD, *et al.* (2018): Study of 300,486 individuals identifies 148 independent genetic loci influencing general cognitive function. *Nat Commun* 9: 2098.
41. Zhu Z, Zheng Z, Zhang F, Wu Y, Trzaskowski M, Maier R, *et al.* (2018): Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun* 9: 224.
42. Batouli SAH, Trollor JN, Wen W, Sachdev PS (2014): The heritability of volumes of brain structures and its relationship to age: A review of twin and family studies. *Ageing Res Rev* 13: 1–9.
43. Giedd JN, Schmitt JE, Neale MC (2007): Structural brain magnetic resonance imaging of pediatric twins. *Hum Brain Mapp* 28: 474–481.

44. Lamballais S, Jansen PR, Labrecque JA, Ikram MA, White T (2021): Genetic scores for adult subcortical volumes associate with subcortical volumes during infancy and childhood. *Hum Brain Mapp* 42: 1583–1593.
45. Vanlandewijck M, He L, Mäe MA, Andrae J, Ando K, Del Gaudio F, *et al.* (2018): A molecular atlas of cell types and zonation in the brain vasculature. *Nature* 554: 475–480.
46. He L, Vanlandewijck M, Mäe MA, Andrae J, Ando K, Del Gaudio F, *et al.* (2018): Single-cell RNA sequencing of mouse brain and lung vascular and vessel-associated cell types. *Sci Data* 5: 180160.
47. Davis JS, Hassanzadeh S, Winitsky S, Lin H, Satorius C, Vemuri R, *et al.* (2001): The Overall Pattern of Cardiac Contraction Depends on a Spatial Gradient of Myosin Regulatory Light Chain Phosphorylation. *Cell* 107: 631–641.
48. Kohda M, Tokuzawa Y, Kishita Y, Nyuzuki H, Moriyama Y, Mizuno Y, *et al.* (2016): A Comprehensive Genomic Analysis Reveals the Genetic Landscape of Mitochondrial Respiratory Chain Complex Deficiencies. *PLoS Genet* 12. <https://doi.org/10.1371/journal.pgen.1005679>
49. Valente L, Tiranti V, Marsano RM, Malfatti E, Fernandez-Vizarra E, Donnini C, *et al.* (2007): Infantile Encephalopathy and Defective Mitochondrial DNA Translation in Patients with Mutations of Mitochondrial Elongation Factors EFG1 and EFTu. *Am J Hum Genet* 80: 44–58.
50. Chauhan G, Adams HHH, Bis JC, Weinstein G, Yu L, Töglhofer AM, *et al.* (2015): Association of Alzheimer disease GWAS loci with MRI-markers of brain aging. *Neurobiol Aging* 36: 1765.e7-1765.e16.



51. Lupton MK, Strike L, Hansell NK, Wen W, Mather KA, Armstrong NJ, *et al.* (2016): The effect of increased genetic risk for Alzheimer's disease on hippocampal and amygdala volume. *Neurobiol Aging* 40: 68–77.
52. Stocker H, Möllers T, Perna L, Brenner H (2018): The genetic risk of Alzheimer's disease beyond APOE  $\epsilon$ 4: systematic review of Alzheimer's genetic risk scores. *Transl Psychiatry* 8. <https://doi.org/10.1038/s41398-018-0221-8>
53. Mormino EC, Sperling RA, Holmes AJ, Buckner RL, De Jager PL, Smoller JW, Sabuncu MR (2016): Polygenic risk of Alzheimer disease is associated with early- and late-life processes. *Neurology* 87: 481–488.
54. Harrison JR, Mistry S, Muskett N, Escott-Price V (n.d.): From Polygenic Scores to Precision Medicine in Alzheimer's Disease: A Systematic Review. *J Alzheimers Dis* 74: 1271–1283.
55. Axelrud LK, Santoro ML, Pine DS, Talarico F, Gadelha A, Manfro GG, *et al.* (2018): Polygenic Risk Score for Alzheimer's Disease: Implications for Memory Performance and Hippocampal Volumes in Early Life. *Am J Psychiatry* 175: 555–563.
56. Kochunov P, Williamson DE, Lancaster J, Fox P, Cornell J, Blangero J, Glahn DC (2012): Fractional anisotropy of water diffusion in cerebral white matter across the lifespan. *Neurobiol Aging* 33: 9–20.

## Figure legends

**Figure 1: Study workflow and samples summary.** Samples of young adults (18-35) are presented in blue, middle-aged (35-65) to older adults (65+) in green, older adults in red and in purple for disease traits. FHS: Framingham Heart Study. GRM: Genetic relationship matrix. GRS: Genetic risk score. TWAS: Transcriptome-wide association study. <sup>15</sup>: Satizabal et al, Nat Genet 2019. <sup>20</sup>: Hibar et al, Nat Commun 2017. <sup>21</sup>: Bis et al, Nat Genet 2012. <sup>38</sup>: Schwartzentruber et al, Nat Genet 2021. <sup>39</sup>: Nalls et al, Lancet Neurol 2019. <sup>40</sup>: Davies et al, Nat Commun 2018. \*: GWAS of subcortical volumes except hippocampal volume after removing ENIGMA and CHARGE cohorts with some young participants (unpublished data).

**Figure 2: Heritability of subcortical volumes in young adults, middle-aged adults and in older adults. Panel A:** the box plots represent the distribution of the heritability of subcortical volumes. The main rectangle represents the interquartile range and the horizontal line is the median. **Panel B:** plot of the estimated SNP heritability for each subcortical volumes, in each age group. For unrelated population, “Young adults” stands for the i-Share cohort (18-35y, N=1,528); “Older adults” stands for the 3C-Dijon cohort (65+, N=1,396). For family-based population, all the groups came from the Framingham Heart Study (three generations). “Middle” or “Middle-aged adults” stands for 36-64y (N=1,999) and “Old” or “Older adults” for 65+ (N=1,828)

**Figure 3: Heatmap of the genetic correlation between subcortical volumes in young adults (“Young”), in older population (Panel A: “Old” and Panel B: “old only”) and between the two populations for equivalent structures.** \*: nominally significant; \*\*: significant after multiple-testing correction ( $p < 1.25 \times 10^{-2}$ ); Larger colored squares correspond to more significant p-values. In each square, the numbers correspond to the coefficient of the genetic correction ( $r_g$ ). “Young” stands for the i-Share cohort; “Middle to old” for subcortical volumes

from the latest largest GWAS (Satizabal et al, Nat Genet 2019 and Hibar et al, Nat Commun 2017); “Old” for the GWAS of subcortical volumes except hippocampal volume after removing ENIGMA and CHARGE cohorts with some young participants and for the CHARGE GWAS of hippocampal volume (Bis et al, Nat Genet 2012)

**Figure 4: Heatmaps of the transcriptome-wide association studies of the caudate nucleus, putamen and pallidum reaching transcriptome wide significance and colocalized in older persons(15,20) and at least nominal significance in young adults (i-Share cohort). \***: TWAS Significant ( $p < 3.30 \times 10^{-6}$ ); \*\*: Conditionally significant ( $p < 0.05$ ); \*\*\*: COLOC PP4  $> 0.75$ ; +: Nominally significant in i-Share; ++: Significant in i-Share (after multiple-testing correction: accumbens:  $p < 0.05$ ; amygdala:  $p < 0.05$ ; caudate:  $p < 2.17 \times 10^{-3}$ ; hippocampus:  $p < 1.67 \times 10^{-2}$ ; pallidum:  $p < 2.94 \times 10^{-3}$ ; putamen:  $p < 3.85 \times 10^{-3}$ ; thalamus:  $p < 0.05$ )

**Figure 5: Association between genetically predicted Alzheimer’s and Parkinson’s diseases and subcortical volumes in young, middle-aged to older and older adults using Mendelian randomization.** “Young adults” stands for the i-Share cohort; “Middle-aged to older adults” for subcortical volumes from the latest largest GWAS (Satizabal et al, Nat Genet 2019 and Hibar et al, Nat Commun 2017); “Older adults” for the GWAS of subcortical volumes except hippocampal volume after removing ENIGMA and CHARGE cohorts with some young participants and for the CHARGE GWAS of hippocampal volume (Bis et al, Nat Genet 2012)

**Table 1: Association of genome-wide significant variants for subcortical volumes in older adults with the same volumes in young adults.**

SNP*	Chr	Position	AI	Freq	Nearest genes	Lead SNP from GWAS	Single variant analysis			TWAS analyses <sup>†</sup>
							i-Share Cohort (n=1,777)			Gene colocalized in the locus <sup>‡</sup>
							Beta	SE	p	
<b>Amygdala</b>										
rs11111293	12q23.2	102,921,296	C	0.18	IGFI,LINC00485	rs11111293	-0.04	0.01	<b>4.50E-03</b>	
<b>Caudate nucleus</b>										
rs10909901	1p36.32	3,131,235	T	0.28	PRDM16	rs2817145	0.13	0.03	<b>8.03E-06</b>	
rs10830894	11q14.3	92,018,778	T	0.41	MIR4490,FAT3	rs3133370	-0.08	0.02	<b>5.98E-04</b>	
rs1953353	14q22.3	56,189,751	A	0.32	KTNI,RPL13AP3	rs148470213	-0.10	0.03	<b>1.92E-04</b>	
rs4115668	16p11.2	28,607,532	A	0.27	SULT1A2	rs1987471	-0.10	0.03	<b>1.45E-04</b>	<b>CCDC101, NPIP7, NPIP9, SULT1A1, TUFM, RP11-1348G14.4, RP11-22P6.2</b>
rs1062794	20q11.21	30,381,758	C	0.34	TPX2	rs6060983	-0.08	0.02	1.59E-03	<b>MYLK2 (++)</b>
<b>Hippocampus</b>										
rs17178006	12q14.3	65,718,299	G	0.09	MSRB3	rs61921502	-0.15	0.03	<b>1.72E-06</b>	
rs113205216	12q24.22	117,326,943	A	0.09	HRK,FBXW8	rs77956314	0.12	0.03	<b>1.41E-04</b>	
<b>Pallidum</b>										
rs945270	14q22.3	56,200,473	G	0.44	KTNI,RPL13AP3	rs10129414	-0.04	0.01	<b>2.29E-04</b>	
rs113818546	20q11.21	30,369,090	T	0.24	TPX2	rs10439607	-0.03	0.01	8.46E-03	<b>ENTPD6, FRG1B (++)</b> , <b>MLLT10P1 (++)</b> , <b>PYGB</b>
<b>Putamen</b>										
rs7445169	5q14.3	87,703,099	G	0.25	TMEM161B-AS1	rs2410767	-0.08	0.03	1.25E-02	<b>CTC-498M16.4, TMEM161B, TMEM161B-AS1</b>
rs12800264	11q23.3	117,396,269	A	0.18	DSCAML1	rs35200015	-0.16	0.04	<b>1.30E-05</b>	
rs8017172	14q22.3	56,199,048	A	0.44	KTNI,RPL13AP3	rs945270	-0.15	0.03	<b>1.43E-07</b>	
rs6060954	20q11.21	30,383,187	T	0.34	TPX2	rs6087771	-0.08	0.03	5.05E-03	<b>FRG1B, MLLT10P1, MYLK2 (++)</b>

Only loci significant in this analysis or in Transcriptome-Wide Association Study in i-Share are presented here.

P-values in **bold** are significant after multiple testing correction (accumbens:  $p < 3.13 \times 10^{-3}$ ; amygdala:  $p < 1.25 \times 10^{-2}$ ; caudate:  $p < 1.25 \times 10^{-3}$ ; hippocampus:  $p < 2.08 \times 10^{-3}$ ; pallidum:  $p < 2.08 \times 10^{-3}$ ; putamen:  $p < 1.39 \times 10^{-3}$ ; thalamus:  $p < 6.25 \times 10^{-3}$ ).

Genes in **bold** are significant in i-Share TWAS and (++) after multiple testing correction (accumbens:  $p < 0.05$ ; amygdala:  $p < 0.05$ ; caudate:  $p < 2.17 \times 10^{-3}$ ; hippocampus:  $p < 1.67 \times 10^{-2}$ ; pallidum:  $p < 2.94 \times 10^{-3}$ ; putamen:  $p < 3.85 \times 10^{-3}$ ; thalamus:  $p < 0.05$ ).

\*For each locus associated at genome-wide significant level with at least one subcortical structure in Satizabal et al and Hibar et al, associations of the lead SNP and nearby variants ( $\pm 250$  kb) in moderate to high LD ( $LD-r^2 > 0.5$ ) with the corresponding phenotype were tested in young adults. Only the top SNP of each locus is presented in this table.

<sup>†</sup>TWAS analyses based on the summary statistics from Satizabal et al, Nat Genet 2019 for subcortical volumes (except hippocampal volume) and from Hibar et al, Nat Commun 2017 for hippocampal volume and results from the i-Share TWAS (n=1,586).

<sup>‡</sup> A gene was considered as in the same locus than the top SNP from the GWAS if at least one of its eQTLs was in LD ( $r^2 > 0.01$ ) with the top SNP

**Table 2: Association of genetic risk score (from genome-wide significant variants) of subcortical volumes in older adults with the same volumes in young adults.**

Genetic Risk Scores (GRS)	i-Share Cohort (n=1,586)		
	Beta	SE	p
<b>Accumbens</b>			
GRS from full GWAS*	-0.04	0.04	3.56E-01
GRS from GWAS old only†	-0.05	0.08	4.69E-01
<b>Amygdala</b>			
GRS from full GWAS*	-0.61	0.14	<b>9.47E-06</b>
GRS from GWAS old only†	-0.23	0.15	1.29E-01
<b>Caudate nucleus</b>			
GRS from full GWAS*	-0.73	0.14	<b>9.76E-08</b>
GRS from GWAS old only†	-0.99	0.21	<b>2.31E-06</b>
<b>Hippocampus</b>			
GRS from full GWAS*	-0.46	0.09	<b>2.08E-07</b>
GRS from GWAS old only†	-1.12	0.21	<b>7.74E-08</b>
<b>Pallidum</b>			
GRS from full GWAS*	-0.28	0.07	<b>6.15E-05</b>
GRS from GWAS old only†	-0.16	0.10	1.20E-01
<b>Putamen</b>			
GRS from full GWAS*	-0.67	0.10	<b>5.04E-11</b>
GRS from GWAS old only†	-1.05	0.21	<b>5.92E-07</b>
<b>Thalamus</b>			
GRS from full GWAS*	-0.21	0.31	5.01E-01
GRS from GWAS old only†	0.07	0.54	8.98E-01

P-values in **bold** significant results after multiple testing correction ( $p < 1.25 \times 10^{-2}$ ).

\* GRS generated using the SNPs with  $p < 5E-08$  from the summary statistics of the GWAS of subcortical volumes from Satizabal et al. Nat Genet 2019 and from Hibar et al, Nat Commun 2017 for hippocampal volume

† GRS generated using the SNPs with  $p < 5E-08$  from the summary statistics of the GWAS of subcortical volumes from Satizabal et al. Nat Genet 2019 after excluding cohorts containing young participants and from Bis et al, Nat Genet 2012 for hippocampal volume

**Table 3: Association of genetic variants associated with subcortical volumes in older and young adults with Alzheimer's and Parkinson's diseases and general cognitive function**


SNP	Locus	AI	Subcortical volumes Young adults (n=1,777)		Alzheimer Disease (n=75,397 / 844,024)		Parkinson Disease (n=33,674 / 449,056)		General cognitive function (n=282,014)	
			Z	p	Z	p	Z	p	Z	p
<b>Amygdala</b>										
rs11111293	12q23.2	C	-2.84	<b>4.50E-03</b>	-0.79	4.28E-01	0.22	8.22E-01	-2.67	7.63E-03
<b>Caudate nucleus</b>										
rs10909901	1p36.32	C	-4.46	<b>8.03E-06</b>	-0.07	9.43E-01	-0.64	5.20E-01	-1.49	1.38E-01
rs10830894	11q14.3	T	-3.43	<b>5.98E-04</b>	0.20	8.43E-01	-2.12	3.40E-02	-0.37	7.10E-01
rs1953353	14q22.3	A	-3.73	<b>1.92E-04</b>	1.46	1.44E-01	-2.60	9.23E-03	-1.60	1.10E-01
rs4115668	16p11.2	A	-3.80	<b>1.45E-04</b>	-1.84	6.64E-02	-2.01	4.52E-02	-7.30	<b>2.88E-13</b>
rs1062794	20q11.21	C	-3.16	1.59E-03	-1.55	1.22E-01	-1.86	6.19E-02	-0.07	9.48E-01
<b>Hippocampus</b>										
rs17178006	12q14.3	G	-4.78	<b>1.72E-06</b>	-2.92	3.49E-03	-1.84	6.53E-02	-0.72	4.75E-01
rs113205216	12q24.22	C	-3.81	<b>1.41E-04</b>	-2.10	3.59E-02	-0.31	7.60E-01	0.73	4.68E-01
<b>Pallidum</b>										
rs945270	14q22.3	G	-3.68	<b>2.29E-04</b>	1.57	1.17E-01	-2.45	1.47E-02	-2.04	4.16E-02
rs113818546	20q11.21	T	-2.63	8.46E-03	-1.02	3.09E-01	-1.60	1.09E-01	0.10	9.18E-01
<b>Putamen</b>										
rs7445169	5q14.3	G	-2.50	1.25E-02	0.81	4.19E-01	-0.12	9.08E-01	3.63	<b>2.88E-04</b>
rs12800264	11q23.3	A	-4.36	<b>1.30E-05</b>	-0.75	4.54E-01	-0.20	8.39E-01	0.95	3.44E-01
rs8017172	14q22.3	A	-5.26	<b>1.43E-07</b>	1.44	1.49E-01	-2.54	1.10E-02	-1.98	4.77E-02
rs6060954	20q11.21	T	-2.80	5.05E-03	-1.53	1.26E-01	-1.84	6.53E-02	0.00	9.98E-01

Only loci significant in single-variant analysis or in Transcriptome-Wide Association Study in i-Share are presented here.

For subcortical volumes, P-values in **bold** are significant after multiple testing correction (accumbens:  $p < 3.13 \times 10^{-3}$ ; amygdala:  $p < 1.25 \times 10^{-2}$ ; caudate:  $p < 1.25 \times 10^{-3}$ ; hippocampus:  $p < 2.08 \times 10^{-3}$ ; pallidum:  $p < 2.08 \times 10^{-3}$ ; putamen:  $p < 1.39 \times 10^{-3}$ ; thalamus:  $p < 6.25 \times 10^{-3}$ ).


For Alzheimer's and Parkinson's diseases and general cognitive function, p-values in **bold** are significant after correction for multiple-testing ( $p < 1.67 \times 10^{-3}$ ).

Heritability




GCTA for population-based cohorts  
 SOLAR for family-based cohorts

Genetic correlation




LD score regression for genetic correlation across age groups

Single-variant and GRS analyses




Associations in young adults of SNPs associated with subcortical volumes in older adults

TWAS




TWAS and COLOC: tissue specific associations of gene expression with subcortical volumes

Clinical correlates




SNP look-ups and Mendelian randomization (GSMR): associations with brain diseases and cognition

A




B








A




B




Tissues



### Alzheimer's Disease



### Parkinson's Disease



◆ Young adults  
◆ Middle-aged to older adults  
◆ Older adults

● NS  
▲  $p < 0.05$   
■  $p < 6.25e-03$