



# To be malnourished or not to be malnourished: that is the question!

Stephane Jouneau, Mallorie Kerjouan, Ronan Thibault

## ► To cite this version:

Stephane Jouneau, Mallorie Kerjouan, Ronan Thibault. To be malnourished or not to be malnourished: that is the question!. European Respiratory Journal, 2022, 59 (1), pp.2102455. 10.1183/13993003.02455-2021 . hal-03467789

**HAL Id: hal-03467789**

**<https://hal.science/hal-03467789>**

Submitted on 28 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Correspondence**

Maitre T, Cottenet J, Godet C, et al. Chronic pulmonary aspergillosis: prevalence, favouring pulmonary diseases and prognosis. Eur Respir J 2021; 58: 2003345 [DOI: 10.1183/13993003.03345-2020].

**To be malnourished or not to be malnourished: that is the question!**

Jouneau Stéphane<sup>1,2</sup>, ORCID 0000-0002-1949-3461, Mallorie Kerjouan<sup>1</sup>, Ronan Thibault<sup>3</sup>.

<sup>1</sup>Department of Respiratory Medicine, Pontchaillou Hospital, Rennes 1 University, France

<sup>2</sup>IRSET UMR1085, Rennes 1 University, Rennes, France.

<sup>3</sup>INSERM, INRAE, Univ Rennes, Nutrition Metabolisms and Cancer, NuMeCan, Nutrition unit, CHU Rennes, Rennes, France

**Correspondence to:**

Prof. Stéphane Jouneau,  
Department of Respiratory Medicine, Pontchaillou Hospital, Rennes 1 University, 2 rue Henri  
Le Guilloux, 35033 Rennes Cedex 9, France.  
Tel: +33.(0)2.99.28.24.78  
Fax: +33.(0)2.99.28.24.80  
[Stephane.jouneau@chu-rennes.fr](mailto:Stephane.jouneau@chu-rennes.fr)

Conflict of interest: none.

Funding: none.

### Take home messages:

Physicians in charge of patients with chronic respiratory diseases, including chronic pulmonary aspergillosis, should be aware of the high prevalence of malnutrition. It is underdiagnosed and yet it is associated with worse prognosis. Malnutrition could be easily diagnosed.

### Correspondence

In the *European Respiratory Journal*, Thomas Maitre and colleagues analysed 17'290 cases of chronic pulmonary aspergillosis (CPA) retrieved between 2009 and 2018 in the French nationwide administrative hospital database (PMSI) (1). They demonstrated high rates of mortality, almost one third at one year. In multivariate analysis, they showed that mortality was increased in patients with malnutrition with the highest hazard ratio (HR): 2.49, 95% CI 2.05–3.04. Malnutrition, present in 33.4% of patients, was extracted from the database using the ICD-10 code. This implies that the malnutrition diagnosis should have been overtly diagnosed and collected in patients' charts or discharge reports. Because the study used the National health database, it was not possible for the authors to report on which criteria was diagnosed malnutrition. We could speculate that body mass index (BMI) below 21 kg/m<sup>2</sup> was probably the main criteria used, since it is the commonest diagnostic tool and the cut-off value for chronic respiratory diseases such as COPD patients (2). The mean BMI has been recently shown as even below this threshold, (20.8 ± 4.9 kg/m<sup>2</sup>) in a recent cohort of CPA patients (3). As it is well known that malnutrition is highly underdiagnosed at hospital as well as in the community setting (4), we could speculate that the prevalence of malnutrition could have even been higher than 33.4% if a systematic and multimodal assessment of malnutrition was performed. In 2019, new international guidelines for malnutrition diagnosis have been published by "The Global Leadership Initiative on Malnutrition" (GLIM) (5) to increase the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

screening and the treatment of malnutrition all over the world. The GLIM guidelines recommend assessing nutritional status by integrating five criteria: BMI, weight loss, loss of muscle mass (by bioimpedance analysis for example) or function, and reduced food intake, using semi-quantitative tools, or the presence of an inflammatory disease such as CPA. Undoubtedly, the use of a full assessment of malnutrition could have made identifying more malnourished patients, and may have strengthened the relation between malnutrition and mortality. Therefore, based on the latter and Thomas Maitre’s paper, and as we proposed in IPF patients (6), the respiratory care community should propose a more systematic and integrative screening of malnutrition in all patients with chronic respiratory diseases (7). It is needed to better understand the relation between malnutrition and clinical outcomes, and optimize patient’s care. It could be as simple as assessing weight and height at the first visit, in order to calculate BMI, assessing food intake (e.g. by using the Simple Evaluation of Food Intake (SEFI®) which is a 10-point visual analogue scale) (8), measuring weight and calculating weight loss at each visit. By improving the awareness of health caregivers and stakeholders towards malnutrition, we would improve patients’ quality of life and clinical outcomes as shown in COPD patients (9, 10).

1. Maitre T, Cottenet J, Godet C, Roussot A, Abdoul Carime N, Ok V, et al. Chronic pulmonary aspergillosis: prevalence, favouring pulmonary diseases and prognosis. *Eur Respir J.* 2021;58(2).
2. Schols AM, Broekhuizen R, Welink-Scheepers CA, Wouters EF. Body composition and mortality in chronic obstructive pulmonary disease. *Am J Clin Nutr.* 2005;82(1):53-9.
3. Godet C, Laurent F, Bergeron A, Ingrand P, Beigelman-Aubry C, Camara B, et al. CT Imaging Assessment of Response to Treatment in Chronic Pulmonary Aspergillosis. *Chest.* 2016;150(1):139-47.
4. Kondrup J, Allison SP, Elia M, Vellas B, Plauth M, Educational, et al. ESPEN guidelines for nutrition screening 2002. *Clin Nutr.* 2003;22(4):415-21.
5. Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. *Clin Nutr.* 2019;38(1):1-9.
6. Jouneau S, Kerjouan M, Rousseau C, Lederlin M, Llamas-Gutierrez F, De Latour B, et al. What are the best indicators to assess malnutrition in idiopathic pulmonary fibrosis patients? A cross-sectional study in a referral center. *Nutrition.* 2019;62:115-21.

- 1
- 2
- 3 7. Jouneau S, Lederlin M, Vernhet L, Thibault R. Malnutrition in idiopathic pulmonary fibrosis:
- 4 the great forgotten comorbidity! *Eur Respir J.* 2019;53(5).
- 5 8. Bouette G, Esvan M, Apel K, Thibault R. A visual analogue scale for food intake as a screening
- 6 test for malnutrition in the primary care setting: Prospective non-interventional study. *Clin Nutr.*
- 7 2021;40(1):174-80.
- 8 9. Collins PF, Elia M, Stratton RJ. Nutritional support and functional capacity in chronic
- 9 obstructive pulmonary disease: a systematic review and meta-analysis. *Respirology.* 2013;18(4):616-
- 10 29.
- 11 10. Pison CM, Cano NJ, Cherion C, Caron F, Court-Fortune I, Antonini MT, et al. Multimodal
- 12 nutritional rehabilitation improves clinical outcomes of malnourished patients with chronic
- 13 respiratory failure: a randomised controlled trial. *Thorax.* 2011;66(11):953-60.
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60