



HAL
open science

Small- and large-scale processes including anthropogenic pressures as drivers of gastropod communities in the NE Atlantic coast: A citizen science based approach

Bruno Serranito, Tristan Diméglio, Frédéric Ysnel, Anne Lizé, Eric Feunteun

► To cite this version:

Bruno Serranito, Tristan Diméglio, Frédéric Ysnel, Anne Lizé, Eric Feunteun. Small- and large-scale processes including anthropogenic pressures as drivers of gastropod communities in the NE Atlantic coast: A citizen science based approach. *Science of the Total Environment*, 2022, 816, pp.151600. 10.1016/j.scitotenv.2021.151600 . hal-03469236

HAL Id: hal-03469236

<https://hal.science/hal-03469236>

Submitted on 18 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Large-scale citizen science-based survey reveals impact of anthropogenic disturbances on gastropod communities of the North-East Atlantic rocky shores

Serranito Bruno^{1*}, Dimeglio Tristan², Ysnel Frederic^{3, 4}, Anne Lizé^{1, 4, 5} & Feunteun Eric^{1,4}

¹ Muséum National d'Histoire Naturelle (MNHN), Centre de Recherche et d'Enseignement sur les Systèmes Côtiers (CRESCO), Station Marine de Dinard, 38 rue du port blanc, 35800 Dinard, France

² Association Planète-mer, 137 avenue Clôt Bey, 13008 Marseille, France

³ Territoires, Urbanisation, Biodiversité, Environnement (EA 7462 G-TUBE), Laboratoire TUBE, Université de Rennes, 263 General Leclerc Avenue, 35042 Rennes, France

⁴ Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), Station Marine de Dinard, 38 rue du port blanc, 35800, Dinard, France

⁵ Department of Evolution, Ecology and Behaviour, University of Liverpool, Liverpool, L69 7ZB, UK

***Corresponding author:**

Email: bruno.serranito@mnhn.fr

Biological data


Citizen Science Program BioLit – ABB



Sampling in intertidal seaweed canopies
5 shore levels investigated
Abundances of 14 gastropod taxa monitored
53 sites across 1000 km of NE Atlantic coast


Environmental data

« natural environment »
descriptors




SST, Salinity, Chloa,
Wave exposure ...

Anthropogenic related
processes descriptors




nutrients, suspended matter...



Seaweed belts -
Shore levels

+




Canopy cover


Large-scale

Small-scale


10 X 10 kfolds
Crossvalidation



Community and individual taxa responses



Predicted mean spatial variability



NE Atlantic Rocky-shore
community predictions

Main Results

- The Influence of small > large scale environmental descriptors
- Anthropogenic related processes highly Impact the intertidal gastropod community
- Responses to environmental perturbations are taxa-dependant
- Anthropogenic disturbances are carried by large river estuaries

Highlights:

- Rocky-shore are subjected to multiple stressors acting at different spatial scales
- Gastropod communities were investigated using a citizen-science scheme
- The relative importance of small/large-scale variables was evaluated with RF models
- Anthropogenic stressors alter the gastropod communities at large scale
- Citizen-science programs represent powerful tools in the context of bioindication

[Click here to view linked References](#)

1
2
3

Abstract

4 Rocky-shores are among the most threatened coastal habitats by human pressures. While
5 rocky-shore communities have been increasingly used to evaluate local anthropogenic
6 perturbations such as water eutrophication, large-scale variability in relation to both natural
7 and anthropogenic stressors was poorly assessed. Here, using a Random Forest modelling
8 approach applied on citizen science-based data, we assessed the relative impact of both
9 small- and large-scale conditions (including natural and anthropogenic stressors) on
10 intertidal gastropod communities as well as taxa-specific gastropod abundances over more
11 than 1000 km of the North-East Atlantic coast. Our results demonstrate that small-scale
12 conditions (*i.e.* within shore) dominate the shaping of gastropod communities. However, this
13 community response is mainly driven by a taxa-specific affinity. Our work also highlights that
14 inorganic nutrient concentrations and suspended matter related to large river inputs are the
15 main drivers among large-scale predictors, impacting the gastropod communities by
16 depleting both taxon abundances and richness. According to models, the expected decline in
17 abundance of the yet most abundant taxa (*Steromphala umbilicalis*, *Patella* spp.,
18 *Steromphala pennanti* and *Phorcus lineatus*) is estimated to be between 65% and 85%, while
19 the common periwinkle *Littorina littorea* was the only taxon being favoured by high
20 anthropogenic activities. Such results provide relevant insights in the context of ever-
21 increasing needs for bioindicators of coastal ecosystems. Finally, this work sheds light on the
22 value of a citizen science program resulting from a consultation between professional and
23 citizen volunteers as a useful and efficient tool to investigate large-scale processes.

24 **Keywords:** intertidal gastropod community, anthropogenic stressors, citizen science, scale-
25 dependant processes, brown macroalgae canopy

26 Rocky-shores, located at the interface between land and sea, are particular habitats
27 concentrating numerous environmental stressors on a very limited surface. As a result, they
28 were considered for a long time as a “natural laboratory” to explore biological and ecological
29 issues (Helmuth et al., 2006; Levin, 1992; Paine, 2010). Owing to the relative accessibility of
30 rocky-shores, variability of the related intertidal communities were extensively documented
31 since the late nineteenth century by marine biologists and naturalists (Benson, 2002;
32 Mieszowska, 2016). Former studies mainly focused on fine-scale patterns of variability (10s
33 cm to 10s m) such as the vertical distribution pattern (the so-called “unidirectional
34 zonation”) resulting from physical and biological stress gradients closely related to tide
35 alternations, wave exposure and biotic interactions (see for instance Little et al. (2010)).
36 While vertical zonation, and other fine-scale patterns (such as ‘patchiness’) represent major
37 and ubiquitous sources of variability for rocky-shore communities (Barnes and Hughes, 1999;
38 Little et al., 2010), other studies pointed out the needs to consider variability at larger spatial
39 scales (from 100s m to 1000s km) as a response to different and entangled environmental
40 processes (*e.g.* Benedetti-Cecchi (2001)). Over the years, there was a paradigmatic shift in
41 the scale-dependant signals recognition from noise to a mean for assessing processes across
42 scales (Cruz-Motta et al., 2010; Dal Bello et al., 2017). Exploring multi-scale variability of
43 species distribution and abundance are important to identify and hierarchize the impact of
44 very broad range of processes on structure assemblage (Levin, 1992; Dungan et al., 2002;
45 Fraschetti et al., 2005).

46 In the context of increased anthropogenic pressures, of which impact on coastal systems are
47 increasingly being recognised (Halpern et al., 2015, 2008), rocky-shores are described as one
48 of the most threatened coastal habitat (Halpern et al., 2007). These concerns led to the
49 development of various indicators to address water or environmental quality status as

50 required respectively by the Water Framework Directive (WFD; Directive 2000/60/EC) and
51 the Marine Strategy Framework Directive (MSFD; 2008/56 EC) for European marine waters
52 (Borja et al., 2008). Small-scale experimental investigations on the effects of anthropogenic
53 disturbances on rocky shore communities yielded mixed results depending on sites and taxa
54 (Crowe et al., 2000). On the whole, intertidal communities such as macroalgae (Neto et al.,
55 2012; Strain et al., 2014; Ar Gall et al., 2016) or, to a lesser extent, molluscan assemblages
56 (Espinosa et al., 2007; Wells et al., 2007; Atalah, 2012) were considered as cost-effective
57 tools to monitor anthropogenic impact on coastal zones (Vinagre, 2016a). Consequently,
58 numerous investigations have reported local intertidal community variations as a response
59 to additive and synergistic impacts from anthropogenic stressors such as nutrient
60 enrichment (Atalah, 2012), sediment deposition (Airoldi, 2007) and even trampling (Micheli
61 et al., 2016) through the comparison between pristine and impacted sites (Kraufvelin et al.,
62 2006; Atalah and Crowe, 2010). However, whether and to what extent human-related
63 stressors shape large-scale rocky-shore assemblages, where many processes are entangled,
64 is still poorly explored (Crowe et al., 2000).

65 Canopy-forming brown macroalgae (*e.g.* Fucal algae) represent a structuring and remarkable
66 element of these sheltered hard-substratum intertidal ecosystems of temperate regions (van
67 den Hoek, 1975). As bioengineer species, the high diversity of seaweed provides food but
68 also protection from desiccation, temperature, wave action and predation for numerous
69 species (Chapman, 1995; Mieszkowska et al., 2013; Watt and Scrosati, 2013a). Gastropods
70 represent a dominant and a key structuring group associated with these intertidal habitats.
71 For instance, some gastropods were shown to benefit from a shelter-effect provided by the
72 macroalgae canopy, in decreasing local temperature and evaporation as compared to
73 neighbour bare rock (Bertness et al., 1999), thus expanding their spatial distribution

74 (Burrows et al., 2020). In turn, gastropods grazers structured seaweed canopies, controlling
75 both cover and diversity of macroalgae assemblages (Bertness et al., 1999) as well as the
76 grazing pressure from other invertebrates (Pavia and Toth, 2000). Therefore, gastropod
77 species, as other sedentary macroinvertebrates, are considered as taxa of interest to explore
78 the impacts of environmental change on intertidal ecosystems in the context of increasing
79 anthropogenic stressors (Atalah, 2012) and climate change (Mieszkowska et al., 2007).
80 Regarding the global decline of canopy macroalgae species (Beck and Airoidi, 2007; Álvarez-
81 Losada et al., 2020), so far only few studies have attempted to investigate large-scale (*e.g.*
82 regional and global) spatial variations of intertidal gastropod communities (Schoch et al.,
83 2006; Dal Bello et al., 2017) and in relation with anthropogenic pressures (Cruz-Motta et al.,
84 2010; Miloslavich et al., 2013). Counterintuitively, the latter studies found a weak
85 relationship between gastropod diversity and anthropogenic variables at large-scales (*i.e.* at
86 Large Marine Ecosystems 'LME' and global scales). According to authors, this could be
87 explained, at least partially, by the monitoring process (Miloslavich et al., 2013). Indeed,
88 despite the recognized value of large-scale monitoring for reporting on the status and trends
89 of biodiversity, the inherent time and financial costs remain one of the main limitations of
90 such program implementation (Caughlan and Oakley, 2001; Pereira and David Cooper,
91 2006).

92 Recent studies have demonstrated how citizen science schemes can be used to address
93 ecological issues over large-scales (Chandler et al., 2017), for instance to improve
94 conservation biogeography (Devictor et al., 2010), test macroecological theories (Vye et al.,
95 2020) or for invasive species monitoring (Lehtiniemi et al., 2020). Although its definition
96 remains unclear in the literature, citizen science involves the contribution of non-scientist
97 volunteers to address scientific issues, often through data collection (Kullenberg and

98 Kasperowski, 2016). As a result, when associated with appropriate protocols and trainings,
99 citizen-science can provide valuable sources of data, whose quality may be equivalent to
100 professional scientific collection form (van der Velde et al., 2017), but whose quantity and at
101 spatial scales generally not affordable by the latest (Tulloch et al., 2013; Theobald et al.,
102 2015). Started in 2012 the “Algues-Brunes et Bigorneaux” sighting scheme (hereafter
103 “ABB”) from the BioLit program from the French NGO Planète-Mer
104 (<https://www.planetemer.org/actions/biolit-et-biolit-junior>) focuses on gastropod
105 communities associated macroalgae canopy, by monitoring the co-occurrence of seaweeds
106 and their gastropod inhabitants. As such, the ABB protocol has the potential to account for
107 local habitat features (small-scale environment) while taking into account large-scale
108 variability. Here, we used this citizen science-based dataset to explore key drivers of
109 variability in gastropod communities associated with canopy-forming brown macroalgae
110 from North-Eastern (NE) Atlantic rocky-shores. More specifically, using an Ecological Niche
111 Modelling (ENM) method on abundance dataset, our aim was threefold: 1) to explore the
112 relative importance of local gradients linked to habitat-forming macroalgae and larger scale
113 processes, including anthropogenic processes, on gastropod communities, 2) to investigate
114 the large-scale and temporal variability on abundances and taxonomic richness of
115 gastropods across a transitional biogeographic area and 3) to evaluate species-specific
116 responses of key gastropod taxa to “natural” and anthropogenic stressors.

117

118

119 **Material and Methods**

120 **Study area**

121 From September 2012 to December 2020 (see details in Fig. SB), the AAB protocol was
122 applied to 55 rocky-shore sites spread over more than 1000 km, including 500 km of latitude,
123 along the French NE Atlantic coast (extending from 45° N to 50° N and from -4° E to 1° W).
124 According to the 'Marine Ecoregion Of the World' (MEOW) biogeographic system (Spalding
125 et al., 2007), the study sites covered 2 different provinces (Northern European Sea and
126 Lusitanian), 3 ecoregions (North Sea, Celtic Sea and South European Atlantic Shelf) (Fig. 1)
127 defining areas characterized by both specific physico-chemical features and biological
128 communities (Dinter, 2001; Ramos et al., 2012). More particularly, the Brittany coast
129 represented a biogeographical transition zone characterized by a gradual change in water
130 temperature, tidal range and wave exposure (Ramos et al., 2012; Gallon et al., 2014).

131 **Data Collection**

132 The ABB protocol focused on 16 of the most common gastropod taxa in the NE Atlantic
133 coast, associated with canopy-forming brown macroalgae and for whom the identification is
134 relatively easy for a non-expert public (see details in Fig. SA). However, among these taxa,
135 some species from the *Littorina* genus are known to exhibit significant morphological
136 variations (as for *L. saxatilis* often confused with *L. compressa*) or a lack reliable
137 morphological features to distinguish sister species (*i.e.* *L. fabalis* and *L. obtusata*). This
138 represents an important source of bias in identification, even for experts. Consequently, we
139 retrospectively decided to group abundances of these pairs into 2 species complexes,
140 namely *L. saxatilis/L. compressa* and *L. fabalis/L. obtusata*. Prior to sampling sessions, each

141 volunteer received a minimum of 4 hours of formation in the use of the protocol and taxa
142 identification. In addition, during sampling session, beginner volunteers were assisted by
143 expert naturalists. Then, taxa abundances and distributions were obtained at low tide using
144 randomly placed 0.1 m² quadrats. When quadrats fell on complex rocky fields including
145 crevices of fissures or puddles, participants were asked to reiterate the throw to limit
146 influence of microhabitats. According to the protocol, sampling was only performed in the
147 presence of main temperate intertidal canopy-forming brown macroalgae species. These
148 seaweeds were known to exhibit a clear vertical zonation pattern (*i.e.* the sequence of
149 *Pelvetia canaliculata*, *Fucus spiralis*, *Fucus vesiculosus*, *Ascophyllum nodosum* and *Fucus*
150 *serratus*, from the top to the bottom of the mediolittoral level) (Schonbeck and Norton,
151 1978; Johnson et al., 1998). Therefore, in order to describe small-scale conditions, the
152 volunteers indicated for each sample the nature of the dominant brown macroalgae species
153 (hereafter 'seaweed belt') and provided a visual estimate of canopy cover using an ordinal
154 scale (0-5; 5-25; 25-50; 50-75; 75-100% of cover). Moreover, to avoid misidentification, the
155 entire sample and each different taxon identified within the sample were photographed.
156 Each sample and taxon identified was then visually validated by experimented naturalists.

157 Prior to analysis, we applied a percentile-based outlier removal method to improve reliability
158 of the data as detailed by Li et al. (2020). For each shore level (defined by the nature of the
159 dominant seaweed species), observations corresponding to the top 1% of total abundance
160 were then removed from the initial dataset (2337 samples). In addition, to reduce influence
161 of small-scale horizontal variability (such as patchiness), taxa abundances and canopy cover
162 estimates were averaged according to the site, date and the shore level. Only samples with a
163 minimum of 2 replicates were considered for further investigations (final dataset of 407
164 averaged samples).

165 **Environmental predictors**

166 Descriptors of small-scale conditions from field observations were coupled with large-scale
167 suited environmental variables to explore both local and larger scale controls on rocky-shore
168 communities (Table 1). Large-scale processes were investigated using mean annual and
169 range of Sea Surface Temperature (SST.M and SST.r), temperature of the coldest and
170 warmest month (SST.cold and SST.warm), mean and standard deviation of coastal Salinity
171 (Sal and Sal.sd), mean and standard deviation of chlorophyll a concentrations (Chloa and
172 chloa.sd), bathymetry, mean annual Kinetic Energy due to Waves (WKE) values and the
173 mean tide amplitude (TAmp) which was defined as the difference between high and low tide
174 during spring tide. We also added predictors related to anthropogenic activities such as
175 mean and standard deviation of inorganic nitrogen ($\text{N-NO}_2^- + \text{N-NO}_3^-$) concentrations (Nit and
176 Nit.sd), mean and standard-deviation of total suspended matter (TSM.M and TSM.sd) and
177 coastal population density estimation (Pop) as proxies of eutrophication, sediment
178 deposition and direct anthropogenic pressure on coastal ecosystems, respectively.

179 For inorganic nitrogen, we produced an interpolated raster layer from monthly data
180 collected between 2012 and 2020 within the framework of the SOMLIT (“Service
181 d’Observation en Milieu LITtoral”) program from the INSU/CNRS (<http://somlit.fr>) and from
182 the REPHY (“REseau de surveillance du PHYtoplancton et des phycotoxines”) program from
183 Ifremer (“French Research Institute for the Sustainable Exploitation of the Sea”), which are
184 both national networks monitoring the hydrological parameters of French waters. Only non-
185 estuarine stations completing a minimum of 3 years of monitoring were considered in order
186 to reduce bias from sampling effort and seasonality effect. Prior to interpolations, time
187 series from each selected station were visually inspected to evaluate presence of abrupt

188 change in phenological signal and suspicious data. Finally, a 1 km resolution layer of coastal
189 waters was interpolated from mean and standard-deviation concentration values using an
190 inverse distance weighting kriging method.

191 All environmental variables were regridded to a 1 km pixel resolution from their native
192 resolution. Then, extraction at each sampling point was performed by averaging values using
193 a buffer of 2 km to deal with the heterogeneity in predictor resolutions and the potential
194 inaccuracies in the spatial localisation of sampled shores. In addition to environmental
195 variables, temporal changes in taxa abundances and taxonomic richness were investigated
196 by considering the 'season' (winter: January to March ; spring: April to June ; summer: July to
197 September ; fall: October to December) and 'year' of each sampling session. We also
198 included the number of replicates as predictor in the modelling step to explicitly account for
199 the impact of sampling effort on mean abundances.

200

201

202 **Modelling framework**

203 Abundance distribution of each taxon, as well as total gastropod abundance (Tot_ab), and
204 taxonomic richness (S) (hereafter 'response variables') were investigated using the Random
205 Forest (RF) method (Breiman, 2001). RF is a supervised tree-based approach relying on a
206 bootstrap-aggregative (bagging) scheme which aggregates predictions from multiple
207 independent decision trees (weak learners) using a subset of the dataset at each iteration
208 (Biau and Scornet, 2016). The RF is considered as a highly suitable method to address
209 ecological issues such as taxon spatial occurrence and abundance distribution (Cutler et al.,
210 2007; Hengl et al., 2018), because it allows to identify complex relationship between the
211 response variable and numerous predictors and is poorly sensitive to multicollinearity as
212 well as outliers or missing value (Li and Wang, 2013). Moreover, the RF algorithm
213 consistently outperforms other classical regression methods (Cutler et al., 2007; Li and
214 Wang, 2013; Prasad et al., 2006), even for rare or occasional taxa (Mi et al., 2017).

215 In the RF approach, two main parameters can be tuned to improve predictions: 'mtry' and
216 the 'ntrees', where 'mtry' characterizes the number of predictors selected at each tree split,
217 and 'ntrees' is the number of trees produced. For each model, 500 trees were established as
218 sufficient to stabilize the error rate in most of cases in a preliminary tests while the optimal
219 'mtry' parameter was assessed using a tuning grid as implemented in the 'caret' R package
220 (Kuhn, 2008). The setting of the 'mtry' parameter was obtained using the overall
221 performance of each model after 10 folds cross-validation repeated 10 times. This procedure
222 randomly split the response variable into training (90%) and testing datasets (remaining
223 10%). Once the models were trained on the corresponding dataset, the performance was
224 assessed using R^2 and the "Root Mean Square Error" (RMSE), which are two goodness-of-fit
225 measures summarizing variations between observed and predicted values on the tested

226 dataset. Additionally, the performance was assessed by plotting predictions as a function of
227 observations to identify model limitations (Fig. S4).
228 Finally, RF also allowed to quantify and rank the importance of each predictor (hereafter
229 'VarImp'), by computing the loss of performance associated with random permutations of
230 predictors. Furthermore, the interpretability of VarImp was supplemented by performing
231 Partial dependence Plot (PdP) and its Individual Conditional Expectations plot (ICE) produced
232 with the 'ime' R package (Molnar et al., 2018). Both PdP and ICE aim to examine how a
233 particular predictor impacts the response variable. While PdP provides an averaged model
234 prediction, ICE allows the response variability to be explored by modulating values from
235 other predictors. In the present case, predictions were performed for each particular shore
236 level (seaweed belt).

237
238
239
240

241 **Collinearity and spatial autocorrelation**

242 To minimize collinearity (Dormann et al., 2013) while maintaining model performances,
243 interpretability and comparison between taxa, predictor selection was performed following
244 a methodology inspired by Georgian et al. (2019). Predictor variables were first clustered
245 based on Pearson's correlation to identify highly correlated clusters (Fig. S3). Subsequently,
246 we produced a RF model and assessed the variable importance for each predictor. Finally,
247 among a cluster of correlated variables ($r > 0.8$), we retained the ones with the highest
248 performance values for at least one taxon and with higher ecological relevance. Thus,
249 despite their high correlation, both Total Suspended Matter and inorganic nitrogen
250 concentrations were kept as they showed high variability in VarImp values according to the
251 response variable (Fig. S3). We also computed the Variance Inflation Factor (VIF) to detect
252 collinearity in selected predictors. Predictors were finally included in models if overall VIF
253 values were lower than a tolerant threshold of 10. As a result, SSTwarm and SST.r were not
254 included in the final models as they tended to hugely inflate the VIF value.

255 Spatial autocorrelation (Legendre, 1993) was evaluated in residuals of model predictions for
256 each response variable using Moran's I test, which compares correlation measures between
257 neighbouring error values with a null distribution using 999 permutations (Moran, 1948).
258 Moran's I test was performed using the 'spdep' R package (Bivand and Wong, 2018). As
259 predictions were made for different shore levels at the same station, the Moran'I test was
260 only computed for the seaweed belt characterizing the more suitable shore level of each
261 response variable.

262

263 **Spatial variability of gastropod community at large scale**

264 Response variables were predicted for the whole French Atlantic rocky shore (from 45 to 51°
265 N; from 5°W to 2°E), but only for models considered as sufficiently robust (i.e. when
266 $R^2 > 0.35$). To delineate rocky shores, high resolution coastline was first extracted from the
267 Global Administration Areas (GADM) using the 'raster' R package (Hijmans, 2020). Then, the
268 nature of the substrate was extracted from a 1:500 000 resolution composite raster layer
269 from The Naval Hydrographic and Oceanographic Service (SHOM, 2020) for each coastline
270 point and rasterized into a 1 km grid cell resolution. Pixels with a 'Rock' feature were
271 selected to distinguish rocky from sandy and muddy shores. Large-scale predictor values
272 were then extracted for selected each pixel. Small-scale predictor variables were fixed (*i.e.*
273 seaweed belt and canopy cover class) for each response variable with the values
274 characterizing the more suitable modality according to the models (Table 2). Spearman's
275 correlation was then performed between predictions to compare the large-scale spatial
276 variability of each response variable.

277

278 **Results**

279 **Model performances**

280 According to the cross-validation, the best models showed high performances ($R^2 > 0.35$) for
281 dominant taxa as well as Tot_ab and S (Table 2). Among these response variables, the
282 highest performance was found for the common and abundant taxon *Steromphala*
283 *umbilicalis* ($R^2 = 0.67 \pm 0.1$; $RMSE = 1.11 \pm 0.2$) while *L. littorea* showed a lower relative accuracy
284 of predictions ($R^2 = 0.37 \pm 0.1$; $RMSE = 1.05 \pm 0.1$). Abundance of rare taxa such as *Nucella*
285 *lapillus* ($R^2 = 0.29 \pm 0.2$; $RMSE = 0.92 \pm 0.2$), *Ocenebra erinaceus* ($R^2 = 0.23 \pm 0.2$; $RMSE = 0.42 \pm$
286 0.1) and *Gibbula magus* ($R^2 = 0.29 \pm 0.1$; $RMSE = 0.13 \pm 0.1$) were inadequately modelled,
287 especially for *Ocenebra inornatus* or *Tritia reticulata* ($R^2 < 0.1$). Moran's I test applied on
288 robust models did not identify any significant spatial autocorrelation (Table 2).

289

290 **Variable importance and interpretation**

291 For most of response variables, small-scale predictors showed higher variable importance
292 than large-scale predictors (Table 3). The nature of the seaweed belt (shore level) was
293 generally a good predictor for gastropod community. Indeed, it represents a critical
294 predictor (rank 1) for *S. umbilicalis*, *Steromphala pennanti*, *L. fabalis/L. obtusata* and *L.*
295 *saxatilis/L. compressa* abundances as well as for Tot_ab (rank2) and S (rank 1). Specifically,
296 the higher bound of the shore (characterized by *P. canaliculata*) turned out to be highly
297 discriminating to predict most of gastropod abundances. Only *S. pennanti* and *S. cineraria*
298 abundances showed important affinity with lower bound (*F. serratus*) (Fig. 2). Moreover, the
299 models captured the relative shore level preference for each taxon as illustrated by the PdP.
300 *Steromphala umbilicalis*, *L. fabalis/L. obtusata*, *Patella* spp., *P. lineatus* and *L. littorea*
301 showed a more generalist behaviour with abundances predicted for several intermediate
302 seaweed belts, as compared to *S. pennanti* and *L. saxatilis/L. compressa*, which were mainly
303 associated with an unique one (*F. serratus* and *P. canaliculata*, respectively).

304 The local estimated canopy cover also represented an important predictor for *P. lineatus*, *L.*
305 *saxatilis/L. compressa*, *S. pennanti* (rank 2) and *Patella* spp., *L. fabalis/L. obtusata* (rank 3),
306 and to another extent for *S. umbilicalis* (rank 4), but PdP and ICE highlighted a taxa-specific
307 response with the canopy cover (Fig. 2). The RF models predicted higher abundance in light
308 cover (0-50%) for *P. lineatus* and *Patella* spp., and in medium cover (25-75%) for *S.*
309 *umbilicalis*, while *S. pennanti*, *L. fabalis/L. obtusata* and *L. saxatilis/L. compressa* were more
310 abundant in dense seaweed canopies (50-100%).

311 The mean of total suspended matter and inorganic nitrogen concentrations were among the
312 most important large-scale predictors for Tot_ab (rank 1 and 4, respectively) and S (rank 4

313 and 2, respectively) as well as for taxon (Fig. 2). The models predicted a clear negative
314 impact of suspended matter on *S. umbilicalis* (rank 2), *Patella* spp. (rank 1) and *P. lineatus*,
315 while *S. pennanti* (rank 3) was negatively impacted by inorganic nitrogen, especially in the *F.*
316 *serratus* belt. More specifically, the mean loss of abundance associated with these stressors
317 was estimated to be 65% for *Patella* spp., 70% for *S. umbilicalis*, and 85% for *S. pennanti*.
318 Conversely, *L. littorea* (rank 4) abundance was the only taxon positively affected, albeit
319 slightly (+25%) by increased inorganic nitrogen concentrations.

320 WKE also appeared to be an important predictor of *Patella* spp. (rank 1) and *P. lineatus* (rank
321 2) with a positive impact on abundances. While the importance of sampling effort logically
322 increased as the taxa became rare (Fig. S5), it nevertheless represents an important
323 predictor for relatively abundant taxa such as the complexes *L. saxatilis/L. compressa* (rank
324 3) and *L. fabalis/L. obtusata* (rank 4). Regarding the PdP and ICE, it turned out that around
325 10 replicates were necessary in most cases to correctly predict the abundance of dominant
326 taxa, albeit not for *S. cineraria*. By contrast, predictors defining temporal scales (*i.e.* ‘season’
327 and ‘year’) displayed a lower importance among predictors, indicating low temporal
328 variability of all response variables (Fig. S5).

329 **Spatial abundance distributions**

330 For robust models ($R^2 > 0.35$), predictions of spatial abundance distributions were generated
331 along the NE Atlantic rocky shores. Taxonomic richness and total gastropod abundance
332 exhibited overall similar spatial variations of predicted values ($r=0.7$) (Fig. 3) in which the
333 lowest values were associated with areas subjected to inputs from the estuaries of major
334 rivers: The Seine, the Vilaine, the Loire and the Charente. Among the predicted taxa *S.*
335 *umbilicalis* showed the highest predicted abundances (from 3.7 to >100 ind.m²) and *S.*

336 *cineraria* represented the less abundant species (from 1 to 5 ind.m²) of the mediolittoral
337 zone (Fig. S6). According to the Pearson's correlation applied on taxa-specific spatial
338 predictions, the pattern showed by the total abundance and the taxonomic richness was
339 logically shared by the abundant *S. umbilicalis* (Tot_ab: r=0.91, p<0.001, S: r=0.76, p<0.001),
340 *Patella* spp. (Tot_ab: r=0.75, p<0.001, S: r=0.43, p<0.001), *S. pennanti* (Tot_ab: r=0.45,
341 p<0.001, S: r=0.28, p<0.001) and to some extent by *S. cineraria* (Tot_ab: r=0.29, p<0.001, S:
342 r=0.34, p<0.001) and *P. lineatus* (S: r=0.4, p<0.001) (Fig. 4). Conversely, *L. littorea* showed a
343 significant opposite pattern (Tot_ab: r=-0.37, p<0.001, S: r=-0.17, p=0.002), with highest
344 abundance values (~7 ind.m⁻²) predicted in southern rocky-shores and particularly near the
345 Vilaine, the Loire and the Charente estuaries (Fig. S6). Finally, weak but significant
346 correlations in spatial predicted values were found for species complexes *L. fabalis/L.*
347 *obtusata* (Tot_ab: r=0.17, p=0.002, S: r=0.25, p<0.001) and *L. saxatilis/L. compressa* (Tot_ab:
348 r=-0.12, p=0.03).

349

350

351 **Discussion**

352 **Small-scale vs large-scale**

353 Based on a citizen science scheme, this study provides a quantification of the relative
354 importance of both local and regional scale processes on NE Atlantic mediolittoral gastropod
355 communities. Our results reveal that small-scale conditions (*i.e.* the shore scale) have a
356 greater impact than large-scale gradients on total abundance and taxonomic richness for
357 most of investigated taxa. Particularly, the seaweed belt proxy of vertical zonation is the
358 most critical factor in determining taxa abundance and diversity. While it was previously
359 considered as an universal pattern (Barnes and Hughes, 1999), the relative importance of
360 horizontal and vertical variations of intertidal rocky-shore community at large-spatial scale is
361 still debated in the scientific community. Indeed, some authors argue that vertical variations
362 have an equivalent or greater impact than horizontal ones at the site level (10s to 100s
363 meters), but not at the shore and regional levels (from 100s m to 1000s km), where
364 horizontal variability is superior in both NW Mediterranean sea (Benedetti-Cecchi, 2001) and
365 NE Atlantic rocky-shores (Valdivia et al., 2011). These former studies were recently
366 challenged by results demonstrating that vertical zonation is determinant even at the
367 regional level in shaping species composition and richness along the Catalan shoreline
368 (NW Mediterranean sea) (Chappuis et al., 2014; Cefalì et al., 2019) but also along the NW
369 Atlantic and SE Pacific (Catalán et al., 2020). Here, our results strengthen these results on a
370 500 km latitudinal gradient along the NE Atlantic coast with a specific focus on gastropods
371 communities associated with canopy habitat. Differences in the definition of the intertidal
372 zone as well as the nature of communities considered in former studies can explain
373 differences of results obtained in NE Atlantic rocky-shores. Our results identify the *Pelvetia*
374 *canaliculata* belt, located at the limit of the supralittoral zone, as the main source of vertical

375 variability in both gastropod total abundance and taxonomic richness. By contrast, in
376 Valdivia et al. (2011), authors have characterized the intertidal range using the *F. spiralis* belt
377 as the upper boundary, thus omitting the upper belts which in our study represent an
378 important part of the observed vertical variability.

379 Additionally, canopy cover also highly shapes the gastropod assemblage driven by an
380 idiosyncratic taxa response. Through experimental studies, brown macroalgae canopy cover
381 was shown to structure the temperate (Watt and Scrosati, 2013a) and sub-arctic (Ørberg et
382 al., 2018) intertidal communities by mediating the temperature, light and desiccation
383 stresses caused by prolonged emergence but only at medium and high shore levels. As a
384 result of this facilitation effect, higher canopy cover was associated with an increase of
385 biomass and diversity (Watt and Scrosati, 2013a, 2013b). Our results on taxonomic richness
386 only partially support this trend, probably due to the limited number of taxa involved.
387 However, following the so-called “space-time” substitution paradigm (Pickett, 1989; Blois et
388 al., 2013), taxa-specific responses bring new insights into how intertidal gastropod
389 communities can shift in the context of macroalgae canopy regression (Airoldi et al., 2008;
390 Álvarez-Losada et al., 2020). According to our model responses, a decrease of macroalgae
391 canopy cover may promote *P. lineatus* and *Patella spp.* as dominating the communities by
392 thriving abundance collapse of more sensitive species such as *S. pennanti*, *L. fabalis*/*L.*
393 *obtusata* and, to a lesser extent, the abundant *S. umbilicalis*, which are all associated with
394 higher macroalgae canopy cover. These predictions are in accordance with results obtained
395 in a recent canopy regression experiment where authors showed that abundances of *P.*
396 *lineatus* are negatively impacted by the *A. nodosum* cover as opposed to *L. obtusata* and *S.*
397 *umbilicalis* ones (Pocklington et al., 2018). The absence of significant response observed for
398 *L. littorea* in the cover regression experiment also supports the weak importance of the

399 canopy cover exhibited by our results. Furthermore, the antagonistic relation between
400 limpets and the canopy cover highlighted by our model is well recognized as an observed
401 pattern of temperate rocky-shore (Johnson et al., 1997; Jenkins et al., 1999a,1999b; Jenkins
402 et al., 2004; Davies et al., 2007).

403 **Large spatial scale processes**

404 At large-scale, our findings reveal the overall dominant influence of both inorganic nitrogen
405 and total suspended matter concentrations on gastropod spatial abundance distribution.
406 These results contrast with environmental gradients frequently mentioned as shaping the
407 intertidal community structure at regional and larger scale such as sea water temperature
408 (Chapman, 1995; Burrows et al., 2008; Ramos et al., 2012; Mieszkowska et al., 2013), wave
409 exposure (Burrows et al., 2008) as well as upwelling or salinity (Schoch et al., 2006). More
410 specifically, our results do not attribute the large-scale variations of gastropod community to
411 a biogeographical gradient induced by changes in water temperature. Located at the
412 intersection of 3 coastal ecoregions (Spalding et al., 2007), the same area was however
413 identified as a transitional zone for macroalgae distribution defining a boundary between
414 colder and warmer temperate waters (Dauvin, 1997; Ramos et al., 2014). In the present
415 study, the studied taxa exhibit a broad spatial distribution, as most of them are found from
416 the North of the British waters to the Lusitanian coast (Hayward et al., 2005) (for details, see
417 Fig. S1). Thus, this macroalgae transitional zone could represent the core of the thermal
418 niche of most of the gastropod taxa considered in this study, where density variations
419 induced by temperature are expected to be small. Furthermore, this apparent difference in
420 driver processes may also be partially explained by the scope of these former studies in
421 which anthropogenic related predictors were generally neglected (Sagarin et al., 2006).
422 Finally, and in a consistent way, Kinetic Energy from Waves was found to be an important

423 predictor positively affecting taxa abundance associated with low brown macroalgae cover,
424 namely *Patella* spp. and *P. lineatus*. This result highlights the overall consistency of our
425 models since high canopy cover is favoured by sheltered shores (van den Hoek, 1975) where
426 the fetch is expected to be lower than in exposed shores.

427 While both nutrient and suspended matter concentrations are related to the influence of
428 large river plumes, they referred to two different anthropogenic stressor processes affecting
429 the rocky shore community: the sedimentary reposition and eutrophication. As a
430 consequence of anthropogenic activities and particularly through soil erosion, increase of
431 sediment loads is pointed out as a major concern for hard substratum coastal communities
432 (Airoldi, 2003), impacting both cover of brown algae such as *A. nodosum* (Airoldi, 2003) or *F.*
433 *vesiculosus* (Eriksson and Johansson, 2005), as well as grazer gastropods (Airoldi, 2007;
434 Atalah and Crowe, 2010). Similarly, impact of coastal eutrophication, defined as
435 anthropogenic excessive nutrient enrichment, is increasingly documented since the mid 90s
436 (Nixon, 1995) as one of the main anthropogenic process impacting coastal community
437 (Cloern, 2001; Crowe et al., 2000; Thompson et al., 2002; Worm and Lotze, 2006). Numerous
438 experimental studies, mainly focussing on small spatial scales, have reported negative
439 impacts of high nutrient concentrations on perennial macroalgae abundance such as canopy-
440 forming brown macroalgae species (Kraufvelin et al., 2006; Worm and Lotze, 2006) and
441 sometimes in conjunction with other stressors such as sedimentation (Atalah and Crowe,
442 2010) or wave action (Kraufvelin et al., 2010). However, less attention was paid for grazer
443 molluscs such as gastropods. For instance, Atalah & Crowe (2012) showed that
444 contaminated sites were also characterized by a reduction of molluscan total abundance,
445 while emphasising a species-specific response. Our results are in agreement with former
446 results demonstrating a decrease of total abundance and taxonomic richness as a result of

447 both total suspended matter and nutrient concentration gradients. Interestingly, significant
448 impacts of these human related stressors are especially found in taxa associated with
449 medium and low mediolittoral levels and appear to be smaller for *L. saxatilis/L. compressa*,
450 located at higher shore levels. Such differences in responses across shore levels are
451 consistent with the hypothesis of water mediated disturbances as lower habitats are more
452 frequently immersed than upper ones. The taxon responses we observed in this study are
453 congruent with recent integration of these taxa as sensitive species to inorganic pollution
454 (Ecological Group I in the AZTI Marine Biotic Index (Borja et al., 2000)) (Vinagre, 2016a,
455 2016b), except for the grey top shell *S. cineraria*. This could be explained by inadequacy of
456 our sampling protocol as *S. cineraria* habitat extent further the subtidal level. Our results
457 also show that *L. littorea* is the only taxon positively affected by the gradient of nutrient
458 concentrations. *Littorina littorea* is widely reported as a species tolerant to contamination
459 since it was unaffected (Atalah, 2012) or even favoured by nutrient enrichment conditions
460 (Kraufvelin et al., 2002) or by microplastic concentrations (Doyle, 2020). Finally, this work
461 demonstrates the ability of a citizen-science program to detect local and regional changes in
462 communities and can therefore be considered as a powerful tool for monitoring intertidal
463 ecosystems, as previously discussed for instance in (Conrad and Hilchey, 2011).

464 **Caveats**

465 Notwithstanding the relevance of our results, it is important to delineate the limitations of
466 our approach in terms of data collection and methods used. One of the main caveats of this
467 study is that rare taxon abundance is zero-inflated in our dataset, and the methodology we
468 used is not sufficiently adapted to deal with it (see fig. S4). Indeed, this study provides robust

469 models for 8 taxa out of 14, therefore excluding taxa for which abundance is zero-inflated (6
470 taxa). Increasing sampling effort may allow overcoming this limitation.

471 To assess large-scale variability of the community, small-scale predictors were fixed,
472 assuming that the key brown macroalgal species defining shore height were found in all
473 shores. However, these canopy-forming brown macro-algae, have their own environmental
474 niche specificity (Chapman, 1995) and may not to be evenly distributed along the NE Atlantic
475 coast. Furthermore, grouping species with similar morphologies into species complexes may
476 also have led to an incorrect assessment of stressor impacts since the species-specific
477 responses were neglected. For instance, the lack of response to nutrient concentrations by
478 the *L. obtusata*/*L. fabalis* complex group, can be related to proven ecological differences
479 between the two merged species (Williams, 1992, 1990). Our results may also inadequately
480 address the specific effect of each stressor related to large river plumes because of potential
481 collinearity between some predictors (*e.g.* inorganic nitrogen and suspended matter
482 concentrations). Moreover, the revealed pattern could encompass other anthropogenic
483 stressors that were not considered in this study such as metallic contamination or
484 microplastic pollution (Gutow et al., 2015; Doyle, 2020). Finally, we found a weak
485 importance of temporal predictors ('season' and 'year'), describing a lesser impact on
486 abundance and species richness than spatial predictors. This low temporal variability could
487 also be due to irregularities in the time series of our dataset or the relative short time period
488 of monitoring (8 years) as compared to the time needed to observe responses to climate
489 change (Hawkins et al., 2008). As such, our findings should be interpreted with caution.

490 **Conclusion**

491 On top of their inherent fundamental interests, our results provide strong supports that
492 citizen science databases can provide robust scientific information to assess and understand
493 responses of biodiversity to environmental variability. Indeed, our modelling study reveals
494 that anthropogenic processes are the main source of variability of gastropod coastal
495 communities at the large-scale along more than 1000 km of coast within a biogeographic
496 transitional zone. Although this spatial scale is one of the most investigated one by citizen
497 science projects, only a small proportion of the provided data ends in peer-reviewed
498 publications, probably highlighting the absence of acknowledgements of citizen science data
499 in publications, which seems particularly true for marine ecosystems (Theobald et al., 2015).
500 Combined efforts of both professional and citizen scientists may achieve relevant outcomes
501 for biodiversity research as well as for the management of marine ecosystems. Finally, rocky-
502 shores community responses to anthropogenic pressures observed in this study, could be
503 used as a bioindicator of coastal water quality at the local scale as previously demonstrated
504 (Borja et al., 2000; Vinagre, 2016a) but also at regional ones where larger anthropogenic
505 processes take place.

506 **Acknowledgements**

507 The authors would like to thank all the citizen scientists having participated to the data
508 collection in the frame of BioLit program. We also acknowledge the French NGO Planète-mer
509 staff and the whole network of associations and citizens who provided support, training and
510 assistance for data validation. Special thanks to Blandine Siegrist for her help in the
511 description of taxa. This research was funded by the Foundation de France.

512

513

515 **References:**

516

517 Airoidi, L., 2007. Negative effects of sediment deposition on grazing activity and survival of the limpet
518 *Patella vulgata*. *Mar Ecol Prog Ser* 332, 235–240.

519 Airoidi, L., 2003. The effects of sedimentation on Rocky Coast Assemblages. *Oceanogr. Mar. Biol.*
520 *Annu. Rev.* 41, 161–236.

521 Airoidi, L., Balata, D., Beck, M.W., 2008. The Gray Zone: Relationships between habitat loss and
522 marine diversity and their applications in conservation. *J. Exp. Mar. Biol. Ecol.* 366, 8–15.
523 <https://doi.org/10.1016/j.jembe.2008.07.034>

524 Álvarez-Losada, Ó., Arrontes, J., Martínez, B., Fernández, C., Viejo, R.M., 2020. A regime shift in
525 intertidal assemblages triggered by loss of algal canopies: A multidecadal survey. *Mar.*
526 *Environ. Res.* 160, 104981. <https://doi.org/10.1016/j.marenvres.2020.104981>

527 Ar Gall, E., Le Duff, M., Sauriau, P.-G., de Casamajor, M.-N., Gevaert, F., Poisson, E., Hacquebart, P.,
528 Joncourt, Y., Barillé, A.-L., Buchet, R., Bréret, M., Miossec, L., 2016. Implementation of a new
529 index to assess intertidal seaweed communities as bioindicators for the European Water
530 Framework Directory. *Ecol. Indic.* 60, 162–173.
531 <https://doi.org/10.1016/j.ecolind.2015.06.035>

532 Atalah, J., 2012. Nutrient enrichment and variation in community structure on rocky shores: The
533 potential of molluscan assemblages for biomonitoring. *Estuar. Coast. Shelf Sci.* 99, 162–170.

534 Atalah, J., Crowe, T.P., 2010. Combined effects of nutrient enrichment, sedimentation and grazer loss
535 on rock pool assemblages. *J. Exp. Mar. Biol. Ecol.* 51–57.

536 Barnes, R.S.K., Hughes, R.N., 1999. *An Introduction to Marine Ecology*, 3rd Edition, Wiley-Blackwell.
537 ed.

538 Beck, M., Airoidi, L., 2007. Loss, Status and Trends for Coastal Marine Habitats of Europe, in: Gibson,
539 R., Atkinson, R., Gordon, J. (Eds.), *Oceanography and Marine Biology - An Annual Review*. CRC
540 Press, pp. 345–405. <https://doi.org/10.1201/9781420050943.ch7>

541 Belin, C., Neaud-Masson, N., Lemoine, M., Daniel, A., Lassus, P., Berthomé, J.-P., Gossel, H., Abadie,
542 E., Allenou, J.-P., Andral, B., Arnaud, C., Auby, I., Bechemin, C., Beliaeff, B., Billard, C.,
543 Bozzozero, L., Blondel, C., Bouchouca, M., Brach-Papa, C., Bruneau, A., Camus, P., 2019.
544 REPHY – French Observation and Monitoring program for Phytoplankton and Hydrology in
545 coastal waters. SEANOE. <https://doi.org/10.17882/47248>

546 Benedetti-Cecchi, L., 2001. Variability in abundance of algae and invertebrates at different spatial
547 scales on rocky sea shores. *Mar. Ecol. Prog. Ser.* 215, 79–92.
548 <https://doi.org/10.3354/meps215079>

549 Benson, K.R., 2002. The Study of Vertical Zonation on Rocky Intertidal Shores—A Historical
550 Perspective^{1,2}. *Integr. Comp. Biol.* 776–779.

551 Bertness, M.D., Leonard, G.H., Levine, J.M., Schmidt, P.R., Ingraham, A.O., 1999. Testing the relative
552 contribution of positive and negative interactions in rocky intertidal communities. *Ecology*
553 80, 2711–2726.

554 Biau, G., Scornet, E., 2016. A random forest guided tour. *TEST* 25, 197–227.
555 <https://doi.org/10.1007/s11749-016-0481-7>

556 Bivand, R.S., Wong, D.W.S., 2018. Comparing implementations of global and local indicators of spatial
557 association. *TEST* 27, 716–748. <https://doi.org/10.1007/s11749-018-0599-x>

558 Blois, J.L., Williams, J.W., Fitzpatrick, M.C., Jackson, S.T., Ferrier, S., 2013. Space can substitute for
559 time in predicting climate-change effects on biodiversity. *Proc. Natl. Acad. Sci.* 110, 9374–
560 9379. <https://doi.org/10.1073/pnas.1220228110>

561 Borja, A., Bricker, S.B., Dauer, D.M., Demetriades, N.T., Ferreira, J.G., Forbes, A.T., Hutchings, P., Jia,
562 X., Kenchington, R., Marques, J.C., Zhu, C., 2008. Overview of integrative tools and methods
563 in assessing ecological integrity in estuarine and coastal systems worldwide. *Mar. Pollut. Bull.*
564 56, 1519–1537. <https://doi.org/10.1016/j.marpolbul.2008.07.005>

565 Boudière, E., Maisondieu, C., Ardhuin, F., Accensi, M., Pineau-Guillou, L., Lepesqueur, J., 2013. A
566 suitable metocean hindcast database for the design of Marine energy converters. *Int. J. Mar.*
567 *Energy* 3–4, 40–52. <https://doi.org/10.1016/j.ijome.2013.11.010>
568 Breiman, L., 2001. Random Forests. *Machine learning* 15–32.
569 Burrows, M., Harvey, R., Robb, L., 2008. Wave exposure indices from digital coastlines and the
570 prediction of rocky shore community structure. *Mar. Ecol. Prog. Ser.* 353, 1–12.
571 <https://doi.org/10.3354/meps07284>
572 Burrows, M.T., Hawkins, S.J., Moore, J.J., Adams, L., Sugden, H., Firth, L., Mieszkowska, N., 2020.
573 Global-scale species distributions predict temperature-related changes in species
574 composition of rocky shore communities in Britain. *Glob. Change Biol.* 26, 2093–2105.
575 <https://doi.org/10.1111/gcb.14968>
576 Catalán, A.M., Valdivia, N., Scrosati, R.A., 2020. Interhemispheric comparison of scale-dependent
577 spatial variation in the structure of intertidal rocky-shore communities. *Ecosphere* 11.
578 <https://doi.org/10.1002/ecs2.3068>
579 Caughlan, L., Oakley, K.L., 2001. Cost considerations for long-term ecological monitoring. *Ecol. Indic.*
580 1, 123–134. [https://doi.org/10.1016/S1470-160X\(01\)00015-2](https://doi.org/10.1016/S1470-160X(01)00015-2)
581 Cefali, M.E., Cebrian, E., Chappuis, E., Terradas, M., Mariani, S., Ballesteros, E., 2019. Community-
582 dependent variability in species composition and richness on rocky shores at a regional scale.
583 *Estuar. Coast. Shelf Sci.* 230, 106425. <https://doi.org/10.1016/j.ecss.2019.106425>
584 Chandler, M., See, L., Copas, K., Bonde, A.M.Z., López, B.C., Danielsen, F., Legind, J.K., Masinde, S.,
585 Miller-Rushing, A.J., Newman, G., Rosemartin, A., Turak, E., 2017. Contribution of citizen
586 science towards international biodiversity monitoring. *Biol. Conserv.* 213, 280–294.
587 <https://doi.org/10.1016/j.biocon.2016.09.004>
588 Chapman, A.R.O., 1995. Functional ecology of furoid algae: twenty-three years of progress.
589 *Phycologia* 34, 1–32. <https://doi.org/10.2216/i0031-8884-34-1-1.1>
590 Chappuis, E., Terradas, M., Cefali, M.E., Mariani, S., Ballesteros, E., 2014. Vertical zonation is the main
591 distribution pattern of littoral assemblages on rocky shores at a regional scale. *Estuar. Coast.*
592 *Shelf Sci.* 147, 113–122. <https://doi.org/10.1016/j.ecss.2014.05.031>
593 Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. *Mar. Ecol.*
594 *Prog. Ser.* 210, 223–253.
595 Conrad, C.C., Hilchey, K.G., 2011. A review of citizen science and community-based environmental
596 monitoring: issues and opportunities. *Env. Monit Assess* 273–291.
597 Crowe, T.P., Thompson, R.C., Bray, S., Hawkins, S.J., 2000. Impacts of anthropogenic stress on rocky
598 intertidal communities. *J. Aquat. Ecosyst. Stress Recovery* 7, 273–297.
599 Cruz-Motta, J.J., Miloslavich, P., Palomo, G., Iken, K., Konar, B., Pohle, G., Trott, T., Benedetti-Cecchi,
600 L., Herrera, C., Hernández, A., Sardi, A., Bueno, A., Castillo, J., Klein, E., Guerra-Castro, E.,
601 Gobin, J., Gómez, D.I., Riosmena-Rodríguez, R., Mead, A., Bigatti, G., Knowlton, A.,
602 Shirayama, Y., 2010. Patterns of Spatial Variation of Assemblages Associated with Intertidal
603 Rocky Shores: A Global Perspective. *PLoS ONE* 5, e14354.
604 <https://doi.org/10.1371/journal.pone.0014354>
605 Cutler, D.R., Edwards, T.C., Beard, K.H., Cutler, A., Hess, K.T., Gibson, J., Lawler, J.J., 2007. Random
606 Forests for classification in ecology. *Ecology* 88, 2783–2792.
607 Dal Bello, M., Leclerc, J.-C., Benedetti-Cecchi, L., Andrea De Lucia, G., Arvanitidis, C., Van Avesaath, P.,
608 Bachelet, G., Bojanic, N., Como, S., Coppa, S., Coughlan, J., Crowe, T., Degraer, S., Espinosa,
609 F., Faulwetter, S., Frost, M., Guinda, X., Jankowska, E., Jourde, J., Juanes De La Pena, J.A.,
610 Kerckhof, F., Kotta, J., Lavesque, N., Magni, P., De Matos, V., Orav-Kotta, H., Pavludi, C.,
611 Pedrotti, M.L., Peleg, O., Pérez-Ruzafa, A., Puente, A., Ribeiro, P., Rigaut-Jalabert, F., Rilov, G.,
612 Rousou, M., Rubal, M., Ruginis, T., Silva, T., Simon, N., Sousa-Pinto, I., Troncoso, J., Warzocha,
613 J., Weslawski, J.M., Hummel, H., 2017. Consistent patterns of spatial variability between NE
614 Atlantic and Mediterranean rocky shores. *J. Mar. Biol. Assoc. U. K.* 97, 539–547.
615 <https://doi.org/10.1017/S0025315416001491>

- 616 Dauvin, J.C., 1997. Les biocénoses marines et littorales françaises des côtes atlantiques, Manche et
617 Mer du Nord. Synthèse, menaces et perspectives., Muséum national d'Histoire naturelle. ed.
618 Paris.
- 619 Davies, A., Johnson, M., Maggs, C., 2007. Limpet grazing and loss of *Ascophyllum nodosum* canopies
620 on decadal time scales. *Mar. Ecol. Prog. Ser.* 339, 131–141.
621 <https://doi.org/10.3354/meps339131>
- 622 Devictor, V., Whittaker, R.J., Beltrame, C., 2010. Beyond scarcity: citizen science programmes as
623 useful tools for conservation biogeography: Citizen science and conservation biogeography.
624 *Divers. Distrib.* 16, 354–362. <https://doi.org/10.1111/j.1472-4642.2009.00615.x>
- 625 Dinter, W.P., 2001. Biogeography of the OSPAR Maritime Area. German Federal Agency for Nature
626 Conservation.
- 627 Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B.,
628 Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking, B.,
629 Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: a review of
630 methods to deal with it and a simulation study evaluating their performance. *Ecography* 36,
631 27–46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
- 632 Doyle, D., 2020. Current environmental microplastic levels do not alter emergence behaviour in the
633 intertidal gastropod *Littorina littorea*. *Mar. Pollut. Bull.* 151, 1–6.
- 634 Dungan, J.L., Perry, J.N., Dale, M.R.T., Legendre, P., Citron-Pousty, S., Fortin, M.-J., Jakomulska, A.,
635 Miriti, M., Rosenberg, M.S., 2002. A balanced view of scale in spatial statistical analysis.
636 *Ecography* 25, 626–640. <https://doi.org/10.1034/j.1600-0587.2002.250510.x>
- 637 Eriksson, B.K., Johansson, G., 2005. Effects of sedimentation on macroalgae: species-specific
638 responses are related to reproductive traits. *Oecologia* 143, 438–448.
639 <https://doi.org/10.1007/s00442-004-1810-1>
- 640 Espinosa, F., Guerra-García, J.M., García-Gómez, J.C., 2007. Sewage pollution and extinction risk: an
641 endangered limpet as a bioindicator? *Biodivers. Conserv.* 16, 377–397.
642 <https://doi.org/10.1007/s10531-005-3014-3>
- 643 Frascchetti, S., Terlizzi, A., Benedetti-Cecchi, L., 2005. Patterns of distribution of marine assemblages
644 from rocky shores: evidence of relevant scales of variation. *Mar. Ecol. Prog. Ser.* 296, 13–29.
645 <https://doi.org/10.3354/meps296013>
- 646 Gallon, R.K., Robuchon, M., Leroy, B., Le Gall, L., Valero, M., Feunteun, E., 2014. Twenty years of
647 observed and predicted changes in subtidal red seaweed assemblages along a
648 biogeographical transition zone: inferring potential causes from environmental data. *J.*
649 *Biogeogr.* 41, 2293–2306. <https://doi.org/10.1111/jbi.12380>
- 650 Georgian, S.E., Anderson, O.F., Rowden, A.A., 2019. Ensemble habitat suitability modeling of
651 vulnerable marine ecosystem indicator taxa to inform deep-sea fisheries management in the
652 South Pacific Ocean. *Fish. Res.* 211, 256–274. <https://doi.org/10.1016/j.fishres.2018.11.020>
- 653 Gohin, F., Druon, J.N., Lampert, L., 2010. A five channel chlorophyll concentration algorithm applied
654 to SeaWiFS data processed by SeaDAS in coastal waters 23, 1639–1661.
- 655 Gutow, L., Eckerlebe, A., Gimén, L., 2015. Experimental Evaluation of Seaweeds as a Vector for
656 Microplastics into Marine Food Webs. *Environ. Sci. Technol.* 50, 915–923.
- 657 Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Rockwood,
658 R.C., Selig, E.R., Selkoe, K.A., Walbridge, S., 2015. Spatial and temporal changes in cumulative
659 human impacts on the world's ocean. *Nat. Commun.* 6, 1–7.
660 <https://doi.org/10.1038/ncomms8615>
- 661 Halpern, B.S., Selkoe, K.A., Micheli, F., Kappel, C.V., 2007. Evaluating and Ranking the Vulnerability of
662 Global Marine Ecosystems to Anthropogenic Threats. *Conserv. Biol.* 21, 1301–1315.
663 <https://doi.org/10.1111/j.1523-1739.2007.00752.x>
- 664 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey,
665 K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H., Madin, E.M., Perry, M.T., Selig,
666 E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global map of human impact on marine
667 ecosystems. *Science* 319, 946–948. <https://doi.org/10.1126/science.1151084>

668 Hawkins, S., Moore, P., Burrows, M., Poloczanska, E., Mieszkowska, N., Herbert, R., Jenkins, S.,
669 Thompson, R., Genner, M., Southward, A., 2008. Complex interactions in a rapidly changing
670 world: responses of rocky shore communities to recent climate change. *Clim. Res.* 37, 123–
671 133. <https://doi.org/10.3354/cr00768>

672 Hayward, P.J., Nelson-Smith, T., Shields, C., 2005. *Guides des bords de mer, Mer du Nord, Manche,*
673 *Atlantique, Méditerranée, Delachaux&Niestlé.* ed.

674 Helmuth, B., Mieszkowska, N., Moore, P., Hawkins, S.J., 2006. Living on the Edge of Two Changing
675 Worlds: Forecasting the Responses of Rocky Intertidal Ecosystems to Climate Change. *Annu.*
676 *Rev. Ecol. Evol. Syst.* 37, 373–404.
677 <https://doi.org/10.1146/annurev.ecolsys.37.091305.110149>

678 Hengl, T., Nussbaum, M., Wright, M.N., Heuvelink, G.B.M., Gräler, B., 2018. Random forest as a
679 generic framework for predictive modeling of spatial and spatio-temporal variables. *PeerJ* 6,
680 e5518. <https://doi.org/10.7717/peerj.5518>

681 Hijmans, R.J., 2020. raster: Geographic Data Analysis and Modeling.

682 Hu, C., Lee, Z., Franz, B., 2012. Chlorophyll *a* algorithms for oligotrophic oceans: A novel approach
683 based on three-band reflectance difference: A NOVEL OCEAN CHLOROPHYLL *a* ALGORITHM.
684 *J. Geophys. Res. Oceans* 117, 1–25. <https://doi.org/10.1029/2011JC007395>

685 Jenkins, S.R., Hawkins, S.J., Norton, T.A., 1999a. Direct and indirect effects of a macroalgal canopy
686 and limpet grazing in structuring a sheltered inter-tidal community.pdf. *Mar. Ecol. Prog. Ser.*
687 188, 81–92.

688 Jenkins, S.R., Hawkins, S.J., Norton, T.A., 1999b. Interaction between a fucoid canopy and limpet
689 grazing in structuring a low shore intertidal community. *J Exp Mar Biol Ecol* 233, 41–63.

690 Jenkins, S.R., Norton, T.A., Hawkins, S.J., 2004. Long term effects of *Ascophyllum nodosum* canopy
691 removal on mid shore community structure. *J. Mar. Biol. Assoc. U. K.* 84, 1–3.

692 Johnson, M., Burrows, M., Hartnoll, R., Hawkins, S., 1997. Spatial structure on moderately exposed
693 rocky shores: patch scales and the interactions between limpets and algae. *Mar. Ecol. Prog.*
694 *Ser.* 160, 209–215. <https://doi.org/10.3354/meps160209>

695 Johnson, M.P., Hawkins, S.J., Hartnoll, R.G., Norton, T.A., 1998. The establishment of fucoid zonation
696 on algal-dominated rocky shores: hypotheses derived from a simulation model: Vertical
697 zonation and algal growth on rocky shores. *Funct. Ecol.* 12, 259–269.
698 <https://doi.org/10.1046/j.1365-2435.1998.00185.x>

699 Kraufvelin, P., Christie, H., Olsen, M., 2002. Littoral macrofauna (secondary) responses to
700 experimental nutrient addition to rocky shore mesocosms and a coastal lagoon, in: Vadstein,
701 O., Olsen, Y. (Eds.), *Sustainable Increase of Marine Harvesting: Fundamental Mechanisms and*
702 *New Concepts.* Springer Netherlands, Dordrecht, pp. 149–166. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-94-017-3190-4_13)
703 [94-017-3190-4_13](https://doi.org/10.1007/978-94-017-3190-4_13)

704 Kraufvelin, P., Lindholm, A., Pedersen, M.F., Kirkerud, L.A., Bonsdorff, E., 2010. Biomass, diversity and
705 production of rocky shore macroalgae at two nutrient enrichment and wave action levels.
706 *Mar. Biol.* 157, 29–47. <https://doi.org/10.1007/s00227-009-1293-z>

707 Kraufvelin, P., Moy, F.E., Christie, H., Bokn, T.L., 2006. Nutrient Addition to Experimental Rocky Shore
708 Communities Revisited: Delayed Responses, Rapid Recovery. *Ecosystems* 9, 1076–1093.

709 Kuhn, M., 2008. Building Predictive Models in *R* Using the **caret** Package. *J. Stat. Softw.* 28.
710 <https://doi.org/10.18637/jss.v028.i05>

711 Kullenberg, C., Kasperowski, D., 2016. What Is Citizen Science? – A Scientometric Meta-Analysis. *PLoS*
712 *ONE* 11, e0147152. <https://doi.org/10.1371/journal.pone.0147152>

713 Legendre, P., 1993. Spatial Autocorrelation: Trouble or New Paradigm? *Ecology* 74, 1659–1673.
714 <https://doi.org/10.2307/1939924>

715 Lehtiniemi, M., Outinen, O., Puntilla-Dodd, R., 2020. Citizen science provides added value in the
716 monitoring for coastal non-indigenous species. *J. Environ. Manage.* 267, 1–8.
717 <https://doi.org/10.1016/j.jenvman.2020.110608>

718 Levin, S.A., 1992. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award
719 Lecture. *Ecology* 73, 1943–1967. <https://doi.org/10.2307/1941447>

720 Li, J.S., Hamann, A., Beaubien, E., 2020. Outlier detection methods to improve the quality of citizen
721 science data. *Int. J. Biometeorol.* 64, 1825–1833. <https://doi.org/10.1007/s00484-020-01968->
722 z

723 Li, X., Wang, Y., 2013. Applying various algorithms for species distribution modelling. *Integr. Zool.* 8,
724 124–135. <https://doi.org/10.1111/1749-4877.12000>

725 Little, C., Williams, G.A., Trowbridge, C.D., 2010. *The Biology of Rocky Shores: 2nd (second) Edition.* p.
726 376.

727 Mi, C., Huettmann, F., Guo, Y., Han, X., Wen, L., 2017. Why choose Random Forest to predict rare
728 species distribution with few samples in large undersampled areas? Three Asian crane
729 species models provide supporting evidence. *PeerJ* 5, 1–22.
730 <https://doi.org/10.7717/peerj.2849>

731 Micheli, F., Heiman, K.W., Kappel, C.V., Martone, R.G., Sethi, S.A., Osio, G.C., Fraschetti, S., Shelton,
732 A.O., Tanner, J.M., 2016. Combined impacts of natural and human disturbances on rocky
733 shore communities. *Ocean Coast. Manag.* 126, 42–50.
734 <https://doi.org/10.1016/j.ocecoaman.2016.03.014>

735 Mieszowska, N., 2016. Intertidal Indicators of Climate and Global Change, in: *Climate Change.*
736 Elsevier, pp. 213–229. <https://doi.org/10.1016/B978-0-444-63524-2.00014-2>

737 Mieszowska, N., Hawkins, S.J., Burrows, M.T., Kendall, M.A., 2007. Long-term changes in the
738 geographic distribution and population structures of *Osilinus lineatus* (Gastropoda:
739 Trochidae) in Britain and Ireland. *J. Mar. Biol. Assoc. U. K.* 87, 537–545.
740 <https://doi.org/10.1017/S0025315407053799>

741 Mieszowska, N., Milligan, G., Burrows, M.T., Freckleton, R., Spencer, M., 2013. Dynamic species
742 distribution models from categorical survey data. *J. Anim. Ecol.* 82, 1215–1226.
743 <https://doi.org/10.1111/1365-2656.12100>

744 Miloslavich, P., Cruz-Motta, J.J., Klein, E., Iken, K., Weinberger, V., Konar, B., Trott, T., Pohle, G.,
745 Bigatti, G., Benedetti-Cecchi, L., Shirayama, Y., Mead, A., Palomo, G., Ortiz, M., Gobin, J.,
746 Sardi, A., Díaz, J.M., Knowlton, A., Wong, M., Peralta, A.C., 2013. Large-Scale Spatial
747 Distribution Patterns of Gastropod Assemblages in Rocky Shores. *PLoS ONE* 8, e71396.
748 <https://doi.org/10.1371/journal.pone.0071396>

749 Molnar, C., Casalicchio, G., Bischl, B., 2018. iml: An R package for Interpretable Machine Learning. *J.*
750 *Open Source Softw.* 3, 1–2.

751 Moran, P.A.P., 1948. The Interpretation of Statistical Maps. *J. R. Stat. Soc. Ser. B Methodol.* 10, 243–
752 251. <https://doi.org/10.1111/j.2517-6161.1948.tb00012.x>

753 Nechad, B., Ruddick, K., Schroeder, T., Oubelkheir, K., Blondeau-Patissier, D., Cherukuru, N., Brando,
754 V., Dekker, A., Clementson, L., Banks, A.C., Maritorena, S., Werdell, P.J., Sá, C., Brotas, V.,
755 Caballero de Frutos, I., Ahn, Y.-H., Salama, S., Tilstone, G., Martinez-Vicente, V., Foley, D.,
756 McKibben, M., Nahorniak, J., Peterson, T., Siliò-Calzada, A., Röttgers, R., Lee, Z., Peters, M.,
757 Brockmann, C., 2015. CoastColour Round Robin data sets: a database to evaluate the
758 performance of algorithms for the retrieval of water quality parameters in coastal waters.
759 *Earth Syst. Sci. Data* 7, 319–348. <https://doi.org/10.5194/essd-7-319-2015>

760 Neto, J.M., Gaspar, R., Pereira, L., Marques, J.C., 2012. Marine Macroalgae Assessment Tool
761 (MarMAT) for intertidal rocky shores. Quality assessment under the scope of the European
762 Water Framework Directive. *Ecol. Indic.* 19, 39–47.
763 <https://doi.org/10.1016/j.ecolind.2011.09.006>

764 Nixon, S.W., 1995. Coastal marine eutrophication: A definition, social causes and future concerns.
765 *Ophelia* 41, 199–219.

766 Ørberg, S.B., Krause-Jensen, D., Mouritsen, K.N., Olesen, B., Marbà, N., Larsen, M.H., Blicher, M.E.,
767 Sejr, M.K., 2018. Canopy-Forming Macroalgae Facilitate Recolonization of Sub-Arctic
768 Intertidal Fauna and Reduce Temperature Extremes. *Front. Mar. Sci.* 5, 1–13.
769 <https://doi.org/10.3389/fmars.2018.00332>

770 Paine, R.T., 2010. Macroecology: Does It Ignore or Can It Encourage Further Ecological Syntheses
771 Based on Spatially Local Experimental Manipulations?: (American Society of Naturalists
772 Address). *Am. Nat.* 176, 385–393. <https://doi.org/10.1086/656273>
773 Pavia, H., Toth, G.B., 2000. Inducible chemical resistance to herbivory in the brown seaweed
774 *Ascophyllum nodosum* 81, 3212–3225.
775 Pereira, H., David Cooper, H., 2006. Towards the global monitoring of biodiversity change. *Trends*
776 *Ecol. Evol.* 21, 123–129. <https://doi.org/10.1016/j.tree.2005.10.015>
777 Pickett, S.T.A., 1989. Space-for-Time Substitution as an Alternative to Long-Term Studies, in: Likens,
778 G.E. (Ed.), *Long-Term Studies in Ecology*. Springer New York, New York, NY, pp. 110–135.
779 https://doi.org/10.1007/978-1-4615-7358-6_5
780 Pocklington, J.B., Jenkins, S.R., Bellgrove, A., Keough, M.J., O’Hara, T.D., Masterson-Algar, P.E.,
781 Hawkins, S.J., 2018. Disturbance alters ecosystem engineering by a canopy-forming alga. *J.*
782 *Mar. Biol. Assoc. U. K.* 98, 687–698. <https://doi.org/10.1017/S0025315416002009>
783 Prasad, A.M., Iverson, L.R., Liaw, A., 2006. Newer Classification and Regression Tree Techniques:
784 Bagging and Random Forests for Ecological Prediction. *Ecosystems* 9, 181–199.
785 <https://doi.org/10.1007/s10021-005-0054-1>
786 Ramos, E., Juanes, J.A., Galván, C., Neto, J.M., Melo, R., Pedersen, A., Scanlan, C., Wilkes, R., van den
787 Bergh, E., Blomqvist, M., Karup, H.P., Heiber, W., Reitsma, J.M., Ximenes, M.C., Silió, A.,
788 Méndez, F., González, B., 2012. Coastal waters classification based on physical attributes
789 along the NE Atlantic region. An approach for rocky macroalgae potential distribution. *Estuar.*
790 *Coast. Shelf Sci.* 112, 105–114. <https://doi.org/10.1016/j.ecss.2011.11.041>
791 Ramos, E., Puente, A., Juanes, J.A., Neto, J.M., Pedersen, A., Bartsch, I., Scanlan, C., Wilkes, R., Van
792 den Bergh, E., Ar Gall, E., Melo, R., 2014. Biological validation of physical coastal waters
793 classification along the NE Atlantic region based on rocky macroalgae distribution. *Estuar.*
794 *Coast. Shelf Sci.* 147, 103–112. <https://doi.org/10.1016/j.ecss.2014.05.036>
795 Sagarin, R.D., Gaines, S.D., Gaylord, B., 2006. Moving beyond assumptions to understand abundance
796 distributions across the ranges of species. *Trends Ecol. Evol.* 21, 524–530.
797 <https://doi.org/10.1016/j.tree.2006.06.008>
798 Sbrocco, E.J., Barber, P.H., 2013. MARSPEC: ocean climate layers for marine spatial ecology:
799 *Ecological Archives* E094-086. *Ecology* 94, 979–979. <https://doi.org/10.1890/12-1358.1>
800 Schoch, G.C., Menge, B.A., Allison, G., Kavanaugh, M., Thompson, S.A., A. Wood, S., 2006. Fifteen
801 degrees of separation: Latitudinal gradients of rocky intertidal biota along the California
802 Current. *Limnol. Oceanogr.* 51, 2564–2585. <https://doi.org/10.4319/lo.2006.51.6.2564>
803 Schonbeck, M., Norton, T.A., 1978. Factors controlling the upper limits of furoid algae on the shore. *J.*
804 *Exp. Mar. Biol. Ecol.* 31, 303–313. [https://doi.org/10.1016/0022-0981\(78\)90065-5](https://doi.org/10.1016/0022-0981(78)90065-5)
805 SHOM, 2020. Information géographique maritime et littorale de référence [WWW Document]. URL
806 See <https://data.shom.fr/>
807 Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern, B.S., Jorge,
808 M.A., Lombana, A., Lourie, S.A., Martin, K.D., McManus, E., Molnar, J., Recchia, C.A.,
809 Robertson, J., 2007. Marine Ecoregions of the World: A Bioregionalization of Coastal and
810 Shelf Areas. *BioScience* 57, 573–583. <https://doi.org/10.1641/B570707>
811 Strain, E.M.A., Thomson, R.J., Micheli, F., Mancuso, F.P., Airoidi, L., 2014. Identifying the interacting
812 roles of stressors in driving the global loss of canopy-forming to mat-forming algae in marine
813 ecosystems. *Glob. Change Biol.* 20, 3300–3312. <https://doi.org/10.1111/gcb.12619>
814 Theobald, E.J., Ettinger, A.K., Burgess, H.K., DeBey, L.B., Schmidt, N.R., Froehlich, H.E., Wagner, C.,
815 HilleRisLambers, J., Tewksbury, J., Harsch, M.A., Parrish, J.K., 2015. Global change and local
816 solutions: Tapping the unrealized potential of citizen science for biodiversity research. *Biol.*
817 *Conserv.* 181, 236–244. <https://doi.org/10.1016/j.biocon.2014.10.021>
818 Thompson, R.C., Crowe, T.P., Hawkins, S.J., 2002. Rocky intertidal communities: past environmental
819 changes, present status and predictions for the next 25 years. *Environ. Conserv.* 29, 168–191.
820 <https://doi.org/10.1017/S0376892902000115>

821 Tonani, M., Sykes, P., King, R.R., McConnell, N., Péquignet, A.-C., O’Dea, E., Graham, J.A., Polton, J.,
822 Siddorn, J., 2019. The impact of a new high-resolution ocean model on the Met Office North-
823 West European Shelf forecasting system. *Ocean Sci.* 15, 1133–1158.
824 <https://doi.org/10.5194/os-15-1133-2019>

825 Tulloch, A.I.T., Possingham, H.P., Joseph, L.N., Szabo, J., Martin, T.G., 2013. Realising the full potential
826 of citizen science monitoring programs. *Biol. Conserv.* 165, 128–138.
827 <https://doi.org/10.1016/j.biocon.2013.05.025>

828 Valdivia, N., Scrosati, R.A., Molis, M., Knox, A.S., 2011. Variation in Community Structure across
829 Vertical Intertidal Stress Gradients: How Does It Compare with Horizontal Variation at
830 Different Scales? *PLoS ONE* 6, e24062. <https://doi.org/10.1371/journal.pone.0024062>

831 van den Hoek, C., 1975. Phytogeographic provinces along the coasts of the northern Atlantic Ocean.
832 *Phycologia* 14, 317–330. <https://doi.org/10.2216/i0031-8884-14-4-317.1>

833 van der Velde, T., Milton, D.A., Lawson, T.J., Wilcox, C., Lansdell, M., Davis, G., Perkins, G., Hardesty,
834 B.D., 2017. Comparison of marine debris data collected by researchers and citizen scientists:
835 Is citizen science data worth the effort? *Biol. Conserv.* 208, 127–138.
836 <https://doi.org/10.1016/j.biocon.2016.05.025>

837 Vinagre, P.A., 2016a. Response of macroalgae and macroinvertebrates to anthropogenic disturbance
838 gradients in rocky shores. *Ecol. Indic.* 61, 850–854.
839 <https://doi.org/10.1016/j.ecolind.2015.10.038>

840 Vinagre, P.A., 2016b. Ability of invertebrate indices to assess ecological condition on intertidal rocky
841 shores. *Ecol. Indic.* 70, 255–268. <https://doi.org/10.1016/j.ecolind.2016.06.004>

842 Vye, S.R., Dickens, S., Adams, L., Bohn, K., Chenery, J., Dobson, N., Dunn, R.E., Earp, H.S., Evans, M.,
843 Foster, C., Grist, H., Holt, B., Hull, S., Jenkins, S.R., Lamont, P., Long, S., Mieszkowska, N.,
844 Millard, J., Morrall, Z., Pack, K., Parry-Wilson, H., Pocklington, J., Pottas, J., Richardson, L.,
845 Scott, A., Sugden, H., Watson, G., West, V., Winton, D., Delany, J., Burrows, M.T., 2020.
846 Patterns of abundance across geographical ranges as a predictor for responses to climate
847 change: Evidence from UK rocky shores. *Divers. Distrib.* 26, 1357–1365.
848 <https://doi.org/10.1111/ddi.13118>

849 Watt, C.A., Scrosati, R.A., 2013a. Bioengineer effects on understory species richness, diversity, and
850 composition change along an environmental stress gradient: Experimental and mensurative
851 evidence. *Estuar. Coast. Shelf Sci.* 123, 10–18. <https://doi.org/10.1016/j.ecss.2013.02.006>

852 Watt, C.A., Scrosati, R.A., 2013b. Regional consistency of intertidal elevation as a mediator of
853 seaweed canopy effects on benthic species richness, diversity, and composition. *Mar. Ecol.*
854 *Prog. Ser.* 491, 91–99. <https://doi.org/10.3354/meps10521>

855 Wells, E., Wilkinson, M., Wood, P., Scanlan, C., 2007. The use of macroalgal species richness and
856 composition on intertidal rocky seashores in the assessment of ecological quality under the
857 European Water Framework Directive. *Mar. Pollut. Bull.* 55, 151–161.

858 Williams, G., 1992. The effects of predation on the life histories of *Littorina obtusata* and *Littorina*
859 *mariae*. *J Mar Biol Assoc UK* 72, 403–412.

860 Williams, G., 1990. The comparative ecology of the flat periwinkles, *Littorina obtusata* (L.) and *L.*
861 *mariae* Sacchi et Rastelli. *Fied Stud.* 7, 469–482.

862 WorldPop, 2020. WorldPop -Global High Resolution Population Denominators Project [WWW
863 Document]. URL www.worldpop.org

864 Worm, B., Lotze, H.K., 2006. Effects of eutrophication, grazing, and algal blooms on rocky shores.
865 *Limnol. Oceanogr.* 51, 569–579. https://doi.org/10.4319/lo.2006.51.1_part_2.0569

866

	Variable		Abb	Units	Native Spatial résolution	Temporal coverage	Source
<i>Selected in final models</i>	<i>Large-scale</i>						
X		Bathymetry	Bath	m	1 km	-	MARSPEC (Sbrocco and Barber, 2013)
X		Mean of annual SST	SST.M	°C	1 km	Mean (2002-2010)	MARSPEC (Sbrocco and Barber, 2013)
X		SST of the coldest month	SST.cold	°C	1 km	Mean (2002-2010)	MARSPEC (Sbrocco and Barber, 2013)
		Annual range of SST	SST.r	°C	1 km		MARSPEC (Sbrocco and Barber, 2013)
		SST of the warmest month	SST.warm	°C	1 km		MARSPEC (Sbrocco and Barber, 2013)
X		Mean Salinity	Sal	-	~3 km x ~2 km	Average value of 2020 daily mean	(Tonani et al., 2019)
		Salinity Standard deviation	Sal.sd	-	~3 km x ~2 km	Average value of 2020 daily mean	(Tonani et al., 2019)
X		Mean Chloa concentrations	Chloa	mg.m ⁻³	1 km	Average value of daily mean (2013-2015)	(Gohin et al., 2010; Hu et al., 2012)
		Chloa concentrations Standard déviation	Chloa.sd	mg.m ⁻³	1 km	Average value of daily mean (2013-2015)	(Gohin et al., 2010; Hu et al., 2012)
X		Kinetic Energy due to Waves	WKE	N.m ² .s ⁻¹	300 m	Mean of 90 th percentile (2010-2015)	(Boudière et al., 2013)
X		Tide amplitude	Tamp	m			(SHOM, 2020)
X		Wave Exposure index	WE	-	90 m		(Burrows et al., 2008)
X		N-NO ₂ ⁻ + N-NO ₃ ⁻ concentrations	Nit	µmol.l ⁻¹	-	Mean (2012-2020)	(Belin et al., 2019)
		N-NO ₂ ⁻ + N-NO ₃ ⁻ concentrations Standard deviation	Nit.sd	µmol.l ⁻¹	-	Mean (2012-2020)	(Belin et al., 2019)
X		Mean of Total Suspended Matter	TSM.M	mg.l ⁻¹	300 m	Average value of daily mean (2008-2012)	(Nechad et al., 2015)
		Total Suspended Matter Standard deviation	TSM.sd	mg.l ⁻¹	300 m	Average value of daily mean (2008-2012)	(Nechad et al., 2015)
X		Coastal population density	Pop			2015	(WorldPop, 2020)
	<i>Small-scale</i>						
X		Nature of seaweed belt (shore level)	SW	-	-	-	BioLit program
X		Sampling effort (N° of replicates)	Rep	-	-	-	BioLit program

X		Canopy cover class	Cover	% class	-	-	BioLit program
---	--	--------------------	-------	---------	---	---	----------------

Table 1 Environmental predictors considered during the modeling process, with abbreviation used, units, native résolution and temporal coverage. Predictors were grouped according to their scale of influence (Small- vs. Large-scale). Predictors included in the final set are identified by X.

Response variables	mtry	RMSE	R ²	Moran's I	Fixed predictors
					SW / Cover class
S	15	0.87±0.1	0.51±0.1	-0.11 NS	Fvesi/ 0-5%
Tot_ab	15	0.75 ±0.1	0.57±0.1	-0.1 NS	Fvesi/ 0-5%
S.umbi	15	1.11 ±0.2	0.67±0.1	-0.11 NS	Fvesi/ 25-50%
L.fab/obt	15	1.07 ±0.1	0.44±0.1	-0.07 NS	Fvesi / 75-100%
Patella.spp	15	1.16 ±0.1	0.45±0.1	-0.02 NS	Fvesi/ 0-5%
S.penn	15	1.22 ±0.1	0.48±0.1	-0.1 NS	Fser/ 75-100%
P.line	15	1.08 ±0.1	0.55±0.1	-0.09 NS	Fspir/ 0-5%
L.litto	2	1.05 ±0.1	0.37±0.1	-0.07 NS	Fvesi/ 0-5%
L.saxa/comp	15	0.87 ±0.1	0.46±0.2	0.01 NS	Pel/ 0-5%
N.lapi	28	0.92 ±0.2	0.29±0.2	-	-
S.cine	25	0.50 ±0.2	0.37±0.2	-0.15 NS	F.ser/ 75-100%
T.reti	28	0.65 ±0.2	0.08±0.1	-	-
O.erin	2	0.42 ±0.1	0.20±0.2	-	-
C.zyzi	2	0.29 ±0.2	0.29±0.2	-	-
O.inor	28	0.15 ±0.1	0.01±0.0	-	-
G.magu	28	0.13 ±0.1	0.29±0.1	-	-

Table 2. Performance metrics of the best RF models (RMSE and R²) according to the 'mtry' parameter from cross-validation procedure for taxonomic richness (**S**), total abundance (**Tot_ab**) and for each taxon. *Steromphala umbilicalis* (**S.umbi**), the complex *Littorina fabalis*/*L. obtusata* (**L.fab/obt**), *Patella spp* (**Patella.spp**), *Steromphala pennanti* (**S.penn**), *Phorcus lineatus* (**P.line**), *Littorina littorea* (**L.litto**), complex *Littorina saxatilis*/*L. compressa* (**L.saxa/comp**), *Nucella lapillus* (**N.lapi**), *Steromphala cineraria* (**S.cine**), *Tritia reticulata* (**T.reti**), *Ocenebra erinaceus* (**O.erin**), *Calliostoma zyziphynum* (**C.zyzi**), *Ocinebrellus inornatus* (**O.inor**) and *Gibbula magus* (**G.magu**). Small-scale fixed predictors (*i.e.* seaweed belt (**SW**) and canopy cover class (**Cover class**)) used for model predictions and spatial

autocorrelation index (Moran's I) were indicated for robust models ($R^2 > 0.35$). SW abbreviations: *Pelvetia canaliculata* (**PeI**), *Fucus spiralis* (**Fspir**), *F. vesiculosus* (**F.vesi**), *F. serratus* (**Fser**). Non significant (NS) low values of Moran'I index indicate an absence of spatial autocorrelation in model residuals.

	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6
S	SW.Pel (100)	Nit (79.8)	Cover (76.9)	TSM.M (72.8)	WKE (70.9)	Pop (62.3)
Tot_ab	TSM.M (100)	SW.Pel (87.3)	WKE (62)	Nit (53.4)	Cover (50.3)	SST.cold (48.7)
S.umbi	SW.Pel (100)	TSM.M (38.4)	Nit (27.8)	Cover (23.1)	WKE (18)	SST.cold (10)
Patella.spp	WKE (100)	TSM.M (96)	Cover (78.5)	SST.cold (49.6)	Rep (38.03)	SW.Pel (37.1)
S.penn	SW.Fser (100)	Cover (30.5)	Nit (30)	TSM.M (21.5)	Pop (20.1)	Rep (19.2)
P.line	Cover (100)	WKE (48.6)	SW.Fser (33)	TSM.M (25.9)	Nit (20.2)	Rep (20.1)
L.fab.obt	SW.Pel (100)	Pop (60.4)	Cover (51.5)	Rep (36.6)	Sal (31.5)	Nit (24.2)
L.litto	Nit (100)	Chloa (97.3)	Sal (94.2)	WKE (92.1)	Pop (87.9.2)	SSTcold(80.3)
L.saxa.comp	SW.Pel (100)	Cover (48.5)	Rep (38.2)	SW.Fspir (35.5)	Nit (30.4)	Bath (25.3)
S.cine	SW.Fser (100)	Rep (97.7)	WKE (72.5)	Cover (46.7)	Pop (39.8)	Sal (39.4)

Table 3. Ranking of the 6 most important predictors according to standardized values of variable importance (in brackets) of each robust model ($R^2 > 0.35$). Standardized values represent the relative importance scaled to 100, identifying the most important predictor. Abbreviations: Mean total suspended matter (**TSM.M**), Canopy cover class (**Cover**), inorganic nitrogen concentrations (**Nit**), Kinetic Energy due to Waves (**WKE**), Sampling effort (**Rep**), Chlorophyll a concentration (**Chloa**), Mean salinity (**Sal**), Coastal population density (**Pop**), Sea Surface Temperature of the coldest month (**SST.cold**), and the nature of seaweed belt (**SW**), namely *Pelvetia canaliculata* (**SW.Pel**), *Fucus spiralis* (**SW.Fspir**), *Fucus serratus* (**SW.Fser**).




Figure 1. Locations of sampled rocky-shores. The study area crossed 3 ecoregions (the North Sea, the Celtic Sea and the South European Atlantic Shelf) according to the 'Marine Ecoregion Of the World' biogeographic system (Spalding et al., 2007). For each point, labels refer to the vernacular shore name: ADG: "Anse Du Gesclin", AUD: "Audresselles", AYT: "Aytre", BAS: "La Bastille", BDM: "Baie des Marsoins", BFB: "Bas Fort Blanc". BOU: "Les Boucholeurs", BRE: "Bretteville", BSM: "Blainville-sur-mer", CAB: "Cabellou", CAL: "Cale", CDB: "Chef de Baie", CDG: "Cale de Gornevez", CDS: "Concheau de Suzac", CFR: "Cap Fréhel", DEL: "Dellec", DJO: "Dame jouanne", EVE: "L'Eventail", FOG: "Fogeo", FON: "Fort National", GC: "Grande Corbière – Chausey »", GLE: "Glénan (Saint-Nicolas)", GRB: "Grand Bé", HSE: "Houston SE – Chausey", IBO: "Ile de Boed", IDB: "Iles des Ebihens", IDU: "Ile Dumet", ILB: "Ile de Bailleron", ILC: "Ilot de la Colombière", ILM: "Iles aux Mouettes", ISL: "L'Islet", ISM: "Ilot Saint-Michel", IVE: "Ile Verte", KCD: "Kercouedo", KER:

“Kermorvan”, KIB: “KignerBell”, LI: “Longue Ile – Chausey”, LOI: “Loix”, MIN: “Les Minimés”, PAG: “Pors-Ar-Goret”, PDA: “Pointe de l’Arcouest”, PDC: “Pointe du Chay”, PDF: “Pointe de la Fumée”, PDL: “Pointe de la Loge”, PDP: “Pointe des Prouards”, PEA: “Petit Ailly”, PEG: “Père Gus”, PEN: “Penvins”, PIE: “Piegu”, PIS: “La Piscine”, PLA: “Platin”, PM: “Port Marie –Chausey”, PNA: “Port Navalo”, POC: “Pont de la Chaume”, PPL: “Pointe du Plomb”, QUI: “Quiberville”, ROP: “Roche Percée”, ROV: “Roche Vilaine”, SEN: “Saint-Enogat”, SGB: “Saint Guirec/Bastille”, SGO: “Saint Goustan”, SJS: “Saint Jean-des-Sables”, SMM: “Sainte Marguerite-sur-mer”, SNE: “Sud Nellière”, TAR: “La Tara”, TAT: “Tatihou”, TRE: “Trenez”, VEC: “Vert-clos”.




Figure 2. Partial dependence Plot (PdP) representing responses with the seaweed belts (**A**; barplot) and PdP in association with Individual Conditional Expectation (ICE) (**B**; yellow curve + coloured curves, respectively) representing relation with other most relevant predictors: Mean total suspended matter (**TSM.M**), inorganic nitrogen concentrations (**Nit**), Kinetic Energy due to Waves (**WKE**), Sampling effort (**Rep**) and Canopy cover class (**Cover**) of each robust models ($R^2 > 0.35$): taxonomic richness (**S**), total gastropod abundance (**Tot_ab**), *Steromphala umbilicalis* (**S.umbi**), *Patella spp* (**Patella.spp**), the complex *Littorina fabalis*/*L. obtusata* (**L.fab.obt**), *Steromphala pennanti* (**S.penn**), *Phorcus lineatus* (**P.line**), *Littorina littorea* (**L.litto**), complex *Littorina saxatilis*/*L. compressa*

(**L.saxa.comp**) and *Steromphala cineraria* (**S.cine**); Cover class refers to the ordinal cover scale (**1**: 0-5%; **2**:5-25%; **3**:25-50%; **4**:50-75; **5**:75-100%). ICE describes the response for each seaweed belt, while PdP shows the overall average response. At the bottom of the subplot panel, black ticks provide the distribution of each predictor.




Figure 3. Predicted spatial variability of mean taxonomic richness (S, left panel) and mean total abundance (right panel) of gastropods along the rocky shores of the French Atlantic coast. Total abundance colour scale was log-transformed. Squares highlight lowest predicted values retrieved for the estuaries (A) The Seine (49°26'02"N; 0°12'24"E), (B) the Loire (47°30'00"N; 2°28'52"W) and the Vilaine (47°17'23"N; 2°07'28"W) and (C) the Charente (45°57'19"N; 1°04'41"W).




Figure 4. Pearson's correlation matrix of spatial abundance predictions for each response variable. Positive correlations (from orange to red) identify similarity in spatial abundance pattern, while negative ones (from violet to blue) highlight opposition.

CRedit authorship contribution statement

Bruno Serranito: Conceptualization, Methodology, Software, Data curation, Investigations, Visualization, Writing – Original Draft

Tristan Dimeglio: Conceptualization, Funding acquisition, Data curation, Ressource, Investigations, Writing – Review and Editing,

Fred Ysnel: Conceptualization, Supervision, Methodology, Investigations, Writing – Review and Editing

Anne Lizé: Supervision, Writing – Review and Editing

Eric Feunteun: Conceptualization, Supervision, Funding acquisition, Methodology, Investigations, Writing – Review and Editing, Project administration

Declaration of interests

- The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: